Quantitative mechanistic model reveals key determinants of placental IgG transfer and informs prenatal immunization strategies

Remziye Rachel Erdogan[1], Sepideh Dolatshahi*[1,2]

[1]Department of Biomedical Engineering, University of Virginia School of Medicine, Charlottesville, VA, 22908

[2]Carter Immunology Center, University of Virginia School of Medicine, Charlottesville, VA, 22908

*Corresponding author: E-mail: sd8us@virginia.edu

Abstract

Transplacental antibody transfer is crucially important in shaping neonatal immunity. Recently,
prenatal maternal immunization has been employed to boost pathogen-specific immunoglobulin G
(IgG) transfer to the fetus. Multiple factors have been implicated in antibody transfer, but how these
key dynamic regulators work together to elicit the observed selectivity is pertinent to engineering
vaccines for mothers to optimally immunize their newborns. Here, we present the first quantitative
mechanistic model to uncover the determinants of placental antibody transfer and inform
personalized immunization approaches. We identified placental FcγRIIb primarily expressed by
endothelial cells as a limiting factor in this receptor-mediated transfer, which plays a key role in
promoting preferential transport of subclasses IgG1, IgG3, and IgG4, but not IgG2. Integrated
computational modeling and *in vitro* experiments reveal that IgG subclass abundance, Fc receptor
(FcR) binding affinity, and FcR abundance in syncytiotrophoblasts and endothelial cells contribute
to inter-subclass competition and potentially inter- and intra-patient antibody transfer heterogeneity.
We utilize this model as an *in silico* immunization testbed, unveiling an opportunity for precision
prenatal immunization approaches that account for a patient's anticipated gestational length,
vaccine-induced IgG subclass, and placental FcR expression. By combining a computational
model of maternal vaccination with this placental transfer model, we identified the optimal
gestational age range for vaccination that maximizes the titer of antibody in the newborn. This
optimum vaccination time varies with gestational age, placental properties, and vaccine-specific
dynamics. This computational approach provides new perspectives on the dynamics of maternal-
fetal antibody transfer in humans and potential avenues to optimize prenatal vaccinations that
promote neonatal immunity.
22
23

Introduction

Neonates are vulnerable to infections due to their tolerogenic immune phenotype; for the same reason, neonatal vaccinations have been met with limited success to date (1). To provide passive immunity while the neonatal immune system adapts to the environment *ex utero*, maternal immunoglobulin G (IgG) is selectively transferred across the placenta during gestation. Not only do these maternal antibodies confer neonates with passive early life immunity against infectious diseases, but also they shape the trajectory of neonatal immune development by priming the cellular immune response (2–4). Despite this crucial role, the molecular underpinnings of IgG transplacental transport are incompletely described to date.

Though the placenta obstructs maternal-fetal transport of most large molecules, maternal IgG is selectively transported by way of Fc receptor (FcR)-mediated transcytosis. To enter fetal circulation, IgG must undergo transcytosis through two key cellular layers of the placenta: syncytiotrophoblasts (STBs) and endothelial cells (ECs). The neonatal Fc receptor (FcRn) is known to mediate IgG transport through STBs, but whether this mechanism is sufficient for IgG transport through ECs remains unclear (5–7). In addition to FcRn, STBs and ECs express several Fc gamma receptor (FcγR) isoforms with heterogeneous affinity for IgG (8–10). These classical FcγRs may contribute selective placental "sieving" of IgG dependent upon fragment crystallizable (Fc) characteristics, e.g., IgG subclass and N-linked glycosylation (11–14), yet there is conflicting evidence surrounding the involvement of classical FcγRs in IgG transport (9,10,15,16). Recent studies hypothesized a role for STB-bound FcγRIIIa or EC-bound FcγRIIb in IgG transfer (9,12,17–20), yet others showed FcRn is sufficient for IgG transport in isolated placental ECs *in vitro* and in mouse placenta *in vivo* (14,21,22). Though informative, the experimental methods underlying these findings do not capture the complexity or time scale of placental antibody transport. To disentangle the involvement of non-canonical FcγRs in IgG sieving, innovative methods that recapitulate the longitudinal dynamics of this fundamental process are needed.

Placental antibody transfer has been leveraged by maternal prenatal vaccines which boost pathogen-specific IgG transport to the neonate (23). Identifying the ideal immunization strategy to maximize neonatal antibody titers is non-trivial due the growth and development of the placenta which dynamically regulates IgG transport efficiency. These complex dynamics coupled with known maternal immune adaptations during pregnancy define a unique immunization design space which poses a challenge to empirical vaccine optimization in clinical trials alone. Predictive kinetic-dynamic modeling can be leveraged to efficiently design vaccines that maximize IgG transfer to the neonate. The ability to make *in silico* predictions of vaccine-induced antibody transfer would enable pre-clinical dosing strategy studies, expediting translation from bench to bedside.

In the United States, expecting mothers are routinely vaccinated against tetanus, diphtheria, and acellular pertussis (Tdap) during the early third trimester (24), but emerging evidence suggests that this recommendation may not optimally protect the entire population. First, it has been shown that maternal Tdap immunization earlier in gestation results in higher pertussis toxin (PT)- and filamentous hemagglutinin (FHA)-IgG in infants regardless of gestational length (24–26). Second, current vaccination efforts are designed to elicit high antibody transfer to term neonates, yet third trimester immunization may not allot sufficient time for the mother to mount a humoral immune response and transfer antibodies to preterm neonates. Collectively, this evidence supports the potential for personalized vaccine approaches to maximize pathogen-specific IgG transfer to the neonate, especially among premature neonates.

To uncover the molecular regulators of placental antibody transport and to inform the development of novel, personalized immunization approaches, we developed the first computational model of placental IgG transfer. We use this model as an *in silico* testbed for prenatal vaccine design and identify potential strategies to improve transfer of vaccine-induced

antibodies, both at a patient-specific and population level. Ultimately, this model-driven
investigation sheds light on the dynamic regulation of maternal-fetal IgG transfer and provides a
foundation to develop precision vaccine approaches which promote neonatal immunity.

Results

Mechanistic model recapitulates IgG subclass-specific placental transfer

To elucidate mechanisms of IgG transfer and selective "sieving", we devised a dynamic model of
IgG transplacental transfer. The model consists of ordinary differential equations (ODEs)
describing IgG mass transport through the distinct layers comprising the maternal-fetal interface:
STB FcRn-mediated transcytosis, diffusion through intervillous stroma, and EC FcγRIIb-mediated
transcytosis (Fig 1A) (see Appendix S1 for model equations). FcRn is widely implicated in placental
IgG transport; we additionally chose to incorporate FcγRIIb into the model due to its expression in
term placental ECs and demonstrated role in EC transcytosis *in vitro* (9,19,27). Model parameters
were derived from literature when available, and unknown parameters were estimated by fitting
predictions of fetal IgG to measurement data obtained from cordocentesis in human pregnancies
(Fig 1B, Table 1). Parameter estimation was performed using CaliPro as described elsewhere
(28,29) (see Methods). With this optimized parameter set, the model recapitulated dynamics of
both bulk and subclass-specific IgG transfer, predicting the IgG subclass transfer hierarchy most
frequently reported in literature (IgG1 > IgG3 > IgG4 > IgG2) (Fig 1C, S1 Fig) (11,29,30).

Figure 1. Mechanistic model captures IgG subclass-specific transfer dynamics. (A) The schematic
depicts the ODE model representing transport of IgG through the placenta, mapped onto a graphical
representation of the maternal-fetal interface. Reaction rate constant associated with each process is
depicted on the corresponding arrow. Each square represents a molar concentration in each compartment.
"C" denotes complexes of bound IgG and Fc receptor either in STB endosomes ($C_{IgG-FcRn}$) or at the EC
surface ($C_{IgG-FcγRIIb}$). For simplicity, the schematic represents only bulk IgG transport, whereas the model
includes separate equations for each IgG subclass. (B) Simulated fetal IgG subclass levels vs. time (lines)
are plotted overlaid with data used for parameter estimation. The overlaid data are the mean and standard
deviation of IgG subclasses in the umbilical cord between 17-41 weeks gestation (n = 107) measured using
cordocentesis. Gray lines represent 500 perturbation simulations resulting from sampling parameters from the
ranges determined by CaliPro. (C) The swarm charts show the fetal/maternal ratio (F:M ratio) for each IgG

subclass at 40 weeks simulation time resulting from 500 simulations sampling from the optimized range of
parameters (filled symbols). Each symbol represents a single simulation. The bold symbols (black outline)
represent the mean of IgG subclass cord/maternal transfer ratio at parturition from the human study used
during parameter optimization. The swarm charts (hollow symbols) show the mean transfer ratio for each IgG
subclass compiled across cohort studies mined from the literature. Black line depicts the sample mean.
Dashed line at F:M (cord/maternal) ratio = 1 indicates equal concentration in maternal and fetal blood at
parturition.

Table 1. List of optimized model parameters.

Parameter	Value	Units	Interpretation	Source
k_{up}	0.0567-0.0938	L/week	Uptake rate into cells	Optimized
k_t	0.0684-0.1071	L/week	Transcytosis rate	Optimized
k_{deg}	4.1635-5.0418	L/week	Lysosomal degradation rate	Optimized
FcRn	6.54-6.794e-6	M	STB FcRn expression	Optimized
FcγRIIb	3.096-3.607e-6	M	EC FcγRIIb expression	Optimized
$IgG1_0$	3.78e-5	M	Maternal IgG1 concentration	(29)
$IgG2_0$	1.81e-5	M	Maternal IgG2 concentration	(29)
$IgG3_0$	2.35e-6	M	Maternal IgG3 concentration	(29)
$IgG4_0$	2.27e-6	M	Maternal IgG4 concentration	(29)
K_D, FcRn-IgG1	1.25e-8	M	FcRn and IgG1 dissociation constant	(31)
K_D, FcRn-IgG2	2e-8	M	FcRn and IgG2 dissociation constant	(31)
K_D, FcRn-IgG3	3.3e-8	M	FcRn and IgG3 dissociation constant	(31)
K_D, FcRn-IgG4	5e-8	M	FcRn and IgG4 dissociation constant	(31)
K_D, FcγRIIb-IgG1	3.1e-6	M	FcγRIIb and IgG1 dissociation constant	(31)
K_D, FcγRIIb-IgG2	6.8e-6	M	FcγRIIb and IgG2 dissociation constant	(31)
K_D, FcγRIIb-IgG3	1.3e-6	M	FcγRIIb and IgG3 dissociation constant	(31)
K_D, FcγRIIb-IgG4	1.7e-6	M	FcγRIIb and IgG4 dissociation constant	(31)
V_M	6.425-6.7255	L	Maternal blood volume	Optimized
V_{STB}	0.0901-0.0924	L	Total STB endosomal volume	Optimized
V_{STR}	0.0752-0.083	L	Stromal volume	Optimized
V_{EC}	0.0235-0.0275	L	Total EC endosomal volume	Optimized
V_F	0.1534-0.1633	L	Fetal blood volume	Optimized
δ_{Ab}	0.02	L/week	IgG decay rate in the fetus	(32)

Transcytosis rate parameters and Fc receptor expression regulate IgG transfer efficiency and selectivity

To determine the contribution of each parameter to the subclass-specific dynamics predicted by the
model, we performed global sensitivity analyses. We generated 500 perturbed parameter sets using
Latin Hypercube Sampling (LHS) from uniform distributions spanning the maximum likelihood
parameter ranges determined by CaliPro (28,33). For each parameter set, the predicted fetal IgG
at 40 weeks was calculated.   To determine the parameters with the greatest influence on total IgG
transfer (Fig 2A-B) and on differential transfer between IgG subclasses defined as entropy of
subclasses (Fig 2C-D), orthogonalized partial least squares regression (OPLSR) models were
developed (34). Variable importance in projection (VIP) scores revealed that total IgG transfer was
most sensitive to the rate of IgG uptake by STBs and ECs (k_{up}) and FcγRIIb expression (Fig 2B).

The volume of interstitial stroma (v_{STR}) and fetal blood volume (v_f) negatively correlated with IgG
transfer, likely because increasing the volume of either compartment dilutes the IgG concentration.
To reveal parameters that contribute to the difference in IgG transfer among IgG subclasses, we
built a separate OPLSR model predicting the entropy among IgG subclasses (Methods) (Fig 2C-D).
Interestingly, VIP score analysis showed FcγRIIb contributes most to the differential transfer rates
among IgG subclasses. FcγRIIb is a low-affinity IgG receptor with highest affinity for subclasses
IgG3 and IgG4, suggesting that it constitutes a rate-limiting factor of IgG transfer which fine-tunes
the subclass transfer hierarchy. The rate of IgG transcytosis across cells (k_t) also contributes to
subclass-specific transfer; this is in part because k_t acts in an FcR-dependent manner, further
amplifying the effects of Fc-mediated selectivity. Thus, the rate of cargo internalization and
fetoplacental size drives the concentration of bulk IgG in the fetus, whereas FcRn and FcγRIIb
collaborate to fine-tune patterns of IgG subclass-specific transfer.

Fig 2. Key parameters associated with transcytosis mechanism and Fc receptor expression regulate
IgG transfer. (A-B) An OPLSR model was built to determine parameters predictive of total fetal IgG. (A) The
scores plot showing X scores on latent variables 1 and 2 (LV1 and LV2) of simulated fetal IgG concentration
at parturition for 500 perturbation parameter sets. Each circle represents a single perturbation experiment
colored proportional to the concentration of IgG in the fetus at delivery. The model was orthogonalized such
that the direction of maximum variance is in the direction of latent variable 1 (LV1), capturing 99% of the Y
variation. Conversely, LV2 captures the variability in parameters that do not contribute to increased IgG
transfer (Y variation = 0%). (B) The bar graph depicts the variable importance in projection (VIP) scores,
which quantifies the contribution of each parameter to total fetal IgG for a term delivery. VIP scores are
artificially oriented in the direction of their loadings on LV1. VIP score > 1 indicates a greater-than-average
influence on the model. The model performed with high predictability power (Q^2 = 99%). The model had a
mean squared error which outperformed 1000 random models with permuted labels ($p < 0.001$). (C-D) An
analogous OPLSR model was built to predict entropy among fetal IgG subclass levels at the end of gestation.
(C) The scores plot showing X scores on LV1 and LV2 of simulated fetal IgG concentration at parturition for
500 perturbation parameter sets where each circle represents a single perturbation experiment colored
proportional to the computed entropy among IgG subclasses in the fetus at delivery. The model was

orthogonalized such that LV1 captures features associated with the highest entropy, accounting for 87% of
the Y variation. Conversely, LV2 captures the variability in parameters that do not contribute to increased
entropy (Y variation = 0%). (D) The VIP scores bar plot shows the contribution of each parameter to the
entropy of IgG subclasses in fetal blood for a term delivery. VIP scores are artificially oriented in the direction
of their loadings on LV1. VIP score > 1 indicates a greater-than-average influence on the model. The model
performed with high predictability power (Q^2 = 87%). The model had a low mean squared error which
outperformed 1000 models based on randomly permutated labels (p < 0.001).

Endothelial cell-bound FcγRIIb is a key driver of subclass-specific IgG transfer

It is well-established that placental IgG transport increases during the third trimester in humans, so
we asked whether increasing FcγRIIb or FcRn expression in ECs drives increased transport
efficiency. To gain insight into the expression dynamics of the genes coding for FcγRIIb and FcRn,
namely *FCGR2B* and *FCGRT* (FcRn heavy chain), in fetal ECs across gestation, we analyzed two
single cell RNA sequencing (scRNA-seq) data sets from published literature spanning three
gestational time points: (1) first trimester samples from elective abortions (< 12 weeks), (2) early
third trimester samples from preeclamptic cesarean sections (30 weeks), and (3) late third trimester
samples from uncomplicated cesarean sections (38 weeks) (35,36). Interestingly, *FCGR2B*
expression was significantly higher in late compared to early third trimester ECs, but we did not
detect a significant difference in *FCGRT* expression between early and late third trimester ECs (Fig
3A). Conversely, *FCGRT* but not *FCGR2B* transcripts were detected in first trimester ECs, and the
ratio of *FCGR2B* to *FCGRT* expression per cell increased across time points (Fig 3B, S2 Fig),
suggesting that FcγRIIb dynamically regulates IgG transcytosis in ECs.
To assess the contribution of endothelial cell FcγRIIb to IgG transport, we focused on the
role of FcγRIIb as a determinant of subclass-specific transfer. Previous studies found that FcRn
and FcγRIIb were co-expressed by placental ECs (9,37), and FcRn has a noticeably higher affinity
for all IgG subclasses (Table 1). This raises the question of whether FcRn alone is responsible for
IgG transfer across fetal ECs. As FcγRIIb binds subclasses IgG1, IgG3, and IgG4 with greater
affinity than IgG2, we hypothesized that FcγRIIb is necessary to predict preferential transfer of
IgG1, IgG3, and IgG4 over IgG2. To test this hypothesis, we compared model-predicted fetal-
maternal IgG ratios from simulations assuming ECs solely express FcRn (EC_{FcRn}) or FcγRIIb
($EC_{FcγRIIb}$). The EC_{FcRn} model did not capture the preferential transfer of IgG1, IgG3, and IgG4 at
any FcRn expression level (Fig 3C, left). Interestingly, in the $EC_{FcγRIIb}$ model the subclass transfer
hierarchy shifted from IgG3 > IgG1 > IgG4 > IgG2 to IgG1 > IgG2 > IgG3 > IgG4 with increasing
FcγRIIb expression (Fig 3C, right). As the transcriptome analysis demonstrated that *FCGRT* and
FCGR2B can be co-expressed in fetal ECs, we questioned whether this FcγRIIb-dependent
mechanism of subclass-specific transfer is present when FcRn and FcγRIIb are co-expressed
($EC_{FcRn,FcγRIIb}$). In a model where ECs express both FcRn and FcγRIIb, we found that the predicted
subclass transfer hierarchy was IgG1 > IgG3 > IgG4 > IgG2 when FcRn expression was less than
40% of FcγRIIb (S3 Fig), demonstrating that higher relative expression of FcγRIIb to FcRn by ECs
drives subclass-specific IgG placental transfer. Together, these data implicate FcγRIIb in ECs as a
key regulator of antibody transfer dynamics across gestation.

Figure 3. Endothelial cell-bound FcγRIIb is a key driver of subclass-specific IgG transfer. (A) Violin
plots show *FCGR2B* and *FCGRT* transcript levels in preterm, preeclamptic ECs (30 weeks) compared to
term, healthy ECs (>37 weeks) from single cell RNA sequencing data. Black circles represent the sample
median, white circles represent the sample mean. (B) Violin plots show the log-transformed ratio of *FCGR2B*
to *FCGRT* counts in ECs from single cell RNA seq data from two separate cohorts spanning 3 gestational
time points: <16 weeks, 30 weeks, and >37 weeks. Black circles represent the sample median, white circles
represent the sample mean. (C) Model-simulated IgG subclass transfer ratios as a function of EC Fc
receptor expression with ECs expressing only FcRn (left, EC$_{FcRn}$) or FcγRIIb (right, EC$_{FcγRIIb}$). The
corresponding schematics depict the simulation conditions. The shaded region shows the optimized range of
FcγRIIb expression as determined by parameter optimization. Statistical significance in (B-C) was
determined by a Wilcoxon rank sum test (* $p < 0.01$, ** $p < 0.001$, n.s. not significant).

IgG subclasses compete for receptor-mediated transcytosis

IgG subclass abundances vary across individuals, which may contribute to patient-specific
differences in IgG subclass transfer. On the other hand, our finding that FcγRIIb limitation plays a
key role in determining the subclass transfer hierarchy suggests that maternal IgG subclasses
compete for receptor-mediated transcytosis. In agreement with this hypothesis, the model
predicted that all four IgG subclasses transferred less efficiently when mixed compared to each
subclass in isolation of the others (Fig 4A), and the effect of mixing was inversely proportional to
subclass abundance in maternal serum at low FcγRIIb expression levels (S4 Fig). To further
elucidate the role of inter-subclass competition in determining transport efficiency, we explored the
effect of IgG1 on IgG4 transcytosis in ECs. We chose to model IgG1 and IgG4 as a case study
because they have the highest and lowest abundance in maternal serum, respectively, yet both

generally exhibit a cord/maternal ratio > 1 despite their different abundances. Based on a simple
kinetic reaction model, the inter-subclass competition can be described by the following closed-form
equation:

230
$$C_{IgG4-FcRn} = \frac{FcRn_0[IgG4]}{K_{D,IgG4}(1 + \frac{[IgG4]}{K_{D,IgG4}} + \frac{[IgG1]}{K_{D,IgG1}})}$$

232 Where $C_{IgG4-FcRn}$ is the concentration of bound IgG4 and FcRn, $FcRn_0$ is the total concentration of
233 receptor, [IgG1] and [IgG4] are concentrations of maternal IgG1 and IgG4, and $K_{D,IgG1}$ and $K_{D,IgG4}$
234 are their respective dissociation constants (see derivation in Methods). Intuitively, $C_{IgG4-FcRn}$ is a
235 function of IgG1 abundance and affinity for FcRn, such that either increasing [IgG1] or decreasing
236 $K_{D,IgG1}$ (i.e., increasing IgG1-FcRn affinity) decrease $C_{IgG4-FcRn}$ and consequently, IgG4 transcytosis.

237
Fig 4. IgG1 competes with Fc IgG4 transcytosis in ECs. (A) The bar plot shows the effect of mixing IgG
subclasses together simulated in the compartmental placental model. Solid filled bars represent IgG1 and
IgG4 isolated (no competition), and patterned bars represent IgG1 and IgG4 mixed (competition). (B)
Schematic showing a cross-sectional view of the Transwell system and corresponding mechanistic model
representation of FcRn-mediated IgG4 and IgG1 transport through HUVEC. Right panel: Arrows indicate the
direction of transfer (apical to basolateral). Center panel and right panel: Reaction rate constant associated
with each process is depicted on the corresponding arrow. (C) Model-predicted IgG4 transcytosis (y-axis)
with increasing apical concentration (x-axis) (solid line) is plotted against experimental data. The
corresponding experimental conditions are displayed in the panel on the left. Experimental results of IgG1
and IgG4 transcytosis in HUVEC in the Transwell assay system from three independent replicates are shown
with their mean and standard deviation. The dashed line represents the model-predicted and experimentally
validated transcytosis saturation point. (D) Model-predicted IgG4 and IgG1 transcytosis (y-axis) with
increasing apical concentration of IgG1 (x-axis) (solid lines) is plotted against experimental data. The
corresponding experimental conditions are displayed in the panel on the left. Experimental results of IgG1
and IgG4 transcytosis in HUVEC in the Transwell assay system from three independent replicates are shown
with their mean and standard deviation. The dashed line represents the model-predicted and experimentally
validated transcytosis saturation point. Statistical comparisons shown are between IgG4 only (far left graph)
in the absence of IgG1 and all combinations of IgG1 and IgG4 mixed using a one-tailed two-sample t-test (* p
< 0.05, ** p < 0.001, n.s. not significant).

To validate and further characterize this hypothesized competition, we used an *in vitro*
system to model human IgG transcytosis through human umbilical vein endothelial cells (HUVEC)
cultured on Transwell permeable membranes (Fig 4B) (Methods). In parallel, we devised a
mechanistic model of receptor-mediated transcytosis and optimized its parameters by fitting to
temporal data of IgG4 transcytosis in the HUVEC experimental system ([IgG4$_{apical}$] = 0.5 or 0.125
mg/ml) (S5 Fig, Table 2). To recapitulate the physiological setting where FcγRIIb receptor is limiting
similar to the physiological conditions, we varied IgG4 levels to determine the range where the
transferred IgG4 stopped increasing with increasing apical IgG4 levels. Although HUVECs express
FcRn but not FcγRIIb, our findings remain valid because the affinities of IgG1 and IgG4 for either
receptor are on the same order of magnitude and thus the dynamics of subclass competition are
comparable. Our mechanistic model predicted that IgG4 transcytosis saturates at [IgG4$_{apical}$] > 0.33
mg/ml, indicating FcRn limitation; we validated this finding in an analogous *in vitro* experiment in the
Transwell system (Fig 4C). To quantify the effect of IgG1 on IgG4 transcytosis, we simulated the
effect of increasing IgG1 concentration when IgG4 was fixed at a concentration just below the FcRn
saturation point ([IgG4$_{apical}$] = 0.2 mg/ml) (Fig 4C). Our model predicted that IgG4 transcytosis
would not be affected by the presence of IgG1 until the total IgG concentration exceeded the
saturation point—that is, [IgG1$_{apical}$] + [IgG4$_{apical}$] > 0.33 mg/ml. In agreement with our
computational model prediction, increasing IgG1 caused IgG4 transcytosis to decrease only when
the total IgG concentration exceeded the FcRn saturation point (Fig 4C). IgG subclass-specific
transfer is thus a function of subclass abundances and FcR affinity, and competition for FcR-
mediated transcytosis occurs when FcR expression is limited.
278
279   **Table 2. Optimized HUVEC transcytosis model parameters.**

Parameter	Value	Units	Interpretation
FcRn	2200	nM	HUVEC FcRn expression level
K$_{deg}$	0.01	1/minute	Lysosomal degradation rate
K$_t$	2.5 x 10^{-5}	1/minute	Transcytosis rate

Optimal Tdap vaccine schedules depend on gestational age and patient-specific Fc receptor expression

Having uncovered new insights into the dynamic regulation of antibody transfer by FcRs, we sought
to uncover key features influencing the transfer of vaccine-induced antibodies which could be
considered in precision immunization approaches. To this end, we developed an *in silico*
immunization testbed by augmenting the placental antibody transfer model with simulated kinetics
of the maternal antibody response post-immunization (Fig 5A). As a case study of a maternal
vaccine currently in use, we adapted an existing model of plasma cell activation and antibody
secretion following malaria infection to recapitulate IgG responses in women of child-bearing age
following Tdap immunization (Methods) (Fig 5B) (38,39). A dose of antigen representing maternal
immunization was applied to the system at three representative gestational ages t$_{vax}$ = 10, 20, and
30 weeks to stimulate antibody secreting cell proliferation and anti-pertussis toxin IgG (α-PT IgG)
secretion and transfer to the fetus (Fig 5C). Though the correlate of protection for pertussis
infection still remains elusive, some possible roles of IgG in pertussis infection include neutralization
and opsonophagocytosis (40,41). For the purposes of this study, we assumed that neutralizing
antibodies confers protection against pertussis toxin in the neonate and therefore the optimal t$_{vax}$ is
defined as the gestational age at immunization resulting in the highest α-PT IgG titer in the fetus at
birth. Interestingly, the interplay of the immunization time and placenta transfer results in complex
temporal changes in fetal IgG levels, emphasizing the importance of model simulations in predicting
the optimal t$_{vax}$. For example, immunization at t$_{vax}$ = 20 weeks results in higher fetal IgG levels
compared to 10 and 30 weeks for very preterm, preterm and term pregnancies (Fig 5C). We then

ran *in silico* simulations over a range of possible immunization times to determine the optimal t_{vax} for
newborns born at different weeks of gestation (Fig 5D).

306
Figure 5. Optimal Tdap immunization time is gestational age dependent. (A) Schematic shows
mechanistic model of maternal plasma cell antibody secretion post-vaccination. Vaccine-induced maternal
IgG kinetics are integrated into the maternal compartment of the placental IgG transport model, shown in the
righthand box. Dashed arrows indicate regulatory processes, solid lines represent processes modeled using
mass action kinetics. L_{ASC}: Long-lived antibody secreting cell, S_{ASC}: short-lived antibody-secreting cell, Ag:
antigen. (B) Simulated maternal anti-pertussis IgG concentrations vs. time is plotted with a solid line and
overlaid with measured IgG concentrations following Tdap vaccination in women of childbearing age from
published literature (n = 105). The 95% confidence intervals are shown with light gray shading. (C)
Simulated fetal anti-pertussis toxin IgG concentrations are plotted against time following maternal vaccination
at three sample vaccination times (indicated along the x-axis): t_{vax} = 10, 20, and 30 weeks gestational age.
Shaded regions correspond to gestational length groups (left to right): extremely preterm, very preterm, late
preterm, term. (D) Simulated anti-pertussis toxin IgG in the fetus at the time of delivery (y-axis) is plotted over
a range of simulated vaccination times (x-axis) and gestational ages (color intensity, color bar on right). The
optimal vaccination time and corresponding fetal IgG levels are marked with a circle. The current CDC-
recommended maternal vaccination window is highlighted in red, and the model-predicted optimal vaccination
range encompassing all gestational age groups is highlighted in green.

324   **Table 3. Pertussis immunization model parameters.**

Parameter	Value	Units	Interpretation	Source
k_{ASC}	0.6	1 / 10^6 PBMCs / day / Ag	Rate of plasma cell generation	Optimized
k_{Ab}	0.0475	mg / ml * (10^6 PBMCs) / days	Rate of antibody production by plasma cells	Optimized
Ag_0	100	mg/ml	Initial dose of antigen	Optimized
ρ	96	%	Proportion of short-lived ASCs (S_{ASC})	(38)
δ_{Ag}	0.15	1 / day	Antigen decay rate	(42)
δ_{Ab}	0.033	1 / day	Antibody decay rate	(38)
δ_{S-ASC}	0.578	1 / day	Short-lived antibody-secreting cell (S_{ASC}) decay rate	(38)
δ_{L-ASC}	6.6×10^{-4}	1 / day	Long-lived antibody secreting cell (L_{ASC}) decay rate	(38)

325
326

Empirical studies have shown that earlier vaccination may be an effective strategy to protect
preterm neonates who typically receive a lower quantity of maternal antibodies (24–26). Using our
immunization testbed, we first asked how the optimal Tdap administration schedule varies for the
average patient delivering at different gestational time points (i.e., term, late preterm, very preterm,
or extremely preterm). In the case of full term gestation, the model-predicted optimal t_{vax} was during
the 26^{th} week of gestation, but Tdap immunization as early as the 10^{th} week still resulted in two-
thirds of the optimal IgG concentration being transferred to term newborns (Fig 5D). Optimal t_{vax}
scaled with increasing gestational length, suggesting that risk for preterm birth should be
considered when determining prenatal immunization schedules (Fig 5D). Strikingly, we found the
optimal t_{vax} fell during the second trimester (between 14-26 weeks) regardless of preterm birth
status, which is earlier than the window recommended by the Centers for Disease Control and
Prevention (CDC) (28-34 weeks) and in agreement with studies reporting higher titer of pertussis
antibodies in the fetus following maternal vaccination during second trimester as opposed to third
trimester (24,26,43). Thus, optimal immunization schedules are gestational length-dependent, but
a specific vaccination window of opportunity during gestation could be an effective population-level
strategy to ensure coverage for preterm and term newborns alike.
Other sources of patient-to-patient variability that could influence optimal immunization
schedules include differences in placental transfer efficiency linked to FcR-mediated transcytosis
and the quality and quantity of IgG induced by the vaccine. To determine key variables that should
be considered in patient-specific immunization approaches, we performed a global sensitivity
analysis to uncover parameters most influential on optimal t_{vax}. We optimized t_{vax} in simulations
from 200 perturbed parameter sets sampled from a uniform distribution plus or minus 10% of each
parameter's optimized value, then built an OPLSR model to predict the optimal t_{vax} given each
parameter set (Fig 6A, 6B). Parameters which were associated with later immunization times were
the rate of IgG internalization (k_{up}), FcγRIIb, and volume of STB endosomes (v_{STB}), whereas
parameters associated with earlier immunization times were the stromal volume (v_{STR}), maternal
plasma volume (v_M), and FcRn (Fig 6B). Intuitively, high maternal and stromal volumes decrease
transfer of vaccine-induced antibodies because IgG concentration is diluted in high volume
compartments. On the other hand, later vaccination is favorable with increased endosomal volume
in STBs and the IgG internalization rate because these parameters increase the overall efficiency of
STB transcytosis, facilitating rapid transport of IgG just prior to delivery and minimizing the time IgG
is subject to degradation in the fetus whilst *in utero*. Therefore, differences in placental size and

function across the population may play a role in determining the ideal patient-specific immunization
program.

Figure 6. Optimal vaccination schedules depend on patient-specific placental and maternal
parameters. (A-B) The OPLSR model predicts the optimal immunization time (t_{vax}, weeks gestational age)
from perturbed parameter sets. (A) The scores plot showing X scores on LV1 and LV2 of the optimal t_{vax} for
200 perturbed parameter sets. Each circle represents a single perturbation experiment colored proportionally
to the predicted optimal t_{vax}. The model was orthogonalized such that LV1 is in the direction of maximum
variance, capturing 76% of the Y variation. Conversely, LV2 captures the variability in parameters that do not
contribute to increased IgG transfer (Y variation = 0%). (B) The bar plot shows VIP scores that quantify the
contribution of each parameter to the optimal t_{vax}. VIP scores are artificially oriented in the direction of their
loadings on LV1. VIP score > 1 indicates a greater-than-average influence on the model. The model
performed with high predictability power (Q^2 = 75%) and with a mean squared error lower than 1000 models
based on randomly permuted labels (p < 0.001). (C) The contour map shows the percent change in anti-
pertussis toxin IgG (α-PT IgG) in a fetus delivered at 40 weeks relative to the baseline simulation with varying
FcRn and FcγRIIb expression. Blue and red correspond to a decrease or increase in fetal IgG relative to the
baseline simulation. FcRn and FcγRIIb are shown on a logarithmic scale. The bold indigo and turquoise lines
indicate the optimized FcRn and FcγRIIb expression levels, and the intersection point indicates the baseline
simulation using parameters estimated by fitting to the average data. The curves plotted to the right are one-
dimensional traces indicated by the bold lines on the two-dimensional contour plot and are indicative of
simulations where either FcγRIIb (left) or FcRn (right) are held at their optimized expression level and the
other receptor is allowed to vary. The dashed line represents the baseline simulation, and the vertical line
shows the concentration of FcγRIIb or FcRn at the baseline simulation. The corresponding circle is marked
with a bold outline. (D,E) The optimal t_{vax} and corresponding anti-pertussis toxin IgG at term delivery is given
for a range of FcγRIIb (D) and FcRn (E) expression levels below their optimized values. The points are
connected to show the trajectory of optimal t_{vax} with increasing FcγRIIb or FcRn. FcγRIIb and FcRn are
indicated by color intensity of the markers. The baseline simulation is marked in both graphs with a bold
outlined circle. (F) A range of t_{vax} are shown for simulations with different compositions of vaccine-induced
IgG subclasses. Simulated conditions were vaccine-induced IgG consisting of entirely IgG1, IgG2, IgG3, or
IgG4, or with equal parts IgG1, IgG2, IgG3, and IgG4. The optimal t_{vax} and corresponding IgG titer for a given
subclass (or mixture of subclasses) for each condition is marked with a circle and labeled on the x-axis.

To investigate the differential impact of FcRn and FcγRIIb on t_{vax} optimization, we simulated
immunization at the optimal t_{vax} for term births determined previously (Fig 5D) for mothers with a
range of FcRn and FcγRIIb expression. By examining the percent change in pertussis toxin IgG
transfer for variations in FcRn and FcγRIIb relative to the average patient, we observed that the

greatest increase in α-PT IgG transfer is achieved when both FcRn and FcγRIIb expression levels
are high, but IgG transfer decreased when either FcRn or FcγRIIb expression decreased (Fig 6C).
To determine whether patients with a deficiency in FcRn or FcγRIIb would benefit from a different
immunization schedule than the average patient, we optimized t_{vax} at FcRn and FcγRIIb levels
below their average values. As seen in the OPLSR analysis in (Fig 6A-B), this confirmed that high
FcγRIIb expression corresponds to later optimal t_{vax}, whereas high FcRn expression corresponds to
earlier optimal t_{vax} (Fig 6D, 6E). Underlying this disparate role of FcRn and FcγRIIb in vaccine
optimization, we found that increasing FcRn causes IgG to build up in the placental villous stroma,
but FcγRIIb is the ultimate rate-limiting factor that determines whether IgG enters fetal circulation.
High FcRn expression increases STB transcytosis efficiency and causes IgG to accumulate in the
stroma "at the ready" for FcγRIIb-mediated transport, enabling FcγRIIb to transfer IgG at its
maximum capacity. Conversely, when FcγRIIb expression is high, it is preferable to vaccinate later
in pregnancy such that the peak maternal IgG post-immunization occurs when STBs express more
FcRn to transport IgG into the stroma. We additionally found that the optimal t_{vax} is further
influenced by the subclass of vaccine-induced IgG, a critical insight given that different pathogens
are known to elicit different IgG subclasses (Fig 6F). These observations indicate that the role of
FcγRIIb as the key rate-determining factor of IgG transfer is pertinent to the design of prenatal
immunization strategies, such that increasing FcRn increases IgG transfer only when FcgRIIb is
sufficiently expressed.
412

Discussion

414
Placental antibody transfer is key to protect neonates against infection and to shape the trajectory
of early-life immune development. Recently, maternal prenatal immunization has been employed to
boost antigen-specific antibody transfer to the fetus, providing targeted immunity to the newborn.
Current prenatal vaccine regimens are applied at the population level, but variation in gestational
length, placental size and function, and maternal immune response presents an opportunity for
patient-specific vaccine approaches. To this end, we developed a computational model to elucidate
the features regulating maternal-fetal IgG transport and to develop novel vaccine strategies for
precision prenatal immunization.
Our modeling approach enabled longitudinal probing of placental antibody transfer
dynamics, shedding light on an otherwise black-boxed system. In contrast to prior studies
attempting to uncover mechanisms of selective antibody transfer through a variety of *in vitro*, *ex*
vivo, and *in vivo* approaches, our computational approach recapitulates this dynamic process as it
occurs in humans. Our dynamic model predicted subclass-specific IgG transfer ratios that agree
with the trend most commonly reported in the literature: IgG1 > IgG3 > IgG4 > IgG2. While the
hierarchy was preserved, the model overestimated IgG3 transfer while underestimating IgG2 and
IgG4 transfer. These subtle discrepancies may point to the involvement of other FcγRs such as
FcγRIIIa (12,44), Hofbauer cells (7,9,16,20,45), or avidity effects that might arise from multivalent
IgG-antigen complexes in fine-tuning antibody transfer dynamics (16,30). Notably, we disentangled
the contributions of FcRn and FcγRIIb to EC transcytosis, revealing that in a baseline average
placenta FcγRIIb drives IgG transcytosis in ECs and may play a key role in dynamic regulation of
IgG transfer across gestation. Together, these observations elucidate the potential role of non-
canonical FcγRs driving antibody transfer dynamics, though the role of FcγRIIb must be validated
mechanistically *in vitro* to bolster our model-driven predictions.
Another key insight emerging from this analysis is the contribution of FcRs to transfer
selectivity for particular IgG subclasses. In particular, subclass-specific IgG transfer is a function of
FcR expression level, maternal IgG subclass abundance, and Fc-FcR interaction affinity. Thus, the
subclass transfer hierarchy of IgG1 > IgG3 > IgG4 > IgG2 can be explained by each subclass'

relative abundance and affinity for FcRn and FcγRIIb. Not only that, but inter-patient variability in
subclass abundances, FcR affinity arising from differential IgG glycosylation, and placental FcR
expression can explain discrepancies in this subclass transfer hierarchy across individuals and
patient cohorts.
The findings of this study have broad implications for prenatal vaccination, both at the
population level and as a basis for patient-specific vaccine approaches. First, the model predicted
that the optimal pertussis immunization time for an average patient is during the second trimester
for term and preterm neonates alike, highlighting an opportunity to amend currently enforced
vaccine regimens recommending Tdap immunization during the late second or early third trimester.
From a precision medicine standpoint, recent studies suggest that careful modulation of vaccine
parameters including the nature of the target antigen, conjugate, number of doses, route of entry,
and choice of adjuvant can fine-tune the pool of maternal vaccine-induced antibodies and the anti-
infection capabilities of the IgG transferred to the newborn. For example, protein antigens tend to
induce an IgG1 or IgG3 response, while polysaccharide antigens tend to induce an IgG2 response
(46); more recent studies suggest that the N-glycan profile of vaccine-induced IgG can be
modulated by adjuvant, providing a means to fine-tune Fc receptor binding potential (47,48). The
model-driven finding that IgG competes for placental transfer contingent upon subclass abundance,
FcR expression, and Fc-FcR affinity provides a basis for improving vaccine formulations and dosing
strategies to optimize vaccine-induced IgG transfer efficiency. Moreover, our model captures
subclass-selectivity and subclass competition in placental transfer and is thus able to quantitatively
assess the effect of a perturbed maternal pool of antibodies due to various immunodeficiency
disorders (49), autoimmune disorders and other disorders that modify the maternal pool of
antibodies and vaccine response. As such, our model has immense potential for devising vaccines
for mothers with specific immunodeficiency disorders and in devising vaccine strategies aimed at
optimizing user-defined selective set of antibodies in contrast with our objective of maximizing
antigen-specific antibody titers in this study.
While many intriguing insights emerged from this model-driven investigation, there are a few
limitations to consider. First, the current model relies on the assumption that all IgG is monovalent
with a uniform glycan profile. This is an oversimplification as IgG glycosylation undergoes a
dynamic shift during pregnancy (50), which can affect the Fc receptor-mediated transfer (12).
Furthermore, in addition to monomeric IgG, IgG can potentially be transferred as antigen-antibody
complexes, which is currently understudied and is not incorporated in our current model (16,30).
Second, there is a stark lack of data surrounding human placental development and dynamic fetal
IgG levels throughout gestation; many model parameters were unavailable in the literature,
necessitating *in silico* parameter optimization. Consequently, we assumed several model
components to be constant, including maternal, fetal, and placental volumes and maternal IgG
levels. In reality, these features undergo rapid remodeling and growth across gestation and
potentially fine-tune antibody transfer dynamics. Future iterations of the model may benefit from
more physiologically accurate dynamics of the developing maternal-fetal interface. To facilitate such
a model, more studies detailing the dynamics of placental growth—especially as it pertains to
vascularization and trophoblast differentiation to form STB layers—and of FcR expression at the
protein level are warranted. To infer the role of these dynamic placental changes in antibody
transfer, placental samples should be analyzed in conjunction with antibody profiling in matched
maternal and umbilical cord serum.
Despite its limitations, a key strength of our computational approach is the ability to simulate
placental transport dynamics which are prohibitively challenging to study *in vivo*. Using tightly
integrated modeling and experimental methods, we probed placental transport across time and size
scales, enabling a more holistic view of the system compared to previous studies in the field.
Traditionally, placental IgG transport efficiency is benchmarked by a simple ratio of maternal to
umbilical cord IgG titer at birth, in part due to its ease of measurement and interpretability. In
addition to masking the nuances of maternal pool of antibodies, this metric insufficiently describes

the highly dynamic and cumulative process of antibody transfer and is thus not capable of
describing: (1) important concepts such as placental FcR limitation and antibody subclass
competition, (2) the interplay between the maternal pool of antibodies and the vaccine-induced
maternal antibodies, and (3) the role of the complex placental functions that rapidly evolve
throughout gestation, among other reasons. Our mathematical model challenges this metric as a
true measure of "efficiency". With the advent and accessibility of state-of-the-art high-throughput
measurements, it is increasingly feasible to probe placental function and define one or more
quantitative measures of transfer efficiency that are conditioned on the maternal pool of antibodies
and are linked with distinct model parameters that account for placental size, perfusion, and Fc
receptor expression.
This study represents a major thrust forward in the field of maternal-fetal immunology and
paves the way towards next generation vaccination programs to fine-tune neonatal immunity
against a broad range of pathogens. As current vaccination regimens are continually revised and
several novel vaccines are currently in clinical trials, now an opportune time to increase the
prevalence of computational systems biology approaches in maternal-fetal medicine to improve
outcomes for vulnerable newborns.
509
510

Methods

Mechanistic model of placental IgG transport

An ordinary differential equation (ODE) model was formulated to simulate mass transport of IgG1, IgG2, IgG3, and IgG4 across the maternal-fetal interface (Appendix S1). A schematic of the model is given in Figure 1A. Fundamental model assumptions are as follows: (i) all processes obey the law of mass action; (ii) IgG binds FcRn within STB endosomes at an acidified pH and FcγRIIb at the EC surface; (iii) binding with either receptor initiates transcytosis; (iv) the effects of FcRn-mediated IgG recycling to the apical STB surface are negligible; (v) unbound IgG in STB endosomes undergoes lysosomal degradation; (vi) all blood volumes, IgG subclass concentrations in maternal blood, and IgG-FcR affinities remain constant throughout gestation; (vii) rate parameters (k_{up}, k_{trans}, and k_{deg}) are equal in STB and EC layers and across IgG subclasses; (viii) stromal cells (e.g., Hofbauer cells, fibroblasts) do not interact with IgG or affect its transfer; (ix) the effect of Fab-antigen interactions is neglected such that all IgG is monovalent; (x) FcR expression increases parabolically across gestation (51) informed by FcRn expression trends in rat placenta; (xi) STBs exclusively express FcRn and ECs exclusively express FcγRIIb. All model development and corresponding *in silico* experiments were performed in MATLAB R2022a (Mathworks).

Parameter estimation

Model parameters were derived from literature when possible, including K_D parameters, maternal IgG concentrations, and all compartment volumes (29,31,52,53). Remaining parameter ranges were determined using CaliPro, a flexible optimization software implemented in MATLAB and described by Joslyn et al (28). A key strength of CaliPro is its ability to home in on a biologically relevant range of parameters that recapitulate the system dynamics rather than minimizing a simple cost function. Briefly, we implemented CaliPro using Latin Hypercube Sampling (LHS) from a physiologically plausible range of parameter values as determined by benchmarking literature studies. We defined the pass criteria as model simulations within the upper standard deviation of the data set at the first time point, and are within 1.5 times the standard deviation at the final time point. Our termination criteria was set to 95% of model runs meeting the pass criteria.

Global sensitivity analysis

Global sensitivity analysis was performed using an orthogonalized partial least squares regression (OPLSR) framework, as described by Surve et al (34). 500 perturbed parameter sets were generated by LHS from a uniform distribution spanning the range for each parameter obtained from CaliPro. An OPLSR model was constructed using the matrix of 500 permuted parameter sets to predict the total concentration of IgG in the fetus at 40 weeks gestation. A second OPLSR model was constructed in the same manner to predict entropy among IgG subclasses in the fetus at 40 weeks gestation, defined as:

$$P_{IgGi} = \frac{IgG_i}{\sum_{i=1}^{4} IgG_i}$$

$$E = -\sum_{i=1}^{4} P_{IgGi}\log\left(P_{IgGi}\right)$$

554 Where E is entropy and P_{IgGi} is the proportion of each subclass in the fetal blood. Model
555 orthogonalization was carried out based on the published method (54). OPLSR model significance
556 was determined empirically by comparing its predictive power (Q^2) against randomly permuted
557 models.

HUVEC transcytosis assay

Antibody preparation

Purified human polyclonal IgG1 and IgG4 (Abcam, AB90283, AB183266) were dialyzed in 1x PBS
using Slide-A-Lyzer mini dialysis units with a 10K molecular weight cutoff (Invitrogen, 88404) to
remove sodium azide. Dialysis units were incubated at 4°C for 2 hours, then dialysis buffer was
changed and incubated again at 4°C for 2 hours. Protein was removed from the dialysis unit using
a micropipette and quantified using a NanoDrop Microvolume Spectrophotometer (Thermo Fisher).

Cell culture

6.5 mm diameter Transwell inserts (Corning, CLS3422) coated in 1.4% gelatin were seeded with
primary human umbilical vein endothelial cells (HUVEC) (Promocell C-12200, kindly provided by
Dan Gioeli at UVA) at a density of 1500 cells/mm^2 (50,000 cells/insert). Cells were cultured in
complete endothelial cell growth media (Promocell, C-22110) and incubated at 37°C and 5% CO_2.
After 24 hours, the cells were washed in sterile PBS and the media was changed.

Transcytosis assay

After 48 hours, the cells were washed in sterile PBS and media containing purified human IgG was
added to the apical chamber. Cells were incubated at 37°C and 5% CO_2 for 120 minutes. IgG1
and IgG4 in the basolateral media were quantified using human enzyme-linked immunosorbent
assay (ELISA) IgG1 and IgG4 quantification kits (Invitrogen, EHIGG1 and BMS2095) according to
the manufacturer's protocol. All conditions were run in triplicate and ELISA measurements were
run in duplicate. A 5-parameter logistic standard curve was generated for each assay. Absorbance
was read at 450 nm with a correction of 590 nm using an Optima microplate reader within 1 hour of
completing the assay. Statistical significance was determined with a one-tailed two-sample t-test
comparing each sample containing IgG1 and IgG4 to the sample with IgG4 but without IgG1.
  To confirm monolayer formation, membrane permeability was quantified for each assay by
addition of FITC-dextran (4000 MW, Sigma Aldrich 46944) (0.5 mg/ml) to the apical chamber of
each insert. FITC-dextran added to inserts without gelatin coating or HUVEC served as a negative
control. The basolateral media was sampled after 120 minutes, and fluorescence was read using
the Optima microplate reader. Monolayer permeability (%) was defined as:

$$Permeability = \frac{[basolat.\,FITC - dextran]_{HUVEC}}{[basolat.\,FITC - dextran]_{blank}} \times 100$$

Samples with FITC-dextran permeability greater than 3 standard deviations from the mean were
excluded based on faulty monolayer formation (S5 Fig).

Mechanistic model of HUVEC transcytosis *in vitro*

To simulate the dynamics of IgG1 and IgG4 receptor-mediated transcytosis in HUVEC, a kinetic-dynamic model of FcRn-IgG binding and transcytosis in endothelial cells was formulated:

$$\frac{dIgG_1}{dt} = -k_{on,1}[IgG_1][FcRn] + k_{off,1}[C_1] \quad (1)$$

$$\frac{dIgG_4}{dt} = -k_{on,4}[IgG_4][FcRn] + k_{off,4}[C_4] \quad (2)$$

$$\frac{dFcRn}{dt} = -[FcRn]\big(k_{on,1}[IgG_1] + k_{on,4}[IgG_4]\big) + \big(k_{off,1} + k_T\big)[C_1] + \big(k_{off,4} + k_T\big)[C_4] \quad (3)$$

$$\frac{dC_1}{dt} = k_{on,1}[IgG_1][FcRn] - \big(k_{off,1} + k_T\big)[C_1] \quad (4)$$

$$\frac{dC_4}{dt} = k_{on,4}[IgG_4][FcRn] - \big(k_{off,4} + k_T\big)[C_4] \quad (5)$$

$$\frac{dIgG_{T,1}}{dt} = k_T[C_1] \quad (6)$$

$$\frac{dIgG_{T,4}}{dt} = k_T[C_4] \quad (7)$$

Where [IgG1] and [IgG4] are apical concentrations of IgG1 and IgG4, [FcRn] is the concentration of FcRn expression in HUVEC, $[C_1]$ and $[C_4]$ are the concentration of bound IgG-FcRn complexes, k_T is the rate of transcytosis, k_{on} is the forward rate of reaction, and k_{off} is the reverse rate of reaction (Table 2). The model parameters were optimized by fitting to dynamic data of IgG4 transcytosis in HUVECs collected at 15, 30, 60, and 120 minutes ([IgG4$_{apical}$] = 0.5 and 0.125 mg/ml) (S5 Fig). To simplify this model and reveal the effect of subclass-subclass competition in Fc receptor-mediated transcytosis, we derived the closed form of equation (5). We assumed that IgG1 and IgG4 are present in excess of FcRn, the system is at quasi-steady state such that

$$\frac{dC_1}{dt} = \frac{dC_4}{dt} = 0 \quad (8)$$

Where C is the concentration of IgG-FcRn complexes. Under these assumptions, equations (4) and (5) can be rearranged as

$$\frac{k_{on,1}[IgG_1][FcRn]}{k_{on,1}[IgG_1][FcRn]\big(k_{off,1} + k_T\big)} = [C_1] \quad (9)$$

$$\frac{k_{on,4}[IgG_4][FcRn]}{k_{on,4}[IgG_4][FcRn]\big(k_{off,4} + k_T\big)} = [C_4] \quad (10)$$

The assumption is then made that the total concentration of receptor does not change such that

$$[FcRn] = [FcRn_0] - [C_1] - [C_4] \quad (11)$$

Substituting (11) into (10):

$$\frac{k_{on,4}[IgG_4]\big([FcRn_0] - [C_1] - [C_4]\big)}{k_{on,4}[IgG_4][FcRn]\big(k_{off,4} + k_T\big)} = [C_4] \quad (12)$$

And (9) into (12):

637
$$\frac{k_{on,4}[IgG_4]\left([FcRn_0] - \frac{k_{on,1}[IgG_1][FcRn]}{k_{on,1}[IgG_1][FcRn]\left(k_{off,1}+k_T\right)} - [C_4]\right)}{k_{on,4}[IgG_4][FcRn]\left(k_{off,4}+k_T\right)} = [C_4] \quad (13)$$

639 Solving (13) for [C_4] yields the closed-form solution for IgG4-FcRn complex formation (55):

641
$$C_4 = \frac{FcRn_0[IgG_4]}{K_{D,4}\left(1 + \frac{[IgG_4]}{K_{D,4}} + \frac{[IgG_1]}{K_{D,1}}\right)} \quad (14)$$

643 Which quantifies the effect of competition between two IgG subclasses undergoing transcytosis in
644 the presence of one Fc receptor.

Pertussis immunization simulations
646

A model of plasma cell activation and IgG secretion in response to antigen stimulus was adapted
from a previous study by White et al in the context of malaria (38). Parameters were tuned to fit
previously published dynamic data of the maternal IgG response to Tdap booster vaccination (39)
(Table 3). We assumed several parameters which depend on the host environment—namely,
systemic cytokine concentrations and availability of FcRn for IgG recycling and half-life extension—
do not depend on antigen-specificity and thus are conserved between malaria and pertussis
contexts (56,57). Such parameters include the proportion of long-lived antibody secreting cells
(L_{ASC}) to short-lived antibody secreting cells (S_{ASC}), the lifespan of L_{ASC} and S_{ASC}, and the antibody
decay rate and were preserved from White et al (38). On the other hand, parameters which depend
on the plasma cell niche and context of the specific antigenic challenge (e.g., rates of plasma cell
proliferation, antibody secretion, and the antigen decay rate) were optimized to fit titers of anti-
pertussis toxin IgG (α-PT IgG) in women of childbearing age following a booster Tdap vaccination
(39) (Table 3). Other important model assumptions include: (i) IgG has a half-life of 31 days in the
fetus; (ii) the dynamics of B cell activation—including antigen presentation by follicular dendritic
cells and cytokine secretion by T cells—and the contribution of memory B cells are neglected; (iii)
mothers have been previously exposed to pertussis toxin (i.e., this is a booster vaccine); (iv) all
mothers have the same dynamic response to immunization; (v) Immunization resulted in equal
parts IgG1, IgG2, IgG3, and IgG4 with uniform glycosylation profile (i.e., glycosylation does not
affect FcRn nor FcγRIIb binding) and immunoglobulin isotype switching was neglected; (vi) L_{ASC}s
and S_{ASC}s secrete IgG with identical subclass and glycosylation profile; (vii) the peak antibody titer
between 14 and 28 days post-immunization; (viii) the IgG response to pertussis is the same in
pregnant and non-pregnant individuals(58). Maternal immunization was simulated by stimulating the
maternal compartment with antigen at a time t_{vax} (given in weeks gestational age) to induce an
antibody response, which directly fed into the maternal compartment of the placental IgG transport
model. To determine the optimal t_{vax} corresponding to maximum pertussis-toxin IgG transfer to the
fetus, iterative simulations were performed over a range of t_{vax} spanning 10-38 weeks gestational
age. Sensitivity of the optimal t_{vax} was determined by parameter perturbation and subsequent
PLSR model construction, as described above. When perturbing the IgG subclass composition
induced by immunization, it was assumed that that vaccine induced solely IgG1, IgG2, IgG3, or
IgG4.

Acknowledgements
679

We thank the members of the Dolatshahi lab, Drs. Donald J. Dudley and Loren D. Erickson for helpful discussion and feedback. We would also like to thank Dr. Daniel Gioeli and Devin Roller for providing cell lines and support with Transwell assays. We would like to acknowledge Drs. Yuk Ming Dennis Lo, Peiyong Jiang, and Xi Hu from the Chinese University of Hong Kong for sharing their published single cell transcriptomic data of placental tissue (59). Research reported in this publication was supported by the Thomas F. and Kate Miller Jeffress Memorial Trust and Bank of America under Jeffress Trust Award in Interdisciplinary Research as well as the National Institute of General Medical Sciences of the National Institutes of Health under Systems & Biomedical Data Science training program (T32-GM145443). The content is solely the responsibility of the authors and does not necessarily represent the official views of the funding sources and National Institutes of Health.

Data and code availability

All data files and codes used to run the model can be found at our GitHub: https://github.com/Dolatshahi-Lab/Transplacental-antibody-transfer-model. Model equations and *in vitro* Transwell assay data are available as Supplemental files.

References

1.   Pichichero ME. Challenges in vaccination of neonates, infants and young children. Vaccine. 2014 Jun 30;32(31):3886–94.

2.   Olin A, Henckel E, Chen Y, Lakshmikanth T, Pou C, Mikes J, et al. Stereotypic Immune System Development in Newborn Children. Cell [Internet]. 2018;174(5):1277-1292.e14. Available from: https://doi.org/10.1016/j.cell.2018.06.045

3.   Simister E, Israel EJ, Patel VK, Taylor SF, Marshak-Rothstein A. neonatal mice. acquisition of maternal IgG by fetal and 2-microglobulin-associated Fc receptor for Requirement for a beta. 1995 [cited 2022 Oct 19]; Available from: http://www.jimmunol.org/content/154/12/6246

4.   Marchant A, Kollmann TR. Understanding the ontogeny of the immune system to promote immune-mediated health for life. Front Immunol. 2015;6(FEB):77.

5.   Brambell FW. THE TRANSMISSION OF IMMUNITY FROM MOTHER TO YOUNG AND THE CATABOLISM OF IMMUNOGLOBULINS. Lancet. 1966 Nov 19;288(7473):1087–93.

6.   Wilcox CR, Holder B, Jones CE. Factors affecting the FcRn-mediated transplacental transfer of antibodies and implications for vaccination in pregnancy [Internet]. Vol. 8, Frontiers in Immunology. Frontiers Media S.A.; 2017 [cited 2021 Jan 15]. p. 1. Available from: www.frontiersin.org

7.   Simister NE. Placental transport of immunoglobulin G. In: Vaccine. Elsevier BV; 2003. p. 3365–9.

8.   Bruhns P, Iannascoli B, England P, Mancardi DA, Fernandez N, Jorieux S, et al. Specificity and affinity of human Fc receptors and their polymorphic variants for human IgG subclasses. 2009 [cited 2021 Mar 16]; Available from: http://ashpublications.org/blood/article-pdf/113/16/3716/1308575/zh801609003716.pdf

9.   Bright NA, Ockleford CD, Anwar M. Ontogeny and distribution of Fc gamma receptors in the human placenta. Transport or immune surveillance? J Anat [Internet]. 1994 Apr [cited 2021 Jun 16];184 (Pt 2(Pt 2):297–308. Available from: http://www.ncbi.nlm.nih.gov/pubmed/8014121

10.   Kameda T, Koyama M, Matsuzaki N, Taniguchi T, Saji F, Tanizawa O. Localization of

728		three subtypes of Fcγ receptors in human placenta by immunohistochemical analysis. Placenta. 1991 Jan 1;12(1):15–26.
730	11.	Clements T, Rice TF, Vamvakas G, Barnett S, Barnes M, Donaldson B, et al. Update on Transplacental Transfer of IgG Subclasses: Impact of Maternal and Fetal Factors. Front Immunol. 2020;11(September):1–17.
733	12.	Jennewein MF, Goldfarb I, Dolatshahi S, Cosgrove C, Noelette FJ, Krykbaeva M, et al. Fc Glycan-Mediated Regulation of Placental Antibody Transfer. Cell [Internet]. 2019;178(1):202-215.e14. Available from: https://doi.org/10.1016/j.cell.2019.05.044
736	13.	Dolatshahi S, Butler AL, Pou C, Henckel E, Bernhardsson AK, Gustafsson A, et al. Selective Transfer of Maternal Antibodies in Preterm and Term Children. 2021 Dec 1 [cited 2022 Jun 7]; Available from: https://www.researchsquare.com
739	14.	Borghi S, Bournazos S, Thulin NK, Li C, Gajewski A, Sherwood RW, et al. FcRn, but not FcγRs, drives maternal-fetal transplacental transport of human IgG antibodies. Proc Natl Acad Sci U S A. 2020;117(23):12943–51.
742	15.	Martinez DR, Fouda GG, Peng X, Ackerman ME, Permar SR. Noncanonical placental Fc receptors: What is their role in modulating transplacental transfer of maternal IgG? Sheppard DC, editor. PLOS Pathog [Internet]. 2018 Aug 30 [cited 2021 Mar 15];14(8):e1007161. Available from: https://dx.plos.org/10.1371/journal.ppat.1007161
746	16.	Simister NE. Human placental Fc receptors and the trapping of immune complexes. Vaccine. 1998;16(14–15):1451–5.
748	17.	Daëron M. FC RECEPTOR BIOLOGY. Annu Rev Immunol [Internet]. 1997 [cited 2022 Jun 2];15:203–37. Available from: www.annualreviews.org
750	18.	Atyeo C, DeRiso EA, Davis C, Bordt EA, De Guzman RM, Shook LL, et al. COVID-19 mRNA vaccines drive differential antibody Fc-functional profiles in pregnant, lactating, and nonpregnant women. Sci Transl Med [Internet]. 2021 Oct 27 [cited 2021 Dec 6];13(617):8631. Available from: https://www.science.org/doi/abs/10.1126/scitranslmed.abi8631
755	19.	Takizawa T, Anderson CL, Robinson JM. A Novel FcγR-Defined, IgG-Containing Organelle in Placental Endothelium. J Immunol [Internet]. 2005 Aug 15 [cited 2021 Apr 29];175(4):2331–9. Available from: http://www.jimmunol.org/content/175/4/2331
758	20.	Saji F, Samejima Y, Kamiura S, Koyama M. Dynamics of immunoglobulins at the feto–maternal interface. Reviews of Reproduction [Internet]. 1999 [cited 2021 Apr 20];81–9. Available from: http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.983.3422&rep=rep1&type=pdf
762	21.	Antohe F, Rădulescu L, Gafencu A, Ghe V, Simionescu M. Expression of functionally active FcRn and the differentiated bidirectional transport of IgG in human placental endothelial cells. Hum Immunol. 2001 Feb 1;62(2):93–105.
765	22.	Kiskova T, Mytsko Y, Schepelmann M, Helmer H, Fuchs R, Miedl H, et al. Expression of the neonatal Fc-receptor in placental-fetal endothelium and in cells of the placental immune system. Placenta. 2019 Mar 1;78:36–43.
768	23.	Niewiesk S. Maternal antibodies: Clinical significance, mechanism of interference with immune responses, and possible vaccination strategies. Front Immunol. 2014;5(SEP):446.
771	24.	CDC, Ncird. DEPARTMENT OF HEALTH AND HUMAN SERVICES CENTERS FOR DISEASE CONTROL AND PREVENTION Advisory Committee on Immunization Practices (ACIP) Summary Report. 2016 [cited 2022 Apr 6]; Available from: https://wwwdev.cdc.gov/vaccines/acip/meetings/downloads/min-archive/min-2016-10.pdf
775	25.	Healy CM, Rench MA, Swaim LS, Smith EOB, Sangi-Haghpeykar H, Mathis MH, et al. Association between Third-Trimester Tdap Immunization and Neonatal Pertussis Antibody Concentration. JAMA - J Am Med Assoc [Internet]. 2018 Oct 9 [cited 2021 Apr 9];320(14):1464–70. Available from: https://jamanetwork.com/

779   26.   Eberhardt CS, Blanchard-Rohner G, Lemaître B, Boukrid M, Combescure C, Othenin-
780   Girard V, et al. Maternal Immunization Earlier in Pregnancy Maximizes Antibody Transfer
781   and Expected Infant Seropositivity Against Pertussis. Clin Infect Dis [Internet]. 2016 Apr 1
782   [cited 2022 Apr 6];62(7):829–36. Available from:
783   https://academic.oup.com/cid/article/62/7/829/2463039
784   27.   Ishikawa T, Takizawa T, Iwaki J, Mishima T, Ui-Tei K, Takeshita T, et al. Fc gamma
785   receptor IIb participates in maternal IgG trafficking of human placental endothelial cells.
786   Int J Mol Med [Internet]. 2015 May 1 [cited 2021 Jul 22];35(5):1273–89. Available from:
787   http://www.spandidos-publications.com/10.3892/ijmm.2015.2141/abstract
788   28.   Joslyn LR, Kirschner DE, Linderman JJ. CaliPro: A Calibration Protocol That Utilizes
789   Parameter Density Estimation to Explore Parameter Space and Calibrate Complex
790   Biological Models. Cell Mol Bioeng. 2021 Feb 1;14(1):31–47.
791   29.   Malek A, Sager R, Kuhn P, Nicolaides KH, Schneider H. Evolution of maternofetal
792   transport of immunoglobulins during human pregnancy. Am J Reprod Immunol.
793   1996;36(5):248–55.
794   30.   Palmeira P, Quinello C, Silveira-Lessa AL, Zago CA, Carneiro-Sampaio M. IgG placental
795   transfer in healthy and pathological pregnancies. Vol. 2012, Clinical and Developmental
796   Immunology. 2012.
797   31.   Bruhns P, Jönsson F. Mouse and human FcR effector functions. Immunol Rev.
798   2015;268(1):25–51.
799   32.   Sarvas H, Sepp I, Kurikka S, Siegberg R, MJi O. Half-Life of the Maternal IgG1 Allotype
800   in Infants. J Clin Immunol. 1993;13(2).
801   33.   McKay MD, Beckman RJ, Conover WJ. A comparison of three methods for selecting
802   values of input variables in the analysis of output from a computer code. Technometrics.
803   2000;42(1):55–61.
804   34.   Surve S V., Myers PJ, Clayton SA, Watkins SC, Lazzara MJ, Sorkina A. Localization
805   dynamics of endogenous fluorescently labeled RAF1 in EGF-stimulated cells. Mol Biol
806   Cell [Internet]. 2019 Feb 15 [cited 2023 Apr 5];30(4):506–23. Available from:
807   https://www.molbiolcell.org/doi/10.1091/mbc.E18-08-0512
808   35.   Vento-Tormo R, Efremova M, Botting RA, Turco MY, Vento-Tormo M, Meyer KB, et al.
809   Single-cell reconstruction of the early maternal–fetal interface in humans. Nature
810   [Internet]. 2018 Nov 15 [cited 2021 Jan 15];563(7731):347–53. Available from:
811   https://pubmed.ncbi.nlm.nih.gov/30429548/
812   36.   Tsang JCH, Vong JSL, Ji L, Poon LCY, Jiang P, Lui KO, et al. Integrative single-cell and
813   cell-free plasma RNA transcriptomics elucidates placental cellular dynamics. Proc Natl
814   Acad Sci U S A [Internet]. 2017 Sep 12 [cited 2021 Mar 29];114(37):E7786–95. Available
815   from: www.pnas.org/cgi/doi/10.1073/pnas.1710470114
816   37.   Antohe F, Rădulescu L, Gafencu A, Ghe V, Simionescu M. Expression of functionally
817   active FcRn and the differentiated bidirectional transport of IgG in human placental
818   endothelial cells. Hum Immunol [Internet]. 2001 [cited 2022 Jun 7];62(2):93–105.
819   Available from: https://pubmed.ncbi.nlm.nih.gov/11182218/
820   38.   White MT, Griffin JT, Akpogheneta O, Conway DJ, Koram KA, Riley EM, et al. Dynamics
821   of the Antibody Response to Plasmodium falciparum Infection in African Children. J Infect
822   Dis [Internet]. 2014 Oct 1 [cited 2022 Sep 22];210(7):1115–22. Available from:
823   https://academic.oup.com/jid/article/210/7/1115/2908670
824   39.   van der Lee S, van Rooijen DM, de Zeeuw-Brouwer ML, Bogaard MJM, van Gageldonk
825   PGM, Marinovic AB, et al. Robust humoral and cellular immune responses to pertussis in
826   adults after a first acellular booster vaccination. Front Immunol. 2018 Apr 4;9(APR):681.
827   40.   Knuutila A, Versteegen P, Barkoff AM, van Gageldonk P, Mertsola J, Berbers G, et al.
828   Pertussis toxin neutralizing antibody response after an acellular booster vaccination in
829   Dutch and Finnish participants of different age groups. Emerg Microbes Infect [Internet].

2022 [cited 2023 Apr 12];11(1):956–63. Available from:
https://www.tandfonline.com/doi/abs/10.1080/22221751.2022.2053364

41. Hovingh ES, Kuipers B, Bonačić Marinović AA, Jan Hamstra H, Hijdra D, Mughini Gras L,
et al. Detection of opsonizing antibodies directed against a recently circulating Bordetella
pertussis strain in paired plasma samples from symptomatic and recovered pertussis
patients. Sci Reports 2018 81 [Internet]. 2018 Aug 13 [cited 2023 Apr 13];8(1):1–11.
Available from: https://www.nature.com/articles/s41598-018-30558-8

42. Dari A, Boulton M, Neyens M, Le Gars M, Valenzuela B, Shukarev G, et al. Quantifying
Antibody Persistence After a Single Dose of COVID-19 Vaccine Ad26.COV2.S in
Humans Using a Mechanistic Modeling and Simulation Approach. Clin Pharmacol Ther.
2022;

43. Vilajeliu A, Goncé A, López M, Costa J, Rocamora L, Ríos J, et al. Combined tetanus-
diphtheria and pertussis vaccine during pregnancy: transfer of maternal pertussis
antibodies to the newborn. Vaccine. 2015 Feb 18;33(8):1056–62.

44. Atyeo C, DeRiso EA, Davis C, Bordt EA, De Guzman RM, Shook LL, et al. COVID-19
mRNA vaccines drive differential antibody Fc-functional profiles in pregnant, lactating,
and nonpregnant women. Sci Transl Med [Internet]. 2021 Oct 27 [cited 2021 Dec
7];13(617):8631. Available from:
https://www.science.org/doi/abs/10.1126/scitranslmed.abi8631

45. Wood GW, King CR. Trapping antigen-antibody complexes within the human placenta.
Cell Immunol. 1982 May 15;69(2):347–62.

46. Vidarsson G, Dekkers G, Rispens T. IgG subclasses and allotypes: From structure to
effector functions. Front Immunol. 2014;5(OCT):520.

47. Alter G, Ottenhoff THM, Joosten SA. Antibody glycosylation in inflammation, disease and
vaccination. Semin Immunol. 2018 Oct 1;39:102–10.

48. Mahan AE, Jennewein MF, Suscovich T, Dionne K, Tedesco J, Chung AW, et al.
Antigen-Specific Antibody Glycosylation Is Regulated via Vaccination. PLoS Pathog.
2016 Mar 1;12(3).

49. Jefferis R, Kumararatne DS. Selective IgG subclass deficiency: quantification and clinical
relevance. Clin exp Immunol. 1990;81:357–67.

50. Einarsdottir HK, Selman MHJ, Kapur R, Scherjon S, Koeleman CAM, Deelder AM, et al.
Comparison of the Fc glycosylation of fetal and maternal immunoglobulin G. Glycoconj J
[Internet]. 2013 Feb 10 [cited 2021 Jan 15];30(2):147–57. Available from: www.ms-
utils.org/Xtractor2D.

51. Wang Y, Jiang X, He J, Diraviyam T, Zhang X. Quantitative Investigation on Correlation
Between IgG and FcRn During Gestation and Lactating Periods in Rat. Am J Reprod
Immunol. 2016;75(2):81–5.

52. Nicolaides KH, Clewell WH, Rodeck CH. Measurement of human fetoplacental blood
volume in erythroblastosis fetalis. Am J Obstet Gynecol [Internet]. 1987 [cited 2021 Dec
1];157(1):50–3. Available from: https://pubmed.ncbi.nlm.nih.gov/3605267/

53. Abdalla AM, Tingari MD, Abdalla MA. Histomorphometric parameters of normal full term
placenta of Sudanese women. Heliyon [Internet]. 2016 Jul 1 [cited 2021 Mar 23];2(7):135.
Available from: /pmc/articles/PMC4971126/

54. Trygg J, Wold S. Orthogonal projections to latent structures (O-PLS). J Chemom.
2002;16(3):119–28.

55. Salahudeen MS, Nishtala PS. An overview of pharmacodynamic modelling, ligand-
binding approach and its application in clinical practice. Vol. 25, Saudi Pharmaceutical
Journal. Elsevier B.V.; 2017. p. 165–75.

56. Wilmore JR, Allman D. Here, There, and Anywhere? Arguments for and against the
Physical Plasma Cell Survival Niche. J Immunol [Internet]. 2017 Aug 1 [cited 2023 Apr
12];199(3):839–45. Available from:

https://journals.aai.org/jimmunol/article/199/3/839/106408/Here-There-and-Anywhere-
Arguments-for-and-against

57. Cyster JG, Allen CDC. B Cell Responses: Cell Interaction Dynamics and Decisions. Cell.
2019 Apr 18;177(3):524–40.

58. Huygen K, Caboré RN, Maertens K, Van Damme P, Leuridan E. Humoral and cell
mediated immune responses to a pertussis containing vaccine in pregnant and
nonpregnant women. Vaccine. 2015 Aug 7;33(33):4117–23.

59. Tsang JCH, Vong JSL, Ji L, Poon LCY, Jiang P, Lui KO, et al. Integrative single-cell and
cell-free plasma RNA transcriptomics elucidates placental cellular dynamics. Proc Natl
Acad Sci U S A [Internet]. 2017 Sep 12 [cited 2022 Jun 30];114(37):E7786–95. Available
from: www.pnas.org/cgi/doi/10.1073/pnas.1710470114