Quantitative mechanistic model reveals key determinants of placental IgG transfer and informs prenatal immunization strategies

Remziye R Erdogan[1], Sepideh Dolatshahi*[1,2]

[1]Department of Biomedical Engineering, University of Virginia School of Medicine, Charlottesville, VA, 22908

[2]Carter Immunology Center, University of Virginia School of Medicine, Charlottesville, VA, 22908

*Corresponding author: E-mail: sd8us@virginia.edu

Abstract

Transplacental antibody transfer is crucially important in shaping neonatal immunity. Recently, prenatal maternal immunization has been employed to boost pathogen-specific immunoglobulin G (IgG) transfer to the fetus. Multiple factors have been implicated in antibody transfer, but how these key dynamic regulators work together to elicit the observed selectivity is pertinent to engineering vaccines for mothers to optimally immunize their newborns. Here, we present the first quantitative mechanistic model to uncover the determinants of placental antibody transfer and inform personalized immunization approaches. We identified placental FcγRIIb expressed by endothelial cells as a limiting factor in receptor-mediated transfer, which plays a key role in promoting preferential transport of subclasses IgG1, IgG3, and IgG4, but not IgG2. Integrated computational modeling and *in vitro* experiments reveal that IgG subclass abundance, Fc receptor (FcR) binding affinity, and FcR abundance in syncytiotrophoblasts and endothelial cells contribute to inter-subclass competition and potentially inter- and intra-patient antibody transfer heterogeneity. We developed an *in silico* prenatal vaccine testbed by combining a computational model of maternal vaccination with this placental transfer model using the tetanus, diphtheria, and acellular pertussis (Tdap) vaccine as a case study. Model simulations unveiled precision prenatal immunization opportunities that account for a patient's anticipated gestational length, placental size, and FcR expression by modulating vaccine timing, dosage, and adjuvant. This computational approach provides new perspectives on the dynamics of maternal-fetal antibody transfer in humans and potential avenues to optimize prenatal vaccinations that promote neonatal immunity.

Introduction

Neonates are vulnerable to infections due to their tolerogenic immune phenotype; for the same reason, neonatal vaccinations have been met with limited success to date (1). To provide passive immunity while the neonatal immune system adapts to the environment *ex utero*, maternal immunoglobulin G (IgG) is selectively transferred across the placenta during gestation. Not only do these maternal antibodies confer neonates with passive early life immunity against infectious diseases, but also they shape the trajectory of neonatal immune development by priming the cellular immune response (2–4). Despite this crucial role, the molecular underpinnings of IgG transplacental transport are incompletely described to date.

Though the placenta obstructs maternal-fetal transport of most large molecules, maternal IgG is selectively transported by way of Fc receptor (FcR)-mediated transcytosis. To enter fetal circulation, IgG must undergo transcytosis through two key cellular layers of the placenta: syncytiotrophoblasts (STBs) and endothelial cells (ECs). The neonatal Fc receptor (FcRn) is known to mediate IgG transport through STBs, but whether this mechanism is sufficient for IgG transport through ECs remains unclear (5–7). In addition to FcRn, STBs and ECs express several Fc gamma receptor (FcγR) isoforms with heterogeneous affinity for IgG (8–10). These classical FcγRs may contribute to observed selective placental "sieving" of IgG dependent upon fragment crystallizable (Fc) characteristics, e.g., IgG subclass and N-linked glycosylation (11–14), yet there is conflicting evidence surrounding the involvement of FcγRs in IgG transport (9,10,15,16). Recent studies hypothesized a role for STB-bound FcγRIIIa or EC-bound FcγRIIb in IgG transfer (9,12,17–21), yet others showed FcRn is sufficient for IgG transport in isolated placental ECs *in vitro* and in mouse placenta *in vivo* (14,22,23). Though informative, the experimental methods underlying these findings do not capture the dynamic nature or time scale of placental antibody transport. To disentangle the involvement of non-canonical FcγRs in IgG sieving, innovative methods that recapitulate the longitudinal dynamics of this fundamental process are needed.

Placental antibody transfer can be leveraged by maternal prenatal vaccines which boost

pathogen-specific IgG transport to the neonate (24). Identifying improved immunization strategies to

maximize neonatal antibody titers is non-trivial. Placental growth and development dynamically

regulating IgG transport coupled with known maternal immune adaptations during pregnancy

together define a unique immunization design space which poses a challenge to empirical vaccine

optimization by clinical trials alone (25). Predictive kinetic-dynamic modeling can be employed to

rationally design vaccines that maximize IgG transfer to the neonate. The ability to make *in silico*

predictions of vaccine-induced antibody transfer would enable pre-clinical dosing strategy studies,

expediting translation of novel therapeutics from bench to bedside.

In the United States, expecting mothers are routinely vaccinated against tetanus, diphtheria,

and acellular pertussis (Tdap) during the early third trimester (26), but emerging evidence suggests

that this recommendation may not optimally protect the entire population. First, it has been shown

that maternal Tdap immunization earlier in gestation results in higher pertussis toxin (PT)- and

filamentous hemagglutinin (FHA)-specific IgG in infants regardless of gestational length (26–28).

Second, current vaccination efforts are designed to elicit high antibody transfer to term neonates,

yet third trimester immunization may not allot sufficient time for the mother to mount a humoral

immune response and subsequently transfer antibodies to preterm neonates (28). Collectively, this

evidence supports the potential for personalized vaccine approaches that account for factors such

as maternal baseline IgG titer, immunization history, placental function, and risk of preterm delivery

to maximize pathogen-specific IgG transfer to the neonate, especially among premature neonates.

To uncover the molecular regulators of placental antibody transport and to inform the

development of personalized immunization approaches, we developed the first computational

model of human placental IgG transfer. In a case study on Tdap immunization, we use this model

as an *in silico* testbed for prenatal vaccine design and identify potential strategies to improve

transfer of vaccine-induced antibodies, both at a patient-specific and population level. Ultimately,

this model-driven investigation sheds light on the dynamic regulation of maternal-fetal IgG transfer

and provides a foundation to develop precision vaccine approaches which promote neonatal

immunity.

Results

Mechanistic model recapitulates IgG subclass-specific placental transfer

To elucidate mechanisms of IgG transfer and selective "sieving", we devised a dynamic model of

IgG transplacental transfer. The model consists of ordinary differential equations (ODEs) describing

IgG mass transport through the distinct layers comprising the maternal-fetal interface: STB FcRn-

mediated transcytosis, diffusion through intervillous stroma, and EC FcRn- and FcγRIIb-mediated

transcytosis (Fig 1A) (see Appendix S1 for model equations). FcRn is widely implicated in placental

IgG transport; we additionally chose to incorporate FcγRIIb into the model due to its expression in

term placental ECs and demonstrated role in EC transcytosis *in vitro* (9,19,29). To account for

increasing Fc receptor expression across gestation as observed in human and rat placenta, the

total bound and unbound Fc receptors were input to the model as time-varying offline variables

(30,31) (Fig S1, Appendix S1). Model parameters were derived from literature when available, and

11 unknown parameters were estimated by fitting predictions of fetal IgG to data obtained from

cordocentesis in human pregnancies (Fig 1B, Table 1) (32). Parameter estimation was performed

using CaliPro as described elsewhere (32,33) (see Methods). With this optimized parameter set,

the model recapitulated dynamics of both bulk and subclass-specific IgG transfer, predicting the IgG

subclass transfer hierarchy most frequently reported in literature (IgG1 > IgG3 > IgG4 > IgG2) (Fig

1C, S2 Fig) (11–13,32,34–36).

Fig 1. Mechanistic model captures IgG subclass-specific transfer dynamics. (A) The schematic depicts

the ODE model representing transport of IgG through the placenta, mapped onto a graphical representation

of the maternal-fetal interface. Reaction rate constants associated with each process are depicted on the

corresponding arrow. Each box represents a molar concentration in each compartment. "C" denotes

complexes of bound IgG and Fc receptors either in endosomes (FcRn) or at the cell surface (FcγRIIb).

$\Delta FcRn_{STB}^{total}$, $\Delta Fc\gamma RIIb_{EC}^{total}$, and $\Delta FcRn_{STB}^{total}$ represent synthesis of new FcRn and FcγRIIb molecules over time

as the placenta grows. For simplicity, the schematic represents bulk IgG transport, whereas the model

includes separate equations for each IgG subclass. (B) Simulated fetal IgG subclass levels vs. time (lines) are

overlaid with the data used for parameter estimation. The overlaid data are the mean and standard deviation

of IgG subclasses in the umbilical cord between 17-41 weeks gestation (n = 107) measured using

cordocentesis by Malek et al (32). Gray lines represent 1000 perturbation simulations resulting from sampling

parameters from the ranges determined by CaliPro. (C) The swarm charts show the fetal/maternal ratio (F:M

ratio) for each IgG subclass at 40 weeks gestation resulting from 1000 simulations sampling from the

optimized range of parameters (filled symbols) compared to the mean cord/maternal ratio for each IgG

subclass compiled across cohort studies mined from the literature (hollow symbols). Each symbol represents

a single simulation or cohort average. Sample means are indicated by black lines. The symbols outlined in

black represent the mean of IgG subclass cord/maternal transfer ratios at parturition from the human study

used during parameter optimization (32). Dashed line at F:M ratio = 1 indicates equal concentration in

maternal and fetal blood at parturition.

116 **Table 1. List of optimized model parameters.**

Parameter	Value	Units	Interpretation	Source
k_{up}	0.0872-0.0982	L/week	Uptake rate into cells	Optimized
k_{trans}	0.0569-0.0743	L/week	Transcytosis rate	Optimized
k_{deg}	8.4733-9.0809	L/week	Lysosomal degradation rate	Optimized
$FcRn_{STB}^{total,end}$	3.33-4.94e-5	M	STB FcRn expression at term	Optimized
$Fc\gamma RIIb_{EC}^{total,end}$	2.73-3.33e-5	M	EC FcγRIIb expression at term	Optimized
$FcRn_{EC}^{total,end}$	2.36-2.86e-6	M	EC FcRn expression at term	Optimized
$IgG1_0$	3.78e-5	M	Maternal IgG1 concentration	(32)
$IgG2_0$	1.81e-5	M	Maternal IgG2 concentration	(32)
$IgG3_0$	2.35e-6	M	Maternal IgG3 concentration	(32)
$IgG4_0$	2.27e-6	M	Maternal IgG4 concentration	(32)
K_D, FcRn-IgG1 (k_{off}/k_{on})	1.25e-8 (0.15/1.2e7)	M	FcRn and IgG1 dissociation constant	(37,38)
K_D, FcRn-IgG2 (k_{off}/k_{on})	2e-8 (0.15/7.5e6)	M	FcRn and IgG2 dissociation constant	(37,38)
K_D, FcRn-IgG3 (k_{off}/k_{on})	3.3e-8 (0.15/4.5e6)	M	FcRn and IgG3 dissociation constant	(37,38)
K_D, FcRn-IgG4 (k_{off}/k_{on})	5e-8 (0.15/3e6)	M	FcRn and IgG4 dissociation constant	(37,38)
K_D, FcγRIIb-IgG1 (k_{off}/k_{on})	1e-5 (0.15/1.5e4)	M	FcγRIIb and IgG1 dissociation constant	(37,38)
K_D, FcγRIIb-IgG2 (k_{off}/k_{on})	5e-5 (0.15/3e3)	M	FcγRIIb and IgG2 dissociation constant	(37,38)
K_D, FcγRIIb-IgG3 (k_{off}/k_{on})	5e-6 (0.15/3e4)	M	FcγRIIb and IgG3 dissociation constant	(37,38)

K_D, FcγRIIb-IgG4 (k_{off}/k_{on})	5e-6 (0.15/3e4)	M	FcγRIIb and IgG4 dissociation constant	(37,38)
V_M	5.552-5.619	L	Maternal blood volume	Optimized
V_{STB}	0.224-0.258	L	Total STB endosomal volume	Optimized
V_{STR}	0.165-0.179	L	Stromal volume	Optimized
V_{EC}	0.247-0.275	L	Total EC endosomal volume	Optimized
V_F	0.254-0.271	L	Fetal blood volume	Optimized
δ_{Ab}	0.02	L/week	IgG decay rate in the fetus	(39)

Transcytosis rate parameters and Fc receptor expression regulate IgG transfer efficiency and selectivity

To determine the contribution of each parameter to the subclass-specific dynamics predicted by the

model, we performed global sensitivity analyses. We generated randomly perturbed parameter sets

using Latin Hypercube Sampling (LHS) from uniform distributions spanning the maximum likelihood

parameter ranges determined by CaliPro (33,40,41). For each parameter set, the predicted fetal

IgG concentration at 40 weeks was calculated. To determine the parameters with the greatest

influence on total IgG transfer (Fig 2A-B) and on differential transfer between IgG subclasses

defined as entropy of subclasses (Fig 2C-D), orthogonalized partial least squares regression

(OPLSR) models were developed using the 1000 perturbed parameter sets to predict either

outcome (Methods) (42). Variable importance in projection (VIP) scores revealed that total IgG

transfer was most sensitive to the rate of IgG transcytosis by STBs and ECs (k_{trans}), FcRn

expressed by STBs, and FcγRIIb expressed by ECs (Fig 2B). On the other hand, all volume

parameters negatively correlated with IgG transfer, likely because increasing the volume of either

compartment dilutes the IgG concentration. To reveal parameters that contribute to the difference in

IgG transfer among IgG subclasses, we built a second OPLSR model predicting the entropy among

IgG subclasses (Methods) (Fig 2C-D). Interestingly, VIP score analysis showed FcγRIIb expressed

by ECs and k_{trans} contribute most to the distinct transfer rates among IgG subclasses. FcγRIIb is a

low-affinity IgG receptor with highest affinity for subclasses IgG3 and IgG4, suggesting that it

constitutes a rate-limiting factor of IgG transfer which fine-tunes the subclass transfer hierarchy

(11,15,29). Likewise, k_{trans} acts in an FcR-dependent manner to further amplify the effects of Fc-

mediated selectivity. Thus, the rate of cargo internalization and fetoplacental size drives the

concentration of bulk IgG in the fetus, whereas FcRn and FcγRIIb collaborate to fine-tune patterns

of IgG subclass-specific transfer.

Fig 2. Key parameters associated with transcytosis mechanism and Fc receptor expression regulate

IgG transfer. (A-B) An OPLSR model was built to determine parameters predictive of total fetal IgG. (A) The

scores plot shows X scores on latent variables 1 and 2 (LV1 and LV2) of simulated fetal IgG concentration at

parturition for 1000 randomly perturbed parameter sets. Each circle represents a single perturbation

experiment. The points are colored proportional to the resulting simulated concentrations of IgG in the fetus at

delivery. The model was orthogonalized such that the direction of maximum variance is in the direction of

latent variable 1 (LV1), capturing 99% of the Y variation. The model performed with high predictability power

(Q^2 = 99%) and a mean squared error which outperformed 1000 random models with permutated labels (p <

0.001). (B) The bar graph depicts the variable importance in projection (VIP) scores, which shows the relative

contribution of each parameter to total fetal IgG for a term delivery. VIP scores are artificially oriented in the

direction of their loadings on LV1. VIP score > 1 indicates greater-than-average contribution of that

parameter. (C-D) An OPLSR model was built to predict entropy among fetal IgG subclass levels at the end of

gestation. (C) The scores plot showing X scores on LV1 and LV2 of simulated fetal IgG concentration at

parturition for 1000 perturbation parameter sets. Each circle represents a single perturbation experiment. The

points are colored proportional to the resulting simulated concentrations of IgG in the fetus at delivery. The

model was orthogonalized such that LV1 captures features associated with the highest entropy, accounting

for 96% of the Y variation. The model performed with high predictability power (Q^2 = 96%) and a mean

squared error which outperformed 1000 models based on randomly permutated labels (p < 0.001). (D) The

VIP scores bar plot shows the relative contribution of each parameter to the entropy of IgG subclasses in fetal

blood for a term delivery. VIP scores are artificially oriented in the direction of their loadings on LV1.

Endothelial cell-bound FcγRIIb is a key driver of subclass-specific IgG transfer

It is well-established that placental IgG transport increases during the third trimester in humans, so

we asked whether increasing FcγRIIb or FcRn expression in ECs drives increased transport

efficiency. To gain insight into the expression dynamics of the genes coding for FcγRIIb and FcRn,

namely *FCGR2B* and *FCGRT* (FcRn heavy chain), in fetal ECs across gestation, we analyzed two

single cell RNA sequencing (scRNA-seq) datasets from published literature spanning three

gestational time points: (1) first trimester samples from elective abortions (< 16 weeks), (2) early

third trimester samples from preeclamptic cesarean sections (30 weeks), and (3) late third trimester

samples from uncomplicated cesarean sections (>37 weeks weeks) (43,44). Interestingly, *FCGR2B*

expression was significantly higher in late compared to early third trimester ECs, but we did not

detect a significant difference in *FCGRT* expression between early and late third trimester ECs (Fig

3A-B). Conversely, *FCGRT* but not *FCGR2B* transcripts were detected in first trimester ECs, and

the ratio of *FCGR2B* to *FCGRT* expression per cell increased across time points (Fig 3C, S3 Fig),

suggesting that FcγRIIb dynamically regulates IgG transcytosis in ECs.

Fig 3. Endothelial cell-bound FcγRIIb is a key driver of subclass-specific IgG transfer. (A-B) Violin plots show *FCGR2B* (A) and *FCGRT* (B) transcript levels in preterm, preeclamptic ECs (30 weeks) compared to term, healthy ECs (>37 weeks) from single cell RNA sequencing data. Black circles represent the sample median, white circles represent the sample mean. (C) Violin plots show the log-transformed ratio of *FCGR2B* to *FCGRT* counts in ECs from single cell RNA seq data from two separate cohorts spanning 3 gestational time points: <16 weeks, 30 weeks, and >37 weeks. Black circles represent the sample median, white circles represent the sample mean. (D-F) Model-simulated IgG subclass transfer ratios as a function of EC Fc receptor expression with ECs expressing FcRn and FcγRIIb (D, EC$_{FcγRIIb-FcRn}$), only FcRn (E, EC$_{FcRn}$), or only FcγRIIb (F, EC$_{FcγRIIb}$). The corresponding schematics depict the simulation conditions. Statistical significance in (A-C) was determined by a Wilcoxon rank sum test (* $p < 0.01$, ** $p < 0.001$, n.s. not significant).

Previous studies showed that FcRn and FcγRIIb are co-expressed by placental ECs at the protein level (9,45). Since FcRn has a greater affinity for all IgG subclasses (Table 1), it is unclear whether FcRn alone is responsible for IgG transfer across fetal ECs (22). To assess the contribution of FcγRIIb to IgG transport, we tested whether FcγRIIb is a determinant of subclass-specific transfer. In the model with FcRn and FcγRIIb co-expressed (EC$_{FcRn,FcγRIIb}$), we found that the predicted subclass transfer hierarchy was IgG1 > IgG3 > IgG4 > IgG2 when FcRn expression was less than 25% of FcγRIIb (Fig 3D), demonstrating that higher relative expression of FcγRIIb to FcRn by ECs drives subclass-specific IgG placental transfer. As FcγRIIb binds subclasses IgG1, IgG3, and IgG4 with greater affinity than IgG2, we hypothesized that FcγRIIb is necessary to predict

preferential transfer of IgG1, IgG3, and IgG4 over IgG2. To test this hypothesis, we compared

model-predicted fetal-maternal IgG ratios from simulations assuming ECs solely express FcRn

(EC_{FcRn}) or FcγRIIb ($EC_{FcγRIIb}$). The EC_{FcRn} model did not capture the preferential transfer of IgG1,

IgG3, and IgG4 at any FcRn expression level (Fig 3E). Interestingly, in the $EC_{FcγRIIb}$ model the

subclass transfer hierarchy shifted from IgG3 > IgG4 > IgG1 > IgG2 to IgG1 > IgG2 > IgG3 > IgG4

with increasing FcγRIIb expression, indicating FcγRIIb drives the observed subclass transfer

hierarchy in humans (Fig 3F). Together, these data implicate FcγRIIb in ECs as a key regulator of

antibody transfer dynamics across gestation.

IgG subclasses compete for receptor-mediated transcytosis

Our finding that FcγRIIb limitation plays a key role in determining the subclass transfer hierarchy

suggests that maternal IgG subclasses compete for receptor-mediated transcytosis. On the other

hand, variation in IgG subclass abundances across individuals may contribute to patient-specific

differences in IgG subclass transfer. In agreement with this hypothesis, the placental transport

model predicted that all four IgG subclasses transferred less efficiently when mixed compared to

each subclass in isolation of the others and the effect of mixing was greater for subclasses with low

abundances in maternal serum at low FcγRIIb expression levels (S4 Fig). To further elucidate the

role of inter-subclass competition in determining transport efficiency, we explored the effect of IgG1

on IgG4 transcytosis in ECs in a separate model of transcytosis across ECs only (see Methods).

We chose to model IgG1 and IgG4 as a case study because they have the highest and lowest

abundance in maternal serum, respectively, yet both generally exhibit a cord/maternal ratio > 1

(11,32). Based on a simple kinetic reaction model, the inter-subclass competition can be described

by the following closed-form equation:

$$C_{IgG4-FcRn} = \frac{FcRn_0 [IgG4]}{K_{D,IgG4}(1 + \frac{[IgG4]}{K_{D,IgG4}} + \frac{[IgG1]}{K_{D,IgG1}})}$$

Where $C_{IgG4-FcRn}$ is the concentration of bound IgG4 and FcRn, $FcRn_0$ is the total concentration of

receptor, [IgG1] and [IgG4] are concentrations of maternal IgG1 and IgG4, and $K_{D,IgG1}$ and $K_{D,IgG4}$

are their respective equilibrium dissociation constants (see derivation in Methods). Intuitively, C_{IgG4-}

$_{FcRn}$ is a function of IgG1 abundance and affinity for FcRn, such that either increasing [IgG1] or

decreasing $K_{D,IgG1}$ (i.e., increasing IgG1-FcRn affinity) decreases $C_{IgG4-FcRn}$ and consequently, IgG4

transcytosis.

To validate and further characterize competition among IgG subclasses, we used an *in vitro*

system to model IgG transcytosis through human umbilical vein endothelial cells (HUVEC) cultured

on Transwell permeable membranes (Fig 4A) (Methods). In parallel, we devised a distinct,

simplified mechanistic model of receptor-mediated transcytosis in ECs and optimized its parameters

by fitting to temporal data of IgG4 transcytosis in the HUVEC experimental system ([IgG4$_{apical}$] = 0.5

or 0.125 mg/ml) (S5 Fig, Table 2). To recapitulate the physiological conditions where FcR is limited,

we varied IgG4 levels to determine the concentration of IgG4 where IgG4 transfer no longer

increases with increasing apical IgG4 levels. Although HUVEC expresses FcRn but not FcγRIIb,

our findings remain valid because the affinities of IgG1 and IgG4 for either receptor are on the same

order of magnitude and thus the dynamics of subclass competition are comparable (37). Our

mechanistic model predicted that IgG4 transcytosis saturates at [IgG4$_{apical}$] > 0.33 mg/ml, indicating

FcRn limitation; we validated this finding in an analogous *in vitro* experiment in the Transwell

system (Fig 4B). To quantify the effect of IgG1 on IgG4 transcytosis, we simulated the effect of

increasing IgG1 concentration when IgG4 was fixed at a concentration just below the FcRn

saturation point ([IgG4$_{apical}$] = 0.2 mg/ml). Our model predicted that IgG4 transcytosis would not be

affected by the presence of IgG1 until the total IgG concentration exceeded the saturation point—

that is, [IgG1$_{apical}$] + [IgG4$_{apical}$] > 0.33 mg/ml. In agreement with our computational model prediction,

increasing IgG1 caused IgG4 transcytosis to decrease only when the total IgG concentration

exceeded the FcRn saturation point (Fig 4C). IgG subclass-specific transfer is thus a function of

subclass abundances and FcR affinity, and competition for FcR-mediated transcytosis occurs when

FcR expression is limited.

Fig 4. IgG1 competes with IgG4 for FcRn-mediated transcytosis in HUVECs. (A) Schematic showing a

cross-sectional view of the Transwell system and corresponding mechanistic model representation of FcRn-

mediated IgG4 and IgG1 transport through HUVEC. Left panel: Arrow indicates the direction of transfer

(apical to basolateral). Center panel and right panel: Reaction rate constant associated with each process is

depicted on the corresponding arrow. (B) Model-predicted IgG4 transcytosis (y-axis) with increasing apical

concentration (x-axis) (solid line) is plotted against experimental data points with the mean and standard

deviation from three independent biological replicates. The corresponding experimental conditions are

displayed in the panel on the left. The dashed line represents the model-predicted and experimentally

validated transcytosis saturation point ($[\text{IgG}_{\text{total}}] = [\text{IgG4}] = 0.33$ mg/ml). (C) Model-predicted IgG4 and IgG1

transcytosis (y-axis) with increasing apical concentration of IgG1 (x-axis) (solid lines) is plotted against

experimental data points with the mean and standard deviation from three independent biological replicates.

The corresponding experimental conditions are displayed in the panel on the left. The dashed line represents

the model-predicted and experimentally validated transcytosis saturation point ($[\text{IgG}_{\text{total}}] = [\text{IgG4}] + [\text{IgG1}] =$

0.33 mg/ml. Statistical comparisons shown are between IgG4 in the absence of IgG1 (far left condition) and

all combinations of IgG1 and IgG4 mixed using a one-tailed two-sample t-test (* $p < 0.05$, ** $p < 0.001$, n.s.

not significant).

Table 2. Optimized HUVEC transcytosis model parameters.

Parameter	Value	Units	Interpretation	Source
FcRn	2200	nM	HUVEC FcRn expression level	Optimized
k_{deg}^{HUVEC}	0.01	1/minute	Lysosomal degradation rate	Optimized
k_{trans}^{HUVEC}	2.5×10^{-5}	1/minute	Transcytosis rate	Optimized
K_D, FcRn-IgG1 (k_{off}/k_{on})	1.25e-8 (0.15/1.2e7)	M	FcRn and IgG1 dissociation constant	(37,38)
K_D, FcRn-IgG4 (k_{off}/k_{on})	5e-8 (0.15/3e6)	M	FcRn and IgG4 dissociation constant	(37,38)

269 Optimal Tdap vaccine strategies depend on gestational age,
270 kinetics of the maternal IgG response to Tdap, and Fc receptor
271 expression

Considering the new insights into the dynamic regulation of antibody transfer by FcRs revealed by

our model, we sought to uncover key features influencing the transfer of vaccine-induced antibodies

which could be considered in precision immunization approaches. To this end, we developed an *in*

silico immunization testbed by layering simulated kinetics of the maternal antibody response post-

immunization into the placental antibody transfer model (Fig 5A). As a case study of a maternal

vaccine currently in use, we adapted an existing model of plasma cell activation and antibody

secretion following malaria infection to recapitulate IgG responses to Tdap immunization in women

of child-bearing age (Methods) (Fig 5B, Table 3) (46,47). A dose of antigen representing maternal

immunization was applied to the system at three representative gestational ages (t_{vax}) of 10, 20, and

30 weeks to stimulate antibody secreting cell proliferation and anti-pertussis toxin IgG (α-PT IgG)

secretion and transfer to the fetus (Fig 5C). Interestingly, the interplay of maternal IgG response

kinetics and placental transfer results in different IgG transfer dynamics depending on t_{vax}. This non-

intuitive, gestational-age dependent effect on transfer dynamics emphasizes the importance of *in*

silico predictions to determine optimal immunization schedules.

Fig 5. Optimal Tdap immunization time is gestational age dependent. (A) Schematic shows mechanistic

model of maternal plasma cell antibody secretion post-vaccination. Vaccine-induced maternal IgG kinetics are

integrated into the maternal compartment of the placental IgG transport model, shown in the righthand box.

Dashed arrows indicate regulatory processes, solid lines represent processes modeled using mass action

kinetics. L_{ASC}: Long-lived antibody secreting cell, S_{ASC}: short-lived antibody-secreting cell, Ag: antigen. (B)

Simulated maternal anti-pertussis IgG concentrations vs. time is plotted with a solid line and overlaid with

measured IgG concentrations following Tdap vaccination in women of childbearing age from published

literature (47) (n = 105). The 95% confidence intervals are shown with light gray shading. (C) Simulated fetal

anti-pertussis toxin IgG concentrations are plotted against time following maternal vaccination at three sample

vaccination times (indicated along the x-axis): t_{vax} = 10, 20, and 30 weeks gestational age. Shaded regions

correspond to gestational length groups (left to right): extremely preterm, very preterm, late preterm, term. (D)

Simulated anti-pertussis toxin IgG in the fetus at the time of delivery (y-axis) is plotted over a range of

simulated vaccination times (x-axis) and gestational ages (color intensity, color bar on right). The optimal

vaccination time and corresponding fetal IgG levels are marked with a circle. The current CDC-recommended

maternal vaccination window is highlighted in red, and the model-predicted optimal vaccination range

encompassing all gestational age groups is highlighted in green.

Table 3. Pertussis immunization model parameters.

Parameter	Value	Units	Interpretation	Source
k_{ASC}	0.6	$1 / 10^6$ PBMCs / day / Ag	Rate of plasma cell generation	Optimized
k_{IgG}	0.0475	mg / ml * (10^6 PBMCs) / days	Rate of antibody production by plasma cells	Optimized
Ag_0	100	mg/ml	Initial dose of antigen	Optimized
ρ	96	%	Proportion of short-lived ASCs (S_{ASC})	(46)
δ_{Ag}	0.15	1 / day	Antigen decay rate	(50)
δ_{IgG}	0.033	1 / day	Antibody decay rate	(46)
δ_{S-ASC}	0.173	1 / day	Short-lived antibody-secreting cell (S_{ASC}) decay rate	(46)
δ_{L-ASC}	6.6×10^{-4}	1 / day	Long-lived antibody secreting cell (L_{ASC}) decay rate	(46)

We next performed simulations over a range of possible immunization times to determine

the optimal t_{vax} for newborns born at different weeks of gestation (Fig 5D). Though the correlate of

protection for pertussis infection still remains elusive, some possible roles of IgG in pertussis

infection include neutralization and opsonophagocytosis (48,49). For the purposes of this study, we

assumed that neutralizing antibodies confer protection against pertussis toxin in the neonate and

therefore the optimal t_{vax} is defined as the gestational age at immunization resulting in the highest α-

PT IgG titer in the fetus at birth. Empirical studies have previously shown that earlier vaccination

may be an effective strategy to protect preterm neonates who typically receive a lower quantity of

maternal antibodies (26–28). Using our immunization testbed, we first asked how the optimal Tdap

administration schedule varies for the average patient delivering at different gestational time points

(i.e., term, late preterm, very preterm, or extremely preterm). In the case of term gestation, the

model-predicted optimal t_{vax} was during the 25th week of gestation, but Tdap immunization as early

as the 10th week still resulted in greater than half of the optimal IgG concentration being transferred

to term newborns (Fig 5D). Optimal t_{vax} scaled with increasing gestational length, suggesting that

risk for preterm birth should be considered when determining prenatal immunization schedules (Fig

5D). Interestingly, we found the optimal t_{vax} fell during the second trimester (between 14-25 weeks)

regardless of preterm birth status, which is earlier than the window recommended by the Center for

Disease Control and Prevention (CDC) (28-34 weeks) and in agreement with studies reporting

higher titer of pertussis antibodies in the fetus following maternal vaccination during second

trimester as opposed to third trimester (26,28,51). This optimized Tdap vaccination window was

unaffected by the magnitude of the maternal α-PT IgG response but did depend on the existence of

a transient spike in maternal α-PT IgG driven by the percentage of S_{ASC} (ρ) (Fig S6-7). While α-PT

IgG responses following Tdap are generally characterized by a short-lived spike in IgG, variation in

the IgG response kinetics for other target antigens should be considered when applying this model

to other vaccines. Thus, the optimal Tdap immunization schedule is gestational length-dependent,

but a revised vaccination window earlier in gestation could be an effective population-level strategy

to ensure coverage for preterm and term newborns alike.

As other sources of inter-patient variability besides gestational length may influence the

effectiveness of prenatal vaccination, we sought to identify further opportunities for patient-specific

immunization approaches. To determine key variables that should be considered in immunization

schedules, we performed a global sensitivity analysis to uncover parameters with the greatest

influence on optimal t_{vax}. We optimized t_{vax} in simulations from 1000 perturbed parameter sets, then

built an OPLSR model to predict the optimal t_{vax} given each parameter set (Fig 6A, 6B, Methods).

Parameters strongly associated with earlier optimal immunization times included $FcRn_{STB}^{total,end}$ and

 v_F; $FcRn_{EC}^{total,end}$ and $Fc\gamma RIIb_{EC}^{total,end}$ similarly had negative loadings on latent variable 1 (LV1), as

did several other volume parameters. Intuitively, higher compartmental volumes decrease transfer

of vaccine-induced antibodies because IgG concentration is diluted in high volume compartments,

and thus earlier vaccination enables more IgG to accumulate in the fetus to compensate for this

dilution. Likewise, higher $FcRn_{STB}^{total,end}$ facilitates IgG transport into the placental stroma making IgG

subsequently available for trans-EC transport. If sufficient $FcRn_{STB}^{total,end}$ is expressed, earlier

immunization corresponds to increased levels of IgG accumulated in the stroma, readily available

for trans-EC transport. On the other hand, k_{trans} corresponds to later optimal immunization timing

because if transcytosis efficiency increases, it confers greater benefit to timing the maternal IgG

spike post-vaccination when FcRn and FcγRIIb are expressed in higher quantities. Therefore,

differences in fetoplacental size and function across the population may play a role in

determining the ideal patient-specific immunization program.

Fig 6. Model reveals opportunities for personalized vaccination strategies. (A-B) The OPLSR model

predicts the optimal immunization time (t_{vax}, weeks gestational age) from perturbed parameter sets. (A) The

scores plot depicts the X scores on LV1 and LV2 of the optimal t_{vax} for 1000 perturbed parameter sets. Each

circle represents a single perturbation experiment colored proportionally to the predicted optimal t_{vax}. The

model was orthogonalized such that LV1 is in the direction of maximum variance. (B) The bar plot shows VIP

scores that quantify the relative contribution of each parameter to the optimal t_{vax}. VIP scores are artificially

oriented in the direction of their loadings on LV1. The model performed with predictive power (Q^2 = 47%) and

with a mean squared error lower than 1000 models based on randomly permutated labels (p < 0.001). (C)

The value of $FcRn_{STB}^{total,end}$ (left), $Fc\gamma RIIb_{EC}^{total,end}$ (center), or $FcRn_{EC}^{total,end}$ (right) was varied from 10-fold lower

to 10 times higher than the mean of their optimal range determined by Calipro. Vaccination was simulated at

the optimal time for a term newborn (t_{vax} = 25 weeks) for each FcR condition and the resulting percent

changes in anti-pertussis toxin IgG (α-PT IgG) in a newborn delivered at 40 weeks is depicted. The dashed

line represents the baseline simulation of an average patient using the optimized values of $FcRn_{STB}^{total,end}$,

$Fc\gamma RIIb_{EC}^{total,end}$, or $FcRn_{EC}^{total,end}$, which are marked by vertical lines. (D) The chart shows the effect of

increasing vaccine dosage on α-PT IgG transfer for patients with a 10-fold deficiency in $FcRn_{STB}^{total,end}$ or

$Fc\gamma RIIb_{EC}^{total,end}$ expression. Dosages are shown as fold-changes from the baseline simulation dosage. (E)

The effect of increasing vaccine-induced α-PT IgG affinity for FcRn or FcγRIIb on α-PT IgG transferred to a

fetus delivered at term.

With the assumption that maternal IgG protects the fetus via neutralization, the effectiveness

of maternal vaccination is ultimately determined by the concentration of maternal antigen-specific

IgG in the neonate. Having observed that FcRn and FcγRIIb are key predictors of IgG transport

efficiency (Fig 2A), we next investigated the relationship between FcR expression and α-PT IgG

transfer efficiency. Consistent with our previous observation that IgG transfer efficiency is highly

sensitive to $FcRn_{STB}^{total,end}$ and $Fc\gamma RIIb_{EC}^{total,end}$, we observed a 10-fold decrease in $FcRn_{STB}^{total,end}$ or

$Fc\gamma RIIb_{EC}^{total,end}$ diminished α-PT IgG transfer by 86 or 69% respectively, whereas $FcRn_{EC}^{total,end}$ had

a minor impact on α-PT IgG in the fetus (Fig 6C). Given that FcR expression is likely variable

across the population, we sought to determine vaccination modifications that could potentially

compensate for deficiency in either receptor. First, we found that patients with a 10-fold decrease in

$FcRn_{STB}^{total,end}$ could transfer comparable levels of α-PT IgG to their fetus by increasing the vaccine

dosage 8-fold, and a 10-fold decrease in $Fc\gamma RIIb_{EC}^{total,end}$ could be compensated for by increasing

the dosage 3-fold (Fig 6D). Naturally, vaccine dosage can only be increased to the extent that it

does not induce toxicity. Rather than increasing the *quantity* of maternal vaccine-induced IgG, we

asked whether modulating the *affinity* of vaccine-induced IgG to FcRs (*e.g.* via Fc glycosylation

modulation) could improve placental transfer. While precision vaccination is a relatively new

concept, recent studies have demonstrated that adjuvants play a key role in driving the glycan

profile of vaccine-induced IgG (52,53). Coincidentally, it has been shown that decreasing core

fucosylation can improve IgG binding to FcγRIIb and terminal galactosylation increases IgG affinity

to FcRn (12,54,55). Our model predicted that decreasing $K_D^{IgG1-4,FcRn}$ by 10-fold increased α-PT

IgG transfer by 30%, and a similar relationship was observed for $K_D^{IgG1-4,FcγRIIb}$ (Fig 6E).

Interestingly, the effect of increasing affinity for both receptors was greater than the sum of its parts,

suggesting affinity for both receptors can be targeted simultaneously to improve prenatal vaccine

efficacy.

Discussion

Placental antibody transfer is key to protect neonates against infection and to shape the trajectory

of early-life immune development (56). Recently, maternal prenatal immunization has been

employed to boost antigen-specific antibody transfer to the fetus, providing targeted immunity to the

newborn (57). Current prenatal vaccine regimens are applied at the population level, but variation in

gestational length, placental size and function, and maternal immune responses present

opportunities for patient-specific vaccine approaches. To this end, we developed a computational

model to elucidate the features regulating maternal-fetal IgG transport and to inform prenatal

immunization strategies. Our modeling approach enabled longitudinal probing of placental antibody

transfer dynamics, shedding light on an otherwise black-boxed system. In contrast to prior studies

attempting to uncover mechanisms of selective antibody transfer through a variety of *in vitro*, *ex*

vivo, and *in vivo* approaches, our computational approach recapitulates the dynamics of this

process as it occurs in humans (14,21,58,59).

Our dynamic model predicted subclass-specific IgG transfer ratios that agree with the trend

most commonly reported in the literature: IgG1 > IgG3 > IgG4 > IgG2 (11). While this phenomenon

of subclass selectivity has long been recognized, our model provides a mechanistic basis for

selective transfer of IgG subclasses based on competition for binding to FcRs that are expressed in

limited quantities. Previous studies have demonstrated that competition for FcRn binding

contributes to the short half-life of IgG3 (58). Our study builds upon this finding by revealing that

FcγRIIb in placental ECs similarly contributes to selective transfer by preferentially binding IgG1,

IgG3, and IgG4 with the highest affinity. Moreover, we built a mathematical model that quantifies

competition between two IgG subclasses as a function of FcR expression level, maternal IgG

subclass abundance, and Fc-FcR interaction affinity. Thus, the subclass transfer hierarchy of IgG1

> IgG3 > IgG4 > IgG2 can be explained by each subclass' relative abundance and affinity for FcRn

and FcγRIIb. Not only that, but inter-patient variability in subclass abundances, IgG-FcR affinity

arising from differential IgG glycosylation, and placental FcR expression can explain discrepancies

in this subclass transfer hierarchy across individuals and patient cohorts.

Previous studies have identified independent mechanisms of both FcγRIIb- and FcRn-

mediated EC transcytosis in placental ECs (29,45), yet the role of FcγRs in placental transport

remains highly contested (14,15,60). Our computational framework enabled us to disentangle the

contributions of FcRn and FcγRIIb to EC transcytosis, revealing that in an average placenta FcγRIIb

drives selectivity of IgG transfer and may play a key role in dynamic regulation of this process

across gestation. Together, these observations support the potential role of non-canonical FcγRs

driving antibody transfer dynamics and provide a mechanistic basis for previously reported

placental IgG "sieving" (12,13,61). Further evidence implicating FcγRIIb in placental transfer include

(1) our analysis of scRNA-seq data revealed an increase in *FCGR2B* transcript counts across

gestation and (2) placental ECs express the splice variant FcγRIIb2—the isoform capable of

endocytosis and expressed by liver sinusoidal endothelial cells, B cells, and macrophages (62–64).

In the same vein, previous studies indicate a potential role for FcγRIIIa in STB transcytosis

(12,15,35). However, the data supporting FcγRIIIa involvement in placental transfer are purely

associative and lack the necessary mechanistic insight to substantiate the inclusion of FcγRIIIa in

this model (12,65). To disentangle the roles of FcRn, FcγRIIb, and FcγRIIIa in IgG transcytosis,

future studies that uncover the expression patterns and dynamics of these receptors in human

placenta and/or experimentally test their effect on IgG transcytosis are warranted.

This unique modeling approach allowed us to predict optimized Tdap immunization

schedules and identify features affecting optimal Tdap timing during pregnancy, with broad

implications for prenatal vaccination both at the population level and as a basis for patient-specific

vaccine approaches. First, the model predicted that the optimal Tdap immunization time for an

average patient to maximize α-PT IgG transfer is during the second trimester for term and preterm

neonates alike, aligning with key findings from other recent studies that Tdap immunization earlier

in gestation provides higher α-PT and α-FHA IgG in the newborn (28). While this optimal Tdap

administration window was not sensitive to the magnitude of the maternal IgG response, it did

depend on the existence of a transient spike in IgG post-vaccination which is not inherent to IgG

responses to all antigens. Therefore, these insights may not necessarily be extrapolated to other

prenatal immunization programs. However, it is worth noting that our modeling framework is

modular, allowing similar optimization routines to be carried out for other prenatal vaccines so long

as there is sufficient data to inform simulations of the maternal antigen-specific IgG response

kinetics.

From a precision medicine standpoint, the model-driven finding that IgG competes for

placental transfer contingent upon subclass abundance, FcR expression, and Fc-FcR affinity

provides a basis for improving vaccine formulations and dosing strategies to optimize vaccine-

induced IgG transfer efficiency. The mathematical model devised in the current study provides a

flexible platform to test these novel vaccination strategies given a set of user-defined inputs. Recent

studies suggest that modulating vaccine parameters such as the nature of the target antigen, conjugate, number of doses, route of entry, and choice of adjuvant can fine-tune the pool of maternal vaccine-induced antibodies and the anti-infection capabilities of the IgG transferred to the newborn. For example, protein antigens tend to induce an IgG1 or IgG3 response, while polysaccharide antigens tend to induce an IgG2 response (66); more recent studies suggest that the N-glycan profile of vaccine-induced IgG can be modulated by adjuvant, providing a means to fine-tune Fc receptor binding potential (52,53,67,68). Using our model, we discovered the efficacy of maternal vaccines to transfer IgG to the fetus depended strongly on expression of FcRn in STBs and FcγRIIb in ECs. Our model predicted that readily tunable vaccine design choices such as timing of administration, dosage, and adjuvant are potential compensatory mechanisms to improve α-PT IgG transfer in patients with a deficiency in either FcRn in STBs or FcγRIIb in ECs. In order to make such recommendations, further research is needed to identify readily ascertainable biomarkers of placental FcR expression and/or transcytosis efficiency. As a more immediately actionable insight, we further uncovered a sensitivity of optimal immunization timing to volume parameters including maternal, fetal, and placental stromal volumes, which can be estimated by ultrasound imaging or alternative methods. Moreover, our model is able to quantitatively assess the effect of a perturbed maternal pool of antibodies due to various immunodeficiency disorders, autoimmune disorders and other disorders that modify the maternal pool of antibodies and vaccine response (69). As such, we envision this model as a tool to devise precision immunization approaches that account for maternal vaccination history, baseline pool of antibodies, autoimmune disorders or other immunodeficiencies, and risk for preterm delivery or other pregnancy complications to determine the optimal immunization approach per patient.

While this model-driven investigation generated many intriguing insights, there are a few limitations to consider. First, the current model relies on the assumption that all IgG is monovalent with a uniform glycan profile. This is an oversimplification as IgG glycosylation undergoes a dynamic shift during pregnancy (70), which can affect Fc receptor-mediated transfer (12,54).

Furthermore, in addition to monomeric IgG, IgG can potentially be transferred as antigen-antibody

complexes, which is currently understudied and is not incorporated in our current model (16,34).

Second, there is a stark lack of data surrounding human placental development and dynamic fetal

IgG levels throughout gestation; many model parameters were unavailable in the literature,

necessitating *in silico* parameter optimization. Consequently, we assumed several model

components to be constant, including maternal, fetal, and placental volumes and maternal IgG

levels. In actuality, these features undergo rapid remodeling and growth across gestation and

potentially fine-tune antibody transfer dynamics. Future iterations of the model may benefit from

more physiologically accurate dynamics of the developing maternal-fetal interface. To facilitate such

a model, further investigations detailing the dynamics of placental growth and FcR expression at the

protein level are warranted.

Despite its limitations, a key strength of our computational approach is the ability to simulate

placental transport dynamics which are prohibitively challenging to study *in vivo*. Using tightly

integrated modeling and experimental methods, we probed placental transport across scales in time

and space, enabling a more holistic view of the system compared to previous studies in the field.

Traditionally, placental IgG transport efficiency is benchmarked by a simple ratio of maternal to

umbilical cord IgG titer at birth, in part due to its ease of measurement and interpretability. Not only

does this metric mask the nuanced impact of the maternal pool of antibodies, but also it

insufficiently describes the highly dynamic and cumulative process of antibody transfer and is thus

not capable of describing: (1) important concepts such as placental FcR limitation and antibody

subclass competition, (2) the interplay between the maternal pool of antibodies and the vaccine-

induced maternal antibodies, and (3) the role of the complex placental functions that rapidly evolve

throughout gestation, among other reasons. Our mathematical model challenges this metric as a

true measure of "efficiency". With the advent and accessibility of state-of-the-art high-throughput

measurements, it is increasingly feasible to probe placental function and define one or more

quantitative measures of transfer efficiency that are conditioned on the maternal pool of antibodies

and are linked with distinct model parameters that account for placental size, perfusion, and Fc

receptor expression.

This study represents a major thrust forward in the field of maternal-neonatal immunology

and paves the way towards next generation vaccination programs to fine-tune neonatal immunity

against a broad range of pathogens. As current vaccination regimens are continually revised and

several novel vaccines are currently in clinical trials, now is an opportune time to increase the

prevalence of computational systems biology approaches in maternal-fetal medicine to improve

outcomes for vulnerable newborns.

519 Methods

520 Mechanistic model of placental IgG transport

An ordinary differential equation (ODE) model was formulated to simulate mass transport of IgG1,

IgG2, IgG3, and IgG4 across the maternal-fetal interface (Appendix S1). A schematic of the model

is given in Figure 1A. Fundamental model assumptions are as follows: (i) all processes obey the

law of mass action; (ii) IgG binds FcRn within STB endosomes at an acidified pH and FcγRIIb at the

EC surface (7,62,71,72); (iii) binding with either receptor initiates transcytosis; (iv) the effects of

FcRn-mediated IgG recycling to the apical STB surface are negligible; (v) unbound IgG in STB

endosomes undergoes lysosomal degradation; (vi) all blood volumes, IgG subclass concentrations

in maternal blood, and IgG-FcR affinities remain constant throughout gestation; (vii) rate

parameters (k_{up}, k_{trans}, and k_{deg}) are equal in STB and EC layers and across IgG subclasses; (viii)

stromal cells (e.g., Hofbauer cells, fibroblasts) do not interact with IgG or affect its transfer; (ix) the

effect of Fab-antigen interactions is neglected such that all IgG is monovalent; (x) FcR expression

increases parabolically across gestation informed by FcRn expression trends in rat placenta (31)

(see Appendix S1); (xi) STBs exclusively express FcRn and ECs express both FcRn and FcγRIIb;

(xii) the effects of Fc or Fab glycosylation on FcR binding are neglected. All model development and

corresponding *in silico* experiments were performed in MATLAB R2022a (Mathworks).

Parameter determination

In total, 17 parameters were derived from literature and 11 were optimized. K_D parameters were

ascertained from Bruhns et al, where the authors determined the equilibrium association constants

(K_A) experimentally by surface plasmon resonance (SPR) and we determined K_D by the formula K_D

= $1/K_A$ (8,37). We used these K_D values to determine parameters k_{on} and k_{off} for each Fc receptor

and IgG subclass pair. The value of k_{off} for IgG1-FcRn was determined experimentally by Suzuki *et*

al using SPR (38). We assumed k_{off} to be constant for all subclasses based on evidence from other

studies comparing multiple subclasses (73–76). We derived the value of k_{on} for each subclass-FcR

interaction from the formula k_{on} = k_{off} / K_D. The rate of IgG decay in the fetus (δ_{AB}) was derived from

the half-life of IgG1 measured in human infants by the formula δ_{AB} = $\ln(2)/t_{1/2}$ (39). Remaining

parameter ranges were determined by calibrating predictions of IgG subclass levels in the fetus to

data from Malek *et al*, where the authors quantified IgG subclass concentrations in umbilical cord

blood samples by enzyme linked immunosorbent assay (ELISA) (32). Initial conditions (maternal

IgG subclass concentrations) were directly ascertained from (32).

We performed parameter estimation for 11 parameters using CaliPro, a flexible optimization

software implemented in MATLAB and described by Joslyn et al (33). A key strength of CaliPro is

its ability to home in on a biologically relevant range of parameters that recapitulate the system

dynamics rather than minimizing a simple cost function. Briefly, we implemented CaliPro using Latin

Hypercube Sampling (LHS) from a physiologically plausible range of parameter values as

determined by benchmarking studies found in the literature. We defined the pass criteria as model

simulations within the upper standard deviation of the data set at the first time point, and are within

1.5 times the standard deviation at the final time point. Our termination criteria was set to 95% of

model runs meeting the pass criteria.

561   Global sensitivity analysis

Global sensitivity analysis was performed using LHS to sample the multi-dimensional parameter

space followed by an orthogonalized partial least squares regression (OPLSR) framework, as

described in (41,42). OPLSR is a multivariate regression technique that predicts a single outcome

(Y) from multiple predictors (X) and is especially suited to model systems with multicollinearity

among predictors (77). Briefly, numerous random perturbations to all *parameters* of the model were

made simultaneously and resulted in individual model runs per random parameter set. We then

leveraged OPLSR to build a model for an output of interest (Y) from the matrix of randomly

perturbed parameter sets (X). Specifically, to produce the matrix of predictors, 1000 perturbed

parameter sets were generated by LHS from a uniform distribution spanning the maximum

likelihood range for each parameter previously obtained from CaliPro. The number of parameter

sets tested was fixed to the lowest sample size needed to achieve stable sensitivity analysis results.

X is therefore a 1000 x 11 matrix (perturbations x parameters) and Y is a 1000 x 1 vector

(perturbations x outcome). First, an OPLSR model was constructed using the matrix of 1000

permuted parameter sets to predict the total concentration of IgG in the fetus at 40 weeks gestation.

A second OPLSR model was constructed in the same manner to predict entropy among IgG

subclasses in the fetus at 40 weeks gestation, defined as:

578
$$P_{IgGi} = \frac{IgG_i}{\sum_{i=1}^{4} IgG_i}$$

579
$$E = -\sum_{i=1}^{4} P_{IgGi}\log(P_{IgGi})$$

580   Where E is entropy and P_{IgGi} is the proportion of each subclass in the fetal blood. Model

581   orthogonalization was performed such that the direction of maximum variance was aligned with

582   latent variable 1 (LV1) (78). A third OPLSR model was constructed to determine the sensitivity of

583   the optimal t_{vax} to each parameter.

584   OPLSR model significance was determined empirically by comparing its predictive

power (Q^2) against randomly permuted models in a 5-fold cross validation scheme, where Q^2 is

defined as:

$$Q^2 = 1 - \frac{\sum_i (y_i - f'(x_i))^2}{\sum_i (y_i - \bar{y})^2}$$

Where x_i are the corresponding randomly perturbed parameter vectors in matrix X, y_i is each

individual outcome in vector Y, and $\bar{y}$ is the mean of all observations in Y.

HUVEC transcytosis assay

Antibody preparation

Purified human polyclonal IgG1 and IgG4 (Abcam, AB90283, AB183266) were dialyzed in 1x PBS

using Slide-A-Lyzer mini dialysis units with a 10K molecular weight cutoff (Invitrogen, 88404) to

remove sodium azide. Dialysis units were incubated at 4°C for 2 hours, then dialysis buffer was

changed and incubated again at 4°C for 2 hours. Protein was removed from the dialysis unit using a

micropipette and quantified using a NanoDrop Microvolume Spectrophotometer (Thermo Fisher).

Cell culture

6.5 mm diameter Transwell inserts (Corning, CLS3422) coated in 1.4% gelatin were seeded with

primary human umbilical vein endothelial cells (HUVEC) (Promocell C-12200, kindly provided by

Dan Gioeli at UVA) at a density of 1500 cells/mm^2 (50,000 cells/insert). Cells were cultured in

complete endothelial cell growth media (Promocell, C-22110) and incubated at 37°C and 5% CO_2.

After 24 hours, the cells were washed in sterile PBS and the media was changed.

Transcytosis assay

After 48 hours, the cells were washed in sterile PBS and media containing purified human IgG1 and IgG4 was added to the apical chamber. Cells were incubated at 37°C and 5% CO_2 for 120 minutes. IgG1 and IgG4 in the basolateral media were quantified using human ELISA) IgG1 and IgG4 quantification kits (Invitrogen, EHIGG1 and BMS2095) according to the manufacturer's protocol. All conditions were run in triplicate and ELISA measurements were run in duplicate. A 5-parameter logistic standard curve was generated for each assay. Absorbance was read at 450 nm with a correction of 590 nm using an Optima microplate reader within 1 hour of completing the assay. Statistical significance was determined with a one-tailed two-sample t-test comparing each sample containing IgG1 and IgG4 to the sample with IgG4 but without IgG1.

To confirm monolayer formation, membrane permeability was quantified for each assay by addition of FITC-dextran (4000 MW, Sigma Aldrich 46944) (0.5 mg/ml) to the apical chamber of each insert. FITC-dextran added to inserts without gelatin coating or HUVEC served as a negative control. The basolateral media was sampled after 120 minutes, and fluorescence was read using the Optima microplate reader. Monolayer permeability (%) was defined as:

$$Permeability = \frac{[basolat.FITC - dextran]_{HUVEC}}{[basolat.FITC - dextran]_{blank}} \times 100$$

Samples with FITC-dextran permeability greater than 3 standard deviations from the mean were excluded based on faulty monolayer formation (S5 Fig).

Mechanistic model of HUVEC transcytosis *in vitro*

To simulate the dynamics of IgG1 and IgG4 receptor-mediated transcytosis in HUVEC, a kinetic-dynamic model of FcRn-IgG binding and transcytosis in endothelial cells was formulated:

$$\frac{dIgG1}{dt} = -k_{on,1}[IgG1][FcRn] + k_{off,1}[C_1] \ (1)$$

$$629 \qquad \frac{dIgG4}{dt} = -k_{on,4}[IgG4][FcRn] + k_{off,4}[C_4] \text{ (2)}$$

$$630 \qquad \frac{dFcRn}{dt} = -[FcRn]\left(k_{on,1}[IgG1] + k_{on,4}[IgG4]\right) + \left(k_{off,1} + k_{trans}^{HUVEC}\right)[C_1] + \left(k_{off,4} + k_{trans}^{HUVEC}\right)[C_4] \text{ (3)}$$

$$631 \qquad \frac{dC_1}{dt} = k_{on,1}[IgG1][FcRn] - \left(k_{off,1} + k_{trans}^{HUVEC}\right)[C_1] \text{ (4)}$$

$$632 \qquad \frac{dC_4}{dt} = k_{on,4}[IgG4][FcRn] - \left(k_{off,4} + k_{trans}^{HUVEC}\right)[C_4] \text{ (5)}$$

$$633 \qquad \frac{dIgG1_T}{dt} = k_{trans}^{HUVEC}[C_1] \text{ (6)}$$

$$634 \qquad \frac{dIgG4_T}{dt} = k_{trans}^{HUVEC}[C_4] \text{ (7)}$$

Where [IgG1] and [IgG4] are apical concentrations of IgG1 and IgG4, [FcRn] is the concentration of

FcRn expression in HUVEC, [C_1] and [C_4] are the concentration of bound IgG-FcRn complexes,

k_{trans}^{HUVEC} is the rate of transcytosis, k_{on} is the forward rate of reaction, and k_{off} is the reverse rate of

reaction (Table 2). The model parameters were optimized by fitting to dynamic data of IgG4

transcytosis in HUVECs collected at 15, 30, 60, and 120 minutes ([IgG4$_{apical}$] = 0.5 and 0.125

mg/ml) (S5 Fig). To simplify this model and reveal the effect of subclass-subclass competition in Fc

receptor-mediated transcytosis, we derived the closed form of equation (5). We assumed that IgG1

and IgG4 are present in excess of FcRn, the system is at quasi-steady state such that

$$644 \qquad \frac{dC_1}{dt} = \frac{dC_4}{dt} = 0 \text{ (8)}$$

645 Where C is the concentration of IgG-FcRn complexes. Under these assumptions, equations (4) and

646 (5) can be rearranged as

$$647 \qquad \frac{k_{on,1}[IgG1][FcRn]}{k_{on,1}[IgG1][FcRn]\left(k_{off,1} + k_{trans}^{HUVEC}\right)} = [C_1] \text{ (9)}$$

$$648 \qquad \frac{k_{on,4}[IgG4][FcRn]}{k_{on,4}[IgG4][FcRn]\left(k_{off,4} + k_{trans}^{HUVEC}\right)} = [C_4] \text{ (10)}$$

649 The assumption is then made that the total concentration of receptor does not change such that

650
$$[FcRn] = [FcRn_0] - [C_1] - [C_4] \quad (11)$$

651 Substituting (11) into (10):

652
$$\frac{k_{on,4}[IgG4]([FcRn_0] - [C_1] - [C_4])}{k_{on,4}[IgG4][FcRn](k_{off,4} + k_{trans}^{HUVEC})} = [C_4] \quad (12)$$

653 And (9) into (12):

654
$$\frac{k_{on,4}[IgG4]([FcRn_0] - \dfrac{k_{on,1}[IgG1][FcRn]}{k_{on,1}[IgG1][FcRn](k_{off,1} + k_{trans}^{HUVEC})} - [C_4])}{k_{on,4}[IgG4][FcRn](k_{off,4} + k_{trans}^{HUVEC})} = [C_4] \quad (13)$$

655 Solving (13) for [C$_4$] yields the closed-form solution for IgG4-FcRn complex formation (79):

656
$$C_4 = \frac{FcRn_0[IgG4]}{K_{D,4}(1 + \dfrac{[IgG4]}{K_{D,4}} + \dfrac{[IgG1]}{K_{D,1}})} \quad (14)$$

657 Which quantifies the effect of competition between two IgG subclasses undergoing transcytosis in

658 the presence of one Fc receptor.

Pertussis (Tdap) immunization simulations

To simulate maternal Tdap immunization, a model of plasma cell activation and IgG secretion in

response to antigen stimulus was adapted from a previous study by White *et al* in the context of

malaria (46). The model consisted of 8 parameters, of which 4 were fitted parameters determined

by White *et al*, 1 was determined experimentally by Dari *et al*, and 3 were optimized in this study

(50). Parameters were tuned to fit previously published dynamic data of the maternal IgG response

to Tdap booster vaccination (47) (Table 3). We assumed several parameters which depend on the

host environment—namely, systemic cytokine concentrations and availability of FcRn for IgG

recycling and half-life extension—do not depend on antigen-specificity and thus are conserved

between malaria and pertussis contexts (80,81). Such parameters were preserved from White *et al*

and include the proportion of long-lived antibody secreting cells (L$_{ASC}$) to short-lived antibody

secreting cells (S$_{ASC}$), the lifespan of L$_{ASC}$ and S$_{ASC}$, and the antibody decay rate (46). On the other

hand, parameters which depend on the plasma cell niche and context of the specific antigenic

challenge (e.g., rates of plasma cell proliferation, antibody secretion, and the antigen decay rate)

were optimized to fit titers of anti-pertussis toxin IgG (α-PT IgG) in women of childbearing age

following a booster Tdap vaccination (47) (Table 3). Other important model assumptions include: (i)

IgG has a half-life of 31 days in the fetus; (ii) the dynamics of B cell activation—including antigen

presentation by follicular dendritic cells and cytokine secretion by T cells—and the contribution of

memory B cells are neglected; (iii) mothers have been previously exposed to pertussis toxin (i.e.,

this is a booster vaccine); (iv) all mothers have the same dynamic response to immunization; (v)

immunization resulted in equal parts IgG1, IgG2, IgG3, and IgG4 with uniform glycosylation profile

and immunoglobulin isotype switching was neglected; (vi) L_{ASC}s and S_{ASC}s secrete IgG with

identical subclass and glycosylation profile; (vii) the peak antibody titer is between 14 and 28 days

post-immunization; (viii) the IgG response to pertussis is the same in pregnant and non-pregnant

individuals (82). Maternal immunization was simulated by stimulating the maternal compartment

with antigen at a time t_{vax} (given in weeks gestational age) to induce an antibody response, which

directly fed into the maternal compartment of the placental IgG transport model. To determine the

optimal t_{vax} corresponding to maximum pertussis toxin-specific IgG transfer to the fetus, iterative

simulations were performed over a range of t_{vax} spanning 10-38 weeks gestational age. Sensitivity

of the optimal t_{vax} was determined by parameter perturbation and subsequent OPLSR model

construction, as described above.

Acknowledgements

We thank the members of the Dolatshahi lab, Drs. Donald J. Dudley and Loren D. Erickson for

helpful discussion and feedback. We would also like to thank Dr. Daniel Gioeli and Devin Roller for

providing cell lines and support with Transwell assays. We would like to acknowledge Drs. Yuk

Ming Dennis Lo, Peiyong Jiang, and Xi Hu from the Chinese University of Hong Kong for sharing

their published single cell transcriptomic data of placental tissue. We acknowledge Research

Computing at The University of Virginia for providing computational resources and technical support

that have contributed to the results reported within this publication. URL: https://rc.virginia.edu.

Research reported in this publication was supported by the Jeffress Trust Award in Interdisciplinary

Research as well as the National Institute of General Medical Sciences of the National Institutes of

Health under Systems & Biomedical Data Science training program (T32-GM145443). The content

is solely the responsibility of the authors and does not necessarily represent the official views of the

funding agencies and the National Institutes of Health.

Data and code availability

707   All data files and codes used to run the model can be found at our GitHub:

708   https://github.com/Dolatshahi-Lab/Transplacental-antibody-transfer-model. Model equations and *in*

709   *vitro* Transwell assay data are available as Supplemental files.

References

1. Pichichero ME. Challenges in vaccination of neonates, infants and young children. Vaccine. 2014 Jun 30;32(31):3886–94.

2. Olin A, Henckel E, Chen Y, Lakshmikanth T, Pou C, Mikes J, et al. Stereotypic Immune System Development in Newborn Children. Cell [Internet]. 2018;174(5):1277-1292.e14. Available from: https://doi.org/10.1016/j.cell.2018.06.045

3. Simister E, Israel EJ, Patel VK, Taylor SF, Marshak-Rothstein A. neonatal mice. acquisition of maternal IgG by fetal and 2-microglobulin-associated Fc receptor for Requirement for a beta. 1995 [cited 2022 Oct 19]; Available from: http://www.jimmunol.org/content/154/12/6246

4. Marchant A, Kollmann TR. Understanding the ontogeny of the immune system to promote immune-mediated health for life. Front Immunol. 2015;6(FEB):77.

5. Brambell FW. THE TRANSMISSION OF IMMUNITY FROM MOTHER TO YOUNG AND THE CATABOLISM OF IMMUNOGLOBULINS. Lancet. 1966 Nov 19;288(7473):1087–93.

6. Wilcox CR, Holder B, Jones CE. Factors affecting the FcRn-mediated transplacental transfer of antibodies and implications for vaccination in pregnancy [Internet]. Vol. 8, Frontiers in Immunology. Frontiers Media S.A.; 2017 [cited 2021 Jan 15]. p. 1. Available from: www.frontiersin.org

7. Simister NE. Placental transport of immunoglobulin G. In: Vaccine. Elsevier BV; 2003. p. 3365–9.

8. Bruhns P, Iannascoli B, England P, Mancardi DA, Fernandez N, Jorieux S, et al. Specificity and affinity of human Fc receptors and their polymorphic variants for human IgG subclasses. 2009 [cited 2021 Mar 16]; Available from: http://ashpublications.org/blood/article-pdf/113/16/3716/1308575/zh801609003716.pdf

9. Bright NA, Ockleford CD, Anwar M. Ontogeny and distribution of Fc gamma receptors in

the human placenta. Transport or immune surveillance? J Anat [Internet]. 1994 Apr [cited

2021 Jun 16];184 (Pt 2(Pt 2):297–308. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/8014121

10.   Kameda T, Koyama M, Matsuzaki N, Taniguchi T, Saji F, Tanizawa O. Localization of

three subtypes of Fcγ receptors in human placenta by immunohistochemical analysis.

Placenta. 1991 Jan 1;12(1):15–26.

11.   Clements T, Rice TF, Vamvakas G, Barnett S, Barnes M, Donaldson B, et al. Update on

Transplacental Transfer of IgG Subclasses: Impact of Maternal and Fetal Factors. Front

Immunol. 2020;11(September):1–17.

12.   Jennewein MF, Goldfarb I, Dolatshahi S, Cosgrove C, Noelette FJ, Krykbaeva M, et al.

Fc Glycan-Mediated Regulation of Placental Antibody Transfer. Cell [Internet].

2019;178(1):202-215.e14. Available from: https://doi.org/10.1016/j.cell.2019.05.044

13.   Dolatshahi S, Butler AL, Pou C, Henckel E, Bernhardsson AK, Gustafsson A, et al.

Selective Transfer of Maternal Antibodies in Preterm and Term Children. 2021 Dec 1

[cited 2022 Jun 7]; Available from: https://www.researchsquare.com

14.   Borghi S, Bournazos S, Thulin NK, Li C, Gajewski A, Sherwood RW, et al. FcRn, but not

FcγRs, drives maternal-fetal transplacental transport of human IgG antibodies. Proc Natl

Acad Sci U S A. 2020;117(23):12943–51.

15.   Martinez DR, Fouda GG, Peng X, Ackerman ME, Permar SR. Noncanonical placental Fc

receptors: What is their role in modulating transplacental transfer of maternal IgG?

Sheppard DC, editor. PLOS Pathog [Internet]. 2018 Aug 30 [cited 2021 Mar

15];14(8):e1007161. Available from: https://dx.plos.org/10.1371/journal.ppat.1007161

16.   Simister NE. Human placental Fc receptors and the trapping of immune complexes.

Vaccine. 1998;16(14–15):1451–5.

17.   Daëron M. FC RECEPTOR BIOLOGY. Annu Rev Immunol [Internet]. 1997 [cited 2022

Jun 2];15:203–37. Available from: www.annualreviews.org

18.   Atyeo C, DeRiso EA, Davis C, Bordt EA, De Guzman RM, Shook LL, et al. COVID-19

763   mRNA vaccines drive differential antibody Fc-functional profiles in pregnant, lactating,

764   and nonpregnant women. Sci Transl Med [Internet]. 2021 Oct 27 [cited 2021 Dec

765   6];13(617):8631. Available from:

766   https://www.science.org/doi/abs/10.1126/scitranslmed.abi8631

767   19.   Takizawa T, Anderson CL, Robinson JM. A Novel FcγR-Defined, IgG-Containing

768   Organelle in Placental Endothelium. J Immunol [Internet]. 2005 Aug 15 [cited 2021 Apr

769   29];175(4):2331–9. Available from: http://www.jimmunol.org/content/175/4/2331

770   20.   Saji F, Samejima Y, Kamiura S, Koyama M. Dynamics of immunoglobulins at the feto–

771   maternal interface. Reviews of Reproduction [Internet]. 1999 [cited 2021 Apr 20];81–9.

772   Available from:

773   http://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.983.3422&rep=rep1&type=pdf

774   21.   Rosenberg YJ, Ordonez T, Khanwalkar US, Barnette P, Pandey S, Backes IM, et al.

775   Evidence for the Role of a Second Fc-Binding Receptor in Placental IgG Transfer in

776   Nonhuman Primates. MBio. 2023 Mar 1;14(2):e0034123.

777   22.   Antohe F, Rădulescu L, Gafencu A, Ghe V, Simionescu M. Expression of functionally

778   active FcRn and the differentiated bidirectional transport of IgG in human placental

779   endothelial cells. Hum Immunol. 2001 Feb 1;62(2):93–105.

780   23.   Kiskova T, Mytsko Y, Schepelmann M, Helmer H, Fuchs R, Miedl H, et al. Expression of

781   the neonatal Fc-receptor in placental-fetal endothelium and in cells of the placental

782   immune system. Placenta. 2019 Mar 1;78:36–43.

783   24.   Niewiesk S. Maternal antibodies: Clinical significance, mechanism of interference with

784   immune responses, and possible vaccination strategies. Front Immunol.

785   2014;5(SEP):446.

786   25.   Faucette AN, Unger BL, Gonik B, Chen K. Maternal vaccination: moving the science

787   forward. Hum Reprod Update [Internet]. 2015 Jan 1 [cited 2023 Aug 31];21(1):119–35.

788    Available from: https://dx.doi.org/10.1093/humupd/dmu041

789  26. CDC, Ncird. DEPARTMENT OF HEALTH AND HUMAN SERVICES CENTERS FOR

790    DISEASE CONTROL AND PREVENTION Advisory Committee on Immunization

791    Practices (ACIP) Summary Report. 2016 [cited 2022 Apr 6]; Available from:

792    https://wwwdev.cdc.gov/vaccines/acip/meetings/downloads/min-archive/min-2016-10.pdf

793  27. Healy CM, Rench MA, Swaim LS, Smith EOB, Sangi-Haghpeykar H, Mathis MH, et al.

794    Association between Third-Trimester Tdap Immunization and Neonatal Pertussis

795    Antibody Concentration. JAMA - J Am Med Assoc [Internet]. 2018 Oct 9 [cited 2021 Apr

796    9];320(14):1464–70. Available from: https://jamanetwork.com/

797  28. Eberhardt CS, Blanchard-Rohner G, Lemaître B, Boukrid M, Combescure C, Othenin-

798    Girard V, et al. Maternal Immunization Earlier in Pregnancy Maximizes Antibody Transfer

799    and Expected Infant Seropositivity Against Pertussis. Clin Infect Dis [Internet]. 2016 Apr 1

800    [cited 2022 Apr 6];62(7):829–36. Available from:

801    https://academic.oup.com/cid/article/62/7/829/2463039

802  29. Ishikawa T, Takizawa T, Iwaki J, Mishima T, Ui-Tei K, Takeshita T, et al. Fc gamma

803    receptor IIb participates in maternal IgG trafficking of human placental endothelial cells.

804    Int J Mol Med [Internet]. 2015 May 1 [cited 2021 Jul 22];35(5):1273–89. Available from:

805    http://www.spandidos-publications.com/10.3892/ijmm.2015.2141/abstract

806  30. Lozano NA, Lozano A, Marini V, Saranz RJ, Blumberg RS, Baker K, et al. Expression of

807    FcRn receptor in placental tissue and its relationship with IgG levels in term and preterm

808    newborns. Am J Reprod Immunol. 2018;80(3):1–7.

809  31. Wang Y, Jiang X, He J, Diraviyam T, Zhang X. Quantitative Investigation on Correlation

810    Between IgG and FcRn During Gestation and Lactating Periods in Rat. Am J Reprod

811    Immunol. 2016;75(2):81–5.

812  32. Malek A, Sager R, Kuhn P, Nicolaides KH, Schneider H. Evolution of maternofetal

813    transport of immunoglobulins during human pregnancy. Am J Reprod Immunol.

1996;36(5):248–55.

33.   Joslyn LR, Kirschner DE, Linderman JJ. CaliPro: A Calibration Protocol That Utilizes

Parameter Density Estimation to Explore Parameter Space and Calibrate Complex

Biological Models. Cell Mol Bioeng. 2021 Feb 1;14(1):31–47.

34.   Palmeira P, Quinello C, Silveira-Lessa AL, Zago CA, Carneiro-Sampaio M. IgG placental

transfer in healthy and pathological pregnancies. Vol. 2012, Clinical and Developmental

Immunology. 2012.

35.   Atyeo C, Pullen KM, Bordt EA, Fischinger S, Burke J, Michell A, et al. Compromised

SARS-CoV-2-specific placental antibody transfer. Cell [Internet]. 2021 Feb 4 [cited 2021

Mar 23];184(3):628-642.e10. Available from: https://doi.org/10.1016/j.cell.2020.12.027

36.   Dolatshahi S, Butler AL, Siedner MJ, Ngonzi J, Edlow AG, Adong J, et al. Altered

Maternal Antibody Profiles in Women With Human Immunodeficiency Virus Drive

Changes in Transplacental Antibody Transfer. Clin Infect Dis [Internet]. 2022 Oct 12

[cited 2023 Aug 31];75(8):1359–69. Available from: https://dx.doi.org/10.1093/cid/ciac156

37.   Bruhns P, Jönsson F. Mouse and human FcR effector functions. Immunol Rev.

2015;268(1):25–51.

38.   Suzuki T, Ishii-Watabe A, Tada M, Kobayashi T, Kanayasu-Toyoda T, Kawanishi T, et al.

Importance of Neonatal FcR in Regulating the Serum Half-Life of Therapeutic Proteins

Containing the Fc Domain of Human IgG1: A Comparative Study of the Affinity of

Monoclonal Antibodies and Fc-Fusion Proteins to Human Neonatal FcR. J Immunol

[Internet]. 2010 [cited 2023 Aug 29];184:1968–76. Available from:

www.jimmunol.org/cgi/doi/10.4049/jimmunol.0903296

39.   Sarvas H, Sepp I, Kurikka S, Siegberg R, MJi O. Half-Life of the Maternal IgG1 Allotype

in Infants. J Clin Immunol. 1993;13(2).

40.   McKay MD, Beckman RJ, Conover WJ. A comparison of three methods for selecting

values of input variables in the analysis of output from a computer code. Technometrics.

2000;42(1):55–61.

41.   Marino S, Hogue IB, Ray CJ, Kirschner DE. A methodology for performing global

uncertainty and sensitivity analysis in systems biology. Vol. 254, Journal of Theoretical

Biology. Academic Press; 2008. p. 178–96.

42.   Surve S V., Myers PJ, Clayton SA, Watkins SC, Lazzara MJ, Sorkina A. Localization

dynamics of endogenous fluorescently labeled RAF1 in EGF-stimulated cells. Mol Biol

Cell [Internet]. 2019 Feb 15 [cited 2023 Apr 5];30(4):506–23. Available from:

https://www.molbiolcell.org/doi/10.1091/mbc.E18-08-0512

43.   Vento-Tormo R, Efremova M, Botting RA, Turco MY, Vento-Tormo M, Meyer KB, et al.

Single-cell reconstruction of the early maternal–fetal interface in humans. Nature

[Internet]. 2018 Nov 15 [cited 2021 Jan 15];563(7731):347–53. Available from:

https://pubmed.ncbi.nlm.nih.gov/30429548/

44.   Tsang JCH, Vong JSL, Ji L, Poon LCY, Jiang P, Lui KO, et al. Integrative single-cell and

cell-free plasma RNA transcriptomics elucidates placental cellular dynamics. Proc Natl

Acad Sci U S A [Internet]. 2017 Sep 12 [cited 2021 Mar 29];114(37):E7786–95. Available

from: www.pnas.org/cgi/doi/10.1073/pnas.1710470114

45.   Antohe F, Rădulescu L, Gafencu A, Ghe V, Simionescu M. Expression of functionally

active FcRn and the differentiated bidirectional transport of IgG in human placental

endothelial cells. Hum Immunol [Internet]. 2001 [cited 2022 Jun 7];62(2):93–105.

Available from: https://pubmed.ncbi.nlm.nih.gov/11182218/

46.   White MT, Griffin JT, Akpogheneta O, Conway DJ, Koram KA, Riley EM, et al. Dynamics

of the Antibody Response to Plasmodium falciparum Infection in African Children. J Infect

Dis [Internet]. 2014 Oct 1 [cited 2022 Sep 22];210(7):1115–22. Available from:

https://academic.oup.com/jid/article/210/7/1115/2908670

47.   van der Lee S, van Rooijen DM, de Zeeuw-Brouwer ML, Bogaard MJM, van Gageldonk

PGM, Marinovic AB, et al. Robust humoral and cellular immune responses to pertussis in

adults after a first acellular booster vaccination. Front Immunol. 2018 Apr 4;9(APR):681.

48.   Knuutila A, Versteegen P, Barkoff AM, van Gageldonk P, Mertsola J, Berbers G, et al.

Pertussis toxin neutralizing antibody response after an acellular booster vaccination in

Dutch and Finnish participants of different age groups. Emerg Microbes Infect [Internet].

2022 [cited 2023 Apr 12];11(1):956–63. Available from:

https://www.tandfonline.com/doi/abs/10.1080/22221751.2022.2053364

49.   Hovingh ES, Kuipers B, Bonačić Marinović AA, Jan Hamstra H, Hijdra D, Mughini Gras L,

et al. Detection of opsonizing antibodies directed against a recently circulating Bordetella

pertussis strain in paired plasma samples from symptomatic and recovered pertussis

patients. Sci Reports 2018 81 [Internet]. 2018 Aug 13 [cited 2023 Apr 13];8(1):1–11.

Available from: https://www.nature.com/articles/s41598-018-30558-8

50.   Dari A, Boulton M, Neyens M, Le Gars M, Valenzuela B, Shukarev G, et al. Quantifying

Antibody Persistence After a Single Dose of COVID-19 Vaccine Ad26.COV2.S in

Humans Using a Mechanistic Modeling and Simulation Approach. Clin Pharmacol Ther.

2022;

51.   Vilajeliu A, Goncé A, López M, Costa J, Rocamora L, Ríos J, et al. Combined tetanus-

diphtheria and pertussis vaccine during pregnancy: transfer of maternal pertussis

antibodies to the newborn. Vaccine. 2015 Feb 18;33(8):1056–62.

52.   Vestrheim AC, Moen A, Egge-Jacobsen W, Reubsaet L, Halvorsen TG, Bratlie DB, et al.

A pilot study showing differences in glycosylation patterns of igg subclasses induced by

pneumococcal, meningococcal, and two types of influenza vaccines. Immunity, Inflamm

Dis. 2014 Aug 1;2(2):76–91.

53.   Francica JR, Zak DE, Linde C, Siena E, Johnson C, Juraska M, et al. Innate

transcriptional effects by adjuvants on the magnitude, quality, and durability of HIV

envelope responses in NHPs.

54.   Subedi GP, Barb AW. The immunoglobulin G1 N-glycan composition affects binding to

each low affinity Fc γ receptor. MAbs [Internet]. 2016 Nov 16 [cited 2023 Aug

28];8(8):1512–24. Available from:

https://www.tandfonline.com/action/journalInformation?journalCode=kmab20

55.   Cambay F, Forest-Nault C, Dumoulin L, Seguin A, Henry O, Durocher Y, et al.

Glycosylation of Fcγ receptors influences their interaction with various IgG1 glycoforms.

Mol Immunol. 2020 May 1;121:144–58.

56.   Brophy-Williams S, Fidanza M, Marchant A, Way SS, Kollmann TR. One vaccine for life:

Lessons from immune ontogeny. J Paediatr Child Health [Internet]. 2021 Jun 1 [cited

2021 Jun 30];57(6):782–5. Available from:

https://onlinelibrary.wiley.com/doi/full/10.1111/jpc.15511

57.   Marchant A, Sadarangani M, Garand M, Dauby N, Verhasselt V, Pereira L, et al. Maternal

immunisation: collaborating with mother nature. Lancet Infect Dis. 2017 Jul 1;17(7):e197–

208.

58.   Stapleton NM, Andersen JT, Stemerding AM, Bjarnarson SP, Verheul RC, Gerritsen J, et

al. Competition for FcRn-mediated transport gives rise to short half-life of human IgG3

and offers therapeutic potential. Nat Commun 2011 21 [Internet]. 2011 Dec 20 [cited

2022 Dec 9];2(1):1–9. Available from: https://www.nature.com/articles/ncomms1608

59.   Jennewein MF, Abu-Raya B, Jiang Y, Alter G, Marchant A. Transfer of maternal immunity

and programming of the newborn immune system.

60.   Rosenberg YJ, Ordonez T, Khanwalkar US, Barnette P, Pandey S, Backes IM, et al.

Evidence for the Role of a Second Fc-Binding Receptor in Placental IgG Transfer in

Nonhuman Primates. MBio [Internet]. 2023 Mar 1 [cited 2023 Jun 26];14(2):e0034123.

Available from: https://journals.asm.org/journal/mbio

61.   Martinez DR, Fong Y, Hang Li S, Yang F, Jennewein MF, Weiner JA, et al. Fc

Characteristics Mediate Selective Placental Transfer of IgG in HIV-Infected Women. Cell

[Internet]. 2019 [cited 2021 Jan 13];178:190-201.e11. Available from:

https://doi.org/10.1016/j.cell.2019.05.046

62.   Mousavi SA, Sporstøl M, Fladeby C, Kjeken R, Barois N, Berg T. Receptor-mediated

endocytosis of immune complexes in rat liver sinusoidal endothelial cells is mediated by

FcγRIIb2. Hepatology. 2007 Sep;46(3):871–84.

63.   Mishima T, Kurasawa G, Ishikawa G, Mori M, Kawahigashi Y, Ishikawa T, et al.

Endothelial Expression of Fc Gamma Receptor IIb in the Full-term Human Placenta.

Placenta. 2007 Feb 1;28(2–3):170–4.

64.   Daëron M. FC RECEPTOR BIOLOGY [Internet]. Vol. 15, Annu. Rev. Immunol. 1997.

Available from: www.annualreviews.org

65.   Atyeo C, DeRiso EA, Davis C, Bordt EA, De Guzman RM, Shook LL, et al. COVID-19

mRNA vaccines drive differential antibody Fc-functional profiles in pregnant, lactating,

and nonpregnant women. Sci Transl Med [Internet]. 2021 Oct 27 [cited 2021 Dec

7];13(617):8631. Available from:

https://www.science.org/doi/abs/10.1126/scitranslmed.abi8631

66.   Vidarsson G, Dekkers G, Rispens T. IgG subclasses and allotypes: From structure to

effector functions. Front Immunol. 2014;5(OCT):520.

67.   Alter G, Ottenhoff THM, Joosten SA. Antibody glycosylation in inflammation, disease and

vaccination. Semin Immunol. 2018 Oct 1;39:102–10.

68.   Mahan AE, Jennewein MF, Suscovich T, Dionne K, Tedesco J, Chung AW, et al.

Antigen-Specific Antibody Glycosylation Is Regulated via Vaccination. PLoS Pathog.

2016 Mar 1;12(3).

69.   Jefferis R, Kumararatne DS. Selective IgG subclass deficiency: quantification and clinical

relevance. Clin exp Immunol. 1990;81:357–67.

70.   Einarsdottir HK, Selman MHJ, Kapur R, Scherjon S, Koeleman CAM, Deelder AM, et al.

Comparison of the Fc glycosylation of fetal and maternal immunoglobulin G. Glycoconj J

[Internet]. 2013 Feb 10 [cited 2021 Jan 15];30(2):147–57. Available from: www.ms-

utils.org/Xtractor2D.

71.   Simister NE, Rees AR. Isolation and characterization of an Fc receptor from neonatal rat

small intestine. Eur J Immunol [Internet]. 1985 Jan 1 [cited 2022 Nov 21];15(7):733–8.

Available from: https://onlinelibrary.wiley.com/doi/full/10.1002/eji.1830150718

72.   Hunziker W, Mellman I. Expression of Macrophage-Lymphocyte Fc Receptors in Madin-

Darby Canine Kidney Cells: Polarity and Transcytosis Differ for Isoforms with or without

Coated Pit Localization Domains. [cited 2023 Aug 31]; Available from:

http://rupress.org/jcb/article-pdf/109/6/3291/1466358/3291.pdf

73.   Zinkhan S, Thoms F, Augusto G, Vogel M, Bachmann MF. On the role of allergen-

specific IgG subclasses for blocking human basophil activation. Front Immunol. 2022 Oct

6;13:892631.

74.   Derebe MG, Nanjunda RK, Gilliland GL, Lacy ER, Chiu ML. Human IgG subclass cross-

species reactivity to mouse and cynomolgus monkey Fcγ receptors. 2018 [cited 2023

Aug 29]; Available from: https://doi.org/10.1016/j.imlet.2018.02.006

75.   Szikora B, Hiripi L, Bender B, Kacskovics I, Iliás A. Characterization of the interactions of

rabbit neonatal Fc receptor (FcRn) with rabbit and human IgG isotypes. PLoS One

[Internet]. 2017 Sep 1 [cited 2023 Aug 29];12(9):e0185662. Available from:

https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0185662

76.   Datta-Mannan A, Witcher DR, Tang Y, Watkins J, Wroblewski VJ. Monoclonal Antibody

Clearance IMPACT OF MODULATING THE INTERACTION OF IgG WITH THE

NEONATAL Fc RECEPTOR *. 2007;

77.   Wold S, Sjostrom M, Eriksson L, Sweden˚˚ S. PLS-regression: a basic tool of

chemometrics [Internet]. Vol. 58, Chemometrics and Intelligent Laboratory Systems.

[cited 2021 May 20]. Available from: www.elsevier.comrlocaterchemometrics

78.   Trygg J, Wold S. Orthogonal projections to latent structures (O-PLS). J Chemom.

2002;16(3):119–28.

79. Salahudeen MS, Nishtala PS. An overview of pharmacodynamic modelling, ligand-binding approach and its application in clinical practice. Vol. 25, Saudi Pharmaceutical Journal. Elsevier B.V.; 2017. p. 165–75.

80. Wilmore JR, Allman D. Here, There, and Anywhere? Arguments for and against the Physical Plasma Cell Survival Niche. J Immunol [Internet]. 2017 Aug 1 [cited 2023 Apr 12];199(3):839–45. Available from: https://journals.aai.org/jimmunol/article/199/3/839/106408/Here-There-and-Anywhere-Arguments-for-and-against

81. Cyster JG, Allen CDC. B Cell Responses: Cell Interaction Dynamics and Decisions. Cell. 2019 Apr 18;177(3):524–40.

82. Huygen K, Caboré RN, Maertens K, Van Damme P, Leuridan E. Humoral and cell mediated immune responses to a pertussis containing vaccine in pregnant and nonpregnant women. Vaccine. 2015 Aug 7;33(33):4117–23.