GABA, glucose and insulin orchestrate CD4[+] T cells effector functions

Jin[1]# Z, Hammoud[1]# H, Bhandage[1] AK, Korol[1] SV, Trujeque-Ramos[1] O, Koreli[1] S., Chowdhury[1]

AI, Sandbaumhüter[2] FA, Jansson[2] ET, Andrén[2] PE, Bergsten[1] P, Kamali-Moghaddam[3] M, Birnir*

B

[1] Department of Medical Cell Biology, Uppsala University, Uppsala, Sweden

[2] Department of Pharmaceutical Biosciences, Uppsala University, Uppsala, Sweden.

[3] Department of Immunology, Genetics and Pathology, Science for Life Laboratory, Uppsala
University, Uppsala, Sweden
10

 # Contributed equally

* Correspondence

Prof Bryndis Birnir

Department of Medical Cell Biology

Uppsala University

Uppsala, 75124, Sweden

Email: bryndis.birnir@mcb.uu.se

Tel: +46765620243

Summary

Metabolic programs of immune cells are closely linked to their effector functions[1], where physiological molecules provide environmental cues and guidance[2-5]. Exactly how it happens is still being unraveled. Insulin maintains normal blood glucose levels[6] and glucose is the main source of energy and a precursor for many biomolecules in T cells[7,8], whereas γ-aminobutyric acid (GABA), best known as a neurotransmitter, is increasingly recognized as a regulatory molecule in the immune system[4,9]. Here, we demonstrate that GABA-mediated reduction of metabolic activity and release of inflammatory molecules, including IFNγ and IL-10, was abolished in human CD4$^+$ T cells, when the glucose concentration was elevated above normal levels. In a glucose concentration-dependent manner, insulin enhanced the GABA$_A$ receptors activated currents and GABA-dependent Ca^{2+} influx. GABA decreased, whereas insulin maintained glycolysis but in a SGLT (Na$^+$-glucose transporter)-dependent manner, revealing expression of SGLTs in activated CD4$^+$ T cells. The SGLTs antagonist phlorizin, alone or together with GABA, restored the inhibition of IFNγ and IL-10 release in presence of high glucose. This study exposes concerted effects of GABA, glucose and insulin on CD4$^+$ T cells metabolic activity and release of inflammatory molecules, and identifies a role for SGLTs in CD4$^+$ T cells function.

Main

GABA inhibition is affected by glucose

Physiological blood glucose concentration 2h after a meal, is normally maintained around 5.6 mM in healthy individuals, while it is higher in diseases such as diabetes (> 7 mM) or Covid-19[10]. Metabolic activity of T cells is closely related to their rate of proliferation. We examined effects of four glucose concentrations on the GABA inhibition of metabolic activity of CD4[+] T cells (N=35–53 human donors; Figure 1a; Extended Data Table 1; Extended Data Figure 1). At glucose concentrations higher than the physiological level (10 mM and 16.7 mM), the inhibition by GABA was abolished (Figure 1a; Extended Data Figure 2a). When the GABA responses were compiled based on low-normal (2.8 mM, 5.6 mM) or high (10 mM, 16.7 mM) glucose concentrations a clear shift was observed for GABA's inhibitory response. At the level 0.9 or lower metabolic activity (Figure 1b left panel), cells from 74% of the donors were inhibited by GABA in the low-normal glucose group. The metabolic activity of cells from these same donors were not inhibited at the higher glucose concentrations (Figure 1b right panel; Extended Data Figure 2b). Intrigued by these results, we sought whether glucose affected other immunological functions regulated by GABA. Since glycolysis feeds substrates into anabolic metabolism, we studied if glucose modulated the GABA-inhibition of release of biomolecules from CD4[+] T cells from the same donors (N=18) cultured in low-normal or high glucose. In each sample, 92 inflammation-related biomolecules commonly associated with inflammation were measured using Olink Target Inflammation protein panel, which is based on the multiplex proximity extension assay (PEA) technology. A total of 59 different proteins were detected. Similar to the results from the metabolic activity, the GABA-inhibition of proteins released from CD4[+] T cells was glucose concentration-dependent (Figure 1c). The GABA inhibition was optimal at the physiological glucose concentration (5.6 mM), where

GABA decreased the release of 37 proteins. In contrast, GABA did not affect release of

proteins in neither 10 mM nor 16.7 mM glucose (Figure 1c). ELISA confirmed the PEA results

for two cytokines; interferon gamma (IFNγ), the principal cytokine of CD4$^+$ T helper1 cells and

interleukin-10 (IL-10), the principal cytokine for CD4$^+$ T helper2 and T regulatory cells (Figure

1d, e). In contrast, in absence of GABA, increasing the glucose concentration alone enhanced

release of IFNγ (Extended Data Figure 2c) but not IL-10 (Extended Data Figure 2d). Thus, GABA-

dependent regulation of CD4$^+$ T cells is linked to the extracellular glucose concentration.

Insulin is normally associated with glucose homeostasis. However, at least in excitable

cells, insulin is also a regulator of GABA signaling[11-13]. The insulin receptor is not expressed in

resting T cells but is prominent at 48 h post-activation[14].GABA activates two types of receptors

in cell membranes; the GABA$_A$ receptor (GABA$_A$R), a pentameric chloride ion channel, that

opens when GABA binds to the receptor and the GABA$_B$ receptor (GABA$_B$R), a G-protein

coupled receptor, that is formed as a dimer. To-date, only one of the two GABA$_B$ isoforms

required for function has been detected in immune cells, whereas several GABA$_A$ receptor

subunits are identified in T cells[4,15]. The GABA$_A$R ρ2 (*GABRR2*) is a highly expressed subunit in

peripheral blood mononuclear cells[15]. Irrespective of glucose concentration, activation of

CD4$^+$ T cells decreased the expression of the *GABRR2* transcript relative to resting cells,

whereas insulin increased the expression of the *GABRR2* (Figure 1f). The protein was detected

by a GABRR2 specific antibody by fluorescent labelling the CD4$^+$ T cells (Figure 1g).

Insulin enhances the GABA-activated currents

We measured the GABA-activated GABA$_A$R currents in intact activated CD4$^+$ cells using

perforated patch-clamp electrophysiology. Since the intracellular environment is unchanged,

intracellular signaling that may take place is expected to function normally. The 500 nM GABA-

activated GABA$_A$R-mediated currents were blocked by the GABA$_A$R antagonist picrotoxin and

enhanced by insulin at 5.6 mM glucose (Figure 2a). The apparent GABA affinity (EC_{50}) for

$GABA_A R$ was similar in 5.6 mM and 16.7 mM glucose, 0.7 nM and 0.9 nM GABA, respectively.

Insulin then shifted the EC_{50} values more than 10-fold, to 0.04 nM and 0.06 nM GABA (Figure

2b, c). The $GABA_A R$ agonists muscimol and TACA evoked currents similar to GABA (Figure 2d).

The currents reversed at −37 mV (Extended Data Figure 3a), close to the calculated reversal

potential for chloride (E_{Cl} = −39 mV). By regulating the membrane potential these very

sensitive $GABA_A R$ channels may affect activity of other ion channels or flow of ions across the

cell membrane[16].

Insulin enhances GABA-activated Ca^{2+} entry

The cytoplasmic concentration of Ca^{2+} ions is tightly regulated in T cells but can be

increased by release from intracellular stores or by entry through plasma membrane Ca^{2+} ion

channels[17]. A representative live cell-Ca^{2+} recording trace measured from activated $CD4^+$ T

cells (Figure 2e) showed that acute GABA (500 nM) application induced transient calcium

signal, which was, in fact, influx of extracellular calcium and not due to release from

intracellular stores as no signal could be detected in absence of extracellular calcium (Figure

2e, left panel). Analyzed live cell-Ca^{2+} recording data from four donors is shown in Figure 2e,

right panel. Transient change in glucose from 5.5 to 16.7 mM did not alter the 500 nM GABA-

activated Ca^{2+} response (Extended Data Figure 3b). Insulin incubation of cells enhanced the

GABA induced Ca^{2+} signals in 5.5 mM but not in 16.7 mM glucose (Figure 2f), whereas no

differences were observed for baseline Ca^{2+} levels in cells with or without insulin incubation

(Extended Data Figure 3c). The GABA-activated Ca^{2+} signal was blocked by the $GABA_A R$

antagonist picrotoxin (Figure 2g), while no effect was observed by the $GABA_B R$ antagonist,

CGP52432 (Extended Data Figure 3d). This prompted us to examine dose-response

relationship between GABA and the Ca^{2+} signals in cells with or without insulin incubation

(Figure 2h, i, j; Extended Data Figure 3e). Representative time-lapse recording of a CD4+ T cell,

16.7 mM glucose, with insulin, demonstrates that raising GABA concentrations induced

increasing Ca^{2+} signals (Figure 2h). Insulin incubation of the cells lowered the EC_{50} values of

the GABA-activated Ca^{2+} signals. The apparent GABA affinity of the Ca^{2+} signal was 55 nM and

113 nM GABA in 5.5 mM and 16.7 mM glucose, and shifted more than 1000-fold to 0.009 nM

and 0.09 nM GABA with insulin, respectively (Figure 2i, j). The Ca^{2+} response to $GABA_AR$

agonists, muscimol and TACA, and antagonists, TPMPA and picrotoxin, further confirmed the

$GABA_AR$ connection in the evoked Ca^{2+} signal (Figure 2k, l, m). Ca^{2+} release-activated

Ca^{2+} (CRAC) channels mediate store operated calcium entry (SOCE) in T cells that is central to

the Ca^{2+} homeostasis and cytoplasmic Ca^{2+} levels[17]. The GABA-activated Ca^{2+} signal was

inhibited by 1 μM YM48583 (Figure 2n), an antagonist of SOCE, revealing the Ca^{2+} ion channel

participating in the cascade. The tight scaling of the Ca^{2+} signal from $GABA_AR$ activation to Ca^{2+}

entry and the sensitization by insulin identifies a specific, focused mechanism regulating Ca^{2+}

signaling in CD4+ T cells.

Insulin is known to regulate intracellular signaling and gene expression, whereas less

is known about effects of GABA. However, out of the nine examined genes encoding

transcription factors or kinases, only *NFATC2* (nuclear factor of activated T cells) was

significantly reduced by GABA at 5.6 mM glucose (Extended Data Table 2). NFATs proteins are

activated by increased Ca^{2+} in the cytoplasm and translocate to the nucleus, where they

regulate transcription of many proteins including inflammatory molecules[18].

GABA regulation of glycolysis is glucose-dependent

That GABA regulated expression of signaling pathways and proteins was revealed by

mass spectrometry (MS) of activated CD4+ T cells cultured in the presence or absence of GABA

(Figure 3a, Extended Data Table 3). Hexokinase 1 (HK1), the first enzyme in the glycolytic

pathway, was significantly down-regulated by GABA in both 5.6 mM and 16.7 mM glucose

(Figure 3b). HK activity was negligible under resting conditions but increased about 60 times

in activated CD4[+] T cells in both 5.6 mM and 16.7 mM glucose. Intriguingly, GABA decreased

the HK activity only at 5.6 mM glucose and insulin had no further effect (Figure 3c). T cells

switch to aerobic glycolysis upon activation as their predominant way to generate adenosine

5´-triphosphate (ATP) and accumulate biomass[19,20], the so-called Warburg effect. It results in

glucose-derived pyruvate being converted to lactate that can be detected by measuring

extracellular acidification rate (ECAR)[21]. Since HK is the gate-keeper for the glycolytic pathway,

we examined if glucose, GABA or insulin modulated glycolysis. Representative ECAR traces

and average values for activated CD4[+] T cells at 5.6 mM and 16.7 mM glucose in the presence

or absence of GABA and insulin are shown in Figure 3d and e. GABA but not insulin inhibited

glycolysis and glycolytic capacity but only in 5.6 mM glucose. No effect of GABA at higher

glucose suggested the possibility that either the intracellular glucose might affect the GABA

inhibition of glycolysis or there were other competing mechanisms at work.

SGLT2 in CD4[+] T cells

Facilitated glucose transport is the main route for glucose entry into activated T cells[3].

We confirmed the dominance of the facilitated transport by examining the effects of an

antagonist of GLUTs, WZB117 (WZB), on the cellular metabolic activity of activated CD4[+] T cells

(Figure 4a), whereas phlorizin, an antagonist of Na[+]-dependent glucose transport (SGLT1 and

SGLT2)[22], had no effect at either glucose concentration (Figure 4a). Nevertheless, *SLC5A2*, the

gene encoding SGLT2, was detected by RNAseq analysis from activated CD4[+] T cells (Figure

4b) and the protein was also expressed (Figure 4c, d). In presence of phlorizin or empagliflozin,

a more specific SGLT2 antagonist and a prescription medicine in type 2 diabetes, a significant

reduction in glucose uptake was observed in insulin incubated cells (Figure 4e). Phlorizin alone

reduced glycolysis and glycolytic capacity at both 5.6 mM and 16.7 mM glucose (Figure 4f),

whereas insulin reduced or abolished the inhibition by phlorizin (Figure 4f). Since phlorizin

reduced glycolysis, it could be expected to reduce synthesis of biomolecules such as IFNγ and

IL-10, similar effects to what we observed for GABA (see Figure 1c, d, e). Indeed, phlorizin

reduced release of IFNγ and IL-10 at both 5.6 mM and 16.7 mM glucose even in the presence

of insulin (Figure 4g, h) with the exception of IFNγ at 16.7 mM glucose with insulin, where only

phlorizin together with GABA reduced IFNγ release, approximately 50%. In general, there was

a trend for GABA to enhance the phlorizin reduction of release of IFNγ and IL-10. The results

demonstrate functional Na$^+$-dependent glucose carriers in CD4$^+$ T cells and identify their

relevance for production of cytokines.

Discussion

T cells modulate their metabolism to adapt to different environments[1,19,20]. When

CD4$^+$ T cells are activated, their metabolic activity is altered to meet the demands of cell

growth, proliferation and effector functions. Here we revealed that GABA, glucose and insulin

together orchestrate functions of CD4$^+$ T cell, to coordinate and to set the activity level of

distinct cellular pathways linked to glucose metabolism and that are central for

immunometabolism and T cells effector functions (Extended Figure 4).

Proliferating T cells shift their metabolic phenotype to aerobic glycolysis at the expense

of mitochondrial oxidative phosphorylation[19,23]. Although this switch is inefficient in terms of

generating ATP, it confers advantage for the biosynthetic pathways that branch out from

glycolysis[19,20]. GABA decreased cellular metabolic activity and release of proteins in a glucose

concentration-dependent manner. Insulin shifted the GABA$_A$R signaling by making it

supersensitive to GABA and consequently enhanced Ca^{2+} entry through CRAC channels, an

elegant way of modulating intracellular Ca^{2+} levels and intracellular processes affecting T cells

effector functions.

It was surprising that extracellular glucose modulated the GABA inhibition to regulate

effector functions of T cells. Although glucose transport in $CD4^+$ T cells is mainly mediated by

facilitated glucose transporters, the present results identified significant Na^+-dependent

glucose transport relevant for the effector functions. HK is the first enzyme in the glycolytic

pathway and by converting glucose to glucose-6-phosphate (G-6-P), traps glucose within the

cell[7]. Human HK has normally maximum activity in 5 mM glucose in the presence of natural

inhibitors, such as G-6-P[24], but HK activity is increased at glucose concentrations > 5 mM[24].

This is in line with our results, where GABA only succeeded decreasing HK enzymatic activity

in 5.6 mM glucose. Na^+-dependent glucose transport drives glucose into cells even at high

intracellular glucose concentrations[22]. Glucose competes with G-6-P for the binding site on

HK[24], decreasing the inhibition of HK[24], which results in increased substrate availability for

biosynthesis[8]. Aerobic glycolysis directly regulates synthesis of IFNγ[25] by engaging

glyceraldehyde-3-phosphate dehydrogenase (GDPDH) and, thereby, relieving its inhibition of

IFNγ mRNA translation. In concordance, activated $CD4^+$ cells secreted more IFNγ in high

glucose. GABA promoted a glycolytic switch to lower values for both glycolysis and the

glycolytic capacity in 5.6 mM glucose that was gone in 16.7 mM glucose, whereas phlorizin

was effective inhibitor at both glucose concentrations and, in particular, in the absence of

insulin. Discovering that $GABA_AR$ activation together with SGLT2 inhibition can dramatically

reduce the levels of IFNγ and IL-10, at both normal and high glucose concentrations, has

implications for regulation of inflammation, as availability of biosynthetic intermediates-

production fuels effector signaling molecules and functions[8].

GABA$_A$Rs are widespread in the brain and are present in many other tissues[26] but

GABA$_A$Rs containing GABRR2 (ρ2 subunit) are rare outside the immune system, with the

exception of the retina[27]. GABA$_A$R-ρ2 might be a valuable and a rather specific drug target[28].

SGLT2 inhibitors such as empagliflozin, are already in clinical use[29] and might be considered

to cover new areas of application. Strengthening the GABA response and inhibiting SGLT2 in

CD4$^+$ T cells might be valuable tools[30-32], when regulating inflammation is desirable.

In conclusion, physiological factors such as glucose, GABA and insulin guide the

adaptation of CD4$^+$ T cells and modify their effector functions. The reduction of metabolic

activity and release of inflammatory molecules in response to GABA or SGLT2 inhibitors,

revealed a switch that can be turned on or off. The study unveils the mechanisms of GABA

modulation of CD4$^+$ T cells and identifies a critical role for SGLTs in immune function. The

results have implications for treatment of a number of diseases including diabetes but also

maladies such as Covid-19 and cytokine storm, inflammatory diseases and cancer.

Acknowledgements: We thank the following for support; The Swedish Children's Diabetes

Foundation, The Swedish Diabetes Foundation, The Swedish Research Council, EXODIAB,

The Ernfors Foundation and The Thurings Foundation. We thank Dr Zhitao Gong for

participation in some of the experiments and Universitetslektor Femke Heindryckx for the gift

of the HepG2 cell line.

Author Contributions: BB, ZJ, HH conceived project; HH, ZJ, AKB, SVK, BB, MKM

designed experiments; HH, ZJ, AKB, SVK, OTR, SK, AIC and FA did experiments; ZJ, HH,

AKB, SVK made figures; all authors analyzed results, BB wrote manuscript that was edited by

ZJ, HH, AKB and then commented on by all authors.

Competing interests: BB has patent applications WO2019/164446A1 and WO2019162403A1

References

Geltink, R. I. K., Kyle, R. L. & Pearce, E. L. Unraveling the Complex Interplay Between T Cell Metabolism and Function. *Annu Rev Immunol* **36**, 461-488 (2018). https://doi.org:10.1146/annurev-immunol-042617-053019

van Niekerk, G., Christowitz, C., Conradie, D. & Engelbrecht, A. M. Insulin as an immunomodulatory hormone. *Cytokine Growth Factor Rev* **52**, 34-44 (2020). https://doi.org:10.1016/j.cytogfr.2019.11.006

Song, W., Li, D., Tao, L., Luo, Q. & Chen, L. Solute carrier transporters: the metabolic gatekeepers of immune cells. *Acta Pharm Sin B* **10**, 61-78 (2020). https://doi.org:10.1016/j.apsb.2019.12.006

Barragan, A., Weidner, J. M., Jin, Z., Korpi, E. R. & Birnir, B. GABAergic signalling in the immune system. *Acta Physiol (Oxf)* **213**, 819-827 (2015). https://doi.org:10.1111/apha.12467

Tian, J. & Kaufman, D. L. The GABA and GABA-Receptor System in Inflammation, Anti-Tumor Immune Responses, and COVID-19. *Biomedicines* **11** (2023). https://doi.org:10.3390/biomedicines11020254

Niswender, K. D. Basal insulin: physiology, pharmacology, and clinical implications. *Postgrad Med* **123**, 17-26 (2011). https://doi.org:10.3810/pgm.2011.07.2300

Macintyre, A. N. *et al.* The glucose transporter Glut1 is selectively essential for CD4 T cell activation and effector function. *Cell Metab* **20**, 61-72 (2014). https://doi.org:10.1016/j.cmet.2014.05.004

Lunt, S. Y. & Vander Heiden, M. G. Aerobic glycolysis: meeting the metabolic requirements of cell proliferation. *Annu Rev Cell Dev Biol* **27**, 441-464 (2011). https://doi.org:10.1146/annurev-cellbio-092910-154237

Zhang, B. *et al.* B cell-derived GABA elicits IL-10(+) macrophages to limit anti-tumour immunity. *Nature* **599**, 471-476 (2021). https://doi.org:10.1038/s41586-021-04082-1

Codo, A. C. *et al.* Elevated Glucose Levels Favor SARS-CoV-2 Infection and Monocyte Response through a HIF-1alpha/Glycolysis-Dependent Axis. *Cell Metab* **32**, 498-499 (2020). https://doi.org:10.1016/j.cmet.2020.07.015

Wan, Q. *et al.* Recruitment of functional GABA(A) receptors to postsynaptic domains by insulin. *Nature* **388**, 686-690 (1997). https://doi.org:10.1038/41792

Xu, E. *et al.* Intra-islet insulin suppresses glucagon release via GABA-GABAA receptor system. *Cell Metab* **3**, 47-58 (2006). https://doi.org:10.1016/j.cmet.2005.11.015

Jin, Z. *et al.* Insulin reduces neuronal excitability by turning on GABA(A) channels that generate tonic current. *Plos One* **6**, e16188 (2011). https://doi.org:10.1371/journal.pone.0016188

Tsai, S. *et al.* Insulin Receptor-Mediated Stimulation Boosts T Cell Immunity during Inflammation and Infection. *Cell Metab* **28**, 922-934 e924 (2018). https://doi.org:10.1016/j.cmet.2018.08.003

Bhandage, A. K. *et al.* GABA Regulates Release of Inflammatory Cytokines From Peripheral Blood Mononuclear Cells and CD4(+) T Cells and Is Immunosuppressive in Type 1 Diabetes. *EBioMedicine* **30**, 283-294 (2018). https://doi.org:10.1016/j.ebiom.2018.03.019

Cahalan, M. D. & Chandy, K. G. The functional network of ion channels in T lymphocytes. *Immunol Rev* **231**, 59-87 (2009). https://doi.org:10.1111/j.1600-065X.2009.00816.x

17 Vaeth, M., Kahlfuss, S. & Feske, S. CRAC Channels and Calcium Signaling in T Cell-
Mediated Immunity. *Trends Immunol* **41**, 878-901 (2020).
https://doi.org:10.1016/j.it.2020.06.012

18 Vaeth, M. *et al.* Store-Operated Ca(2+) Entry Controls Clonal Expansion of T Cells
through Metabolic Reprogramming. *Immunity* **47**, 664-679 e666 (2017).
https://doi.org:10.1016/j.immuni.2017.09.003

19 Pearce, E. L., Poffenberger, M. C., Chang, C. H. & Jones, R. G. Fueling immunity: insights
into metabolism and lymphocyte function. *Science* **342**, 1242454 (2013).
https://doi.org:10.1126/science.1242454

20 Jones, W. & Bianchi, K. Aerobic glycolysis: beyond proliferation. *Front Immunol* **6**, 227
(2015). https://doi.org:10.3389/fimmu.2015.00227

21 Traba, J., Miozzo, P., Akkaya, B., Pierce, S. K. & Akkaya, M. An Optimized Protocol to
Analyze Glycolysis and Mitochondrial Respiration in Lymphocytes. *J Vis Exp* (2016).
https://doi.org:10.3791/54918

22 Hummel, C. S. *et al.* Glucose transport by human renal Na+/D-glucose cotransporters
SGLT1 and SGLT2. *Am J Physiol Cell Physiol* **300**, C14-21 (2011).
https://doi.org:10.1152/ajpcell.00388.2010

23 Michalek, R. D. *et al.* Cutting edge: distinct glycolytic and lipid oxidative metabolic
programs are essential for effector and regulatory CD4+ T cell subsets. *J Immunol* **186**,
3299-3303 (2011). https://doi.org:10.4049/jimmunol.1003613

24 Fujii, S. & Beutler, E. High glucose concentrations partially release hexokinase from
inhibition by glucose 6-phosphate. *Proc Natl Acad Sci U S A* **82**, 1552-1554 (1985).
https://doi.org:10.1073/pnas.82.5.1552

25 Chang, C. H. *et al.* Posttranscriptional control of T cell effector function by aerobic
glycolysis. *Cell* **153**, 1239-1251 (2013). https://doi.org:10.1016/j.cell.2013.05.016

26 Young, S. Z. & Bordey, A. GABA's control of stem and cancer cell proliferation in adult
neural and peripheral niches. *Physiology (Bethesda)* **24**, 171-185 (2009).
https://doi.org:10.1152/physiol.00002.2009

27 Bailey, M. E., Albrecht, B. E., Johnson, K. J. & Darlison, M. G. Genetic linkage and
radiation hybrid mapping of the three human GABA(C) receptor rho subunit genes:
GABRR1, GABRR2 and GABRR3. *Biochim Biophys Acta* **1447**, 307-312 (1999).
https://doi.org:10.1016/s0167-4781(99)00167-0

28 Naffaa, M. M., Hibbs, D. E., Chebib, M. & Hanrahan, J. R. Pharmacological Effect of
GABA Analogues on GABA-varrho2 Receptors and Their Subtype Selectivity. *Life (Basel)*
12 (2022). https://doi.org:10.3390/life12010127

29 Packer, M. *et al.* Cardiovascular and Renal Outcomes with Empagliflozin in Heart
Failure. *N Engl J Med* **383**, 1413-1424 (2020).
https://doi.org:10.1056/NEJMoa2022190

30 Espes, D. *et al.* GABA induces a hormonal counter-regulatory response in subjects with
long-standing type 1 diabetes. *BMJ Open Diabetes Res Care* **9** (2021).
https://doi.org:10.1136/bmjdrc-2021-002442

31 Masoodi, M. *et al.* Disturbed lipid and amino acid metabolisms in COVID-19 patients. *J
Mol Med (Berl)* (2022). https://doi.org:10.1007/s00109-022-02177-4

32 Alshnbari, A. & Idris, I. Can sodium-glucose co-transporter-2 (SGLT-2) inhibitor reduce
the risk of adverse complications due to COVID-19? - Targeting hyperinflammation.
Curr Med Res Opin **38**, 357-364 (2022).
https://doi.org:10.1080/03007995.2022.2027141

Figure 1

327
328
329

Figure 1. GABA effects on metabolic activity and cytokine release are glucose
concentration-dependent in activated CD4$^+$ T cells.
a, Cellular metabolic activity, glucose (mM): 2.8 (N=36), 5.6 (N=54), 10 (N=35) and 16.7 (N=53).
Violin plots where horizontal black and grey lines indicate the median and quartiles.
b, Left panel. Histograms of normalized cellular metabolic activity at low-normal (2.8 and 5.6)
and high (10 and 16.7) glucose concentrations (mM). Solid lines: Gaussian fits, vertical broken
line: 10% inhibition by GABA (500 nM). Right panel. Metabolic activity of cells from donors
where ≥10% GABA inhibition was observed at 5.6 mM glucose; 2.8 (N=28), 5.6 (N=37), 10
(N=27) and 16.7 (N=36).
c, Heatmap and hierarchical clustering of biomolecules by PEA 72 h post-stimulation of cells
(N=17). Data represent the mean log2 (fold change) for samples cultured in presence and
compared to absence of GABA.
d, e, GABA effect on IFNγ **(d)** and IL-10 **(e)** release. ELISA, 72 h post-stimulation, glucose (mM):
2.8 (N=7;7), 5.6 (N=17;17), 10 (N=6;10) and 16.7 (N=16;16).
f, Relative expression of *GABRR2* mRNA in resting and activated cells in 5.6- or 16.7-mM glucose
(N=6) analyzed by $2^{-\Delta\Delta Ct}$ method.
g, Immunofluorescence co-staining of GABRR2 (green) and DAPI (blue), 5.6- or 16.7-mM
glucose. Bar graph; normalized GABRR2 fluorescence intensity (n>50 cells/group, N=3).
Data were normalized to values of activated cells in the absence of drugs at each glucose
concentration (**a, c, e, f, g**). Data represent geometric mean ± SEM in (**f**), mean ± SEM in (**g**). Box
and whisker (**c, e, f**): 10-90 percentiles, means: +, medians: black lines in the boxes. Statistics:
one sample t-test when compared to activated cell group (#, p<0.05; ##, p<0.01; ###, p<0.001),
paired groups by repeated measures ANOVA or fitting a mixed-effects model, or ordinary one-
way ANOVA followed by Tukey (**a, c, e, f**) or Fisher's LSD test (**g**) for multiple comparisons (*,
p<0.05; **, p<0.01; ***, p<0.001). Statistical significance in **c** (p ≤ 0.05 and adjusted p-value q <
0.12) are marked with *. **N**: donors.
356
357
358
359

Figure 2

Figure 2. Insulin enhances GABA-activated GABA$_A$R currents and Ca^{2+} entry in activated CD4$^+$ T cells.

a, GABA-activated whole-cell currents (n=3-8, N=11). Insert: a patch-pipette attached to an activated CD4$^+$ T cell.

b, c, GABA dose-response curves in cells without (b) or treated with (c) insulin. The currents were normalized to 100 nM (b) or 100 pM (c) GABA responses (n=3-11, N=27).

d, Whole-cell currents evoked by GABA (n=11, N=9), muscimol (n=4, N=3) or TACA (n=5, N=4), 5.6 mM glucose.

e, Left panel: Mean Ca^{2+} signal intensity trace (black dots) and SEM (gray whiskers) recorded 72 h after activation of cells (n=14). Colored bars: Perfusion time of drugs (GABA 500 nM and thapsigargin, Tg 1 μM) in 0 or 2 mM Ca^{2+}-containing perfusion medium. Right panel: Relative Ca^{2+} signal (mean ± SD) (n=14, 20, 41, 47, N= 4).

f, GABA-activated relative Ca^{2+} signal (n=58-232, N=3-5) without or treated with insulin.

g, GABA-activated relative Ca^{2+} signal in cells (n=131, N=3) in absence or presence of picrotoxin, with insulin.

h, Representative time-lapsed micrographs of a live Ca^{2+} imaged cell (16.7 mM glucose, with insulin). GABA concentrations were sequentially perfused and inter-spaced with medium applications. Color scale: Relative fluorescence intensity.

i, j, Dose-response relationship for GABA-induced Ca^{2+} signals (n=63-279, N=3-8), see also ~~Figure~~ Extended Data 2e.

k, l, m, n, Effects of GABA$_A$R agonists and antagonists on the relative Ca^{2+} signal: muscimol, TACA, picrotoxin, TPMPA and SOCE antagonist YM48583, 5.5 mM glucose. Box plots; black without and red with insulin, 5.5 mM glucose, (n=191-265, N=3-5).

Curve fitting (**b, c, i, j**): nonlinear regression with sigmoidal polynomial (4PL) model. Data presented as mean $\pm$ SEM (**a, b, c, d, i, j**). Statistics: ordinary one-way ANOVA (**a, d**) or repeated measures ANOVA followed by Tukey multiple comparison test (**k, l, m**) or unpaired (**f**) or paired (**g, n**) student's t-test. ns, $p > 0.05$; **, $p < 0.01$; ***, $p < 0.001$. #, $p < 0.05$; ###, $p < 0.001$ – picrotoxin vs. respective group before picrotoxin application. **n**: cells, **N**: donors.

Figure 3

398
Figure 3. GABA inhibits hexokinase and glycolysis in activated CD4+ T cells.
a, The volcano plot summarizes pathways that are significantly up- (green) or down-regulated
(blue and red (HK1)) by GABA treatment in cells 72 h post activation (N=5), 5.6 mM glucose. The
x-axis represents the average $\log_2$ fold-change of all proteins measured by mass spectrometry
within that pathway and every dot is one pathway. The y-axis represents the adjusted p-value (–
log10 transformation). The horizontal line represents p = 0.05, and the vertical lines represent
average log2 (fold change) <–0.5 or >+0.5.
b, Western blot images and relative expression of HK1 protein (N=7 at 5.6 mM and 3 at 16.7 mM).
c, *In vitro* HK activity analyzed in resting and activated cells, 5.6 mM (N=12) or 16.7 mM (N=6)
glucose. HK activity (µmol NADH/min/ml) was normalized to the activity of activated cells in the
absence of drugs.
d, Representative ECAR over time in cells 72 h post-activation treated with or without GABA (500
nM) or insulin at 5.6 mM (left panel) or 16.7 mM (right panel). Dashed lines indicate injections of
glucose (Glu.), oligomycin (Oligo) or 2-deoxyglucose (2-DG) into media.
e. ECAR results (mpH/min/protein) were normalized to controls in absence of drugs for average
glycolysis, glycolytic capacity and glycolytic reserve in cells 72 h post-activation treated with or
without GABA or insulin, 5.6 mM (N=5) or 16.7 mM (N=5) glucose.
Data are normalized to values of activated cells in the absence of drugs at each glucose
concentration (**b, c, e**). Data represent mean ± SEM (**b, c, d, e**). Statistics by one sample t-test when
compared to activated cell group (#, p<0.05; ##, p<0.01; ###, p<0.001), repeated measures
ANOVA or fitting a mixed-effects model followed by Tukey multiple comparison test. **N**=donor.
420

Figure 4

Figure 4. SGLT2 is expressed, functional and modulates glycolysis and release of IFNγ and IL-10 in activated CD4⁺ T cell.

a, The cellular metabolic activity of resting and activated cells in presence of GLUT1 inhibitor, WZB 117 or SGLTs inhibitor, phlorizin was normalized to that of activated cells without drug treatment at 5.6 mM or 16.7 mM glucose (N=4).

b, Relative gene expression levels of Na⁺-dependent glucose transporters in activated cells at 5.6 mM glucose (N=6) determined by RNAseq analysis. TPM, transcripts per million. ND, not detected.

c, Representative Western blot image of SGLT2 protein in activated CD4⁺ T cells and HEPG2 cell line. The stain-free blot image represents total proteins loaded for each sample, insulin (3 nM), 16.7 mM glucose.

d, Representative immunofluorescence co-staining of SGLT2 (green) and DAPI (blue) in CD4⁺ T cells, 5.6 mM or 16.7 mM glucose.

e, Glucose uptake in absence or presence of phlorizin (N=9) and SGLT2 selective inhibitor, empagliflozin (N=6) in cells 72 h post-activation, insulin, 5.6 mM or 16.7 mM glucose.

f, ECAR results (mpH/min/protein) normalized to untreated cells for average glycolysis, glycolytic capacity and glycolytic reserve in cells, 72 h post-activation, incubated with phlorizin, with or without insulin in 5.6 mM (N=5) or 16.7 mM (N=5) glucose.

g, h, IFNγ **(g)** and IL-10 **(h)** release from activated cells 72 h post-activation in with or without GABA, phlorizin or insulin, at 5.6 mM or 16.7 mM glucose (N=17).

443   Data represent mean ± SEM. Data are normalized to values of activated cells in the absence of
444   drugs at each glucose concentration (**a, e, f, g, h**). Statistics by one sample t-test when compared
445   to activated cell group (#, $p < 0.05$; ##, $p < 0.01$; ###, $p < 0.001$), repeated measures ANOVA or
446   fitting a mixed-effects model followed by Tukey multiple comparisons test, *, $p < 0.05$. **N**=donor.

Extended Data

Extended Data Fig. 1: Schematic diagram of the experimental design.
Created with BioRender.com.

1
2

3
4
Extended Data Fig. 2: The effect of glucose on cellular metabolic activity and release of
IFNγ and IL-10 from CD4$^+$ T cells.
a, Cellular metabolic activity from donors where <10% GABA inhibition at 5.6 mM glucose was
observed. **b,** Cellular metabolic activity of CD4$^+$ T cells 72 h after activation in the presence of
different glucose concentrations. **c, d,** IFNγ (**c**) and IL-10 (**d**) released from CD4$^+$ T cells 72 h
after activation in presence of different glucose concentrations. Data are normalized to values
at 5.6 mM glucose and presented as mean ± SEM. Statistics by one sample t-test when compared
to 5.6 mM glucose group (#, $p<0.05$; ##, $p<0.01$), repeated measures ANOVA or fitting a mixed-
effects model followed by Tukey test for multiple comparisons. N = number of donors.

Extended Data Fig. 3: GABA-activated current and Ca²⁺ signaling in CD4⁺ T cells.
a, A representative current-voltage relationship recorded in an activated CD4⁺ T cell in response to 500 nM GABA application by perforated patch-clamp, 5.6 mM glucose. **b,** GABA (500 nM)-activated relative Ca²⁺ signals (left panel) and mean intensity (right panel) recorded from activated CD4⁺ T cells (n=128 cells/ N=3 donors) in sequentially applied 5.6 mM and 16.7 mM glucose-containing media. Cells were activated in 5.5 mM glucose-containing media for 72 h. **c,** The ratio of Ca²⁺ signals in the absence and presence of calcium-ionophore calcimycin (1 M) in CD4⁺ T cells (n=85 cells/ N=4 donors) 72 h after activation with or without insulin treatment. **d,** GABA-activated relative Ca²⁺ signal recorded in CD4⁺ T cells (n=131 cells/ N=3 donors) in the absence or presence of GABA_B receptor antagonist CGP 52432 (1 μM) at 5.5 mM glucose. **e,** Relative Ca²⁺ signals induced by different concentrations of GABA in activated CD4⁺ T cells treated without or with insulin in 5.5 mM or 16.7 mM glucose (n=63-279 cells/ N=3-8 donors). Statistics by two-tailed student's t-test. ns, p > 0.05, **, p<0.01; ***, p<0.001.

34
35
36
37

38
39
Extended Data Fig. 4: GABA, glucose and insulin regulate CD4+ T cell effector functions.
In human CD4+ T cells, the activation of T cell receptor (TCR)/CD3 complex by anti-CD3
antibodies binding, initiates store-operated Ca2+ entry (SOCE) through Ca2+ release-activated
Ca2+ (CRAC) channels. It regulates intracellular Ca2+ signals, drives changes in transcription
factors, upregulates glycolysis and increases T cell effector functions. Physiological concentration
of GABA (pM-nM) activates GABAA receptors (GABAAR) to modulate T cells metabolic activity
and levels of cytokines (e.g., IFNγ and IL-10) by 1) enhancing function of CRAC channels and Ca2+
entry; 2) reducing expression of Nuclear Factor of Activated T cell 2 (NFATC2 or NFAT1); 3)
decreasing expression and activity of hexokinase (HK), the 1st rate-limiting enzyme in aerobic
glycolysis. In addition, insulin-insulin receptor (IR) signaling promotes glycolysis but
paradoxically reduce glycolysis by enhancing GABAAR function. The fluctuation of blood glucose
concentration affects glycolysis and cytokine production. Glucose uptake is mediated by not only
the facilitated glucose transporters GLUT1/3 but also the sodium-dependent glucose transporter
2 (SGLT2). Phlorizin, a SGLT inhibitor, reduces glucose uptake, attenuates glycolysis and reduces
IFNγ and IL-10 levels.
Created with BioRender.com.
56

63 **Extended Data Table 1 | Donors' demographic information**

Parameter	Normal Donors (N=160)
Age (years)	26.2 ± 0.25
Sex (N, %)	89 males (55.6%)
	71 females (44.4%)

64
65 Data are presented as means ± SEM.

Extended Table 2 I Relative gene quantification (normalized to activated cells) by qPCR (N=7-11 donors).

Gene name	5.6 mM glucose					16.7 mM glucose				
	Resting	Activated	Activated+Insulin	GABA	GABA+Insulin	Resting	Activated	Activated+Insulin	GABA	GABA+Insulin
NFATC1	0.088	1***	1.129***	1.086**	1.140***	1.902	1**	1.017	1.072*	0.923**
NFATC2	0.996	1	0.980	0.884##	0.942	1.333	1	0.961&	1.121	0.876*, &
NFATC3	0.512	1**	0.929**	0.836*	0.773**	0.484	1*	0.895*, &	1.057**	0.922*
JAK1	1.808	1***	1.156	1.096***	1.144***, #	2.123	1	1.049	1.329	1.027
JAK2	0.405	1***	1.060***	1.085***	1.133***	0.291	1	0.802	1.041	0.932
TYK2	1.639#	1	1.278	1.016	0.997*	1.254	1	0.716	0.826	0.596
HIF1A	0.310	1***	1.005***	0.961***	0.960***	0,287	1***	0.919***	0.968***	0.794***
STAT1	0.088	1***	1.129***, #	1.086***	1.140***	0.067	1***	0.939***	0.925***	0.981***
MYC	0.626	1**	0.925**	0.918*	0.946**	0.368	1**	0.662#	0.805*	0.787*

*p<0.05, **p<0.01, ***p<0.001, as compared to resting cells; #p<0.05, ##p<0.01, as compared to activated cells: &p<0.05, as compared to GABA-treated cells. The statistical analysis was performed on ΔCq values with repeated measures one-way ANOVA with Fisher's LSD test for multiple comparison.

2 **Extended Data Table 3 │Reactome analysis of MS data (N=5 at 5.6 mM glucose). Proteins are**
3 **named after UniProtKB.**
4 **Top up-regulated proteins**

Identifier	logFC	AveExpr	t	P.Value
K2C79_HUMAN	0.67266061	13.2926208	2.34394122	0.04308325
CHM4A_HUMAN	0.55827697	13.7149206	2.20925349	0.05380739
HLAH_HUMAN	1.03346983	14.8113954	2.03947592	0.07106636
RAB2A_HUMAN	1.37200668	14.1750267	2.03797883	0.07123998
SC11A_HUMAN	5.70132617	10.587487	1.91122932	0.08749562
K22O_HUMAN	0.48972535	12.2546633	1.88242372	0.09164997
DYL1_HUMAN	5.54957599	10.7818438	1.81112175	0.10274055
PLXA4_HUMAN	0.69449352	13.706242	1.7195177	0.11881622
RL36_HUMAN	6.02123532	12.32297	1.68555819	0.12534011
IF4G2_HUMAN	0.82748497	13.4787932	1.68221061	0.12600056

5
6 **Top down-regulated proteins**

Identifier	logFC	AveExpr	t	P.Value
TBB1_HUMAN	-7.55965074	8.58464831	-2.74841088	0.02204275
STA5B_HUMAN	-7.60692039	6.04448902	-2.44305026	0.03656351
EXOS3_HUMAN	-6.6014009	5.67159622	-2.16624065	0.05775108
BGAL_HUMAN	-6.67048214	5.77846898	-2.13711593	0.06057899
AL1B1_HUMAN	-6.67577438	5.87276979	-2.08793829	0.06566002
HXK1_HUMAN	-0.73989819	14.3697968	-2.06803575	0.06783079
HLAG_HUMAN	-6.34860408	5.64345434	-2.03980339	0.07102844
COPG1_HUMAN	-5.15586276	10.0461233	-1.80136877	0.10435116
BCL2_HUMAN	-4.91388319	9.86964223	-1.73134501	0.11661798
PIP_HUMAN	-4.68190741	9.67505331	-1.66669489	0.12910318

Methods

Study individuals and collection of samples

Human blood buffy coat and blood samples were obtained from Uppsala University Hospital (Uppsala Akademiska sjukhuset), protocol approved by the Regional Research Ethical Committee in Uppsala. All donors were voluntary recruited and informed consent were signed. In total, samples from 160 normal donors, 89 men and 71 women with mean age 26.2 ± 0.25 years were included in the study.

Chemicals and reagents

Drugs, buffers and salts unless otherwise specified were purchased from Sigma-Aldrich/Merck (Steinheim, Germany). TACA, muscimol, TPMPA, YM-58483, WZB 117 were purchased from Tocris (UK, 7A/109031). Phlorizin and empagliflozin were obtained from AdooQ Bioscience. Urea and NH_4HCO_3 were from Acros Organics, Geel, Belgium and trypsin and dimethyl sulfoxide (DMSO) from Thermo Scientific (Waltham, MA). Tris(hydroxymethyl) aminomethane hydrochloride (Tris), LC-MS grade water, acetonitrile and formic acid were obtained from Fisher Scientific (Geel, Belgium).

CD4$^+$ T cells isolation and activation

Peripheral blood mononuclear cells (PBMCs) were isolated by density-gradient centrifugation using Ficoll-Paque™ PLUS (Sigma Aldrich, Sweden). CD4$^+$ T cells were then purified from the isolated PBMCs by negative selection using humanCD4$^+$ T cell isolation kits (Miltenyi Biotec, Germany) according to the manufacturer's instructions. The freshly isolated CD4$^+$ T cells ($1x\ 10^6$/ml) were suspended in RPMI 1640 glucose-free medium (Gibco, Fisher Scientific, Sweden) supplemented with 2 mM glutamine, 10% heat-inactivated dialyzed fetal bovine serum, 100 U/ml penicillin, 10 mg/ml streptomycin, and 5 µM β-mercaptoethanol (Gibco, Fisher Scientific, Sweden). The activation of the cells was performed in either 96-well (10^5 cell/well) or 24-well (10^6 cell/well) plates based on the purpose of the experiment. CD4$^+$ T cells were activated with 3 µg/ml plate-bound anti-CD3 (BD Biosciences, USA, 555329) in the presence of different concentrations of glucose (2.8 mM, 5.6 mM, 10 mM and 16.7 mM) for 66–72 h and indicated drugs were added based on the experimental design.

MTT Assay

The metabolic activity of CD4$^+$ T cells was assessed in 96-well plates with colorimetric MTT (3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide) assay. After 66–72 h of activation, a PBS-soluble tetrazolium dye MTT (final concentration 1mM) was added and incubated for additional 4 h at 37 °C. Thereafter, the cells were centrifuged at 2000 rpm for 10 min to concentrate the insoluble purple formazan pellets. The supernatants were collected and stored at −80 °C for later biomolecules analysis. The formazan crystal pellets were dissolved in DMSO and the plate was read 10 min later using Multiskan FC Microplate Photometer (Thermo Fisher, USA) at 550 nm.

Multiplex Proximity Extension Assay (PEA)

The supernatants collected at the end of the MTT assay were sent on-ice to the SciLifeLab Affinity Proteomics Unit in Uppsala, Sweden, for analysis. Inflammatory proteins were measured using high-throughput multiplex PEA with an Olink Target 96 Inflammatory panel (OlinkProteomics, Uppsala, Sweden). PEA measures simultaneously 92 proteins/biomolecules in different sample matrices where it

combines a dual-detection immunoassay and quantitative PCR. Normalized protein expression (NPX) values, an arbitrary unit on the log2 scale, were used to present the data. A protein expression (NPX value) is considered acceptable if the value is above the limit of detection (LOD) for each particular protein. Proteins that were detected in more than 75% of the samples were analyzed in this study: 59 out of 92 proteins had detectable levels. Data for resting, activated and GABA-treated activated CD4$^+$ cells supernatants were analyzed separately. The biomolecules in the supernatants were assessed for each donor at each glucose concentration. The data was computed, processed and analyzed with support from NBIS (National Bioinformatics Infrastructure Sweden). The log2FC (FC =fold change) is the average increase in NPX in GABA as compared to activated alone, this value is both estimated based on the linear mixed effects model and computed as the mean over the GABA-activated difference and plotted in heatmaps. Paired t-test was used followed by multiple testing correction using false discovery rate (FDR) method (Two-stage step-up method of Benjamini, Krieger and Yekutieli) with a threshold 12%.

Determination of IFNγ and IL-10 concentrations in supernatants

The concentrations of IFNγ and IL-10 were measured with enzyme-linked immunosorbent assay (ELISA). The supernatants were diluted when necessary to be in the optimal linear optical range and analyzed using commercial kits for IFNγ (Human IFN-gamma DuoSet ELISA, R&D Systems, USA and Bio legend, UK), and IL-10 (Biolegend, UK) according to manufacturers´ instructions. Optical densities were measured using FC Microplate Photometer (Thermo Fisher, USA) at 450 nm, and 540 nm was used as a correction wavelength. The levels of IFNγ or IL-10 were normalized to controls (activated CD4$^+$ T cells in the absence of drugs).

Total RNA and protein extraction and, bulk RNA sequencing

Total RNA and protein were extracted from activated CD4$^+$ T cells, in the absence or presence of 500 nM GABA, using RNA/Protein Purification Plus Kit (Norgen Biotek, Ontario, Canada). During the extraction process, total RNA was treated with RNase-Free DNase I Kit (Norgen Biotek, Ontario, Canada) to eliminate contamination of genomic DNA. The extracted proteins were quantified using DC protein assay kit (Bio-Rad, USA), and immediately frozen at −80°C. Total RNA samples were submitted to the GENEWIZ NGS facility (Germany) and quality control analysis showed that all RNA samples had an RNA integrity number (RIN) >9 (9.48 ± 0.1, mean ± SEM). Strand-specific RNAseq cDNA libraries with polyA selection were generated and sequenced using Illumina HiSeq sequencing platform (12 samples, 20.8-31.1 million reads, 150 bp paired end per sample). Sequence reads were filtered and trimmed using Trimmomatic V.0.36 to remove low quality reads. The trimmed reads were further mapped and aligned to the Homo Sapiens GRCh38 reference genome (ENSEMBL) using the STAR aligner v.2.5.2b. Unique gene hit counts were calculated using feature Counts from the Subread package v.1.5.2.

Real-Time Quantitative Reverse Transcription (qPCR)

The cDNA was synthesized using SuperScript reverse transcriptase IV (Invitrogen Thermo Scientific, USA) in a 20 μl reaction following the standard protocol from manufacturer. The qPCR reaction was then performed using qPCRBIO SYBR or Probe Mix low-ROX (PCR Biosystems, UK; PB20.11-05), and gene-specific primers (Table S1) or Taqman probes (*GABRR2* Hs00266703_m1; *IPO8* Hs00914057_m1, ThermoFisher Scientific, USA). Changes in the abundance of each transcript were normalized to the expression of reference genes (*IPO8*). Relative expression levels

were calculated by normalizing to non-treated activated CD4$^+$ T cells obtained from
the same donor at the same glucose concentration using $2^{-\Delta\Delta Ct}$ method. All samples
were run in duplicate. The real-time qPCR amplification was performed in 10 µl
reaction mix in a 384-well plate usings a QuantStudio™ 5 instrument (ThermoFisher
Scientific). All runs started with an initial denaturation step of 2 min at 95 °C, followed
by 45 cycles of 95 °C for 5 s and 60 °C for 30 s.
118
119

Mass spectrometry (MS)-based bottom-up proteomics
The lysates were purified and tryptic digested for MS-based bottom-up proteomic
analysis using Filter Aided Sample Preparation (FASP)[1, 2]. Briefly, 20 µg of total protein
were transferred onto centrifugal filter units (Microcon-30 kDa; Merck, Darmstadt,
Germany) and washed with a buffer containing 8 M urea and 100 mM Tris (pH 8.5).
The washing step was repeated after reduction with 8 mM dithiothreitol (DTT),
alkylation with 50 mM iodoacetamide (IAA) and removal excess of IAA with 8 mM DTT.
Before tryptic digestion (enzyme-to-protein ratio 1:50 (w/w)) the filter was washed
three times with NH$_4$HCO$_3$. After incubation in a wet chamber at 37 °C for 16 h the
resulting peptides were washed twice from the filter by adding 50 mM NH$_4$HCO$_3$.
Trifluoroacetic acid was then added to a final concentration of 1% (v/v) and the
samples were dried at 45 °C. Finally, the samples were reconstituted in 3% acetonitrile
and 0.1% formic acid in water to a final concentration of 150 ng protein/µl. The tryptic
peptides were analyzed on a nanoAcquity UPLC system equipped with a C18, 5 µm,
180 µm x 20 mm trap column and an HSS-T3 C18 1.8 µm, 75 µm x 250 mm analytical
column (Waters Corporation, Manchester, UK) coupled to a Synapt G2 Si HDMS mass
spectrometer with an electrospray ionization source (Waters Corporation, Manchester,
UK). The UDSM^E approach with positive ionization was used[1, 2]. Mobile phase A
contained 3 % DMSO and 0.1 % formic acid in water and mobile phase B 3 % DMSO
and 0.1 % formic acid in acetonitrile. After loading 300 ng of protein on the column in
trapping mode, peptide separation was performed with a gradient run from 3 - 40%
(v/v) mobile phase B over 120 min. The flow rate was set to 0.3 µl/min and the column
oven was set to 40 °C. Method performance was controlled with a commercially
available HeLa digest (Thermo Scientific, Waltham, MA). UPLC-MS data processing
and label-free quantification: ProteinLynx Global Server (version 3.0.3, Waters
Corporation, Milford, MA) was used for raw data processing. With FDR of 10%, a
database search against a randomized UniProt human database (Uni-ProtKB version
14/01/2020) was performed. Carbamidomethyl cysteine was set as fixed modification,
acetyl lysine, C-terminal amidation, asparagine deamidation, glutamine deamidation
and methionine oxidation as variable modifications and trypsin as digest reagent.
Minimum peptide matches per protein were 2 and minimum fragment ion matches per
peptide and protein were 1 and 3, respectively. For label-free quantification analysis,
the identified proteins were further processed using ISOQuant 1.8 as described
elsewhere[2]. Following the TOP3 quantification approach the average intensity of the
three most intense peptides of each protein were used for relative protein
quantification. ISOQuant settings are provided in Table S2.
156

Immunoblotting
158 CD4$^+$ T cells were lysed in RIPA buffer (Sigma-Aldrich, R0278) containing protease
159 inhibitor cocktail (Roche, 11 836 153 001) and phosphatase inhibitors (Sodium
160 orthovanadate, Sigma-Aldrich, S6508). Proteins were heated at 100 °C for 5 min in
161 SDS loading buffer (275 mM Tris. HCL, 9% SDS, 50% glycerol, 0.03% bromophenol

blue) containing 9% beta-mercaptoethanol and separated by Mini-Protean TGX gel 4 - 15% (BioRad, stain-free gels #4568084, 4561035) and blotted onto Amersham Hybond-P PVDF membrane. Membranes were blocked with blocking buffer (EveryBlot BioRad, 12010020) for 20 min at room temperature (RT) and incubated overnight at 4 °C with the following primary antibodies: HK1 (GenTex, GTX105248, 1:3000), SGLT2 (ProteinTech, 24654-1-AP, 1:500) and GABRR2 (Invitrogen, PA5-41008, 1:1000). After immunoblotting, membranes were washed in TBST and incubated with Peroxidase-conjugated AffiniPure secondary antibodies (Jackson ImmunoResearch, 111-035-045, 1:5000). Protein detection was done using chemiluminescent ECL reagent (Thermo scientific, 32209). Quantitative densiometric analysis of the immunoblotted bands was performed using Image Labs software (BioRad Laboratories, V6.0.0) and data were normalized for each sample using stain-free technique.

Immunofluorescence staining

CD4$^+$ T cells were harvested after 72 h of anti-CD3 activation on cytoslides (Thermo scientific, 5991056) using a Shandon cytospin centrifuge (400 rpm for 4 min at RT). After drying, cells were fixed with 4% PFA in PBS for 15 min and washed 3 times, followed by membrane permeabilization with 0.1% Triton-X for 5 min. Un-specific binding was blocked with 5% bovine serum albumin and 10% donkey serum (Jackson ImmunoResearch, 017-000121) in 1X PBS for 60 min. Cells were then washed and incubated with primary antibodies diluted in blocking solution overnight at 4 °C followed by incubation with an AlexaFluor 488-conjugated AffiniPure Donkey Anti-Rabbit IgG secondary antibody (Jackson ImmunoResearch, 711-545-152, 1:500) for 1h at RT. Finally, cells were washed again and counterstained with DAPI for 5 min and slides were mounted in ProLongTM Gold antifade reagent (Invitrogen). Images were acquired using confocal microscopy (LSM700, Zeiss) with 63X objective. For statistics, ≥ 50 cells from 3 donors were quantified using ImageJ (NIH) software. Primary antibodies and concentrations included rabbit anti-insulin receptor, InsR (Invitrogen, PA5-29286, 1:800), rabbit anti-GABRR2 (Invitrogen, PA5-41008, 1:600), and rabbit anti-SGLT2 (ProteinTech, 24654-1-AP, 1:100).

In vitro **Hexokinase activity assay**

Hexokinase activity was measured at RT indirectly as a readout of NADH production, a by-product of the reaction converting glucose-6-phosphate to 6 phosphogluconolacton by the enzyme glucose-6-phosphate dehydrogenase (GPDH). The cells were harvested, centrifugated and washed twice with PBS. Washed cells were resuspended in 1 ml assay buffer solution, on ice, included in the hexokinase enzymatic assay kit (Bio Vision, K789-100). After centrifugation at 4 °C for 5 min, the resulting supernatant was used as the final cell extract for the enzymatic assay.

Electrophysiological recordings

The electrophysiological recordings from the isolated CD4$^+$ T cells were done using the perforated whole-cell patch-clamp configuration. In order to keep the intracellular chloride concentration intact, gramicidin was used to obtain the perforated configuration. Gramicidin was diluted in DMSO at the stock concentration 2 mg/ml and the final concentration of the gramicidin in the pipette solution was 2.6 µg/ml. The pipette solution with the gramicidin was protected from light, kept cold and used within 2 hours after the preparation. After making the cell-attached configuration, the access resistance was monitored until characteristic slow and low-amplitude capacitance

transients were observed and the access resistance decreased from GΩ range to the
range of tens of MΩ. The extracellular solution (in mM) contained: 157 NaCl, 4.5 KCl,
0.5 $CaCl_2$, 1 $MgCl_2$, 5 HEPES and 5.6 or 16.7 glucose (pH 7.4, adjusted with NaOH).
The pipette solution consisted of (mM): 149 KCl, 2 $CaCl_2$, 1 $MgCl_2$, 1 NaCl, 10 HEPES,
5.6 or 16.7 glucose (pH 7.3 adjusted with KOH). The patch-clamp pipettes were made
from borosilicate glass and had a resistance of 12–14 MΩ when filled with the pipette
solution and immersed in the extracellular solution. Recordings were done using
Axopatch 200B amplifier, filtered at 2 kHz and digitized on-line at 10 kHz using an
analog-to-digital converter. To record the electrophysiological data Clampex 10.5
(Molecular Devices, San Jose, CA, USA) software was used. Currents were recorded
at +30 mV or by a ramp protocol, where the potential was changed from −80 to +80
mV in 1 s, the holding potential (Vh) was kept −40 mV. The single sweep duration was
20 s with a break between sweeps 5 s. After application of a drug, the chamber was
perfused 2-6 min with extracellular solution only before next application of a drug.
Currents in the absence of drugs or in the presence of picrotoxin were subtracted from
the test-drug current response.
228
Live cell time-lapse real-time Ca^{2+} imaging
Cells were loaded with a calcium indicator, 3 µM Fluo-8H AM (AAT Bioquest,
Pleasanton, CA) for 15 min at 37 °C and seeded on 5% 3-aminopropyltriethoxysilane-
coated coverslip for 15 min at 37 °C. Time lapse images were acquired by confocal
microscopy (LSM700, Zeiss) with 40X objective at an interval of 1-1.5 second per
image. Drugs were diluted in RPMI 1640 without phenol red and perfused at indicated
time and concentration by a peristaltic pump. After application of each drug, cells were
perfused with the extracellular solution (RPMI 1640) before next application. Area for
single cells was marked and absolute fluorescence intensity values (F) were extracted
using ZENBlue software. Relative intensity (F/Fmedian) for each cell was calculated
for every recorded image and plotted against time. Further, maximum relative intensity
of the cells under drug application time is considered as drug's response. Data is
represented for all cells and as box-whiskers plot.
242
Metabolism Assay
Isolated $CD4^+$ T cells were plated in a XF^e96 cell plates that were coated with Cell-
Tak (#354240, Corning, New York, USA) according to manufacturer's protocol. The
metabolic function of activated $CD4^+$ T cells cultured for 3 days *in vitro* was analyzed
by measuring the extracellular acidification rate (ECAR) using an XF^e96 extracellular
flux analyzer (Seahorse Bioscience). The cells were kept in XF media (Seahorse
Bioscience) supplemented with 5 mM glucose (Sigma Aldrich), and subjected to
glycolysis stress-test protocol using sequential injection of 25 mM glucose, 2 µM
oligomycin, and finally 15 mM 2-deoxyglucose (2DG). Measurements for each
experiment were conducted at least in triplicates and glycolysis, glycolytic capacity,
and glycolytic reserve calculated and values normalized to protein content [3, 4].
254
Glucose uptake assay
2x 10^6 activated $CD4^+$ T cells (72 h activation) with insulin treatment from 48 h after
activation were pelleted, resuspended in glucose-free RPMI-1640 media, incubated
for 10 min in the incubator at 37 °C, and washed once with the same media. Glucose-
free media (500 µl) containing 1µCi/ml $[^{14}C]$ glucose (Perkin Elmer) were added to
cells in the absence or presence of phlorizin (100 µM) or empagliflozin (0.5 µM). Cells
were incubated for 40 min at 37 °C, pelleted by centrifugation at 300x g for 5 min and

washed three times in cold PBS. Cells were then lysed in RIPA buffer and ^{14}C content
was measured with Gold XR scintillation fluid using a liquid scintillation counter (Perkin
Elmer). The glucose uptake was calculated by normalizing counts to protein contents
for each sample.
266
Statistical analysis
Statistical analysis was performed using GraphPad Prism 9 GraphPad Prism
(GraphPad Software Inc., La Jolla, CA, Version 9.3.1). Normalized PEA data were
assessed by two-tailed paired t-test followed by Benjamini-Hochberg's FDR (False
Discovery Rate) method. The significance level is set to 10% FDR. Cellular metabolic
activity, cytokine levels, mRNA, protein expression and ECAR of CD4$^+$ T cells in the
presence of indicated drugs were normalized to controls (activated cells without drug
treatment) from each donor and for each glucose concentration. Normalized data were
analyzed by one-sample t-test when compared to controls. A comparison between
paired groups was performed using repeated measures or mixed-model effects one-
way ANOVA followed by Tukey or Fisher's LSD multiple comparisons tests. Unpaired
multiple-group comparison was performed using ordinary one-way ANOVA followed
by Tukey test. Changes in protein levels in MS data were analyzed using Quantitative,
multi-dataset Pathway Analysis (ReactomeGSA,
https://reactome.org/PathwayBrowser/#TOOL=AT)[5] and volcano plot was
subsequently generated. Error bars in the bar graphs represent SEM unless otherwise
specified and the statistical significance level was set to 0.05.
284
285 **Additional references**

1. Rofo, F. *et al.* Wide-Ranging Effects on the Brain Proteome in a Transgenic Mouse Model of Alzheimer's Disease Following Treatment with a Brain-Targeting Somatostatin Peptide. *ACS Chem Neurosci* **12**, 2529-2541 (2021).
2. Distler, U., Kuharev, J., Navarro, P. & Tenzer, S. Label-free quantification in ion mobility-enhanced data-independent acquisition proteomics. *Nat Protoc* **11**, 795-812 (2016).
3. Pelletier, M., Billingham, L.K., Ramaswamy, M. & Siegel, R.M. Extracellular flux analysis to monitor glycolytic rates and mitochondrial oxygen consumption. *Methods Enzymol* **542**, 125-149 (2014).
4. Traba, J., Miozzo, P., Akkaya, B., Pierce, S.K. & Akkaya, M. An Optimized Protocol to Analyze Glycolysis and Mitochondrial Respiration in Lymphocytes. *J Vis Exp* (2016).
5. ReactomeGSA - Efficient Multi-Omics Comparative Pathway AnalysisJohannes Griss, Guilherme Viteri, Konstantinos Sidiropoulos, Vy Nguyen, Antonio Fabregat, Henning Hermjakob. bioRxiv 2020.04.16.044958;
   doi: https://doi.org/10.1101/2020.04.16.044958