<u>Title</u>: Rab11 suppresses neuronal stress signaling by localizing Dual leucine zipper kinase to

axon terminals for protein turnover.

Seungmi Kim, Yaw Quagraine, Monika Singh, Jung Hwan Kim*

Department of Biology, University of Nevada, Reno, Reno, NV 89557

*Correspondence: jungkim@unr.edu

<u>Summary</u>

Dual Leucine Zipper Kinase (DLK) mediates multiple neuronal stress responses, and its

expression levels are constantly suppressed to prevent excessive stress signaling. We found

that Wallenda (Wnd), the *Drosophila* ortholog of DLK, is highly enriched in the axon terminals of

Drosophila sensory neurons in vivo and that this subcellular localization is necessary for

Highwire-mediated Wnd protein turnover under normal conditions. Our structure-function

analysis found that Wnd palmitoylation is essential for its axon terminal localization.

Palmitoylation-defective Wnd accumulated in neuronal cell bodies, exhibited dramatically

increased protein expression levels, and triggered excessive neuronal stress responses.

Defective intracellular transport is implicated in neurodegenerative conditions. Comprehensive

dominant-negative Rab protein screening identified Rab11 as an essential factor for Wnd

localization in axon terminals. Consequently, *Rab11* loss-of-function increased the protein levels

of Wnd and induced neuronal stress responses. Inhibiting Wnd activity significantly ameliorated

neuronal loss and c-Jun N-terminal kinase signaling triggered by *Rab11* loss-of-function. Taken

together, these suggest that DLK proteins are constantly transported to axon terminals for

protein turnover and a failure of such transport can lead to neuronal loss. Our study

demonstrates how subcellular protein localization is coupled to protein turnover for neuronal

stress signaling.

<u>Keywords</u>

<u>Highlights</u>

- Wnd is highly enriched in axon terminals.

- Wnd protein turnover by Hiw is restricted in the axon terminals.

- Protein palmitoylation of Wnd and Rab11 activity is essential for Wnd axonal localization.

- *Rab11* mutations and defective Wnd palmitoylation impair Wnd protein turnover leading

to increased Wnd protein levels and neuronal loss.

- Inhibiting Wnd activity mitigates neuronal stress response caused by *Rab11* loss-of-

function.

 Introduction

A typical neuron is composed of distinct somatodendritic and axonal compartments that express

unique compositions of membrane and cytoplasmic proteins, which are precisely established

and maintained through a complex network of intracellular transport processes (Guedes-Dias

and Holzbaur, 2019; Guillaud et al., 2020). Consistently, impaired neuronal intracellular

transport has been recognized as an early cellular event in various neurodegenerative disorders

including amyotrophic lateral sclerosis (Collard et al., 1995; De Vos and Hafezparast, 2017),

Alzheimer's (Calkins et al., 2011; Wang et al., 2015), Parkinson's (Saha et al., 2004; Chu et al.,

2012), and Huntington's diseases (Lee et al., 2004; Reddy and Shirendeb, 2012; Smith et al.,

2014). The intricate regulation of intracellular transport relies on a family of small G proteins

known as Rab (Ras-related in brain) proteins (Stenmark, 2009). Studies have demonstrated

altered Rab activity or expression levels in Huntington's disease model (Li et al., 2009b, 2009a),

Alzheimer's disease model (Pensalfini et al., 2020) and from the amyotrophic lateral sclerosis

patients (Mitra et al., 2019). However, how defective neuronal intracellular transport leads to

neuronal loss is not fully understood.

Dual Leucine Zipper Kinase (DLK), also known as MAP3K12 is a cytoplasmic protein kinase

essential for stress signaling and mediates diverse neuronal responses including axon

regeneration, axon degeneration, and cell death (Miller et al., 2009; Xiong et al., 2010; Shin et

al., 2012; Xiong and Collins, 2012; Watkins et al., 2013; Li et al., 2021). Under normal

conditions, neurons tightly control DLK activity to prevent excessive stress signaling primarily

through protein turnover by an evolutionarily conserved PHR1/RPM-1/Highwire (Hiw) ubiquitin

ligases (Nakata et al., 2005; Collins et al., 2006; Xiong et al., 2010; Huntwork-Rodriguez et al.,

2013). Notably, recent studies have shown potential widespread involvement of DLK in neuronal

death observed in neurotrophic factor deprivation, glaucoma, amyotrophic lateral sclerosis, and

Alzheimer's disease models (Pozniak et al., 2013; Welsbie et al., 2013, 2017; Larhammar et al.,

2017a; Le Pichon et al., 2017; Patel et al., 2017). These observations suggest the possibility of

dysregulated or elevated DLK protein/activity levels in these conditions. Interestingly, DLK

proteins are found within axonal compartments (Xiong et al., 2010; Klinedinst et al., 2013;

Holland et al., 2016; Niu et al., 2020, 2022). However, there is limited understanding of the

mechanisms underlying DLK protein localization in axonal compartments and how such axonal

localization of DLK plays a role in neuronal stress responses.

Here, we report a novel coupling between axon terminal localization and protein turnover of

DLK. We further found that Rab11 is essential for this coupling, which when disrupted, leads to

neuronal loss by excessive stress signaling. We found that Wallenda (Wnd), *Drosophila* DLK is

highly enriched in the axon terminals of *Drosophila* sensory neurons in vivo. Furthermore, we

found that Wnd protein turnover by Hiw is confined to axon terminals. We identified a protein

palmitoylation on Wnd as an essential motif for Wnd protein localization in axon terminals. By

taking advantage of this unique behavior of *Drosophila* DLK, Wnd in its reliance to protein

palmitoylation for axonal localization, we demonstrated that Wnd protein turnover is greatly

diminished when Wnd proteins are not transported to axon terminals, which in turn causes

excessive Wnd signaling and neuronal cell death. We further uncovered the selective role of

Rab11 in Wnd axonal localization and protein turnover. Importantly, neuronal stress responses

caused by *Rab11* loss-of-function were significantly mitigated by inhibiting Wnd. Together, our

findings suggest that Rab11 suppresses DLK protein levels by coupling axonal localization and

protein turnover of DLK.

Result

Wnd is highly enriched in axon terminals and Wnd protein turnover is restricted to axon

terminals.

DLK proteins are found in the axonal compartments of diverse animals (Holland et al., 2016; Niu

et al., 2020, 2022). To understand the mechanism of DLK axonal localization, we first set out to

investigate the extent of axonal localization of *Drosophila* DLK, Wallenda (Wnd). We took

advantage of a *wnd* protein-trap allele, *Mi{MIC}wnd^{GFSTF}*, which allows us to visualize

endogenous Wnd protein localization via an EGFP tag (Venken et al., 2011; Li et al., 2017).

The levels of Wnd-GFSTF proteins in the third instar larva stage were extremely low (**Figure**

1A, left). To enhance detection of Wnd-GFSTF, the experiment was performed in *highwire* (*hiw*)

mutants, *hiw^{ΔN}* (Wu et al., 2005). Hiw is a neuronal ubiquitin ligase that selectively targets Wnd

for protein degradation (Collins et al., 2006). Immunostaining was conducted using an anti-GFP

antibody, and the larval samples were examined using a confocal microscope. The result

showed a strikingly polarized expression pattern of Wnd. While Wnd-GFSTF expression was

evident in the neuropil region in the ventral nerve cord (VNC) (**Figure 1A and 1B**), we could not

detect any meaningful signal from neuronal cell bodies, especially from the peripheral nervous

system neurons. Neuropils are enriched in neuronal dendrites and axons, while neuronal cell

bodies are mostly absent in this region. This suggests that Wnd protein expression levels are

kept extremely low in neuronal cell bodies but enriched in neuronal processes likely in axon

terminals.

Next, we employed a transgene approach. We generated Wnd-GFP transgenes with and

without *wnd* mRNA's untranslated region (UTR), namely 5'UTR-Wnd::GFP-3'UTR and

Wnd::GFP to test a role of *wnd*-UTRs in Wnd protein localization. These transgenes were

expressed specifically in larval class IV dendritic arborization (C4da) neurons using the *ppk*-

GAL4 driver (Grueber et al., 2007). C4da neurons have their cell bodies positioned in the larval

body wall, where they extend dendrites and send axons to the VNC of the larval central nervous

system. This sensory neuron system provides an excellent opportunity for studying protein

localization in cell bodies, dendrites, and axon terminals (**Figure 1B**) (Grueber et al., 2007). We

quantified the expression levels of Wnd transgenes in the cell bodies and axon terminals of

C4da neurons using quantitative confocal microscope imaging. The expression levels were

normalized using a general membrane marker, mCD8::RFP. The expression levels of

mCD8::RFP in axon terminals may represent passive diffusion of membrane proteins, which

reports the absence of axonal enrichment or equals to the axonal enrichment of 1. We observed

a significant enrichment of Wnd in the axon terminals of C4da neurons, regardless of *wnd*-UTRs

(**Figure 1C** and **1D**). Overactive Wnd signaling impairs *ppk*-GAL4 activity, compounding our

analysis (Wang et al., 2013). To address this, we treated larvae with DLKi/GNE-3511, a specific

inhibitor of DLK/Wnd kinase activity (Patel et al., 2017; Russo and DiAntonio, 2019). The

treatment with DLKi did not affect larval viability or basal Cd4a morphology. Additionally, a

kinase-dead version of Wnd, Wnd-KD::GFP, exhibited similar axonal enrichment in comparison

to Wnd::GFP in the presence and absence of DLKi (**Figure 1 - supplement**).

What might be the biological significance of such high axon terminal enrichment of Wnd

proteins? Interestingly, a previous work has shown that nerve injury in *Drosophila* larvae

increased Wnd protein levels in proximal axons but not in motor neuron cell bodies (Xiong et al.,

2010), which suggests that injury signal does not affect Wnd protein levels in neuronal cell

bodies. Thus, we wondered whether Wnd protein degradation under normal conditions is also

compartment specific. To test this possibility, we expressed Wnd-KD::GFP in C4da neurons of

wild-type and *hiw* mutant animals ($hiw^{\Delta N}$). The expression levels of Wnd-KD::GFP was

measured in C4da cell bodies and C4da axon terminals by quantitative confocal microscope

imaging (**Figure 1E**). The result showed that *hiw* mutations did not change Wnd-KD::GFP

expression in C4da cell bodies as compared to a wild-type control. In contrast, there was a

remarkable 7-fold increase in Wnd-KD::GFP expression levels specifically in the axon terminals

in the presence of *hiw* mutations.

Taken together, these suggest that Wnd proteins are highly enriched in axon terminals and that

this axon terminal localization is a necessary step for Hiw-mediated Wnd protein turnover. DLK

proteins act as a sensor in neuronal stress responses, and under normal conditions, their

protein levels are constantly suppressed to prevent undesired stress responses (Nakata et al.,

2005; Collins et al., 2006; Xiong et al., 2010; Huntwork-Rodriguez et al., 2013). In this context, if

Wnd localization in axon terminals is disrupted, it may lead to dysregulated Wnd protein

expression levels, potentially resulting in exacerbated stress signaling. To test the model, we

decided to identify the mechanisms underlying Wnd localization in axon terminals.

The protein palmitoylation of Wnd is essential for axonal localization of Wnd.

The coding region of *wnd* was sufficient for Wnd axon terminal localization (**Figure 1**). Wnd

proteins are found on axonally transporting vesicles (Xiong et al., 2010). Together these further

suggest that Wnd protein localization in axon terminals are through axonal anterograde

transport of Wnd proteins, and that Wnd protein degradation is dependent on this axonal

anterograde transport. Protein transport often involves small protein domains or motifs. To find

the motifs necessary for axonal transport of Wnd, we performed a structure-function analysis by

progressively deleting 100 amino acids (aa) from Wnd transgenes (**Figure 2A**). These Wnd

transgenes were expressed in C4da neurons, and the axonal enrichment of Wnd was measured

as described in Figure 1D. Deletion of the first 100 aa of Wnd (WndΔ100::GFP) did not have an

impact on enrichment of Wnd in axon terminals. However, deleting the first 200 aa of Wnd

(WndΔ200::GFP) resulted in a dramatic 43-fold reduction of Wnd enrichment. Similarly, deleting

aa 101-200 of Wnd (WndΔ101-200::GFP) led to a dramatic decrease in Wnd localization in

axon terminals (**Figure 2B**). These findings suggest that the 101-200 aa region of Wnd contains

a crucial motif for axonal anterograde transport of Wnd.

Interestingly, this region contains an evolutionarily conserved palmitoylation motif, Cysteine-130

(Wnd-C130) (**Figure 2C**). Previous studies have shown that palmitoylation is required for DLK

association with neuronal vesicles and DLK signaling in cultured mammalian neurons and *C.*

elegans (Holland et al., 2016). However, whether DLK palmitoylation changes DLK protein

localization in axon terminals remains unknown. Additionally, whether Wnd undergoes protein

palmitoylation is not known. To examine the protein palmitoylation of Wnd, we introduced a

mutation changing the Cysteine to Serine, resulting in Wnd-C130S. We expressed wild-type

Wnd (Wnd::GFP) and the palmitoylation site mutant Wnd (Wnd-C130S::GFP) in cultured

Drosophila Schneider 2 (S2) cells using PEI-mediated transfection (Longo et al., 2013). The

Acyl-Biotin Exchange (ABE) assay (Drisdel et al., 2006) was conducted to measure protein

palmitoylation of Wnd proteins after pulldown using a GFP-tag. The results showed a 50%

reduction in protein palmitoylation in Wnd-C130S::GFP compared to Wnd::GFP (**Figure 2D**),

providing strong evidence that Wnd-C130 is a palmitoylation site. Subsequently, we expressed

Wnd::GFP and Wnd-C130S::GFP in larval C4da neurons and measured their axonal

enrichment. The findings revealed a dramatic 30-fold reduction in Wnd axonal localization for

Wnd-C130S::GFP compared to Wnd::GFP (**Figure 2E**). Taken together, our data demonstrates

that Wnd-C130 is palmitoylated in *Drosophila,* which is essential for Wnd localization in axon

terminals.

Interestingly, we noticed that unlike diffuse cytoplasmic expression patterns of palmitoylation-

defective DLK observed in mammalian cells (Holland et al., 2016), Wnd-C130S::GFP exhibited

discrete localization patterns in C4da cell bodies (**Figure 2E**). These structures extensively

colocalized with somatic Golgi complex (**Figure 2 – supplement 1**). In mammalian cultured

cells, DLK was found in somatic Golgi complex where its palmitoyltransferase, HIP14 localizes

(Niu et al., 2020). Consistently, we found that Wnd is localized to somatic Golgi complex in

C4da cell bodies, which required the 101-200 aa region of Wnd (**Figure 2 – supplement 2**).

Drosophila HIP14, dHIP14 showed somatic Golgi localization and extensive colocalization with

Wnd proteins in C4da cell bodies (**Figure 2 – supplement 3**). Furthermore, mutating *dHIP14*

significantly decreased Wnd enrichment in axon terminals (**Figure 2 – supplement 3**). These

suggest that Wnd protein palmitoylation is catalyzed by dHIP14 in somatic Golgi complex like

mammalian cells. However, unlike mammalian and *C. elegans* DLK proteins, Wnd

palmitoylation is not necessary for Golgi localization since Wnd-C130S::GFP retained Golgi

localization (**Figure 2 – supplement 1**). Moreover, Wnd palmitoylation plays a unique role in

localizing Wnd proteins in axon terminals in *Drosophila*.

Palmitoylation-defective Wnd evades protein turnover and accumulates in neuronal cell

bodies.

Wnd-C130S does not localize in the axon terminals but rather retained in neuronal cell bodies.

This may prevent protein turnover of Wnd-C130S, which may result in increased protein

expression levels of Wnd-C130S. To test the idea, we expressed Wnd::GFP and Wnd-

C130S::GFP using a pan-neuronal driver, *Elav*-GAL4. The larvae were treated with DLKi/GNE-

3511 to prevent lethality caused by overactive Wnd signaling. The total protein expression levels

of Wnd::GFP and Wnd-C130S::GFP were measured in larval brain lysates using Western blot

analysis with GFP antibody (**Figure 3A**). The results revealed a dramatic 7.1-fold increase in

protein levels of Wnd-C130S::GFP compared with Wnd::GFP. To determine the cellular

localization of increased Wnd-C130S::GFP expression, we expressed Wnd::GFP and Wnd-

C130S::GFP along with mCD8::RFP in larval C4da neurons using *ppk*-GAL4 and directly

compared the protein expression levels of the Wnd transgenes in C4da axon terminals and cell

bodies as measured by GFP fluorescent intensity (**Figure 3B**). We observed a dramatic 10-fold

increase in protein expression levels of Wnd-C130S::GFP in C4da cell bodies compared with

Wnd::GFP. On the other hand, Wnd-C130S::GFP exhibited greatly reduced protein expression

levels in C4da axon terminals (3-fold reduction) compared with Wnd::GFP. Together, these

findings strongly support the notion that Wnd localization in axon terminals is crucial for protein

turnover of Wnd.

Palmitoylation-defective Wnd triggers excessive stress signaling and neuronal loss.

DLK triggers neuronal stress signaling (Nakata et al., 2005; Collins et al., 2006; Karney-Grobe

et al., 2018). A failure to suppress Wnd protein levels in Wnd-C130S::GFP may lead to

excessive neuronal stress signaling. While the essential role of palmitoylation in DLK signaling

has been demonstrated in *C. elegans* and cultured mammalian cells (Holland et al., 2016;

Martin et al., 2019; Niu et al., 2020, 2022), its impact on the signaling capacity of Wnd remains

unknown. In contrast to the diffuse cytoplasmic expression pattern of palmitoylation-deficient

DLK proteins in worms and mammalian cells, Wnd-C130S::GFP exhibited clear somatic Golgi

localization (**Figure 2 – supplement 1**). A recent study suggested that the membrane

localization of DLK proteins, rather than palmitoylation itself, is crucial for DLK signaling (Tortosa

et al., 2022). Therefore, we investigated whether Wnd-C130S::GFP retains signaling capacity.

We expressed Wnd::GFP and Wnd-C130S::GFP in larval C4da neurons using *ppk-GAL4*. We

used *puc-lacZ,* a transcription reporter downstream of c-Jun N-terminal Kinase (JNK) as a proxy

for Wnd signaling (Martin-Blanco et al., 1998). Expression levels of *puc-lacZ* were evaluated by

immunostaining and its fluorescent intensity was measured in C4da nucleus. The results

showed robust induction of *puc-lacZ* by Wnd::GFP as well as by Wnd-C130S::GFP (**Figure 4 –**

supplement 1A). Additionally, the excessive axon arborization induced by Wnd signaling was

also observed both in Wnd::GFP and Wnd-C130S::GFP expressing C4da neurons. These were

dependent on Wnd kinase activity since there was no exuberant axon arborization by Wnd-

KD::GFP nor by Wnd-C130S-KD::GFP – the kinase dead versions of Wnd and Wnd-C130S

respectively (**Figure 4 – supplement 1B**). These strongly suggest that unlike DLK proteins in *C.*

elegans and mammalian cells, the palmitoylation of Wnd is not essential for signaling in

Drosophila.

Wnd-C130S::GFP exhibited 7-fold increase in protein expression levels compared to Wnd::GFP

(**Figure 3A**) while showing similar signaling capacity both in axon arborization and *puc-lacZ*

expression. We wondered if excessive transgene expression levels caused signaling pathway

saturation. Thus, we decided to reduce transgene expression levels. We recently have

generated a plasmid system for reduced transgene expression (Singh and Kim, 2022). Using

these plasmids, additional Wnd transgenes containing two UAS sites (2XUAS) rather than five

UAS sites (5XUAS) were generated and were expressed in larval C4da neurons. Then, *puc-*

lacZ expression levels were measured. The result showed that Wnd-C130S::GFP induced a

3.7-fold more *puc-lacZ* expression than Wnd::GFP when expressed under 2XUAS (**Figure 4A**).

This strongly suggests that Wnd-C130S::GFP triggers stronger stress signaling because it

cannot be transported towards axon terminals thus, evades protein turnover.

To determine whether this excessive Wnd signaling from Wnd-C130S::GFP results in neuronal

loss, we expressed Wnd-C130S::GFP and Wnd::GFP under *GMR-GAL4*, eye specific driver

(Seong et al., 2001). Expressing Wnd transgenes with 5XUAS caused complete lethality under

a standard growth condition for both Wnd::GFP and Wnd-C130S::GFP. Growing flies at a lower

temperature partially rescued lethality - likely due to lower transgene expression levels. Severe

eye loss phenotypes were observed from Wnd-C130S::GFP and Wnd::GFP expressing flies in

this experimental condition. We observed, albeit to a small extent, more severe eye loss

phenotypes in Wnd-C130S::GFP flies as compared to Wnd::GFP flies (**Figure 4 – supplement**

2A and 2B). To substantiate our findings, we expressed Wnd-C130S::GFP and Wnd::GFP

under 2XUAS using *GMR-GAL4*. Even with 2XUAS transgenes, animals expressing Wnd-

C130S::GFP showed significant lethality as compared to Wnd::GFP expressing animals (**Figure**

4B). Moreover, 100% of surviving flies of Wnd-C130S::GFP exhibited rough eye and eye

pigmentation loss phenotypes, which indicates loss of photoreceptors. On the other hand, flies

expressing 2XUAS-Wnd::GFP did not show any discernable phenotypes as compared to the

controls (**Figure 4C**). Wnd transgenes alone did not exhibit any defects in eye development.

These strongly suggest that axonal anterograde transport of Wnd is essential for Wnd protein

turnover and that a failure of this transport triggers excessive stress signaling.

Selective involvement of Rab proteins in Wnd axonal localization.

Protein palmitoylation enhances the hydrophobicity of target proteins, leading to increased

membrane association. This suggests that Wnd proteins are transported to axons through an

intracellular vesicular transport system. The regulation of intracellular vesicular transport is

governed by Rab proteins (Stenmark, 2009). To gain insights on axonal anterograde transport of

DLK proteins, we conducted a screening of Rab proteins to identify those that contribute to Wnd

axonal localization. We tested a total of 26 Rab proteins out of the 33 annotated *Drosophila* Rab

proteins (Zhang et al., 2007). YFP-tagged dominant-negative (DN) Rab proteins were

expressed along with RFP-fused Wnd-KD (Wnd-KD::mRFP) in C4da neurons. The axonal

enrichment of Wnd-KD::mRFP was analyzed by measuring the fluorescence intensities of Wnd-

KD::mRFP in C4da cell bodies and axon terminals (**Figure 5**). Among the tested DN-Rab

proteins, only Rab1 and Rab11 showed a significant reduction in Wnd axonal enrichment. These

findings suggest that Wnd targeting to axon terminals is a highly specific process. Rab1 is

involved in ER to Golgi traffic. The dramatic decrease in Wnd axonal localization observed upon

Rab1 inhibition is interesting as Wnd palmitoylation likely occurs at the somatic Golgi complex.

Rab11 regulates membrane traffic at the recycling endosome (Stenmark, 2009).

Rab11 inhibition caused dysregulated Wnd protein levels.

Rab11-positive recycling endosomes (REs) play a role in mediating anterograde Golgi-to-plasma membrane trafficking (Ang et al., 2004; Satoh et al., 2005; Taguchi, 2013). This raises an interesting possibility that once palmitoylated at the Golgi, Wnd is sorted at the REs for axonal transport. To investigate this, we examined whether Wnd proteins are present at the REs in C4da cell bodies. Wnd-KD::mRFP was expressed in C4da neurons, and the Rab11-positive REs were labeled using endogenously tagged YFP::Rab11 (TI-YFP::Rab11) (Dunst et al., 2015). We analyzed the portion of Wnd-KD::mRFP that colocalizes with YFP::Rab11 using the Manders' Overlap Coefficient. The results showed extensive colocalization between Wnd-KD::mRFP and YFP::Rab11 in C4da cell bodies (Manders' coefficient 0.60 ± 0.14) (**Figure 6A**). Similar results were obtained with transgenic expression of YFP::Rab11 and Wnd-KD::mRFP (Manders' coefficient 0.81 ± 0.11) (**Figure 6B**). On the other hand, we did not detect any meaningful colocalization between YFP::Rab11 and Wnd-KD::mRFP in C4da axon terminals or in axons (Manders' coefficient 0.34 ± 0.14 and 0.41 ± 0.10 respectively) (**Figure 6 – supplement 1**). Scale bar = 3 µm. These suggest that Rab11 is involved in Wnd protein sorting at the somatic REs rather than transporting Wnd directly.

If Rab11 is necessary for Wnd anterograde transport, inhibiting Rab11 may lead to dysregulated Wnd protein turnover and increased Wnd protein levels. To test the idea, we expressed DN-YFP::Rab11 along with Wnd-KD::mRFP in C4da neurons. We directly compared the expression levels of Wnd-KD::mRFP in C4da cell bodies and axon terminals. A mCD8::GFP transgene was used as a control for DN-Rab11. The results revealed a dramatic 4.2-fold increase in Wnd-KD::mRFP expression levels in C4da cell bodies by DN-Rab11, but not in the axon terminals (**Figure 6C**), which is reminiscent of Wnd-C130S behavior (**Figure 3B**).

We further examined the role of Rab11 in Wnd axonal localization using *Rab11* mutants, *Rab11*[93Bi], which is a strong loss-of-function allele (Giansanti et al., 2007). Wnd-KD::GFP was

expressed along with mCD8::RFP in wild-type and homozygous *Rab11[93Bi]* C4da neurons, and

the axonal enrichment of Wnd-KD::GFP was measured. The results showed a dramatic

reduction in Wnd axonal enrichment by *Rab11* mutations (**Figure 6D**). Next, we measured the

total expression levels of endogenous Wnd using Western blot analysis on brain lysates from

wild-type control and homozygous *Rab11[93Bi]* larvae (**Figure 6E**). The results showed a

significant 2.8-fold increase in total Wnd protein levels by *Rab11* mutations. This increase in

Wnd protein levels was not due to increased *wnd* gene expression because *wnd* mRNA levels

were not increased by *Rab11* mutations (**Figure 6 - supplement 2**). Rather, it showed a

significant reduction in *wnd* mRNA, which may be caused by a possible negative feedback from

excessive Wnd signaling. Together, these findings strongly suggest that Rab11 mediates Wnd

protein transport out of neuronal cell bodies towards axon terminals and that a failure in this

transport leads to dysregulated Wnd protein turnover.

Neuronal loss caused by Rab11 loss of function requires Wnd activity.

A previous work has shown that *Rab11* mutations caused photoreceptor loss in *Drosophila* eye

and JNK signaling activation (Tiwari and Roy, 2009). However, how *Rab11* mutations induce

JNK signaling is not clear. One of the pathways leading to JNK activation is DLK/Wnd (Collins et

al., 2006; Xiong et al., 2010; Ghosh et al., 2011). Thus, we hypothesized that increased Wnd

expression contributes to the neuronal loss and JNK activation by *Rab11* loss-of-function. To

test the idea, we expressed DN-Rab11 along with a membrane marker mCD8::RFP in the larval

C4da sensory neurons using *ppk-GAL4* (Grueber et al., 2007). JNK activation was monitored by

puc-lacZ induction (Martin-Blanco et al., 1998). The result showed a 2.8-fold increase of *puc-*

lacz expression in C4da neurons by DN-Rab11 (**Figure 7A**), which was completely abolished by

treating larvae with DLKi/GNE-3511, Wnd kinase inhibitor. This strongly suggests that increased

Wnd protein levels, hence Wnd activity mediates JNK activation induced by DN-Rab11.

Expressing DN-Rab11 caused severe eye loss phenotypes (**Figure 7B**). To test whether

increased Wnd expression contributes to the neuronal loss caused by DN-Rab11, we treated

DN-Rab11 expressing flies with Wnd kinase inhibitor, DLKi/GNE-3511. The treatment partially

but significantly mitigated eye loss phenotype caused by DN-Rab11 expression (Elav-GAL4/DN-

Rab11) while having no effect on eye development in control flies (Elav-GAL4 and DN-Rab11)

(**Figure 7B**). Furthermore, genetically reducing Wnd activity by introducing a loss of *wnd* allele,

wnd^3, in DN-Rab11 expressing photoreceptors significantly ameliorated the eye loss phenotypes

by DN-Rab11 (**Figure 7C**).

Taken together, these support a model in which Wnd proteins are transported to the axon

terminals for Hiw-mediated protein degradation under normal conditions, and that Rab11 is

essential for anterograde sorting of Wnd towards axon terminals. In this model, Rab11

inactivation leads to defective anterograde transport of Wnd, thus a deficit in Wnd protein

degradation. Subsequent increase in Wnd protein levels triggers excessive stress signaling

leading to neuronal death (**Figure 8**).

Discussion

Here, we report that axonal localization and protein turnover of Wnd are tightly coupled and that

failure in this coupling leads to dysregulated Wnd protein turnover and triggers excessive stress

signaling. We further uncover the role of Rab11 in this coupling.

Many neuronal proteins exhibit polarized protein expression patterns. DLK proteins are found in

the axonal compartments of diverse animals (Xiong et al., 2010; Klinedinst et al., 2013; Holland

et al., 2016; Niu et al., 2020, 2022). However, the extent of DLK axonal localization has not

been thoroughly studied, presumably because of technical limitations. We employed *Drosophila*

larval sensory neurons to answer this question. Our data suggests that Wnd proteins are highly

enriched in the axon terminals of these neurons (**Figure 1**). Given that Wnd encodes a

cytoplasmic protein, the extent of axonal enrichment was rather surprising. Wnd protein levels

are constantly suppressed by Hiw in the axon terminals. Without such suppression, Wnd

proteins would show well over 70-fold enrichment in axons. The UTRs of *wnd* are dispensable

for Wnd axonal enrichment (**Figure 1C**), which suggests that axonal anterograde transport of

Wnd is responsible for such axonal enrichment. This is supported by previous observations of

DLK proteins on membrane vesicles undergoing both anterograde and retrograde transport in

axons (Xiong et al., 2010; Holland et al., 2016; Li et al., 2017). Taken together, these indicate

that Wnd proteins are synthesized in the neuronal cell body and subsequently transported into

axon terminals through a highly active and efficient anterograde transport process.

What might be such a transport mechanism? DLK genes encode cytoplasmic proteins. Protein

palmitoylation is an important post-translational mechanism for protein transport (Aicart-Ramos

et al., 2011; Tortosa and Hoogenraad, 2018). It increases hydrophobicity of target proteins to aid

membrane association so that palmitoylated proteins are transported along with the membrane

vesicles they associate. In our study, we found that protein palmitoylation on Wnd at the

Cysteine residue 130 (Wnd-C130) is essential for its axonal localization. Interestingly, the

palmitoylation-defective mutant Wnd-C130S exhibited large punctate structures in the cell body

(**Figure 2E**), which colocalized with the somatic Golgi complex (**Figure 2 – supplement 1**). This

suggests that Wnd-C130 palmitoylation is not solely responsible for membrane association. The

presence of residual palmitoylation in Wnd-C130S (**Figure 2D**) may indicate the existence of an

additional palmitoylation site, which may explain how Wnd-C130S associates with the Golgi.

Consistent with our findings, previous studies have shown that DLK proteins in mammalian cells

and *C. elegans* are palmitoylated on a single Cysteine residue, and when this residue is

mutated, the proteins exhibit a diffuse cytoplasmic expression pattern (Holland et al., 2016).

These observations raise the question of why Wnd-C130S, with its residual palmitoylation and

membrane association, fails to be transported to axon terminals. Interestingly, recent research

has proposed a role for protein palmitoylation in anterograde transport within the Golgi complex

(Ernst et al., 2018). We envision that Wnd-C130 palmitoylation is a necessary sorting step

towards the axonal anterograde transport from the somatic Golgi complex rather than merely

acting as a membrane anchor.

The role of Rab11 in Wnd axonal targeting and protein turnover uncovered by our DN-Rab

screening suggests a role of RE in these processes (**Figure 5 and 6**). Rab11 plays a critical role

in membrane trafficking events at the RE and is often used as a marker for the RE (Taguchi,

2013). It is involved in the anterograde transport of cargo from the Golgi apparatus to the

plasma membrane through the RE (Ang et al., 2004; Satoh et al., 2005; Taguchi, 2013). We

envision that once Wnd is palmitoylated, it is sorted at the peri-nuclear RE for subsequent

axonal anterograde transport. Our observation of significant colocalization between Wnd

proteins and Rab11-positive puncta in C4da cell bodies (**Figure 6**) but not in axon terminals or

in axons (**Figure 6 - supplement 1**) supports the idea that Wnd proteins are associated with the

RE during their anterograde transport. These further suggest that Rab11 is not directly involved

in the anterograde long-distance transport of Wnd proteins, rather is responsible for sorting Wnd

into the axonal anterograde transporting vesicles.

Inhibiting Rab11 activity led to increased Wnd protein levels (**Figure 6E**). Reduced activity or

expression levels of Rab11 have been reported in experimental models of Huntington's disease,

Parkinson's disease and in patients with amyotrophic lateral sclerosis (Li et al., 2009a, 2010;

Breda et al., 2015; Mitra et al., 2019; Rai and Kumar Roy, 2022). Furthermore, recent studies

point to the involvement of DLK in multiple neurodegenerative conditions (Pozniak et al., 2013;

Welsbie et al., 2013, 2017; Larhammar et al., 2017a; Le Pichon et al., 2017; Patel et al., 2017).

These raise a tantalizing possibility that disrupted anterograde transport of DLK, and

subsequent dysregulated DLK protein turnover, could be a contributing factor in these

neurodegenerative disorders. In this regard, it is highly encouraging that inhibiting Wnd activity

mitigated the neuronal loss and JNK activation by *Rab11* loss-of-function (**Figure 7**). Further

investigations are needed to elucidate the causal role of dysregulated DLK protein turnover in

neurodegenerative conditions and to better understand the interplay between Rab11-mediated

transport, protein turnover, and DLK-mediated stress signaling.

The mechanism by which Hiw performs compartmentalized Wnd protein turnover in axons is an

interesting question. There are several possibilities to consider. One possibility is that Hiw

protein itself is specifically present in the axon terminals and not in the cell body. Studies have

shown that Phr1, the mammalian homolog of Hiw, is expressed in axonal segments of motor

neuron explants and in the axon shaft of sensory neurons, but its expression in the growth cone

is low or absent (Lewcock et al., 2007). However, it is not known whether Hiw proteins are

present in the cell body and to what extent. Unfortunately, the lack of an immunostaining

compatible Hiw antibody has limited our ability to test this possibility. Nevertheless, transgenic

expression of GFP::Hiw in *Drosophila* has shown significantly higher expression in the motor

neuron cell bodies (Xiong et al., 2010). Moreover, a study has demonstrated that Hiw controls

the expression of dNMNAT2, a substrate of Hiw, both in the cell body and the neuromuscular

junction (Xiong et al., 2012). These observations suggest that Hiw expression itself may not be

restricted to the axon terminals. Another possibility is that DLK proteins are protected from

PHR1/RMP-1/Hiw-mediated protein turnover in the cell body. DLK protein stability can be

enhanced by Hsp90 (Karney-Grobe et al., 2018), protein phosphorylation by JNK (Huntwork-

Rodriguez et al., 2013), and PKA (Hao et al., 2016). These protective pathways may collaborate

with PHR1/RMP-1/Hiw to execute axon-specific protein turnover of DLK proteins.

Understanding the interplay between these protective mechanisms and PHR1/RMP-1/Hiw in

regulating DLK protein turnover will provide valuable insights on subcellular specific Wnd protein

turnover.

The coupling between axonal transport and protein turnover of DLK proteins may have

important implications in the context of neurodegenerative disorders. Recent studies have

implicated DLK as a central player in neuronal death, including in diseases such as Alzheimer's

disease and amyotrophic lateral sclerosis (Pozniak et al., 2013; Welsbie et al., 2013, 2017;

Larhammar et al., 2017a, 2017b; Le Pichon et al., 2017; Patel et al., 2017). Defective axonal

transport is a common feature observed in many neurodegenerative disorders (Chevalier-

Larsen and Holzbaur, 2006; De Vos et al., 2008; Guo et al., 2020), and it has been proposed to

contribute to disease pathogenesis. Interestingly, studies have reported increased DLK protein

levels in neuronal cell bodies following neuronal insults (Watkins et al., 2013; Welsbie et al.,

2013). Furthermore, DLK signaling has been shown to be activated by cytoskeletal disruptions

(Valakh et al., 2013, 2015), which may impair cytoskeleton-based axonal transport. This raises

an intriguing possibility that disrupted axonal transport, leading to the accumulation of DLK

proteins, could be a contributing factor for DLK activation and neuronal loss in

neurodegenerative diseases.

    Materials and Methods

Drosophila Strains, rearing, DNA Constructs for Generating Transgenic Flies

The following *Drosophila* strains were reared at 25°C in a humidified chamber. *ppk-GAL4* (Grueber et al., 2007); *UAS-dHIP14::Tdtomato* and *Dhip14^{ex11}* (Stowers and Isacoff, 2007); *hiw$^{\Delta N}$* (Wu et al., 2005); *UAS-ManII::GFP* (Ye et al., 2007); *UAS-ManII-Tag::RFP* (Zhou et al., 2014); *Mi{MIC}wnd ^{GFSTF}* (Venken et al., 2011); *GMR-Gal4* (Seong et al., 2001); *elav-Gal4^{C155}* (BDSC #459); *w^{1118}* (BDSC #3605); *TI{TI}Rab11^{EYFP}* (BDSC #62549)(Dunst et al., 2015); *UASp-YFP.Rab11* (BDSC #50782)(Zhang et al., 2007); *Rab11^{93Bi}* (BDSC #4158)(Giansanti et al., 2007); *puc-lacZ* (Martin-Blanco et al., 1998). For DN-Rab screening, the fly lines (Zhang et al., 2007) were obtained from the Bloomington Drosophila Stock Center and the full list of fly lines is provided in Table 1.

To generate DNA constructs of *Wnd* with its natural 5'- and 3'-UTRs and the serial deletion constructs of Wnd, the genomic sequences of the 5'-UTR and 3'-UTR of *Wnd* were obtained from *w^{1118}* genomic DNA by PCR. The coding sequence of *Wnd* was recovered from UAS-Wnd (Collins et al., 2006) via PCR and inserted into the pUASTattB plasmid using NotI and XbaI restriction sites. UAS-Wnd-KD::GFP, Wnd-KDΔ100::GFP, and WndC130S-KD::GFP were generated by site-directed mutagenesis using In-Fusion® Snap Assembly Master Mix (Takara Bio USA, Inc, San Jose, CA). UAS-Wnd-C130S::GFP and UAS-WndΔ101-200::GFP were generated by site-directed mutagenesis. To generate UAS-Wnd-KD::mRFP, the coding sequence of Wnd-KD was obtained from UAS-Wnd-KD::GFP via PCR and inserted into the pUASTattB plasmid using NotI and XbaI sites. To generate 2XUAS-Wnd::GFP and 2XUAS-Wnd-C130S::GFP, the coding sequences of Wnd and Wnd-C130S were obtained from UAS-Wnd::GFP and UAS-Wnd-C130S::GFP via PCR respectively. Subsequently, they were inserted

into the p2xUASTattB plasmid (Addgene #189860)(Singh and Kim, 2022) using NotI and XbaI

restriction sites.

Transgenic flies were generated by PhiC31-mediated germline transformation using an identical

landing site (Bischof et al., 2007).

To dose larvae with DLKi (GNE-3511, Sigma Aldrich), DLKi was dissolved in DMSO and was

added to a final concentration of 35 µM in fly food. Fly larvae were raised directly on either

control (DMSO) or DLKi containing food.

Drosophila Schneider 2 cell culture, transfection, Palmitoylation assay, and Western blot.

Drosophila Schneider 2 (S2) cells were prepared as previously described (Singh et al., 2022).

Cells were transfected with indicated DNA constructs and tubulin-Gal4 (Lee and Luo, 1999)

using polyethyleneimine (Polysciences, Warrington, PA) (Longo et al., 2013).

Two days after transfection, the Acyl-Biotin Exchange (ABE) assay was performed (Drisdel et

al., 2006). Cells were lysed in the ice-cold Lysis buffer (LB, 50 mM HEPES/pH 7.4, 150 mM

NaCl, 1 mM EDTA, 1% NP40, 5% glycerol, 1 mM PMSF, 1 mM Na_3VO_4, 5 mM NaF) with Halt^{TM}

protease inhibitor cocktail EDTA- free (ThermoFisher Scientific, Waltham, MA) and 25 mM N-

ethylmaleimide (Sigma, St. Louis, MO), incubated for 1hr at 4˚C and centrifuged at 1000 xg for 5

min. The resulting post-nuclear lysates were centrifuged at 20,000 x g for 30 min at 4˚C. The

supernatant was incubated with the GFP selector (NanoTag Biotechnologies, Göttingen,

Germany) for 2 hr at 4˚C to pulldown Wnd-GFP proteins. The GFP selector beads along with

bound proteins were recovered by a brief centrifugation and washed two times with LB

containing 0.1% SDS before incubating with 1 M hydroxylamine hydrochloride (HAM, Sigma, St.

Louis, MO) in LB (pH 7.2) at room temperature for 1 hr. Half of the protein-bound GFP selector

beads were incubated with LB without HAM and used as a "- HAM" control. Then, the protein

GFP selector beads were washed with pH 6.2 LB twice and treated with 1.6 µM EZ-Link™

BMCC-Biotin (ThermoFisher Scientific, Waltham, MA) in pH 6.2 LB for 45 min at 4˚C. Beads

were washed twice with LB without NP40 before being processed for SDS-PAGE. Proteins were

separated by SDS-PAGE and were subjected to Western blot analysis with mouse monoclonal

anti-Biotin antibody (Jackson Immunoresearch, West Grove, PA) and chicken polyclonal anti-

GFP antibody (Aves Labs, Tigard, OR).

To measure Wnd expression in larva brain lysates, larvae were grown on fly food containing 35

µM DLKi food in a humidified chamber. The larval central nervous system (the ventral nerve

cord and brain lobes) was dissected out from the 3rd instar larvae and collected in ice-cold PBS

before recovery by a brief centrifugation at 4°C. The total brain lysates were prepared by directly

adding SDS-PAGE sample buffer and homogenization before being resolved on 8% SDS-PAGE

gel and subsequent Western blot analysis. The samples were blotted using anti-Wnd antibody

(Collins et al., 2006) to estimate total Wnd proteins. Anti-Elav antibody (DSHB - 9F8A9)(O'Neill

et al., 1994) was used as a loading control.

Immunostaining and Imaging

Drosophila 3rd instar larvae were prepared for immunostaining as previously described (Singh et

al., 2022). Primary antibodies used were rabbit polyclonal anti-RFP (Rockland

Immunochemicals, Limerick, PA), chicken polyclonal anti-GFP (Aves Labs, Tigard, OR), and

goat anti-horseradish peroxidase conjugated with Cy5 (Jackson ImmunoResearch, West Grove,

PA). The secondary antibodies used were Cy2- or Cy5-conjugated goat anti-chicken and Cy2-

or Cy5-conjugated goat anti-Rabbit (Jackson ImmunoResearch, West Grove, PA).

Confocal imaging was done with a Leica SP8 confocal system, or a custom-built spinning disk

confocal microscope equipped with a 63x oil-immersion objective with a 0.3 µm step-sizes.

For Wnd axonal enrichment analysis, the C4da neuron cell bodies from abdominal segments 4-

6 and the corresponding segments in the ventral nerve cord were imaged. The resulting 3D

images were projected into 2D images using a maximum projection method in the imageJ

software. A region of interest was drawn in the cell body of C4da neurons, and the mean

fluorescence intensity was measured using ImageJ software. The mean fluorescence intensity

of UAS-mCD8::mRFP transgene expression was measured from the same region of interest

and used as a normalization control. The mean intensities of the C4da cell bodies from a single

larva were averaged out to calculate the axonal enrichment index. The axonal enrichment index

was defined as Wnd expression (mean intensity) in the axon terminals divided by an average

Wnd expression in cell bodies (mean intensity).

To analyze Golgi and Wnd colocalization, the resulting 3D images were projected into 2D

images using a maximum projection method in the ImageJ software. The Manders' Overlap

Coefficient (Pike et al., 2017) was measured as the percentage of total GFP signal overlapping

with RFP signal using the ImageJ plugin, JACoP (Bolte and Cordelières, 2006). To analyze

puncta colocalization, we used the Manders' coefficient (Pike et al., 2017) that is correlation

between each channel signal using JACop. For Wnd colocalization with TI-YFP::Rab11, a single

confocal slice was selected.

Drosophila eye degeneration

The flies that are 24 – 48 hr after eclosion were imaged using the Basler ace acA2440-75 µm

CCD-camera or Moticam A5 camera installed on a stereomicroscope. The region of interest was

drawn on the fly eye area from the images and expressed as pixel numbers divided by 1000.

Flies were raised at 27°C in a humidified incubator except for Figure 4 – supplement 2.

Drosophila Viability and Lethality Analysis

Flies were raised at 27°C. Viability was determined by counting the number of filial pupae generation (adult flies after eclosion) and pupae lethality by counting the number of dead pupae per vial.

Quantitative reverse transcription PCR

Total RNAs were extracted from the wondering third instar larval brains from a wild-type (w^{1118}) or homozygous *Rab11* ^{93bi} mutants using the Quick-RNA MicroPrep kit (Zymo Research, Irvine, CA), and were subjected to reverse transcription using a mixture of gene-specific primers as follows, GAPDH1 (CTTCATTCGATGCACAAGTTTTATTTTTCAAATAGCTGG) and wnd (CCATACAGTTTAGGTCGATTTCTC, CAACTTCGGGCCTAATCAAGG, GCGTAACTTGGACAACTCCAA, GGAGATAGTTTTTGCTGCGTAAC, and GGCCCCCACCTAGACTAGAT).

The quantitative PCR was conducted using SYBR MasterMix (ThermoFisher Scientific, Waltham, MA). The following gene specific primer pairs were used.

GAPDH1, ATGTCTCCGTTGTGGATCTTAC and CCTCGACCTTAGCCTTGATTT; wnd primer1, TCTGCGGAGAAGAAGCATTGG and TGGATGGCAATCTTCTGGGAG; wnd primer 2, GTTCGATAGGCAATGCCAAC and CGCAGAGCGTACTTGTAGTC.

The mRNA abundance of *wnd* was calculated by the 2ΔΔCT method using GAPDH1 as a normalizing control.

Experimental design and statistical analysis.

All statistical analysis was performed as two-tailed using GraphPad Prism software version

7.04. The Mann–Whitney test was used for immunostaining results. A ratio paired t-test was

used for Western blot analysis (Figure 3A and 6E). For multigroup comparisons, one-way

ANOVA was used followed by post hoc Tukey's multiple comparisons tests. The Chi-square test

was performed for Drosophila lethality analysis (Figure 4B). A p-value smaller than 0.05 was

considered statistically significant. All p values are indicated as NS; non-significant, *; $P < 0.05$,

; $P < 0.01$, *; $P < 0.001$, and ****; $P < 0.0001$

Acknowledgments

We thank Drs. Steve Stowers and Catherine Collins for sharing reagents. Research reported in

this study used the Cellular and Molecular Imaging Core facility at the University of Nevada

Reno, which was supported by NIH P30 GM145646. This study was supported by NIH

R01NS116463 and P20GM103440 (NV INBRE) to J.K. The content is solely the responsibility of

the authors and does not necessarily represent the official views of the National Institutes of

Health.

Author contributions

S.K., Y.Q., M.S., and J.K. designed the experiments. S.K., Y.Q., and M.S. experiments. S.K. and

J.K. wrote the manuscript.

Declaration of interests

The authors declare no competing financial interests.

<u>References</u>

Aicart-Ramos, C., Valero, R.A., and Rodriguez-Crespo, I. (2011). Protein palmitoylation and
subcellular trafficking. Biochim Biophys Acta 1808, 2981-2994. 10.1016/j.bbamem.2011.07.009.

Ang, A.L., Taguchi, T., Francis, S., Fölsch, H., Murrells, L.J., Pypaert, M., Warren, G., and
Mellman, I. (2004). Recycling endosomes can serve as intermediates during transport from the
Golgi to the plasma membrane of MDCK cells. J Cell Biol 167, 531-543.
10.1083/jcb.200408165.

Bischof, J., Maeda, R.K., Hediger, M., Karch, F., and Basler, K. (2007). An optimized
transgenesis system for Drosophila using germ-line-specific phiC31 integrases. Proc Natl Acad
Sci U S A 104, 3312-3317. 10.1073/pnas.0611511104.

Bolte, S., and Cordelières, F.P. (2006). A guided tour into subcellular colocalization analysis in
light microscopy. J Microsc 224, 213-232. 10.1111/j.1365-2818.2006.01706.x.

Breda, C., Nugent, M.L., Estranero, J.G., Kyriacou, C.P., Outeiro, T.F., Steinert, J.R., and
Giorgini, F. (2015). Rab11 modulates α-synuclein-mediated defects in synaptic transmission and
behaviour. Hum Mol Genet 24, 1077-1091. 10.1093/hmg/ddu521.

Calkins, M.J., Manczak, M., Mao, P., Shirendeb, U., and Reddy, P.H. (2011). Impaired
mitochondrial biogenesis, defective axonal transport of mitochondria, abnormal mitochondrial
dynamics and synaptic degeneration in a mouse model of Alzheimer's disease. Hum Mol Genet
20, 4515-4529. 10.1093/hmg/ddr381.

Chevalier-Larsen, E., and Holzbaur, E.L. (2006). Axonal transport and neurodegenerative
disease. Biochim Biophys Acta 1762, 1094-1108. 10.1016/j.bbadis.2006.04.002.

Chu, Y., Morfini, G.A., Langhamer, L.B., He, Y., Brady, S.T., and Kordower, J.H. (2012).
Alterations in axonal transport motor proteins in sporadic and experimental Parkinson's disease.
Brain 135, 2058-2073. 10.1093/brain/aws133.

Collard, J.F., Côté, F., and Julien, J.P. (1995). Defective axonal transport in a transgenic mouse
model of amyotrophic lateral sclerosis. Nature 375, 61-64. 10.1038/375061a0.

Collins, C.A., Wairkar, Y.P., Johnson, S.L., and DiAntonio, A. (2006). Highwire restrains synaptic
growth by attenuating a MAP kinase signal. Neuron 51, 57-69. 10.1016/j.neuron.2006.05.026.

De Vos, K.J., Grierson, A.J., Ackerley, S., and Miller, C.C. (2008). Role of axonal transport in
neurodegenerative diseases. Annu Rev Neurosci 31, 151-173.
10.1146/annurev.neuro.31.061307.090711.

De Vos, K.J., and Hafezparast, M. (2017). Neurobiology of axonal transport defects in motor
neuron diseases: Opportunities for translational research? Neurobiol Dis 105, 283-299.
10.1016/j.nbd.2017.02.004.

Dunst, S., Kazimiers, T., von Zadow, F., Jambor, H., Sagner, A., Brankatschk, B., Mahmoud, A.,
Spannl, S., Tomancak, P., Eaton, S., and Brankatschk, M. (2015). Endogenously tagged rab
proteins: a resource to study membrane trafficking in Drosophila. Dev Cell 33, 351-365.
10.1016/j.devcel.2015.03.022.

Drisdel, R.C., Alexander, J.K., Sayeed, A., and Green, W.N. (2006). Assays of protein
palmitoylation. Methods 40, 127-134. 10.1016/j.ymeth.2006.04.015.

Ernst, A.M., Syed, S.A., Zaki, O., Bottanelli, F., Zheng, H., Hacke, M., Xi, Z., Rivera-Molina, F.,
Graham, M., Rebane, A.A., et al. (2018). S-Palmitoylation Sorts Membrane Cargo for
Anterograde Transport in the Golgi. Dev Cell 47, 479-493.e477. 10.1016/j.devcel.2018.10.024.

Ghosh, A.S., Wang, B., Pozniak, C.D., Chen, M., Watts, R.J., and Lewcock, J.W. (2011). DLK
induces developmental neuronal degeneration via selective regulation of proapoptotic JNK
activity. J Cell Biol 194, 751-764. 10.1083/jcb.201103153.

Giansanti, M.G., Belloni, G., and Gatti, M. (2007). Rab11 is required for membrane trafficking
and actomyosin ring constriction in meiotic cytokinesis of Drosophila males. Mol Biol Cell 18,
5034-5047. 10.1091/mbc.e07-05-0415.

Grueber, W.B., Ye, B., Yang, C.H., Younger, S., Borden, K., Jan, L.Y., and Jan, Y.N. (2007).
Projections of Drosophila multidendritic neurons in the central nervous system: links with
peripheral dendrite morphology. Development 134, 55-64. 10.1242/dev.02666.

Guedes-Dias, P., and Holzbaur, E.L.F. (2019). Axonal transport: Driving synaptic function.
Science 366. 10.1126/science.aaw9997.

Guillaud, L., El-Agamy, S.E., Otsuki, M., and Terenzio, M. (2020). Anterograde Axonal Transport
in Neuronal Homeostasis and Disease. Front Mol Neurosci 13, 556175.
10.3389/fnmol.2020.556175.

Guo, W., Stoklund Dittlau, K., and Van Den Bosch, L. (2020). Axonal transport defects and
neurodegeneration: Molecular mechanisms and therapeutic implications. Semin Cell Dev Biol
99, 133-150. 10.1016/j.semcdb.2019.07.010.

Hao, Y., Frey, E., Yoon, C., Wong, H., Nestorovski, D., Holzman, L.B., Giger, R.J., DiAntonio, A.,
and Collins, C. (2016). An evolutionarily conserved mechanism for cAMP elicited axonal
regeneration involves direct activation of the dual leucine zipper kinase DLK. Elife 5.
10.7554/eLife.14048.

Holland, S.M., Collura, K.M., Ketschek, A., Noma, K., Ferguson, T.A., Jin, Y., Gallo, G., and
Thomas, G.M. (2016). Palmitoylation controls DLK localization, interactions and activity to
ensure effective axonal injury signaling. Proc Natl Acad Sci U S A 113, 763-768.
10.1073/pnas.1514123113.

Huntwork-Rodriguez, S., Wang, B., Watkins, T., Ghosh, A.S., Pozniak, C.D., Bustos, D.,
Newton, K., Kirkpatrick, D.S., and Lewcock, J.W. (2013). JNK-mediated phosphorylation of DLK
suppresses its ubiquitination to promote neuronal apoptosis. J Cell Biol 202, 747-763.
10.1083/jcb.201303066.

Karney-Grobe, S., Russo, A., Frey, E., Milbrandt, J., and DiAntonio, A. (2018). HSP90 is a
chaperone for DLK and is required for axon injury signaling. Proc Natl Acad Sci U S A 115,
E9899-E9908. 10.1073/pnas.1805351115.

Klinedinst, S., Wang, X., Xiong, X., Haenfler, J.M., and Collins, C.A. (2013). Independent
pathways downstream of the Wnd/DLK MAPKKK regulate synaptic structure, axonal transport,
and injury signaling. J Neurosci 33, 12764-12778. 10.1523/JNEUROSCI.5160-12.2013.

Larhammar, M., Huntwork-Rodriguez, S., Jiang, Z., Solanoy, H., Sengupta Ghosh, A., Wang, B., Kaminker, J.S., Huang, K., Eastham-Anderson, J., Siu, M., et al. (2017). Dual leucine zipper kinase-dependent PERK activation contributes to neuronal degeneration following insult. Elife 6. 10.7554/eLife.20725.

Larhammar, M., Huntwork-Rodriguez, S., Rudhard, Y., Sengupta-Ghosh, A., and Lewcock, J.W. (2017). The Ste20 Family Kinases MAP4K4, MINK1, and TNIK Converge to Regulate Stress-Induced JNK Signaling in Neurons. J Neurosci 37, 11074-11084. 10.1523/JNEUROSCI.0905-17.2017.

Le Pichon, C.E., Meilandt, W.J., Dominguez, S., Solanoy, H., Lin, H., Ngu, H., Gogineni, A., Sengupta Ghosh, A., Jiang, Z., Lee, S.H., et al. (2017). Loss of dual leucine zipper kinase signaling is protective in animal models of neurodegenerative disease. Sci Transl Med 9. 10.1126/scitranslmed.aag0394.

Lee, T., and Luo, L. (1999). Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. Neuron 22, 451-461. 10.1016/s0896-6273(00)80701-1.

Lee, W.C., Yoshihara, M., and Littleton, J.T. (2004). Cytoplasmic aggregates trap polyglutamine-containing proteins and block axonal transport in a Drosophila model of Huntington's disease. Proc Natl Acad Sci U S A 101, 3224-3229. 10.1073/pnas.0400243101.

Lewcock, J.W., Genoud, N., Lettieri, K., and Pfaff, S.L. (2007). The ubiquitin ligase Phr1 regulates axon outgrowth through modulation of microtubule dynamics. Neuron 56, 604-620. 10.1016/j.neuron.2007.09.009.

Li, J., Zhang, Y.V., Asghari Adib, E., Stanchev, D.T., Xiong, X., Klinedinst, S., Soppina, P., Jahn, T.R., Hume, R.I., Rasse, T.M., and Collins, C.A. (2017). Restraint of presynaptic protein levels by Wnd/DLK signaling mediates synaptic defects associated with the kinesin-3 motor Unc-104. Elife 6. 10.7554/eLife.24271.

Li, X., Sapp, E., Chase, K., Comer-Tierney, L.A., Masso, N., Alexander, J., Reeves, P., Kegel, K.B., Valencia, A., Esteves, M., et al. (2009). Disruption of Rab11 activity in a knock-in mouse model of Huntington's disease. Neurobiol Dis 36, 374-383. 10.1016/j.nbd.2009.08.003.

Li, X., Standley, C., Sapp, E., Valencia, A., Qin, Z.H., Kegel, K.B., Yoder, J., Comer-Tierney, L.A., Esteves, M., Chase, K., et al. (2009). Mutant huntingtin impairs vesicle formation from recycling endosomes by interfering with Rab11 activity. Mol Cell Biol 29, 6106-6116. 10.1128/MCB.00420-09.

Li, X., Valencia, A., Sapp, E., Masso, N., Alexander, J., Reeves, P., Kegel, K.B., Aronin, N., and Difiglia, M. (2010). Aberrant Rab11-dependent trafficking of the neuronal glutamate transporter EAAC1 causes oxidative stress and cell death in Huntington's disease. J Neurosci 30, 4552-4561. 10.1523/JNEUROSCI.5865-09.2010.

Li, Y., Ritchie, E.M., Steinke, C.L., Qi, C., Chen, L., Zheng, B., and Jin, Y. (2021). Activation of MAP3K DLK and LZK in Purkinje cells causes rapid and slow degeneration depending on signaling strength. Elife 10. 10.7554/eLife.63509.

Longo, P.A., Kavran, J.M., Kim, M.S., and Leahy, D.J. (2013). Transient mammalian cell transfection with polyethylenimine (PEI). Methods Enzymol 529, 227-240. 10.1016/B978-0-12-418687-3.00018-5.

Martin, D.D.O., Kanuparthi, P.S., Holland, S.M., Sanders, S.S., Jeong, H.K., Einarson, M.B.,
Jacobson, M.A., and Thomas, G.M. (2019). Identification of Novel Inhibitors of DLK
Palmitoylation and Signaling by High Content Screening. Sci Rep 9, 3632. 10.1038/s41598-019-
39968-8.

Martín-Blanco, E., Gampel, A., Ring, J., Virdee, K., Kirov, N., Tolkovsky, A.M., and Martinez-
Arias, A. (1998). puckered encodes a phosphatase that mediates a feedback loop regulating
JNK activity during dorsal closure in Drosophila. Genes Dev 12, 557-570.
10.1101/gad.12.4.557.

Miller, B.R., Press, C., Daniels, R.W., Sasaki, Y., Milbrandt, J., and DiAntonio, A. (2009). A dual
leucine kinase-dependent axon self-destruction program promotes Wallerian degeneration. Nat
Neurosci 12, 387-389. 10.1038/nn.2290.

Mitra, J., Hegde, P.M., and Hegde, M.L. (2019). Loss of endosomal recycling factor RAB11
coupled with complex regulation of MAPK/ERK/AKT signaling in postmortem spinal cord
specimens of sporadic amyotrophic lateral sclerosis patients. Mol Brain 12, 55. 10.1186/s13041-
019-0475-y.

Nakata, K., Abrams, B., Grill, B., Goncharov, A., Huang, X., Chisholm, A.D., and Jin, Y. (2005).
Regulation of a DLK-1 and p38 MAP kinase pathway by the ubiquitin ligase RPM-1 is required
for presynaptic development. Cell 120, 407-420. 10.1016/j.cell.2004.12.017.

Niu, J., Holland, S.M., Ketschek, A., Collura, K.M., Hesketh, N.L., Hayashi, T., Gallo, G., and
Thomas, G.M. (2022). Palmitoylation couples the kinases DLK and JNK3 to facilitate
prodegenerative axon-to-soma signaling. Sci Signal 15, eabh2674. 10.1126/scisignal.abh2674.

Niu, J., Sanders, S.S., Jeong, H.K., Holland, S.M., Sun, Y., Collura, K.M., Hernandez, L.M.,
Huang, H., Hayden, M.R., Smith, G.M., et al. (2020). Coupled Control of Distal Axon Integrity
and Somal Responses to Axonal Damage by the Palmitoyl Acyltransferase ZDHHC17. Cell Rep
33, 108365. 10.1016/j.celrep.2020.108365.

Patel, S., Meilandt, W.J., Erickson, R.I., Chen, J., Deshmukh, G., Estrada, A.A., Fuji, R.N.,
Gibbons, P., Gustafson, A., Harris, S.F., et al. (2017). Selective Inhibitors of Dual Leucine Zipper
Kinase (DLK, MAP3K12) with Activity in a Model of Alzheimer's Disease. J Med Chem 60, 8083-
8102. 10.1021/acs.jmedchem.7b00843.

Pensalfini, A., Kim, S., Subbanna, S., Bleiwas, C., Goulbourne, C.N., Stavrides, P.H., Jiang, Y.,
Lee, J.H., Darji, S., Pawlik, M., et al. (2020). Endosomal Dysfunction Induced by Directly
Overactivating Rab5 Recapitulates Prodromal and Neurodegenerative Features of Alzheimer's
Disease. Cell Rep 33, 108420. 10.1016/j.celrep.2020.108420.

Pike, J.A., Styles, I.B., Rappoport, J.Z., and Heath, J.K. (2017). Quantifying receptor trafficking
and colocalization with confocal microscopy. Methods 115, 42-54. 10.1016/j.ymeth.2017.01.005.

Pozniak, C.D., Sengupta Ghosh, A., Gogineni, A., Hanson, J.E., Lee, S.H., Larson, J.L.,
Solanoy, H., Bustos, D., Li, H., Ngu, H., et al. (2013). Dual leucine zipper kinase is required for
excitotoxicity-induced neuronal degeneration. J Exp Med 210, 2553-2567.
10.1084/jem.20122832.

Rai, P., and Kumar Roy, J. (2022). Endosomal recycling protein Rab11 in Parkin and Pink1
signaling in Drosophila model of Parkinson's disease. Exp Cell Res 420, 113357.
10.1016/j.yexcr.2022.113357.

Reddy, P.H., and Shirendeb, U.P. (2012). Mutant huntingtin, abnormal mitochondrial dynamics,
defective axonal transport of mitochondria, and selective synaptic degeneration in Huntington's
disease. Biochim Biophys Acta 1822, 101-110. 10.1016/j.bbadis.2011.10.016.

Russo, A., and DiAntonio, A. (2019). Wnd/DLK Is a Critical Target of FMRP Responsible for
Neurodevelopmental and Behavior Defects in the Drosophila Model of Fragile X Syndrome. Cell
Rep 28, 2581-2593.e2585. 10.1016/j.celrep.2019.08.001.

Saha, A.R., Hill, J., Utton, M.A., Asuni, A.A., Ackerley, S., Grierson, A.J., Miller, C.C., Davies,
A.M., Buchman, V.L., Anderton, B.H., and Hanger, D.P. (2004). Parkinson's disease alpha-
synuclein mutations exhibit defective axonal transport in cultured neurons. J Cell Sci 117, 1017-
1024. 10.1242/jcs.00967.

Satoh, A.K., O'Tousa, J.E., Ozaki, K., and Ready, D.F. (2005). Rab11 mediates post-Golgi
trafficking of rhodopsin to the photosensitive apical membrane of Drosophila photoreceptors.
Development 132, 1487-1497. 10.1242/dev.01704.

Seong, K.H., Matsuo, T., Fuyama, Y., and Aigaki, T. (2001). Neural-specific overexpression of
drosophila plenty of SH3s (DPOSH) extends the longevity of adult flies. Biogerontology 2, 271-
281. 10.1023/a:1013249326285.

Shin, J.E., Cho, Y., Beirowski, B., Milbrandt, J., Cavalli, V., and DiAntonio, A. (2012). Dual
leucine zipper kinase is required for retrograde injury signaling and axonal regeneration. Neuron
74, 1015-1022. 10.1016/j.neuron.2012.04.028.

Singh, M., and Kim, J.H. (2022). Generation and validation of pX-UASTattB for dose-dependent
misexpression studies in. MicroPubl Biol 2022. 10.17912/micropub.biology.000626.

Singh, M., Ye, B., and Kim, J.H. (2022). Dual Leucine Zipper Kinase Regulates Dscam
Expression through a Noncanonical Function of the Cytoplasmic Poly(A)-Binding Protein. J
Neurosci 42, 6007-6019. 10.1523/JNEUROSCI.0543-21.2022.

Smith, G.A., Rocha, E.M., McLean, J.R., Hayes, M.A., Izen, S.C., Isacson, O., and Hallett, P.J.
(2014). Progressive axonal transport and synaptic protein changes correlate with behavioral and
neuropathological abnormalities in the heterozygous Q175 KI mouse model of Huntington's
disease. Hum Mol Genet 23, 4510-4527. 10.1093/hmg/ddu166.

Stenmark, H. (2009). Rab GTPases as coordinators of vesicle traffic. Nat Rev Mol Cell Biol 10,
513-525. 10.1038/nrm2728.

Stowers, R.S., and Isacoff, E.Y. (2007). Drosophila huntingtin-interacting protein 14 is a
presynaptic protein required for photoreceptor synaptic transmission and expression of the
palmitoylated proteins synaptosome-associated protein 25 and cysteine string protein. J
Neurosci 27, 12874-12883. 10.1523/JNEUROSCI.2464-07.2007.

Taguchi, T. (2013). Emerging roles of recycling endosomes. J Biochem 153, 505-510.
10.1093/jb/mvt034.

Tiwari, A.K., and Roy, J.K. (2009). Mutation in Rab11 results in abnormal organization of
ommatidial cells and activation of JNK signaling in the Drosophila eye. Eur J Cell Biol 88, 445-
460. 10.1016/j.ejcb.2009.02.188.

Tortosa, E., and Hoogenraad, C.C. (2018). Polarized trafficking: the palmitoylation cycle
distributes cytoplasmic proteins to distinct neuronal compartments. Curr Opin Cell Biol 50, 64-
71. 10.1016/j.ceb.2018.02.004.

Tortosa, E., Sengupta Ghosh, A., Li, Q., Wong, W.R., Hinkle, T., Sandoval, W., Rose, C.M., and
Hoogenraad, C.C. (2022). Stress-induced vesicular assemblies of dual leucine zipper kinase
are signaling hubs involved in kinase activation and neurodegeneration. EMBO J 41, e110155.
10.15252/embj.2021110155.

Valakh, V., Frey, E., Babetto, E., Walker, L.J., and DiAntonio, A. (2015). Cytoskeletal disruption
activates the DLK/JNK pathway, which promotes axonal regeneration and mimics a
preconditioning injury. Neurobiol Dis 77, 13-25. 10.1016/j.nbd.2015.02.014.

Valakh, V., Walker, L.J., Skeath, J.B., and DiAntonio, A. (2013). Loss of the spectraplakin short
stop activates the DLK injury response pathway in Drosophila. J Neurosci 33, 17863-17873.
10.1523/JNEUROSCI.2196-13.2013.

Venken, K.J., Schulze, K.L., Haelterman, N.A., Pan, H., He, Y., Evans-Holm, M., Carlson, J.W.,
Levis, R.W., Spradling, A.C., Hoskins, R.A., and Bellen, H.J. (2011). MiMIC: a highly versatile
transposon insertion resource for engineering Drosophila melanogaster genes. Nat Methods 8,
737-743. 10.1038/nmeth.1662.

Wang, X., Kim, J.H., Bazzi, M., Robinson, S., Collins, C.A., and Ye, B. (2013). Bimodal control of
dendritic and axonal growth by the dual leucine zipper kinase pathway. PLoS Biol 11, e1001572.
10.1371/journal.pbio.1001572.

Wang, Z.X., Tan, L., and Yu, J.T. (2015). Axonal transport defects in Alzheimer's disease. Mol
Neurobiol 51, 1309-1321. 10.1007/s12035-014-8810-x.

Watkins, T.A., Wang, B., Huntwork-Rodriguez, S., Yang, J., Jiang, Z., Eastham-Anderson, J.,
Modrusan, Z., Kaminker, J.S., Tessier-Lavigne, M., and Lewcock, J.W. (2013). DLK initiates a
transcriptional program that couples apoptotic and regenerative responses to axonal injury. Proc
Natl Acad Sci U S A 110, 4039-4044. 10.1073/pnas.1211074110.

Welsbie, D.S., Yang, Z., Ge, Y., Mitchell, K.L., Zhou, X., Martin, S.E., Berlinicke, C.A., Hackler,
L., Fuller, J., Fu, J., et al. (2013). Functional genomic screening identifies dual leucine zipper
kinase as a key mediator of retinal ganglion cell death. Proc Natl Acad Sci U S A 110, 4045-
4050. 10.1073/pnas.1211284110.

Welsbie, D.S., Mitchell, K.L., Jaskula-Ranga, V., Sluch, V.M., Yang, Z., Kim, J., Buehler, E.,
Patel, A., Martin, S.E., Zhang, P.W., et al. (2017). Enhanced Functional Genomic Screening
Identifies Novel Mediators of Dual Leucine Zipper Kinase-Dependent Injury Signaling in
Neurons. Neuron 94, 1142-1154.e1146. 10.1016/j.neuron.2017.06.008.

Wu, C., Wairkar, Y.P., Collins, C.A., and DiAntonio, A. (2005). Highwire function at the
Drosophila neuromuscular junction: spatial, structural, and temporal requirements. J Neurosci
25, 9557-9566. 10.1523/JNEUROSCI.2532-05.2005.

Xiong, X., and Collins, C.A. (2012). A conditioning lesion protects axons from degeneration via
the Wallenda/DLK MAP kinase signaling cascade. J Neurosci 32, 610-615.
10.1523/JNEUROSCI.3586-11.2012.

Xiong, X., Hao, Y., Sun, K., Li, J., Li, X., Mishra, B., Soppina, P., Wu, C., Hume, R.I., and Collins,
C.A. (2012). The Highwire ubiquitin ligase promotes axonal degeneration by tuning levels of
Nmnat protein. PLoS Biol 10, e1001440. 10.1371/journal.pbio.1001440.

Xiong, X., Wang, X., Ewanek, R., Bhat, P., Diantonio, A., and Collins, C.A. (2010). Protein
turnover of the Wallenda/DLK kinase regulates a retrograde response to axonal injury. J Cell
Biol 191, 211-223. 10.1083/jcb.201006039.

Ye, B., Zhang, Y., Song, W., Younger, S.H., Jan, L.Y., and Jan, Y.N. (2007). Growing dendrites
and axons differ in their reliance on the secretory pathway. Cell 130, 717-729.
10.1016/j.cell.2007.06.032.

Zhang, J., Schulze, K.L., Hiesinger, P.R., Suyama, K., Wang, S., Fish, M., Acar, M., Hoskins,
R.A., Bellen, H.J., and Scott, M.P. (2007). Thirty-one flavors of Drosophila rab proteins. Genetics
176, 1307-1322. 10.1534/genetics.106.066761.

Zhou, W., Chang, J., Wang, X., Savelieff, M.G., Zhao, Y., Ke, S., and Ye, B. (2014). GM130 is
required for compartmental organization of dendritic golgi outposts. Curr Biol 24, 1227-1233.
10.1016/j.cub.2014.04.008.

Figure 1

Fig 1. Wnd protein turnover by Hiw is restricted in the axon terminals of *Drosophila* sensory neurons.

Wnd is highly enriched in axon terminals (A-D).

(A) A representative image of *Mi{MIC}wnd* [GFSTF] in the neuropil of larval ventral nerve cords (top) and C4da neuron cell body (bottom) from wild-type male (*w[1118]*) or hemizygous male *hiw* mutant (*hiw[ΔN]*) were shown. Wnd-GFP (Wnd-GFSTF, green) and anti-HRP staining (magenta). Scale bar = 10 μm.

(B) Schematic of *Drosophila* larval C4da neurons. C4da cell bodies and dendrites are located in the body wall (green) while their axons (red) terminate in the ventral nerve cord (VNC) collectively forming the ladder-like structure of axon terminals (yellow).

(C) The Wnd::GFP (with or without *wnd*-UTRs, green) along with mCD8::RFP (magenta) were expressed in the larval C4da neurons in the presence of DLK inhibitor, DLKi. The expression levels of transgenes were shown in C4da cell body and the axon terminals.

(D) The axonal enrichment index was expressed as median ± 95% CI. Sample numbers were 5'-Wnd::GFP-3' (*w[1118]; UAS-wnd-5'UTR-Wnd::GFP-wnd-3'UTR/+;ppk-GAL4, UAS-mCD8::RFP/+*) (n = 26), Wnd::GFP (*w[1118]; UAS-Wnd::GFP-SV40 3'UTR /+; ppk-GAL4, UAS-mCD8::RFP/+*) (n = 22), Wnd-KD::GFP (*w[1118]; UAS-Wnd-KD::GFP/+; ppk-GAL4, UAS-mCD8::RFP/+*) (n = 6). All samples were treated with DLKi ; One-way ANOVA ($F_{(2,51)}$ = 0.5176) followed by post hoc Tukey's multiple comparison test.

(E) The Wnd-KD::GFP transgene along with mCD8::RFP was expressed in C4da neurons with the *ppk-Gal4* driver, in wild-type (*wt*) (*w[1118];UAS-Wnd-KD::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*, sample n = 8) or in *hiw* mutant (*hiw[ΔN]; UAS-Wnd-KD::GFP/+;ppk-Gal4,UAS-mCD8::RFP/+*, sample n = 9) larvae. Representative images of Wnd-KD::GFP in C4da cell bodies and axon terminals, stained for anti-GFP (green). The average Wnd-KD::GFP intensity from cell bodies and axon terminals were quantified and expressed as median ± 95% CI; (Cell body: U=25, P=0.3213, Axon terminal: U=0, P<0.0001, two-tailed Mann-Whitney test). Scale bar = 10 μm.

Figure 1 - supplement

Figure 1- Supplement. The axon terminal enrichment of Wnd is independent of Wnd kinase activity.

(A) A Wnd kinase-dead (Wnd-KD::GFP, green) along with mCD8::RFP (magenta) were expressed in the larval C4da neurons in either regular food, food containing DMSO (vehicle control) or DLKi. The expression levels of transgenes were shown in C4da cell body and the axon terminals. Scale bar = 10 μm.

(B) The axonal enrichment index from Figure 1 and supplement 1 were expressed as median ± 95% CI. Genotypes and sample numbers were 5'-Wnd::GFP-3' (*w^{1118} ; UAS-wnd-5'UTR-Wnd::GFP-wnd-3'UTR/+;ppk-GAL4, UAS-mCD8::RFP/+*) with DLKi (n = 26), Wnd::GFP (*w^{1118} ; UAS-Wnd::GFP-SV40 3'UTR /+; ppk-GAL4, UAS-mCD8::RFP/+*) with DLKi (n = 22), Wnd-KD::GFP (*w^{1118} ; UAS-Wnd-KD::GFP/+; ppk-GAL4, UAS-mCD8::RFP/+*) (n = 10), with DLKi (n = 6), with DMSO (n = 6); One-way ANOVA ($F_{(4,65)}$ = 1.017) followed by post hoc Tukey's multiple comparison test.

Figure 2

Figure 2. Wnd palmitoylation is essential for Wnd axonal localization.

(A) Schematic of Wnd deletion mutants. Deletion of the N-terminal 1-100aa of Wnd (WndΔ100), Deletion of the N-terminal 1-200 aa of Wnd (WndΔ200), and Deletion of the N-terminal 101-200 aa of Wnd (WndΔ101-200).

(B) The transgenes of serially deleted Wnd (green) along with mCD8::mRFP were expressed in the larval C4da neurons. Note that WndΔ100 was further mutated to remove kinase activity (Wnd-KDΔ100). The expression levels of Wnd::GFP proteins in C4da cell body and the axon terminals were shown (left). Scale bars = 10 μm. The axonal enrichment of each Wnd transgenes was measured using mCD8::RFP (not shown in figure) as a normalization control and expressed as median ± 95% CI. One-way ANOVA ($F_{(3,59)}$ = 103.8) followed by post hoc Tukey's multiple comparison test. The p values from Tukey's test are indicated in the graph. Genotypes and sample numbers were Wnd-KD (*UAS-Wnd-KD::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*, n=22), Wnd-KDΔ*100* (*UAS-Wnd-KDΔ100::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*, n=18), WndΔ*200* (*UAS-WndΔ200::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*, n=13), and WndΔ*101-200* (*UAS-WndΔ101-200::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*, n=10).

(C) Wnd-C130 is an evolutionarily conserved palmitoylation site. The protein sequence alignment of DLK proteins from the indicated species is shown. The conserved Cysteine residue is shown in red.

(D) GFP-tagged Wnd proteins were expressed in S2 cells before pulled down by GFP nanobody beads. The Acyl Biotin Exchange assay was performed on the beads coupled Wnd proteins. Wnd-C130S::GFP is a palmitoylation defective version of Wnd. "No HAM" serves as a negative control since Biotin exchange does not occur in the absence of HAM. Biotin blot shows protein palmitoylation levels while total Wnd shows Wnd protein levels in the pulldown. The numbers indicate the total intensity of palmitoylation blot that were normalized by GFP blot from Wnd::GFP and Wnd-C130S::GFP.

(E) A wild-type Wnd::GFP transgene (Wnd::GFP) and a palmitoylation-defective Wnd transgene (Wnd-C130S::GFP) along with mCD8::RFP were expressed in the larval C4da neurons in the presence of DLKi. GFP immunostaining from Wnd::GFP proteins in C4da cell body and the axon terminals were shown (top). Scale bar = 10 μm. The axonal enrichment of the Wnd transgenes was measured using mCD8::RFP (not shown in figure) as a normalization control and expressed as median ± 95% CI (U=0, P<0.0001, two-tailed Mann-Whitney test). Genotypes and samples numbers were Wnd::GFP (*UAS-Wnd::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+* , n = 12) and Wnd-C130S::GFP (*UAS-WndC130S::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+* , n=10).

Figure 2 – supplement 1

Figure 2 - Supplement 1. Wnd palmitoylation is not required for Wnd localization in somatic Golgi complex.

Wnd-C130S::GFP (green) along with a Golgi marker, ManII-Tag::RFP (magenta) were expressed in the larval C4da neurons. The expression patterns of the two transgenes in C4da cell bodies were shown (left). Scale bar = 10 µm. The Manders' overlap coefficient between Wnd transgenes and ManII-Tag::RFP was measured and expressed as median ± 95% CI (right). The Genotype and sample numbers were Wnd-C130S::GFP (*w1118; UAS-Wnd-C130S-KD::GFP/+;ppk-GAL4/UAS-ManII-Tag-RFP*, n=22).

Figure 2 - Supplement 2. Somatic Golgi localization is necessary for Wnd localization in axon terminals.

Wnd transgenes (green) along with a Golgi marker, ManII-Tag::RFP (magenta) were expressed in the larval C4da neurons. The expression patterns of the two transgenes in C4da cell bodies were shown (left). Scale bar = 5 μm. The Manders' overlap coefficient between Wnd transgenes and ManII-Tag::RFP was measured and expressed as median ± 95% CI (right). Genotypes and sample numbers were Wnd-KD::GFP (*w1118; UAS-Wnd-KD::GFP/+;ppk-GAL4/UAS-ManII-Tag-RFP*, n = 23), Wnd-KDΔ100::GFP (*w1118; UAS-Wnd-Δ100-KD::GFP/+;ppk-GAL4/UAS-ManII-Tag-RFP*, n=26), Wnd-Δ200::GFP (*w1118;UAS-Wnd-Δ200::GFP/+;ppk-GAL4/UAS-ManII-Tag-RFP*, n=28) and Wnd-Δ101-200::GFP (*w1118;UAS-WndΔ101-200::GFP/+;ppk-GAL4/UAS-ManII-Tag-RFP*, n = 15). One-way ANOVA ($F_{(3,88)}$ = 129.8) followed by post hoc Tukey's multiple comparison test. The p values from Tukey's test are indicated in the graph.

Figure 2 – supplement 3

Figure 2 - Supplement 3. dHIP14 is required for Wnd localization in axon terminals.

(A) A Golgi marker, ManII::GFP (green) was expressed in the larval C4da neurons along with dHIP14::Tdtomato (magenta). The image was taken from a C4da cell body. Scale bar = 10 μm.
(B) Wnd-KD::GFP (green) was expressed in the larval C4da neurons along with dHIP14::Tdtomato (magenta). The image was taken from a C4da cell body. Scale bar = 10 μm.

(C) Wnd-KD::GFP (green) along with mCD8::RFP (magenta) were expressed in the larval C4da neurons from wild-type (*wt*) or homozygous *dHIP14* mutants (*dHIP14^{ex11}*). Scale bar = 10μm. The axonal enrichment of Wnd-KD::GFP was measured and expressed as a median ± 95% CI (Mann-Whitney U=13, P<0.0001, two-tailed Mann-Whitney test). Genotypes and sample numbers were *wt* (*UAS-Wnd-KD::GFP/ppk-Gal4, U-mCD8::RFP;+/+*, n=17) and *dHIP14^{ex11}*(*UAS-Wnd-KD::GFP/ppk-Gal4, U-mCD8::RFP;dHIP14^{ex11}*, n=16).

Figure 3

Figure 3. Increased protein expression levels of palmitoylation-defective Wnd.

(A) Wnd::GFP and Wnd-C130S::GFP were expressed in the larval nervous system using a pan-neuronal driver, *Elav*-GAL4 in the presence of DLKi. Larval brain lysates were subjected to Western blot analysis using a GFP antibody. Elav blot was used as a loading control. The GFP blots were normalized by Elav blots and expressed as mean ± SEM (n = 4, t = 9.409, df =3, p=0.0025, two-tailed ratio paired t-test). Genotypes were Wnd::GFP (*elav-Gal4/+; UAS-Wnd::GFP/+;*) and WndC130S::GFP (*elav-Gal4/+; UAS-WndC130S::GFP/+;*).

(B) Wnd transgenes (green) along with mCD8::RFP were expressed in the larval C4da neurons to directly compare the expression levels of Wnd::GFP and Wnd-C130S::GFP in C4da cell bodies and axon terminals. The larvae were treated with DLKi. Scale bar = 10 μm. The average fluorescent intensities from GFP staining of each larva sample (C4da cell bodies and axon terminals) were measured and expressed as median ± 95% CI (Axon terminal: Mann-Whitney U=2, P=0.0162, two-tailed Mann-Whitney test, Cell body: Mann-Whitney U=0, P<0.0001, two-tailed Mann-Whitney test). Genotypes and sample numbers were Wnd::GFP (*UAS-Wnd::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*, n = 8) and Wnd-C130S::GFP (*UAS-WndC130S::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*, n = 4).

Figure 4

A

B

C

Figure 4. **Palmitoylation-defective Wnd caused exacerbated stress responses.**

(A) *puc*-LacZ were expressed along with *2XUAS-Wnd::GFP* and *2XUAS-Wnd-C130S::GFP* in the larval C4da neurons. The average fluorescent intensities from *puc*-LacZ staining of each larva sample in C4da cell body were measured and expressed as median ± 95% CI (Mann-Whitney U=56, P<0.0001, two-tailed Mann-Whitney test). Genotypes and sample numbers were 2XUAS-Wnd::GFP (*2XUAS-Wnd::GFP/ppk-Gal4,UAS-mCD8::RFP ;puc-LacZ/+*, n = 28) and 2XUAS-Wnd-C130S::GFP (*2XUAS-Wnd-C130S::GFP/ ppk-Gal4,UAS-mCD8::RFP ;puc-LacZ/+*, n = 28) Scale bar = 10 µm.

(B) *2XUAS-Wnd::GFP* and *2XUAS-Wnd-C130S::GFP* were expressed under an eye specific driver, *GMR-GAL4*. Adult fly lethality was scored and analyzed using the Chi-square (df =

41.57,1, z= 6.448, p<0.0001, two-sided contingency test). Genotypes and sample numbers were Wnd::GFP (*2XUAS-Wnd::GFP/GMR-Gal4*, n = 188), Wnd-C130S::GFP (*2XUAS-Wnd-C130S::GFP/GMR-Gal4*, n = 195).

(C) Representative images of fly eyes from Wnd::GFP (*2XUAS-Wnd::GFP/+*), Wnd-C130S::GFP (*2XUAS-Wnd-C130S::GFP/+*), Wnd::GFP under GMR-GAL4 (*2XUAS-Wnd::GFP/GMR-Gal4*), Wnd-C130S::GFP under GMR-GAL4 (*2XUAS-Wnd-C130S::GFP/GMR-Gal4*).

Figure 4 – supplement 1

Figure 4 - Supplement 1. Wnd-C130S::GFP retains Wnd signaling capacity.

(A) *puc*-LacZ were expressed along with *5XUAS-Wnd-KD::GFP, 5XUAS-Wnd::GFP* and *5XUAS-Wnd-C130S::GFP* in the larval C4da neurons. The average fluorescent intensities from *puc*-LacZ staining of each larva sample in C4da cell body were measured and expressed as median ± 95% CI (One-way ANOVA, F (2, 114) = 108.3). Genotypes and sample numbers were 5XUAS-Wnd-KD::GFP (*;ppk-Gal4,UAS-mCD8::RFP/5XUAS-Wnd-KD::GFP;puc-LacZ/+*, n = 42), 5XUAS-Wnd::GFP (*;ppk-Gal4,UAS-mCD8::RFP/5XUAS-Wnd::GFP;puc-LacZ/+*, n = 49), and 5XUAS-Wnd-C130S::GFP (*;ppk-Gal4,UAS-mCD8::RFP/5XUAS-Wnd-C130S::GFP;puc-LacZ/+*, n = 26) Scale bar = 10 μm.

(B) Wnd transgenes along with mCD8::RFP were expressed in the larval C4da neurons using *ppk-GAL4*. The axon terminal morphology was visualized by mCD8::RFP. Wnd-KD::GFP (*UAS-Wnd-KD::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*), Wnd::GFP(*UAS-Wnd::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*), WndC130S::GFP (*UAS-WndC130S::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*) and WndC130S-KD::GFP (*UAS-WndC130S-KD::GFP/+;ppk-Gal4, UAS-mCD8::RFP/+*). Scale bar = 10 μm.

Figure 4 – supplement 2

Figure 4 - Supplement 2. Wnd-C130S::GFP exhibits higher toxicity levels than Wnd::GFP.

(A) Representative images of fly eyes from Wnd::GFP under GMR-GAL4 (*GMR>Wnd::GFP*), Wnd-C130S::GFP under GMR-GAL4 (*GMR>Wnd-C130::GFP*), and Wnd-KD::GFP under GMR-GAL4 (*GMR>Wnd-KD::GFP*). Flies were reared at 18˚C.

(B) Eye area was quantified and expressed as median ± 95% CI. One-way ANOVA ($F(6, 136) = 362.0$) followed by post hoc Tukey's multiple comparison test (****, $P < 0.0001$). Genotypes and sample numbers were GMR (*GMR-GAL4/+*, n = 21), Wnd::GFP (*UAS-Wnd::GFP/+*, n = 22), Wnd-C130S::GFP (*UAS-Wnd-C130S::GFP/+*, n = 20), Wnd-KD::GFP (*UAS-Wnd-KD::GFP/+*, n = 24), GMR>Wnd::GFP (*GMR-GAL4/UAS-Wnd::GFP*, n = 17), GMR>Wnd-C130S::GFP (*GMR-GAL4/UAS-Wnd-C130S::GFP*, n = 16), and GMR>Wnd- KD::GFP (*GMR-GAL4/UAS-Wnd-KD::GFP*, n = 23). Note more severe eye phenotypes in GMR>Wnd-C130S::GFP as compared to GMR>Wnd::GFP.

Figure 5

Figure 5. Screening of Dominant negative Rab proteins for Wnd axonal enrichment.

(A) YFP-tagged dominant negative-Rab (DN-Rab) transgenes along with Wnd-KD::mRFP were expressed in the larval C4da neurons. The axonal enrichment of Wnd-KD::mRFP was measured by Wnd-KD::mRFP expression levels from C4da cell bodies and axon terminals and expressed as mean ± sem of a fold change of control (mCD8::GFP) (n ≥10). One way ANOVA followed by post hoc Dunnett test ($F_{(27,284)}$ = 7.146). A mCD8::GFP transgene was used as a transgene control (*UAS-Wnd-KD::mRFP/+;ppk-Gal4, UAS-mCD8::GFP/+*). Note that YFP::DN-Rab1 and YFP::DN-Rab11 reduced Wnd axonal enrichment by more than 50% compared to the control (gray bar).

(B) Representative images of Wnd-KD::mRFP expression in the axon terminals and cell bodies from the C4da neurons that express mCD8::GFP, DN-Rab1, and DN-Rab11 were shown. The Wnd axonal enrichment was expressed as median ± 95% CI (Mann-Whitney U=0, P < 0.0001 for DN-Rab1, U=13, P<0.0001 for DN-Rab11, two-tailed Mann-Whitney test). Genotypes and sample numbers were mCD8::GFP (*UAS-Wnd-KD::mRFP/+;ppk-Gal4, UAS-mCD8::GFP/+*, n= 19) DN-Rab1 (*UAS-Wnd-KD::mRFP/+;ppk-Gal4/ UAST-YFP.Rab1 S25N*, n=12) and DN-Rab11 (*UAS-Wnd-KD::mRFP/ UASp-YFP.Rab11 S25N; ppk-Gal4/+*, n=10) Scale bar = 10 μm.

Figure 6

Figure 6. Rab11 is necessary for Wnd localization in axon terminals and Wnd protein turnover.

(A) and (B). Wnd localizes at the Rab11-positive endosomes in C4da cell bodies.

Wnd-KD::mRFP was expressed in C4da neurons of *TI-YFP::Rab11* larvae (A). *TI-YFP::Rab11* is endogenously YFP-(Yellow fluorescent protein) tagged *Rab11* allele. Wnd-KD::mRFP was expressed in C4da neurons along with YFP::Rab11 (B). Wnd-KD::mRFP (magenta) and YFP::Rab11 (green) were visualized in C4da cell bodies using an RFP and GFP antibodies, respectively. Genotypes and sample numbers were *TI-YFP::Rab11* (*UAS-Wnd-KD::mRFP/+;TI{TI}Rab11[EYFP]/ppk-Gal4,* n=28) and *YFP::Rab11* (*UAS-Wnd-KD::mRFP/UAS-YFP::Rab11;ppk-Gal4/+*; , n=30). Scale bar = 10 μm.

(C) Wnd transgenes (magenta) along with mCD8::GFP or YFP::DN-Rab11 were expressed in the larval C4da neurons to directly compare the expression levels of Wnd-KD::mRFP in C4da cell bodies and axon terminals. Scale bar = 10 μm. The average fluorescent intensities from mRFP staining of each larva sample were measured and expressed as median ± 95% CI (Cell body: Mann-Whitney U=0, P <0.0001, two-tailed Mann-Whitney test; Axon terminal: Mann-Whitney U=12, P=0.0160, two-tailed Mann-Whitney test). Genotypes and sample numbers were *wt* (*UAS-Wnd-KD::mRFP/+;ppk-Gal4, UAS-mCD8::GFP/+,* n = 9) and *YFP::DN-Rab11* (*UAS-Wnd-KD::mRFP/UASp-YFP.Rab11.S25N ;ppk-Gal4, UAS-mCD8::RFP/+*, n = 9).

(D) Wnd-KD::GFP (green) along with mCD8::RFP (magenta) were expressed in the larval C4da neurons from wild-type (*wt*) or from homozygous *Rab11* mutants (*Rab11[93bi]*). Images from C4da axon terminals and cell bodies were shown. Scale bar = 10 μm. The axonal enrichment of Wnd-KD::GFP was measured using mCD8::RFP as a normalization control and expressed as median ± 95% CI (Mann-Whitney U=0, P<0.0001, two-tailed Mann-Whitney test). Genotypes and sample numbers were *wt* (*UAS-Wnd-KD::GFP/ppk-Gal4, U-mCD8::RFP*, n=17) and *Rab11[93bi]* (*UAS-Wnd-KD::GFP/ppk-Gal4, U-mCD8::RFP; Rab11[93bi]*, n=12).

(E) Western blot was performed on the larval brain lysates from wild-type (*wt, w[1118]*) or homozygous *Rab11[93bi]* using Wnd antibody to measure total Wnd protein levels. Elav blot was used as a loading control. The Wnd blots were normalized by Elav blots and expressed as mean ± SEM (n = 3, t = 6.06, df =2, p=0.026, two-tailed ratio paired t-test).

Figure 6 - Supplement 1. Wnd does not localize in the Rab11-positive endosomes in axons and axon terminals.

(A) Wnd-KD::mRFP and YFP::Rab11 were expressed in the larval C4da neurons using *ppk-GAL4*. Wnd-KD::mRFP (magenta) and YFP::Rab11 (green) were visualized in the segmental nerve/axons (top) and in C4da axon terminals (bottom) using RFP and GFP antibodies, respectively. Scale bar = 10 µm.

(B) The Manders' overlap coefficient between Wnd-KD::mRFP and YFP::Rab11 was measured in cell bodies, axons (segmental nerve), and axon terminals. They are expressed as median ± 95% CI. One-way ANOVA (F(2, 62) = 109.8) followed by post hoc Tukey's multiple comparison test. Sample numbers were; Cell body n=30, axons n = 14, neuropils n=21.

Figure 6 – supplement 2

Figure 6 - Supplement 2. *Rab11* mutations did not increase *wnd* mRNA abundance.

Total RNAs extracted from the wondering third instar larval brains from a wild-type (*wt*) control (*w^1118^*) and *Rab11^93bi^* were subjected to RT-qPCR to measure *wnd* mRNA abundance. The *wnd* mRNA levels were normalized by *GAPDH1* mRNA levels via the 2ΔΔCT method. Two independent wnd primer sets were used (Primer-1 and Primer-2). The data were expressed as mean ± SEM of five technical replicates. Welch's t-test (Primer-1: t=8.778, df=4.772, Primer-2: t=6.576, df=7.875).

Figure 7

Figure 7. Wnd mediates stress signaling induced by Rab11 loss-of-function.

(A) *puc*-LacZ were expressed along with YFP::DN-Rab11 and mCD8::GFP in the larval C4da neurons in the presence of DMSO (vehicle) or DLK inhibitor (DLKi). The average fluorescent intensities from LacZ staining of each larva sample in C4da cell bodies were measured and quantified as median ± 95% CI. One-way ANOVA ($F_{(3,171)}$ = 11.48) followed by post hoc Tukey's multiple comparison test (****, $P < 0.0001$). Genotypes and sample numbers were *wt (; ppk-Gal4, UAS-mCD8::RFP/+;puc-LacZ/+*, in DMSO, n = 36, in DLKi n=42) and DN-Rab11 (*; ppk-Gal4, UAS-mCD8::RFP/+;puc-LacZ/UASp-YFP.Rab11.S25N*, in DMSO, n = 51, in DLKi, n=46) Scale bar = 10 μm.

(B) Representative images of fly eyes from *Elav-GAL4* only, *DN-Rab11* only, and *DN-Rab11* expression under a pan-neuronal driver, *Elav-GAL4 (Elav-GAL4/DN-Rab11)*. Flies were reared in the presence of DMSO and DLKi (down). Eye area was quantified and expressed as median ± 95% CI. One-way ANOVA ($F_{(5,114)}$ = 290.3) followed by post hoc Tukey's multiple comparison test (****, $P < 0.0001$). Genotypes and sample numbers were Elav-Gal4 (*Elav-GAL4/+*, n = 20), DN-Rab11 (*;;UASp-YFP.Rab11.S25N/+*, n = 20), Elav-Gal4/DN-Rab11 (*ElavGal4/+;;UASp-YFP.Rab11.S25N /+*, n = 20).

(C) Representative images of fly eyes from *Elav-GAL4* only, *DN-Rab11* only in heterozygous *wnd* mutant (*DN-Rab11, wnd[3]*), DN-Rab11 expression under a pan-neuronal driver, *Elav-GAL4* (*Elav-GAL4/DN-Rab11*)*, and DN-Rab11 expression under a pan-neuronal driver in heterozygous *wnd* mutant (*Elav-GAL4/DN-Rab11, wnd[3]*). Eye area was quantified and expressed as median ± 95% CI. One-way ANOVA ($F_{(4,95)}$ = 530.4) followed by post hoc Tukey's multiple comparison test.

Figure 8

Figure 8. Rab11 suppresses neuronal stress signaling by localizing Dual leucine zipper kinase to axon terminals for protein turnover.

Under normal conditions, the newly synthesized Wnd proteins are recruited to the Golgi apparatus in neuronal cell body. Wnd is palmitoylated at somatic Golgi apparatus, which enables it to be sorted into the Recycling Endosomes via Rab11. Wnd is actively transported out of neuronal cell bodies towards axon terminals where Hiw-mediated degradation of Wnd occurs. When Rab11 function is lost, Wnd proteins cannot be transported towards axon terminals. This prevents Wnd protein turnover resulting in higher Wnd protein expression, which in turn triggers cell death pathway.

Table 1

REAGENT or RESOURCE	SOURCE	RRID
UAST-YFP.Rab1.S25N	Zhang et al., 2007	BDSC_9757
UAST-YFP.Rab2.S20N	Zhang et al., 2007	BDSC_23640
UASp-YFP.RabX2.S21N	Zhang et al., 2007	BDSC_9843
UASp-YFP.Rab3.T35N	Zhang et al., 2007	BDSC_9766
UASp-YFP.Rab4.S22N	Zhang et al., 2007	BDSC_9769
UAST-YFP.RabX4.T40N	Zhang et al., 2007	BDSC_9849
UASp-YFP.Rab5.S43N	Zhang et al., 2007	BDSC_9771
UAST-YFP.Rab6.T26N	Zhang et al., 2007	BDSC_23249
UAST-YFP.RabX6.S22N	Zhang et al., 2007	BDSC_9856
UASp-YFP.Rab7.T22N	Zhang et al., 2007	BDSC_9778
UASp-YFP.Rab8.T22N	Zhang et al., 2007	BDSC_9780
UASp-YFP.Rab9.S26N	Zhang et al., 2007	BDSC_23642
UASp-YFP.Rab10.T23N	Zhang et al., 2007	BDSC_9786
UASp-YFP.Rab11.S25N	Zhang et al., 2007	BDSC_9792
UAST-YFP.Rab18.S19N	Zhang et al., 2007	BDSC_23238
UAST-YFP.Rab19.T35N	Zhang et al., 2007	BDSC_9799
UAST-YFP.Rab21.T27N	Zhang et al., 2007	BDSC_23241
UASp-YFP.Rab23.S51N	Zhang et al., 2007	BDSC_9804
UAST-YFP.Rab26.T204N	Zhang et al., 2007	BDSC_9807
UASp-YFP.Rab27.T25N	Zhang et al., 2007	BDSC_23267
UASp-YFP.Rab32.T33N	Zhang et al., 2007	BDSC_23281
UAST-YFP.Rab35.S22N	Zhang et al., 2007	BDSC_9819
UAST-YFP.Rab39.S23N	Zhang et al., 2007	BDSC_23247
UAST-YFP.Rab40.H25N	Zhang et al., 2007	BDSC_9829
UAST-YFP.CG9807.T21N	Zhang et al., 2007	BDSC_23257
UAST-YFP.CG9807.T21N	Zhang et al., 2007	BDSC_23258
UAST-YFP.CG32673.T21N	Zhang et al., 2007	BDSC_23254
UASp-YFP.Rab11	Zhang et al., 2007	BDSC_50782
TI{TI}Rab11[EYFP]	Dunst et al., 2015	BDSC_62549
Rab11[93Bi]	Giansanti et al., 2007	BDSC_4158