Chronic exposure of mosquito larvae to pesticide residues endangers a new generation of

agrochemicals repurposed for malaria prevention

Marilene M. Ambadiang[1, 2], Caroline Fouet[3], Fred A. Ashu[1, 2], Véronique Penlap-Beng[2], and

Colince Kamdem[3]*

[1]Centre for Research in Infectious Diseases (CRID), P.O. Box 13591, Yaoundé, 9 Cameroon

[2]Department of Biochemistry, Faculty of Science, University of Yaoundé 1, P.O. Box 11 812,

Yaoundé, Cameroon

[3]Department of Biological Sciences, The University of Texas at El Paso, 500 W. University

Ave. El Paso, TX 79968

* Correspondence: gckamdemdjo@utep.edu

Abstract

Agrochemicals have been successfully repurposed to control mosquitoes worldwide, but

pesticides used in agriculture challenge their effectiveness by contaminating surface waters

and helping larval populations develop resistance. Thus, knowledge of the lethal and sublethal

effects of residual pesticide exposure on mosquitoes is critical for selecting effective

insecticides. Here we implemented a new experimental approach to predict the efficacy of

agricultural pesticides newly repurposed for malaria vector control. We mimicked insecticide

resistance selection as it occurs in contaminated aquatic habitats by rearing field-collected

mosquito larvae in water containing a dose of insecticide capable of killing individuals from a

susceptible strain within 24 h. We then simultaneously monitored short-term lethal toxicity

within 24 h and sublethal effects for 7 days. We found that due to chronic exposure to

agricultural pesticides, some mosquito populations are currently pre-adapt to resist

neonicotinoids if those were used in vector control. Larvae collected from rural and

agricultural areas where neonicotinoid formulations are intensively used for insect pest

management were able to survive, grow, pupate and emerge in water containing a lethal dose

of acetamiprid, imidacloprid or clothianidin. These results emphasize the importance of

addressing exposure of larval populations to formulations applied in agriculture prior to using

agrochemicals against malaria vectors.

Keywords: pesticide, neonicotinoids, *Anopheles*, malaria, urbanization.

1. Introduction

Prevention through chemical control of vector populations has lead to a drastic reduction in

malaria burden in sub-Saharan Africa over the last two decades (World Health Organization,

2022a). The prevention strategy relies heavily on repurposing agrochemicals, which provides

a rapid and cost-effective approach to identify new active ingredients against vector

populations (Hemingway, 2017; Hoppé et al., 2016). In order to effectively control *Anopheles*

mosquito populations that have become resistant to available insecticides, dozens of

agrochemicals have recently been tested against adult populations, and some promising

candidates have been identified (Hoppé et al., 2016; Lees et al., 2019; N'Guessan et al., 2007;

Oxborough et al., 2010; Portwood et al., 2022). However, concomitant use of the same class

of agrochemicals for crop protection can provide sufficient selection pressure to engineer

resistant mosquito populations before any application of the pesticide in public health

(Georghiou, 1972; Georghiou et al., 1973; Lines, 1988). For example, prior to the scale up of

pyrethroid-impregnated bed nets across the continent, resistance driven by crop protection

spraying was already detected in some *Anopheles* populations in West Africa (Curtis et al.,

1998). There is a concern history may repeat itself with the new generation of agrochemicals

that are being adopted for malaria vector control (World Health Organization, 2023).

Africa's pesticide use for agriculture stood at nearly 108 thousand metric tons in 2019

(Kramer, 2022). Lax regulations allow the use of hundreds of formulations for agricultural

pest management in some African countries (Akpesse, 2018; Katambo, 2018; Ngamo Tinkeu,

2018; Okolle et al., 2022; Urio et al., 2022). A large part of pesticide application occurs in

small farms held by non-professional operators that are not trained and not supervised for safe

and effective use of chemicals. As a result, too frequent and untargeted application provides

suitable conditions for chronic exposure of mosquito larvae to pesticide residues. In tropical

regions, rain and human activities create ponds in farmlands, which become ideal breeding

sites for the most dangerous malaria mosquito species such as *Anopheles gambiae* and *An.*

coluzzii in agricultural areas (Antonio-Nkondjio et al., 2011; Mattah et al., 2017; Zogo et al.,

2019). When these breeding sites are contaminated with pesticides, chronic residual exposure

pre-adapts larval populations to resist synthetic chemicals (Lines, 1988). The consequence of

this adaptation is a decrease in efficacy of agrochemicals, which in turn undermines their

suitability for mosquito control.

Neonicotinoids are among the most widely used pesticides in agriculture worldwide (Matsuda

et al., 2020; Simon-delso et al., 2015). These chemicals are highly water-soluble and

environmentally persistent, and thus may leach into surface waters and become a risk for

aquatic insects (Morrissey et al., 2015; Ramadevi et al., 2022; Schaafsma et al., 2015). Two

new formulations of clothianidin, a neonicotinoid, were recently approved for indoor residual

spraying targeting malaria vectors, but have yet to be applied in endemic countries (Agossa et

al., 2018; Kweka et al., 2018; Uragayala et al., 2018; World Health Organization, 2023).

Additionally, clothianidin is normally not registered for crop protection in most African

countries (Katambo, 2018; Ngamo Tinkeu, 2018). So far, agricultural spraying of other

registered neonicotinoids such as acetamiprid, imidacloprid, thiamethoxam and thiacloprid

remain the main source of selection pressure that could drive resistance to clothianidin in

Anopheles mosquitoes through cross-resistance mechanisms. This hypothesis has yet to be

supported by empirical evidence, but observational studies have described suboptimal levels

of neonicotinoid susceptibility in adults of at least two *Anopheles* species, populations from

agricultural areas being the most tolerant (Ashu et al., 2023a; Fouet et al., 2020; Mouhamadou

et al., 2019). Importantly, there is evidence that neonicotinoids used for crop protection can

confer cross-resistance to clothianidin, raising concerns about the efficacy the new

agrochemicals repurposed for malaria prevention (Ashu et al., 2023a, 2023b; Fouet et al.,

2020).

Large African cities are the venue of dynamic coevolution between humans and disease

vectors, which has sometimes reveled textbook examples of rapid adaptation of mosquitoes to

synthetic chemicals (Antonio-Nkondjio et al., 2015; Kamdem et al., 2017, 2012b). In

Yaoundé the capital of Cameroon, agricultural activities associated with intensive use of

pesticides are pervasive in suburban and rural settings in the outskirts of the city. Mixtures

containing neonicotinoids such as acetamiprid, imidacloprid, and thiamethoxam, as well as

pyrethroids and fungicides are frequently sprayed on diverse crops, creating ideal conditions

for contamination of surface waters and resistance selection (Ngamo Tinkeu, 2018; Okolle et

al., 2022). By contrast, the center of the city provides an island where surface waters are less

exposed to pesticide contamination and aquatic species are therefore less likely to develop

resistance. Indeed, a survey of susceptibility in adult mosquitoes using standard testing

procedures confirmed this prediction. *An. coluzzii*, the only species found in densely

urbanized areas of Yaoundé remains susceptible to neonicotinoids while its sibling species

An. gambiae, which occurs in the countryside, has developed resistance (Ashu et al., 2023a,

2023b; Fouet et al., 2020). This geographic area thus provides a unique setting to dissect the

environmental drivers of neonicotinoid resistance in *Anopheles* mosquitoes.

The objective of the present study was to assess the interplay between residual pesticide

exposure and the development of resistance to new insecticides in malaria vectors. It has been

known for nearly a half century that exposure to agricultural pesticides can lead to the spread

of resistance to public health insecticides in *Anopheles* mosquitoes (Georghiou, 1972;

Georghiou et al., 1973). However, we still lack a holistic approach that could allow to

simultaneously deciphering the lethal (mortality) and sub-lethal effects (survival, growth,

emergence, reproduction, swimming behavior, mobility etc.) which contribute to resistance

selection in agricultural settings. To fill this gap, we implemented a bioassay protocol where

we mimicked pesticide exposure in the wild by rearing filed-collected *Anopheles* larvae in

water containing field-realistic doses of chemicals. Using larval populations collected from

urban and rural areas of Yaoundé, we compared mortality within 24 h as well as some life

table parameters in order to infer the role of neonicotinoid contaminants in resistance

selection. Our findings indicated that resistance selection in *An. gambiae* is due to chronic

exposure to neonicotinoids in rural and agricultural settings and translates into low mortality

but also high survival, growth, pupation and emergence in lethal concentrations of pesticides.

The present study highlights concerns regarding the efficacy of some agrochemicals

repurposed for malaria vector control, especially active ingredients that are concomitantly

used in agricultural pest management.

2. Material and methods

2.1 Study sites

The study was carried out in Yaoundé, the capital of Cameroon, and some of its neighboring

rural   areas.   Approval   to   conduct   a   study   in   the   Center   region   (N°:   1-

140/L/MINSANTE/SG/RDPH-Ce), ethical clearance (N°: 1-141/CRERSH/2020) and

research permit (N°: 000133/MINRESI/B00/C00/C10/C13) were granted by the ministry of

public health and the ministry of scientific research and innovation of Cameroon. We

surveyed four sites which included a farming area located in the suburbs (Nkolondom,

3°56'43" N, 11°3'01" E), two densely urbanized neighborhoods (Etoa Meki, 3°52'53" N,

11°31'40" E and Combattant, 3°52'53" N, 11°31'40" E) and another suburban site

(Nkolnkoumou, 3°52'29" N, 11°23'2" E) (Figure 1). The average distance between sites was

4-5 km. The most abundant malaria vector populations in this area belong to the species

complex *An. gambiae sensu lato (s.l.)* (Antonio-Nkondjio et al., 2011; Kamdem et al., 2012a,

2012b). Previous surveys have revealed that the nominal species *An. gambiae sensu stricto*

(hereafter *An. gambiae*) is the only member of the complex present in the agricultural site

Nkolondom while its sibling species *An. coluzzii* prefers the most urbanized areas of Yaoundé

(Bamou et al., 2019; Kamdem et al., 2012b; Tene-Fossog et al., 2022). Larvae from the semi-

rural neighborhood, Nkolnkoumou, are a mixture of ~ 80% *An. gambiae* and 20% *An. coluzzii*

(Ashu et al., 2023a, 2023b; Fouet et al., 2020). In rural and urban settings, larvae thrive in

rain-dependent puddles, which may contain high concentrations of organic pollutants in some

densely urbanized areas of the city (Antonio-Nkondjio et al., 2011). In the agricultural site

Nkolondom, *An. gambiae* larvae occur in waters that accumulate between furrows and ridges

in farms due to rain and/or irrigation and are likely contaminated with pesticide residues

(Figure 1).

2.2 Larval collection and insecticides

Typical *Anopheles gambiae s.l.* breeding sites were inspected during rainy seasons in 2022 to

collect larvae using the standard dipping method (Service, 1993). Larvae were transported in

plastic containers to the insectary where they were immediately tested in a controlled room

(27°C, 80% relative humidity, light:dark = 12:12 h).

Three neonicotinoids were tested in this study: in addition to acetamiprid and imidacloprid

that are used in local farming activities, we also tested clothianidin, which remains an exotic

insecticide in Cameroon. Deltamethrin, a first-generation malaria-control insecticide, was

used to compare the efficacy of clothianidin. The following commercial formulations were

tested: acetamiprid (Aceplant 40EC, 40g/L, emulsifiable concentrate, JACO, Yaoundé,

Cameroon), imidacloprid (Plantima 30SC, 30g/L, concentrated suspension, JACO, Yaoundé,

Cameroon), clothianidin (Pestanal^{TM}, analytical standard, Sigma-Aldrich) and deltamethrin

(Decis 25EC, 25 g/L, emulsifiable concentrate, Bayer Cropscience SL, Valencia, Spain).

Stock solutions were prepared by diluting the formulation in absolute ethanol or distilled

water.

2.3 Larval bioassays

In order to assess the lethal and sublethal effects of each insecticide, we reared larvae

collected from the field in water containing a dose of the insecticide that was lethal to larvae

of a susceptible strain and we measured mortality as well as some life history parameters. The

mosquito life cycle comprises four larval stages that precede pupation and emergence of a

male or female adult (Service, 1993). The four stages include the 1st (L1), 2nd (L2), 3rd (L3)

and 4th (L4) instars and can last between one and two weeks depending on the mosquito

species, the feeding regime and the environmental conditions. We monitored mortality and

life table parameters starting from 3rd instars until emergence. 3rd instar larvae of *Anopheles*

are abundant in the wild, are easy to identify morphologically and typically complete

transformation into L4, pupation and emergence in approximately 7 days.

The doses tested for each insecticide were determined as follows: we exposed 3rd instar larvae

of the susceptible strain *An. gambiae* Kisumu to increasing concentrations of the insecticide,

starting from 0.001 mg/L and we selected the lowest concentration at which 100% of

individuals died within 24 h. Batches of 25 larvae were placed in 500 mL plastic trays filled

with 200 mL borehole water containing the desired concentration of insecticide and covered

with a net. Four replicates were tested for each insecticide dose in addition to a control

without insecticide (i.e. containing only water). The lowest lethal concentrations were 0.035

mg/L clothianidin, 0.075 mg/L imidacloprid, 0.15 mg/L acetamiprid and 1.5 mg/L

deltamethrin. According to studies conducted worldwide, the concentrations of neonicotinoids

tested fell in the upper limits of average values detected in contaminated waters and soils

(Morrissey et al., 2015; Ramadevi et al., 2022; Schaafsma et al., 2015). However, in some

agricultural areas, these concentrations can be 100 times higher (Ramadevi et al., 2022).

Larvae were collected from the field in the morning and brought to the insectary. 3rd instars

were sorted immediately and rinse in a tray containing borehole before being transferred into

test trays. Four replicates of 25 larvae were tested in 500 mL plastic trays that were filled with

200 mL borehole water containing the above-determined lethal dose of the insecticide. Water

was not changed throughout the experiment, and 10 mg of TetraMin® fish food was added to

each tray daily. Every 24 h, the number of L3, L4, pupae and adults was counted in each tray.

Larvae were considered dead if they were unable to move when touched with a dropper. Dead

larvae were removed from the test containers and were not replaced. Adults were also

removed using a mouth aspirator. Survival, growth, pupation and emergence were assessed

daily for 7 days, which was sufficient for 3rd instar larvae to reach the adult stage.

2.4 Data analysis

The lethal endpoint of insecticide exposure was assessed using the mortality rate at 24 h. The

sublethal effects of the different insecticides were addressed using survival probability as well

as L4 transformation rate, pupation rate and emergence rate. L4 transformation rate was

defined as the percentage of L3 larvae transformed into L4. Pupation rate represented the

percentage of L3 that reached the pupa stage and emergence rate the percentage of L3 that

became adults. Line plots of mean and standard error, computed with the packages *plyr* and

ggplot2 in R V. 4.2.2, were used to compare L4 transformation rate, pupation rate and

emergence rate at 24 h intervals. The ratio between the number of dead larvae and the initial

number of larvae was calculated every 24 h and provided an estimate of survival probability.

Kaplan Meier survival curves were plotted using the packages *ggplot 2* and *ggfortify* in R (R

Core Team, 2016). Larvae reaching the adult stage before the end of the test were treated as

censored data. Confidence intervals were computed for the four replicates, and a log-rank test

was used to determine if survival was significantly different between treatments.

3. Results

3.1 Lethal toxicity in 24 h

We first tested a gradient of insecticide doses on the laboratory strain *An. gambiae* Kisumu,

which is susceptible to common classes of insecticides used in mosquito control such as

pyrethroids, organophosphates, carbamates and organochlorines. The objective was to

determine a dose yielding 100% mortality in 24 h. We then used this dose as a discriminating

concentration to study the lethal and sublethal effects of neonicotinoids on field collected

Anophleles larvae. We compared the efficacy of neonicotinoids to that of deltamethrin. As

opposed to the susceptible strain, 100% mortality was not achieved with field populations

after 24 h (Figure 2). Mortality rates were very low in larvae of both *An. gambiae* and *An.*

coluzzii reared in water containing deltamethrin. Pyrethroid resistance is widespread in

anopheline mosquitoes, and parallel application of deltamethrin in malaria prevention and in

agricultural pest management has drastically reduced its efficacy against disease vectors. 24-h

mortality in water containing deltamethrin was typically below 30% and was not significantly

different between the four tested populations (Wilcoxon rank sum test, p>0.05). In contrast with deltamethrin, susceptibility to neonicotinoids varied between species and geographic areas. Consistent with findings obtained using adult bioassays (Ashu et al., 2023a; Fouet et al., 2020), *An. coluzzii* larvae collected from the two urban neighborhoods, Combattant and Etoa meki, were more susceptible, with mortality rates between 70-80%, except clothianidin for which only about 50% of larvae were killed within 24 h. The species present in suburban and rural areas of Yaoundé, *An. gambiae*, has developed strong resistance to neonicotinoids. Particularly, larvae collected from the Nkolondom farm where pesticide spraying has selected for resistance to acetamiprid and imidacloprid and conferred cross-resistance to clothianidin (Ashu et al., 2023a; Fouet et al., 2020), displayed less than 10% mortality in water containing a neonicotinoid. Mortality rate of larvae collected from the agricultural site was <10% in water containing the new malaria-control chemical, clothianidin, or the first-generation insecticide, deltamethrin, which suggested that both active ingredients currently have similar efficacy against some *An. gambiae* populations.

3.2 L4 transformation rate

Some neonicotinoid insecticides act slowly and are also known to have a wide range of sublethal effects in addition to acute toxicity on non-target species (Morrissey et al., 2015; Schuhmann et al., 2022). Thus, beside lethal toxicity within the first 24 h of the experiment, we used life table analysis to monitor some of the sublethal effects of neonicotinoids on *Anopheles* larvae, starting with L4 transformation rate. The three neonicotinoids tested had little effect on the ability of *An. gambiae* L3 larvae to become L4 (Figure 3A). The highest transformation rate was observed in populations from the agricultural site, Nkolondom: $90 \pm 3\%$ in water containing acetamiprid followed by $75 \pm 9\%$ in imidacloprid in two days and $60 \pm 3\%$ in clothianidin after 3 days. Similarly, in Nkolnkoumou, a semi-rural site harboring ~ 80% *An. gambiae*, transformation rates varied from $60 \pm 6\%$ for acetamiprid, to $35 \pm 16\%$

and 23 ± 7% for imidacloprid and clothianidin, respectively after 2 to 4 days. Conversely,

even if based on 24-h mortality, *An. coluzzii* larvae showed a degree of tolerance, they were

strongly inhibited in water containing neonicotinoids leading to very low transformation rates

between 0 and 10% in both Etoa Meki and Combattant populations. In control tests,

transformation rates were 100% after 2 to 3 days.

3.3 Pupation rate

Pupation rate was consistent with L4 transformation and highlighted the low susceptibility of

agricultural and semi-rural populations of *An. gambiae* to neonicotinoids (Figure 3B). The

agricultural population of *An. gambiae* (Nkolondom) had the highest pupation rate, 77.5 ± 4%

in water containing acetamiprid, and 45 ± 11% in imidacloprid reflecting their tolerance to the

both neonicotinoid pesticides that are sprayed in local farms. The exotic insecticide,

clothianidin, at 0.035 mg/L had the most important inhibitory power among neonicotinoids,

but 15 to 20% of *An. gambiae* larvae were able to achieve pupation. *Anopheles coluzzii* larvae

had low pupation rates with less than 5% pupae obtained in all the neonicotinoids tested.

Notably, none of the *An. coluzzii* populations were able to pupate in clothianidin.100% pupae

were obtained in all the control tests after 72 h.

3.4 Emergence rate

100% of *An. gambiae* and *An. coluzzii* pupae emerged into adults in control water without

insecticide between the first and the sixth day in standard laboratory conditions. The total

emergence rate of *An. coluzzii* larvae was 0%, 2% and 4% in water containing clothianidin,

acetamiprid or imidacloprid, respectively after 7 days (Figure 3C). Meanwhile, at least half of

An. gambiae L3 larvae tested on average emerged between 2-7 days in acetamiprid and

imidacloprid. Only *An. gambiae* larvae from the agricultural site were able to emerge in

clothianidin albeit at lower rate (7 ± 3%) compared with the other neonicotinoids. However,

emergence rate was also very low in deltamethrin, around 10%, indicating that for any

insecticide, the ability to inhibit emergence can be compatible with strong resistance in adult

mosquitoes and loss of efficacy of the insecticide. In addition to lethal toxicity in 24 h,

emergence rate revealed another similarity between clothianidin, a new insecticide and

deltamethrin whose effectiveness in malaria vector control has become very limited.

3.5 Survival probability

Rearing field-collected larvae in water containing an agricultural pesticide affected their

survival probability as revealed by Kaplan Meier survival curves (Figure 4). Larvae that

survived lethal toxicity within the first 24 h of the experiment were monitored for seven days.

Although *An. coluzzii* showed potential to resist the 24-h lethal effect of some of the

agricultural pesticides, larvae of this species had low survival probability in neonicotinoids.

By contrast, *An. gambiae* populations from agricultural and semi-rural sites displayed high

survival potential in water containing lethal doses of neonicotinoids. A log-rank test revealed

that regardless of the neonicotinoid that was added to water, larvae collected from

Nkolondom were super-resistant and had significantly higher survival probability compared

with any of the three other populations tested ($p < 0.05$). 100% survival was obtained in all

control experiments without insecticide.

4. Discussion

Here we have used an innovative approach to simultaneously assess the lethal and sublethal

effects of pesticide exposure in mosquito larvae. We found that monitoring some sublethal

effects via persistent exposure to chemicals in a controlled aquatic environment provides

critical information on the level of tolerance of larval populations. Such information could be

complementary to routine quantification of mortality in adults via standard bioassays (CDC,

2012; World Health Organization, 2022b), and combining both approaches may offer a more

robust framework to better evaluate pre-existing levels of adaptation to agrochemicals in

malaria mosquitoes. We conducted a laboratory experiment mimicking the selection process,

which is promoting the emergence of *Anopheles* populations with increased tolerance to

neonicotinoids. While analyzing mortality and life table in wild larvae reared in artificial

media containing a lethal dose of the active ingredient, we observed gradients of tolerance

reflecting past exposure to neonicotinoids in nature. Control tests supported the validity of our

experimental approach as 3rd instar larvae unexposed to insecticides showed standard

survival, growth and emergence in controlled insectary conditions.

Previous studies have suggested that resistance to neonicotinoids observed in anopheline

populations from some agricultural areas could be due to chronic exposure of larvae to

sublethal doses of agricultural pesticides (Ashu et al., 2023a, 2023b; Fouet et al., 2020;

Mouhamadou et al., 2019). Our findings provide the first experimental support to this

hypothesis. Indeed, we found that *An. gambiae* larvae had exceptional fitness in field-realistic

concentrations of acetamiprid and imidacloprid, two neonicotinoids whose formulations are

extensively used in farming activities in Cameroon (Fouet et al., 2020). As anopheline

populations have not yet been exposed to clothianidin in sub-Saharan Africa, this exotic

neonicotinoid strongly inhibited larval emergence in both *An. gambiae* and *An. coluzzii*.

However, likely due to cross-resistance selected by intensive exposure to acetamiprid and

imidacloprid, larvae from the agricultural site, Nkolondom, were able to survive in water

contaminated with clothianidin, albeit at lower frequency.

Exposure of non-target insects to sublethal doses of neonicotinoids affect functions such as

motility, behavior, growth, fecundity and survival (Morrissey et al., 2015; Řezáč et al., 2021;

Shepherd et al., 2021). Our study reveals that notwithstanding these sublethal effects whose

magnitude remains unknown, *An. gambiae* populations have evolved adaptation to chronic

exposure to neonicotinoid residues in contaminated aquatic habitats. *An. gambiae* larvae from

rural areas around Yaoundé exhibited high survivorship, L4 transformation, pupation and

emergence rate in water containing neonicotinoids contrary to urban *An. coluzzii* mosquitoes

whose growth and survival were inhibited. This result corroborates susceptibility profiles of

adult populations: *An. coluzzii* from Yaoundé is more susceptible to neonicotinoids while *An.*

gambiae is resistant to at least four different active ingredients (Ashu et al., 2023a, 2023b;

Fouet et al., 2020). Larval populations whose adults were resistant to an agrochemical (i.e.

mortality against the discriminating dose < 90%) typically displayed less than 50% mortality

in 24 h and more than 5% emergence in 6 days in water containing a lethal dose of the active

ingredient. Testing a larger number of chemicals on diverse *Anopheles* species and

populations will provide more robust guidelines on the interpretation of mortality, growth and

emergence of larvae reared in pesticide-laced water. However, based on the current study and

complimentary evidence from adults testing (Ashu et al., 2023a, 2023b; Fouet et al., 2020),

we could establish that any agrochemical yielding less that 50% mortality after 24 h and more

than 5% emergence within 6 days is likely to have limited efficacy against *Anopheles*

mosquitoes.

The first evidence of insecticide resistance driven by agricultural pesticide application dates

back to the 1960s. Based on observations made in El Salvador, it was clearly established that

aerial spraying of carbamates and organophosphates led to the development of broad-

spectrum resistance to both insecticide classes in the vector *Anopheles albimanus* (Georghiou,

1972; Georghiou et al., 1973). Since then, a number of observational studies have argued that

intensive use of crop protection chemicals is a major driver of resistance to public health

insecticides in anopheline mosquitoes (Chouaïbou et al., 2016; Diabate et al., 2002; Müller et

al., 2008; Nwane et al., 2009; Urio et al., 2022; Yadouleton et al., 2009). However, in the

majority of cases, it was difficult to determine the guilty part between agricultural chemicals

and malaria vector control insecticides. In a review of evidence, Lines (1988) (Lines, 1988)

proposed a set of conditions that should be met for a crop protection chemical to be the source

of insecticide resistance selection in malaria vectors. Appearance of resistance in mosquitoes

to a particular chemical should occur before it has been used for mosquito control; there

should be correlation in space and time, e.g. higher levels of resistance in areas with

agricultural spraying and seasonality of resistance overlapping with spraying periods; the

cross-resistance spectrum should show that the chemicals used in agriculture may confer

cross-resistance to public health insecticides and that both adults and larvae can be killed by

the agricultural pesticides. Agricultural application of neonicotinoids fits all of the criteria in

Sub-Saharan Africa and provides the best example of human activities accidentally

engineering insecticide resistance in mosquitoes.

In our study, both lethal and sublethal effects point to the existence of levels of resistance that

could be a major obstacle to using neonicotinoids in malaria mosquito control. We noted that

lethal toxicity in 24 h, pupation rate and emergence rate were very similar in water containing

either deltamethrin or clothianidin. This finding suggested that, just like deltamethrin, the

efficacy of clothianidin in *An. gambiae* might already be substantially reduced. In line with

this prediction, it has been shown that neonicotinoid resistant mosquitoes from Nkolondom

were significantly less susceptible to SumiShield® 50WG, a formulation of clothianidin

recently approved for indoor residual spraying (Fouet et al., 2020). Moreover, a laboratory

experiment recently demonstrated that exposure of *Anopheles* larvae to sublethal

concentrations of a mixture containing several herbicides, pesticides and fungicides resulted

in ~ 2.5 increase in tolerance to clothianidin and Fludora® Fusion, a formulation combining

clothianidin and deltamethrin (Zoh et al., 2022). These studies and ours highlight the

vulnerability of some agrochemicals repurposed for malaria vector control, as resistance

driven by agricultural activities could become a major thread to their efficacy.

Although our findings are based on only two *Anopheles* species sampled from a relatively

small geographic area, these results confirm concerns about the efficacy of neonicotinoids

against malaria vectors (Ashu et al., 2023a; Fouet et al., 2020; Hoppé et al., 2016;

Mouhamadou et al., 2019; Oxborough et al., 2019). Repurposing agrochemicals has thus far

provided a rapid mechanism to identifying new candidate insecticides for malaria prevention.

However, the case of neonicotinoids emphasizes the crucial role of evaluating prior exposure

and cross-resistance in areas where mosquito larval populations are likely to be in contact

with agricultural pesticides.

Author Contributions

MA: Conceptualization, Formal analysis, Investigation, Methodology, Writing – original

draft; CF: Conceptualization, Formal analysis, Investigation, Methodology; FA: Investigation,

Methodology; VP-B, Resources, Supervision. CK: Conceptualization, Formal analysis,

Funding acquisition, Investigation, Project administration, Writing – review & editing.

Funding

This study was supported by a National Institutes of Health grant (R01AI150529) to C K.

The funders had no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Informed Consent Statement

Not applicable.

Data Availability Statement

The data for this study have been presented within this article.

Conflicts of Interest

The authors declare that they have no competing interests.

References

Agossa, F.R., Padonou, G.G., Fassinou, A.J.Y.H., Odjo, E.M., Akuoko, O.K., Salako, A.,

Koukpo, Z.C., Nwangwu, U.C., Akinro, B., Sezonlin, M., Akogbeto, M.C., 2018. Small-

scale field evaluation of the efficacy and residual effect of Fludora® Fusion (mixture of

clothianidin and deltamethrin) against susceptible and resistant Anopheles gambiae

populations from Benin, West Africa. Malar. J. 17, 484. doi:10.1186/s12936-018-2633-6

Akpesse, A., 2018. Neonicotinoids and their impact on ecosystem service for agriculture and

biodiversity in Africa [WWW Document].

Antonio-Nkondjio, C., Fossog, B.T., Ndo, C., Djantio, B.M., Togouet, S.Z., Awono-Ambene,

P., Costantini, C., Wondji, C.S., Ranson, H., 2011. Anopheles gambiae distribution and

insecticide resistance in the cities of Douala and Yaoundé (Cameroon): influence of

urban agriculture and pollution. Malar. J. 10, 154. doi:10.1186/1475-2875-10-154

Antonio-Nkondjio, C., Tene Fossog, B., Kopya, E., Poumachu, Y., Menze Djantio, B., Ndo,

C., Tchuinkam, T., Awono-Ambene, P., Wondji, C.S., 2015. Rapid evolution of

pyrethroid resistance prevalence in Anopheles gambiae populations from the cities of

Douala and Yaoundé (Cameroon). Malar. J. 14, 1–9. doi:10.1186/s12936-015-0675-6

Ashu, F.A., Fouet, C., Ambadiang, M.M., Penlap-Beng, V., Kamdem, C., 2023a. Enhancing

the efficacy of neonicotinoids against mosquitoes and overcoming resistance issues.

bioRxiv. doi:https://doi.org/10.1101/2023.04.18.537427

Ashu, F.A., Fouet, C., Ambadiang, M.M., Penlap-Beng, V., Kamdem, C., 2023b. Vegetable

oil surfactants are synergists that can bias neonicotinoid susceptibility testing in adult

mosquitoes. bioRxiv. doi:https://doi.org/10.1101/2023.04.18.537421

Bamou, R., Sonhafouo-Chiana, N., Navridis, K., Tchuinkam, T., Wondji, C.S., Vontas, J.,

Antonio-Nkondjio, C., 2019. Status of Insecticide Resistance and Its Mechanisms in

Anopheles gambiae and Anopheles coluzzii Populations from Forest Settings in South

Cameroon. Genes (Basel). 10, doi:10.3390/genes10100741. doi:10.3390/genes10100741

CDC, 2012. Guideline for Evaluating Insecticide Resistance in Vectors Using the CDC Bottle

Bioassay, CDC Methods.

Chouaïbou, M.S., Fodjo, B.K., Fokou, G., Allassane, O.F., Koudou, B.G., David, J.P.,

Antonio-Nkondjio, C., Ranson, H., Bonfoh, B., 2016. Influence of the agrochemicals

used for rice and vegetable cultivation on insecticide resistance in malaria vectors in

southern Côte d'Ivoire. Malar. J. 15, 1–12. doi:10.1186/s12936-016-1481-5

Curtis, G.F., Miller, J.E., Hodjati, M.H., Kolaczinski, J.H., Kasumba, I., 1998. Can anything

be done to maintain the effectiveness of pyrethroid-impregnated bednets against malaria

vectors? Philos. Trans. R. Soc. B Biol. Sci. 353, 1769–1775. doi:10.1098/rstb.1998.0329

Diabate, A., Baldet, T., Chandre, F., Akogbeto, M., Guiguemde, R.T., Darriet, F., Brengues,

C., Guillet, P., Hemingway, J., Small, G.J., Hougard, J.M., 2002. The Role Of

Agricultural Use Of Insecticides In Resistance To Pyrethroids In Anopheles gambiae s . l

. In Burkina Faso. Am. J. Trop. Med. Hyg. 67, 617–622.

Fouet, C., Ashu, F.A., Ambadiang, M.M., Tchapga, W., Wondji, C.S., Kamdem, C., 2020.

Resistance to clothianidin reduces the efficacy of SumiShield 50WG®, a neonicotinoid

formulation for indoor residual spraying, against Anopheles gambiae. bioRxiv.

doi:https://doi.org/10.1101/2020.08.06.239509

Georghiou, G.P., 1972. Studies on Resistance to Carbamate and Organophosphorus

Insecticides in Anopheles albimanus. Am. J. Trop. Med. Hyg. 21, 797–806.

Georghiou, G.P., Breeland, S.G., V. Ariaratnam, 1973. Seasonal escalation of

organophosphorus and carbamate resistance in Anophelesalbimanus by agricultural

sprays. Env. Entomol 2, 369–374.

Hemingway, J., 2017. The way forward for vector control. Science (80-.). 358, 998–999.

Hoppé, M., Hueter, O.F., Bywater, A., Wege, P., Maienfisch, P., 2016. Evaluation of

Commercial Agrochemicals as New Tools for Malaria Vector Control. Chimia (Aarau).

70, 721–729. doi:10.2533/chimia.2016.721

Kamdem, C., Fouet, C., Etouna, J., Etoa, F.-X., Simard, F., Besansky, N.J., Costantini, C.,

2012a. Spatially explicit analyses of anopheline mosquitoes indoor resting density:

implications for malaria control. PLoS One 7, e31843.

doi:10.1371/journal.pone.0031843

Kamdem, C., Fouet, C., Gamez, S., White, B.J., 2017. Pollutants and insecticides drive local

adaptation in African malaria mosquitoes. Mol. Biol. Evol. 34, 1261–1275. doi:DOI:

10.1093/molbev/msx087

Kamdem, C., Tene Fossog, B., Simard, F., Etouna, J., Ndo, C., Kengne, P., Boussès, P., Etoa,

F.-X., Awono-Ambene, P., Fontenille, D., Antonio-Nkondjio, C., Besansky, N.J.,

Costantini, C., 2012b. Anthropogenic Habitat Disturbance and Ecological Divergence

between Incipient Species of the Malaria Mosquito Anopheles gambiae. PLoS One.

doi:10.1371/journal.pone.0039453

Katambo, M., 2018. Neonicotinoids and their impact on ecosystem services in Tanzania

[WWW Document].

Kramer, L., 2022. Pesticide use for agriculture in Africa from 2000 to 2019 [WWW

Document]. doi:https://www.statista.com/statistics/1291464/pesticide-use-for-

agriculture-in-africa/

Kweka, E., Mahande, A., Ouma, J., Msangi, S., Karanja, W., Temba, V., Lyaruu, L.,

Himeidand, Y., 2018. Novel Indoor Residual Spray Insecticide With Extended Mortality

Effect : A Case of SumiShield 50WG Against Wild Resistant Populations of Anopheles

arabiensis in Northern Tanzania. Glob. Heal. Sci. Pract. 6, 758–765.

Lees, R., Praulins, G., Davies, R., Brown, F., Parsons, G., White, A., Ranson, H., Small, G.,

Malone, D., 2019. A testing cascade to identify repurposed insecticides for next-

generation vector control tools: Screening a panel of chemistries with novel modes of

action against a malaria vector. Gates Open Res. 3, 1–25.

doi:10.12688/gatesopenres.12957.2

Lines, J.D., 1988. Do agricultural insecticides select for insecticide resistance in mosquitoes?

A look at the evidence. Parasitol. Today 4. doi:10.1016/0169-4758(88)90083-X

Matsuda, K., Ihara, M., Sattelle, D.B., 2020. Neonicotinoid insecticides: Molecular targets,

resistance, and toxicity. Annu. Rev. Pharmacol. Toxicol. 60, 241–255.

doi:10.1146/annurev-pharmtox-010818-021747

Mattah, P.A.D., Futagbi, G., Amekudzi, L.K., Mattah, M.M., De Souza, D.K., Kartey-

Attipoe, W.D., Bimi, L., Wilson, M.D., 2017. Diversity in breeding sites and distribution

of Anopheles mosquitoes in selected urban areas of southern Ghana. Parasites and

Vectors 10, 1–15. doi:10.1186/s13071-016-1941-3

Morrissey, C.A., Mineau, P., Devries, J.H., Sanchez-Bayo, F., Liess, M., Cavallaro, M.C.,

Liber, K., 2015. Neonicotinoid contamination of global surface waters and associated

risk to aquatic invertebrates: A review. Environ. Int. 74, 291–303.

doi:10.1016/j.envint.2014.10.024

Mouhamadou, C.S., Souza, S.S. De, Fodjo, B.K., Zoh, M.G., Bli, N.K., Koudou, B.G., 2019.

Evidence of insecticide resistance selection in wild Anopheles coluzzii mosquitoes due

to agricultural pesticide use. Infect. Dis. Poverty 8, 1–8.

Müller, P., Chouaïbou, M., Pignatelli, P., Etang, J., Walker, E.D., Donnelly, M.J., Simard, F.,

Ranson, H., 2008. Pyrethroid tolerance is associated with elevated expression of

antioxidants and agricultural practice in Anopheles arabiensis sampled from an area of

cotton fields in Northern Cameroon. Mol. Ecol. 17, 1145–1155. doi:10.1111/j.1365-

294X.2007.03617.x

N'Guessan, R., Boko, P., Akogbeto, M., Yates, A., Rowland, M., 2007. Chlorfenapyr : A

pyrrole insecticide for the control of pyrethroid or DDT resistant Anopheles gambiae (

Diptera : Culicidae) mosquitoes. Acta Trop. 102, 69–78.

doi:10.1016/j.actatropica.2007.03.003

Ngamo Tinkeu, L., 2018. Current situation on the registration and use of neonicotinoids in

Cameroon [WWW Document].

Nwane, P., Etang, J., Chouaibou, M., Toto, J.C., Kerah-hinzoumbé, C., Mimpfoundi, R.,

Awono-ambene, H.P., Simard, F., 2009. Trends in DDT and pyrethroid resistance in

Anopheles gambiae s.s. populations from urban and agro-industrial settings in southern

Cameroon. BMC Infect. Dis. 9, doi:10.1186/1471-2334-9-163. doi:10.1186/1471-2334-

9-163

Okolle, N.J., Monono, E.Y., Tabikam, Amungwa Ivan Mambo, Stephania Kinge Magwell,

P.F.R., 2022. Insecticide Use and Application in Cameroon, in: Insecticides. IntechOpen.

Oxborough, R.M., Kitau, J., Matowo, J., Mndeme, R., Feston, E., Boko, P., A., O., Metonnou,

C.G., Irish, S., N'Guessan, R., Mosha, F.W., Rowland, M., 2010. Evaluation of indoor

residual spraying with the pyrrole insecticide chlorfenapyr against pyrethroid-susceptible

Anopheles arabiensis and pyrethroid-resistant Culex quinquefasciatus mosquitoes. Trans.

R. Soc. Trop. Med. Hyg. 104, 639–645. doi:10.1016/j.trstmh.2010.07.008

Oxborough, R.M., Seyoum, A., Yihdego, Y., Dabire, R., Gnanguenon, V., Wat, F., Agossa,

F.R., Yohannes, G., Coleman, S., Samdi, L.M., Diop, A., Faye, O., Magesa, S.,

Manjurano, A., Okia, M., Alyko, E., Masendu, H., Baber, I., Sovi, A., Rakotoson, J.,

Varela, K., Abong, B., Lucas, B., Fornadel, C., Dengela, D., 2019. Susceptibility testing

of Anopheles malaria vectors with the neonicotinoid insecticide clothianidin; results

from 16 African countries, in preparation for indoor residual spraying with new

insecticide formulations. Malar. J. 18, https://doi.org/10.1186/s12936-019-2888-6.

doi:10.1186/s12936-019-2888-6

Portwood, N.M., Mbewe, N.J., Tungu, P.K., Mlay, G., Small, G., Snetselaar, J., Kristan, M.,

Levy, P., Walker, T., Kirby, M.J., Kisinza, W., Mosha, F.W., Rowland, M., Messenger,

L.A., 2022. Multi-centre discriminating concentration determination of broflanilide and

potential for cross- resistance to other public health insecticides in Anopheles vector

populations. Sci. Rep. 1–10. doi:10.1038/s41598-022-26990-6

R Core Team, 2016. R: A language and environment for statistical computing.

Ramadevi, R., Ramachandraiah, C., Reddy, G.V.S., 2022. A Review on Contamination of

Soil and Water by Neonicotinoid Pesticides and Trends it's in Soil and Water Samples

with Chromatographic Analytical Techniques. Orient. J. Chem. 38, 259–267.

doi:10.13005/ojc/380205

Řezáč, M., Gloríková, N., Wilder, S.M., Heneberg, P., 2021. The sublethal effects of

neonicotinoids on spiders are independent of their nutritional status. Sci. Rep. 11, 1–11.

doi:10.1038/s41598-021-87935-z

Schaafsma, A., Limay-Rios, V., Baute, T., Smith, J., Xue, Y., 2015. Neonicotinoid insecticide

residues in surface water and soil associated with commercial maize (corn) fields in

Southwestern Ontario. PLoS One 10, 1–21. doi:10.1371/journal.pone.0118139

Schuhmann, A., Schmid, A.P., Manzer, S., Schulte, J., Scheiner, R., 2022. Interaction of

Insecticides and Fungicides in Bees. Front. Insect Sci. 1, 1–14.

doi:10.3389/finsc.2021.808335

Service, M.W., 1993. Mosquito ecology: field sampling methods, 2nd ed. Springer,

Netherlands. doi:10.1007/978-94-015-8113-4

Shepherd, S., Lima, M.A.P., Oliveira, E.E., Sharkh, S.M., Aonuma, H., Jackson, C.W.,

Newland, P.L., 2021. Sublethal neonicotinoid exposure attenuates the effects of

electromagnetic fields on honey bee flight and learning. Environ. Adv. 4, 100051.

doi:10.1016/j.envadv.2021.100051

Simon-delso, N., Amaral-rogers, V., Belzunces, L., Bonmatin, J.M., Chagnon, M., Downs, C.,

Furlan, L., Gibbons, D.W., Giorio, C., Girolami, V., 2015. Systemic insecticides

(neonicotinoids and fipronil): trends , uses , mode of action and metabolites. Environ.

Sci. Pollut. Res. 22, 5–34. doi:10.1007/s11356-014-3470-y

Tene-Fossog, B., Fotso-Toguem, Y.G., Amvongo-Adjia, N., Ranson, H., Wondji, C.S., 2022.

Temporal variation of high-level pyrethroid resistance in the major malaria vector

Anopheles gambiae s.l. in Yaoundé, Cameroon, is mediated by target-site and metabolic

resistance. Med. Vet. Entomol. 36, 247–259. doi:10.1111/mve.12577

Uragayala, S., Kamaraju, R., Tiwari, S.N., Sreedharan, S., Ghosh, S.K., Valecha, N., 2018.

Village-scale (Phase III) evaluation of the efficacy and residual activity of SumiShield

50® WG (Clothianidin 50 %, w/w) for indoor spraying for the control of pyrethroid-

resistant Anopheles culicifacies Giles in Karnataka state , India. Trop. Med. Int. Heal.

23, 605–615. doi:10.1111/tmi.13056

Urio, N.H., Pinda, P.G., Ngonzi, A.J., Muyaga, L.L., Msugupakulya, B.J., Finda, M.,

Matanila, G.S., Mponzi, W., Ngowo, H.S., Kahamba, N.F., Nkya, T.E., Okumu, F.O.,

2022. Effects of agricultural pesticides on the susceptibility and fitness of malaria vectors

in rural south-eastern Tanzania. Parasites and Vectors 15, 1–14. doi:10.1186/s13071-

022-05318-3

World Health Organization, 2023. List of WHO Prequalified Vector Control Products [WWW

Document]. URL https://extranet.who.int/pqweb/vector-control-products/prequalified-

product-list

World Health Organization, 2022a. World malaria report 2022. Geneva.

World Health Organization, 2022b. Standard operating procedure for testing insecticide

susceptibility of adult mosquitoes in WHO tube tests, World Health Organization.

Yadouleton, A.W.M., Asidi, A., Djouaka, R.F., Brama, J., Agossou, C.D., Akogbeto, M.C.,

2009. Development of vegetable farming: A cause of the emergence of insecticide

resistance in populations of Anopheles gambiae in urban areas of Benin. Malar. J. 8, 1–8.

doi:10.1186/1475-2875-8-103

Zogo, B., Koffi, A.A., Alou, L.P.A., Fournet, F., Dahounto, A., Dabiré, R.K., Baba-Moussa,

L., Moiroux, N., Pennetier, C., 2019. Identification and characterization of Anopheles

spp. breeding habitats in the Korhogo area in northern Côte d'Ivoire: A study prior to a

Bti-based larviciding intervention. Parasites and Vectors 12, 1–10. doi:10.1186/s13071-

019-3404-0

Zoh, M.G., Tutagata, J., Fodjo, B.K., Mouhamadou, C.S., Sadia, C.G., McBeath, J., Schmitt,

F., Horstmann, S., David, J.P., Reynaud, S., 2022. Exposure of Anopheles gambiae

larvae to a sub-lethal dose of an agrochemical mixture induces tolerance to adulticides

used in vector control management. Aquat. Toxicol. 248.

doi:10.1016/j.aquatox.2022.106181

Figure legends

Figure 1: Examples of breeding sites where wild mosquito larvae were sampled. *An. gambiae*

larvae were collected from furrows between ridges in the agricultural site (A) or from puddles

created by human activities in Nkolnkoumou (suburban area) (B). *An. coluzzii* larvae were

sampled from human-made puddles in densely urbanized settings (C).

Figure 2: 24-h mortality of mosquito larvae reared in water containing a lethal dose of

agrochemical. 100% of larvae of the susceptible strain *An. gambiae* Kisumu were killed

within 24 h. Wild populations showed varying levels of tolerance to four insecticides.

Mortality values were lowest in larvae collected from the agricultural area (Nkolondom).

Error bars represent the standard error of the mean.

Figure 3: Sublethal effects of three neonicotinoids on life table parameters of *Anopheles*

larvae. Rate of transformation of 3rd instars (L3) into 4th instars (L4) (A), pupation rate (B)

and emergence rate (C). Larvae that survived 24-h lethal toxicity were monitored for 7 days

under standard laboratory conditions. Controls were larvae reared in water without

insecticide. Vertical bars represent the standard error of the mean.

610 **Figure 4:** Kaplan Meier survival curves (lines) with 95% confidence intervals (colored bands)

611 of 3^{rd} instar larvae reared in water containing a lethal dose of neonicotinoid. Survival

612 probabilities of four wild populations were compared under controlled laboratory conditions.

613 (+) indicates emergence that occurred before the end of the experiment and was treated as

614 censored data.

620