Leveraging a new data resource to define the response of *C. neoformans* to environmental signals

How host-like signals drive gene expression and capsule expansion in *Cryptococcus neoformans*

Yu Sung Kang[1,2,3], Jeffery Jung[1,2,3], Holly Brown[3], Chase Mateusiak[3], Tamara L. Doering[5], and Michael R. Brent[2,3,4,6]

1. These authors contributed equally.

2. Center for Genome Sciences and Systems Biology, Washington University School of Medicine, St. Louis, MO 63110

3. Department of Computer Science and Engineering, Washington University, St. Louis, MO 63108

4. Department of Genetics, Washington University School of Medicine, St. Louis, MO 63110

5. Department of Molecular Microbiology, Washington University School of Medicine, St. Louis, MO 63110

6. Corresponding author

Michael Brent

brent@wustl.edu

http://mblab.wustl.edu

314-660-2205

Campus Box 8510

Washington University

Saint Louis, MO 63130

ABSTRACT

Cryptococcus neoformans is an opportunistic fungal pathogen with a polysaccharide capsule that becomes greatly enlarged in the mammalian host and during *in vitro* growth under host-like conditions. To understand how individual environmental signals affect capsule size and gene expression, we grew cells in all combinations of five signals implicated in capsule size and systematically measured cell and capsule sizes. We also sampled these cultures over time and performed RNA-Seq in quadruplicate, yielding 881 RNA-Seq samples. Analysis of the resulting data sets showed that capsule induction in tissue culture medium, typically used to represent host-like conditions, requires the presence of either CO_2 or exogenous cyclic AMP (cAMP). Surprisingly, adding either of these pushes overall gene expression in the opposite direction from tissue culture media alone, even though both are required for capsule development. Another unexpected finding was that rich medium blocks capsule growth completely. Statistical analysis further revealed many genes whose expression is associated with capsule thickness; deletion of one of these significantly reduced capsule size. Beyond illuminating capsule induction, our massive, uniformly collected dataset will be a significant resource for the research community.

IMPORTANCE

Cryptococcus neoformans is an opportunistic yeast that kills ~150,000 people each year. This major impact on human health makes it imperative to understand the basic biology of *C. neoformans* and the factors that mediate its virulence. One key virulence factor is a polysaccharide capsule that expands greatly during infection. To help define capsule synthesis and fungal biology, we provided cells with many different combinations of host-like signals and sampled the cultures over time for transcriptional analysis. The resulting time resolved data set is by far the largest gene expression resource ever produced for *C. neoformans* (881 RNA-seq samples), further enriched by accompanying capsule images and measurements. It revealed surprising findings, including that rich medium suppresses capsule size regardless of other signals. This landmark data resource will be enormously valuable to the research community as it continues to define the relationships between environmental signals and cryptococcal gene expression, biology, and virulence.

Late in the 19[th] century, several scientific articles described a budding yeast with a dis-
tinctive capsule, now called *Cryptococcus neoformans*. Today, we know *C. neofor-*
mans as the main causative agent of a deadly meningitis that kills close to 150,000
people each year worldwide[1]. We also know much about the capsule that surrounds this
pathogen, including the chemical structure of the polysaccharides that compose it, its
key role in disease, and the fact that it is exquisitely sensitive to environmental condi-
tions.[2] Upon entry to a mammalian host, the capsule dramatically increases in thick-
ness, from a barely perceptible structure to a distinctive shell whose thickness can ex-
ceed the cell's diameter.[3] Enlarged capsules inhibit phagocytosis of the yeast by host
immune cells and shed capsule polysaccharides inhibit host defenses.[4] The importance
of this material in cryptococcal pathogenesis is amply supported by the reduced viru-
lence of strains in which capsule is altered or dysregulated.[2]

Multiple *in vitro* conditions induce the growth of cryptococcal capsule.[5-10] Conditions that
reflect aspects of the mammalian host environment are of particular interest, such
as those that incorporate tissue culture medium (TCM) or mammalian serum[9], human
body temperature (37 °C), physiological pH (7.35-7.45),[6,9-12] host-like CO_2 concentra-
tions (~5%)[6], or nutrient limitations typical of the host environment.[7]

Cyclic AMP (cAMP) signaling is required for capsule growth and virulence of *C. neofor-*
mans.[3,13] This pathway has been well described in *Saccharomyces cerevisiae*[14,15] and
much of the machinery is also found in *C. neoformans*.[13] Cellular levels of cAMP reflect
its formation from ATP by Cac1 (adenylyl cyclase) and degradation by Pde1 and Pde2
(phosphodiesterases). cAMP binds the repressive subunit of the protein kinase A (PKA)
complex, Pkr1, causing it to separate from the catalytic subunit, Pka1. This activates
Pka1, allowing it to phosphorylate transcription factors that are central in capsule regu-
lation, including Nrg1[16] and Rim101[17]. In growth conditions where capsule is normally
induced, strains lacking Cac1[3,18] or Pka1[17,19] fail to do so. In the same conditions Pkr1-
deficient mutants make larger capsules than wild-type (WT) cells[3,20].

In this paper, we assess key features of capsule inducing conditions, which we call *sig-*
nals, focusing on five of them: tissue-culture medium (DMEM or RPMI), temperature (37
°C), CO_2 (5%), exogenous cAMP, and the addition of buffer. We previously showed that
1,10-phenanthroline, which inhibits transcription, completely blocks capsule growth.[21,22]
We have now systematically explored the effects of these signals on gene expression,
cell size, and capsule size, both individually and in combination, over time. We also as-
sessed the effects of deleting several cAMP pathway genes. Our analysis of almost 900
RNA-seq samples and over 5,000 micrographs has allowed us to trace capsule-
inducing signals through their effects on gene expression to their ultimate effects on
capsule size. This large, uniformly collected dataset, which will be a significant resource
for the research community, also enabled us to identify the changes in gene expression
that are most consistently associated with capsule growth across a wide range of
growth conditions and genetic perturbations. We further discovered that capsule induc-
tion is blocked by rich medium under multiple growth conditions and identified new
genes that are required for normal capsule thickness.

103
104 RESULTS
105
106 <u>Effects of signals on capsule size.</u>
107
To tease out the effects of various environmental signals on capsule size, we examined
all possible combinations of the following growth condition variables: medium (YPD,
DMEM, RPMI), CO_2 (room air, 5%), temperature (30 °C, 37 °C), HEPES buffer pH 7.0
(none, 25 mM), and cAMP (none, 20 mM). For these studies we used *C. neoformans*
strain KN99α, which was derived from the reference strain H99.[23] For rigor of the stud-
ies, we grew at least four replicates in each combination of conditions and measured
capsule sizes for fifteen fields of view per replicate (totaling 96 annotated cells on aver-
age).
116
We first examined how each individual variable affects capsule size, adjusting for the
independent effects of all other variables. To do this, we built a linear regression model
with the environmental variables as predictors of capsule size. The resulting regression
coefficients showed that the factor with the biggest impact on capsule size was RPMI
medium, which increased capsule width an average of 1.12 μm, followed by DMEM
(0.67 μm), cAMP (0.53μm) and CO_2 (0.51 μm), (Fig. 1A).
123
We observed no statistically significant effect of increasing temperature from 30 °C to
37 °C or adding HEPES buffer (Fig. 1A). We therefore combined replicates, regardless
of temperature or HEPES, and plotted capsule size for all possible combinations of me-
dium, CO_2, and cAMP (Fig. 1B). Strikingly, there was no capsule induction in any condi-
tion with YPD, regardless of CO_2 or cAMP addition. DMEM alone yielded negligible in-
duction, but adding cAMP, CO_2, or both yielded progressively thicker capsules. Results
in RPMI were similar, except that it produced larger capsules than DMEM in every con-
dition. In RPMI, cAMP also had a larger effect than in DMEM, such that its effect was
similar to that of CO_2.
133
134 To further explore how the signals that affect capsule size interact, we identified the
135 conditions under which each signal had the greatest impact. DMEM and RPMI each had
136 their biggest effects in the presence of both CO_2 and cAMP. cAMP had its biggest effect
137 in RPMI without CO_2 at 30°, while CO_2 had its biggest effect in DMEM with cAMP at 37
138 °C (Fig. 1C). These observations guided our choice of media in the analyses below.
139
Taking these results together, the most surprising result was that no combination of
temperature, CO_2, cAMP, or HEPES buffer induced capsule at all in YPD (Fig. 1B; all P-
values > 0.2). This suggested to us the novel idea that this rich medium has a repres-
sive effect on capsule (see Discussion). Our data set also allowed us to make several
other observations. First, RPMI generally led to greater capsule width than DMEM, hold-
ing all other variables constant (linear model, $P<10^{-8}$). Second, CO_2 or cAMP each in-
creased capsule size in either tissue culture medium (DMEM ± CO_2, $P<10^{-3}$; RPMI ±

CO_2, P<10^{-3}; DMEM ± cAMP, P<0.058; RPMI ± cAMP, P<10^{-3}), and the two together
yielded the largest capsules.

Effects of signals on gene expression

To examine the transcriptional response to various combinations of signals, we collect-
ed samples for RNA-Seq upon initial exposure to the signals and after 30, 90, 180, and
1440 min. After quality control (see Methods), we were left with RNA-Seq data for 720
samples. These included at least four replicates for each combination of environmental
signals, which had been grown and libraries prepared on separate days to account for
day-to-day variability in both processes. Each growth batch and library preparation
batch also included a control culture grown in our standard non-inducing laboratory
conditions: YPD, 30°, no CO_2, no cAMP, no added buffer. This design allowed us to re-
gress out batch (date) effects, which often dominate experimental factors in large-scale
gene expression studies.

To assess the magnitude of signal effects on overall gene expression, we computed
principal components and plotted PC1 and PC2 for each sample. Time of growth had
the biggest effect on both PC1 and PC2 (Fig. 2A), as can be seen by the progression
towards the upper right of data points from successive sampling times. The next great-
est effect was due to medium, as shown for the 24 h timepoints in Fig. 2B, where YPD
samples (blue) are clearly separated from those in TCM (green and orange). Samples
grown in TCM with various combinations of cAMP or CO_2 were also distinct in terms of
transcriptional response (Fig. 2C, showing PC1 and PC3). Together, these results show
that the primary differences in gene expression are driven by experimental factors, ra-
ther than batch effects, demonstrating the quality of the data. Furthermore, TCM, which
has the greatest effect on capsule size at 24 h, also has the greatest effect on gene ex-
pression at 24 h.

Effects of DMEM and CO_2 on gene expression

In the analyses above, capsule growth required both a tissue culture medium and either
CO_2 or cAMP. To identify the biological functions of the genes that were most respon-
sive to these signals, we started with the response to CO_2 in DMEM, a combination that
had large effects on capsule width (Fig. 1). To assess transcriptional changes we ran
DESeq2[24] on these datasets with a linear model that predicts log normalized gene ex-
pression levels from signals and extracted shrunken log fold changes in response to
each signal (see Methods).

Next, we carried out over-representation analysis on the most responsive genes
at each time point using Gene Ontology (GO) biological process terms together with
additional functional categories of interest. Examples of enriched biological functions
are shown in Figure 2D, as a heatmap showing the average expression levels of all
genes annotated with each term (not just the significantly differentially expressed
genes). For each time point, we show the effects of DMEM relative to YPD, the effects

of DMEM+CO_2 relative to DMEM alone, and the combined effects of DMEM+CO_2 rela-
tive to YPD alone.

We first examined genes that are upregulated by DMEM alone. The genes that
were most strongly induced are involved in phosphate transport (Fig. 2D, top row), con-
sistent with recent reports that these genes are induced by another capsule-inducing
growth condition, 10% Sabouraud's dextrose medium.[25] Other biological functions that
were strongly upregulated in DMEM include environmental stress response (Fig. 2D,
fourth row) and the functions designated as Group 1 (Fig. 2D; see gene lists in Supple-
mental File S5). This group includes several functions that are notable for their involve-
ment in cell wall synthesis (cell wall organization and chitin metabolism) as well as car-
bohydrate biosynthesis. The individual expression patterns of each gene in this catego-
ry are shown in Fig. 2E, plotted as the ratio of expression in the indicated conditions at
each time point (so they may reflect differences in either or both conditions). The ex-
pression ratio for many of these genes was upregulated by at least 2-fold (colored
lines), including for those involved in gluconeogenesis (*PCK1*, *TPI1*, *PYC2*), glycogen
synthesis (*GLC3*, *GSY1*), and trehalose synthesis (*TSP1* and *TSP2*, both of which are
required for virulence in *C. neoformans*[26]). All of these genes showed peak expression
difference from YPD at 30 minutes with a slow convergence back to YPD levels over
time (see Discussion).

Another category of interest was genes implicated in capsule synthesis (genes
annotated with a capsule phenotype in FungiDB, shown in the third row of Fig. 2D).
These were, on average, upregulated by DMEM and barely affected by CO_2 (see Dis-
cussion). Finally, we noted that homologs of yeast genes expressed specifically during
the M/G phases of the cell cycle (identified in ref.[27]) were upregulated by DMEM and
slightly down regulated by CO_2, while those expressed at other cell cycle phases were
generally downregulated by DMEM, relative to YPD (see Discussion).

Overall, the gene sets that were most strongly down regulated in response to
DMEM (Fig. 2D, group 2) are growth related: ribosome biogenesis, steroid biosynthesis,
cell division, and chromosome segregation. Expression of each individual gene in the
last category (Fig. 2F) shows that, like upregulated genes, the most dramatic differ-
ences from YPD levels occurred by the first time point (30 min), with expression in the
two cultures becoming more similar by the end of the 24-hour period. The category of
carbohydrate transport was also strongly down-regulated relative to YPD for most time
points, but then up-regulated at 24 hours (Fig. 2D, second row). Among these carbohy-
drate transport genes are *LPI8*, whose product promotes cryptococcal uptake by phag-
ocytes[28], and *GMT2*, which encodes a GDP-mannose transporter required for the syn-
thesis of capsule and other glycoconjugates.[29]

Interestingly, the effect of CO_2, when added to DMEM, was in the opposite direc-
tion from the effect of DMEM relative to YPD for most gene sets in the heat map. How-
ever, the CO_2 response was weaker, so the overall effect of DMEM+CO_2 was in the

same direction as that of DMEM alone. Exceptions to this pattern include ribosome bio-
genesis and mRNA splicing, both of which were consistently reduced.

Effects of cyclic AMP on capsule and cell size

Cyclic AMP signaling is essential for capsule growth.[3,13,18] To investigate the effects of
exogenous cAMP on capsule size and gene expression, we performed experiments in
conditions in which capsule is normally small, but addition of cAMP generates much
larger capsules: RPMI, 30°, room air, no HEPES (Fig. 1B). We found that increasing
cAMP yielded a consistent dose response in capsule width (Fig. 3A), with the logarithm
of exogenous cAMP concentration a highly significant predictor of average capsule
thickness ($P<10^{-10}$). On average, increasing cAMP by a factor of 1.8 increased the cap-
sule width by 0.2 um, although the final step from 11 mM to 20 mM had a much bigger
effect (Figure 3A).

Exogenous cAMP also increased the average cell body radius, excluding the
capsule, (Fig. 3B; ANOVA $P < 3*10^{-5}$), with a 1.8 fold increase in cAMP causing an av-
erage increase of 0.11 um (Fig. 3B). Unlike the effect on capsule width, this effect was
greater at lower concentrations. Notably, the effect of cAMP on cell size does not ex-
plain its effect on capsule width, since the capsule index (capsule width as a fraction of
total radius) increased with cAMP concentration (Fig. S1).

We took advantage of mutants in the cAMP pathway to further investigate the
role of cAMP in capsule enlargement. We first measured the capsules of cells lacking
PDE1, which encodes a phosphodiesterase capable of degrading intracellular cAMP.
We observed little effect on capsule width in either YPD or RPMI (both at 30° with no
CO_2 or cAMP; Fig 3C), consistent with previous reports[30]. We next examined cells lack-
ing PKR1, which encodes the repressive moiety of the Protein Kinase A (PKA) complex.
cAMP causes Pkr1 to dissociate from the complex, activating the kinase moiety, so a
pkr1 deletion mutant might be expected to have an enlarged capsule. Indeed, we saw
that, when grown in RPMI alone, the pkr1 mutant's capsule was similar to that of WT
grown in RPMI with 20 mM cAMP (Fig. 3C). We also saw an increase in capsule width
of pkr1 grown in YPD. No other perturbation we tested, including 20 mM exogenous
cAMP, yielded increased capsule size in YPD (Fig. 1B; see Discussion). Meanwhile, cell
body size was only modestly changed in the mutants in each growth condition (Fig. 3D).

Effects of cAMP on gene expression

In our experiments, cAMP had the greatest impact on WT capsule size in RPMI
with no CO_2 (Figure 1C, right-most panel). For this reason, we characterized gene ex-
pression changes over time in response to these conditions (Fig. 3E), examining the
same functional categories we had for DMEM and CO_2 (Fig. 2D). The responses to
RPMI and DMEM were broadly similar (compare Figs. 3E and 2D). Furthermore, adding
cAMP or CO_2 tended to moderate the responses to TCM more often than it reinforced
them. However, we did observe some differences. For example, while RPMI and DMEM
both reduced the expression of ribosome biogenesis genes, adding CO_2 reinforced that
effect at all time points, while adding cAMP moderated it at later time points. We also

noted that genes involved in carboxylic acid metabolic processes (mainly nitrogen as-
similation and amino acid metabolism) responded differently to RMPI+cAMP and
DMEM+CO_2, especially at early time points.

Nitrogen source is another factor that has been implicated in capsule regulation.
[6] Nitrogen catabolite repression (NCR) is a process that represses the expression of
certain genes when preferred nitrogen sources, such as glutamine (the main nitrogen
source in DMEM and RPMI) are available.[31] In RPMI, the genes in this category were
indeed slightly repressed compared to YPD (Fig. 3E, left columns). However, adding
cAMP released NCR rapidly, most notably at 30 min (Fig. 3E, middle columns), sug-
gesting that the cells responded to cAMP as though they had been moved to a less pre-
ferred nitrogen source (despite the presence of glutamine in the medium; see Discus-
sion). This is consistent with the general role of cAMP in cryptococcal stress response.
cAMP addition also overcame the RPMI effect, so that their combined effect was to re-
lieve NCR (Fig. 3E, right columns). The detailed effect of adding cAMP to RPMI is
shown in Fig. 3F.

Finally, to evaluate the effects of signals across all genes, rather than on specific
processes, we calculated the global correlations between pairs of signals at each time
point. As we had observed for selected groups of genes (Fig. 3E), the effects on gene
expression of the two tissue culture media (DMEM and RPMI) are highly correlated (Fig.
3G). The effects of CO_2 and cAMP were positively correlated with each other, consistent
with previous evidence that CO_2/HCO_3 stimulate adenylyl cyclase, which generates
cAMP.[32]

Expression of multiple genes is associated with capsule size

Next, we set out to identify genes whose expression is strongly associated with capsule
development. Our approach was to dichotomize capsule size into "induced" or "not in-
duced" and the expression of each gene into "high" or "low" and analyze the relationship
between expression and capsule induction for each gene separately. We considered a
sample to be "induced" if its median capsule width after 24 hours exceeded 1.13 um,
three standard deviations above the mean of all WT cells in YPD. For each gene at
each RNA-Seq time point, we then searched for a threshold to divide high and low ex-
pression for that gene that was most associated with induction status. Specifically, we
calculated the χ^2 statistic for association between induction status and gene expression
status (Fig. 4A; see Methods for details). We were not concerned with P-values here,
but with using the statistic itself to rank genes by their association with capsule size.
The maximum χ^2 value, over all time points, was assigned to the gene and the genes
with high χ^2 values were identified as potentially involved in capsule induction (Table
S1). However, we were mindful that these associations can arise in many ways and do
not necessarily reflect a causal role in capsule induction.

New genes whose deletion affects capsule size

As noted above, the same conditions that induce capsule and stimulate changes in ex-
pression of genes essential for capsule growth also affect expression of genes involved

in multiple other processes, including cell division, nutrient utilization, or other cellular
processes (see Figs. 2D and 3E). We therefore expected only a small subset of the
genes whose expression is correlated with capsule thickness to be essential for capsule
development. To seek such genes, we examined deletion strains. Mutants correspond-
ing to 24 of the 33 genes in Table S1 were available in a publicly available deletion col-
lection[33] or in our own collections. We grew each mutant in the conditions we found to
induce large capsules (RPMI, 37 °C, 5% CO_2, no cAMP) and measured their capsules
using our rigorous protocols (see Methods).

We were excited to identify a mutant that was not previously known to influence
capsule width. Deletion of CNAG_00368, which encodes a homolog of *S. cerevisiae*
Vps53, reduced mean capsule width by 0.44 μm (Fig. 4B,C). A scatter plot of capsule
width versus expression for this gene is shown in Fig. 4A.

An adventitious finding in these studies was that a strain deleted for
CNAG_05977 had mean capsule width 0.54 um smaller than wild-type cells (see Dis-
cussion). This strain was not chosen as described above – we had intended to test
CNAG_06050, but routine genome sequencing for quality control showed that the strain
we tested in fact had a deletion of CNAG_05977. We verified the genotypes of both de-
letion strains by whole-genome sequencing.

DISCUSSION

This work makes significant contributions in three areas. First, we have generat-
ed a major dataset that will be a tremendous resource for researchers in the field. Sec-
ond, we have used these data to make interesting observations about how cells react to
environmental conditions that induce capsule. Third, we identified a set of genes whose
expression correlates with capsule size and discovered two new genes that influence
capsule thickness.

When we undertook this project, it was known that capsule growth requires both
new transcription[6,22] and an intact cAMP/PKA pathway.[3,13,18] It was also known that, in
tissue culture medium (TCM), capsule can be induced by increasing the concentration
of dissolved CO_2/HCO_3^-, achieved by growth in a high CO_2 atmosphere, addition of
$NaHCO_3$, or both.[6] We set out to determine how various combinations of capsule induc-
ing signals would affect gene expression and capsule size. This led to a massive, freely
available dataset consisting of RNA-Seq in biological quadruplicate in 42 combinations
of potentially capsule-inducing signals, as well as cAMP titrations and mutants in the
cAMP signaling pathway (881 total RNA-Seq samples). Each RNA-Seq experiment
comes with matched India ink images and capsule width measurements. The dataset
also includes images and capsule-width measurements for selected gene deletion mu-
tants, yielding a total of 47,458 annotated cells in 5,175 images of 392 biological sam-
ples.

We found that either a 5% CO_2 atmosphere or cAMP could induce capsule in
TCM, but YPD, a rich medium containing yeast extract and peptone, completely blocks
capsule growth. This repressive effect is upstream of PKA activation, since deleting

PKR1 increases capsule sizes of cells growing in YPD (Fig. 3C). However, adding 20
mM exogenous cAMP, which is thought to act directly on the Pka1/Pkr1 complex, does
not enlarge the capsules of cells growing in YPD (Fig. 1B). It may be that cAMP cannot
completely inactivate Pkr1 or that cells do not maintain a high enough internal concen-
tration for long enough to fully inactivate it.

RPMI alone caused capsules to enlarge slightly, whereas DMEM alone produces
a barely detectable enlargement (Fig. 1B). RPMI contains less of most nutrients than
DMEM, making it a generally more stressful condition. One possible explanation for
these observations is the absence of iron in RPMI (DMEM contains 10^{-4} g/L ferric ni-
trate). These results are reminiscent of observations concerning titan cells, extremely
large cells that form during infection and specific *in vitro* conditions.[34] Interestingly, titan
cell formation can be induced by low-density inoculation into RPMI at 37 °C with 5%
CO_2, but substitution of DMEM for RPMI prevents this. The critical differences in this
case were identified as the presence of iron in DMEM and the presence of para-
aminobenzoic acid in RPMI.[35]

In addition to the conditions we studied, capsule growth is known to be stimulat-
ed by specific stressors, some of which resemble stresses encountered in a mammalian
host. For example, growth in mammalian serum without added nutrients induces cap-
sule growth at 30 °C or 37 °C, with or without CO_2, although this does not occur in
Sabouraud dextrose broth, a rich medium like YPD.[9] Sabouraud medium alone, diluted
to 10% normal concentration and buffered to pH 7.4 (termed CAP medium), is reported
to induce capsule.[8] Iron deprivation in LIM medium, which consists of 5 g/L glucose, 5
g/L asparagine, minerals, and 55 mM EDDA at pH 7.4, can also induce capsule.[7] Thus,
rich media consistently block capsule induction while several forms of nutrient depriva-
tion induce it. Nutrient deprivation is likely the normal state of Cryptococcus, whether in
a mammalian host or in the environment, so we suggest that it makes more sense to
think of rich media as blocking capsule growth than to think of TCM as inducing it.

Low pH (6.1) can also block capsule induction.[6] High osmolarity appears to re-
duce capsule thickness, although there has been speculation about whether this is a
cellular response or simply physical compression of capsule.[12] A cellular response is
consistent with the observation that deletion of *HOG1*, a key component of the high os-
molarity response, increases capsule size in DMEM+CO_2 induction.[3,36] Studying the ef-
fects of these stimuli on gene expression may provide additional insights about which
changes in gene expression are essential for capsule growth.

Turning to gene expression, tissue culture media (TCM) have several interesting
effects. First, they decrease the expression of genes associated with growth and in-
crease that of genes associated with stress. Further, gene expression in TCM suggests
an accumulation of cells in the M/G1, post-mitotic phase of the cell cycle. Finally, TCM
tends to engage nitrogen catabolite repression, probably because of its glutamine con-
tent.

Surprisingly, both CO_2 and cAMP moderate or reduce most of the effects of TCM
on gene expression. The fact that either of these components can act as the necessary
partner with TCM for capsule induction, but their overall effects on gene expression are
opposite that of TCM, poses intriguing questions: When CO_2 is added to TCM, which of

the resulting gene expression changes are critical for capsule induction and do those
changes work by dialing down the effects of TCM on key genes, by reinforcing the ef-
fects of TCM on the few genes that respond in the same direction, or by acting on
genes that do not respond to TCM?

Both phosphate and nitrogen availability have been suspected as regulators of
capsule size. Both nutrients activate sensors that stimulate the PKA pathway.[37-39] It has
also been reported that expression of phosphate acquisition genes is strongly induced
in CAP medium and that subsequent addition of KH_2PO_4 reduces capsule width as well
as cell size.[10] We found that phosphate transporters were also strongly induced by
TCM, although this effect was much reduced by addition of CO_2 or cAMP.

Regarding nitrogen availability, it has also been reported that YNB medium (trace
nutrients, $(NH_4)_2SO_4$, and glucose) with bicarbonate and 5% CO_2 induces capsule, but
only if the $(NH_4)_2SO_4$ is replaced by arginine.[6] Both are considered preferred nitrogen
sources for *S. cerevisiae*[31], so the cells may be responding to amino acids, rather than
the quality of the nitrogen source. Further linking amino acids and capsule, several
genes with expression levels that are associated with capsule size encode proteins that
act in transport of amino acids or oligopeptides, such as CNAG_01119 and
CNAG_02539, which are orthologs of the *S. cerevisiae* oligopeptide transporter *PTR2*
and amino acid transporter *DIP5*, respectively (Tables S1).

As mentioned earlier, conditions that induce capsule also impact multiple cellular
response pathways. Nonetheless, we hypothesized that we might be able to identify
genes required for normal capsule size among those genes whose expression was best
correlated with capsule thickness and tested this using deletion mutants. Unfortunately,
there is no deletion mutant in the Madhani collection for many of the genes whose ex-
pression is most predictive of capsule thickness (Tables S1). Our own attempts to de-
lete five of these genes also failed, suggesting that they may be essential. Studying
their effects on capsule thickness, therefore, may require construction of over-
expression strains or development of more robust tunable promoter systems than are
currently available for *C. neoformans*.

When a mutant was available for a gene whose expression was statistically as-
sociated with capsule thickness, we assayed it for this characteristic. Unsurprisingly,
most mutants did not show a significant change in capsule size. Some of the corre-
sponding genes may indeed have no role in capsule elaboration, their correlation with
capsule size resulting from parallel pathways activated by the same stressful conditions
that lead to capsule growth. Others may play a role in capsule growth that can also be
filled by other proteins. Deletion of still others may affect capsule in ways that our meth-
ods do not detect, such as changes in glycan composition, structure, or other features.
An alternative approach that hypothesizes capsule phenotypes for genes whose ex-
pression pattern is similar to those of known capsule genes was recently reported to be
effective, with 6 of 12 tested mutants having altered capsule width.[40]

We did identify two genes, not previously known to influence capsule, whose de-
letion affects capsule size in inducing conditions. Deletion of CNAG 00368 reduced
mean capsule thickness by 0.44 um. This gene encodes an ortholog of *S. cerevisiae*
Vps53, which is involved in recycling proteins from endosomes to the late Golgi. Vps53

not been studied in Cryptococcus, but a protein that is part of a different complex involved in recycling proteins from endosomes, Vps23, is known to be required for capsule elaboration.[41] Reinforcing the theme of protein and peptide recycling, deletion of CNAG_05977 (proteasome activator subunit 4) reduced capsule size by 0.54 um. Future work will be needed to define the specific mechanisms by which these genes influence capsule.

Our dataset of matched RNA-Seq time courses and capsule thickness measurements has yielded insights into the environmental signals and gene expression changes that affect capsule size. However, this is only the beginning. We expect that future analyses by our group and other researchers will yield additional insights, not only into capsule biology but into numerous aspects of cryptococcal physiology, many with importance for disease.

METHODS

Additional details for all methods can be found in the online supplement.

Cell growth

To maximize reproducibility of RNA-seq and capsule imaging studies, cell recovery from frozen stocks, initial culture in YPD, inoculation into preconditioned media, and growth followed strictly controlled protocols. These methods are detailed in the Supplemental Material.

Microscopy and manual image annotation

1-ml samples were collected for imaging from the stock cell suspension prior to inoculation of flasks or from cultures at 1440 min, fixed, resuspended in PBS, and mixed with India ink (5 parts cells:2 parts ink) for brightfield microscopy. Images were manually annotated using a custom annotation interface written in Mathematica / Wolfram Language (available on request). Fifteen fields were annotated for each replicate of each combination, yielding an average of 96 annotated cells per replicate (min 31; max 361; standard deviation 54).

Analysis of capsule thickness in gene deletion mutants

Effects of deletion mutants on capsule thickness were evaluated using a Linear Mixed Models framework with individual cells' thicknesses as the datapoints. The model had fixed effects for intercept and genotype; each specific biological sample was used as a grouping variable with a random intercept for each group.

RNA-Seq

Libraries were constructed using the NEBNext Ultra Directional RNA Library Prep Kit from Illumina and samples were pooled at 10 nM. The pools were sequenced on a NextSeq 500 using the High75v2 kit as 1x75 (single end).

RNA-Seq Computational Pipeline and Analysis

Documented code implementing the process described below is available on our public github repository: https://github.com/BrentLab/brentlabRnaSeqTools. Reads were aligned with Novoalign (version 4.03.02) and quantified with HTSeq (0.9.1) using the

FungiDB KN99α genome sequence (FungiDB accession ASM221672v1). Custom
scripts were used to verify the strain in each sample by calculating coverage over the
open reading frame of the putatively deleted genes and over marker genes, ensuring
that the former were absent and the latter present. Documented code and parameters
for strain validation and QC can be found here:
https://brentlab.github.io/brentlabRnaSeqTools/articles/QC_Library_Quality.html#crypto
coccus. RNA-Seq samples that passed strain validation were subjected to two phases
of QC. In Phase 1, files were labeled as 'passing' if they contained at least 10^6 reads
aligned to protein coding regions and less than 7% of all reads failed to align.
In Phase 2, we evaluated replicate agreement using the Regularized Log Ex-
pression (RLE) [42]. First, we used DESeq2 (version 1.34.0) to estimate the effect of the
library date (the known batch effect). We removed the batch effect using the DESeq co-
efficients for the library dates such that the data were standardized to a single date, re-
sulting in adjusted expression levels on a $\log_2$ normalized scale. To compute the RLE
value for a given gene, the median expression level of that gene, across all samples in
a replicate set, was subtracted from the expression level of the gene in each sample.
For each sample, we then calculated the interquartile range of the distribution of these
RLE values across genes. If the interquartile range of RLEs of a given sample was
greater than 1, indicating that more than half of genes deviated from their respective
medians by a factor of 2 in adjusted count, the sample was considered an outlier and
failed for replicate agreement. 22 samples (2.75%) failed replicate agreement and were
discarded.
501

Gene set overrepresentation analysis
Differential expression analysis was conducted using DESeq2 with a model containing
the main effect of time, represented as a categorical variable, and the interaction effects
of time with environmental signals. The interaction effects indicate how the normal effect
of time in standard laboratory conditions (YPD, no CO_2, 30 °C, no cAMP, no HEPES) is
modulated by the presence of other factors at each time point. For each time-
point:signal interaction, a gene was considered to be differentially expressed if the ab-
solute value of its shrunken $\log_2$ fold change (LFC) reached 1.0. (DESeq shrinkage re-
duces the estimated LFCs of factors with greater replicate-to-replicate variance).
Overrepresentation analysis was carried out using GOTermFinder (see online supple-
ment for additional details).[43] Gene annotations came from UniProtKB, release
2020_01, and were supplemented with several annotations derived from literature (see
Online Supplement and File S5).

Selection of potential capsule-associated genes: Chi-squared method
The following procedure was carried out separately for samples at each time point. To
calculate the χ^2 statistic for each gene, we first classified all samples with average cap-
sule width greater than 1.13 µm (3 S.D. above the mean of all YPD conditions) as in-
duced. Other samples were classified as uninduced. Samples were also classified by
high or low expression of each gene, yielding a 2x2 contingency table. For each gene,
we tried all possible thresholds of high versus low expression, constructed the contin-
gency table, and calculated the χ^2 statistic. Finally, we chose the threshold that yielded
largest χ^2 statistic. These statistics were only used to characterize the degree of associ-

ation between the expression of a gene in a sample and that sample's induction status. They were not used for hypothesis testing.

Genes were ranked for likelihood of the corresponding deletion strain exhibiting a capsule phenotype by the maximum of their χ^2 values, across all time points. Genes with high χ^2 statistics were selected for testing. For genes that were positively correlated with capsule size, the gene deletion mutant was tested in inducing conditions (RPMI, 37 °C, 5% CO_2) with the prediction that deletion would reduce capsule size compared to WT grown in the same conditions. For negatively correlated genes, the gene deletion mutant was tested in non-inducing conditions (YPD, 37 °C, 5% CO_2) and in 'almost-inducing' conditions (RPMI, 37 °C, room air); the rationale for the latter was that the conditions were quite close to inducing conditions and perhaps favor capsule synthesis.

Forming metagenes as features for machine learning

From the original gene expression data matrix, we sought to decrease the number of features by filtering and combining genes. We first removed low variance genes by filtering out genes whose expression in >= 95% samples was within one $\log_2$ of their mean, indicating that less than 5% of samples showed substantial changes. We next combined highly correlated genes into metagenes (correlation threshold > 0.8). The "expression level" of a metagene was then set to the mean of the expression levels of its constituent genes. We repeated this process to combine genes into metagenes until we had combined all highly correlated genes into metagenes.

COMPETING INTERESTS

The authors declare that they have no competing interests.

AVAILABILITY OF DATA AND MATERIALS

All RNA-Seq data are available from NCBI GEO under accession numbers GSE226255, GSE226637, or GSE226651, which are subseries of GSE226656. All capsule size measurements are available as supplemental files. Corresponding India ink images and mutant strains are available on request.

FUNDING

This work was supported by NIH awards AI087794 (Doering) and GM141012 (Brent).

REFERENCES CITED

1.   Denning, D.W. Global incidence and mortality of severe fungal disease. *Lancet Infect Dis* (2024).
2.   Kwon-Chung, K.J. et al. *Cryptococcus neoformans* and *Cryptococcus gattii*, the Etiologic Agents of Cryptococcosis. *Cold Spring Harb Perspect Med* **4** (2014).
3.   Maier, E.J. et al. Model-driven mapping of transcriptional networks reveals the circuitry and dynamics of virulence regulation. *Genome Res* **25**, 690-700 (2015).

4.   O'Meara, T.R. & Alspaugh, J.A. The *Cryptococcus neoformans* capsule: a sword and a shield.
Clinical microbiology reviews **25**, 387-408 (2012).
5.   Littman, M.L. Capsule synthesis by Cryptococcus neoformans. *Trans N Y Acad Sci* **20**, 623-48
(1958).
6.   Granger, D.L., Perfect, J.R. & Durack, D.T. Virulence of *Cryptococcus neoformans*. Regulation of
capsule synthesis by carbon dioxide. *The Journal of clinical investigation* **76**, 508-16 (1985).
7.   Vartivarian, S.E. et al. Regulation of cryptococcal capsular polysaccharide by iron. *J Infect Dis*
167, 186-90 (1993).
8.   Zaragoza, O. & Casadevall, A. Experimental modulation of capsule size in *Cryptococcus*
neoformans. *Biol Proced Online* **6**, 10-15 (2004).
9.   Zaragoza, O., Fries, B.C. & Casadevall, A. Induction of capsule growth in *Cryptococcus*
neoformans by mammalian serum and CO_2. *Infect Immun* **71**, 6155-64 (2003).
10.   Denham, S.T. et al. A dissemination-prone morphotype enhances extrapulmonary organ entry
by *Cryptococcus neoformans*. *Cell Host Microbe* **30**, 1382-1400 e8 (2022).
11.   Farhi, F., Bulmer, G.S. & Tacker, J.R. *Cryptococcus neoformans* IV. The Not-So-Encapsulated
Yeast. *Infect Immun* **1**, 526-31 (1970).
12.   Dykstra, M.A., Friedman, L. & Murphy, J.W. Capsule size of *Cryptococcus neoformans*: control
and relationship to virulence. *Infect Immun* **16**, 129-35 (1977).
13.   Caza, M. & Kronstad, J.W. The cAMP/Protein Kinase a Pathway Regulates Virulence and
Adaptation to Host Conditions in *Cryptococcus neoformans*. *Front Cell Infect Microbiol* **9**, 212
(2019).
14.   Zaman, S., Lippman, S.I., Zhao, X. & Broach, J.R. How Saccharomyces responds to nutrients.
Annu Rev Genet **42**, 27-81 (2008).
15.   Conrad, M. et al. Nutrient sensing and signaling in the yeast *Saccharomyces cerevisiae*. *FEMS*
microbiology reviews **38**, 254-99 (2014).
16.   Cramer, K.L., Gerrald, Q.D., Nichols, C.B., Price, M.S. & Alspaugh, J.A. Transcription factor Nrg1
mediates capsule formation, stress response, and pathogenesis in *Cryptococcus neoformans*.
Eukaryot Cell **5**, 1147-56 (2006).
17.   O'Meara, T.R. et al. Interaction of *Cryptococcus neoformans* Rim101 and protein kinase A
regulates capsule. *PLoS Pathogens* **6**, e1000776 (2010).
18.   Alspaugh, J.A. et al. Adenylyl cyclase functions downstream of the Galpha protein Gpa1 and
controls mating and pathogenicity of *Cryptococcus neoformans*. *Eukaryot Cell* **1**, 75-84 (2002).
19.   Hu, G. et al. Transcriptional regulation by protein kinase A in *Cryptococcus neoformans*. *PLoS*
Pathog **3**, e42 (2007).
20.   D'Souza, C.A., Hagen, F., Boekhout, T., Cox, G.M. & Heitman, J. Investigation of the basis of
virulence in serotype A strains of *Cryptococcus neoformans* from apparently immunocompetent
individuals. *Curr Genet* **46**, 92-102 (2004).
21.   Haynes, B.C. et al. Toward an Integrated Model of Capsule Regulation in *Cryptococcus*
neoformans. *PLoS Pathogens* **7**, e1002411 (2011).
22.   Gish, S.R. et al. Computational Analysis Reveals a Key Regulator of Cryptococcal Virulence and
Determinant of Host Response. *MBio* **7** (2016).
23.   Nielsen, K. et al. Sexual cycle of *Cryptococcus neoformans var. grubii* and virulence of congenic a
and alpha isolates. *Infect Immun* **71**, 4831-41 (2003).
24.   Love, M.I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. *Genome Biol* **15**, 550 (2014).
25.   Denham, S.T. et al. Regulated Release of Cryptococcal Polysaccharide Drives Virulence and
Suppresses Immune Cell Infiltration into the Central Nervous System. *Infect Immun* **86** (2018).

26. Petzold, E.W. et al. Characterization and regulation of the trehalose synthesis pathway and its
importance in the pathogenicity of *Cryptococcus neoformans*. *Infect Immun* **74**, 5877-87 (2006).
27. Spellman, P.T. et al. Comprehensive identification of cell cycle-regulated genes of the yeast
Saccharomyces cerevisiae by microarray hybridization. *Mol Biol Cell* **9**, 3273-97 (1998).
28. Santiago-Tirado, F.H., Peng, T., Yang, M., Hang, H.C. & Doering, T.L. A Single Protein S-acyl
Transferase Acts through Diverse Substrates to Determine Cryptococcal Morphology, Stress
Tolerance, and Pathogenic Outcome. *PLoS Pathog* **11**, e1004908 (2015).
29. Cottrell, T.R., Griffith, C.L., Liu, H., Nenninger, A.A. & Doering, T.L. The pathogenic fungus
Cryptococcus neoformans expresses two functional GDP-mannose transporters with distinct
expression patterns and roles in capsule synthesis. *Eukaryot Cell* **6**, 776-85 (2007).
30. Hicks, J.K., Bahn, Y.S. & Heitman, J. Pde1 phosphodiesterase modulates cyclic AMP levels
through a protein kinase A-mediated negative feedback loop in *Cryptococcus neoformans*.
Eukaryot Cell **4**, 1971-81 (2005).
31. Ljungdahl, P.O. & Daignan-Fornier, B. Regulation of amino acid, nucleotide, and phosphate
metabolism in *Saccharomyces cerevisiae*. *Genetics* **190**, 885-929 (2012).
32. Klengel, T. et al. Fungal adenylyl cyclase integrates CO2 sensing with cAMP signaling and
virulence. *Current biology : CB* **15**, 2021-6 (2005).
33. Liu, O.W. et al. Systematic genetic analysis of virulence in the human fungal pathogen
Cryptococcus neoformans. *Cell* **135**, 174-88 (2008).
34. Zaragoza, O. & Nielsen, K. Titan cells in *Cryptococcus neoformans*: cells with a giant impact. *Curr*
Opin Microbiol **16**, 409-13 (2013).
35. Saidykhan, L. et al. An in vitro method for inducing titan cells reveals novel features of yeast-to-
titan switching in the human fungal pathogen *Cryptococcus gattii*. *PLoS Pathog* **18**, e1010321
(2022).
36. Bahn, Y.S., Kojima, K., Cox, G.M. & Heitman, J. Specialization of the HOG pathway and its impact
on differentiation and virulence of *Cryptococcus neoformans*. *Mol Biol Cell* **16**, 2285-300 (2005).
37. Xue, C., Bahn, Y.S., Cox, G.M. & Heitman, J. G protein-coupled receptor Gpr4 senses amino acids
and activates the cAMP-PKA pathway in *Cryptococcus neoformans*. *Molecular biology of the cell*
17, 667-79 (2006).
38. Steyfkens, F., Zhang, Z., Van Zeebroeck, G. & Thevelein, J.M. Multiple Transceptors for Macro-
and Micro-Nutrients Control Diverse Cellular Properties Through the PKA Pathway in Yeast: A
Paradigm for the Rapidly Expanding World of Eukaryotic Nutrient Transceptors Up to Those in
Human Cells. *Front Pharmacol* **9**, 191 (2018).
39. Samyn, D.R. et al. Mutational analysis of putative phosphate- and proton-binding sites in the
Saccharomyces cerevisiae Pho84 phosphate:H(+) transceptor and its effect on signalling to the
PKA and PHO pathways. *Biochem J* **445**, 413-22 (2012).
40. O'Meara, M.J. et al. CryptoCEN: A Co-Expression Network for Cryptococcus neoformans reveals
novel proteins involved in DNA damage repair. *PLoS Genet* **20**, e1011158 (2024).
41. Hu, G. et al. *Cryptococcus neoformans* requires the ESCRT protein Vps23 for iron acquisition
from heme, for capsule formation, and for virulence. *Infect Immun* **81**, 292-302 (2013).
42. Gandolfo, L.C. & Speed, T.P. RLE plots: Visualizing unwanted variation in high dimensional data.
PLoS One **13**, e0191629 (2018).
43. Boyle, E.I. et al. GO::TermFinder--open source software for accessing Gene Ontology
information and finding significantly enriched Gene Ontology terms associated with a list of
genes. *Bioinformatics* **20**, 3710-5 (2004).

        FIGURE LEGENDS

Figure 1 (A) Average effects of environmental signals on capsule size, relative to YPD, in a linear model without interaction terms (N = 232). *** P<0.001 compared to YPD. (B) Capsule sizes for all combinations of media, cAMP, and CO_2 signals. Horizontal line: median; box: quartile 1 to quartile 3; fence: 1.5 times quartile range; diamonds: individual outlier points. (C) Conditions in which individual signals had their biggest effects on capsule size. None of these conditions includes HEPES buffer. India ink-stained micrographs of cells grown in each condition, chosen to match the mean capsule width, are shown below. All images are to the same scale; scale bar, 5mm.

Figure 2 (A-C) Principal Components (PC) plots. (A) PC1 and PC2. (B) 24 h samples only, PC1 and PC2. (C) 24 h samples in tissue culture medium (TCM) only, PC1 and PC3. (D) Log_2 fold changes (LFCs) for all genes in the indicated categories in DMEM relative to YPD at the same time point (DMEM / YPD), in DMEM with CO_2 relative to DMEM without CO_2 (DMEM + CO_2 / DMEM), or in DMEM with CO_2 relative to YPD (DMEM + CO_2 / YPD). Colors indicate the average LFC for all genes in the category, including those that are not significantly differentially expressed. Most LFCs were in the -1 to +1 range; in order to maximize visual discriminability those outside that range were truncated at -1 or +1. Lower case: GO terms; Capitalized terms: other gene sets (see Supplemental File S5 and Methods). Group 1: processes that are strongly upregulated by DMEM; Group 2: processes that are strongly downregulated by DMEM. Green text highlights categories that are expanded in subsequent panels. N=695 samples (E-F) LFCs of individual genes (denoted by the numeric part of CKF44 gene IDs) that are annotated as being involved in carbohydrate synthesis (C) or chromosome segregation (D). Colored lines indicate genes with absolute LFC > 1 at some time point. Asterisk: Gene name provided is that of the *S. cerevisiae* ortholog. N=695.

Figure 3 (A) Capsule width or (B) cell body radius as a function of added cAMP. Mean -/+ SD is shown. (C) Capsule width or (D) cell body radius in the indicated strains. Each bar represents 2-3 independent experiments done on different days. (E) Log_2 fold changes (LFC) for all genes in the indicated functional categories in RPMI (no cAMP) relative to YPD at the same time point (RPMI / YPD), in RPMI with 20 mM cAMP relative to RPMI without cAMP at the same time point (RPMI + cAMP / RPMI), or in RPMI with 20 mM cAMP relative to YPD at the same time point (RPMI + cAMP / YPD). Colors indicate the average LFC for all genes in the category, including those that are not significantly differentially expressed. LFCs were truncated at -1 and +1 to maximize visual discriminability in that range. (F) LFCs of individual genes (denoted as in Fig. 2) that are repressed by the Nitrogen Catabolite Repression (NCR) pathway. Colored lines indicate genes with absolute LFC > 1 at some time point. (G) Correlations between the effects of tissue culture media, CO_2, and cAMP on expression of all genes. DMEM and RPMI are relative to YPD, all in room air. CO_2 and cAMP are added to DMEM and RPMI, respectively, and compared to the tissue culture medium alone. Panels E-G are based on a model with 695 RNA-Seq samples.

Figure 4 (A) Example gene with high χ^2, reflecting the visual relationship between its expression level and capsule size. Cells with capsules larger than the horizontal lines are considered induced while genes with expression levels to the right of the vertical red

704 lines are considered highly expressed. Expression levels are $\log_2$ of counts, +1 pseudo
705 count, after normalization by DESeq2 and batch effect removal. (B-C) Example micro-
706 graphs of cells of the indicated strain, stained with India ink. All images are to the same
707 scale; scale bar, 5 mm.

A Capsule Size: OLS coefficients for signals

B

C cAMP, 5% CO_2, 37° | cAMP, 5% CO_2, 30° | DMEM, cAMP, 37° | RPMI, no CO_2, 30°

A χ^2=113

capsule width

CNAG_00368 expression level

B CNAG_00368 deletion strain

C Wild type KN99α strain