Structural rather than catalytic role for mitochondrial respiratory chain supercomplexes

Michele Brischigliaro[1,2,*,¶], Alfredo Cabrera-Orefice[3], Susanne Arnold[3,4,#], Carlo Viscomi[1,#], Massimo Zeviani[2,5,†,#], Erika Fernández-Vizarra[1,2,#,*, §]

[1] Department of Biomedical Sciences, University of Padova, Padova, Italy
[2] Veneto Institute of Molecular Medicine, Padova, Italy
[3] Radboud Institute for Molecular Life Sciences, Radboud University Medical Center, Nijmegen, The Netherlands
[4] Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated Diseases (CECAD), University of Cologne, Cologne, Germany
[5] Department of Neurosciences, University of Padova, Padova, Italy

* Correspondence to: Michele Brischigliaro (michele.brischigliaro@unipd.it) and Erika Fernández-Vizarra (erika.fernandezvizarra@unipd.it)
[#] Senior author
[§] Lead contact: Erika Fernandez-Vizarra
[¶] Present address: Department of Neurology, University of Miami Miller School of Medicine, Miami, FL, USA
[†] Present address: Institute for Maternal and Child Health, IRCCS "Burlo Garofolo", Trieste, Italy

ABSTRACT

Mammalian mitochondrial respiratory chain (MRC) complexes are able to associate into

quaternary structures named supercomplexes (SCs), which normally coexist with non-

bound individual complexes. The functional significance of SCs has not been fully clarified

and the debate has been centered on whether or not they confer catalytic advantages to

the non-bound individual complexes. Mitochondrial respiratory chain organization does not

seem to be conserved in all organisms. In fact, and differently from mammalian species,

mitochondria from insect tissues are characterized by low amounts of SCs, despite the

high metabolic demands and MRC activity shown by these mitochondria. Here, we show

that attenuating the biogenesis of individual respiratory chain complexes was

accompanied by increased formation of stable SCs, which are missing in *Drosophila*

melanogaster in physiological conditions. This phenomenon was not accompanied by an

increase in mitochondrial respiratory activity. Therefore, we conclude that SC formation is

necessary to stabilize the complexes in suboptimal biogenetic conditions, but not for the

enhancement of respiratory chain catalysis.

KEYWORDS: Mitochondrial respiratory chain, supercomplexes, respiratory chain complex

biogenesis, mitochondrial respiratory activity

INTRODUCTION

Mitochondria are the organelles providing most of the cellular energy in form of adenosine triphosphate (ATP) in aerobic eukaryotes. The molecular machinery responsible for energy transformation is the oxidative phosphorylation (OXPHOS) system, which is canonically composed of five multiprotein complexes embedded in the inner mitochondrial membrane. OXPHOS consists of two tightly regulated processes: electron transport and ATP synthesis. Electron transport takes place between complexes I-IV and two mobile electron carriers (coenzyme Q and cytochrome *c*). During electron transport, complexes I, III and IV pump protons from the mitochondrial matrix to the intermembrane space, generating a proton gradient that provides the protonmotive force exploited by complex V to synthesize ATP. In mammalian mitochondria, mitochondrial respiratory chain (MRC) complexes I, III and IV can interact with each other forming supramolecular structures described generally by the term 'supercomplexes' [1,2]. MRC supercomplexes (SCs) can have different stoichiometries and compositions, ranging from the binding of only two complexes, such as the I_1III_2 and III_2IV_1 SCs [3,4], to higher-order associations between complexes I, III and IV, with the SC of $I_1III_2IV_1$ stoichiometry known as the 'respirasome' [1,5-9]. Now that the association of individual MRC complexes into supramolecular structures is well-established, with structures of several SC species being resolved, the debate is centered on what the functional significance of these structures might be.

Several possible roles have been proposed for SCs. First, it was suggested that the association between complex I (CI) and the obligate dimer of complex III (CIII$_2$) would allow substrate channeling, sequestering a dedicated coenzyme Q (CoQ) pool and allowing a more efficient electron transfer between the two complexes, while separating this electronic route from those of $FADH_2$-linked dehydrogenases (e.g. complex II) to the CIII$_2$ not bound to CI [1,10-15]. This increased efficiency would in turn decrease electron leak and, as a consequence, produce less reactive oxygen species (ROS) than the individual free complexes [16,17]. However, the available high-resolution respirasome structures show that the distance between the CoQ binding sites in CI and CIII$_2$ are too far apart to allow substrate channeling [4,18]. In addition, exogenously added CoQ was necessary to sustain CI activity in the purified I_1III_2 SC [3], arguing against a tightly bound and segregated CoQ pool as a functional component of the SC. A large body of work from the late 1960's to the 1980's, resulted in the 'random collision model' to explain electron transfer between the respiratory chain complexes, and in the evidence that CoQ is present as an undifferentiated pool[19-23]. More recently, additional proof of the non-compartmentalized

electronic routes from CI to $CIII_2$ and from complex II (CII) to $CIII_2$, came from kinetic

measurements in sub-mitochondrial particles. In these systems, MRC organization in SCs

was conserved but *b*- and *c*-type cytochromes in $CIII_2$ were equally accessible to CI-linked

and CII-linked substrates [24], and CoQ reduced by CI in the respirasomes was able to

reach and readily reduce external enzymes to the SCs [25]. In addition, growing evidence

supports the notion that different MRC organizations exist *in vivo*, where varying

proportions of SC vs. free complexes do not result in separate and distinct CI-linked and

CII-linked respiratory activities [26-29]. This is in contrast with an idea that segregation into

different types of SCs and in individual complexes to explains their functional interplay for

the adaptation of OXPHOS activity [13]. The physical proximity of $CIII_2$ and CIV has also

been suggested to promote faster electron transfer kinetics via cytochrome *c* [4,30,31],

although this is a matter of debate as well [32,33].

The second main explanation to justify the existence of SCs is that they play a structural

function, stabilizing the individual complexes [34,35] and/or serving as a platform for the

efficient assembly of the complexes, with a special relevance for the biogenesis of

mammalian CI [36-38].

Notably, the MRC structural organization, especially the stoichiometry, arrangements and

stability of the SCs, may not be conserved in all eukaryotic species [39-42]. This is the case

even within mammals, as human cells and tissues barely contain free CI, which is rather

contained in SC I_1+III_2 and the respirasome [29,37]. In contrast, other mammalian

mitochondria (from bovine, ovine, rat or mouse) contain larger amounts of CI in its free

form, even if the majority is still mostly in the form of SCs [2,17,28,43-45]. The distribution of the

MRC complexes between free complexes and SCs seems to differ even more in non-

mammalian animal species. Several reptile species contain a very stable SC $I+III_2$ that

lacks CIV [28], and in *Drosophila melanogaster* practically all of CI is free, SCs being almost

completely absent [46,47]. However, comparative studies of MRC function in diverse animal

species suggest that higher amounts and stability of the SCs do not correlate with

increased respiratory activity/efficiency and/or reduced ROS production [28,47].

Here we show that SCs can be stably formed in *D. melanogaster* mitochondria upon mild

perturbations of individual CIV, $CIII_2$ and CI biogenesis. This finding enabled us to test

whether increased SC formation translated into enhanced respiration proficiency.

However, MRC performance of fruit fly mitochondria did not change regardless of the

presence or absence of SCs. These observations have led us to conclude that: 1) the

efficiency in the assembly of the individual complexes is likely to be the main determinant

117 of SC formation and 2) these supramolecular complexes play a more relevant role in

118 maintaining the stability and/or supporting the biogenesis of the MRC than in promoting

119 catalysis.

RESULTS

D. melanogaster MRC organization does rely on SC formation under physiological conditions

To obtain a detailed characterization of MRC organization in _D. melanogaster,_ we isolated mitochondria from wild-type adults and, after solubilization with digitonin, we performed Blue-Native Gel Electrophoresis (BNGE) followed by mass spectrometry analysis of the gel lanes, using 'Complexome Profiling (CP)' [48] and thus obtaining a profile of peptide intensities from most OXPHOS subunits along the electrophoresis lane (Figure 1A). This allowed us to unequivocally determine the identity of the main bands that can be visualized by Coomassie staining of BNGE gels (Figure 1B). The identity of the bands corresponding to complex I (CI) and complex IV (CIV) was also confirmed using specific in-gel activity (IGA) assays (Figure 1B). These analyses verified that _D. melanogaster_ mitochondria contain extremely low amounts of high-molecular weight CI-containing SCs [46,47], using the same solubilization and electrophoresis conditions in which the SCs are readily detectable in mammalian cells and tissues [29,43,49]. As previously described [47], dimeric complex V (CV_2) is easily visualized by BNGE and present in similar amounts as monomeric CV in _D. melanogaster_ mitochondrial membranes solubilized with digitonin (Figure 1A, B). This CV_2 species is not a strongly bound dimer, as it disappears when the samples are solubilized using a harsher detergent such as dodecylmaltoside (DDM) [47] (Figure 1C). Conversely, CI is mainly found as a free complex in the native gels irrespective of whether the mitochondria had been solubilized with digitonin or DDM (Figure 1A, B and C). The other minor CI-containing band corresponding to the fraction associated with $CIII_2$, accounts for about 3% of the total amounts of CI and $CIII_2$, according to label-free mass spectrometry quantifications in the CP analysis and densitometric quantifications of the CI-IGA signals (Figure 1A and Figure S1A). CIV activity is absent in this band both in digitonin- and DDM-solubilized samples (Figure 1B, C), whereas it is present in the bands that correspond to individual CIV and dimeric CIV (CIV_2) detected both in digitonin- and DDM-treated samples, as well as in SC III_2IV_1, which is present only in the digitonin-solubilized samples. This is different from mammalian mitochondria in which SC III_2IV_1 is present also in DDM-solubilized mitochondria, probably due to the tight binding of $CIII_2$ to CIV through COX7A2L/SCAF1 [4,29,50]. The latter does not have a homolog in _D. melanogaster_ even though this species has three different COX7A isoforms (named COX7A, COX7AL and COX7AL2) that exhibit a tissue-specific expression pattern. CP analysis of _Drosophila_

mitochondria only detected COX7A (mammalian COX7A1 homolog) and COX7AL2

(mammalian COX7A2 homolog), whereas COX7AL, that is solely expressed in testis, was

not found. Therefore, SC I_1III_2 can be considered the only stable SC species in

physiological conditions in *D. melanogaster*, yet containing a minute fraction of the total CI

and $CIII_2$.

Perturbations of CIV assembly result in increased formation of SC I_1III_2

COA8 is a CIV assembly factor the defects of which cause isolated mitochondrial CIV

deficiency in human and mouse [51,52], as well as in *Drosophila melanogaster* [53,54].

Consistent with the role of Coa8 in CIV biogenesis, CP analysis of mitochondria from *Coa8*

knockout (*Coa8^KO*) flies showed a clear decrease in fully assembled CIV and in all the CIV-

containing species (Figure 2A, B) when compared to the corresponding wild-type (WT)

individuals (Figure 1A and Figure 2B). Curiously, CP also showed that in the *Coa8^KO*

mitochondria the amounts of SC I_1III_2 were noticeably increased (Figure 2A, B). In these

samples, complexes I and III_2 build a stable SC species containing ~16% of the total

amount of CI, as visualized by CI-IGA, as well as by western blot (WB) and

immunodetection of specific CI and $CIII_2$ subunits after BNGE in DDM-solubilized

mitochondria (Figure 2C, D, Figure S1A and S2A). We initially speculated that the ~5-fold

increase of SC I_1III_2 formation could be linked to the release of III_2 from SC III_2IV_1 induced

by the strong reduction in CIV amounts when Coa8 is absent.

To test this hypothesis, we modulated Coa8 expression via UAS/GAL4 system using RNAi

driven by a 'mild' ubiquitous GAL4 driver (*da-gal4*). With this system, the *Coa8* mRNA

levels were reduced to ~60% of the control (Figure 3A). However, these flies showed

comparable levels of fully assembled CIV (Figure 3B, Figure S2B). Interestingly, in this

case there was also an increased formation of SC I_1III_2 from ~3% in the control to ~10% in

the mild *Coa8^RNAi* (Figure 3B, C, Figure S1B). Therefore, both strong and weak

perturbations of CIV assembly produce an increased formation of CI-containing SCs in *D.*

melanogaster, irrespective of whether they result in CIV deficiency or not.

Enhanced formation of SC I_1III_2 does not result in increased respiratory rates.

SC formation was proposed to serve as a means to favor electron transfer between the

complexes and therefore increase the efficiency of CI-fueled respiration [1]. With this in

mind, the complete *Coa8^KO* and mild *Coa8^RNAi* fly mitochondria, which show increased

amounts of SC I_1III_2 compared to the WT controls, provide an excellent opportunity to test

this possibility. Oxygen consumption activities of fly homogenates in presence of different

substrates and inhibitors were analyzed by high-resolution respirometry (Figure 4A, B).

The significant decrease in CIV enzymatic activity in the $Coa8^{KO}$ (Figure 4C) was not

reflected by reduced oxygen consumption (Figure 4A). This could be explained as a result

of a high CIV excess in fly mitochondria, in which the observed 60% reduction in CIV

enzyme activity is still above the threshold at which the CIV defect determines lower

respiratory rates [55,56]. In contrast, the mild reduction in $Coa8$ mRNA levels did not result in

CIV enzymatic deficiency but produced a slight elevation in CI amounts and activity

(Figure S1B, Figure 4D), which could be related to the increased SC I_1III_2 formation.

However, the $Coa8\text{-}KD$ mitochondria did not show any differences in respiration with either

CI-linked or CII-linked substrates. Also, the increased and stable interactions between

complexes I and III_2 in the Coa8 deficient models did not produce a preferential utilization

of electrons coming from CI, which would be the prediction if SC formation increased

electron transfer efficiency [13].

Mild perturbation of $CIII_2$ biogenesis also enhances SC formation in *D. melanogaster*

To determine whether increased SC I_1III_2 formation was specific for CIV deficient flies, we

targeted $CIII_2$ by knocking down the expression of $Bcs1$. BCS1L, the human homolog, is

fundamental for a correct $CIII_2$ biogenesis, being responsible for the incorporation of the

catalytic subunit UQCRFS1 in the last steps of $CIII_2$ maturation [57,58]. To obtain a severe

$CIII_2$ defect in *D. melanogaster*, we crossed a 'strong' ubiquitous GAL4 driver (*act5c-gal4*)

line with a *UAS-Bcs1* RNAi responder line [59]. The knockdown efficiency was high, with a

~75% decrease in $Bcs1$ expression at the mRNA level (Figure 5A). In this model, *D.*

melanogaster development was severely impaired causing an arrest at the larval stage [59].

The strong $Bcs1^{RNAi}$ caused also a significant decrease in fully assembled $CIII_2$ levels

(Figure 5B, C, Figure S2C) and in $CIII_2$ enzymatic activity of about 50% (Figure 5D).

Consistent with the observed $CIII_2$ deficiency, both the CI- and CII- linked respiration rates

were significantly decreased by around 40% (Figure 5E).

In contrast, less pronounced decreases in $Bcs1$ expression (Figure 5F) by using the mild

da-gal4 driver instead, did not produce a noticeable $CIII_2$ enzymatic defect (Figure 5I).

However, the mild $Bcs1\text{-}KD$ mitochondria showed a very different pattern of CI distribution

than the controls (Figure 5G, H, Figure S1D), showing much higher levels of SC I_1III_2 as

well as the appearance of a new higher molecular weight band containing also CIV, as

revealed by WB and immunodetection analyses (Figure 5G, H), similarly to the

mammalian respirasome (SC $I_1III_2IV_1$). At the functional level, this was associated with a

~1.5-fold increase in CI enzymatic activity (Figure 5I), which is proportional to the increase

in total CI amounts (Figure S1D), and in higher CI-linked respiration rates but only by ~1.2-

fold (Figure 5J). CII-linked respiration was the same in the mild $Bcs1^{RNAi}$ samples as in the

controls. Therefore, the formation of respirasome-like SCs in these mitochondria did

neither increase the efficiency of electron transfer from CI nor determine a diversion of the

electronic routes giving preference to the SC-bound CI.

Mild perturbation of CI biogenesis also leads to increased SC assembly

To understand the effect of the strong and mild perturbations in CI biogenesis on MRC

organization in D. melanogaster, we employed a similar strategy as that for CIII (see

above). Crossing the strong ubiquitous *act5c-gal4* driver fly line with the *UAS-Ndufs4* RNAi

responder line, produced a decrease in *Ndufs4* mRNA expression of ~90% (Figure 6A).

Defects in *NDUFS4* are a major cause of CI deficiency-associated mitochondrial disease

in humans [60] and the mouse and *D. melanogaster* animal models display CI deficiency and

pathological phenotypes [61,62]. Accordingly, the strong reduction in Ndufs4 expression

observed in our models resulted in developmental arrest and a significant decrease in fully

assembled CI levels by ~40% (Figure 6B, C, Figure S1E) and in a proportional decrease in

NADH:CoQ oxidoreductase enzyme activity in the larvae (Figure 6D). BNGE analysis of

the strong $Ndufs4^{RNAi}$ *D. melanogaster* mitochondria, revealed the presence of a CI

subassembly, containing the core Ndufs3 subunit (Figure 6C, Figure S2E) but lacking

NADH-dehydrogenase activity (Figure 6B). This is similar to what is observed in NDUFS4-

deficient human and mouse, which accumulate the so-called ~830 kDa intermediate

lacking the N-module and stabilized by the NDUFAF2 assembly factor [63-65]. In contrast,

CIV levels and enzyme activity were significantly increased by 1.5-fold in the strong

Ndufs4-KD (Figure 6C, D, Figure S2E). CI-linked respiration measured in isolated

mitochondria from these flies was significantly lower than in the controls, whereas the CII-

linked respiration was comparable to the control (Figure 6E). These observations are

compatible with the isolated CI defect displayed by the strong *Ndufs4-KD* flies.

Conversely, when *Ndufs4* mRNA expression was reduced to about half of the control

levels (Figure 6F), a milder defect in CI abundance (Figure S1F, Figure S2F), which did

not affect either NADH:CoQ oxidoreductase enzymatic activity or respiratory capacity was

observed (Figure 6G, H, I, J) in the adult flies. However, smaller amounts of the inactive

sub-CI were still detectable (Figure 6H). Interestingly enough, this milder perturbation of CI

biogenesis by reducing the amounts of Ndufs4 also produced an increase in the formation

of SC I_1III_2 (Figure 6G, H), without any changes in respiratory performance compared to

the controls (Figure 6J).

DISCUSSION

The first description of MRC SCs in the early 2000's led to opposite opinions on whether these were real and functionally relevant entities. On the one hand, researchers argued that the random collision model and diffusion of the individual MRC complexes was well established experimentally and the idea of the SCs did not fit with these observations. On the other hand, others considered that these SCs were real entities and therefore they must have a functional relevance, mainly as a means to enhance electron transfer between the individual complexes. Presently, the existence of the SCs is not debated anymore, especially after the determination of the high resolution structures by cryo-electron microscopy (EM), first of the mammalian respirasomes (reviewed in [66]), followed by that of other mammalian SCs [3,4], and of mitochondrial respiratory chain SCs from other eukaryotic species [39-42,67,68]. However, whether SCs provide any catalytical advantage to the MRC or not, is still being debated and opposing views continue to exist [18,69-72]. The recent resolution of MRC SCs from different eukaryotic species has revealed that the relative arrangement of the complexes within the SCs and the bridging subunits varies substantially depending on the species. Therefore, given the conservation of the structures of the individual complexes, one could argue that if SC formation was of capital importance for MRC function, the way the complexes interact should be strictly conserved as well. Importantly, neither in the mammalian respirasomes, nor in the mammalian and plant SC I_1III_2 there is any evidence of substrate channeling, as the CoQ binding sites in CI and $CIII_2$ are too far apart to allow for this phenomenon to happen [3,4,18,41]. This is in agreement with different sets of functional data indicating that CoQ is interchangeable between the CI-containing SCs and the rest of the MRC [24,25,29,37]. Therefore, this contrasts with the possible segmentation of the CoQ pool - one dedicated to the respirasome and the other to the $FADH_2$-linked enzymes – which has been proposed to take place as a consequence of respirasome formation [12-14]. A further element against the notion that SC formation is essential for mitochondrial function, is the fact that in normal conditions the MRC of *D. melanogaster* is predominantly organized based on individual complexes, as shown by us in this work and by others [46,47]. The *Drosophila* organization was justified by a tighter packing within the mitochondrial cristae and a higher concentration of the mobile electron transporters, as a way to compensate for the lack of SCs [47]. However, inter-species differences in MRC organization do not appear to be of great relevance to determine the level of functionality [28]. For example, disaggregation of CI from the SCs in *A. thaliana*, where there are normally present, did not affect plant viability [73]. Altogether, these

observations argue against the strict requirement of SC formation to maintain MRC

function.

The second main proposed role for SC is that of stabilizing and/or favoring the assembly of

the individual complexes, especially that of CI [37,38,74]. Very recently, the *D. melanogaster*

CI structure has been solved [75,76]. This has provided a structural explanation as to why this

complex is majorly found as an individual entity, which seems to be due to a more stable

association of the NDUFA11 subunit in the *D. melanogaster* complex [76]. Subunit

NDUFA11 stabilizes the transmembrane helix anchoring the lateral helix of subunit MT-

ND5, bridging the two parts of the membrane arm of CI. In other systems, $CIII_2$ and, thus,

SC formation seem to be important for stabilizing the interaction of NDUFA11 in the CI

membrane domain instead [3,5,41,77]. However, in this work we show that there are ways to

induce the formation of CI-containing SCs in fruit fly mitochondria where they normally do

not exist. For example, whereas a strong impairment of $CIII_2$ biogenesis resulted in $CIII_2$

deficiency, decreased respiration and equal amounts of assembled CI, a mild perturbation

of $CIII_2$ biogenesis did not result in any MRC deficiency but rather in increased CI total

amounts, mainly due to enhanced SC formation. Similarly, strong decreases in Ndufs4

expression resulted in low CI amounts and activity, but milder decreases did not affect CI

function although CI was redistributed into supramolecular species. In contrast, both the

complete KO and mild KD of *Coa8*, induced a significantly increased formation of CI-

containing SCs. Even though the *Coa8^KO* flies display a significant reduction in CIV

amounts and activity, the loss of COA8 is associated with milder phenotypes in humans,

mouse and flies compared with the lack or dysfunction of other CIV assembly factors

[51,52,54]. Therefore, we propose that a partial loss of CIV assembly also induces the

formation of SCs in *D. melanogaster*. These observations are in line with the idea that in

situations of suboptimal MRC complex biogenesis, formation of CI-containing SCs could

be a way to structurally stabilize the system and preserve its function [65,78]. In the context of

the cooperative assembly model, where partially assembled complexes get together

before completion forming SC precursors [38], one could envision that slower assembly

kinetics would increase the chance of interactions at early stages, letting SC assembly

occur in a stable way. Difference in the kinetics of CIV assembly have been observed

between human and mouse fibroblasts, which is slower in the human cells [79]. Therefore,

differences in assembly kinetics could also explain the observed different amounts and

stoichiometries of the MRC SCs in different organisms. This is an interesting open

question that will deserve further future investigation. In any case, this enhanced SC

formation did not translate in an increase in respiratory function nor in a change in
substrate preference in any of the tested models. If CI-containing SC formation enhanced
respiratory activity significantly, we should have had detected a noticeable increase in CI-
linked respiration in all the models of mild perturbation of CIV, CIII$_2$ and CI. Although we
did observe higher respiration with CI-linked substrates in the mild *Bcs1*-KD, this was even
lower to the increase in CI enzymatic activity and total abundance.
Therefore, we conclude that the main role of SC formation is to provide structural stability
to the MRC, principally for CI, rather than to enhance electron transfer between the
complexes during respiration.
340
341

343 **KEY RESOURCES TABLE**

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse monoclonal anti-NDUFS3	Abcam	Ab14711; RRID:AB_301429
Rabbit polyclonal anti-COXIV	Cell Signaling Technology	#4844; RRID:AB_2085427
Rabbit polyclonal anti-UQCR-C2	Dr. Edward Owusu-Ansah, Columbia University, NY	N/A
Rabbit polyclonal anti-SdhA	Dr. Edward Owusu-Ansah, Columbia University, NY	N/A
Chemicals, peptides, and recombinant proteins		
3,30-Diaminobenzidine tetrahydrochloride hydrate	Merck	D5637
6-Aminocaproic acid	Merck	A2504
Acetyl coenzyme A lithium salt	Merck	A2181
Antimycin A	Merck	A8674
BSA (fatty acid free)	Merck	A6003
Catalase from bovine liver	Merck	C9322
Coenzyme Q1	Merck	C7956
Cytochrome c from equine heart	Merck	C7752
D-Mannitol	Merck	M4125
DCIP (2,6-Dichloroindophenol Sodium Salt Hydrate)	Merck	D1878
Decylubiquinone	Merck	D7911
Digitonin, High Purity	Calbiochem	300410
DTNB (5,5′-Dithiobis(2-nitrobenzoic acid))	Merck	D218200
EDTA (Ethylenediaminetetraacetic acid disodium salt dihydrate)	Merck	E1644
EGTA (Ethylene glycol-bis(2-aminoethylether)-N,N,N′,N′-tetraacetic acid)	Merck	E3889
HEPES (N-(2-Hydroxyethyl)piperazine-N′-(2-ethanesulfonic acid), 4-(2-Hydroxyethyl)piperazine-1-ethanesulfonic acid)	Merck	H3375
KCN (Potassium cyanide)	Merck	31252
Magnesium chloride hexahydrate	Merck	M2670
Malonic acid	Merck	M1296
n-Dodecyl-beta-maltoside (DDM)	Merck	D4641
NADH (β-Nicotinamide adenine dinucleotide, reduced dipotassium salt)	Merck	N4505
NativePAGE™ Cathode Buffer Additive	Thermo Fisher Scientific	BN2002
NativePAGE™ Running Buffer	Thermo Fisher Scientific	BN2001
Nitrotetrazolium Blue chloride	Merck	N6876
Oligomycin from Streptomyces diastatochromogenes	Merck	O4876
Oxaloacetic acid	Merck	O4126
Phospho(enol)pyruvic acid monopotassium salt	Merck	860077
Potassium borohydride	Merck	438472

Potassium phosphate dibasic	Merck	P2222
Potassium phosphate monobasic	Merck	P5655
Rotenone	Merck	R8875
Sodium hydrosulphite	Merck	157953
Sodium succinate	Merck	S2378
Sucrose	Merck	S7903
TRIS base	Merck	T1503
Tween-20	Merck	P7949
Critical commercial assays		
GoScript® Reverse Transcriptase Kit	Promega	A5001
GoTaq® qPCR Master Mix	Promega	A6001
Experimental models: Organisms/strains		
D. melanogaster strain *act5c-gal4*	BDSC	4414
D. melanogaster strain *da-gal4*	BDSC	8641
D. melanogaster strain UAS-Ndufs4 RNAi	VDRC	101489
D. melanogaster strain UAS-Bcs1 RNAi	BDSC	51863
D. melanogaster strain UAS-Coa8 RNAi	VDRC	100605
D. melanogaster strain Coa8 KO	Wellgenetics Inc.	N/A
Oligonucleotides		
Bcs1-Fw-qPCR: CTGAATGTTGCGCCAGAG	Thermo Fisher Scientific	N/A
Bcs1-Rv-qPCR: GACGAATGCTGCGTCGAT	Thermo Fisher Scientific	N/A
Coa8-Fw-qPCR: CAATAAGCGCTTCTACGAGGA	Thermo Fisher Scientific	N/A
Coa8-Rv-qPCR: CCAGTTCTTGTCGAGGAACG	Thermo Fisher Scientific	N/A
Ndufs4-Fw-qPCR: AAGATCACCGTGCCGACTG	Thermo Fisher Scientific	N/A
Ndufs4-Rv-qPCR: GACAATGGGTCGCCGCTG	Thermo Fisher Scientific	N/A
Rp49-Fw-qPCR: ATCGGTTACGGATCGAACAA	Thermo Fisher Scientific	N/A
Rp49-Rv-qPCR: GACAATCTCCTTGCGCTTCT	Thermo Fisher Scientific	N/A
Software and algorithms		
CFX Manager 3.0	Bio-Rad	1845000
Excel 16.69.1	Microsoft	https://www.microsoft.com/en-us/microsoft-365/excel
Graphpad Prism 8	GraphPad Software	https://www.graphpad.com/scientific-software/prism/
Fiji v2.0.0-rc-69/1.52p	ImageJ	https://imagej.net/software/fiji/downloads
GelAnalyzer v19.1	GelAnalyzer	http://www.gelanalyzer.com
MaxQuant v1.6.10.43	[80]	https://www.maxquant.org/
Other		
NativePAGE 3-12% Bis-Tris Gels	Thermo Fisher Scientific	BN1001BOX

PVDF Transfer Membrane, 0.45 µm, 26.5 cm x 3.75 m	Thermo Fisher Scientific	88518
XCell SureLock™ Mini-Cell	Thermo Fisher Scientific	EI0001

METHOD DETAILS

Fly stocks and maintenance

Fly stocks were raised on standard cornmeal medium and kept at 23°C, 70% humidity on

a 12:12 hours light/dark cycle. Strains used in this study were obtained from Bloomington

Drosophila Stock Center (BDSC) and Vienna Drosophila Resource Center (VDRC).

Genotypes used in this study were: *act5c-gal4* (BDSC 4414), *da-gal4* (BDSC 8641), *UAS-*

Ndufs4 RNAi (VDRC 101489), *UAS-Bcs1 RNAi* (BDSC 51863), *UAS-Coa8 RNAi* (VDRC

100605). Control strains were obtained in each experiment by crossing the specific *gal4*

driver line with the genetic background flies *w1118*. *Coa8* KO flies were generated by

Wellgenetics Inc. by using CRISPR/Cas9 technology, generating a 676bp deletion, from

the -49th nucleotide relative to ATG to the -34th nucleotide relative to the stop codon of

Coa8.

RNA isolation, reverse transcription, and qRT-PCR

Total RNA was extracted from 10 individuals for each genotype using TRIzol method

(Thermo Fisher Scientific) according to the manufacturer's protocol. Reverse transcription

was performed with the GoScript Reverse Transcriptase kit (Promega). qRT-PCRs were

performed using GoTaq qPCR SYBR Green chemistry (Promega) and a Bio-Rad CFX 96

Touch System (Bio-Rad). The $2^{-\Delta\Delta Ct}$ method was used to calculate the expression levels

of the targets (*Bcs1, Ndufs4, Coa8*) using *Rp49* as endogenous control. The

oligonucleotides used are listed in the Key Resource Table

Isolation of Mitochondria

Mitochondria from *D. melanogaster* larvae and adults were prepared by homogenization

and differential centrifugation as described in [81]. Protein concentration of mitochondrial

extracts was measured with the Bio-Rad protein assay, based on the Bradford method.

Blue-native polyacrylamide gel electrophoresis (BN-PAGE) and in-gel activity (IGA)

assays

Isolated mitochondria were solubilized in 1.5M aminocaproic acid, 50 mM Bis-Tris/HCl pH
7.0. The samples were solubilized with 4 mg of digitonin (Calbiochem) or 4 mg n-dodecyl
β-D-maltoside (Sigma) per mg of protein. After 5 min of incubation on ice, samples were
centrifuged at 18,000 X g at 4 °C for 30 min. The supernatant was collected and
resuspended with Sample Buffer (750 mM aminocaproic acid, 50 mM Bis-Tris/HCl pH 7.0,
0.5 mM EDTA and 5% Serva Blue G). Native samples were separated using NativePAGE
3-12% Bis-Tris gels (Thermo Fisher Scientific) according to the manufacturer's protocol.
For Coomassie staining, gels were stained with Coomassie R 250 for 20 minutes and
destained/fixed using 20% methanol, 7% acetic acid. For in-gel activity assays, gels were
stained with the following solutions: complex II (succinate dehydrogenase): 5 mM Tris–HCl
pH 7.4, 0.2 mM phenazine methosulfate (Sigma), 20 mM succinate, and 1 mg/ml
nitrotetrazolium blue chloride; Complex IV (cytochrome c oxidase): 50 mM potassium
phosphate pH 7.4, 1 mg/ml 3′,3′-diaminobenzidine tetrahydrochloride hydrate (Sigma), 24
units/ml catalase from bovine liver (Sigma), 1 mg/ml cytochrome c from equine heart
(Sigma), and 75 mg/ml sucrose [82].

Complexome profiling

Mitochondria from *D. melanogaster* were analyzed by complexome profiling [83]. Isolated
mitochondria (0.2 mg) were solubilized with 6 g digitonin/g protein in 50 mM NaCl, 5 mM 6-
aminohexanoic acid, 1 mM EDTA, 50 mM imidazole/HCl, pH 7.0. After centrifugation at
22,000 X g for 20 min at 4°C, the supernatant was supplemented with Coomassie brilliant
blue G250 and proteins were separated by 4-16% gradient BN-PAGE. Digitonin-
solubilized mitochondrial proteins from bovine heart were loaded as molecular mass
standards. Gel lanes were cut into 60 slices, transferred to a 96-well filter microtiter plate
(Millipore), and destained in 50% (v/v) methanol, 50 mM ammonium bicarbonate. After
destaining, in-gel digestion with trypsin was performed. Tryptic peptides were separated by
liquid chromatography and analyzed by tandem mass spectrometry (LC-MS/MS) in a Q-
Exactive 2.0 Orbitrap Mass Spectrometer (2.8 SP1) equipped with an Easy nLC1000
nano-flow ultra-high-pressure liquid chromatography system (Thermo Fisher Scientific) at
the front end. Thermo Scientific Xcalibur 3.1 Software Package was used for data
recording. MS RAW data files were analyzed using MaxQuant (version 1.5.0.25). The
extracted spectra were matched against the *Drosophila melanogaster* Uniprot Reference
Sequence database (release 2020_04). Database searches were done with 20 ppm match
tolerances. Trypsin was selected as the protease with two missed cleavages allowed.

Dynamic modifications included N-terminal acetylation and oxidation of methionine.
Cysteine carbamidomethylation was set as a fixed modification. Keratins, and trypsin were
removed from the list. The abundance of each protein was determined by label-free
quantification using the composite intensity based absolute quantification (iBAQ) values
determined by MaxQuant analysis and was corrected for loading and MS sensitivity
variations between samples based on the total iBAQ value for all detected complex V
subunits. Gel migration profiles were created for each protein and normalized to the
maximum abundance. Profiles of the identified mitochondrial proteins were hierarchically
clustered by distance measures based on Pearson correlation coefficient (uncentered) and
the average linkage method using the NOVA software package v0.5 [84]. The visualization
and analysis of the heatmaps representing the normalized abundance in each gel slice by
a three-color code gradient (black/yellow/red) were done using Microsoft Excel 2019 and
Graph Pad Prism 8.4.3. The mass calibration for the BN-PAGE gels was performed as
previously described [85]. Membrane proteins were calibrated using the well-known
molecular masses of respiratory chain complexes and supercomplexes from bovine heart
mitochondria. The soluble proteins were, however, calibrated using the following set of
Drosophila proteins: ATPB (51 kDa), malate dehydrogenase (72 kDa, dimer), citrate
synthase (100 kDa, dimer), ETFA/B (122, heterodimer), heat shock protein 60 (410 kDa,
heptamer), ALDH7A1 (675 kDa, dodecamer).

Western blot and immunodetection
BN-PAGE gels were transferred to PVDF membranes in Dunn carbonate buffer (10 mM
NaHCO$_3$, 3 mM Na$_2$CO$_3$) applying a constant current of 300 mA at 4°C for 1 hour using a
Mini Trans-Blot® Cell (Bio-Rad). For the immunodetection of specific protein targets,
blotted PVDF membranes were blocked in 5% skimmed milk in PBS-T (0.1% Tween-20) at
room temperature for 1 hour and then incubated overnight with primary antibodies diluted
in 3% BSA in PBS-T overnight at 4°C. PVDF membranes were washed three times with
PBS-T for 10 minutes, incubated with the secondary HRP- conjugated antibody for 1 hour
at room temperature and washed three times with PBS-T for 10 minutes.
Chemiluminescent signals were recorded using an Alliance Mini HD9 (UVITEC).
Antibodies used are listed in the Key Resource Table. The primary antibodies against *D.*
melanogaster UQCR-C2 and SdhA were a kind gift of Dr. Edward Owusu-Ansah
(Columbia University, NY).

High-resolution respirometry

To measure oxygen consumption individuals were homogenized on ice in respiration buffer (120 mM sucrose, 50 mM KCl, 20 mM Tris-HCl, 4 mM KH_2PO_4, 2 mM $MgCl_2$, 1 mM EGTA, 1% fatty acid-free BSA, pH 7.2). Homogenates were loaded in the chamber of an O2k-HRR (High Resolution Respirometer, Oroboros Instruments) Complex I-linked respiration was measured at saturating concentrations of malate (2 mM), glutamate (10 mM), proline (10 mM) and ADP (2.5 mM). Afterwards, complex II-linked respiration was assessed using 10 mM succinate to the reaction after inhibition of complex I with rotenone (1.25 μM).

Analysis of MRC enzymatic activities

The activities of mitochondrial respiratory chain complexes and citrate synthase (CS) were measured using kinetic spectrophotometric assays as described [81].

Statistical analysis

Statistical analysis was performed with GraphPad Prism Software, version 8.2.1. Statistical tests and significance are described in the figure captions.

Data availability

Complexome profiling data will be deposited in the ComplexomE profiling DAta Resource (CEDAR) repository [86]. Data will be available upon manuscript acceptance.

ACKNOWLEDGEMENTS

We are grateful to Prof. Rodolfo Costa (CNR institute of Neuroscience, Padova, Italy) for providing the Coa8 KO and Bcs1 and Ndufs4 RNAi lines, Dr. Edward Owusu-Ansah (Columbia University, NY) for sharing the antibodies against *D. melanogaster* UQCR-C2 and SdhA and to Prof. Paolo Bernardi (Dept. of Biomedical Sciences, University of Padova) for critically reading the manuscript.

This research was funded by Fondazione Telethon-Cariplo Alliance GJC21014 (to E.F.-V.), Telethon Foundation GGP19007 (to M.Z.) and GGP20013 (to C.V.), AFM-Telethon 23706 (to C.V.), Department of Biomedical Sciences (University of Padova) FERN_FAR22_01 (to E.F.-V.) and SID2022- VISC_BIRD2222_01 (to C.V.), and Associazione Luigi Comini Onlus (MitoFight2, to M.Z. and C.V.).

AUTHOR CONTRIBUTIONS

Conceptualization, M.B., M.Z. and E.F.-V.; Methodology, M.B., A.C.-O., S.A. and E.F.-V.; Formal analysis, M.B., A.C.-O., S.A. and E.F.-V.; Investigation, M.B. and A.C.-O.; Data Curation, M.B., A.C.-O., S.A. and E.F.-V.; Writing-Original Draft, M.B. and E.F.-V.; Writing-Review & Editing, M.B., A.C.-O., S.A., M.Z. and E.F.-V.; Visualization, M.B. and E.F.-V.; Supervision, S.A., M.Z., C.V. and E.F-V.; Project Administration, M.B., M.Z., C.V. and E.F.-V.; Funding Acquisition, M.Z., C.V. and E.F.-V.

CONFLICT OF INTEREST

The authors declare no competing interests.

REFERENCES

1. Schagger, H., and Pfeiffer, K. (2000). Supercomplexes in the respiratory chains of yeast and mammalian mitochondria. The EMBO journal *19*, 1777-1783. 10.1093/emboj/19.8.1777.

2. Schagger, H., and Pfeiffer, K. (2001). The ratio of oxidative phosphorylation complexes I-V in bovine heart mitochondria and the composition of respiratory chain supercomplexes. J Biol Chem *276*, 37861-37867.

3. Letts, J.A., Fiedorczuk, K., Degliesposti, G., Skehel, M., and Sazanov, L.A. (2019). Structures of Respiratory Supercomplex I+III2 Reveal Functional and Conformational Crosstalk. Molecular cell *75*, 1131-1146 e1136. 10.1016/j.molcel.2019.07.022.

4. Vercellino, I., and Sazanov, L.A. (2021). Structure and assembly of the mammalian mitochondrial supercomplex CIII2CIV. Nature *598*, 364-367. 10.1038/s41586-021-03927-z.

5. Letts, J.A., Fiedorczuk, K., and Sazanov, L.A. (2016). The architecture of respiratory supercomplexes. Nature *537*, 644-648. 10.1038/nature19774.

6. Gu, J., Wu, M., Guo, R., Yan, K., Lei, J., Gao, N., and Yang, M. (2016). The architecture of the mammalian respirasome. Nature *537*, 639-643. 10.1038/nature19359.

7. Sousa, J.S., Mills, D.J., Vonck, J., and Kuhlbrandt, W. (2016). Functional asymmetry and electron flow in the bovine respirasome. Elife *5*. 10.7554/eLife.21290.

8. Wu, M., Gu, J., Guo, R., Huang, Y., and Yang, M. (2016). Structure of Mammalian Respiratory Supercomplex I1III2IV1. Cell *167*, 1598-1609 e1510. 10.1016/j.cell.2016.11.012.

9. Guo, R., Zong, S., Wu, M., Gu, J., and Yang, M. (2017). Architecture of Human Mitochondrial Respiratory Megacomplex I2III2IV2. Cell *170*, 1247-1257 e1212. 10.1016/j.cell.2017.07.050.

10. Bianchi, C., Genova, M.L., Parenti Castelli, G., and Lenaz, G. (2004). The mitochondrial respiratory chain is partially organized in a supercomplex assembly: kinetic evidence using flux control analysis. J Biol Chem *279*, 36562-36569.

11. Lenaz, G., and Genova, M.L. (2007). Kinetics of integrated electron transfer in the mitochondrial respiratory chain: random collisions vs. solid state electron

channeling. Am J Physiol Cell Physiol *292*, C1221-1239. 10.1152/ajpcell.00263.2006.

12. Lenaz, G., and Genova, M.L. (2009). Mobility and function of coenzyme Q (ubiquinone) in the mitochondrial respiratory chain. Biochim Biophys Acta *1787*, 563-573. 10.1016/j.bbabio.2009.02.019.

13. Lapuente-Brun, E., Moreno-Loshuertos, R., Acin-Perez, R., Latorre-Pellicer, A., Colas, C., Balsa, E., Perales-Clemente, E., Quiros, P.M., Calvo, E., Rodriguez-Hernandez, M.A., et al. (2013). Supercomplex assembly determines electron flux in the mitochondrial electron transport chain. Science *340*, 1567-1570. 10.1126/science.1230381.

14. Calvo, E., Cogliati, S., Hernansanz-Agustin, P., Loureiro-Lopez, M., Guaras, A., Casuso, R.A., Garcia-Marques, F., Acin-Perez, R., Marti-Mateos, Y., Silla-Castro, J.C., et al. (2020). Functional role of respiratory supercomplexes in mice: SCAF1 relevance and segmentation of the Qpool. Sci Adv *6*, eaba7509. 10.1126/sciadv.aba7509.

15. Garcia-Poyatos, C., Cogliati, S., Calvo, E., Hernansanz-Agustin, P., Lagarrigue, S., Magni, R., Botos, M., Langa, X., Amati, F., Vazquez, J., et al. (2020). Scaf1 promotes respiratory supercomplexes and metabolic efficiency in zebrafish. EMBO Rep *21*, e50287. 10.15252/embr.202050287.

16. Maranzana, E., Barbero, G., Falasca, A.I., Lenaz, G., and Genova, M.L. (2013). Mitochondrial respiratory supercomplex association limits production of reactive oxygen species from complex I. Antioxid Redox Signal *19*, 1469-1480. 10.1089/ars.2012.4845.

17. Lopez-Fabuel, I., Le Douce, J., Logan, A., James, A.M., Bonvento, G., Murphy, M.P., Almeida, A., and Bolanos, J.P. (2016). Complex I assembly into supercomplexes determines differential mitochondrial ROS production in neurons and astrocytes. Proceedings of the National Academy of Sciences of the United States of America *113*, 13063-13068. 10.1073/pnas.1613701113.

18. Hirst, J. (2018). Open questions: respiratory chain supercomplexes-why are they there and what do they do? BMC Biol *16*, 111. 10.1186/s12915-018-0577-5.

19. Green, D.E., and Tzagoloff, A. (1966). The mitochondrial electron transfer chain. Arch Biochem Biophys *116*, 293-304. 10.1016/0003-9861(66)90036-1.

20.   Kröger, A., and Klingenberg, M. (1973). Further evidence for the pool function of
ubiquinone as derived from the inhibition of the electron transport by antimycin. Eur
J Biochem *39*, 313-323. 10.1111/j.1432-1033.1973.tb03129.x.

21.   Kröger, A., and Klingenberg, M. (1973). The kinetics of the redox reactions of
ubiquinone related to the electron-transport activity in the respiratory chain. Eur J
Biochem *34*, 358-368. 10.1111/j.1432-1033.1973.tb02767.x.

22.   Hackenbrock, C.R., Chazotte, B., and Gupte, S.S. (1986). The random collision
model and a critical assessment of diffusion and collision in mitochondrial electron
transport. Journal of bioenergetics and biomembranes *18*, 331-368.

23.   Chazotte, B., and Hackenbrock, C.R. (1988). The multicollisional, obstructed, long-
range diffusional nature of mitochondrial electron transport. J Biol Chem *263*,
14359-14367.

24.   Blaza, J.N., Serreli, R., Jones, A.J., Mohammed, K., and Hirst, J. (2014). Kinetic
evidence against partitioning of the ubiquinone pool and the catalytic relevance of
respiratory-chain supercomplexes. Proceedings of the National Academy of
Sciences of the United States of America *111*, 15735-15740.
10.1073/pnas.1413855111.

25.   Fedor, J.G., and Hirst, J. (2018). Mitochondrial Supercomplexes Do Not Enhance
Catalysis by Quinone Channeling. Cell Metab *28*, 525-531 e524.
10.1016/j.cmet.2018.05.024.

26.   Mourier, A., Matic, S., Ruzzenente, B., Larsson, N.G., and Milenkovic, D. (2014).
The respiratory chain supercomplex organization is independent of COX7a2l
isoforms. Cell Metab *20*, 1069-1075. 10.1016/j.cmet.2014.11.005.

27.   Lobo-Jarne, T., Nyvltova, E., Perez-Perez, R., Timon-Gomez, A., Molinie, T., Choi,
A., Mourier, A., Fontanesi, F., Ugalde, C., and Barrientos, A. (2018). Human
COX7A2L Regulates Complex III Biogenesis and Promotes Supercomplex
Organization Remodeling without Affecting Mitochondrial Bioenergetics. Cell reports
25, 1786-1799 e1784. 10.1016/j.celrep.2018.10.058.

28.   Bundgaard, A., James, A.M., Harbour, M.E., Murphy, M.P., and Fago, A. (2020).
Stable mitochondrial CICIII2 supercomplex interactions in reptiles versus
homeothermic vertebrates. The Journal of experimental biology *223*, jeb223776.
10.1242/jeb.223776.

29.   Fernandez-Vizarra, E., Lopez-Calcerrada, S., Sierra-Magro, A., Perez-Perez, R.,
Formosa, L.E., Hock, D.H., Illescas, M., Penas, A., Brischigliaro, M., Ding, S., et al.

(2022). Two independent respiratory chains adapt OXPHOS performance to
glycolytic switch. Cell Metab *34*, 1792-1808. 10.1016/j.cmet.2022.09.005.

30. Berndtsson, J., Aufschnaiter, A., Rathore, S., Marin-Buera, L., Dawitz, H., Diessl, J.,
Kohler, V., Barrientos, A., Buttner, S., Fontanesi, F., and Ott, M. (2020). Respiratory
supercomplexes enhance electron transport by decreasing cytochrome c diffusion
distance. EMBO Rep *21*, e51015. 10.15252/embr.202051015.

31. Stuchebrukhov, A., Schafer, J., Berg, J., and Brzezinski, P. (2020). Kinetic
advantage of forming respiratory supercomplexes. Biochim Biophys Acta Bioenerg
1861, 148193. 10.1016/j.bbabio.2020.148193.

32. Trouillard, M., Meunier, B., and Rappaport, F. (2011). Questioning the functional
relevance of mitochondrial supercomplexes by time-resolved analysis of the
respiratory chain. Proceedings of the National Academy of Sciences of the United
States of America *108*, E1027-1034. 10.1073/pnas.1109510108.

33. Nesci, S., and Lenaz, G. (2021). The mitochondrial energy conversion involves
cytochrome c diffusion into the respiratory supercomplexes. Biochim Biophys Acta
Bioenerg *1862*, 148394. 10.1016/j.bbabio.2021.148394.

34. Acin-Perez, R., Bayona-Bafaluy, M.P., Fernandez-Silva, P., Moreno-Loshuertos, R.,
Perez-Martos, A., Bruno, C., Moraes, C.T., and Enriquez, J.A. (2004). Respiratory
Complex III Is Required to Maintain Complex I in Mammalian Mitochondria.
Molecular cell *13*, 805-815. 10.1016/s1097-2765(04)00124-8.

35. Diaz, F., Fukui, H., Garcia, S., and Moraes, C.T. (2006). Cytochrome c oxidase is
required for the assembly/stability of respiratory complex I in mouse fibroblasts.
Molecular and Cellular Biology *26*, 4872-4881. 10.1128/Mcb.01767-05.

36. Moreno-Lastres, D., Fontanesi, F., Garcia-Consuegra, I., Martin, M.A., Arenas, J.,
Barrientos, A., and Ugalde, C. (2012). Mitochondrial complex I plays an essential
role in human respirasome assembly. Cell Metab *15*, 324-335. S1550-
4131(12)00022-8 [pii]
10.1016/j.cmet.2012.01.015.

37. Protasoni, M., Perez-Perez, R., Lobo-Jarne, T., Harbour, M.E., Ding, S., Penas, A.,
Diaz, F., Moraes, C.T., Fearnley, I.M., Zeviani, M., et al. (2020). Respiratory
supercomplexes act as a platform for complex III-mediated maturation of human
mitochondrial complexes I and IV. The EMBO journal *39*, e102817.
10.15252/embj.2019102817.

38.   Fernandez-Vizarra, E., and Ugalde, C. (2022). Cooperative assembly of the
mitochondrial respiratory chain. Trends Biochem Sci *47*, 999-1008.
10.1016/j.tibs.2022.07.005.

39.   Maldonado, M., Guo, F., and Letts, J.A. (2021). Atomic structures of respiratory
complex III2, complex IV, and supercomplex III2-IV from vascular plants. Elife *10*,
e62047. 10.7554/eLife.62047.

40.   Zhou, L., Maldonado, M., Padavannil, A., Guo, F., and Letts, J.A. (2022). Structures
of Tetrahymena's respiratory chain reveal the diversity of eukaryotic core
metabolism. Science *376*, 831-839. 10.1126/science.abn7747.

41.   Maldonado, M., Fan, Z., Abe, K.M., and Letts, J.A. (2023). Plant-specific features of
respiratory supercomplex I + III(2) from Vigna radiata. Nat Plants *9*, 157-168.
10.1038/s41477-022-01306-8.

42.   Klusch, N., Dreimann, M., Senkler, J., Rugen, N., Kuhlbrandt, W., and Braun, H.P.
(2023). Cryo-EM structure of the respiratory I + III(2) supercomplex from
Arabidopsis thaliana at 2 A resolution. Nat Plants *9*, 142-156. 10.1038/s41477-022-
01308-6.

43.   Acin-Perez, R., Fernandez-Silva, P., Peleato, M.L., Perez-Martos, A., and Enriquez,
J.A. (2008). Respiratory active mitochondrial supercomplexes. Molecular cell *32*,
529-539. 10.1016/j.molcel.2008.10.021.

44.   Letts, J.A., and Sazanov, L.A. (2017). Clarifying the supercomplex: the higher-order
organization of the mitochondrial electron transport chain. Nat Struct Mol Biol *24*,
800-808. 10.1038/nsmb.3460.

45.   Davies, K.M., Blum, T.B., and Kuhlbrandt, W. (2018). Conserved in situ
arrangement of complex I and III2 in mitochondrial respiratory chain
supercomplexes of mammals, yeast, and plants. Proceedings of the National
Academy of Sciences of the United States of America *115*, 3024-3029.
10.1073/pnas.1720702115.

46.   Garcia, C.J., Khajeh, J., Coulanges, E., Chen, E.I., and Owusu-Ansah, E. (2017).
Regulation of Mitochondrial Complex I Biogenesis in Drosophila Flight Muscles. Cell
reports *20*, 264-278. 10.1016/j.celrep.2017.06.015.

47.   Shimada, S., Oosaki, M., Takahashi, R., Uene, S., Yanagisawa, S., Tsukihara, T.,
and Shinzawa-Itoh, K. (2018). A unique respiratory adaptation in Drosophila
independent of supercomplex formation. Biochim Biophys Acta Bioenerg *1859*, 154-
163. 10.1016/j.bbabio.2017.11.007.

48. Cabrera-Orefice, A., Potter, A., Evers, F., Hevler, J.F., and Guerrero-Castillo, S.
(2022). Complexome Profiling-Exploring Mitochondrial Protein Complexes in Health
and Disease. Front Cell Dev Biol *9*, 796128. 10.3389/fcell.2021.796128.

49. Wittig, I., Braun, H.P., and Schagger, H. (2006). Blue native PAGE. Nat Protoc *1*,
418-428.

50. Perales-Clemente, E., Fernandez-Vizarra, E., Acin-Perez, R., Movilla, N., Bayona-
Bafaluy, M.P., Moreno-Loshuertos, R., Perez-Martos, A., Fernandez-Silva, P., and
Enriquez, J.A. (2010). Five entry points of the mitochondrially encoded subunits in
Mammalian complex I assembly. Mol Cell Biol *30*, 3038-3047. MCB.00025-10 [pii]
10.1128/MCB.00025-10.

51. Melchionda, L., Haack, T.B., Hardy, S., Abbink, T.E., Fernandez-Vizarra, E.,
Lamantea, E., Marchet, S., Morandi, L., Moggio, M., Carrozzo, R., et al. (2014).
Mutations in APOPT1, Encoding a Mitochondrial Protein, Cause Cavitating
Leukoencephalopathy with Cytochrome c Oxidase Deficiency. Am J Hum Genet *95*,
315-325. 10.1016/j.ajhg.2014.08.003.

52. Signes, A., Cerutti, R., Dickson, A.S., Beninca, C., Hinchy, E.C., Ghezzi, D.,
Carrozzo, R., Bertini, E., Murphy, M.P., Nathan, J.A., et al. (2019). APOPT1/COA8
assists COX assembly and is oppositely regulated by UPS and ROS. EMBO
molecular medicine *11*. 10.15252/emmm.201809582.

53. Brischigliaro, M., Corra, S., Tregnago, C., Fernandez-Vizarra, E., Zeviani, M.,
Costa, R., and De Pitta, C. (2019). Knockdown of APOPT1/COA8 Causes
Cytochrome c Oxidase Deficiency, Neuromuscular Impairment, and Reduced
Resistance to Oxidative Stress in Drosophila melanogaster. Front Physiol *10*, 1143.
10.3389/fphys.2019.01143.

54. Brischigliaro, M., Badocco, D., Costa, R., Viscomi, C., Zeviani, M., Pastore, P., and
Fernandez-Vizarra, E. (2022). Mitochondrial Cytochrome c Oxidase Defects Alter
Cellular Homeostasis of Transition Metals. Front Cell Dev Biol *10*, 892069.
10.3389/fcell.2022.892069.

55. Villani, G., Greco, M., Papa, S., and Attardi, G. (1998). Low reserve of cytochrome c
oxidase capacity in vivo in the respiratory chain of a variety of human cell types. J
Biol Chem *273*, 31829-31836. 10.1074/jbc.273.48.31829.

56. Villani, G., and Attardi, G. (2000). In vivo control of respiration by cytochrome c
oxidase in human cells. Free Radic Biol Med *29*, 202-210.

57.   Fernandez-Vizarra, E., Bugiani, M., Goffrini, P., Carrara, F., Farina, L., Procopio, E.,
Donati, A., Uziel, G., Ferrero, I., and Zeviani, M. (2007). Impaired complex III
assembly associated with BCS1L gene mutations in isolated mitochondrial
encephalopathy. Hum Mol Genet *16*, 1241-1252. 10.1093/hmg/ddm072.
58.   Fernandez-Vizarra, E., and Zeviani, M. (2018). Mitochondrial complex III Rieske Fe-
S protein processing and assembly. Cell cycle *17*, 681-687.
10.1080/15384101.2017.1417707.
59.   Brischigliaro, M., Frigo, E., Corra, S., De Pitta, C., Szabo, I., Zeviani, M., and Costa,
R. (2021). Modelling of BCS1L-related human mitochondrial disease in Drosophila
melanogaster. J Mol Med (Berl) *99*, 1471-1485. 10.1007/s00109-021-02110-1.
60.   Ortigoza-Escobar, J.D., Oyarzabal, A., Montero, R., Artuch, R., Jou, C., Jimenez,
C., Gort, L., Briones, P., Muchart, J., Lopez-Gallardo, E., et al. (2016). Ndufs4
related Leigh syndrome: A case report and review of the literature. Mitochondrion
28, 73-78. 10.1016/j.mito.2016.04.001.
61.   Kruse, S.E., Watt, W.C., Marcinek, D.J., Kapur, R.P., Schenkman, K.A., and
Palmiter, R.D. (2008). Mice with mitochondrial complex I deficiency develop a fatal
encephalomyopathy. Cell Metab *7*, 312-320. 10.1016/j.cmet.2008.02.004.
62.   Foriel, S., Beyrath, J., Eidhof, I., Rodenburg, R.J., Schenck, A., and Smeitink,
J.A.M. (2018). Feeding difficulties, a key feature of the Drosophila NDUFS4
mitochondrial disease model. Dis Model Mech *11*. 10.1242/dmm.032482.
63.   Vogel, R.O., van den Brand, M.A., Rodenburg, R.J., van den Heuvel, L.P.,
Tsuneoka, M., Smeitink, J.A., and Nijtmans, L.G. (2007). Investigation of the
complex I assembly chaperones B17.2L and NDUFAF1 in a cohort of CI deficient
patients. Mol Genet Metab *91*, 176-182. 10.1016/j.ymgme.2007.02.007.
64.   Assouline, Z., Jambou, M., Rio, M., Bole-Feysot, C., de Lonlay, P., Barnerias, C.,
Desguerre, I., Bonnemains, C., Guillermet, C., Steffann, J., et al. (2012). A constant
and similar assembly defect of mitochondrial respiratory chain complex I allows
rapid identification of NDUFS4 mutations in patients with Leigh syndrome. Biochim
Biophys Acta *1822*, 1062-1069. 10.1016/j.bbadis.2012.01.013.
65.   Calvaruso, M.A., Willems, P., van den Brand, M., Valsecchi, F., Kruse, S., Palmiter,
R., Smeitink, J., and Nijtmans, L. (2012). Mitochondrial complex III stabilizes
complex I in the absence of NDUFS4 to provide partial activity. Human molecular
genetics *21*, 115-120. 10.1093/hmg/ddr446.

66. Caruana, N.J., and Stroud, D.A. (2020). The road to the structure of the mitochondrial respiratory chain supercomplex. Biochemical Society transactions *48*, 621-629. 10.1042/BST20190930.

67. Rathore, S., Berndtsson, J., Marin-Buera, L., Conrad, J., Carroni, M., Brzezinski, P., and Ott, M. (2019). Cryo-EM structure of the yeast respiratory supercomplex. Nat Struct Mol Biol *26*, 50-57. 10.1038/s41594-018-0169-7.

68. Hartley, A.M., Lukoyanova, N., Zhang, Y., Cabrera-Orefice, A., Arnold, S., Meunier, B., Pinotsis, N., and Marechal, A. (2019). Structure of yeast cytochrome c oxidase in a supercomplex with cytochrome bc1. Nat Struct Mol Biol *26*, 78-83. 10.1038/s41594-018-0172-z.

69. Milenkovic, D., Blaza, J.N., Larsson, N.G., and Hirst, J. (2017). The Enigma of the Respiratory Chain Supercomplex. Cell Metab *25*, 765-776. 10.1016/j.cmet.2017.03.009.

70. Hernansanz-Agustin, P., and Enriquez, J.A. (2021). Functional segmentation of CoQ and cyt c pools by respiratory complex superassembly. Free Radic Biol Med *167*, 232-242. 10.1016/j.freeradbiomed.2021.03.010.

71. Cogliati, S., Cabrera-Alarcon, J.L., and Enriquez, J.A. (2021). Regulation and functional role of the electron transport chain supercomplexes. Biochemical Society transactions *49*, 2655-2668. 10.1042/BST20210460.

72. Vercellino, I., and Sazanov, L.A. (2022). The assembly, regulation and function of the mitochondrial respiratory chain. Nature reviews. Molecular cell biology *23*, 141-161. 10.1038/s41580-021-00415-0.

73. Rohricht, H., Przybyla-Toscano, J., Forner, J., Boussardon, C., Keech, O., Rouhier, N., and Meyer, E.H. (2023). Mitochondrial ferredoxin-like is essential for forming complex I-containing supercomplexes in Arabidopsis. Plant Physiol. 10.1093/plphys/kiad040.

74. Lobo-Jarne, T., and Ugalde, C. (2018). Respiratory chain supercomplexes: Structures, function and biogenesis. Semin Cell Dev Biol *76*, 179-190. 10.1016/j.semcdb.2017.07.021.

75. Agip, A.A., Chung, I., Sanchez-Martinez, A., Whitworth, A.J., and Hirst, J. (2023). Cryo-EM structures of mitochondrial respiratory complex I from Drosophila melanogaster. Elife *12*. 10.7554/eLife.84424.

76. Padavannil, A., Murari, A., Rhooms, S.-K., Owusu-Ansah, E., and Letts, J.A. (2022). Resting mitochondrial complex I from Drosophila melanogaster adopts a helix-locked state. bioRxiv, 2022.2011.2001.514701. 10.1101/2022.11.01.514701.

77. Padavannil, A., Ayala-Hernandez, M.G., Castellanos-Silva, E.A., and Letts, J.A. (2022). The Mysterious Multitude: Structural Perspective on the Accessory Subunits of Respiratory Complex I. Front Mol Biosci *8*, 798353. 10.3389/fmolb.2021.798353.

78. Tropeano, C.V., Aleo, S.J., Zanna, C., Roberti, M., Scandiffio, L., Loguercio Polosa, P., Fiori, J., Porru, E., Roda, A., Carelli, V., et al. (2020). Fine-tuning of the respiratory complexes stability and supercomplexes assembly in cells defective of complex III. Biochim Biophys Acta Bioenerg *1861*, 148133. 10.1016/j.bbabio.2019.148133.

79. Kovarova, N., Pecina, P., Nuskova, H., Vrbacky, M., Zeviani, M., Mracek, T., Viscomi, C., and Houstek, J. (2016). Tissue- and species-specific differences in cytochrome c oxidase assembly induced by SURF1 defects. Biochim Biophys Acta *1862*, 705-715. 10.1016/j.bbadis.2016.01.007.

80. Cox, J., and Mann, M. (2008). MaxQuant enables high peptide identification rates, individualized p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol *26*, 1367-1372. 10.1038/nbt.1511.

81. Brischigliaro, M., Frigo, E., Fernandez-Vizarra, E., Bernardi, P., and Viscomi, C. (2022). Measurement of mitochondrial respiratory chain enzymatic activities in Drosophila melanogaster samples. STAR Protoc *3*, 101322. 10.1016/j.xpro.2022.101322.

82. Fernandez-Vizarra, E., and Zeviani, M. (2021). Blue-Native Electrophoresis to Study the OXPHOS Complexes. In Mitochondrial Gene Expression: Methods and Protocols, M. Minczuk, and J. Rorbach, eds. (Springer US), pp. 287-311. 10.1007/978-1-0716-0834-0_20.

83. Brischigliaro, M., Cabrera-Orefice, A., Sturlese, M., Elurbe, D.M., Frigo, E., Fernandez-Vizarra, E., Moro, S., Huynen, M.A., Arnold, S., Viscomi, C., and Zeviani, M. (2022). CG7630 is the Drosophila melanogaster homolog of the cytochrome c oxidase subunit COX7B. EMBO Rep *23*, e54825. 10.15252/embr.202254825.

84. Giese, H., Ackermann, J., Heide, H., Bleier, L., Drose, S., Wittig, I., Brandt, U., and Koch, I. (2015). NOVA: a software to analyze complexome profiling data. Bioinformatics *31*, 440-441. 10.1093/bioinformatics/btu623.

789   85.   Guerrero-Castillo, S., Baertling, F., Kownatzki, D., Wessels, H.J., Arnold, S., Brandt,
790   U., and Nijtmans, L. (2017). The Assembly Pathway of Mitochondrial Respiratory
791   Chain Complex I. Cell Metab *25*, 128-139. 10.1016/j.cmet.2016.09.002.
792   86.   van Strien, J., Haupt, A., Schulte, U., Braun, H.P., Cabrera-Orefice, A., Choudhary,
793   J.S., Evers, F., Fernandez-Vizarra, E., Guerrero-Castillo, S., Kooij, T.W.A., et al.
794   (2021). CEDAR, an online resource for the reporting and exploration of
795   complexome profiling data. Biochim Biophys Acta Bioenerg *1862*, 148411.
796   10.1016/j.bbabio.2021.148411.
797
798

 FIGURE CAPTIONS

Figure 1. *D. melanogaster* **mitochondrial respiratory chain does not rely on SC formation under physiological conditions. (A)** Complexome profiling of wild-type *D. melanogaster* mitochondria. Heatmaps show relative abundance of MRC subunits belonging to complex I (CI), complex II (CII), complex III$_2$ (CIII) and complex IV (CIV). Color scale of normalized peptide intensities are 0 (black), 96[th] percentile (yellow) and 1 (red). **(B)** BN-PAGE separation of mitochondria from wild-type *D. melanogaster* solubilized with digitonin. Native gels were either stained with Coomassie R250 or analyzed by in-gel activity (IGA) for complex I (CI) and complex IV (CIV). **(C)** BN-PAGE separation of mitochondria from wild-type *D. melanogaster* solubilized with dodecylmaltoside (DDM). Native gels were either stained with Coomassie R250 or analyzed by in-gel activity assay (IGA) for complex I (CI) and complex IV (CIV).

Figure 2. Severely perturbed CIV assembly results in increased formation of SC I$_1$III$_2$. **(A)** Complexome profiling of Coa8 KO *D. melanogaster* mitochondria. Heatmaps show relative abundance of MRC subunits belonging to complex I (CI), complex II (CII), complex III$_2$ (CIII) and complex IV (CIV). Color scale of normalized peptide intensities are 0 (black), 96° percentile (yellow) and 1 (red). **(B)** Average MS profiles depicted as relative abundance of MRC enzymes in natively separated complexes from wild-type (top) and Coa8 KO (bottom) fly mitochondria. Profiles of complexes I, III$_2$ and V (CI, CIII and CV) are plotted as average peptide intensity of the specific subunits identified by MS for each complex vs. apparent molecular weight. The increase in the relative abundance of SC I$_1$III$_2$ in Coa8 KO mitochondria is indicated by a black arrow. **(C)** In gel-activity assays for MRC complex I (CI), complex II (CII) and complex IV (CIV) in DDM-solubilized mitochondria from wild-type (w^{1118}) and Coa8 KO ($Coa8^{KO}$) flies. **(D)** BN-PAGE, western blot immunodetection of MRC complexes from wild-type (w^{1118}) and Coa8 KO ($Coa8^{KO}$) fly mitochondria using antibodies against specific subunits: anti-UQCRC2 (complex III), anti-NDUFS3 (complex I), anti-COX4 (complex IV) and anti-SDHA (complex II).

Figure 3. Mildly perturbed CIV assembly results in increased formation of SC I$_1$III$_2$. **(A)** Relative quantification (RQ) of *Coa8* mRNA expression in control (*da-gal4>+*) and *Coa8* KD (*da-gal4>Coa8^{RNAi}*) flies measured by qPCR. Data are plotted as mean ± SD (n = 3 biological replicates, Student's t test *p ≤ 0.05). **(B)** In gel-activity assays for MRC complex I (CI), complex II (CII) and complex IV (CIV) in DDM-solubilized mitochondria

from control (*da-gal4>+*) and *Coa8* KD (*da-gal4>Coa8^{RNAi}*) flies. **(C)** BN-PAGE, western
blot immunodetection of MRC complexes from a pool of three control (*da-gal4>+*) and
three Coa8 KD (*da-gal4>Coa8^{RNAi}*) fly mitochondria samples, using antibodies against
specific subunits: anti-UQCRC2 (complex III), anti-NDUFS3 (complex I), anti-COX4
(complex IV) and anti-SDHA (complex II).

Figure 4. Enhanced formation of SC I$_1$III$_2$ does not result in increased respiration. (A-
B) High-resolution respirometry (HRR) analyses of whole-fly homogenates. Respiration is
represented by oxygen flux (JO$_2$) measured by oxygen consumption rates (OCR –
pmol/s*fly). OCR have been measured via substrate-driven respiration under saturating
concentrations of substrates inducing either complex I (CI) or complex II (CII) -linked
respiration. Rotenone was used to block CI-linked respiration before measuring CII-linked
respiration. HRR was performed on **(A)** *Coa8* KO and **(B)** *Coa8* KD fly homogenates
compared to relative controls. Data are plotted as mean ± SD (n = 4 biological replicates).
(C-D) Kinetic enzyme activity of individual MRC complexes in **(C)** *Coa8* KO and **(D)** *Coa8*
KD compared with the relative control individuals, normalized by citrate synthase (CS)
activity. Data are plotted as mean ± SD (n = 3 biological replicates, two-way ANOVA with
Sidak's multiple comparisons, **p ≤ 0.01, ****p ≤ 0.0001).

Figure 5. Mild perturbation of CIII$_2$ biogenesis enhances SC formation in *D.*
***melanogaster.* (A)** Relative quantification (RQ) of *Bcs1* mRNA expression in control
(*act5c-gal4>+*) and *Bcs1* KD (*act5c-gal4>Bcs1^{RNAi}*) larvae measured by qPCR. Data are
plotted as mean ± SD (n = 3 biological replicates, Student's t test ***p ≤ 0.001). **(B)** In gel-
activity assays for MRC complex I (CI), complex II (CII) and complex IV (CIV) in DDM-
solubilized mitochondria from control (*act5c-gal4>+*) and *Bcs1* KD (*act5c-gal4>Bcs1^{RNAi}*)
larvae. **(C)** BN-PAGE, western blot immunodetection of MRC complexes from a pool of
three control (*act5c-gal4>+*) and three *Bcs1* KD (*act5c-gal4>Bcs1^{RNAi}*) larvae mitochondria
samples, using antibodies against specific subunits: anti-UQCRC2 (complex III), anti-
NDUFS3 (complex I), anti-COX4 (complex IV) and anti-SDHA (complex II). **(D)** Kinetic
enzyme activity of individual MRC complexes in control (*act5c-gal4>+*) and *Bcs1* KD
(*act5c-gal4>Bcs1^{RNAi}*) larvae mitochondria normalized by citrate synthase (CS) activity.
Data are plotted as mean ± SD (n = 3 biological replicates, two-way ANOVA with Sidak's
multiple comparisons, **p ≤ 0.01, ****p ≤ 0.0001). **(E)** High-resolution respirometry (HRR)
analyses of whole-fly homogenates. Respiration is represented by oxygen flux (JO$_2$)

measured by oxygen consumption rates (OCR – pmol/s*fly). OCR have been measured

via substrate-driven respiration under saturating concentrations of substrates inducing

either complex I (CI) or complex II (CII) -linked respiration. Rotenone was used to block CI-

linked respiration before measuring CII-linked respiration. HRR was performed on control

(*act5c-gal4>+*) and *Bcs1* KD (*act5c-gal4>Bcs1^{RNAi}*) homogenates compared to relative

controls. Data are plotted as mean ± SD (n = 3 biological replicates, two-way ANOVA with

Sidak's multiple comparisons, ****p ≤ 0.0001). **(F)** Relative quantification (RQ) of *Bcs1*

mRNA expression in control (*da-gal4>+*) and *Bcs1* KD (*da-gal4>Bcs1^{RNAi}*) larvae

measured by qPCR. Data are plotted as mean ± SD (n = 3 biological replicates, Student's t

test ***p ≤ 0.001). **(G)** In gel-activity assays for MRC complex I (CI), complex II (CII) and

complex IV (CIV) in DDM-solubilized mitochondria from control (*da-gal4>+*) and *Bcs1* KD

(*da-gal4>Bcs1^{RNAi}*) larvae. **(H)** BN-PAGE, western blot immunodetection of MRC

complexes from a pool of three control (*da-gal4>+*) and three *Bcs1* KD (*da-gal4>Bcs1^{RNAi}*)

larvae mitochondria samples using antibodies against specific subunits: anti-UQCRC2

(complex III), anti-NDUFS3 (complex I), anti-COX4 (complex IV) and anti-SDHA (complex

II). **(I)** Kinetic enzyme activity of individual MRC complexes in control (*da-gal4>+*) and

Bcs1 KD (*da-gal4>Bcs1^{RNAi}*) larvae mitochondria normalized by citrate synthase (CS)

activity. Data are plotted as mean ± SD (n = 3 biological replicates, two-way ANOVA with

Sidak's multiple comparisons, **p ≤ 0.01). **(J)** High-resolution respirometric (HRR)

analyses of whole-fly homogenates. Respiration is represented by oxygen flux (JO$_2$)

measured by oxygen consumption rates (OCR - pmol/s*fly). OCR have been measured via

substrate-driven respiration under saturating concentrations of substrates inducing either

complex I (CI) or complex II (CII) -linked respiration. Rotenone was used to block CI-linked

respiration before measuring CII-linked respiration. HRR was performed on control (*act5c-*

gal4>+) and *Bcs1* KD (*act5c-gal4>Bcs1^{RNAi}*) homogenates compared to relative controls.

Data are plotted as mean ± SD (n = 3 biological replicates, two-way ANOVA with Sidak's

multiple comparisons, ****p ≤ 0.0001).

Figure 6. Mild perturbation of CI biogenesis enhances SC formation in *D.*

***melanogaster.* (A)** Relative quantification (RQ) of *Ndufs4* mRNA expression in control

(*act5c-gal4>+*) and *Ndufs4* KD (*act5c-gal4>Ndufs4^{RNAi}*) larvae measured by qPCR. Data

are plotted as mean ± SD (n = 3 biological replicates, Student's t test **p ≤ 0.01). **(B)** In gel-

activity assays for MRC complex I (CI), complex II (CII) and complex IV (CIV) in DDM-

solubilized mitochondria from control (*act5c-gal4>+*) and *Ndufs4* KD (*act5c-*

gal4>Ndufs4^{RNAi}) larvae. **(C)** BN-PAGE, western blot immunodetection of MRC complexes

from a pool of three control (*act5c-gal4>+*) and three *Ndufs4* KD (*act5c-gal4>Ndufs4^{RNAi}*)

larvae mitochondria samples, using antibodies against specific subunits: anti-UQCRC2

(complex III), anti-NDUFS3 (complex I), anti-COX4 (complex IV) and anti-SDHA (complex

II). **(D)** Kinetic enzyme activity of individual MRC complexes in control (*act5c-gal4>+*) and

Ndufs4 KD (*act5c-gal4>^{RNAi}*) larvae mitochondria normalized by citrate synthase (CS)

activity. Data are plotted as mean ± SD (n = 3 biological replicates, two-way ANOVA with

Sidak's multiple comparisons, **$p \leq 0.01$, ***$p \leq 0.001$). **(E)** High-resolution respirometry

(HRR) analyses of whole-fly homogenates. Respiration is represented by oxygen flux

(JO_2) measured by oxygen consumption rates (OCR – pmol/s*fly). OCR have been

measured via substrate-driven respiration under saturating concentrations of substrates

inducing either complex I (CI) or complex II (CII) -linked respiration. Rotenone was used to

block CI-linked respiration before measuring CII-linked respiration. HRR was performed on

control (*act5c-gal4>+*) and *Ndufs4* KD (*act5c-gal4>Ndufs4^{RNAi}*) homogenates compared to

relative controls. Data are plotted as mean ± SD (n = 3 biological replicates, two-way

ANOVA with Sidak's multiple comparisons, ****$p \leq 0.0001$). **(F)** Relative quantification (RQ)

of *Ndufs4* mRNA expression in control (*da-gal4>+*) and *Ndufs4* KD (*da-gal4>Ndufs4^{RNAi}*)

larvae measured by qPCR. Data are plotted as mean ± SD (n = 3 biological replicates,

Student's t test ***$p \leq 0.001$). **(G)** In gel-activity assays for MRC complex I (CI), complex II

(CII) and complex IV (CIV) in DDM-solubilized mitochondria from control (*da-gal4>+*) and

Ndufs4 KD (*da-gal4>Ndufs4^{RNAi}*) larvae. **(H)** BN-PAGE, western blot immunodetection of

MRC complexes from a pool of three control (*da-gal4>+*) and three *Ndufs4* KD (*da-*

gal4>Ndufs4^{RNAi}) larvae mitochondria samples, using antibodies against specific subunits:

anti-UQCRC2 (complex III), anti-NDUFS3 (complex I), anti-COX4 (complex IV) and anti-

SDHA (complex II). **(I)** Kinetic enzyme activity of individual MRC complexes in control (*da-*

gal4>+) and *Ndufs4* KD (*da-gal4>Ndufs4^{RNAi}*) larvae mitochondria normalized by citrate

synthase (CS) activity. Data are plotted as mean ± SD (n = 3 biological replicates, two-way

ANOVA with Sidak's multiple comparisons, *$p \leq 0.05$). **(J)** High-resolution respirometry

(HRR) analyses of whole-fly homogenates. Respiration is represented by oxygen flux

(JO_2) measured by oxygen consumption rates (OCR - pmol/s*fly). OCR have been

measured via substrate-driven respiration under saturating concentrations of substrates

inducing either complex I (CI) or complex II (CII) -linked respiration. Rotenone was used to

block CI-linked respiration before measuring CII-linked respiration. HRR was performed on

control (*act5c-gal4>+*) and *Ndufs4* KD (*act5c-gal4>^{RNAi}*) homogenates compared to

relative controls. Data are plotted as mean $\pm$ SD ($n = 4$ biological replicates, two-way

ANOVA with Sidak's multiple comparisons).