Reversing protonation of weakly basic drugs greatly enhances intracellular diffusion and decreases lysosomal sequestration

Debabrata Dey,[1] Shir Marciano,[1] Anna Poryvai,[2] Ondřej Groborz,[2] Lucie Wohlrábová,[2] Tomás Slanina,[2] and Gideon Schreiber*,[1]

[1]Department of Biomolecular Sciences, Weizmann Institute of Science, Israel

[2]Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo náměstí 542/2, Prague 6, 160 00 Czech Republic

*Corresponding author: gideon.schreiber@weizmann.ac.il

KEYWORDS: Small molecule drugs, diffusion, in-cell, lysosome

ABSTRACT: For drugs to be active they have to reach their targets. Within cells this requires free diffusion, distribution, and availability. Here, we explored the intracellular diffusion rates and distribution of a series of small molecular fluorescent drugs, in comparison to proteins, by microscopy and fluorescence recovery after photobleaching (FRAP). While all proteins diffused freely, we found a strong correlation between pK_a and the intracellular diffusion and distribution of small molecule drugs. Weakly basic, small-molecule drugs displayed lower fractional recovery after photobleaching and 10-to-20-fold slower diffusion rates in cells than in aqueous solutions. As, more than half of pharmaceutical drugs are weakly basic, they, are protonated at pH 7.4, resulting in their sequestration in the cell cytoplasm. Protonation, facilitates the formation of membrane impermeable ionic form of the weak base small molecules. This results in ion trapping, further reducing diffusion rates of weakly basic small molecule drugs under macromolecular crowding conditions where other nonspecific interactions become more relevant and dominant. Our imaging studies showed that acidic organelles, particularly the lysosome, captured these molecules. Surprisingly, blocking lysosomal import only slightly increased diffusion rates and fractional recovery. Conversely, blocking protonation by *N*-acetylated analogues, greatly enhanced their diffusion and fractional recovery after FRAP. Based on these results, *N*-acetylation of small molecule drugs may improve the intracellular availability and distribution of weakly basic, small molecule drugs.

INTRODUCTION

Drug molecules alter many biochemical reaction pathways inside the cell by interacting with proteins, DNAs, RNAs, or others. By this, they influence one or more specific cellular activities (Ong et al., 2009; Schreiber, 2005; Stockwell, 2004). Due to their bioactivity, stability, rather simple chemical synthesis allowing for industrial bulk scale production and the easiness of administration, most drugs are small molecules (< 1 kDa) (Ngo and Garneau-

Tsodikova, 2018; Veber et al., 2002). Over half of these drugs have a significant degree of hydrophobicity and are weakly basic at intracellular pH of 7.4 (Charifson and Walters, 2014; Lipinski et al., 1997). Hydrophobicity is required to cross the lipid bilayer of the cell membranes, and polar functional groups solubilize the drug in the cellular medium (Lipinski et al., 1997). The balance between pK_a and logP (partition coefficient of small molecules as measured from their partition between octanol/water) is critical for successful drug design (Manallack, 2007; Meanwell, 2011). Lipinski's "Rules of 5" suggests that logP should be between 1 and 6 to be a candidate for successful oral administration. Small molecule drugs can be acidic or basic depending on their pK_a values (Lipinski et al., 1997). Drugs with pK_a ~ 0-7 are considered acidic, and pK_a ~ 8-14 are regarded as basic or alkaline. Basic drugs with pKa >8 are more common than their acidic counterparts (pKa < 7, ~60:40 ratios). In some cases, lowering their pK_a to the range between 6 and 7, has proven beneficial (Charifson and Walters, 2014). For a long time, mechanisms such as passive diffusion (Di et al., 2012) and carrier-mediated drug delivery (Dobson and Kell, 2008) or a combination of both (Sugano et al., 2010) have been used to explain intracellular drug delivery. However, transport of a drug across the cellular membrane is only one part of a successful delivery. Intracellular diffusion and bio-distribution needs to be taken in account when assessing the drug mode of action.

In previous work, we studied the diffusion of fluorescent drug molecules in crowded environments *in vitro* (Dey et al., 2022). However, the cellular cytoplasm of eukaryotic cells is by far more complex (Model et al., 2021). Intracellular membranes, multiple organelles (Ellis, 2001; Fulton, 1982) and a large diversity of macromolecules, in terms of size, charge and hydrophobicity, render the cellular interior extremely complex and heterogeneous (Milo and Phillips, 2015; Neurohr and Amon, 2020). Moreover, the pH inside cellular organelles can vastly differ from the cytosolic pH (Madshus, 1988). Organelles like lysosomes, late endosomes, Golgi, and secretory vesicles are acidic and have been known for sequestering weakly basic drugs (Asokan and Cho, 2002; Proksch, 2018). Therefore, predicting diffusion of a particular drug molecule in the dense cytoplasm is a complex task and requires sub-organelle resolution. For example, positively charged (basic) drugs designed to target nuclear DNA may bind cytoplasmic matter through nonspecific interactions (Long et al., 2022; Tiwari et al., 2018), they could be sequestrated within the lysosome, or diffuse slower than expected due to crowding and nonspecific interactions (Gotink et al., 2011; Jansen et al., 1999; Zhitomirsky and Assaraf, 2015).

The mobility of a drug in the cellular interior has been shown to be a good proxy for intracellular drug delivery and other processes, such as nonspecific binding or aggregation (Dey et al., 2022). The primary tool for measuring small molecule diffusion inside the cell is fluorescence recovery after photobleaching (FRAP), which inherently requires fluorescent drug molecules. FRAP with a confocal microscope is a sufficiently sensitive, versatile, and easy-to-use method suitable for measuring the diffusion of weakly fluorescent molecules. Unfortunately, single-molecule fluorescence correlation spectroscopy (FCS), a gold-standard method for measuring diffusion of fluorescent particles, cannot be applied for most small molecule drugs due to their weak fluorescence (Dey et al., 2022, 2021; Zotter et al., 2017).

In this work, we determined the intracellular diffusion coefficients of a series of fluorescent dyes and drugs compared to protein diffusion by FRAP (Figure 1 and Table 1). Specifically, we used Line-FRAP, which has a much higher time resolution and therefore can measure faster diffusion rates in comparison to standard FRAP (800

Hz versus 50 Hz, for detailed explanations, see (Dey et al., 2022) (Dey et al., 2021)). The apparent diffusion coefficients derived from FRAP measurements are termed $D_{confocal}$ (Equation 1, materials and method sections). Importantly, $D_{confocal}$ values for proteins determined by Line-FRAP agreed with the diffusion rates measured by FCS (Dey et al., 2021). For small molecules, Line-FRAP only provides relative 3D diffusion coefficients (Dey et al., 2022). We studied the following fluorescent drugs: the antimalarial drugs quinacrine dihydrochloride (Ehsanian et al., 2011), primaquine phosphate (Camarda et al., 2019) and amidoquine (Olliaro and Mussano, 2009). The anti-cancer drug mitoxantrone (Shenkenberg and Von Hoff, 1986), and the glycogen synthase kinase GSK3 inhibitor SB216763 (Wagman et al., 2005). In addition, we monitored the diffusion of the diagnostic staining agent fluorescein disodium salt (Jampol and Cunha-Vaz, 1984), the fluorogenic substrate CCF2 (Zuverink and Barbieri, 2015) and boron dipyrromethene (BODIPY) analogues, which are used as fluorescent markers for lipids, membranes, and other lipophilic compounds (Poryvai et al., 2022).

Table 1: LogP, pKa and lysosomal sequestration of drugs used in this study
(Predicted by ChemAxon software, and obtained from drug bank data base (Zhitomirsky and Assaraf, 2015)**)**

Therapeutic Indication	Drug/Small Molecules	Calculated LogP	Calculated pKa (Strongest basic)	Lysosomal sequestration
Anti-cancer	mitoxantrone	1.19	9.08	+
Anti-Malarial	quinacrine	5.15	10.33	+
	primaquine-biphosphate	1.64	10.4	+
	amodiaquine	4.83	8.66	+
Other Small molecules	5-Amino fluorescein	3.02	9.34	+
	fluorescein-Sodium Salt	2.65	2.85/3.7	-
	BODIPY-NH$_2$	_	9.00	_
	BODIPY-NHAc	_	_	_
	CCF2	_	5.1	_
	CF514	_	~3.0 (from UV spectrum)	_

Values of pKa were estimated using ChemAxon. Lysosomal sequestration was taken from the in-cell micrographs in this work.

The diffusion coefficients and their cellular distributions showed very slow diffusion of the basic compounds, i.e., low FRAP recovery, combined with an accumulation of the molecules in lysosomes. Surprisingly, while inhibiting the V-ATPase H$^+$ pump using Bafilomycin A1 (Spugnini et al., 2010; Wang et al., 2021) or sodium

azide (Hiruma et al., 2007) inhibited accumulation in lysosomes, cellular diffusion and fractional recovery increased only slightly. Contrary, blocking protonatable amino groups by acetylation greatly enhanced diffusion and FRAP recovery of studied small molecules. As over half the small molecule drugs are basic, our findings are highly important in future drug development, as slow diffusion and low FRAP recovery is directly related to reduced cellular activity. Contrary to small molecules, we show here that diffusion rates within the cytoplasm of HeLa cells of 16 different *E.coli* proteins with p*K*a values of 4.5-8, are within the expected range, suggesting that hindered diffusion is mainly a problem of small molecules.

RESULTS

Acidic Proteins diffuse freely in the HeLa cytoplasm. We have previously shown that diffusion in the cellular cytoplasm of E-fts and baeR, two bacterial globular proteins, aligns with the expected values considering the ~2-3-fold higher macro-viscosity (Dey et al., 2021). Here, we extended this study to include 16 additional proteins (mostly from *E. coli*) whose oligomeric states in solution were previously characterized (Table S1 and (Marciano et al., 2022)). The idea was to see and compare how a considerable population of proteins, heterologous to the cell (HeLa) diffuse inside the living cell. The proteins showed $D_{confocal}$ values of 18-30 $\mu m^2 s^{-1}$ in HeLa cell cytoplasm, which is typical for a cytoplasmic protein diffusion coefficient (Figure 2). Moreover, a complete FRAP recovery was observed, indicating free diffusion. Table S1 provides the predicted isoelectric points and net charges at pH 7.4 of the purified proteins as calculated by Prot Pi, showing p*K*a values of 4.5-8, as are the majority of *E.coli* cytosolic proteins. The results show that acidic and neutral proteins, even if not in their native environment (*E.coli* proteins in HeLa cells), do not stick to the HeLa cell cytoplasm. Next, we studied diffusion of small molecule drugs in HeLa cells.

Negatively charged small molecules rapidly diffuse in live cells. Fluorescein is a negatively charged small organic molecule (pK_a's = 2.85, 3.7, M.W. = 332 Da, Figure 1 and Table 1). We showed in previous work that fluorescein diffusion is slower in the presence of crowder proteins like BSA, lysozyme and, myoglobin, even at relatively low protein concentrations (Dey et al., 2022). Here, the Fluorescein dye was micro-injected in live HeLa cells, followed by FRAP measurements. For comparison, we also incubated HeLa cells for 24 hrs with fluorescein, which is a sufficient time for fluorescein to accumulate in the cells by diffusion. We found that the diffusion coefficient of fluorescein in PBS ($D_{confocal}$ of 56.5±2.4 $\mu m^2 s^{-1}$) was reduced to $D_{confocal}$ of 38.5±2.0 $\mu m^2 s^{-1}$ in HeLa cytoplasm (Figure 3B). In the cell nucleus, a value of $D_{confocal}$ of 43.2±2.2 $\mu m^2 s^{-1}$ was determined, while in concentrated HeLa cell extract (equivalent protein concentration of ~100 mg/mL), $D_{confocal}$ of 49.6±1.9 $\mu m^2 s^{-1}$ was measured. The faster $D_{confocal}$ in cell extract compared to live-cell cytoplasm may result from the higher crowding density of the latter. The lowering of diffusion coefficients of negatively charged fluorescein correlates with the increasing crowding density of the medium moving from simple buffer to cells. While micrographs did not show dye aggregation inside the cell cytoplasm (Figure S1A-B) and diffusion rates are the same as determined by microinjection (Figure S1C and S1D), the fractional recovery dropped from 0.95 to 0.65. This suggests that a fraction of fluorescein is immobile in the cell after 24 h incubation.

CCF2 is a fluorogenic substrate used for monitoring β-lactamase activity, with a pKa of ~5.1 (Zlokarnik et al., 1998). It is composed of cephalosporin core linking B7-hydroxycoumarin to fluorescein (Figure 1). Its calculated $D_{confocal}$ values in PBS and inside cells are about half as fast as those measured for fluorescein (Figure 3D-F), and its fractional recovery is ~0.7. No sign of aggregation is seen in micrographs inside the cell cytoplasm or nucleus (Figure 3F). The substantial slower rate of recovery, and lower fractional recovery of CCF2 relative to Fluorescein suggests its diffusion to be obstructed within the cell. This is in line with our prediction, that the observed substrate limited enzymatic degradation of CCF2 in HeLa cells is a consequence of its occlusion within the cellular cytoplasm (Zotter et al., 2017).

A third molecule for which we determined the diffusion coefficient is the labelling dye CF514. It is used to label proteins and has high quantum yield. As its structure has not been published, we estimated the pK_a value from the pH dependent UV-spectrophotometry measurements. The sharp changes of the UV- spectrum at pH~3 (Figure S1E-F), as well as the redshift confirms a pK_a value of ~ 3. The FRAP curves look similar to fluorescein (Figure 3G-I), with ~90% recovery after FRAP. $D_{confocal}$ in PBS was ~ 61.1±3.0 $\mu m^2 s^{-1}$, and 26.3±1.9 and 29.2±2.0 $\mu m^2 s^{-1}$ in the cell cytoplasm and nucleus of HeLa cells, respectively (Figure 3H). The drop of $D_{confocal}$ is attributed to cellular crowding. The results so far show that acidic small molecules diffuse well inside cells. Next, we focus on basic small molecules with higher pKa values.

High FRAP recovery but slow diffusion measured for the glycogen synthase kinase (GSK3) inhibitor SB216763. The GSK3 inhibitor SB216763 (pKa=7.0, M.W.=371 Da, Figure 1) aggregates when dissolved in PBS buffer solution. However, in our previous study we have seen that aggregation of this particular small molecule drug can be reversed by adding BSA and the extent of de-aggregation depends on the amount of BSA added (Dey et al., 2022). Here, our aim is to compare diffusion behavior of the GSK3 inhibitor in solution phase to that found in live cell cytoplasm. Since the molecule has some inherent solubility issues in PBS buffer, we compared its diffusion also in BSA protein solution and in cell culture media DMEM (Dulbecco's modified eagle medium). Comparative FRAP profiles and $D_{confocal}$ rates of SB216763 in different mediums are shown in Figure 4A-C. The FRAP profiles and relative $D_{confocal}$ values show very slow diffusion of SB216763 in HeLa cell cytoplasm of ~0.61±0.03 $\mu m^2 s^{-1}$ with a fraction recovery of 0.8 after 10s (Figure 4A-C). This is a 40-fold reduction in $D_{confocal}$ compared to measurements in 100 mg/mL BSA protein solution, where SB216763 is not aggregating ($D_{confocal}$~24.3±0.5 $\mu m^2 s^{-1}$ and fractional recovery of ~0.9 - Figure 4A-C). In cell culture DMEM media (containing 5% serum protein) fractional recovery after FRAP is very low (Figure 4A), indicating that the drug is still aggregating (Dey et al., 2022). Interestingly, the $D_{confocal}$ value in HeLa cell extract was higher ~ 11.3±0.7 $\mu m^2 s^{-1}$ as is its fractional recovery (0.9), showing that HeLa cell extract, like BSA is disaggregating SB216763. Co-localization with Lysotracker-633 (Figure 4D-E) shows that SB216763 colocalizes and is sequestered in acidic chambers of the cell lysosomes. Moreover, SB216763 accumulation into the lysosome increased over time (Figure 4D and E, 45-minute and 2-hour treatment). Varying drug dosages resulted in $D_{confocal}$ values of 0.42-0.61 $\mu m^2 s^{-1}$, suggesting that the slow diffusion is not related to SB216763 concentration (Figure S2A-B).

Despite its high solubility in aqueous solution, Quinacrine diffusion in HeLa cells is slow. Quinacrine DHC (M.W.= 472.9 Da, Figure 1) is a compound highly soluble in PBS (Dey et al., 2022) with a fast $D_{confocal}$ rate of 41.9±3.0 $\mu m^2 s^{-1}$ (Figure 5A-C). This rate dropped to $D_{confocal} \sim 2.5 \pm 0.1$ $\mu m^2 s^{-1}$ in HeLa cell cytoplasm, with fraction recovery of ~0.8, in comparison to one in buffer. In HeLa cell extract a $D_{confocal}$ rate of 15.1±0.6 $\mu m^2 s^{-1}$ was determined, which lays between PBS and HeLa cytoplasm. Micrographs of live HeLa cells show that the quinacrine DHC molecules aggregate in the cytoplasm. Co-localization with Lysotracker-633 (Figure 5D-E) show that quinacrine colocalizes and is sequestered in acidic chambers of the cell lysosomes. Moreover, quinacrine aggregation into the lysosome increased over time (Figure 5D and E, 45-minute and 2 hours treatment). Therefore, we measured diffusion instantaneously following micro-injection, and compared it to 2 and 24 hours of incubation (Figure S3A-B). Quinacrine accumulation in the lysosome was observed also immediately after micro-injection, with aggregation increasing over time. $D_{confocal}$ of 4.2±0.2 $\mu m^2 s^{-1}$ was calculated from line-FRAP immediately after micro-injection, slowing to 2.2±0.1 $\mu m^2 s^{-1}$ following 2 hours incubations, with fractional recoveries of 0.63 and 0.57 respectively. After 24 hours of incubation the fractional recovery is less than 0.25, and thus the diffusion value is meaningless (Figure S3C and D). Next, we compared FRAP of the cells treated with 2 and 6 µM quinacrine solutions after 2- and 24-hours incubations (Figure S4A-F). The concentration of quinacrine had only a marginal effect on the results. Again, following 24 hours of incubation, most of the quinacrine is in the lysosomes.

Mitoxantrone, primaquine and amidoquine all aggregate within cells. Mitoxantrone (anti-cancer) and primaquine phosphate (antimalarial) fluorescence with low quantum yields and weak photo-stability. However, due to their slow diffusion, the temporal resolution of standard XY-FRAP was sufficient for FRAP measurements (Figure 6A-H). Figures 6A and 6E show very low fraction recovery for these two drugs, suggesting that these weakly basic drugs are also sequestered in the cellular cytoplasm, as indeed shown in Figures 6B and 6F. Colocalization with LysoTracker red-633 show that Primaquine (S5A-B) and Amidoquine (S5C-D) are sequestered in the lysosomes. A summary of the measured FRAP rates in buffer and in cells are shown in Figure S6. The results shown above for the 5 basic drugs led us to investigate whether that their trapping in lysosomes is the reason for their slow diffusion.

Inhibition of lysosomal sequestration is only slightly increasing diffusion in cells. Lysosome internal pH of ~4.5 is maintained by vacuolar proton ATPase (V-H+-ATPase) (de Duve et al., 1974; Kaufmann and Krise, 2007). The low pKa of weakly basic small molecule drugs drive their accumulation in the lysosome. There, they become protonated (ionic form is less-permeable), which hinders their back-diffusion to the cytosol. The phenomenon is known as cationic ion trapping (Asokan and Cho, 2002; Kaufmann and Krise, 2007) and results in drug accumulation in lysosomes up to millimolar concentrations (de Duve et al., 1974). We blocked lysosomal acidification using either Bafilomycin A1 or sodium azide prior to drug administration. Bafilomycin A1 is a specific V-ATPase inhibitor and thus inhibits the supply of protons to the lysosomal chamber (Wang et al., 2021). Sodium azide, blocks ATP generation, resulting in the disruption of cellular activities which require ATP (Hiruma et al.,

2007; Ishii et al., 2014). As a result, cationic ion trapping is inhibited. These two compounds were used to investigate whether lysosomal trapping is the cause for the 20-40-fold lower diffusion and lower fraction recovery in HeLa cells compared to buffer. Figures S7A and B, and Figure S8A and B show colocalization images for GSK3 inhibitor and quinacrine, respectively (with Lysotracker-633 used as marker) after addition of 100 nM Bafilomycin A1 (Yoshimori et al., 1991). The micrographs show that Bafilomycin A1 strongly reduces lysosomal volumes following quinacrine and GSK3 treatment (Figure S7A *vs* S7B and S8A *vs* S8B). However, $D_{confocal}$ values were not altered by Bafilomycin A1, as shown in Figures S7C-D and S8C-D for GSK3 inhibitor and Quinacrine, respectively. $D_{confocal}$ for the GSK3 inhibitor is ~ 0.60 ± 0.02 $\mu m^2 s^{-1}$, and for Quinacrine ~ 2.4 ± 0.1 $\mu m^2 s^{-1}$. This suggests that Lysosomal accumulation of these two drugs is not the reason for the slow diffusion measured in the HeLa cell cytoplasm.

To verify the Bafilomycin results, HeLa cells were pre-treated with Sodium azide at 1 and 50 mM concentrations (Figure 7A and D), followed by treatment with GSK3 inhibitor or Quinacrine (Poste and Papahadjopoulos, 1976). The micrographs in Figures 7A and 7D, show that sodium azide inhibited lysosomal accumulation of these drugs. Comparative FRAP profiles and diffusion coefficients (Figure 7B-C and 7E-F) were not much changed for the two drugs and stayed low ($D_{confocal}$ are 1.8 ± 0.1-1.9 ± 0.1 $\mu m^2 s^{-1}$ for quinacrine and 0.4-0.5 $\mu m^2 s^{-1}$ for GSK3 inhibitor), similarly to what was observed with Bafilomycin.

5-amino fluorescein diffuses slowly in HeLa cells and is accumulated within lysosomes. To experimentally test the relation between the pKa of small molecules and their in cell diffusion, we measured the diffusion of 5-amino fluorescein (AM-Fluorescein, M.W.= 347.3 Da, Figure 8A), which is structurally similar to fluorescein disodium salt, but with an additional -NH$_2$ group that increases its pKa from ~3 to ~9 (Figure 8A). Figures 8B and C show comparative FRAP traces for these molecules, and for CCF2. While fluorescein disodium salt has a fast FRAP rate, and is completely recovering, the cationic charged AM-fluorescein recovers much slower with a fractional recovery of 0.4. Moreover, while fluorescein disodium salt is not sequestered in lysosomes, 5-Amino fluorescein is. This sequestration is alleviated by adding sodium azide to the cells (Figure 8D), but has only a small effect on the FRAP rate and fractional recovery (Figure 8B). This experiment strongly suggests that the functional primary amine group is the main contributor towards the slow diffusion and low FRAP recovery of AM-fluorescein in the cell cytoplasm.

Acidification of primaquine and BODIPY increases their diffusion and FRAP recovery. We substituted the NH$_2$ of primaquine (antimalarial drug) and BODIPY compounds for the charge-neutral-NHAc functional group (Figure S9A and S10A). Due to their low quantum yield, we could not implement the faster Line-FRAP protocol, therefore, we used standard FRAP, which suits the slow diffusion of both compounds. As seen from the FRAP profiles (time-dependent monographs) in Figures S9B-D for primaquine derivatives and in Figures S10B-C for BODIPY derivatives, the recovery is much slower for free amine-containing moieties. Figure S9I-J shows that the fractional recovery for primaquine is only ~0.27, whereas, for primaquine-NHAc, the recovery is ~0.6. Comparing the measurements using micro-injecting versus incubation of the cells for FRAP measurements for primaquine-

NHAc gave similar results (Figure S9E-H). Unfortunately, the classical FRAP is not fast enough to calculate the diffusion coefficients, as the dead time was 150 ms. For the BODIPY analogues, the observations were similar, but to a smaller extend. For BODIPY-NH$_2$, the recovery percentage is ~57%, whereas, for BODIPY-NHAc, the recovery is ~67% (Figure S10H). The estimated recovery half-life also supports the faster diffusion of BODIPY-NHAc, as shown in Figure S10I.

Basic small molecules do not colocalize with lipid droplets, the ER or nucleic acids. To identify the location of colocalization of basic small molecules, we tested colocalization of GSK3 inhibitor in lipid droplets using the Nile red dye (Figure S11) and in the ER using a specific BFP/mCherry tagged fluorescent antibody markers (Figure S12). Next, we looked for colocalization of Mitoxantrone with nucleic acids present in cytoplasm using SYTO blue (Figure S13). No colocalization was found between the drugs and either lipid droplets, the ER or nucleic acids. Super resolution images of GSK3 inhibitor treated HeLa cells were taken to improve the spatial distribution (Figure S14), however, this did not contribute new information to the puzzle. Therefore, we could not pinpoint the location of sequestration of weakly basic small molecule drugs within the cellular cytoplasm, except the lysosome.

DISCUSSION

In this study, we selected well-known and widely used small molecule drugs to study their behavior in aqueous solutions and living cells. All drug molecules used here are fluorescent, allowing for their FRAP measurements. Previously, we found that small-molecule drugs diffuse differently in crowded media than in a simple buffer solution, even if aggregation is not an issue (Dey et al., 2022). Here, we show that many small molecules get trapped within the cellular cytoplasm, resulting in extremely slow diffusion and lower fractional recovery after FRAP. This is particularly observed for cationic-charged small molecules, with D$_{confocal}$ values of 20-40-fold lower than observed for fluorescein or CF514, which, in many cases is accompanied by low recovery after photobleaching. The later suggests that most molecules are occluded to components of the HeLa cell cytoplasm. Even dense cell extract from HeLa cells is not mimicking the *in vivo* results (Figure 4A and C). This is very different from protein diffusion, where none of the 16 proteins measured here showed slow diffusion or low fractional recovery, despite their *E.coli* origin, expressed in HeLa cells. Slow diffusion and lower fractional recovery were limited to basic small molecules, showing D$_{confocal}$ values of 0.2-2 $\mu m^2 s^{-1}$. While the fractional recovery for all proteins is ~0.9, for many basic small molecules it is 0.2-0.5.

As primary culprit for the slow diffusion of the basic small molecule drugs we suspected sequestration in the lysosomes. Indeed, high level of accumulation in the lysosomes were found for all the basic drugs. However, pre-treatment with sodium azide or Bafilomycin A1, which we showed to inhibit lysosomal accumulation, had only a small effect on diffusion or fractional recovery after FRAP, which stayed low. Therefore, we evaluated colocalization of GSK3 inhibitor with lipid droplets, the ER or nucleic acids present in cytoplasm (Figure S13), but did not detect any.

The conclusive evidence that slow diffusion and lower FRAP recovery results from the drugs being basic came by altering three molecules from being basic to acidic and vice versa. The best example here is fluorescein,

where transformation to 5-amino fluorescein drastically reduced its diffusion as well as its fractional recovery after FRAP. Two additional examples here were Primaquine that was modified to Primaquine Acetate and BODIPY-NH$_2$ that was modified to BODIPY-NHAc. In both cases, the diffusion became faster and the fractional recovery increased significantly. This finding is important, as the activity of a drug within the cell is dictated by its active concentration. When the drug is sequestered, its active concentration is reduced. Even, if it is not fully sequestered, but only diffuses much slower, it will result in lower association rate constant, and thus lower affinity towards the drug target. As a result, to keep the drug active, its dose has to be increased, which can have negative implications on side effects, due to off-target binding.

CONCLUSIONS

Our study shows that *in vitro* biophysical crowding studies for small molecule drugs are of limited value if one wants to understand the biophysical behavior of the same drug within the cell. While it is true that availability of a drug within a cell is only one factor dictating its biological activity, it is a crucial one. Our findings here raise an important limitation on the standard rules for drug design, as these do not consider the stickiness of basic small molecules within the cell. We were able to directly address this question by using fluorescent molecule drugs, and measuring their location and diffusion within cells. Most importantly, we also show that by blocking protonation of a number of basic compounds we were able to increase substantially their diffusion and recovery after photobleaching, and thus potentially their activity. These findings may be consequential in future drug development.

MATERIALS AND METHODS

All the reagents used are described in a table format (Table S2, Key resources). The supplementary method sections describe the synthetic procedures, characterizations, and purity of the synthesized compounds.

Mammalian Cell Culture. HeLa cells were grown in 35-mm glass-bottomed dishes (MatTek Corporation) in DMEM (1X) Gibco (Life Technologies Limited) supplemented with 1X pyruvate, penicillin/streptomycin (BioIndustries), and 10% fetal bovine serum (Life Technologies Limited). The cells were subcultured when 80% confluence was reached using trypsin-EDTA for cell detachment. 2x10^5 HeLa cells in 2.5 ml of DMEM were pipetted into glass-bottomed dishes and incubated overnight. The cells were cultured in a humid atmosphere at 37 °C and 5% CO$_2$. The cells were imaged 24–30 h after seeding. Before the microinjection, the medium was aspirated, and fresh medium was supplemented with 25 mM HEPES, pH 7.4. For drug treatment, HeLa cells were incubated with 10 µM of drug (diluted 1:1000 from stock) for indicated times at 37 °C, followed by three times PBS 1X wash prior to imaging. For dose-dependent studies, the drug concentrations varied typically between 2-12 µM. Sodium azide concentration was 1 and 50 µM. Bafilomycin A1 was used at 100 nM concentration. Similarly, during the colocalization study, HeLa cells were treated with LysoTracker dye (1X) with 1:1000 dilution for 30 mins to 1 h at 37 °C, followed by 3x washing with PBS. Quinacrine, GSK3 inhibitor, Mitoxantrone, 5-amino Fluorescein, Amidoquine, Primaquine and BODIPY analogues were incubated with cells, while Fluorescein, Quinacrine, Primaquine-Ac, and CF514 dye were also micro-injected inside the HeLa cells.

Microinjection into HeLa cells. Microinjections were performed using the Eppendorf FemtoJet microinjector attached to the Eppendorf InjectMan NI2 micromanipulator. The fluorescein sodium salt, CCF2, CF514, quinacrine, primaquine analogues, BODIPY analogues were dissolved in DMSO and highly concentrated stock aliquots were made. Diluted PBS solutions of small molecules were injected into cells using glass capillaries from Warner instruments and pulled by a vertical puller (Narishige). For every measurement, a single pressure pulse was applied to deliver the sample into the cell. Air was administrated at 15– 25 hPa for 0.1-0.3 s. For injections, single cells containing morphologically healthy and well-connected HeLa cells were selected. Before and after the microinjection, cell morphology and membrane integrity were confirmed by visually inspecting the injected cells.

HeLa cell extract preparation. Cytoplasmic HeLa cell extracts were prepared as described previously with slight modifications (Zotter et al., 2017). HeLa cells from 4×10 cm plates at 80% confluency were washed, treated with trypsin and collected. Pellets were mixed with 500μl of RIPA buffer, IMP40, 5 μL of protease inhibitor and stored in ice. After 20 min, the mixture was centrifuged at 13k×g. This process was repeated twice to obtain concentrated extract. A BSA calibration curve determined the total protein concentration; the final concentration was up to 100 mg/mL. After liquid nitrogen freezing, the final cell extract solution was stored at -80° C.

Protein purification and dye labeling. Bacterial proteins used in this study were purified as described by us previously (Marciano et al., 2022). The dye labelling procedure with CF514 for purified proteins and removal of excess dye was described in detail (Dey et al., 2021).

Confocal microscopy and FRAP Analysis. *Confocal microscopy.* Images were collected with an Olympus IX81 FluoView FV1000 Spectral/SIM Scanner confocal laser-scanning microscope, using 60X DIC oil-immersion objective, N.A. 1.35. For fluorescein sodium salt fluorescence measurements, excitation was done at 440 nm, using a diode laser at an output power of 1-4 % of maximal intensity for high to low concentrations. In contrast, emission was recorded from 520 to 550 nm using the spectral detection system. For CCF2-FA, excitation was done with the laser at 440 nm using 1-2 % of the maximal intensity, while emission was collected from 470-560 nm with SDM560 emission dichromator as cut-off filter. For quinacrine DHC and GSK3 inhibitor, excitation was done at 488 nm laser using 1-2% of the maximal intensity, while emission was collected from 502-560 nm with SDM560 emission dichromator as cut-off filter. During colocalization experiments of quinacrine DHC and GSK3 inhibitor with LysoTracker-633 dye, the same setting was used for detecting the small molecule drugs in Channel 1 (green). LysoTracker-633 dye was detected in Channel 2 (red) by exciting at 635 nm laser using 1-2 % of the maximal intensity, while emission was collected from 655-755 nm. For mitoxantrone and BODIPY analogues, excitation was done at 635 nm laser using 1-2 % of the maximal power, and emission was collected from 655-755 nm in Channel 2 (red). In the blue channel, excitation was done by 400 nm laser using 2-4% of maximal power, whereas the emission was collected from 420-480 nm with SDM480 emission dichromator as cut-off filter. The same setting is used in the blue channel for the SYTO blue maker (to visualize free nucleic acids) and blue fluorescent protein marker (BFP to visualize the ER). For the mcherry protein marker (to visualize the ER), the same setting for the red channel (Channel 2) is used. For every colocalization experiments, proper blank experiments were performed separately in all the respective channels with the same microscope settings to avoid any bleeding/leakage of emission signals from one Channel to another Channel, as shown in Figure S15. Image analyses were performed

using FluoView/Imaris software, and data analyses were performed using Kaleidagraph software version 4.1 (Synergy). Colocalization coefficients as a percentage of material colocalized are calculated using Imaris software version 7.

Line-FRAP and Classical XY-FRAP. Line-FRAP was carried out in liquid drops and HeLa cells. For photobleaching, "Tornado" of 4 pixels (2x2) diameter (Dey et al., 2021). It is the smallest area achievable using Tornado. The bleach circle was kept precisely in the middle of the scanning line. The lines we scanned were unidirectional with time intervals of 1.256 ms 1000 times (equivalent to 1.256 s) in the majority of the measurements. The number of scans prior to, during, and after the photobleaching was 10, 42, and 948, respectively. Only in some specific cellular measurements (for GSK3 inhibitor and Quinacrine DHC, where recovery is very slow) the unidirectional lines were scanned 5000 times. Photobleaching was achieved by using the laser at 405-nm or with 635-nm excitations for the 63-millisecond duration at full intensity (100%). The simultaneous scanner moved at a 100 μs/pixel speed to perform an efficient photo-bleach. We have used two simultaneous scanners during the Line-FRAP experiments: one scanner (at 405 nm with the full intensity of 100%) for photobleaching and another scanner (at 440/515/635 nm with weak intensity) for data acquisition.

For all the drugs except Mitoxantrone and BODIPY analogues (emission in red wavelengths), photo-bleach was performed by 405 nm laser. For fluorescence signal detection: fluorescein 440 nm laser (1-4%); CCF2 440 nm laser (1-2%); GSK3 inhibitor 440 nm laser (0-2%); quinacrine (1-2%) of maximal intensity were used. Emission collections were done from 520-550 nm for quinacrine DHC and GSK3 inhibitor.

Line FRAP was not possible for primaquine analogues, as it suffers from strong photo-bleach during recovery. Here classical XY FRAP was performed, with photo-bleach done by 405 nm laser with full intensity. The frame rate is maintained at 172 ms with a bleach pulse duration was 150 ms. Due to microscope setting limitations, Line FRAP is also not feasible for Mitoxantrone and BODIPY analogues. A classical XY-FRAP using two simultaneous scanners is done for these molecules. Here, bleach was performed by simultaneous 635 nm laser with the full intensity of 100%, and the main excitation was done by 635 nm laser (with the power of 4%). Emission collections were done from 655-755 nm. For BODIPY analogues FRAP recoveries, the bleach pulse time was 150 ms. The bleach pulse time was also 150 ms for primaquine analogues for comparative FRAP profile collections. The frame rate is maintained at 172 ms. Using the Olympus IX81 FluoView FV1000 Spectral/SIM Scanner confocal laser-scanning microscope, using Tornado (which requires SIM scanner to be loaded) greatly enhances bleaching efficiency. In addition, it shortens the time to obtain the first measurement after bleach (which is immediate in this mode). This property is highly beneficial for Line-FRAP measurements, where the time scale of data acquisition plays an important role. The Fluoview SIM scanner unit synchronizes laser light simulation with confocal and multiphoton imaging to avoid interruption to image observation during laser stimulation or manipulation. We have varied the intensity of the lasers to achieve an optimal signal/noise ratio. Fluorescence recovery plots were fitted to a double exponent growth curve. FRAP experiments were also performed inside the PBS buffer drops and crowding conditions. A coverslip was introduced at the top of the drop to stop evaporation during measurements. Our previously developed method on Line FRAP protocol has been employed to calculate diffusion coefficients from the FRAP rates and averaged bleach sizes. Diffusion rates cannot be calculated using XY-FRAP. However, FRAP

parameters were kept constant so that qualitatively we could compare the average half-life and percentage of recoveries.

Determination of $D_{confocal}$ from FRAP results. We previously developed (Dey et al., 2021) the FRAP in Line mode (Line FRAP) method to monitor the diffusion rates of proteins in various environments. Line-FRAP allows a much faster data acquisition rate compared to classical XY FRAP. It is more suitable for measuring diffusion rates for fast diffusing molecules. The apparent diffusion coefficients calculated from Line FRAP measurements are $D_{confocal}$, which are calculated according to equation 1:

$$D_{confocal} = \frac{r_e^2 + r_n^2}{8\tau_{1/2}} \qquad \text{eq. 1}$$

where $t_{1/2}$ is the half-time of recovery and r_e and r_n is the effective and nominal bleach radii values (for details see (Dey et al., 2021)).

To calculate $D_{confocal}$ at least 30 independent measurements on different cells (n) were binned, and the curve fit of the progression curve was used to obtain $t_{1/2}$, and r_e values and their associated errors as determined from the fit of the curve. r_n was obtained from bleaching a fixed sample. The standard errors (SE) of the individual parameters were combined to obtain $D_{confocal} \pm$SE. To verify SE values, we repeated the 30 measurements independently multiple times, which gave the same SE as obtained from individual curve fits of n = 30 cells.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge on the e.life website.

AUTHOR INFORMATION

Corresponding Author
* Gideon Schreiber, Department of Biomolecular Sciences, Weizmann Institute of Science, Israel

AUTHORS

Debabrata Dey, Department of Biomolecular Sciences, Weizmann Institute of Science, Israel.

Shir Marciano, Department of Biomolecular Sciences, Weizmann Institute of Science, Israel**.**

Anna Poryvai, Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo náměstí 542/2, Prague 6, 160 00 Czech Republic

Ondrej Groborz, Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo náměstí 542/2, Prague 6, 160 00 Czech Republic

Lucie Wohlrábová, Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo náměstí 542/2, Prague 6, 160 00 Czech Republic

Tomás Slanina, Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo náměstí 542/2, Prague 6, 160 00 Czech Republic

Author Contributions

D.D. performed all the experiments. S.M. purified all the bacterial proteins. A.P, O.G., L.W., and T.S. designed the synthetic targets and synthesized primaquine acetate and BODIPY derivatives. D.D. and G.S. conceptualized the idea, analyzed the data and wrote the original draft of the manuscript. D.D., G.S., and T.S. reviewed and edited the paper. G.S. supervised the project. The manuscript was written through contributions of all authors. All authors have given approval to the final version of the manuscript.

Funding Sources

The Israel Science Foundation grant number 1268/18 (GS). The Czech Science Foundation (project No. 22-20319S) for funding this project (T.S.)

ACKNOWLEDGMENT

We would like to thank Dr. Ori Avinoam from Department of Biomolecular Sciences, and Prof. Hagen Hofmann from Department of chemical & structural Biology, Weizmann Institute of Science for their valuable discussions and advices. We want to acknowledge Dr. Reinat Nevo for her help with Imaris software to analyze the microscopy images.

ABBREVIATIONS

FRAP, fluorescence recovery after photo-bleaching; CCF2, coumarin cephalosporin fluorescein; BODIPY, boron dipyrromethene analogues; NMR, Nuclear magnetic resonance.

REFERENCES

1. Schreiber SL (2005) **Small Molecules: The Missing Link in The Central Dogma** *Nat. Chem. Biol* **1**:64–66
2. Stockwell BR (2004) **Exploring Biology with Small Organic Molecules** *Nature* **432**:846–854
3. Ong SE, Schenone M, Margolin AA, Li X, Do K, Doud MK, Mani DR, Kuai L, Wang X, Wood JL, Tolliday NJ, Koehler AN, Marcaurelle LA, Golub TR, Gould RJ, Schreiber SL, Carr SA (2009) **Identifying the proteins to which small-molecule probes and drugs bind in cells** *Proc. Natl. Acad. Sci. U. S. A.* **106**:4617-4622
4. Ngo HX, Garneau-Tsodikova S (2018) **What are the drugs of the future?** *Med chem comm* **9**:757-758
5. Veber DF, Johnson SR, Cheng HY, Smith BR, Ward KW, Kopple KD (2002) **Molecular Properties That Influence the Oral Bioavailability of Drug Candidates** *J. Med. Chem* **45**:2615-2623
6. Charifson PS, Walters WP (2014) **Acidic and Basic Drugs in Medicinal Chemistry: A Perspective** *J. Med. Chem* **57**:9701-9717
7. Lipinski CA, Lombardo F, Dominy BW, Feeney PJ (1997) **Experimental and Computational Approaches to Estimate Solubility and Permeability in Drug Discovery and Development Settings** *Adv. Drug Deliv. Rev* **23**:3–25
8. Manallack DT (2007) The pK(a) **Distribution of Drugs: Application to Drug Discovery** *Perspect Medicin Chem* **1**:25–38
9. Meanwell NA (2011) **Improving drug candidates by design: a focus on physicochemical properties as a means of improving compound disposition and safety** *Chem Res Toxicol* **24**:1420-1456

10. Di L, Artursson P, Avdeef A, Ecker GF, Faller B, Fischer H, Houston JB, Kansy M, Kerns EH, Krämer SD, Lennernäs H, Sugano K (2012) **Evidence-Based Approach to Assess Passive Diffusion and Carrier-Mediated Drug Transport**. *Drug Discov. Today* **17**:905–912

11. Dobson PD, Kell DB (2008) **Carrier-Mediated Cellular Uptake of Pharmaceutical Drugs: An Exception or the Rule?** *Nat. Rev. Drug Discov* **7**:205–220

12. Sugano K, Kansy M. Artursson P. Avdeef A. Bendels S. Di L, Ecker GF, Faller B, Fischer H, Gerebtzoff G, Lennernaes H, Senner F (2010) **Coexistence of Passive and Carrier-Mediated Processes in Drug Transport** *Nat. Rev. Drug Discov* **9**:597–614.

13. Dey D, Nunes-Alves, A, Wade RC, Schreiber G (2022) **Diffusion of small molecule drugs is affected by surface interactions and crowder proteins** *iScience* **25**:105088

14. Model MA, Hollembeak JE, Kurokawa M (2021) **Macromolecular crowding: A hidden link between cell volume and everything else** *Cell. Physiol. Biochem* **55**:25-40.

15. Fulton AB (1982) **How crowded is the cytoplasm?** *Cell* **30**:345-347

16. Ellis RJ (2001) **Macromolecular crowding: Obvious but underappreciated** *Trends in Biochemical Sciences* **26**:597-604

17. Milo R, Phillips R (2015) **Cell Biology by the Numbers** *Cell Biology by the Numbers*

18. Neurohr GE, Amon A (2020) **Relevance and Regulation of Cell Density** *Trends Cell Biol* **30**:213-225

19. Madshus IH (1988) **Regulation of intracellular pH in eukaryotic cells** *The Biochem. J* **250***:1*–8

20. Asokan A. Cho MJ (2002) **Exploitation of intracellular pH gradients in the cellular delivery of macromolecules** *J. Pharm. Sci* **91**:903-913.

21. Proksch E (2018) **pH in nature, humans and skin** *J. Dermatol* **45**:1044-1052

22. Long X, Zhang X, Chen Q, Liu M, Xiang Y, Yang Y, Xiao Z, Huang J, Wang X, Liu C, Nan Y, Huang Q (2022) **Nucleus-Targeting Phototherapy Nanodrugs for High-Effective Anti-Cancer Treatment** *Front. pharmacol* **13**:1-8

23. Tiwari R, Jain P, Asati S, Haider T, Soni V, Pandey V (2018) **State-of-art based approaches for anticancer drug-targeting to nucleus** *J. Drug Deliv. Sci. Technol* **48***:383*-392

24. Jansen G, Barr H, Kathmann I, Bunni MA, Priest DG, Noordhuis P, Peters GJ, Assaraf, YG (1999) **Multiple mechanisms of resistance to polyglutamatable and lipophilic antifolates in mammalian cells: Role of increased folylpolyglutamylation, expanded folate pools, and intralysosomal drug sequestration** *Mol. Pharmacol* **55**:761-769

25. Gotink KJ, Broxterman HJ, Labots M, de Haas RR, Dekker H, Honeywell RJ, Rudek MA, Beerepoot LV, Musters RJ, Jansen G, Griffioen AW, Assaraf YG, Pili R, Peters GJ, Verheul HMW (2011) **Lysosomal sequestration of sunitinib: A novel mechanism of drug resistance** *Clin. Cancer Res* **17**:7337–7346

26. Zhitomirsky B, Assaraf YG (2015) **Lysosomal sequestration of hydrophobic weak base chemotherapeutics triggers lysosomal biogenesis and lysosome dependent cancer multidrug resistance** *Oncotarget* **6**:1143-1156

27. Zotter A, Bäuerle, F, Dey D, Kiss V, Schreiber G (2017) **Quantifying enzyme activity in living cells** *J. Biol. Chem* **292**:15838–15848

28. Dey D, Marciano S, Nunes-Alves A, Kiss V, Wade RC, Schreiber G (2021) **Line-FRAP, A Versatile Method to Measure Diffusion Rates In Vitro and In Vivo** *J. Mol. Biol* **433**:166898

29. Ehsanian R, van Waes C, Feller SM (2011**) Beyond DNA binding - A review of the potential mechanisms mediating quinacrine's therapeutic activities in parasitic infections, inflammation, and cancers** *Cell Commun. Signal* **9**:1-18

30. Camarda G, Jirawatcharadech P, Priestley RS, Saif A, March S, Wong MHL, Leung S, Miller AB, Baker DA, Alano P, Paine MJI, Bhatia SN, O'Neill PM, Ward, SA, Biagini GA (2019) **Antimalarial activity of primaquine operates via a two-step biochemical relay** *Nat. Commun* **10:**1-9

31. Olliaro P, Mussano P (2003) **Amodiaquine for treating malaria** *Cochrane Database of Systematic Reviews* **2**:1-69

32. Shenkenberg TD, von Hoff DD (1986) **Mitoxantrone: A new anticancer drug with significant clinical activity** *Ann. Intern. Med* **105**:67-81

33. Wagman A, Johnson K, Bussiere D (2005) **Discovery and Development of GSK3 Inhibitors for the Treatment of Type 2 Diabetes** *Curr. Pharm. Des* **10**:1105–1137

34. Jampol LM, Cunha-Vaz J (1984) **Diagnostic Agents in Ophthalmology: Sodium Fluorescein and Other Dyes** *Pharmacology of the Eye Handbook of Experimental Pharmacology* **69**:699–714

35. Zuverink M. Barbieri JT (2015) **From GFP to β-lactamase: advancing intact cell imaging for toxins and effectors** *Pathog. Dis.* **73**: ftv097

36. Poryvai A, Galkin M, Shvadchak V, Slanina T (2022) **Red-Shifted Water-Soluble BODIPY Photocages for Visualisation and Controllable Cellular Delivery of Signaling Lipids** *Angew. Chemie - Int. Ed* **61**: e202205855

37. Wang R, Wang J, Hassan A, Lee CH, Xie XS, Li X (2021) **Molecular basis of V-ATPase inhibition by bafilomycin A1** *Nat. Commun* **12**:1-8

38. Spugnini EP, Citro G, Fais S (2010) **Proton pump inhibitors as anti-vacuolar-ATPases drugs: A novel anticancer strategy** *J. Exp. Clin. Cancer Res* **29**:44

39. Hiruma H, Katakura, T, Takenami T, Igawa S, Kanoh M, Fujimura T, Kawakami T (2007) **Vesicle disruption, plasma membrane bleb formation, and acute cell death caused by illumination with blue light in acridine orange-loaded malignant melanoma cells** *J. Photochem. Photobiol. B Biol* **86**:1-8.

40. Marciano S, Dey D, Listov D, Fleishman SJ, Sonn-Segev A, Mertens H, Busch F, Kim Y, Harvey SR, Wysocki VH, Schreiber G (2022) **Protein quaternary structures in solution are a mixture of multiple forms** *Chem Sci* **13**:11680-11695

41. Zlokarnik G, Negulescu PA, Knapp TE, Mere L, Burres N, Feng L, Whitney M, Roemer K, Tsien RY (1998) **Quantitation of transcription and clonal selection of single living cells with beta-lactamase as reporter** *Science* **279**:84-88

42. de Duve C, de Barsy T, Poole B, Trouet A, Tulkens P, van Hoof F (1974) **Commentary. Lysosomotropic agents**. *Biochem. Pharmacol* **23**:2495-2531

43. Kaufmann AM, Krise JP (2007) **Lysosomal sequestration of amine-containing drugs: Analysis and therapeutic implications** *J. Pharm. Sci* **96**:729-746

44. Ishii H, Shirai T, Makino C, Nishikata T (2014) Mitochondrial inhibitor sodium azide inhibits the reorganization of mitochondria-rich cytoplasm and the establishment of the anteroposterior axis in ascidian embryo *Dev. Growth Differ* **56**:175-188

45. Yoshimori T, Yamamoto A, Moriyama Y, Futai M, Tashiro Y (1991) **Bafilomycin A1, a specific inhibitor of vacuolar-type H+-ATPase, inhibits acidification and protein degradation in lysosomes of cultured cells** *J. Biol. Chem* **266**:17707-17712

46. Poste G, Papahadjopoulos D. (1976) **Lipid vesicles as carriers for introducing materials into cultured cells: influence of vesicle lipid composition on mechanism(s) of vesicle incorporation into cells** *Proc. Natl. Acad. Sci. U. S. A.* **73**:1603-1607

Insert Table of Contents artwork here

Figure 1. Chemical structures and pKa values (prediction from ChemAxon software) of the small molecules and drugs used in this study.

Figure 2. Diffusion coefficients of proteins and small molecules in the HeLa cells. Comparison of diffusion coefficients (Dconfocal in blue circles) and percentage of recoveries (in red squares) of bacterial proteins and BSA as measured in HeLa cell cytoplasm are shown. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure 3. Fluorescein, CCF2 and CF514 diffusion in PBS and inside HeLa cells. Comparative averaged FRAP profiles (N= 30; R= 0.99 for each of the fits) in PBS and inside HeLa cells with exponential fits are shown for (A) Fluorescein, (D) CCF2 and (G) CF514. Comparative $D_{confocal}$ values for (B) Fluorescein (E) CCF2 and (H) CF514 are also shown. HeLa cells after Micro-injection are shown in (C) for Fluorescein (F) for CCF2 and (I) for CF514. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure 4. Diffusion of GSK3 inhibitor in PBS and inside HeLa cells. Comparative (A) averaged FRAP recovery profiles with exponential fits for GSK3 inhibitor (SB216763) (N= 30; R= 0.99 for each of the fits). Dots show prebleach fluorescence, Black line for DMEM, blue for PBS buffer + 50 mg/mL of BSA, green for HeLa cells and orange for HeLa extract. (B) Averaged bleach size profiles with gaussian fits (N= 30; R= 0.99 for each of the fits) and (C) Calculated $D_{confocal}$ values. (D-E) Colocalization of GSK3 inhibitor in lysosomal compartments. HeLa cells were treated with Lyso Tracker 633 for 30 mins. GSK3 inhibitor treatment for (D) 45 mins and (E) 2 hours respectively at 10 µM concentrations are also shown. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure 5. Diffusion of Quinacrine in PBS and in HeLa cells. Comparative (A) averaged FRAP recovery profiles with exponential fits for Quinacrine dihydrochloride (N= 30; R= 0.99 for each of the fits), (B) averaged bleach size profile with gaussian fits (N= 30; R= 0.99 for each of the fits) and (C) diffusion coefficients in PBS buffer, in 50mg/mL of BSA, HeLa cell extract and in live HeLa cells are shown. (D-E) Colocalization of Quinacrine DHC in lysosomal compartments are shown. HeLa cells were treated with Lyso Tracker 633 for 30 mins. Quinacrine DHC treatment was done for (D) 45 mins and (E) 2 hours respectively at 10 μM concentration. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure 6. Mitoxantrone and Primaquine diffusion inside HeLa cells. Averaged XY- FRAP recovery profiles with exponential fits (N= 20; R= 0.98 for each of the fits) for (A) Mitoxantrone and (E) Primaquine in HeLa cells. Fluorescent and Transmission channel images of HeLa cells after 45 mins incubations with (B-C) Mitoxantrone and (F-G) Primaquine are shown. Time lapse micrographs of a portion of HeLa cell going through a classical rectangular XY-FRAP protocol for (D) Mitoxatrone and, (H) Primaquine are also shown.

Figure 7. Effect of Sodium Azide on the diffusion of GSK3 inhibitor and Quinacrine inside HeLa cells. (A) GSK3 inhibitor and (D) Quinacrine dihydrochloride treated HeLa cells with or without sodium azide. Comparison of averaged FRAP profiles with exponential fits (N= 30; R= 0.99 for each of the fits) and calculated $D_{confocal}$ values for (B-C) GSK3 inhibitor and (E-F) Quinacrine dihydrochloride. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure 8. FRAP of Fluorescein analogues inside HeLa cells. (A) Structures of Fluorescein analogues with pKa values. (B) FRAP recoveries with exponential fits for Fluorescein, CCF2 and 5-amino Fluorescein, with/without Na-Azide treatments in HeLa cells (N= 30; R= 0.99 for each of the fits) . (C) Line FRAP profiles with time lapses. (D) Micrograph images of treated HeLa cells after small molecule incubations.

Reversing protonation of weakly basic drugs greatly enhances intracellular diffusion and decreases lysosomal sequestration

Debabrata Dey,[1] Shir Marciano,[1] Anna Poryvai,[2] Ondřej Groborz,[2] Lucie Wohlrábová,[2] Tomáš Slanina[2] and Gideon Schreiber*,[1]

[1]Department of Biomolecular Sciences, Weizmann Institute of Science, Israel

[2]Institute of Organic Chemistry and Biochemistry of the Czech Academy of Sciences, Flemingovo náměstí 542/2, Prague 6, 160 00 Czech Republic

*Corresponding author: gideon.schreiber@weizmann.ac.il

Table S1: Oligomeric state, Isoelectric point and Net charge at pH 7.4

Protein entry	Isoelectric point: pI	Net charge at pH **7.4**: Z	$D_{confocal}$	Oligomeric state	Uniprot (kDa) Monomer
NfuA	4.534	-15.767	24.8±1.1	M/D	20.93
NadK	6.387	-4.808	29.8±1.4	M/D/T	32.57
FabG	6.250	-1.66	30.33±1.4	D/T/H	25.56
E-fts	5.228	-9.527	22.5±1.0	M/D	30.29
SodA	6.511	-2.481	25.5±1.2	D	22.97
tpiA	5.688	-7.851	27.9±1.7		
GpmA	5.913	-4.262	21.7±1.0	D	28.43
Can	6.230	-5.908	18.7±1.4	T	25.1
Upp	5.335	-6.829	23.0±1.1	M/D	22.53
SpeB	5.160	-14.788	30.4±1.4	Tri/H	33.56
IspD	6.257	-5.445	24.1±1.2	D/T/H	25.61
BaeR	5.554	-6.032	21.9±1.1	M/D	27.66
AcuI	5.669	-5.986	28.0±1.3	M/D/T/H	34.73
PyrF	5.874	-3.9	26.0±1.20	M/D	26.35
ThiD	5.793	-6.842	28.7±1.3	M/D	28.64
NadE	5.603	-4.993	23.2±1.1	D	27.16
CRP	7.979	+1.388	30.1±1.3	D/T	23.64
DES3im-4rp4	5.041	-5.659	29.4±1.4		
DES7im-3rk6	4.599	-10.654	28.8±1.3		
6E5C_denovobeta30	6.336	-0.714	30.0±1.4		
BSA	4.5–4.8	-15	29.6±1.4	M/D	66.5

Isoelectric points and net-charges were predicted using Prot Pi. Oligomeric states were taken from (Marciano et al., 2022) native mass spectrometry measurements.

Table S2. Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
Dulbecco's phosphate-buffered saline (PBS 1X)	Biological Industries	Cat# 02-023-1A
HEPES buffer	Fisher BioReagents	LOT# 170358
DMEM(1X) Gibco	Life Technologies Limited	REF# 41965-039
Fetal Bovine Serum (Gibco)	Life Technologies Limited	REF# 12657-029
Trypsin-EDTA Solution A	Biological Industries	REF# 03-050-1B
Penicillin/ Streptomycin	Biological Industries	REF# 03-031-1B
Sodium Pyruvate Solution	Biological Industries	REF# 03-042-1B
35-mm glass-bottomed dishes	MatTek Corporation	P35G-0-14-C
Pierce™ Dye Removal Columns	Thermo-Fisher scientific	Cat# 22858
Gebaflex tubes 3.5KDa (GeBa)	TIVAN BIOTECH	Cat# MIDI3-100
Capillary Glass Tubing	Warner Instruments	Model No. G120TF-4 203-776-0664
96- well Black F-Bottom Plate	Greiner-bio-one	REF# 655076
Disposable cuvettes	Fisher Scientific	ZEN0040
BSA (Albumin Bovine, fraction V)	MP Biomedicals, LLC	CAS# 9048-46-8
HEWL (Lysozyme from chicken egg white)	Merck	CAS# 12650-88-3
Myoglobin (from equine heart)	Merck	CAS# 100684-32-0
Doxorubicin	AdooQ Bioscience	Cat# A14403
Fluorescein disodium salt	chemcruz	Cat# sc-206026
GSK3 inhibitor SB216763	Abcam	Cat# ab120202
Quinacrine dihydrochloride	Abcam	Cat# ab120749
Primaquine biphosphate	Merck	CAS# 63-45-6
CCF2-FA	Thermo-Fisher Scientific	Cat# K1039a
CF514 Labelling dye	Biotium	CF514 Dye Alternative green fluorescent dye
Amodiaquine	AdooQ Bioscience	Cat# A17660
Sodium Azide	Merck	CAS# 26628-22-8
Mitoxantrone	Abcam	Cat# ab141041
Bafilomycin A1	Abcam	Cat# ab120497
5-amino Fluorescein	Abcam	Cat#ab145305

Characterization of Prmaquine (Prim) and N-acetylated Primaquine (PrimAc).

Primaquine (Prim)   Primaquine acetate (PrimAc)

The absorption and emission spectra of Primaquine analogues are shown in Figure S16 A& B respectively. They were measured in the mixture of 10% DMSO in PBS or in Glycine/NaOH buffer.

Materials & Reagent: Glycine/NaOH (pH = 10) (preparation: 80 mL of distilled water, 0.6 g of glycine, 0.25 g of NaOH, pH adjusted to 10 using 35% HCl, diluted to 100 mL)

Synthesis of PrimAc: To the stirred solution of Primaquine bisphosphate (100 mg, 220 µmol, 1 eq) and TEA (92 µL, 660 µmol, 3 eq) in DMF acetic anhydride (21 µL, 220 µmol, 1 eq) was added at RT. Reaction mixture was stirred for 5 minutes and the reaction mixture was diluted with dichloromethane (50 mL) and extracted 2x with water. Water phase was extracted with 20 mL of DCM and combined organic layers were extracted with brine. The organic phase was evaporated to the volume of approximately 1 mL and the solution was liquid loaded to the column. Gradient chromatography was done using cyclohexane, ethyl acetate and methanol (grad.: 100% cyclohexane – 30% cyclohexane, 60% ethyl acetate, 10% methanol). The chromatography was repeated to obtain the target compound as yellowish oil in 94% yield and 98.5% purity.

¹H NMR (401 MHz, CDCl3) δ 8.53 (dd, J = 4.2, 1.6 Hz, 1H), 7.92 (dd, J = 8.3, 1.6 Hz, 1H), 7.31 (dd, J = 8.2, 4.2 Hz, 1H), 6.34 (d, J = 2.5 Hz, 1H), 6.28 (dd, J = 2.6, 0.6 Hz, 1H), 5.99 (d, J = 8.4 Hz, 1H), 5.50 (s, 1H), 3.69 – 3.58 (m, 1H), 3.36 – 3.18 (m, 2H), 1.92 (s, 3H), 1.72 – 1.63 (m, 5H), 1.30 (d, J = 6.4 Hz, 3H). **¹³C NMR** (101 MHz, CDCl3) δ 170.03, 159.45, 144.92, 144.37, 135.36, 134.84, 121.91, 96.84, 91.75, 55.24, 47.85, 39.61, 34.01, 26.30, 23.33, 20.65. (Figure S17 A&B). HRMS spectrum of Primaquine-Ac is shown in Figure S18.

Hydrolytic experiment: The compound PrimAc (1 mg) was dissolved in 20 µL DMSO and diluted with 1.98 ml 0.1M PBS. The solution was measured by HPLC-MS (0-day, 1 day, 7 days, 1 month). The sample was stored in the dark at 25 °C between measurements. (Figure S16 C&D)

Characterization of BODIPY-NH₂ (BDP-NH₂) and BODIPY-NHAc (BDP-NHAc).

The Structures and photo-physical properties of BODIPY analogues are shown in Figure S19A. The absorption and emission spectra of BODIPY-NH₂ and BODIPY-NHAc in PBS/DMSO are shown in Figures S19 B&C, respectively.

BDP-NH₂: To a solution of a starting **BDP-N₃** (60 mg, 0.1 mmol, 1 eq.) and PPh₃ (47 mg, 0.2 mmol, 2 eq.) in THF (8 mL) water (1 mL) was added. The reaction mixture was stirred for 24 h and transferred on 20 g of silica. First fractions were eluted using a MeOH/DCM, 6/1 mixture followed by elution of the product using MeOH/DCM/NH₃ aq, 90/7/3, mixture. Organic solvents were removed under vacuum and an aqueous layer was extracted using DCM. The final crude product was reprecipitated from DCM using pentane, yielding blue crystals (34 mg, 61 % yield).

^{1}H NMR (401 MHz, CD₂Cl₂) δ 7.64 – 7.49 (m, 11H), 7.26 (d, J = 16.3 Hz, 2H), 7.01 – 6.92 (m, 4H), 6.66 (s, 2H), 4.03 (t, J = 5.2 Hz, 4H), 3.07 (t, J = 5.2 Hz, 4H), 1.64 (s, 4H), 1.46 (s, 6H). ^{13}C NMR (101 MHz, CD₂Cl₂) δ 160.5, 152.9, 142.6, 136.2, 135.5, 133.6, 132.2, 130.1, 129.9, 129.5, 129.3, 129.0, 117.8, 117.3, 115.3, 114.9, 71.0, 41.9, 14.8. HR-MS (ESI+), m/z: [M+H]+, calcd for [C₃₇H₃₈O₂N₄BF₂]⁺ 619.30571, found 619.30504. ^{1}H NMR and ^{13}C NMR of **BODIPY-NH₂** are shown Figures S20 A,B, respectively. HRMS spectrum of **BODIPY-NH₂** is shown in Figure S21.

BDP-NHAc: To a solution of **BDP-NH₂** (5 mg, 0.008 mmol, 1 eq.) in THF (2.0 ml), Et₃N (15 mg, 0.14 mmol, 17 eq.) and CH₃COCl (11 mg, 0.15 mmol, 18 eq) were added. The reaction mixture was stirred at r.t. for 2 h. Water was added and a product was extracted using DCM. The solvent was removed under

reduced pressure and the crude product was 3 times reprecipitated from DCM using pentane, yielding blue crystals (4.5 mg, 79 %).

^{1}H NMR (401 MHz, CD$_2$Cl$_2$) δ 7.78 – 7.46 (m, 10H), 7.40 – 7.31 (m, 3H), 7.26 (d, J = 16.3 Hz, 2H), 7.01 – 6.89 (m, 4H), 6.65 (s, 2H), 4.09 (t, J = 5.1 Hz, 4H), 3.65 (q, J = 5.4 Hz, 4H), 1.97 (s, 3H), 1.46 (s, 6H). ^{13}C NMR (101 MHz, CDCl$_3$) δ 160.0 (C), 152.9 (C), 142.7 (C), 139.1 (C), 136.0 (CH), 135.5 (C), 130.2 (C), 129.5 (CH), 129.3 (CH), 128.9 (CH), 117.8 (CH), 117.5 (CH), 115.2 (CH), 67.5 (CH$_2$), 39.3 (CH$_2$), 19.1 (CH$_3$), 14.8 (CH$_3$). HR-MS (ESI+), m/z: [M+H]+, calcd for [C$_{41}$H$_{41}$O$_4$N$_4$BF$_2$Na]$^+$ 725.30841, found 725.30811.

^{1}H NMR and ^{13}C NMR of **BODIPY-NHAc** are shown Figures S22A,B respectively. HRMS spectrum of **BODIPY-NHAc** is shown in Figure S2

Figure S1: FRAP measurements for Fluorescein following different incubation times. Micrographs of HeLa cells (A) instantaneously after micro-injection *vs* (B) FRAP after 24 hours incubation are shown. Comparative (C) averaged FRAP profiles with fits (N= 30; R= 0.99 for each of the fits) and (D) their $D_{confocal}$ values. (E-F) Comparative UV spectrum of CF514 labelling dye in 20 mM sodium phosphate buffer at different pH values, from which a pKa value of 3 was calculated. Error bars represent SE calculated from fitting the FRAP progression curves averaged over at least 30 independent measurements.

Figure S2: Diffusion rates at different doses of GSK3 inhibitor inside HeLa. (A-B) Drug dose dependency of GSK3 inhibitor treated HeLa cells. Effect of drug dosage (2-10 μM) on (A) averaged FRAP profiles with fits (N= 30; R= 0.99 for each of the fits) and (B) $D_{confocal}$ values. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure S3: Quinacrine diffusion in HeLa after micro-injection or incubation. Comparison of FRAP immediately following micro-injection (instant FRAP) and FRAP following 2-24 hr incubation. HeLa cell micrographs following (A) micro-injection and (B) 2 hours of incubation. Comparative (C) averaged FRAP profiles with exponential curve fits (N= 30; R= 0.99 for each of the fits) and (D) calculated $D_{confocal}$ values. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure S4: Dose and time dependency of Quinacrine diffusion in HeLa cells. HeLa cell micrographs after 2 hours of incubation with (A) 2 µM and (B) 6 µM of Quinacrine. (C) and (D) are micrographs of HeLa cells after 24 hours of incubation with 2 µM and 6 µM Quinacrine respectively. (E) Effect of drug dose and time dependency on comparative FRAP profiles, with fits (N= 30; R= 0.99 for each of the fits) and (F) Calculated $D_{confocal}$ values from the fits in (E). Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure S5: Colocalization of Primaquine and Amidoqiuine with Lyso-tracker. Colocalization of Primaquine (A-B) and Amidoquine (C-D) with Lyso Tracker 633 in HeLa cells. Time dependent measurements show higher aggregation of both drugs in the lysosomes over time.

Figure S6: Diffusion coefficients of small molecules in PBS buffer and inside HeLa cells.
Comparison of $D_{confocal}$ values of small molecule drugs in (A) PBS buffer and in (B) HeLa cells. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure S7: Inhibition of lysosomal accumulation of GSK3 inhibitor by Bafilomycin A. (A-B) Colocalization of GSK3 inhibitor with Lyso Tracker 633 in HeLa cells. (A) Untreated and (B) treated with 100 nM Bafilomycin A1 and corresponding colocalizations are shown. (C) Comparative averaged FRAP profiles with fits (N= 30; R= 0.99 for each of the fits) and (D) $D_{confocal}$ diffusion values with and without the Bafilomycin A1 treatment. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure S8: Inhibition of lysosomal accumulation of Quinacrine by Bafilomycin A. (A-B) Colocalization of Quinacrine with Lyso Tracker 633 in HeLa cells. HeLa cells (A) untreated and (B) treated with 100 nM Bafilomycin A1 and corresponding colocalizations. (C) Comparative averaged FRAP profiles with fits (N= 30; R= 0.99 for each of the fits) and (D) $D_{confocal}$ values with and without the Bafilomycin A1 treatment. Error bars represent SE calculated from fitting the FRAP progression curves, which are averaged over at least 30 independent measurements.

Figure S9: Comparative FRAP of Primaquine analogues inside HeLa cells. (A) Structures of Primaquine analogues with pKa values. Comparative (B-D) XY FRAP micrographs with time lapses in HeLa cell. (E-H) Micrographs of treated HeLa cells either with micro-injection or after drug incubations with Prim or Prim-Ac. Comparative (I) XY-FRAP recovery profiles with fits (N= 20; R= 0.98 for each of the fits) and (J) percentage of recoveries for Primaquine analogues in HeLa cytoplasm (N= 20).

Figure S10: Comparative FRAP of BODIPY analogues inside HeLa cells. (A) Structures of BODIPY analogues. (B-C) Comparative XY FRAP micrographs with time lapses in HeLa cell cytoplasm. (D-G) Micrographs of treated HeLa cells after incubations (with BDP/BDP-Ac) and, Comparative (H) averaged XY-FRAP recovery profiles with fits (N= 30; R= 0.99 for each of the fits), (I) averaged half-life and percentage of recoveries for BODIPY analogues in HeLa cytoplasm (H) (N= 30).

Figure S11: Time dependent colocalization of GSK3 inhibitor and Nile Red inside HeLa cells. Panels (A-C) show Nile red accumulating in lipid droplets, while GSK3 inhibitor does not colocalize.

Figure S12: Time dependent colocalization of GSK3 inhibitor and ER marker inside HeLa cells. Panels (A-D) show BFP/mCherry fused antibody marker accumulating in ER, but GSK3 inhibitor does not colocalize. HeLa cells were first transiently transfected with BFP/mCherry fused antibody markers, then treated with GSK3 inhibitor.

Figure S13: Time dependent colocalization study of Mitoxantrone and SYTO blue marker inside HeLa cells. Panels (A-B) show SYTO blue marker accumulating in nucleic acids, but Mitoxantrone is not colocalized. HeLa cells were first treated with SYTO blue, then treated with Mitoxantrone.

Figure S14: Super resolution images of GSK3 inhibitor treated HeLa cells. Micrographs were taken using a Zeiss LSM900 microscope.

Figure S15: Evaluating channel leakage. Control experiments showing that channel leakage did not occur during colocalization studies. (A) Lyso-Tracker only, (B) Quinacrine only and (C) Mitoxantrone only.

Figure S16: Characterization of Primaquine (Prim) and N-acetylated Primaquine (PrimAc). (A) Absorption (c ≈ 1 · 10-3) and (B) emission (c ≈ 3 · 10-4) spectra. Both experiments were measured in the mixture of 10% DMSO in PBS or in Glycine/NaOH buffer. (C) Measurement of hydrolytic stability by HPLC. (D) HPLC-MS measurement of PrimAc are shown. According to HPLC-MS data the peak at 7.125 corresponds to 302 m/z (ESI +) so as the main peak at 7.296. The molar mass of PrimAc corresponds to the measured value.

A   ¹H NMR spectrum.

B   ¹³C NMR spectrum

Figure S17: N-acetylated Primaquine. (A) ¹H NMR spectrum is shown. The presence of impurities is caused by the degradation in the used solvent. (B) ¹³C NMR spectrum is shown.

HRMS spectrum

C17 H23 N3 O2: +EI Scan (rt: 18.9-19.0 min, 7 scans) Frag=70.0V Wohlrabova_TSLW-222.D Subtract

301.1790

302.1816

300.1716

303.1836

Counts vs. Mass-to-Charge (m/z)

Figure S18: N-acetylated Primaquine. HRMS spectrum (+EI) is shown

Figure S19: (A) Structure and photo-physical properties of BODIPY analogues. (B) Absorption (full line) and emission spectra (dashed line) of BDP-NH₂ in DMSO (black) and PBS (red) are shown. (C) Absorption (full line) and emission spectra (dashed line) of BDP-NHAc in DMSO (black) and PBS (red) are shown.

A ¹H NMR spectrum

B ¹³C NMR spectrum

Figure S20: Characterization of BODIPY-NH₂. (A) ¹H NMR spectrum and (B) ¹³C NMR spectrum are shown.

071122_servisHR_20_221107143510 #127 RT: 3.52 AV: 1 SB: 21 0.47-0.69 , 0.53-0.84 NL: 8.34E5
T: FTMS + p ESI Full ms [220.00-2000.00]

Figure S21: BODIPY-NH₂. HRMS spectrum (+EI) is shown.

A ¹H NMR spectrum

I57-AP-272-PREC-P1-DN1.1.fid

7.60, 7.60, 7.59, 7.59, 7.58, 7.57, 7.55, 7.55, 7.54, 7.53, 7.52, 7.52, 7.51, 7.50, 7.37, 7.36, 7.36, 7.35, 7.35, 7.28, 7.24, 6.97, 6.97, 6.96, 6.96, 6.95, 6.66, 6.65

5.32 CD2Cl2

4.10, 4.09, 4.08, 3.67, 3.65, 3.64, 3.63

1.97, 1.46

10.17, 5.05, 4.06, 2.00, 3.97, 3.97, 3.26, 6.73, 6.25

f1 (ppm)

B ¹³C NMR spectrum

I57-AP-272-PREC-P1-DN1.2.fid

160.0, 152.9, 142.7, 139.1, 136.0, 135.5, 130.2, 129.5, 129.3, 128.9, 117.8, 117.5, 115.2

67.5

53.8 CD2Cl2

39.3

19.1, 14.8

f1 (ppm)

Figure S22: Characterization of BODIPY-NHAc. (A) ¹H NMR spectrum and (B) ¹³C NMR spectrum are shown.

Figure S23: BODIPY-NHAc. HRMS spectrum (+EI) is shown