1

Title: Plant genome evolution in the genus *Eucalyptus* driven by structural rearrangements

that promote sequence divergence

Running title: Architectural evolution of plant genomes

Scott ferguson[1*], Ashley Jones[1*], Kevin Murray[1,2], Rose Andrew[3], Benjamin Schwessinger[1], and

Justin Borevitz[1]

1. Research School of Biology, Australian National University, Canberra, Australian

10   Capital Territory, Australia

11   2. Weigel Department, Max Planck Institute for Biology Tübingen, Tübingen,Germany

12   3. Botany & N.C.W. Beadle Herbarium, School of Environmental and Rural Science,

13   University of New England, Armidale, NSW 2351, Australia.

14   *. First authors

16   Corresponding author

17   Scott Ferguson

18   scott.ferguson@anu.edu.au

22 Abstract

Genomes have a highly organised architecture (non-random organisation of functional and

non-functional genetic elements within chromosomes) that is essential for many biological

functions, particularly, gene expression and reproduction. Despite the need to conserve

genome architecture, a surprisingly high level of structural variation has been observed

within species. As species separate and diverge, genome architecture also diverges,

becoming increasingly poorly conserved as divergence time increases. However, within

plant genomes, the processes of genome architecture divergence are not well described.

Here we use long-read sequencing and *de novo* assembly of 33 phylogenetically diverse,

wild and naturally evolving *Eucalyptus* species, covering 1-50 million years of diverging

genome evolution to measure genome architectural conservation and describe architectural

divergence. The investigation of these genomes revealed that following lineage divergence

genome architecture is highly fragmented by rearrangements. As genomes continue to

diverge, the accumulation of mutations and subsequent divergence beyond recognition of

rearrangements becomes the primary driver of genome divergence. The loss of syntenic

regions also contribute to genome divergence, but at a slower pace than rearrangements.

We hypothesise that duplications and translocations are potentially the greatest contributors

to *Eucalyptus* genome divergence.

41 Keywords

42   1. Plant genome evolution

43   2. Synteny

44   3. Genome rearrangements

45   4. Genome assembly

46   5. Eucalyptus

Introduction

Genomes from all kingdoms are highly organised, but vary greatly in their structural

architecture (Koonin, 2009). Within eukaryotic genomes, genome architecture refers to the

non-random organisation of functional and non-functional genetic elements within

chromosomes (genes, regulatory regions, small RNAs, transposons, pseudogenes, introns,

centromeres, telomeres, etc.), and is critical for many biological functions, in particular

reproduction and gene expression. However, the conservation and divergence of genome

architecture or structure among a group of radiating plant species that share a common

karyotype has not been well described.

For effective recombination during meiosis and the production of viable reproducing

offspring, the genome architecture of both parental haplotypes must be highly similar.

Changes to the genetic architecture can result in reproductive isolation/incompatibility or

non-viable gametes (Hardigan *et al.*, 2020; Simakov *et al.*, 2020). Therefore, a common

genome architecture within individuals of a breeding population tends to be highly

conserved, except at some loci with high diversity (Jiao and Schneeberger, 2020). Similarly,

for expression of a gene to be correctly regulated, it must be placed on a chromosome

alongside the required promoters, enhancers, and inhibitors. The 3D organisation of the

surrounding chromatin must permit physical access to allow transcription (Heng *et al.*, 2004;

Dixon, Gorkin and Ren, 2016; Oudelaar and Higgs, 2021).

Despite this functional need for structural conservation, some structural differences are

known to exist between genomes within species. The extent to which reproductively

compatible genomes are structurally different is an open area of research; however, several

studies have shown genomes with a surprising amount of structural differences to be

reproductively compatible (Lin and Gokcumen, 2019; Alonge *et al.*, 2020; Jiao and

Schneeberger, 2020; Tang *et al.*, 2022). Between diverged species, genomes share less of

their architecture than genomes within species, but typically genome architecture is

conserved in proportion to phylogenetic distance (Luo *et al.*, 2020; Weissensteiner *et al.*,

2020; Derežanin *et al.*, 2022; Ruggieri *et al.*, 2022), and becomes poorly conserved at larger

evolutionary distances (Koonin, 2009).

However, genomes have often, but not always, been viewed as containers to hold genes

(Heng, 2009; Marques, Meier and Seehausen, 2019). The legacy of a gene-centric genome

has persisted due to the Modern Synthesis (Crkvenjakov and Heng, 2022) and the highly

influential work of Dawkins (Dawkins, 1976) and others. Guided by an evolutionary view

dominated by genes and gene variants, many genomes from various species have been

sequenced, and by identifying their genes and gene variants have provided us with a better

understanding of the processes of evolution, divergence, and speciation (Rellstab *et al.*,

2015; Schumer *et al.*, 2015; Meier *et al.*, 2017). However, a heavily gene-centric view may

also limit our understanding (Heng, 2009). This heavily gene variant-based view of evolution

was common until recent advances in long-read sequencing technologies enabled genome-

wide investigations into genome architecture (Amarasinghe *et al.*, 2020). Larger structural

genome changes were thought to be rare, and as such genomes have been treated as

largely structurally static, with individuals typically conceptualised as differing mostly by

single nucleotide polymorphisms (SNPs) (Feulner and De-Kayne, 2017). Pan-genome

studies by using a collection of genes or sequences in a population or species (Bayer *et al.*,

2020; Lei *et al.*, 2021), have revealed a significant amount of structural variation within

genomes (Torkamaneh, Lemay and Belzile, 2021; Tang *et al.*, 2022; Li *et al.*, 2023).

Shared genome architecture is measured by synteny. Synteny is the conservation of both

the order and sequence of homologous chromosomes between genomes (Passarge,

Horsthemke and Farber, 1999; Dawson *et al.*, 2007; Heger and Ponting, 2007). Synteny can

refer both to individual genome regions, or in aggregate when comparing whole genomes. A

genome pair with a large proportion of syntenic loci can be said to be more syntenic than a

genome pair with a small proportion of syntenic loci. Synteny can become disrupted by the

loss, gain, duplication, rearrangement, or divergence of existing sequences.

Rearrangements can occur as inversions, translocations, and duplications; altering the order

of sequences within chromosomes while maintaining gene content and are often labelled as

structural variants (SVs) (Rieseberg, 2001). Species-specific sequences resulting from the

insertion, deletion, or localised divergence of sequence, appear as unaligned regions when

genomes are analysed. The true origin of unaligned regions is more difficult to infer than

rearrangements or syntenic regions (Weisman, Murray and Eddy, 2020).

Crucial to the study of plant genome evolution is study group choice. The ideal study group

would be naturally evolving, have low prezygotic reproductive barriers, highly specious, and

exist over a wide and variable evolutionary range. *Eucalyptus* with over 800 wild and

undomesticated species that exist across a wide geographic and environmental range (Potts

and Wiltshire, 1997; Booth *et al.*, 2015; Supple *et al.*, 2018), retain a conserved karyotype

(Grattapaglia *et al.*, 2015; Butler *et al.*, 2017), are pollinated by generalist pollinators

(Pfeilsticker *et al.*, 2023), are capable of wide-ranging dispersal of genetic material (Bezemer

et al., 2016; Murray *et al.*, 2019), and span 50 million years of divergent evolution (Thornhill

et al., 2019) make an ideal genus to study plant genome evolution.

Continuing our study into plant genome evolution (Ferguson *et al.*, 2022), we generated long

read sequences and assembled the genomes of 30 undomesticated *Eucalyptus* genomes

and outgroups from two closely related genera, *Angophora floribunda* and *Corymbia*

maculata. Combined with three previously and identically assembled *Eucalyptus* genomes

(Ferguson *et al.*, 2022), we create a dataset covering approximately 1-50 million years of

diverging genome evolution, including all eight *Eucalyptus* subgenera (Thornhill *et al.*, 2019;

Nicolle, 2022). Identifying all syntenic and rearranged regions between all species pairs we

demonstrate the rapid pace at which ancestral genome architecture is lost. We further

analysed our results to determine if ancestral genome architecture was being lost to

sequence rearrangement, divergence beyond recognition, or insertions and deletions.

Additionally, by framing synteny, rearrangement, and unaligned loss or gain with

phylogenetic distance we sought to describe the overall pattern of genome evolution.

135 Results

136 Sequencing and Assembly

To investigate genome architecture, we performed nanopore long-read native DNA

sequencing and *de novo* genome assembly for 32 Eucalypt species (30 *Eucalyptus*, one

Angophora, and one *Corymbia*; Table 1). All read libraries were trimmed and filtered in

preparation of assembly. Curated read libraries had an average haploid coverage of 42.8x

(range: 24.7x to 78.0x). For details of read libraries and sequence length distributions, see

Supplementary Table S1 and S2, and Supplementary Figures S1 and S2. *Eucalyptus*

pauciflora fast5 files were obtained from Wang et al (Wang et al., 2020), processed and

assembled as per our datasets, using a randomly selected 60x coverage of reads.

After assembling our trimmed and filtered read libraries, we curated our genomes, removing

contigs identified as contamination and assembly artefacts. Additionally we filtered haplotigs

from our primary assemblies to form pseudo-haploid genomes (Supplementary Table S3).

Our genomes, which have a known and conserved haploid karyotype of 11 chromosomes

(Ribeiro et al., 2016), assembled into an average of 517 contigs (range: 120 to 1,755) (Table

1). At the completion of our assembly pipeline our genomes had an average contig N50 of

3.65 Mbp (range: 614.30 Kbp to 11.10 Mbp). Scaffolding contigs against *E. grandis* (Myburg

et al., 2014) greatly increased our genome contiguity, placing on average 99.69% of our

genomes into pseudo-chromosomes (range: 98.83% to 99.99%). On average each pseudo-

chromosomes was comprised 46 contigs. We have found syntenic scaffolding within

Eucalyptus to be suitable in the absences of chromosome conformation data (Ferguson *et*

al., 2022) as Eucalypts have a conserved karyotype (Healey *et al.*, 2021; Low *et al.*, 2022).

Additionally, within other genera closely related genomes have been found suitable for

scaffolding (Burns *et al.*, 2021). Additionally, RaGOO provides confidence scores for

assigning contigs to a scaffold, ordering contigs within scaffolds, and orienting contigs within

scaffolds. Confidence scores achieved by our genomes indicated scaffolding was

satisfactory, Supplementary Figure S3.

The completeness of our genomes was evaluated with benchmarking universal single-copy

orthologs (BUSCO) (Manni *et al.*, 2021) and LTR Assembly Index (LAI) (Ou, Chen and

Jiang, 2018). A more complete genome will contain a high proportion of single-copy BUSCO

genes, and all our genomes were found to be highly BUSCO complete (average: 97.01%;

range: 95.44% to 98.11). LAI searches a genome for long terminal repeat (LTR) sequences,

and reports on the proportion that are intact. The LAI scores achieved by our genomes

indicate that they are highly complete (average: 18.17; range: 14.50 to 23.85). Quality

scores for all our genomes indicate that our genomes are of high quality, contiguity and

completeness (Table 1 and Supplementary Table S4). For statistics and sequence

distribution plots describing our genomes during and at the completion of assembly (see

Supplementary Tables S5 and S6, and Supplementary Figures S4 to S6).

Table 1. Summary of de novo genome assembly, quality assessment, and annotation of 33 Eucalypt genomes. Alphabetically ordered list of genomes assembled, listing: scaffolded genome size, proportion of contigs that were placed in a scaffold, scaffold N50 score, contig N50 score, the number of contigs in unscaffolded genomes, BUSCO scores for scaffolded genomes, LAI score, and the proportion of genome occupied TEs. *Eucalyptus albens*, *E. melliodora*, and *E. sideroxylon*, have been previously reported, being assembled using the same pipeline (Ferguson *et al.*, 2022).

Species	Scaffolded genome size (Mbp)	% of genome in scaffolds	Scaffold N50 (Mbp)	Contig N50 (Mbp)	Contig count	BUSCO complete	LAI	TE %
A. floribunda	388.21	99.73%	36.02	4.02	224	96.82%	14.5	34.55%
C. maculata	403.82	99.90%	40.55	4.69	173	97.25%	15.92	36.26%
*E. albens	606.89	99.79%	56.93	2.55	674	96.47%	17.3	46.57%
E. ANBG9806169	507.93	99.65%	49.61	2.40	476	96.86%	22.16	44.00%
E. brandiana	507.08	99.82%	45.47	7.28	168	98.11%	23.85	44.21%
E. caleyi	589.32	99.53%	59.52	4.77	276	96.47%	18.24	46.00%
E. camaldulensis	558.45	99.87%	52.65	2.48	418	96.73%	16.99	45.31%
E. cladocalyx	544.08	99.68%	51.92	2.80	390	97.59%	18.53	45.85%
E. cloeziana	480.07	99.75%	44.75	1.74	625	97.12%	19.06	42.57%
E. coolabah	606.31	99.53%	53.56	1.29	935	95.44%	15.9	45.89%
E. curtisii	435.26	99.96%	40.29	2.96	288	97.29%	18.34	41.66%
E. dawsonii	706.90	99.35%	67.73	0.99	1,342	97.51%	17.01	45.88%
E. decipiens	590.95	99.50%	60.20	1.99	552	96.99%	18.87	46.95%
E. erythrocorys	539.20	99.99%	50.47	4.02	250	97.55%	20.18	47.07%
E. fibrosa	589.91	99.85%	55.66	6.45	192	96.73%	17.49	45.10%
E. globulus	545.02	99.28%	51.39	0.64	1,747	96.69%	17.46	44.29%
E. grandis	615.89	99.44%	58.49	0.61	1,747	96.09%	17.11	46.53%
E. guilfoylei	472.36	99.97%	44.61	4.25	209	98.02%	16.39	41.22%
E. lansdowneana	633.52	99.92%	59.67	2.35	489	97.12%	19.46	46.10%
E. leucophloia	568.48	99.38%	54.41	2.66	382	96.99%	17.91	44.37%
E. marginata	512.89	98.83%	50.56	1.01	989	96.17%	19.58	43.43%
*E. melliodora	639.15	99.30%	60.83	1.87	564	98.67%	18.32	47.20%
E. melliodora x E. sideroxylon	603.57	99.80%	57.05	6.22	281	97.72%	17.96	46.71%
E. microcorys	440.91	99.92%	41.20	4.00	233	97.21%	16.2	41.39%
E. paniculata	588.85	99.66%	55.38	3.70	330	97.12%	18.58	44.92%
E. pauciflora	494.03	99.88%	50.46	6.58	209	97.25%	20.29	43.10%
E. polyanthemos	603.28	99.56%	57.46	4.66	300	96.82%	17.52	45.55%
E. pumila	529.75	99.70%	48.19	2.49	473	97.38%	17.74	44.17%
E. regnans	494.97	99.84%	47.06	5.26	205	97.25%	20.18	43.06%
E. shirleyi	597.18	99.88%	56.34	6.91	181	97.29%	19.89	45.85%
*E. sideroxylon	592.133	99.87%	62.13	5.22	297	96.65%	18.68	46.57%
E. tenuipes	397.78	99.99%	35.74	3.43	207	96.39%	15.07	37.82%

E. victrix	557.16	99.85%	53.19	11.10	120	96.65%	18.71	44.34%
E. viminalis	558.71	99.11%	52.93	0.65	1,755	96.47%	16.5	44.57%
E. virginea	532.79	99.97%	56.15	2.39	376	97.08%	17.69	43.78%

Genome annotation

As masking of repeats within genomes aids in gene annotation, we annotated our genomes

for both transposable elements (TEs) and simple repeats. Repeat annotation was performed

using *de novo* repeat libraries built for each genome. Repeat annotation resulted in the

classification of an average of 43.78% (range: 34.55% to 47.07%) of our genomes as TEs,

and an average of 1.25% (range: 1.14% to 1.39%) as simple repeats, Table 1 and

Supplementary Table S7. After soft-masking all genomes, we trained species-specific gene

HMM models and subsequently annotated all genomes for genes. Gene models were

trained on all available gene transcripts for *A. thaliana* (Taxonomy ID: 3702) and Myrtaceae

(Taxonomy ID: 3931) found within the NCBI (Sayers *et al.*, 2021). Annotation predicted an

average of 53,390 (range: 41,623 to 77,764) gene candidates within our genomes,

Supplementary Table S8. While the number of annotated genes is consistent with plant gene

number estimates (Sterck *et al.*, 2007) there is a wide variation between genomes. It is

important to note the genes annotated within these genomes will contain both false positives

and false negatives and are gene candidates, which in addition to real gene number

variations will contribute to the variation in the number of annotated genes.

Eucalyptus pan-genome

Due to the shared evolutionary history of our genomes, many gene candidates will be

homologues that have arisen pre-speciation (orthologs) or post-speciation (paralogs)

(Jensen, 2001). To examine the evolutionary relationship between *Eucalyptus* gene

candidates we placed all highly similar primary (longest) gene transcripts into orthogroups

(OG). Of the 1,761,851 identified gene candidates across our 33 *Eucalyptus* genomes,

1,726,511 (97.99%) were placed into one of 68,248 orthogroups (OG). The remaining

35,340 (2.01%) unique genes were not placed within an OG as their sequences were too

dissimilar (>40% transcript identity and e-value < 0.001) to all other genes. On average each

genome had 98.03% (range: 94.62% to 98.03%) of its gene candidates placed within an

OG. 0.26% (4,551) of all gene candidates were found to occur within a genome-specific OG.

For detailed statistics on orthogrouping see Supplementary Tables S8 and S9. Additionally,

OGs were classified as core (present in all species), dispensable (present in at least two

species), and private (present in a single species) (Figure 1). A total of 21.33% (14,552) of

OG were core, likely representing key *Eucalyptus* genes. Most OGs were dispensable,

76.00% (51,858), which may be a source of phenotypic and adaptive variation within the

species. Only a very small number were private, 2.67% (1,821), potentially representing

highly species-specific genes and newly evolved genes.

Figure 1. Pangenome of 33 species of *Eucalyptus*. Shows the number of orthogroups
shared by an increasing number of genomes. Private orthogroup are orthogroups that exist
within a single genome, core exist in all, and dispensable orthogroups are those that exist in
2 to 32 (n-1) genomes.
225

226 *Eucalyptus* phylogeny

To describe the evolutionary patterns between our genomes we built a phylogenetic tree

from single-copy BUSCO genes. We additionally included the genome of *C. calophylla*

genome, which was identically assembled (Ahrens *et al.*, 2021). From the initial BUSCO set

of 2,326 genes, we selected only genes present within 30 or more genomes, leaving 2,106

BUSCO genes across our 36 genomes or 72,516 total genes. For each gene, we generated

a multi-sequence alignment (MSA) with MAFFT, which we then trimmed and filtered,

removing low abundance regions and genes with overall poor alignments, leaving 1,674

gene MSAs. Each MSA was used to construct a gene tree, subsequently all gene trees were

combined into a consensus species tree. The species tree was manually rooted using the

established relationship between *Angophora*, *Corymbia* and *Eucalyptus* (Thornhill *et al.*,

2019), Figure 2 and Supplementary Figure S7. The species tree in Newick format is

available within Supplementary results.

After constructing the species tree, *E. salubris* was unexpectedly found to be grouped with

E. pauciflora and *E. regnans*. If correctly placed, *E. salubris* would be a sister lineage to the

Adnataria group (*E. victrix* to *E. sideroxylon*) (Thornhill *et al.*, 2019). Morphological

examination of the sample tree revealed that the tree was incorrectly labelled. The correct

species name is currently unknown, as such we use its NCBI name *E. ANBG9806169*.

246 Genome conservation and loss

To resolve the syntenic and non-syntenic regions of our *Eucalyptus* genomes, we performed

one-to-one genome comparisons for all genome pairs. Whole genome alignments for all

comparisons were analysed with SyRI (Goel *et al.*, 2019), and subsequently, all genomic

regions within both genomes of an alignment pair were annotated as syntenic, rearranged

(inversion, translocation, or duplication), or unaligned (sequence that only exists in one

genome, resulting from either an insertion, deletion or sequence divergence). As repeat

masking would inflate the unaligned proportion of genome alignments and bias results, all

genomes remained unmasked. This analysis resulted in all genomes being annotated for

syntenic, rearranged, and unaligned regions 32 times, giving a total of 1,056 annotated

genomes. A visual summary of shared synteny was plotted using our phylogenetic ordering,

Figure 2 and Supplementary Figures S8 to S17. Inspection of synteny plots indicated that

syntenic regions exist across the length of all chromosomes, however, synteny has become

highly fragmented, and is differently maintained.

261
262   **Figure 2. Synteny karyotype of chromosome 1.** Blue ribbons between karyotypes indicate
263   the presence of syntenic sequences between species pairs. In all other regions synteny has
264   become lost. Synteny is lost to either rearrangements (inverted, translocated, or duplicated),
265   sequence divergence, loss or gain. Chromosomes are ordered by our phylogenetic tree.
266

Next, we calculated the proportion of sequences shared between genomes, how sequences

were shared (syntenic, inverted, translocated, and duplicated), and the frequency at which

rearrangements occurred between genomes. For this analysis, we excluded all events less

than 200 bp in length (the majority of which were small unaligned annotations). The majority

of sequence was shared (syntenic and rearranged) between genomes, averaging 69.35%

(range 46.67% to 91.86%). Only four pairwise alignments had <50% shared sequence: *E.*

coolabah, E. dawsonii, E. grandis, and E. melliodora, all when compared to *E. erythrocorys*.

Synteny was the major contributor to shared sequence, averaging 39.32% (range: 21.34% to

60.44%). Rearrangements averaged: 30.24% (range: 16.97% to 49.49%). The remainder of

sequence was annotated as unaligned, averaging 30.43% (range: 8.08% to 53.32%), Table

2. For a per-species comparison breakdown of the percent of genome shared, syntenic,

rearranged, and unaligned see Supplementary Tables S10 to S13.

Examination of the size and frequency of syntenic regions indicates that synteny between

the 11 chromosomes of all genome pairs has, on average, fragmented into 12,153 (range:

6,657 to 18,810) regions with an average size of 17.97 kbp (range: 16.27 kbp to 24.26 kbp)

(see Supplementary Figures S18 and S19 for per genome average event size and frequency

plots). Rearrangements in total (inversions + duplications + translocations) contributed more

to synteny loss than did unaligned regions, however, unaligned contributed more than any

single rearrangement type. A more detailed examination of the relative size and frequency of

syntenic, rearrangement, and unaligned events showed that syntenic regions were long and

common, unaligned regions were short and common, inversions were long and very rare,

duplications were shortest and very common, and translocations occurred at a moderate

frequency and size. Syntenic regions are distributed over the entire length of all

chromosomes between all genome pairs, however, synteny has become highly fragmented

by rearrangements and unaligned regions.

Table 2. Summary of synteny, rearranged, and unaligned statistics of all pairwise
genome analyses. For each pairwise comparison the number and total size of syntenic,
unaligned, inverted, translocated, duplicated, rearranged (sum of all inverted, translocated,
duplicated regions), and total shared was calculated. Additionally, the proportion of the

genome occupied by all event types was calculated. Shown here are the average and range
of these statistics.

Alignment type	Average event size within each pairwise alignment (Kbp)			Event counts			Percent of genome		
	Least	Most	Average	Least	Most	Average	Least	Most	Average
Syntenic	14.03	29.45	18.34	6,657	18,810	12,153	21.34%	60.44%	39.32%
Unaligned	3.28	15.62	7.58	11,808	35,983	22,365	8.08%	53.32%	30.43%
Inverted	29.48	477.33	125.18	81	209	148	0.87%	10.46%	3.28%
Translocated	6.70	22.98	11.65	4,322	15,975	9,521	9.26%	41.14%	19.92%
Duplicated	2.48	6.14	3.60	3,807	32,286	15,350	3.60%	38.97%	14.03%
Rearranged	-	-	-	8,274	46,066	25,020	16.97%	49.49%	30.24%
Total shared	-	-	-	-	-	-	46.67%	91.86%	69.35%

Divergence time and genome conservation/loss

To examine these trends of architecture change over increasing divergence time, we

examined the relationship between phylogenetic distance and genome conservation and

divergence, Figure 3. Not unexpectedly, we find that as phylogenetic distances increase, the

proportion of syntenic ($R^2 = 0.261$) and rearranged ($R^2 = 0.356$) sequence decreased as

lineages acquire unique genomic variation. Similarly, as phylogenetic distances increase, the

proportion of genomes within duplications ($R^2 = 0.189$) and translocations ($R^2 = 0.240$)

decreased. Whereas, the portion of genomes unaligned quickly increases with increasing

phylogenetic distance ($R^2 = 0.536$). Inversions consistently occupied a small proportion of

genomes across all phylogenetic distances ($R^2 = 0.000$).

Unaligned sequences accumulate through the loss, gain, or divergence of sequences. As

genome sizes are similar (average: 552.75 Mbp; standard deviation: 65.62 Mbp) sequence

loss and gain are unlikely to fully explain the rapid accumulation of unaligned sequences.

Divergence beyond recognition is likely the largest contributing factor. To test which regions

were contributing to the growth of unaligned sequences we gathered all alignment identity

scores for all syntenic, inverted, translocated, and duplicated regions, in each pairwise

alignment. Plotting identities against phylogenetic distance, we examined the rate at which

sequences diverge. Syntenic was observed to lose sequence homology more rapidly (R^2:

0.516), than duplicated (R^2: 0.236), translocated (R^2: 0.303), and inverted (R^2: 0.260),

Supplementary Figure SX. However, in all cases the regression spanned a very small

interval (syntenic: 91.58% - 93.14%, duplicated: 91.48% - 92.63%, translocated: 91.56% -

92.81%, and inverted: 91.49% - 92.72%) and none approached our 80% sequence similarity

threshold for alignments.

Overall, we find that the syntenic proportion of the genome decreases slowly with increasing

divergence time, while the proportion rearranged as duplications and translocations

decrease faster. The loss of homology between synteny, duplicated and translocated

regions leads to a strong increase in the unaligned portion of the genome (insertions,

deletions, and diverged sequences) as divergence time increases. The loss of duplications

and translocations contribute more to the growth of unaligned than does synteny.

333
Figure 3. Pairwise genome conservation and loss, as phylogenetic distance
increases. The proportion of both *Eucalyptus* genomes with an alignment pair that was
identified as syntenic, rearranged, or unaligned, plotted against the phylogenetic distance of

the two genomes. The unaligned proportion is the species-specific fraction of the genome
between genome pairs, resulting from either an insertion, deletion, differential inheritance, or
sequence divergence. When combined, the proportion of sequence that is syntenic,
unaligned, and rearranged equals 100% for each genome within an alignment pair. The
rearranged fraction is further broken down into inverted, translocated, and duplicated
regions. Phylogenetic distance was calculated as the sum of branch lengths between each
genome pair within phylogeny. P-value tests if the slope of the regression line is nonzero.
344

345 Genome-specific and group-wide sequences

Unaligned sequences occupied on average 30.65% of each *Eucalyptus* genome within each

pairwise alignment. To determine if these sequences were unique to a single genome or

shared between multiple, all pairwise alignments for each species were combined and the

number of species sharing each base calculated. Subsequently, genome regions that were

unique to a genome, shared by multiple genomes or shared by all genomes identified,

Figure 4.

Genome-specific (unique) sequences occupied an average of 1.36% (241.55 Mbp) of the 33

Eucalyptus genomes, the remaining 98.64% of sequence was shared by one or more

genomes. The proportion of each genome shared by all others averaged 22.83%. This

finding mirrors our OG analysis where 2.67% of groups were private, 76.00% dispensable,

and 21.33% were core.

Of note is *E. erythrocorys*, whose genome had a significantly lower proportion of genome-

specific sequence and higher proportion of sequence shared by all other genomes.

Eucalyptus erythrocorys is the sister taxon of all our other genomes within our *Eucalyptus*

dataset. Given the age of the divergence between the *E. erythrocorys* lineage and its sister

lineage, this genome was expected to display a unique pattern in this analysis, however, the

extent to which *E. erythrocorys* is different from all others was surprising.

The table in the figure:

	Proportion of genome		
	Present in all others	Unique to genome	Shared with 1 or more
Minimum	18.57%	0.57%	90.61%
Average	22.83%	1.36%	98.64%
Maximum	29.17%	9.39%	99.43%

Figure 4. Proportion of *Eucalyptus* genomes unique and shared by all others.
Sequence unique to species is the union of the genome that was classified as unaligned within all pairwise alignments. Sequence within all is the union of the genome that was classified syntenic and rearranged (i.e. common between genomes) within all pairwise alignments.

Lineage-conserved rearrangements

The one-to-one analysis of our *Eucalyptus* genomes has described a genome structure that has become highly fragmented by frequently occurring rearrangements and unaligned regions. As genome structure is inherited by offspring, some of the rearrangements discovered during our analysis are assumed to exist within the genomes of monophyletic groups, i.e. a group of species that have descended from a single ancestral species. Rearrangements found within multiple genomes also help to confirm their validity. To search for evidence of inherited rearrangements we analysed the *Adnataria* section, for which we have the best coverage of genomes (13 genomes, as listed in Figure 5). Additionally, using only *Adnataria* genomes should maximise the occurrence of retained rearrangements, as the phylogenetic distances within the *Adnataria* group are relatively low, with many species still hybridising (Delaporte, Conran and Sedgley, 2001).

As all alignments and subsequent annotations are relative to the two species involved, directly comparing the breakpoints of annotations to find common rearrangements is not

possible. Therefore, an outgroup genome, *E. leucophloia*, a sister species of the *Adnataria*

group, is used for comparison with each of the 13 selected genomes. The outgroup genome

imposes a single set of genetic coordinates and genome architecture, enabling comparisons

of rearrangement breakpoints and subsequent identification of shared rearrangements.

Shared rearrangements will contain the same sequence. While this method allows us to find

common inversions, translocations, and duplications, it doesn't allow us to find unaligned

(insertions, deletions, and highly diverged) regions between our ingroup genomes, as

genomes are not being directly compared.

Comparing the start and end breakpoints (± 50 bp) for events >1 Kbp (250,693 total

rearrangements across all *Adnataria* genomes) identified 58,388 (23.29%) common

rearrangements (rearrangements that exist within two or more genomes). Of the 58,388

common rearrangements, 28,059 (48.06%) were shared by two genomes, and 391 (0.67%)

were shared by all. The number of common rearrangements quickly decreased as the

number of genomes increased, (Supplementary Figure S20). Lineage-conserved

rearrangements were identified by tracing common rearrangements through *Adnataria's*

phylogeny, Figure 5. As expected, more closely related genomes shared the largest number

of rearrangements, while more distant genomes shared less. Additionally, as the number of

descendant genomes of nodes increased, the number of shared rearrangements also

decreased. Inherited rearrangements were identified within the *Adnataria* group. We

repeated this analysis twice using *E. brandiana* and *E. cladocalyx* as the outgroup genome

achieving similar results, Supplementary Figures S21 and S22.

Figure 5. Lineage-conserved rearrangements. Using an outgroup genome, *E. leucophloia*, rearrangements were identified that were shared among members of a lineage, i.e. rearrangements with the same start and end points within the outgroup genome. At each branch in the dendrogram, the number of rearrangements shared by all taxa within that clade are labelled. e.g. *E. victrix* and *E. coolabah* share 6,029 rearrangements.

Gene content of synteny, rearrangements, and unaligned

To assess whether rearrangements that encompass genes are selected against, we

calculated the proportion of genic (contains a gene/s) and non-genic (contains no gene/s)

rearrangements, as well as syntenic and unaligned events per genome. Initially, all events

too small to contain a gene and genes unplaced within an orthogroup were removed. A

conservative event length of 1 kbp was used to filter out events, as events smaller than this

are unlikely to contain a gene (Xu *et al.*, 2006). Genes unplaced within an orthogroup are

highly dissimilar to all other gene candidates and may be false positives resulting from

incorrect annotation. The remaining rearrangement, synteny, and unaligned events were

examined for the presence of genes placed within an OG and subsequently classed as genic

or non-genic.

For each genome, when compared to all other genomes, we calculated the average

proportion of genic syntenic, inverted, translocated, duplicated, and unaligned events, and

plotted the results (Figure 6). An average of 88.80% (range: 82.52% to 95.57%) genic

syntenic events were observed across our genomes. Genic unaligned averaged 41.13%

(range: 19.76% to 73.48%), genic inversions averaged 94.93% (range: 81.65% to 99.13%),

genic translocations averaged 65.70% (range: 48.77% to 83.98%), and genic duplications

averaged 45.71% (range: 30.59% to 79.20%).

Additionally, we analysed the effects of divergence time on the proportion of genic events for

all rearrangement types, synteny, and unaligned, Supplementary Figures S23. Results

indicated that phylogenetic distance has little to no impact on the proportion of genic

syntenic events (R^2 = 0.162), inverted events (R^2 = 0.000), translocation events (R^2 = 0.004),

or duplication events (R^2 = 0.002). Unaligned was the only event type whose genic

proportion was affected by phylogenetic distance. As phylogenetic distance increases,

unaligned events become more genic (R^2 = 0.292; p-value = 0.000).

Figure 6. Average proportion of genic events for each species genome. The proportion of genic events was calculated for each pairwise alignment, and averaged. Error bars indicate the minimum and maximum proportion of genic events found when aligned to all other genomes.

Discussion

In this study we created a large collection of wild and naturally evolving high-quality

Eucalyptus genomes covering 1-50 million years of divergent evolution. Using these

genomes we find a pattern of genome evolution led by an initial rapid accumulation of

rearrangements and subsequently a slow loss of both rearranged and syntenic sequences

as lineage-specific mutations erode sequence homology. Rearrangements, likely due to their

recombination effects and subsequent fixation/reduction of alleles (Faria and Navarro, 2010),

are lost more rapidly than syntenic regions. Translocations and duplications were the major

disruptors of synteny and were rapidly lost as divergence times increased. Inversions did not

contribute substantially to the loss of synteny or the loss of rearrangements, instead

occurring at consistently low rates across all divergence times. As genome sizes remain

constant and little species-specific sequence exists across our dataset, loss of existing

sequence or gain of new sequence provides an unlikely explanation for the growth of

unaligned sequences as divergence times increase.

Duplications are a major contributor to functional and genome divergence (Lynch and

Conery, 2000; Adams and Wendel, 2005) especially within plant lineages (Hanada *et al.*,

2008; Van de Peer, Maere and Meyer, 2009). We found that duplications were highly

abundant between all genomes, and that at smaller divergence times, duplications

contributed strongly to genome divergence. As time since divergence increased, the

contribution of duplications to genome divergence lessened, becoming overshadowed by

unaligned portions of the genome. However, at all phylogenetic distances duplications were

a major contributor to genome divergence (occupying on average 14.03% of genomes

across an average of 15,350 events). The observed pattern of duplication loss as time since

divergence increased was unsurprising, as duplications, while highly important to adaptation

and evolution, are rarely conserved (Inoue *et al.*, 2015; Naseeb *et al.*, 2017). Why some

duplications are preserved while the majority are lost is speculative; however, theory centres

on neofunctionalisation, subfunctionalisation, and novel function evolution (Freeling, Scanlon

and Fowler, 2015; Braasch *et al.*, 2016; Lien *et al.*, 2016; Wu and Cox, 2019). These

hypotheses rely upon the genic properties of duplications, i.e. if duplications don't gain novel

function or retain ancestral function, purifying selection will likely result in their removal (Wu

and Cox, 2019). While duplications were the least genic of all rearrangement types, a

significant number (45.71%) were found to contain genes, likely contributing to their

preservation. Non-genic duplications, being less visible to selection, are likely to experience

increased evolutionary rates (mutations) and genetic drift (Scannell and Wolfe, 2008),

eventually mutating beyond recognition, and ultimately contributing to the unaligned

proportion of alignments. Duplications, both preserved and unpreserved, are likely one of the

greatest sources of genome divergence.

Chromosomal inversions, which are known to be associated with the development of

complex phenotypes, local adaptation, and speciation (Lowry and Willis, 2010; Twyford and

Friedman, 2015; Arostegui *et al.*, 2019), were extremely rare between all genomes (average

148 between genomes) and contributed less than all other types of rearrangement to

genome divergence. This observation was consistent at all phylogenetic distances: as time

since divergence increased, the number of inversions remained constant. A similar finding

was made by Hirabayashi and Owens (Hirabayashi and Owens, 2023). Inversions likely

occur at a high rate within plant genomes (Huang and Rieseberg, 2020), however, a low

number of inversions was identified, suggesting that inversions are strongly selected against

and rarely maintained. To survive underdominant selection, a novel inversion must provide

enough selective advantages to outweigh its disadvantages. Inversions may provide a

selective advantage by rearranging recombination loci, and linking alleles captured within

their bounds. Inversion-linked alleles can be strongly selected for, if adaptive, and rise to

high frequencies within populations (Rieseberg, 2001; Harringmeyer and Hoekstra, 2022).

Additionally, adaptive alleles linked by inversions can be protected from strong gene flow

(Yeaman, 2013). Alternatively, inversions may instead hinder adaptation. If selective

conditions were to alter, previously adaptive inversions could prevent recombination from

producing new allele combinations suitable for the new conditions (Rieseberg, 2001).

Inversions, due to recombination suppression, also reduce effective population size and

increase genetic load, as purifying selection can not purge linked deleterious mutations (Jay

et al., 2021). The inversions identified here, which are assumed to have survived selection,

were all very large, as expected (Wellenreuther and Bernatchez, 2018), with the majority

(94.93%) containing genes. Inversions are rare and contribute little to genome divergence,

but are highly genic and likely play a significant role in adaptation, evolution, and speciation

processes.

Translocations can have similar genomic effects to inversions (Ortiz-Barrientos, Engelstädter

and Rieseberg, 2016), contributing to the development of complex phenotypes, local

adaptation, and speciation by disrupting recombination (Martin *et al.*, 2020). As for

inversions, novel translocations that survive drift must provide enough selective advantages

to outweigh their disadvantages or be removed by purifying or underdominant selection.

Translocations were highly abundant between recently diverged, phylogenetically close

genomes. As time since divergence increased, translocations reduced in frequency but

remained common. Translocations were the most common type of large rearrangement

(average size: 11.2 Kbp), mirroring results obtained by Martin et al. (Martin *et al.*, 2020).

Translocations were much more abundant than inversions, especially when genomes have

recently diverged, suggesting that translocations are less strongly selected against than

inversions, despite having a similar effect on recombination. Additionally, translocations,

while highly genic (65.70%) were less genic than inversions (94.93%). The different genomic

pattern observed for translocations and inversions is possibly due to the effects of local

versus non-local changes to recombination. Meiotic recombination may be more disrupted

when reordered recombination loci are close to their location of origin. If true, purifying

selection acts more strongly on inversions than on translocations. Translocations are

common and along with duplications are a major contributor to genome divergence, possibly

aiding in adaptation, evolution, and speciation processes. However, as the effects and

mechanisms of translocations have been less studied than other rearrangements

(Robberecht *et al.*, 2013), it remains to be seen if they are more likely to have

functional/adaptive significance.

To further investigate the potential importance of the syntenic, rearranged, and unaligned

genome regions identified in our study further research using genome-wide association

studies (GWAS) of phenotypes measured on seedlings in pots or field trials, as well as

landscape and genome-wide genotyping for Genome Environment Association (GEA) scans

for adaptive rearrangements are needed. Within species derived rearrangements are

predicted to be predominantly neutral and exist at low frequencies, while others rising to

higher frequencies could be true lineage-specific adaptive rearrangements. With additional

genomes from populations, the frequency of rearrangements within each species could be

assessed. This would provide insight into the functional significance of the widespread

genomic rearrangements we have found and potentially identify rearrangements conferring

adaptive traits across the landscape.

The *de novo* assembled *Eucalyptus* genomes used in this study were scaffolded based on

homology to *E. grandis* (Myburg *et al.*, 2014). Using non-species-specific scaffolding risks

incorrectly joining contigs, resulting in the inflation or deflation of synteny and introduction of

false positive and false negative rearrangements. However, *Eucalyptus* are known to have a

largely conserved karyotype (Potts and Wiltshire, 1997; Booth *et al.*, 2015; Grattapaglia *et*

al., 2015; Butler *et al.*, 2017; Supple *et al.*, 2018; Healey *et al.*, 2021). Furthermore, a

previous study has demonstrated the viability of such a scaffolding strategy for *Eucalyptus*

genomes in the absence of species-specific scaffolding data such as Hi-C, genetic maps, or

optical maps (Ferguson *et al.*, 2022). Lastly, our pseudo-chromosomes were observed to

consist of very few contigs, suggesting that scaffolding is unlikely to be a significant source

of bias.

Eucalyptus contains over 800 species that exist across a wide geographic and

environmental range, while retaining a largely conserved karyotype (Potts and Wiltshire,

1997; Booth *et al.*, 2015; Grattapaglia *et al.*, 2015; Butler *et al.*, 2017; Supple *et al.*, 2018),

which makes the genus ideal to study plant genome evolution. Here we assembled

representative genomes of 33 species, creating one of the most comprehensive datasets to

study plant genome evolution. These genomes provide a genus-wide resource to study

genome rearrangements, and support future *Eucalyptus* research that require genomic

references. Our findings suggest that following divergence, genome architecture is highly

fragmented predominantly by rearrangements. As genomes continue to diverge, genome

architecture continues to be slowly lost. Additionally, as genomes diverge they increasingly

become unalignable due to the divergence of duplications and translocations. Syntenic

regions also contribute to the growing unalignable proportion of genomes, but at a slower

rate than rearrangements. Duplications and translocations are potentially the greatest

contributors to functional and genome divergence, aiding in the development of complex

phenotypes, and local adaptation. Inversions occur at consistently low rates, contributing

little to genome architecture loss or accumulation of unalignable sequences. However,

inversions were highly genic, much more so than either duplications or translocations, and

likely also play a crucial role in the development of complex phenotypes, and local

adaptation. Genome architecture results from a complex interaction of positive, neutral, and

negative forces, all of which contribute to the evolution, divergence, and adaptability of

species (Koonin, 2009; Huang and Rieseberg, 2020; Mérot *et al.*, 2020). However, owing to

technical limitations, the evolution of genome architecture and its role within biology is not

well understood (Lynch *et al.*, 2011; Cortés *et al.*, 2018; Jiggins, 2019). Here, by describing

the pattern of genome architecture as time since divergence increases of 33 *Eucalyptus*

genomes, we contribute to a better understanding of the evolution of plant genomes.

Rearrangements, along with polyploidy, TEs, and other genome evolutionary mechanisms,

play an important role in plant genome evolution (Galindo-González *et al.*, 2017; Marques,

Meier and Seehausen, 2019; Meudt *et al.*, 2021). Further research in other plant lineages is

required to assess the prominence of rearrangements upon genome evolution.

Materials and Methods

Sampling

Eucalyptus species used in this study were collected throughout multiple locations in

Australia, which are detailed in supplementary results. The majority of collected species are

living collections with accession numbers at the Australian National Botanic Gardens

(Canberra, Australian Capital Territory (ACT)) and Currency Creek Arboretum (Currency

Creek, South Australia). Additional samples were sourced from the Australian National

University (Acton, ACT) the National Arboretum Canberra (Molonglo Valley, ACT), the

University of Tasmania Herbarium (Sandy Bay, Tasmania) and within Eucalyptus woodlands

of southern Tasmania. Leaves were placed in plastic zip-lock bags, lightly sprayed with

water to keep them moist and transported to the lab as soon as possible, where they were

washed with water and stored at -80°C until DNA extraction.

DNA extraction, sequencing, and basecalling

To extract high-molecular weight DNA from recalcitrant Eucalyptus samples, we developed

two methods. Initially we combined a protocol to purify nuclei with hexylene glycol (Bolger et

al., 2014) with a magnetic bead-based DNA extraction protocol (Mayjonade et al., 2016),

which was further developed and is available on Protocols.io in detail (Jones and Borevitz,

2019). This was further optimised and developed, which led to the second method of

adopting a sorbitol pre-wash of homogenate (Inglis et al., 2018) to wash crude nuclei instead

of isolating pure nuclei, followed by a magnetic bead-based DNA extraction, according to

(Jones et al., 2021). We found this method to be more time and resource efficient, hence we

switched to this method for all subsequent high-molecular weight DNA extractions. For each

Eucalyptus sample, the method which was used is listed within the supplementary material,

Table Supplementary Table S1, with the two methods being referred to as nuclei and sorbitol

respectively.

After isolating high-molecular weight DNA, we further purified and size selected the DNA by

using a PippinHT (Sage Science). The DNA was size selected for fragments ≥ 20 kb or ≥ 40

kb depending on DNA yield and molecular weight, which is listed in the supplementary

material, Table S1, for each sample. Two Oxford Nanopore Technologies long-read native

DNA sequencing libraries were prepared for each species according to the manufacturer's

protocol 1D genomic DNA by ligation (SQK-LSK109). E. marginata was an exception, which

had one ligation library as described but the second was a transposome library prep,

according to the manufacturer's protocol for rapid sequencing (SQK-RAD004). Sequencing

was performed on MinION Mk1B devices using two FLO-MIN106D R9.4.1 flow cells per

species. Sequencing output was improved when ONT Flow Cell Wash Kits (EXP-WSH003

and EXP-WSH004) were made available, whereby flow cells were washed when sequencing

declined, primed again and more library was loaded, according to the manufacturer's

instructions. After sequencing was complete, the fast5 reads were basecalled with ONT

Guppy (versions: 3.3.0, 4.0.11, 4.0.14, and 4.0.15; See Supplementary Table S14 for per

species versions).

We complemented the long-read sequencing with highly accurate Illumina short-read

sequencing for later use in genome polishing of the long-read de novo assemblies. Illumina

short-read, whole-genome DNA sequencing libraries were generated using a cost-optimised,

transposome protocol based on Illumina Nextera DNA prep methods (Jones et al., 2023).

The pooled libraries were then size selected for fragments with insert sizes between 350 and

600 bp with a PippinHT (Sage Science). Multiplexed sequencing with other projects was

performed on a NovaSeq 6000 (Illumina), using a lane of an S4 flow cell with a 300 cycle kit

(150 bp paired-end sequencing), at the Biomolecular Resource Facility, Australian National

University, Australian Capital Territory, Australia.

De novo assembly

De novo assembly and annotation was performed using the long-read *de novo* plant

assembly protocol developed by Ferguson et al. (Ferguson, Jones and Borevitz, 2022).

Briefly, fastq reads are quality screened, removing DNA control strand, sequencing adaptors

and low quality read ends (the first and last 200 bp), short reads (>1 Kbp in length), and low

quality reads (average quality <Q7), using the NanoPack set of tools (De Coster *et al.*,

2018). Curated reads are next assembled using the long-read assembler Canu (versions:

1.9 and 2.0) (Koren *et al.*, 2017), which assembles high-quality *Eucalyptus* genomes

(Ferguson *et al.*, 2022). Assemblies were filtered of contamination (non-plant contigs),

assembly artefact, plasmid, and haplotig (contigs that span the same genomic region but

originate from different parental chromosomes) contigs using Blobtools (Laetsch and Blaxter,

2017) and Purge Haplotigs (version: 1.1.0) (Roach, Schmidt and Borneman, 2018). Next, all

assemblies were long-read and then short-read polished, using assembly reads and Illumina

reads originating from the same individual as used for assembly. Long-read polishing was

performed with Racon (Vaser *et al.*, 2017), and short-read with Pilon (version: 1.3.1) (Walker

et al., 2014). Long-read polishing made use of the long-read aligner minimap2 (version:

2.17) (Li, 2018, p. 2), while short-read polishing used (version: 0.7.17) (Li, 2013). Next,

assemblies were filtered to remove all contigs less than 1 Kbp in length. We chose this

contig length threshold so as to maximise genome contiguity while removing all contigs too

small to contain a gene. Finally, assemblies were scaffolded using homology to *E. grandis*

(Myburg *et al.*, 2014). Scaffolding was performed with RaGOO (version 1.1) (Alonge *et al.*,

2019) and minimap2.

After assembly all genomes were quality assessed using Benchmarking Universal Single-

Copy Orthologs (BUSCO; version 5; database: eudicots_odb10.2020-09-10) (Manni *et al.*,

2021), long terminal repeat assembly index (version: 2.9.0; LAI) (Ou, Chen and Jiang,

2018), and assembly statistics.

Transposon and gene annotation, and gene ortho grouping

Genome repeat and gene annotation was also performed using the long-read *de novo* plant

assembly protocol developed by Ferguson et al. (Ferguson, Jones and Borevitz, 2022).

Firstly, *de novo* repeat libraries were created for each genome using EDTA (version: 1.9.6)

(Ou *et al.*, 2019), subsequently all genomes were repeat annotated with RepeatMasker

(version: 4.0.9) (Smit, Hubley and Green, 2020). All genomes were repeat soft-masked and

subsequently annotated for genes. Gene annotation was performed with BRAKER (version

2.1.5; Brůna *et al.*, 2021, p. 2) using GeneMark-EP (version: 4; Brůna, Lomsadze and

Borodovsky, 2020). Gene transcript sequences for model training were obtained from the

National Center for Biotechnology Information (NCBI) (Sayers *et al.*, 2021). Included in gene

training data were all Myrtaceae (Taxonomy ID: 3931) and Arabidopsis thaliana (Taxonomy

ID: 3702) transcripts. All gene candidates were grouped into orthogroups using Orthofinder

(version: 2.5.4) (Emms and Kelly, 2019). Using DIAMOND (Buchfink, Reuter and Drost,

2021) Orthofinder aligned all gene transcripts, grouping those with >40% identity and

achieving an e-score < 0.001.

691 Genome synteny, rearrangement and unaligned annotation

Identification of all shared sequences began by aligning all pairwise combinations of

genomes with the MUMmer (version 3) (Kurtz *et al.*, 2004) tool nucmer (parameters: --

maxmatch -l 40 -b 500 -c 200). Nucmer first identifies all shared 40-mers between genomes

and their locations. Next, 40-mers within 500bp are clustered, creating a list of collinear

blocks or alignments. Lastly, using MUMmer's delta-filter tool alignments are filtered,

removing all alignments less than 200 bp in length and less than 80% similar. A low 80%

alignment similarity score was used as *Eucalyptus* are highly heterozygous (Murray *et al.*,

2019), and a more stringent similarity score may incorrectly filter out real alignments.

Having identified all shared sequences we next annotated all syntenic, rearranged (inverted,

translocated, and duplicated), and unaligned (sequence that only exists in one genome,

resulting from either an insertion, deletion or sequence divergence) sequence between

pairwise genomes using SyRI (version: 1.5) (Goel *et al.*, 2019).

SyRI's use of a directed acyclic graph results in genomes being annotated for smaller

regions, which when occurring in an unbroken series of a single type, get combined. The

resulting output includes both levels of annotations, smaller more fragmented, and larger

and more contiguous. We make use of the larger and more continuous alignments.

Additionally, we combined inverted duplications with duplications, and inverted

translocations with translocations.

712 Phylogeny

Using highly conserved and single-copy genes, BUSCO genes, we built a eucalypt

phylogenetic tree describing the evolutionary relationships between all genomes included in

this study. The phylogenetic tree included four previously and identically assembled

genomes for *E. albens*, *E. melliodora*, *E. sideroxylon*, and *C. calophylla*, creating a dataset

of 36 genomes. To begin, fasta sequences for all single-copy BUSCO genes found within

30+ genomes were collected. Using masce (version 2.03; Ranwez *et al.*, 2018, p. 2) multi-

sequence alignment was performed individually on all genes. As errors within gene MSAs

will subsequently lead to errors in phylogenetic inferences, we trimmed and filtered all gene

MSAs. Gene sequence errors were detected and removed using HmmCleaner version:

0.180750; (Di Franco *et al.*, 2019). HmmCleaner uses a profile-hidden Markov model to

identify sequence segments that poorly fit the gene MSA and subsequently removes them.

Errors resulting from poor alignments were removed using report2AA (parameters: -

min_NT_to_keep_seq 30, -min_seq_to_keep_site 4,-dist_isolate_AA 3, -

min_homology_to_keep_seq 0.5, -min_percent_NT_at_ends 0.7) from the macse program.

report2AA removed sites within MSAs that included less than 30 genomes, had less than 4

informative characters, or had isolated sites (site was more than 3 characters away from the

next non-gap character). Additionally, report2AA removed genome from MSAs that had less

than 50% homology to another genome within the MSA, and trimmed both MSA ends that

had less than 70% of aligned sites as nucleotides (i.e. 26+ genomes had to have a non-gap

character). Additionally, as a result of filtering and trimming MSAs of low quality are

removed.

Individual gene trees were constructed for all filtered and trimmed MSAs using iqtree

(version: 1.6.12; Nguyen *et al.*, 2015) . Finally, all gene trees were concatenated into a

single file, from which a species tree was generated using Astral III version 5.7.3; (Zhang *et*

al., 2018). The resulting species tree was manually rooting at the *Angophora/Corymbia* and

Eucalyptus branch, using Figtree (version: 1.4.4; Rambaut, no date).

Data access

Sequencing data and reference genomes generated in this project are publicly available on

the Sequence Read Archive (SRA) and NCBI genome repository under BioProject

PRJNA509734. Gene predictions and SyRI annotations have been deposited in FigShare

and are available at:

https://figshare.com/projects/Plant_genome_evolution_in_the_genus_Eucalyptus_driven_by

_structural_rearrangements_that_promote_sequence_divergence/97010. All analysis scripts

created and used by this project have been deposited within our github repository:

https://github.com/fergsc/33-Eucs.

Competing interest statement

752 The authors declare that they have no competing interests.

Funding

754 This research was supported by the Australian Research Council (Project code:

755 CE140100008 and DP150103591).

Authors' contributions

758 Scott Ferguson led the project and ran all the analysis. Ashley Jones managed, developed,

759 and performed DNA sampling and sequencing. The project was conceived and designed by

760 all authors. Scott Ferguson wrote the first manuscript draft. All authors contributed to writing

761 and review of the final manuscript.

Acknowledgements

We would like to thank the Australian National Botanic Gardens in Canberra, Australia for

providing plant samples and associated metadata. This research acknowledges the support

provided by the Director of National Parks, the park staff of the Australian National Botanic

Gardens, and Parks Australia. The views expressed in this document do not necessarily

represent the views of the Australian Government.

We thank Dean Nicolle, owner of the Currency Creek Arboretum, South Australia, for

providing samples and support for this project. We also thank Yoav Daniel Bar-Ness, Giant

Tree Expeditions, for collecting E. regnans (the Centurion). We also thank Tamera Beath,

David Stanley, Cynthia Torkel, Rob Lanfear, Wang Weiwen and Brad Potts for their support

and collecting samples.

We would like to thank the ACRF Biomolecular Resource Facility at the John Curtin School

of Medical Research, ANU in Canberra, Australia, where Oxford Nanopore Technologies

PromethION and Illumina NovaSeq 6000 sequencing was conducted. This research

acknowledges the support provided by NCRIS-enabled Bioplatforms Australia infrastructure.

Computational resources were provided by the Australian Government through the National

Computational Infrastructure (NCI) under the ANU Merit Allocation Scheme.

This research is funded by The Australian Research Council Centre of Excellence in Plant

Energy Biology and an Australian Research Council Discovery Grant.

Kevin Murray and Scott Ferguson were supported by Australian Government Research

Training Program scholarships.

References

Adams, K.L. and Wendel, J.F. (2005) 'Polyploidy and genome evolution in plants', *Current*
Opinion in Plant Biology, 8(2), pp. 135–141. Available at:
https://doi.org/10.1016/j.pbi.2005.01.001.

Ahrens, C.W. *et al.* (2021) 'Genomic constraints to drought adaptation', *bioRxiv* [Preprint].
Available at: https://doi.org/10.1101/2021.08.07.455511.

Alonge, M. *et al.* (2019) 'RaGOO: fast and accurate reference-guided scaffolding of draft
genomes', *Genome Biology*, 20(1), p. 224. Available at: https://doi.org/10.1186/s13059-019-
1829-6.

Alonge, M. *et al.* (2020) 'Major Impacts of Widespread Structural Variation on Gene
Expression and Crop Improvement in Tomato', *Cell*, 182(1), pp. 145-161.e23. Available at:
https://doi.org/10.1016/j.cell.2020.05.021.

Amarasinghe, S.L. *et al.* (2020) 'Opportunities and challenges in long-read sequencing data
analysis', *Genome Biology*, 21(1), p. 30. Available at: https://doi.org/10.1186/s13059-020-
1935-5.

Arostegui, M.C. *et al.* (2019) 'Retention of a chromosomal inversion from an anadromous
ancestor provides the genetic basis for alternative freshwater ecotypes in rainbow trout',
Molecular Ecology, 28(6), pp. 1412–1427. Available at: https://doi.org/10.1111/mec.15037.

Bayer, P.E. *et al.* (2020) 'Plant pan-genomes are the new reference', *Nature Plants*, 6(8), pp.
914–920. Available at: https://doi.org/10.1038/s41477-020-0733-0.

Bezemer, N. *et al.* (2016) 'Paternity analysis reveals wide pollen dispersal and high multiple
paternity in a small isolated population of the bird-pollinated Eucalyptus caesia (Myrtaceae)',
Heredity, 117(6), pp. 460–471. Available at: https://doi.org/10.1038/hdy.2016.61.

Bolger, A. *et al.* (2014) 'The genome of the stress-tolerant wild tomato species Solanum
pennellii', *Nature Genetics*, 46(9), pp. 1034–1038. Available at:
https://doi.org/10.1038/ng.3046.

Booth, T.H. *et al.* (2015) 'Native forests and climate change: Lessons from eucalypts', *Forest*
Ecology and Management, 347, pp. 18–29. Available at:
https://doi.org/10.1016/j.foreco.2015.03.002.

Braasch, I. *et al.* (2016) 'The spotted gar genome illuminates vertebrate evolution and
facilitates human-teleost comparisons', *Nature Genetics*, 48(4), pp. 427–437. Available at:
https://doi.org/10.1038/ng.3526.

Brůna, T. *et al.* (2021) 'BRAKER2: automatic eukaryotic genome annotation with GeneMark-
EP+ and AUGUSTUS supported by a protein database', *NAR Genomics and Bioinformatics*,
3(1). Available at: https://doi.org/10.1093/nargab/lqaa108.

Brůna, T., Lomsadze, A. and Borodovsky, M. (2020) 'GeneMark-EP+: eukaryotic gene
prediction with self-training in the space of genes and proteins', *NAR Genomics and*
Bioinformatics, 2(2), p. lqaa026. Available at: https://doi.org/10.1093/nargab/lqaa026.

Buchfink, B., Reuter, K. and Drost, H.-G. (2021) 'Sensitive protein alignments at tree-of-life

scale using DIAMOND', *Nature Methods*, 18(4), pp. 366–368. Available at:
https://doi.org/10.1038/s41592-021-01101-x.

Burns, R. *et al.* (2021) 'Gradual evolution of allopolyploidy in Arabidopsis suecica', *Nature*
Ecology & Evolution, 5(10), pp. 1367–1381. Available at: https://doi.org/10.1038/s41559-
021-01525-w.

Butler, J.B. *et al.* (2017) 'Comparative genomics of Eucalyptus and Corymbia reveals low
rates of genome structural rearrangement', *BMC Genomics*, 18(1), p. 397. Available at:
https://doi.org/10.1186/s12864-017-3782-7.

Cortés, A.J. *et al.* (2018) 'Does the Genomic Landscape of Species Divergence in
Phaseolus Beans Coerce Parallel Signatures of Adaptation and Domestication?', *Frontiers in*
Plant Science, 9. Available at: https://www.frontiersin.org/articles/10.3389/fpls.2018.01816
(Accessed: 27 March 2023).

Crkvenjakov, R. and Heng, H.H. (2022) 'Further illusions: On key evolutionary mechanisms
that could never fit with Modern Synthesis', *Progress in Biophysics and Molecular Biology*,
169–170, pp. 3–11. Available at: https://doi.org/10.1016/j.pbiomolbio.2021.10.002.

Dawkins, R. (1976) *The selfish gene*. Oxford University Press.

Dawson, D.A. *et al.* (2007) 'Gene Order and Recombination Rate in Homologous
Chromosome Regions of the Chicken and a Passerine Bird', *Molecular Biology and*
Evolution, 24(7), pp. 1537–1552. Available at: https://doi.org/10.1093/molbev/msm071.

De Coster, W. *et al.* (2018) 'NanoPack: visualizing and processing long-read sequencing
data', *Bioinformatics*. Edited by B. Berger, 34(15), pp. 2666–2669. Available at:
https://doi.org/10.1093/bioinformatics/bty149.

Delaporte, K.L., Conran, J.G. and Sedgley, M. (2001) 'Interspecific Hybridization within
Eucalyptus (Myrtaceae): Subgenus *Symphyomyrtus* , Sections *Bisectae* and *Adnataria*',
International Journal of Plant Sciences, 162(6), pp. 1317–1326. Available at:
https://doi.org/10.1086/323276.

Derežanin, L. *et al.* (2022) 'Multiple types of genomic variation contribute to adaptive traits in
the mustelid subfamily Guloninae', *Molecular Ecology*, 31(10), pp. 2898–2919. Available at:
https://doi.org/10.1111/mec.16443.

Di Franco, A. *et al.* (2019) 'Evaluating the usefulness of alignment filtering methods to
reduce the impact of errors on evolutionary inferences', *BMC Evolutionary Biology*, 19(1), p.
21. Available at: https://doi.org/10.1186/s12862-019-1350-2.

Dixon, J.R., Gorkin, D.U. and Ren, B. (2016) 'Chromatin Domains: The Unit of Chromosome
Organization', *Molecular Cell*, 62(5), pp. 668–680. Available at:
https://doi.org/10.1016/j.molcel.2016.05.018.

Emms, D.M. and Kelly, S. (2019) 'OrthoFinder: phylogenetic orthology inference for
comparative genomics', *Genome Biology*, 20(1), p. 238. Available at:
https://doi.org/10.1186/s13059-019-1832-y.

Faria, R. and Navarro, A. (2010) 'Chromosomal speciation revisited: rearranging theory with
pieces of evidence', *Trends in Ecology & Evolution*, 25(11), pp. 660–669. Available at:
https://doi.org/10.1016/j.tree.2010.07.008.

Ferguson, S. *et al.* (2022) 'Interspecies genome divergence is predominantly due to frequent
small scale rearrangements in Eucalyptus', *Molecular Ecology*, n/a(n/a). Available at:
https://doi.org/10.1111/mec.16608.

Ferguson, S., Jones, A. and Borevitz, J. (2022) *Plant assemble - Plant de novo genome*
assembly, scaffolding and annotation for genomic studies, *protocols.io*. Available at:
https://dx.doi.org/10.17504/protocols.io.81wgb6zk3lpk/v1 (Accessed: 4 August 2022).

Feulner, P.G.D. and De-Kayne, R. (2017) 'Genome evolution, structural rearrangements and
speciation', *Journal of Evolutionary Biology*, 30(8), pp. 1488–1490. Available at:
https://doi.org/10.1111/jeb.13101.

Freeling, M., Scanlon, M.J. and Fowler, J.E. (2015) 'Fractionation and subfunctionalization
following genome duplications: mechanisms that drive gene content and their
consequences', *Current Opinion in Genetics & Development*, 35, pp. 110–118. Available at:
https://doi.org/10.1016/j.gde.2015.11.002.

Galindo-González, L. *et al.* (2017) 'LTR-retrotransposons in plants: Engines of evolution',
Gene, 626, pp. 14–25. Available at: https://doi.org/10.1016/j.gene.2017.04.051.

Goel, M. *et al.* (2019) 'SyRI: finding genomic rearrangements and local sequence differences
from whole-genome assemblies', *Genome Biology*, 20(1), p. 277. Available at:
https://doi.org/10.1186/s13059-019-1911-0.

Grattapaglia, D. *et al.* (2015) 'A novel genome-wide microsatellite resource for species of
Eucalyptus with linkage-to-physical correspondence on the reference genome sequence',
Molecular Ecology Resources, 15(2), pp. 437–448. Available at:
https://doi.org/10.1111/1755-0998.12317.

Hanada, K. *et al.* (2008) 'Importance of lineage-specific expansion of plant tandem
duplicates in the adaptive response to environmental stimuli', *Plant Physiology*, 148(2), pp.
993–1003. Available at: https://doi.org/10.1104/pp.108.122457.

Hardigan, M.A. *et al.* (2020) 'Genome Synteny Has Been Conserved Among the Octoploid
Progenitors of Cultivated Strawberry Over Millions of Years of Evolution', *Frontiers in Plant*
Science, 10, p. 1789. Available at: https://doi.org/10.3389/fpls.2019.01789.

Harringmeyer, O.S. and Hoekstra, H.E. (2022) 'Chromosomal inversion polymorphisms
shape the genomic landscape of deer mice', *Nature Ecology & Evolution*, 6(12), pp. 1965–
1979. Available at: https://doi.org/10.1038/s41559-022-01890-0.

Healey, A.L. *et al.* (2021) 'Pests, diseases, and aridity have shaped the genome of Corymbia
citriodora', *Communications Biology*, 4(1), p. 537. Available at:
https://doi.org/10.1038/s42003-021-02009-0.

Heger, A. and Ponting, C.P. (2007) 'Evolutionary rate analyses of orthologs and paralogs
from 12 Drosophila genomes', *Genome Research*, 17(12), pp. 1837–1849. Available at:
https://doi.org/10.1101/gr.6249707.

Heng, H.H.Q. *et al.* (2004) 'Chromatin loops are selectively anchored using scaffold/matrix-
attachment regions', *Journal of Cell Science*, 117(7), pp. 999–1008. Available at:
https://doi.org/10.1242/jcs.00976.

Heng, H.H.Q. (2009) 'The genome-centric concept: resynthesis of evolutionary theory',
BioEssays, 31(5), pp. 512–525. Available at: https://doi.org/10.1002/bies.200800182.

Hirabayashi, K. and Owens, G.L. (2023) 'The rate of chromosomal inversion fixation in plant
genomes is highly variable', *Evolution*, 77(4), pp. 1117–1130. Available at:
https://doi.org/10.1093/evolut/qpad027.

Huang, K. and Rieseberg, L.H. (2020) 'Frequency, Origins, and Evolutionary Role of
Chromosomal Inversions in Plants', *Frontiers in Plant Science*, 11. Available at:
https://www.frontiersin.org/articles/10.3389/fpls.2020.00296 (Accessed: 21 December 2022).

Inglis, P.W. *et al.* (2018) 'Fast and inexpensive protocols for consistent extraction of high
quality DNA and RNA from challenging plant and fungal samples for high-throughput SNP
genotyping and sequencing applications', *PLOS ONE*, 13(10), p. e0206085. Available at:
https://doi.org/10.1371/journal.pone.0206085.

Inoue, J. *et al.* (2015) 'Rapid genome reshaping by multiple-gene loss after whole-genome
duplication in teleost fish suggested by mathematical modeling', *Proceedings of the National
Academy of Sciences*, 112(48), pp. 14918–14923. Available at:
https://doi.org/10.1073/pnas.1507669112.

Jay, P. *et al.* (2021) 'Mutation load at a mimicry supergene sheds new light on the evolution
of inversion polymorphisms', *Nature Genetics*, 53(3), pp. 288–293. Available at:
https://doi.org/10.1038/s41588-020-00771-1.

Jensen, R.A. (2001) 'Orthologs and paralogs - we need to get it right', *Genome Biology*, 2(8),
p. interactions1002.1-interactions1002.3.

Jiao, W.-B. and Schneeberger, K. (2020) 'Chromosome-level assemblies of multiple
Arabidopsis genomes reveal hotspots of rearrangements with altered evolutionary
dynamics', *Nature Communications*, 11(1), p. 989. Available at:
https://doi.org/10.1038/s41467-020-14779-y.

Jiggins, C.D. (2019) 'Can genomics shed light on the origin of species?', *PLOS Biology*,
17(8), p. e3000394. Available at: https://doi.org/10.1371/journal.pbio.3000394.

Jones, A. *et al.* (2021) 'High-molecular weight DNA extraction, clean-up and size selection
for long-read sequencing', *PLOS ONE*. Edited by M. Eppinger, 16(7), p. e0253830. Available
at: https://doi.org/10.1371/journal.pone.0253830.

Jones, A. *et al.* (2023) 'Cost-conscious generation of multiplexed short-read DNA libraries for
whole-genome sequencing', *PLOS ONE*, 18(1), pp. 1–7. Available at:
https://doi.org/10.1371/journal.pone.0280004.

Jones, A. and Borevitz, J. (2019) *Nuclear DNA purification from recalcitrant plant species for
long-read sequencing*, *protocols.io*. Available at: https://www.protocols.io/view/nuclear-dna-
purification-from-recalcitrant-plant-s-28bghsn (Accessed: 11 January 2022).

Koonin, E.V. (2009) 'Evolution of genome architecture', *The International Journal of
Biochemistry & Cell Biology*, 41(2), pp. 298–306. Available at:
https://doi.org/10.1016/j.biocel.2008.09.015.

Koren, S. *et al.* (2017) 'Canu: scalable and accurate long-read assembly via adaptive *k* -mer
weighting and repeat separation', *Genome Research*, 27(5), pp. 722–736. Available at:
https://doi.org/10.1101/gr.215087.116.

Kurtz, S. *et al.* (2004) 'Versatile and open software for comparing large genomes', *Genome
Biology*, p. 9.

Laetsch, D.R. and Blaxter, M.L. (2017) 'BlobTools: Interrogation of genome assemblies',
F1000Research, 6, p. 1287. Available at: https://doi.org/10.12688/f1000research.12232.1.

Lei, L. *et al.* (2021) 'Plant Pan-Genomics Comes of Age', *Annual Review of Plant Biology*,
72(1), pp. 411–435. Available at: https://doi.org/10.1146/annurev-arplant-080720-105454.

Li, H. (2013) 'Aligning sequence reads, clone sequences and assembly contigs with BWA-
MEM', *arXiv:1303.3997 [q-bio]* [Preprint]. Available at: http://arxiv.org/abs/1303.3997
(Accessed: 5 August 2021).

Li, H. (2018) 'Minimap2: pairwise alignment for nucleotide sequences', *Bioinformatics*. Edited
by I. Birol, 34(18), pp. 3094–3100. Available at:
https://doi.org/10.1093/bioinformatics/bty191.

Li, N. *et al.* (2023) 'Super-pangenome analyses highlight genomic diversity and structural
variation across wild and cultivated tomato species', *Nature Genetics*, 55(5), pp. 852–860.
Available at: https://doi.org/10.1038/s41588-023-01340-y.

Lien, S. *et al.* (2016) 'The Atlantic salmon genome provides insights into rediploidization',
Nature, 533(7602), pp. 200–205. Available at: https://doi.org/10.1038/nature17164.

Lin, Y.-L. and Gokcumen, O. (2019) 'Fine-Scale Characterization of Genomic Structural
Variation in the Human Genome Reveals Adaptive and Biomedically Relevant Hotspots',
Genome Biology and Evolution, 11(4), pp. 1136–1151. Available at:
https://doi.org/10.1093/gbe/evz058.

Low, Y.W. *et al.* (2022) 'Genomic insights into rapid speciation within the world's largest tree
genus Syzygium', *Nature Communications*, 13(1), p. 5031. Available at:
https://doi.org/10.1038/s41467-022-32637-x.

Lowry, D.B. and Willis, J.H. (2010) 'A Widespread Chromosomal Inversion Polymorphism
Contributes to a Major Life-History Transition, Local Adaptation, and Reproductive Isolation',
PLOS Biology, 8(9), p. e1000500. Available at: https://doi.org/10.1371/journal.pbio.1000500.

Luo, X. *et al.* (2020) 'An ultra-high-density genetic map provides insights into genome
synteny, recombination landscape and taproot skin colour in radish (Raphanus sativus L.)',
Plant Biotechnology Journal, 18(1), pp. 274–286. Available at:
https://doi.org/10.1111/pbi.13195.

Lynch, M. *et al.* (2011) 'The Repatterning of Eukaryotic Genomes by Random Genetic Drift',
Annual Review of Genomics and Human Genetics, 12(1), pp. 347–366. Available at:
https://doi.org/10.1146/annurev-genom-082410-101412.

Lynch, M. and Conery, J.S. (2000) 'The Evolutionary Fate and Consequences of Duplicate
Genes', 290(5494), pp. 1151–1155. Available at:
https://doi.org/10.1126/science.290.5494.1151.

Manni, M. *et al.* (2021) 'BUSCO Update: Novel and Streamlined Workflows along with
Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral
Genomes', *Molecular Biology and Evolution*. Edited by J. Kelley, 38(10), pp. 4647–4654.
Available at: https://doi.org/10.1093/molbev/msab199.

Marques, D.A., Meier, J.I. and Seehausen, O. (2019) 'A Combinatorial View on Speciation
and Adaptive Radiation', *Trends in Ecology & Evolution*, 34(6), pp. 531–544. Available at:
https://doi.org/10.1016/j.tree.2019.02.008.

Martin, G. *et al.* (2020) 'Chromosome reciprocal translocations have accompanied
subspecies evolution in bananas', *The Plant Journal*, 104(6), pp. 1698–1711. Available at:
https://doi.org/10.1111/tpj.15031.

Mayjonade, B. *et al.* (2016) 'Extraction of high-molecular-weight genomic DNA for long-read
sequencing of single molecules', *BioTechniques*, 61(4), pp. 203–205. Available at:
https://doi.org/10.2144/000114460.

Meier, J.I. *et al.* (2017) 'Ancient hybridization fuels rapid cichlid fish adaptive radiations',
Nature Communications, 8(1), p. 14363. Available at: https://doi.org/10.1038/ncomms14363.

Mérot, C. *et al.* (2020) 'A Roadmap for Understanding the Evolutionary Significance of
Structural Genomic Variation', *Trends in Ecology & Evolution*, 35(7), pp. 561–572. Available
at: https://doi.org/10.1016/j.tree.2020.03.002.

Meudt, H.M. *et al.* (2021) 'Polyploidy on Islands: Its Emergence and Importance for
Diversification', *Frontiers in Plant Science*, 12. Available at:
https://www.frontiersin.org/articles/10.3389/fpls.2021.637214 (Accessed: 12 October 2022).

Murray, K.D. *et al.* (2019) 'Landscape drivers of genomic diversity and divergence in
woodland Eucalyptus', *Molecular Ecology*, 28(24), pp. 5232–5247. Available at:
https://doi.org/10.1111/mec.15287.

Myburg, A.A. *et al.* (2014) 'The genome of Eucalyptus grandis', *Nature*, 510(7505), pp. 356–
362. Available at: https://doi.org/10.1038/nature13308.

Naseeb, S. *et al.* (2017) 'Rapid functional and evolutionary changes follow gene duplication
in yeast', *Proceedings of the Royal Society B: Biological Sciences*, 284(1861), p. 20171393.
Available at: https://doi.org/10.1098/rspb.2017.1393.

Nguyen, L.-T. *et al.* (2015) 'IQ-TREE: A Fast and Effective Stochastic Algorithm for
Estimating Maximum-Likelihood Phylogenies', *Molecular Biology and Evolution*, 32(1), pp.
268–274. Available at: https://doi.org/10.1093/molbev/msu300.

Nicolle, D. (2022) 'Classification of the eucalypts (Angophora, Corymbia and Eucalyptus)
Version 6'. Available at: http://www.dn.com.au/Classification-Of-The-Eucalypts.pdf.

Ortiz-Barrientos, D., Engelstädter, J. and Rieseberg, L.H. (2016) 'Recombination Rate
Evolution and the Origin of Species', *Trends in Ecology & Evolution*, 31(3), pp. 226–236.
Available at: https://doi.org/10.1016/j.tree.2015.12.016.

Ou, S. *et al.* (2019) 'Benchmarking transposable element annotation methods for creation of
a streamlined, comprehensive pipeline', *Genome Biology*, 20(1), p. 275. Available at:
https://doi.org/10.1186/s13059-019-1905-y.

Ou, S., Chen, J. and Jiang, N. (2018) 'Assessing genome assembly quality using the LTR
Assembly Index (LAI)', *Nucleic Acids Research* [Preprint]. Available at:
https://doi.org/10.1093/nar/gky730.

Oudelaar, A.M. and Higgs, D.R. (2021) 'The relationship between genome structure and
function', *Nature Reviews Genetics*, 22(3), pp. 154–168. Available at:
https://doi.org/10.1038/s41576-020-00303-x.

Passarge, E., Horsthemke, B. and Farber, R.A. (1999) 'Incorrect use of the term synteny',
Nature Genetics, 23(4), pp. 387–387. Available at: https://doi.org/10.1038/70486.

Pfeilsticker, T.R. *et al.* (2023) 'Molecular insights into the dynamics of species invasion by
hybridisation in Tasmanian eucalypts', *Molecular Ecology*, 32(11), pp. 2913–2929. Available
at: https://doi.org/10.1111/mec.16892.

Potts, B.M. and Wiltshire, R.J.E. (1997) 'Eucalypt genetics and genecology', in J. Williams
and J. Woinarski (eds). Cambridge University Press, pp. 56–91. Available at:
https://eprints.utas.edu.au/7460/ (Accessed: 30 November 2021).

Rambaut, A. (no date) *FigTree*. Available at: http://tree.bio.ed.ac.uk/software/figtree/
(Accessed: 28 November 2022).

Ranwez, V. *et al.* (2018) 'MACSE v2: Toolkit for the Alignment of Coding Sequences
Accounting for Frameshifts and Stop Codons', *Molecular Biology and Evolution*, 35(10), pp.
2582–2584. Available at: https://doi.org/10.1093/molbev/msy159.

Rellstab, C. *et al.* (2015) 'A practical guide to environmental association analysis in
landscape genomics', *Molecular Ecology*, 24(17), pp. 4348–4370. Available at:
https://doi.org/10.1111/mec.13322.

Ribeiro, T. *et al.* (2016) 'Advancing Eucalyptus Genomics: Cytogenomics Reveals
Conservation of Eucalyptus Genomes', *Frontiers in Plant Science*, 7. Available at:
https://doi.org/10.3389/fpls.2016.00510.

Rieseberg, L.H. (2001) 'Chromosomal rearrangements and speciation', p. 8.

Roach, M.J., Schmidt, S.A. and Borneman, A.R. (2018) 'Purge Haplotigs: allelic contig
reassignment for third-gen diploid genome assemblies', *BMC Bioinformatics*, 19(1), p. 460.
Available at: https://doi.org/10.1186/s12859-018-2485-7.

Robberecht, C. *et al.* (2013) 'Nonallelic homologous recombination between
retrotransposable elements is a driver of de novo unbalanced translocations', *Genome
Research*, 23(3), pp. 411–418. Available at: https://doi.org/10.1101/gr.145631.112.

Ruggieri, A.A. *et al.* (2022) 'A butterfly pan-genome reveals that a large amount of structural
variation underlies the evolution of chromatin accessibility', *Genome Research*, 32(10), pp.
1862–1875. Available at: https://doi.org/10.1101/gr.276839.122.

Sayers, E.W. *et al.* (2021) 'Database resources of the National Center for Biotechnology
Information', *Nucleic Acids Research*, 49(D1), pp. D10–D17. Available at:
https://doi.org/10.1093/nar/gkaa892.

Scannell, D.R. and Wolfe, K.H. (2008) 'A burst of protein sequence evolution and a
prolonged period of asymmetric evolution follow gene duplication in yeast', *Genome
Research*, 18(1), pp. 137–147. Available at: https://doi.org/10.1101/gr.6341207.

Schumer, M. *et al.* (2015) 'Reproductive Isolation of Hybrid Populations Driven by Genetic
Incompatibilities', *PLoS Genetics*, 11(3), p. e1005041. Available at:
https://doi.org/10.1371/journal.pgen.1005041.

Simakov, O. *et al.* (2020) 'Deeply conserved synteny resolves early events in vertebrate
evolution', *Nature Ecology & Evolution*, 4(6), pp. 820–830. Available at:
https://doi.org/10.1038/s41559-020-1156-z.

Smit, A., Hubley, R. and Green, P. (2020) *RepeatMasker Open-4.0*. Available at:
<http://www.repeatmasker.org> (Accessed: 11 February 2020).

Sterck, L. *et al.* (2007) 'How many genes are there in plants (… and why are they there)?',
Current Opinion in Plant Biology, 10(2), pp. 199–203. Available at:
https://doi.org/10.1016/j.pbi.2007.01.004.

Supple, M.A. *et al.* (2018) 'Landscape genomic prediction for restoration of a *Eucalyptus*
foundation species under climate change', *eLife*. Edited by D.J. Kliebenstein, 7, p. e31835.
Available at: https://doi.org/10.7554/eLife.31835.

Tang, D. *et al.* (2022) 'Genome evolution and diversity of wild and cultivated potatoes',
Nature, 606(7914), pp. 535–541. Available at: https://doi.org/10.1038/s41586-022-04822-x.

Thornhill, A.H. *et al.* (2019) 'A dated molecular perspective of eucalypt taxonomy, evolution
and diversification', *Australian Systematic Botany*, 32, pp. 29–48. Available at:
https://doi.org/10.1071/SB18015.

Torkamaneh, D., Lemay, M.-A. and Belzile, F. (2021) 'The pan-genome of the cultivated
soybean (PanSoy) reveals an extraordinarily conserved gene content', *Plant Biotechnology*
Journal, 19(9), pp. 1852–1862. Available at: https://doi.org/10.1111/pbi.13600.

Twyford, A.D. and Friedman, J. (2015) 'Adaptive divergence in the monkey flower Mimulus
guttatus is maintained by a chromosomal inversion', *Evolution*, 69(6), pp. 1476–1486.
Available at: https://doi.org/10.1111/evo.12663.

Van de Peer, Y., Maere, S. and Meyer, A. (2009) 'The evolutionary significance of ancient
genome duplications', *Nature Reviews Genetics*, 10(10), pp. 725–732. Available at:
https://doi.org/10.1038/nrg2600.

Vaser, R. *et al.* (2017) 'Fast and accurate de novo genome assembly from long uncorrected
reads', *Genome Research*, 27(5), pp. 737–746. Available at:
https://doi.org/10.1101/gr.214270.116.

Walker, B.J. *et al.* (2014) 'Pilon: An Integrated Tool for Comprehensive Microbial Variant
Detection and Genome Assembly Improvement', *PLoS ONE*. Edited by J. Wang, 9(11), p.
e112963. Available at: https://doi.org/10.1371/journal.pone.0112963.

Wang, W. *et al.* (2020) 'The draft nuclear genome assembly of Eucalyptus pauciflora: a
pipeline for comparing de novo assemblies', *GigaScience*, 9(1), p. giz160. Available at:
https://doi.org/10.1093/gigascience/giz160.

Weisman, C.M., Murray, A.W. and Eddy, S.R. (2020) 'Many, but not all, lineage-specific
genes can be explained by homology detection failure', *PLOS Biology*, 18(11), p. e3000862.
Available at: https://doi.org/10.1371/journal.pbio.3000862.

Weissensteiner, M.H. *et al.* (2020) 'Discovery and population genomics of structural variation
in a songbird genus', *Nature Communications*, 11(1), p. 3403. Available at:
https://doi.org/10.1038/s41467-020-17195-4.

Wellenreuther, M. and Bernatchez, L. (2018) 'Eco-Evolutionary Genomics of Chromosomal
Inversions', *Trends in Ecology & Evolution*, 33(6), pp. 427–440. Available at:
https://doi.org/10.1016/j.tree.2018.04.002.

Wu, B. and Cox, M.P. (2019) 'Greater genetic and regulatory plasticity of retained duplicates
in Epichloë endophytic fungi', *Molecular Ecology*, 28(23), pp. 5103–5114. Available at:
https://doi.org/10.1111/mec.15275.

Xu, L. *et al.* (2006) 'Average Gene Length Is Highly Conserved in Prokaryotes and
Eukaryotes and Diverges Only Between the Two Kingdoms', *Molecular Biology and*
Evolution, 23(6), pp. 1107–1108. Available at: https://doi.org/10.1093/molbev/msk019.

Yeaman, S. (2013) 'Genomic rearrangements and the evolution of clusters of locally
adaptive loci', *Proceedings of the National Academy of Sciences*, 110(19), pp. E1743–
E1751. Available at: https://doi.org/10.1073/pnas.1219381110.

Zhang, C. *et al.* (2018) 'ASTRAL-III: polynomial time species tree reconstruction from
partially resolved gene trees', *BMC Bioinformatics*, 19(6), p. 153. Available at:
https://doi.org/10.1186/s12859-018-2129-y.