Evidence for a role of human blood-borne factors in mediating age-associated changes in molecular circadian rhythms

Authors: Jessica E. Schwarz[1,2], Antonijo Mrčela[3], Nicholas F. Lahens[3], Yongjun Li[2], Cynthia T. Hsu[1,2], Gregory Grant[3,4], Carsten Skarke[2,3], Shirley L. Zhang[1,2,5*], and Amita Sehgal[1,2*]

Affiliations:

[1]Howard Hughes Medical Institute, Perelman School of Medicine, University of Pennsylvania; Philadelphia, PA 19104, United States.

[2]Chronobiology and Sleep Institute, Perelman School of Medicine, University of Pennsylvania; Philadelphia, PA 19104, United States.

[3]Institute for Translational Medicine and Therapeutics (ITMAT), Perelman School of Medicine, University of Pennsylvania; Philadelphia, PA 19104, United States.

[4]Department of Genetics, Perelman School of Medicine, University of Pennsylvania; Philadelphia, PA 19104, United States.

[5]Current Institution: Department of Cell Biology, Emory University School of Medicine; Atlanta, GA 30323, United States.

*Email: Amita Sehgal, amita@pennmedicine.upenn.edu ; Shirley L. Zhang, shirley.zhang2@emory.edu

Author Contributions: Conceptualization, J.E.S. and A.S.; Methodology, J.E.S, A.M., N.F.L., Y.L., C.T.H., G.G., C.S., S.L.Z, and A.S.; Formal Analysis, J.E.S, S.L.Z, A.M., N.F.L., Y.L., and C.T.H; Investigation, J.E.S, S.L.Z., and C.S.; Writing – Original Draft Preparation, J.E.S. and A.S.; Writing – Review & Editing, J.E.S, A.M., N.F.L., Y.L., C.T.H., G.G., C.S., S.L.Z, and A.S.; Visualization, J.E.S., A.M, N.F.L.; Supervision, S.L.Z and A.S.

ABSTRACT

Aging is associated with a number of physiologic changes including perturbed circadian

rhythms; however, mechanisms by which rhythms are altered remain unknown. To test the idea

that circulating factors mediate age-dependent changes in peripheral rhythms, we compared the

ability of human serum from young and old individuals to synchronize circadian rhythms in

culture. We collected blood from apparently healthy young (age 25-30) and old (age 70-76)

individuals and used the serum to synchronize cultured fibroblasts. We found that young and old

sera are equally competent at driving robust ~24h oscillations of a luciferase reporter driven by

clock gene promoter. However, cyclic gene expression is affected, such that young and old sera

drive cycling of different genes. While genes involved in the cell cycle and

transcription/translation remain rhythmic in both conditions, genes identified by STRING and

IPA analyses as associated with oxidative phosphorylation and Alzheimer's Disease lose

rhythmicity in the aged condition. Also, the expression of cycling genes associated with

cholesterol biosynthesis increases in the cells entrained with old serum. We did not observe a

global difference in the distribution of phase between groups, but find that peak expression of

several clock controlled genes (*PER3, NR1D1, NR1D2, CRY1, CRY2,* and *TEF*) lags in the

cells synchronized with old serum. Taken together, these findings demonstrate that age-

dependent blood-borne factors affect peripheral circadian rhythms in cells and have the

potential to impact health and disease via maintaining or disrupting rhythms respectively.

INTRODUCTION

Circadian rhythms are known to regulate homeostatic physiology including sleep:wake,

hormone production and body temperature, and their dysregulation with aging is accompanied

by adverse health consequences[1–3], raising the possibility that health decline with age is caused

in part by circadian dysfunction. Although the mechanisms responsible for age effects on

circadian rhythms are unknown, signals from the central clock in the suprachiasmatic nucleus

(SCN) dampen with age[3–5] and rhythms change in peripheral tissues in different ways[5,6]. Here

we aimed to develop a culture model to study the effect of aging on human rhythms of

peripheral tissues. Given that serum can reset the clock in peripheral fibroblasts[7], we

questioned the extent to which serum factors normally contribute to peripheral rhythms of gene

expression and how they might affect rhythms with age, given that blood-borne factors can

influence other aspects of aging[8].

We collected blood samples from young (age 25-30) and old (age 70-76) apparently

healthy individuals following observations of their behavior and physiology, and tested the

hypothesis that age-dependent factors in the sera affect the core clock or outputs of the core

clock in cultured fibroblasts. In support of this theory, here we show that genes associated with

oxidative phosphorylation and mitochondrial functions lose rhythmicity in the presence of aged

serum factors. We also find that the expression of several molecular clock genes (*PER3,*

NR1D1, NR1D2, CRY1, CRY2 and *TEF*) is phase delayed when synchronized with aged serum.

These novel findings suggest that it may be possible to treat impaired circadian physiology and

the associated disease risks by targeting blood borne factors.

RESULTS

We enrolled 8 old and 7 young human subjects (Fig.1A), whose demographics are in

Table S1. Midline Estimating Statistic of Rhythm: a rhythm-adjusted mean (MESOR) data

obtained from wearable EKG demonstrated that heart rate in old subjects trended lower with

increased variability (Fig. 1B&C). Young subjects displayed a trend of higher activity in the

sympathetic nervous system (SNS), and lower activity in the parasympathetic nervous system

(PNS) (Fig S1A), with no significant difference in cortisol levels (Fig S1B).

To monitor rhythms in culture, *BMAL1-luciferase* transfected BJ-5TA fibroblasts[9] were

synchronized with old/young serum[7,10,11], and circadian effects were assessed by luciferase

assay[9] over 4 days (Fig S2A). No significant differences were observed in the period, amplitude,

and phase of the *BMAL1-luciferase* rhythm with young versus old serum treatment (Fig S2B).

To assess serum effects on circadian gene expression, we first performed RNA-seq

around-the-clock on fibroblasts synchronized with serum from a single old or single young

individual and found that, compared to the first day of synchronization (CT12-34), the second

day (CT 36-58) showed greater differences in MESORs between young and old serum-treated

groups (Fig. S3). This is not surprising because the first day includes acute responses of the

fibroblasts to serum, which can mask circadian rhythms, and so the second day is expected to

reveal differences in endogenous rhythms between samples. Day 2 also revealed different

phases of cyclic expression between young and old groups for a larger number of genes. We

proceeded to collect fibroblasts synchronized by sera from eight different subjects (four old, four

young with two male and two female per group) at two-hour intervals, from 32 to 58 hours post

serum addition. This added an extra two timepoints to the second day to facilitate the calculation

of rhythmicity. Using CircaCompare[12] and a weighted BIC>0.75 cutoff, a significant number of

genes were found to lose (1519 genes) or gain (637 genes) rhythmicity with age, underscoring

the impact of age on the ability of serum to synchronize circadian rhythms, while 1209 genes

were rhythmic in both groups (Fig 2A). Additionally, we used CircaCompare to estimate

MESORs, amplitudes, and phases of gene oscillatory patterns in young and old groups, and to

compare these cosinor parameters between groups (Fig 2B). Of the genes that were rhythmic in

both groups, many also showed changes with age. For instance, 568 cyclically expressed

genes showed a change in MESOR with age (q < 0.05 for MESOR differences), with MESOR

values increasing for 163 genes and decreasing for 405 genes in the old serum-treated

samples. Using p-values provided by CircaCompare we were able to detect changes in

amplitude (Fig 2C) for only a small number of genes (39 genes had decreased amplitude in old

and 2 had increased amplitude). Using CircaCompare and a q < 0.05 cutoff for phase

differences in genes rhythmic in young and old, we detected 20 genes with advanced phase,

and 34 with delayed phase (Fig 2D).

Search Tool for the Retrieval of Interacting Genes (STRING)[13] and Ingenuity Pathway

Analysis (IPA)[14] were used for functional genomics, and both approaches indicate a

maintenance of cycling of cell cycle genes, and a loss of rhythmicity of genes associated with

oxidative phosphorylation in the aged serum (Fig 3 and S4). STRING analysis revealed that the

dominant pathways that are rhythmic in both young and old conditions, cell cycle and DNA

replication demonstrate a decrease in MESOR with old serum. Checkpoint control and

chromosomal replication pathways were expressed cyclically in both young and old conditions;

however, several chromosomal replication pathway genes exhibit decreased MESORs in the

aged sera (Fig S4 D,E, Table S2). MESORs of steroid biosynthesis genes, particularly those,

relating to cholesterol biosynthesis, were also increased in the old sera condition (Fig 3, Table

S3).

As noted, the pathways that lost rhythmicity in the old sera included oxidative

phosphorylation, implicating mitochondrial dysfunction. By STRING analysis, 24 out of the 26

genes associated with oxidative phosphorylation are also among the Alzheimer's Disease

STRING network highlighting the disease relevance of this class of genes that loses cycling in

older individuals. Given that Alzheimer's pathology is closely associated with the accumulation

of oxidative stress[15], it is possible that loss of cycling contributes to oxidative damage. An

additional 31 genes in the Alzheimer's Disease STRING network lose rhythmicity with aged

serum; these include amyloid precursor protein (APP) and apolipoprotein E (APOE). APP is by

definition the precursor of amyloid beta, which accumulates in AD[16]. While APOE plays an

important role in lipid transport, specific variants of this gene are strongly associated with AD

risk as APOE interacts with amyloid beta in amyloid plaques, a hallmark of the disease [17].

Lastly, several clock genes showed differences in expression with the aged serum, most

notably genes in the Circadian Rhythm KEGG pathway (Fig 4). In particular, expression of

CRY1, *CRY2*, *NR1D1*, *NR1D2*, *PER3*, and *TEF* was significantly phase delayed after

synchronization with old serum (Fig 4). Importantly, the RNA-seq did not reveal a difference in

the phase or amplitude of *BMAL1* expression with age, supporting our *BMAL1-luciferase*

findings, although the MESOR significantly increased with age.

DISCUSSION

Although studies of the aged SCN have revealed persistent cycling of clock gene

expression but a breakdown of output signals[4,5], whether age-related changes in systemic

signaling impact organismal clocks has yet to be elucidated. We demonstrate here that while

the core clock continues to cycle in cultured fibroblasts synchronized with serum from old

volunteers, the circadian transcriptome is different from that seen in cells treated with serum

from young individuals. Through this analysis of the role of serum in age-induced changes in

circadian rhythms, we suggest a potential mechanism for other studies, which showed that

specific genes lose, gain, or maintain rhythmicity with age[18–22]. This phenomenon, known as

circadian reprogramming, may illuminate which pathways are affected by or become more

impactful for cellular maintenance through aging in a tissue specific manner[23]. Interestingly,

many of the genes in the circadian transcriptome that exhibit age-related changes are

independently implicated in aging. For instance, some of the genes that lose rhythmicity in the

aged condition are involved in oxidative phosphorylation and mitochondrial function, both of

which decrease with age[24,25]. We suggest that the age-related inefficiency of oxidative

phosphorylation derives at least in part from loss of cycling due to changes in serum signals.

In order to study the effect of circulating factors on age-related changes of the circadian

transcriptome, we utilized the well-established serum starvation-serum addition protocol[7] to

synchronize cells in culture. Fibroblasts from old and young subjects have robust clocks and

respond similarly to synchronization with dexamethasone[10]; however, when synchronized with

dexamethasone in a media containing old serum, they exhibited shortened circadian periods[10].

This effect was reversed by heat inactivating old serum. We did not observe a significant

difference in period length between young and old serum synchronization in our *BMAL1-*

luciferase experiment, perhaps because of the absence of dexamethasone. Our use of serum to

synchronize allowed us to more closely simulate *in vivo* conditions where blood is an important

mode by which central clocks can entrain peripheral clocks to maintain synchrony with the

day:night cycle. Previously, serum factors were shown to activate the immediate early

transcription factor, SRF, in a diurnal manner to confer time of day signals to both cells in

culture and to mouse liver[7]. We show here that effects of age are also mediated by serum. We

find that while the number of rhythmic transcripts in the young serum condition (~10%) is higher

than the old serum condition (~4%), both conditions demonstrate a much larger number of

cycling transcripts in these cultured fibroblasts than in other cell culture studies[26–29]. However, in

mammals, up to 20% of transcripts can cycle in a given tissue[30,31], and the low number of

cycling transcripts in culture has been cited as a major limitation of the culture model[28]. A major

contributing factor to robust in vivo cycling might be human serum as the synchronization signal.

In this way, our model of mimicking signaling to peripheral tissues by using serum directly from

humans may more accurately recapitulate the human condition.

Together, these findings indicate that at least some of the age-related changes in the

circadian clock and circadian transcriptome are derived from signals circulating in the serum

and not the age of the tissue. This has profound implications for understanding and treating

circadian disruption with age, and could also be relevant for other age-related pathology, given

established links between circadian disruption and diseases of aging. Notably, many of the

genes whose cycling is affected by old serum contribute to age-associated disorders.

FIGURES

Fig 1. Healthy elderly individuals tend to have lower heart rate, increased heart rate

variability, and phase advanced activity patterns relative to young subjects.

Experimental protocol for enrolment and monitoring of human subjects **(A)**. As assessed by

Zyphyr BioPatch, EKG measurements suggest that MESOR of average heart rate decreases

with age (p=0.056) and that MESOR of heart rate variability increases with age (p=0.056). N=5

per group. MESOR and amplitude were tested using two-sided Wilcoxon rank sum exact test,

while phase was tested by Kuiper's two-sample test **(B)**. Average activity counts across three

axes (vector magnitude), as recorded by the actigraph device plotted throughout the day (top)

and analyzed for circadian rhythm (bottom) **(C)**. While the amplitudes of activity did not differ

between the age groups, the older individuals trended towards an early phase (p~0.055,

Kupier's two-sample test) compared to young individuals. N=7 for young and N=8 for old. Lines

in the top panel of C are smoothed means (fit with penalized cubic regression splines) for data

from each age group. Dots in the bottom panel of C are subject-level cosinor parameter

estimates derived from cosinor fits to the actigaraphy data. Summary Figure (A) was made with

BioRender.com. Boxplot midlines correspond to median values, while the lower and upper

hinges correspond to the first and their quartiles, respectively. Boxplot whiskers extend to the

smallest/largest points within 1.5 * IQR (Inter Quartile Range) of the lower/upper hinge.

Fig 2. The circadian transcriptome is differentially affected by entrainment with sera from young and old human subjects.

A visual representation of the serum starvation-serum addition protocol to synchronize the *BMAL1-luciferase* BJ-5TA fibroblasts **(A**, made with BioRender.com**)**. When comparing the circadian transcriptome entrained by either young or old sera (n=4 sera per group), 1519 genes lost rhythmicity with age (B, top) while only 637 genes gained rhythmicity with age (B, bottom). Weighted BIC criterion with a threshold of 0.75 was used to assess rhythmicity **(B)**. The number of genes rhythmic in young and old (according to weighted BIC > 0.75 criterion) that show a detectable change in MESOR (q < 0.05 criterion for MESOR difference) is 568, with MESOR increasing in 163 and decreasing in 405 genes. Out of these, 40 genes with increased MESOR (red) and 98 genes with decreased MESOR (blue) also satisfy the condition $|\log_2 FC| > 0.25$ **(C)**. We were only able to detect change in amplitude for a small number of genes, 39 genes had decreased amplitude in old and 2 had increased amplitude using CircaCompare. Only 30 genes with decreased amplitude also satisfy the condition $|\log_2 FC| > 0.1$ (blue) **(D)**. For phase, using a test provided by CircaCompare, under q < 0.05 cutoff for age-related phase differences in genes rhythmic in young and old (with BIC>0.75 cycling criteria), we detected 20 genes with advanced phase, and 34 with delayed phase **(E)**.

Fig 3. Each type of circadian change is associated with different KEGG pathways by

STRING analysis

Entrainment of the fibroblasts in culture with old serum significantly altered the circadian

transcriptome despite use of the same cells in both young and old conditions, suggesting the

aged serum itself affects the regulation of specific pathways in the cell. Significant KEGG

Pathways (FDR $<1\times10^{-4}$ unless specified) are indicated next to each category of genes. Age

related pathways such as Alzheimer's Disease/oxidative phosphorylation are associated with a

loss of rhythms in the old condition. Cell cycle and DNA replication pathways remain rhythmic in

the old serum condition, but cycle with a decreased MESOR. Rhythmic genes were determined

using CircaCompare. Rhythmicity was determined using the weighted BIC > 0.75 criterion while

p-values for difference in MESOR and phase were determined using CircaCompare.

230
Fig 4. Synchronizing with old serum phase delays the expression profile of several core

clock genes

Traces of molecular clock mRNA transcripts, the average curve (bold) of results of individual

sera (faded). P-values for the difference in MESOR (P_**Δ**M), amplitude (P_**Δ**A), and phase

(P_**Δ**P) are shown for the comparison of young and old conditions. Several clock genes are

significantly phase delayed (*CRY1, CRY2, NR1D1, NR1D2, PER3, TEF*) in response to

synchronization with old serum. While *BMAL1* is not phase delayed, the MESOR significantly

increases with age. N=4 subjects per timepoint for both young and old groups.

MATERIALS AND METHODS

Clinical research study:

This clinical research study enrolled apparent healthy participants from the volunteer pool maintained by the Institute for Translational Medicine and Therapeutics (ITMAT), University of Pennsylvania. The Institutional Review Board of the University of Pennsylvania (Federal wide Assurance FWA00004028; IRB Registration: IORG0000029) approved the clinical study protocol (Penn IRB#832866). The study was registered on ClinicalTrials.gov with identifier NCT04086589. After obtaining informed consent from all volunteers, study assessments were conducted in the Center for Human Phenomic Science (Penn CHPS#3002) in accordance with relevant GCP guidelines and regulations. We originally intended on recruiting n=20 per group; however, patient recruitment was halted due to the COVID-19 pandemic. Participants met criteria for inclusion (in general good health, either 70-85 years of age for the elderly cohort or 20-35 years of age for the young cohort, and a wrist-actigraphy-based average TST (total sleep time) ≥ 6 hours per night occurring between 22:00 – 08:00). Participants were excluded due to pregnancy or nursing, shift work (defined as recurring work between 22:00-05:00), a history of clinically significant obstructive sleep apnea, transmeridian travel across ≥2 time zones in the two weeks prior to study assessments and one week after, > 2 drinks of alcohol per day, and use of illicit drugs. Blood collections were done from the median cubital vein via venipuncture using a 22 G butterfly needle (BD, Franklin, Lakes, NJ, USA). All blood draws occurred at 14:00, the same time as Pagani et al.[32]. One subject returned for a repeat clinical assessment including biosampling to provide additional sample.

Acquisition of accelerometry data streams:

Participants wore a triaxial actigraph device (wGT3X-BT, ActiGraph, Pensacola, FL) on the non-dominant wrist. The devices were initialized using the following parameters: start date and time were synchronized with atomic server time without pre-defined stop date/time, at 60

Hz sampling rate for the three accelerometer axes, enabled for delay modus, steps, lux,

inclinometer, and sleep while active. Raw data were downloaded from the device in AGD and

GT3X file format in one second epochs using ActiLife software (version 6, ActiGraph,

Pensacola, FL) and submitted for further analyses. For visualization and cosinor analysis,

actigraphy data were aggregated into 1-minute intervals by summing ActiGraph counts across

each minute. Cosinor analyses of the data were adapted from the single component cosinor

analysis reviewed by Cornelissen[33], as well as the cosine fit described by Refinetti et al. [34].

Briefly, the measurement times for the actigraphy data were recalculated as hours since

midnight on the first day of measurement, within each participant's data. For each actigraphy

variable and each participant, the lm() function in R (v4.2.0) was used to perform a cosinor fit

with a fixed 24-hour period. The two cosinor coefficients from these fits were used to calculate

participant-level amplitudes and circadian phases, while the intercepts provided MESOR

estimates. The two-sided Wilcoxon rank sum exact test, as implemented by R's wilcox.test()

function, was used to test for significant differences in amplitude and MESOR between the age

groups. A two-sample Kuiper's test, as implemented by the kuiper_test(nboots = 10000)

function from the twosamples R package (v2.0.0), was used to test for significant differences in

circadian phase between the age groups.

Acquisition of EKG data:

The Zephyr BioPatch devices (Zephyr Technology, Annapolis, MD) were deployed as

previously established[35]. All subjects included in this analysis wore the BioPatch for at least

24hrs. EKG recordings were analyzed by Kubios HRV Premium (ver. 3.5.0, Kubios Team,

Kuopio, Finland) to report time-of-day-specific measures of cardiovascular function consisting of

heart rate, RR intervals, sympathetic and parasympathetic nervous activity (SNS and PNS).

Preprocessing in Kubios was set to automatic beat correction to remove artifacts and Smoothn

priors for detrending. Cosinor fits and tests for differences in circadian parameters between age

groups were performed as described for the actigraphy data. The circadian parameters MESOR

and amplitude were tested for significant differences between the age groups by Wilcoxon rank

sum exact test (two-sided), while phase was tested by Kuiper's two-sample test to account for

the circular measurement.

Cortisol Measurements:

Cortisol was measured in human serum by coated tube RIA (MP Bio, Solon OH) in

duplicate. Tubes were counted on Perkin Elmer gamma counter and data reduced by STATLia

software.

Generation of stable BJ-5TA cell line expressing *BMAL1-dLuc-GFP*[9]:

Virus was generated and cells were infected as we've previously described[36]. Briefly,

LentiX 293T cells (Clonetech) were transfected with Lipofectamine 3000 PLUS (Life Tech) using

manufacturer instructions. The transfection included 18ug of DNA per reaction and the plasmid

(BMAL1-dLuc-GFP) to packaging vector (DVPR, Addgene) to envelope (VSV-G, Addgene) ratio

was 10:1:0.5. Media was changed 24hrs post-transfection after the cells were checked using a

fluorescence scope to make sure cells were >50% GFP positive. Supernatant with the virus was

collected at 48hrs and 72hrs post transfection and spun down at 3000 RPM for five minutes (to

eliminate any cells/debris). BJ-5TA cells were infected with fresh virus upon virus collection.

Polybrene (Sigma-Aldrich, 10 mg/mL) was also added to aid with infection. Transduced BJ-5TA

cells (>2000 cells) were sorted (FACSMelody, BD Biosciences) for high GFP expression. Once

the cell line was established, blasticidin was added to the culture at 2ug/ml. Due to fragility of

the cell line in the presence of antibiotic, BJ-5TA *Bmal1-luciferase* cells with differing expression

levels of GFP were sorted. The line with highest stable luminescence oscillation was used for all

experiments reported here.

Serum entrainment and bioluminescent recording:

BJ-5TA *BMAL1-dLuc-GFP* cells in 24-well plates (~confluent) were washed (2x) with

DPBS and given serum free media for 24hrs. After the starvation, cells were given media with

10% human serum and 200uM beetle luciferin potassium salt. Each well of cells was given

media with the serum from a single patient. The cells were continuously monitored by

LumiCycle luminometer (Actimetrics) for 4.5 days from the point of serum addition. LumiCycle

raw data was exported using LumiCycle software (Actimentrics). The data were analyzed by

BioDare2 (biodare2.ed.ac.uk[37]) using FFT NLLS with baseline detrending. Any replicates that

were not cycling or had a period outside of the 20-28hr range was excluded from analysis. Both

the serum free media and serum added media used the recipe from[9] at pH 7.4; however, the

serum free media had no serum and the serum media had 10% human serum instead of FBS.

Sample preparation, RNA extraction, and RNA sequencing:

BJ-5TA *BMAL1-dLuc-GFP* cells in 24-well plates (~confluent) were washed (2x) with

DPBS and given serum free media (DMEM with Penn Strep) for 24hrs. After 24hrs cells were

given media with human serum (DMEM, Penn Strep, 10% human serum). Serum starvation was

staggered every 12 hours over two days to allow for samples to be collected on the same day.

Upon sample collection wells were place on ice and rinsed with cold DPBS and then put in cold

RLT buffer with 2-Mercaptoethanol (10uL/mL). Samples were frozen at -80 overnight and sent

to Admera for RNA extraction and sequencing. Total RNA was extracted with RNeay mini kit

(Qiagen). Isolated RNA sample quality was assessed by High Sensitivity RNA Tapestation

(Agilent Technologies Inc., California, USA) and quantified by Qubit 2.0 RNA HS assay

(ThermoFisher, Massachusetts, USA). Paramagnetic beads coupled with oligo d(T)25 are

combined with total RNA to isolate poly(A)+ transcripts based on NEBNext® Poly(A) mRNA

Magnetic Isolation Module manual (New England BioLabs Inc., Massachusetts, USA). Prior to

first strand synthesis, samples are randomly primed (5´ d(N6) 3´ [N=A,C,G,T]) and fragmented

based on manufacturer's recommendations. The first strand is synthesized with the Protoscript

II Reverse Transcriptase with a longer extension period, approximately 40 minutes at 42°C. All

remaining steps for library construction were used according to the NEBNext® UltraTM II Non-

Directional RNA Library Prep Kit for Illumina® (New England BioLabs Inc., Massachusetts,

USA). Final libraries quantity was assessed by Qubit 2.0 (ThermoFisher, Massachusetts, USA)

and quality was assessed by TapeStation D1000 ScreenTape (Agilent Technologies Inc.,

California, USA). Final library size was about 430bp with an insert size of about 300bp.

Illumina® 8-nt dual-indices were used. Equimolar pooling of libraries was performed based on

QC values and sequenced on an Illumina® NovaSeq S4 Illumina, California, USA) with a read

length configuration of 150 PE for 40 M PE reads per sample (20 M in each direction).

RNA-seq and statistical analysis

Raw RNA-seq reads were aligned to the GRCh38 build of the human genome by STAR

version 2.7.10a[38]. The dataset contained an average of 19,265,220 paired-end non-stranded

150 bp reads mapping uniquely to genes, per sample. Data were normalized and quantified at

both gene and exon-intron level, using a downsampling strategy implemented in PORT

(Pipeline Of RNA-seq Transformations, available at https://github.com/itmat/Normalization),

version 0.8.5f-beta_hotfix1. Both STAR and PORT were provided with gene models from

release 106 of the Ensembl annotation[39].

MESOR, amplitude, and phase estimates, as well as p-values for the difference in

MESOR, amplitude, and phase, were calculated with CircaCompare[12], version 0.1.1. Only

rhythmic genes were taken into consideration in pathway and other analyses involving MESORs

and phases. The criterion for rhythmicity was either based on (BH adjusted) p-values reported

by CircaCompare, or weighted BIC values, obtained by an approach similar to dryR[40]. In the

latter approach we fitted four models of rhythmicity, one modeling gene expression that is

rhythmic in both cells treated with young or old sera, another modeling gene

expression rhythmic in neither cells treated with young nor old sera, and two modeling gene

expression rhythmic in either young or old sera treated cells respectively. The BIC values were

calculated for the four models and were weighted to obtain numbers between 0 and 1. All

methods were provided with $\log_{10}(1 + \text{PORT normalized count})$ values and were run on R,

version 4.1.2, accessed through Python, version 3.9.9, via rpy2, an interface to R running

embedded in a Python process, https://rpy2.github.io/, version 3.4.5. Additionally, we used

Nitecap[41] to visualize and explore circadian profiles of gene expression.

STRING pathway analysis

We performed STRING[13] pathway analyses using STRING API version 11.5.

Enrichment analyses were performed on the sets of genes rhythmic in both groups (weighted

BIC > 0.75) with observed decrease in MESOR, increase in MESOR, advance in phase, and

delay in phase according to CircaCompare q < 0.05 criterion. We also performed analyses on

the sets consisting of genes rhythmic only in young group (weighted BIC > 0.75), only in old

group (weighted BIC > 0.75), and the set of genes rhythmic in both groups (weighted BIC >

0.75). Finally, two additional analyses were performed, on the sets of genes with observed

decrease and increase of amplitude (CircaCompare q < 0.05).

Ingenuity Pathway Analysis:

QIAGEN IPA[14] was used to identify pathways enriched in various subsets of cycling

genes. The following subsets of genes were identified using a combination of CircaCompare

stats and BIC cutoffs: (1) Decreased MESOR in old sera (929 genes) – CircaCompare rhythmic

q < 0.05 in old, CircaCompare rhythmic q < 0.05 in young, CircaCompare MESOR difference q

< 0.05, MESOR difference (old – young) < 0. (2) Increased MESOR in old sera (515 genes) –

same selection criteria as 'Decreased MESOR in old sera,' except MESOR difference (old –

young) > 0. (3) Phase advance in old sera (148 genes) - CircaCompare rhythmic q < 0.05 in old,

CircaCompare rhythmic q < 0.05 in young, CircaCompare Phase difference q < 0.05, phase

difference (old – young) < 0. (4) Phase delay in old sera (156 genes) – same selection criteria

as 'Phase advance in old sera,' except phase difference (old – young) > 0. (5) Loss of

rhythmicity in old sera (1519 genes) – weighted BIC > 0.75 for 'Rhythmic in Young but not Old'

model. (6) Gain of rhythmicity in old sera (637 genes) – weighted BIC > 0.75 for 'Rhythmic in

Old but not Young' model. (7) Rhythmic in old and young sera (1209 genes) – weighted BIC >

0.75 for 'Rhythmic in both Old and Young' model. Each of these gene subsets were processed

separately with IPA's core analysis, using default parameters.

For visualization via heatmap, we perform three rounds of normalization within each

gene. First, we mean-normalize the read counts within each age group and serum treatment

group (A, B, C, D). This is to account for baseline differences between sera collected from the

different subjects. Second, we collapse replicates at each timepoint by calculating their means.

Third, we calculate Z-Scores across all timepoints, within each age group. Note, these

normalization procedures are to aid with visualization of the data and were not used as part of

the statistical analyses.

Data and materials availability:

Sequencing data are deposited in Gene Expression Omnibus (NCBI) under accession

number TBD. All additional data files are available upon request.

Acknowledgments:

We wish to extend our sincere gratitude to the study volunteers. Ms. LaVenia Banas

provided excellent logistical study support. We thank Dr. Andrew Liu for the BMAL1-luc plasmid.

We thank the RIA Biomarker Core of the Penn Diabetes Research Center, P30-DK19525 for

cortisol measurements. We thank Sara Bernardez-Noya and Rebecca Moore for input on analysis and data presentation. The project described was supported by the National Center for Research Resources and the National Center for Advancing Translational Sciences, National Institutes of Health, through Grant 5UL1TR001878 (A.S.) and National Heart, Blood, and Lung Institutes of Health, through Grant R00HL1472 (S.L.Z.). C.S. is the Robert L. McNeil Jr. Fellow in Translational Medicine and Therapeutics. A.S. is an investigator of the Howard Hughes Medical Institute. J.E.S. was supported by a training grant in Neuroscience (NIH T32-NS105607), an National Institutes of Health Diversity Supplement (NIH NS48471), and by a grant to the University of Pennsylvania from the Howard Hughes Medical Institute through the James H. Gilliam Fellowship for Advanced Study program. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

References

1. Monk, T. H., Buysse, D. J., Reynolds, C. F. & Kupfer, D. J. Inducing jet lag in older people: Adjusting to a 6-hour phase advance in routine. *Experimental Gerontology* **28**, 119–133 (1993).

2. Monk, T. H., Buysse, D. J., Carrier, J. & Kupfer, D. J. Inducing jet-lag in older people: Directional asymmetry. *Journal of Sleep Research* **9**, 101–116 (2000).

3. Hood, S. & Amir, S. The aging clock: circadian rhythms and later life. *Journal of Clinical Investigation* **127**, 437–446 (2017).

4. Farajnia, S. *et al.* Evidence for Neuronal Desynchrony in the Aged Suprachiasmatic Nucleus Clock. *J. Neurosci.* **32**, 5891–5899 (2012).

5. Nakamura, T. J. *et al.* Age-Related Decline in Circadian Output. *J Neurosci* **31**, 10201–10205 (2011).

6. Yamazaki, S. *et al.* Effects of aging on central and peripheral mammalian clocks. *Proc Natl Acad Sci U S A* **99**, 10801–10806 (2002).

7. Gerber, A. *et al.* Blood-Borne Circadian Signal Stimulates Daily Oscillations in Actin

Dynamics and SRF Activity. *Cell* **152**, 492–503 (2013).

8. Katsimpardi, L. *et al.* Vascular and Neurogenic Rejuvenation of the Aging Mouse Brain by

Young Systemic Factors. *Science* **344**, 630–634 (2014).

9. Ramanathan, C., Khan, S. K., Kathale, N. D., Xu, H. & Liu, A. C. Monitoring Cell-

autonomous Circadian Clock Rhythms of Gene Expression Using Luciferase

Bioluminescence Reporters. *JoVE (Journal of Visualized Experiments)* e4234 (2012)

doi:10.3791/4234.

10. Pagani, L. *et al.* Serum factors in older individuals change cellular clock properties.

Proceedings of the National Academy of Sciences of the United States of America (2011)

doi:10.1073/pnas.1008882108.

11. Balsalobre, A., Damiola, F. & Schibler, U. A Serum Shock Induces Circadian Gene

Expression in Mammalian Tissue Culture Cells. *Cell* **93**, 929–937 (1998).

12. Parsons, R., Parsons, R., Garner, N., Oster, H. & Rawashdeh, O. CircaCompare: a method

to estimate and statistically support differences in mesor, amplitude and phase, between

circadian rhythms. *Bioinformatics* **36**, 1208–1212 (2020).

13. Szklarczyk, D. *et al.* STRING v11: protein-protein association networks with increased

coverage, supporting functional discovery in genome-wide experimental datasets. *Nucleic*

Acids Res **47**, D607–D613 (2019).

14. Krämer, A., Green, J., Pollard, J. & Tugendreich, S. Causal analysis approaches in

Ingenuity Pathway Analysis. *Bioinformatics* **30**, 523–530 (2014).

15. Holubiec, M. I., Gellert, M. & Hanschmann, E. M. Redox signaling and metabolism in

Alzheimer's disease. *Front Aging Neurosci* **14**, 1003721 (2022).

16. Sehar, U., Rawat, P., Reddy, A. P., Kopel, J. & Reddy, P. H. Amyloid Beta in Aging and

Alzheimer's Disease. *Int J Mol Sci* **23**, 12924 (2022).

17. Raulin, A.-C. *et al.* ApoE in Alzheimer's disease: pathophysiology and therapeutic

strategies. *Mol Neurodegener* **17**, 72 (2022).

18. Chen, C.-Y. *et al.* Effects of aging on circadian patterns of gene expression in the human

prefrontal cortex. *Proc Natl Acad Sci U S A* **113**, 206–211 (2016).

19. Kuintzle, R. C. *et al.* Circadian deep sequencing reveals stress-response genes that adopt

robust rhythmic expression during aging. *Nat Commun* **8**, 14529 (2017).

20. Sato, S. *et al.* Circadian Reprogramming in the Liver Identifies Metabolic Pathways of Aging.

Cell **170**, 664-677.e11 (2017).

21. Solanas, G. *et al.* Aged Stem Cells Reprogram Their Daily Rhythmic Functions to Adapt to

Stress. *Cell* **170**, 678-692.e20 (2017).

22. Blacher, E. *et al.* Aging disrupts circadian gene regulation and function in macrophages. *Nat

Immunol* **23**, 229–236 (2022).

23. Wolff, C. A. *et al.* Defining the age-dependent and tissue-specific circadian transcriptome in

male mice. *Cell Reports* **42**, 111982 (2023).

24. Lesnefsky, E. J. & Hoppel, C. L. Oxidative phosphorylation and aging. *Ageing Research

Reviews* **5**, 402–433 (2006).

25. Greco, M. *et al.* Marked aging-related decline in efficiency of oxidative phosphorylation in

human skin fibroblasts. *The FASEB Journal* **17**, 1706–1708 (2003).

26. Duffield, G. E. *et al.* Circadian programs of transcriptional activation, signaling, and protein

turnover revealed by microarray analysis of mammalian cells. *Curr Biol* **12**, 551–557 (2002).

27. Grundschober, C. *et al.* Circadian regulation of diverse gene products revealed by mRNA

expression profiling of synchronized fibroblasts. *J Biol Chem* **276**, 46751–46758 (2001).

28. Hughes, M. E. *et al.* Harmonics of Circadian Gene Transcription in Mammals. *PLOS

Genetics* **5**, e1000442 (2009).

29. Jang, C., Lahens, N. F., Hogenesch, J. B. & Sehgal, A. Ribosome profiling reveals an

important role for translational control in circadian gene expression. *Genome Res* **25**, 1836–

1847 (2015).

30. Patel, V. R., Eckel-Mahan, K., Sassone-Corsi, P. & Baldi, P. CircadiOmics: integrating

circadian genomics, transcriptomics, proteomics and metabolomics. *Nat Methods* **9**, 772–

773 (2012).

31. Zhang, R., Lahens, N. F., Ballance, H. I., Hughes, M. E. & Hogenesch, J. B. A circadian

gene expression atlas in mammals: Implications for biology and medicine. *Proc Natl Acad*

Sci U S A **111**, 16219–16224 (2014).

32. Pagani, L. *et al.* Serum factors in older individuals change cellular clock properties.

Proceedings of the National Academy of Sciences **108**, 7218–7223 (2011).

33. Cornelissen, G. Cosinor-based rhythmometry. *Theor Biol Med Model* **11**, 16 (2014).

34. Refinetti, R., Lissen, G. C. & Halberg, F. Procedures for numerical analysis of circadian

rhythms. *Biol Rhythm Res* **38**, 275–325 (2007).

35. Lahens, N. F. *et al.* Time-specific associations of wearable sensor-based cardiovascular

and behavioral readouts with disease phenotypes in the outpatient setting of the Chronic

Renal Insufficiency Cohort. *Digit Health* **8**, 20552076221107904 (2022).

36. Zhang, S. L. *et al.* A circadian clock regulates efflux by the blood-brain barrier in mice and

human cells. *Nat Commun* **12**, 617 (2021).

37. Zielinski, T., Moore, A. M., Troup, E., Halliday, K. J. & Millar, A. J. Strengths and Limitations

of Period Estimation Methods for Circadian Data. *PLOS ONE* **9**, e96462 (2014).

38. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

39. Yates, A. D. *et al.* Ensembl 2020. *Nucleic Acids Research* **48**, D682–D688 (2020).

40. Weger, B. D. *et al.* Systematic analysis of differential rhythmic liver gene expression

mediated by the circadian clock and feeding rhythms. *Proceedings of the National Academy*

of Sciences **118**, e2015803118 (2021).

41. Brooks, T. G. *et al.* Nitecap: An Exploratory Circadian Analysis Web Application. *J Biol*

Rhythms **37**, 43–52 (2022).

516
517

518
Fig S1. There are no significant differences in grip strength, sympathetic,

parapsympathetic nervous system indices, or cortisol levels between young and old

individuals

Sympathetic nervous system (SNS, A, top) and parasympathetic nervous system (PNS, A,

bottom) indices were measured using EKG. Amplitude, MESOR, and Phase were calculated

from the resulting oscillations and the Median, quartiles, and SEMs are shown. Dots represent

individuals. P>0.05. N=5 per group. MESOR and amplitude were tested using Wilcoxon rank

sum exact test, while phase was tested by Kuiper's two-sample test. Boxplot midlines

correspond to median values, while the lower and upper hinges correspond to the first and third

quartiles, respectively. Boxplot whiskers extend to the smallest/largest points within 1.5 * IQR

(inter Quartile Range) of the lower/upper hinge **(A)**. Serum cortisol levels are not statistically

different by unpaired t-test **(B**, N=7 young, 8 old **)**.

Table S1. Demographic information for subjects in the study.
533

Age Group	Sex	Age (y)	BMI (kg/m^2)	Total Sleep Time (TST) [h]
	Male	30	25.9	6.4
	Female	30	28.7	7.4
	Female	25	23.7	6.6
20-35	Male	28	25.0	9.7
	Female	28	21.2	6.9
	Male	28	25.2	7.4
	Male	28	25.9	9.3
Young	*Mean*	*28.1*	*25.1*	*7.7*
	SD	*1.7*	*2.3*	*1.3*
	Male	76	22.3	5.8
	Male	75	28.8	7.8
	Female	73	32.4	7.7
70-85	Female	74	26.4	7.5
	Male	73	21.5	6.6
	Male	71	26.7	11.3
	Male	74	27.2	7.0
	Male	70	26.8	6.7
Old	*Mean*	*73.3*	*26.5*	*7.6*
	SD	*2.0*	*3.4*	*1.7*

534 Total sleep time (TST) was calculated from ActiLife 6 determined in-bed and out-of-bed times
535 averaged across nights with available actigraphy data (≥7 nights). Old (N=8), Y-young (N=7).

Fig S2: Young and old serum contain equivalent levels of cortisol and are equally effective at entraining cells in culture

(A) Cells synchronized with human serum from either young (N=7 subjects) or old (N=8 subjects) individuals did not show differences in amplitude (as measured in luminescence counts per second, bottom, left), period (bottom, middle), or phase (bottom, right). Data points represent average values for a specific patient's serum over 2 experiments with 2-3 replicates per experiment. A visual representation of the individual replicates averaged in A **(B)** Summary statistics are displayed as mean +/- SEM. Means compared by unpaired t-test. Despite individual and run to run variability, cells synchronized with human serum from either young (N=7 subjects) or old (N=8 subjects) individuals showed similar amplitude (top, left), period (top, right), and phase (bottom) of BMAL1-luciferase rhythms. Data points represent individual replicates within an experiment. Wells run in the same experiment are displayed in the same color. Each subject's sample was run in 2-3 replicates.

Fig S3. The circadian transcriptomes of cells synchronized with young or old sera deviate significantly on day two.

On day two of serum entrainment, young and old transcriptional rhythms differentiated.

CircaCompare analysis of RNA sequencing revealed that on Day 2 of serum entrainment

analysis (CT 36-58) the MESOR differences of cycling genes are larger between the cells

entrained with young or old serum **(A**, top**)**. Additionally, more genes were phase shifted in the

old serum condition compared to young serum on Day 2 **(B**, top**)**. The distribution of p-values

shows an enrichment of low p-values on Day 2 for both MESOR and phase differences (A,

bottom, B, bottom). The size of each circle in the scatter plots is proportional to $-\log_{10}q$, so the

bigger the circle, the smaller the q-value for the difference between young and old for each

metric.

Fig S4: Oxidative phosphorylation genes and chromosomal replication genes are

affected by entrainment with old serum

Additional pathway analysis employing IPA further emphasizes finding with STRING analysis.

Genes identified by IPA analysis to be associated with oxidative phosphorylation/mitochondrial

dysfunction and eIF2 genes lose rhythmicity with age (BIC >0.75) **(A, B, C)**. Cell cycle

checkpoint control (CHK proteins) **(D)** and chromosomal replication genes **(E)** maintain their

cycling in the aged condition, which is consistent with the continued division of cells.

A

Cycling genes phase-advanced in old samples
eIF2 Signaling Pathway

B

Cycling genes phase-advanced in old samples
eIF2 Signaling Pathway

574
575
576
577

578 **Fig S5: eIF2 signaling pathway genes that maintain cycling with age phase advance in**
579 **the old serum condition**
580
From the IPA analyses, the eIF2 signaling pathway was enriched (BIC >0.75) in two datasets:

1) genes cycling in the young sera and not the old and 2) cycling genes phase-advanced in the

old sera (q<0.05). A significant number of genes that maintain rhythmicity in the old sera

condition are phase advanced as visualized in the heat map **(A)** or plot of peak phase **(B)**. As

the eIF2 pathway is involved in protein translation, this suggests that the timing of protein

translation is phase advanced in the older serum and/or less rhythmic.

Gene	Symbol	Old_mesor_estimate	Young_mesor_estimate
ENSG00000014138	POLA2	1.942220257	2.030676546
ENSG00000062822	POLD1	2.727523056	2.828026817
ENSG00000073111	MCM2	3.003984606	3.115331295
ENSG00000076003	MCM6	3.124116273	3.203138512
ENSG00000085840	ORC1	2.177440998	2.305154417
ENSG00000091651	ORC6	2.415062674	2.513336383
ENSG00000093009	CDC45	2.428834807	2.540049538
ENSG00000094804	CDC6	2.566892644	2.679401971
ENSG00000097046	CDC7	2.279003902	2.377425914
ENSG00000100297	MCM5	3.17577594	3.288560637
ENSG00000101868	POLA1	2.640718445	2.728951631
ENSG00000104738	MCM4	3.310670684	3.413600036
ENSG00000112118	MCM3	3.168296215	3.269321473
ENSG00000117748	RPA2	2.754645539	2.798914718
ENSG00000131747	TOP2A	3.722304752	3.798890616
ENSG00000132383	RPA1	3.238584015	3.287556646
ENSG00000132646	PCNA	3.232760966	3.321439626
ENSG00000135446	CDK4	3.143303063	3.172936739
ENSG00000138346	DNA2	2.113453067	2.238476228
ENSG00000146143	PRIM2	2.620994795	2.675363852
ENSG00000164815	ORC5	2.361538047	2.414675015
ENSG00000166508	MCM7	3.381394362	3.482444492
ENSG00000167258	CDK12	3.10241713	3.129181459
ENSG00000167513	CDT1	2.474002764	2.590319547
ENSG00000198056	PRIM1	2.116803844	2.231962095

589 **Table S2: Genes with decreased MESOR in the IPA cell cycle control of chromosomal**

590 **replication pathway**

591 Table of genes that are rhythmic in both conditions with a decreased MESOR with old serum

592 treatment compared to young serum treatment by CircaCompare. All genes are involved in the

593 cell cycle/DNA replication pathway.

Gene	Symbol	Old_mesor_estimate	Young_mesor_estimate
ENSG00000001630	CYP51A1	2.040377336	1.871207095
ENSG00000079459	FDFT1	3.330291993	3.19001773
ENSG00000104549	SQLE	3.116822033	2.950386727
ENSG00000110921	MVK	2.530172998	2.4121956
ENSG00000112972	HMGCS1	2.93111747	2.696086286
ENSG00000113161	HMGCR	3.011982868	2.833316048
ENSG00000116133	DHCR24	3.506294106	3.343492054
ENSG00000120437	ACAT2	2.71791125	2.58109012
ENSG00000132196	HSD17B7	1.940436315	1.782521965
ENSG00000147383	NSDHL	2.540302415	2.468852126
ENSG00000160285	LSS	3.486802994	3.411431601
ENSG00000160752	FDPS	3.143592562	3.081359846
ENSG00000167508	MVD	2.819963909	2.68386468
ENSG00000172893	DHCR7	2.990620862	2.797567922

600 **Table S3: Genes with increased MESOR in the IPA cholesterol biosynthesis pathway**

601 Table of genes that are rhythmic in both conditions with a decreased MESOR with old serum

602 treatment compared to young serum treatment by CircaCompare. All genes are involved in the

603 cholesterol biosynthesis pathway.