AIDS virus-neutralizing antibody induction reciprocal to a PI3K gain-of-function disease

Hiroyuki Yamamoto[*,1,2] and Tetsuro Matano[*,1,3,4]

[1]AIDS Research Center, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashi-Murayama City, Tokyo 208-0011, Japan

[2]Department of Biomedicine, University Hospital Basel, Hebelstrasse 20, Basel 4031, Switzerland

[3]The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku, Tokyo 108-8639, Japan

[4]Joint Research Center for Human Retrovirus Infection, Kumamoto University, 2-2-1 Honjo, Kumamoto 860-0811, Japan

*Correspondence:

Hiroyuki Yamamoto, AIDS Research Center, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashi-Murayama City, Tokyo 208-0011, Japan. Phone: +81-42-848-7154; Fax: +81-42-848-7155. Email: h-yamato@niid.go.jp

Tetsuro Matano, AIDS Research Center, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo 162-8640, Japan. Phone: +81-3-4582-2811; Fax: +81-3-5285-1165. Email: tmatano@niid.go.jp

Running title: SIV neutralization post-PI3K diminution

Abstract

HIV and simian immunodeficiency virus (SIV) infections are known for impaired neutralizing antibody (NAb) responses. While sequential virus-host B cell interaction appears to be basally required for NAb induction, driver molecular signatures predisposing to NAb induction still remain largely unknown. Here we describe SIV-specific NAb induction following a virus-host interplay reciprocal to a congenital human antibody-deficient disease. Screening of seventy neutralization-resistant SIV_{mac239}-infected macaques found nine NAb-inducing animals, with seven selecting for a specific $CD8^+$ T-cell escape mutation in viral *nef* before NAb induction. This mutation reduced aberrant Nef interaction-mediated drive of B-cell maturation-limiting phosphoinositide 3-kinase (PI3K)/mammalian target of rapamycin complex 2 (mTORC2). *In vivo* imaging cytometry depicted preferential Nef perturbation of cognate Envelope-specific B cells, corroborating cognate B-cell maturation post-mutant selection up to NAb induction. Results exemplify a NAb induction pattern extrinsically reciprocal to human PI3K gain-of-function antibody deficiency, shaped through two-front, sequential virus interaction with both wings of adaptive immunity.

Keywords

neutralizing antibody; B cells; $CD8^+$ T cells; PI3 kinase; inborn errors of immunity; HIV/SIV

44   **Graphical Abstract**

Introduction

Virus-specific neutralizing antibody (NAb) responses by B cells are induced by a tight cooperation of adaptive immune cells (Kumar et al., 2010; Shulman et al., 2014; Giltin et al., 2014; Wang et al., 2014) and often play a central role in clearance of acute viral infections (Junt et al., 2007). In contrast, persistence-prone viruses such as human immunodeficiency virus type 1 (HIV-1), simian immunodeficiency virus (SIV) and lymphocytic choriomeningitis virus (LCMV) variously equip themselves with B cell/antibody-inhibitory countermeasures (Moir et al., 2001; Mattapallil et al., 2005; Sommerstein et al., 2015; Sammicheli et al., 2016; Fallet et al., 2016; Mason et al., 2016) and NAb responses are impaired (Levesque et al.,2009; Mikell et al., 2011). These viruses successfully suppress elicitation of potent NAb responses particularly in acute infection (Hunziker et al., 2003; Tomaras et al., 2008) and establish viral persistence, posing considerable challenge for developing protective strategies.

Reciprocal to this detrimental acute-phase NAb absence, we and others have previously described that acute-phase passive NAb infusion can trigger an endogenous T-cell synergism resulting in robust SIV/SHIV (simian/human immunodeficiency virus) replication control (Haigwood et al., 1996; Yamamoto et al., 2007; Ng et al., 2010; Iseda et al., 2016; Nishimura et al., 2017). This shows that virus-specific NAbs not only confer sterile protection but also can evoke T-cell-mediated non-sterile viral control, suggesting the importance of inducing endogenous NAb responses during an optimal time frame as well as humoral-cellular response synergisms. Therefore, identifying mechanisms driving NAb induction against such viruses is an important step to eventually design NAb-based HIV control strategies.

As an approach, analysis of *in vivo* immunological events linked with NAb induction against difficult-to-neutralize SIVs in a non-human primate model can provide important insights into this goal. HIV-specific NAb induction patterns and correlates, such as antibody-NAb coevolution (Moore et al., 2012), autoimmune-driven induction (Moody et al., 2016) and natural killer cell-related host polymorphisms (Bradley et al., 2018) have been variously reported to date; the wealth of knowledge on this topic collectively, and interestingly, rather

suggests that pathways to NAb induction against neutralization-resistant viruses including

HIV/SIV can be highly redundant and there may be as-yet-unknown *in vivo* mechanisms that

can robustly drive NAb induction. For example, the neutralization resistance of certain SIV

strains apparently is not explanatory by the aforementioned, posing such models as attractive

analytic targets of NAb induction mechanisms.

In the present study, we examined virus-specific antibody responses in rhesus

macaques infected with a highly neutralization-resistant SIV strain, SIV_{mac239}. This virus is

highly pathogenic, shows persistent viremia and hosts lack NAb responses throughout

infection (Kestler et al., 1991). Yet, upon performing a large-scale screening of SIV_{mac239}-

infected rhesus macaques for up to 100 weeks, we noticed a subgroup inducing NAbs in the

chronic phase. Interestingly, these animals commonly selected for a specific $CD8^+$ cytotoxic T

lymphocyte (CTL) escape mutation in the viral Nef-coding gene before NAb induction. The

mutant Nef, compared with wild-type (WT) Nef, manifested a decrease in aberrant interaction

with phosphoinositide 3-kinase (PI3K)/mammalian target of rapamycin complex 2 (mTORC2),

resulting in decreased downstream hyperactivation of the canonical B-cell negative regulator

Akt (Omori et al., 2006; Limon et al., 2014). Nef was graphically identified via machine

learning-comprising imaging cytometry to preferentially target Env-specific B cells *in vivo*, and

in those macaques inducing NAbs, the NAb induction was linked with sustained Env-specific

B-cell responses after the mutant Nef selection. Results describe a B-cell functional boosting

reciprocal to human PI3K gain-of-function antibody deficiency disease, and suggest that

intervening PI3K/mTORC2 signaling can potentially result in harnessing NAb induction against

difficult-to-neutralize viruses.

     Results

Identification of macaques inducing SIV$_{mac239}$-neutralizing antibodies

We performed a retrospective antibody profile screening in rhesus macaques infected

with NAb-resistant SIV$_{mac239}$ (n = 70) (**Figure 1A**) and identified a group of animals inducing

anti-SIV$_{mac239}$ NAb responses (n = 9), which were subjected to characterization. These NAb

inducers showed persistent viremia with no significant difference in viral loads compared with

a profile-clarified naïve, major histocompatibility complex class I (MHC-I) haplotype-balanced

subgroup of NAb non-inducers (n = 19) representatively utilized for comparison hereafter

(**Figure 1B** and **Figure 1-figure supplement 1**). Plasma SIV$_{mac239}$-NAb titers measured by 10

TCID$_{50}$ SIV$_{mac239}$ virus-killing assay showed an average maximum titer of 1:16, being induced

at an average of 48 weeks post-infection (p.i.) (**Figure 1C**). Two of the nine NAb inducers

showed detectable NAb responses before six months p.i., while the remaining seven induced

NAbs after six months. Anti-SIV$_{mac239}$ neutralizing activity was confirmed in immunoglobulin G

(IgG) purified from plasma of these NAb inducers (**Figure 1-figure supplement 2**). SIV Env-

binding IgGs developed from early infection in NAb inducers as well as NAb non-inducers

(**Figure 1-figure supplement 3**); titers were not higher but rather lower at year 1 p.i. in the

NAb inducers (**Figure 1-figure supplement 3B** and **3C**), consistent with other reports

(Havenar-Daughton et al., 2016). NAb inducers and non-inducers showed similar patterns of

variations in viral *env* sequences (Burns et al., 1993), mainly in variable regions 1, 2 and 4

(V1, V2 and V4) (**Figure 1-figure supplement 4**).

Selection of a viral *nef* mutation (Nef-G63E) precedes chronic-phase SIV$_{mac239}$-specific

NAb induction

Viral nonsynonymous polymorphisms outside *env* were next assessed. Strikingly, we

found selection of a viral genome mutation resulting in G (glycine)-to-E (glutamic acid)

substitution at residue 63 of Nef (Nef-G63E) in seven of the nine NAb inducers (**Figure 2A**).

This 63rd residue is in the unstructured N-terminus of Nef, flanked by conserved α-helices

(**Figure 2B**). This region is conserved in HIV-2/SIV but not in HIV-1, occasionally being deleted

in laboratory HIV-1 strains (Geyer et al., 2001). AlphaFold2-based structure prediction did not

derive palpable change except for low-probability disruption of the alpha helices (data not

shown). The Nef G63E mutation was not observed in two early NAb inducers (with detectable

NAbs before 6 months p.i.) but selected in all of the remaining seven inducing detectable NAb

responses after 6 months. In contrast, this mutation was found only in two of the nineteen NAb

non-inducers, including one rapid progressor (**Figure 1-figure supplement 1**) and selection

was significantly enriched in the NAb inducers (**Figure 2C**). Analysis of Nef-G63E mutation

frequencies in plasma viruses showed that Nef-G63E selection preceded or at least paralleled

NAb induction (**Figure 2D**).

Nef-G63E is a CD8$^+$ T-cell escape mutation

These seven NAb inducers selecting this mutation elicited CD8$^+$ T-cell responses

specific for a 9-mer peptide Nef$_{62-70}$ QW9 (QGQYMNTPW) (**Figure 3A**). This Nef$_{62-70}$ QW9

epitope is restricted by MHC-I molecules Mamu-B*004:01 and Mamu-B*039:01 (Evans et al.,

1999; Sette et al., 2012). Possession of these accounted for six cases of Nef-G63E selection

(**Figure 3A**), and the remaining one animal possessed Mamu-A1*032:02 predicted to bind to

Nef$_{63-70}$ GW8 (NetMHCpan). Ten of nineteen NAb non-inducers also had either of these alleles

(**Figure 1-figure supplement 1**), indicating that NAb induction is not associated with these

MHC-I genotypes (P = 0.25 by Fisher's exact test, data not shown) but with the Nef-G63E

mutation itself (**Figure 2C**). Nef-G63E was confirmed to be an escape mutation from Nef$_{62-70}$-

specific CD8$^+$ T-cell responses (**Figure 3B**). NAb non-inducers possessing these alleles

elicited little or no Nef$_{62-70}$-specific CD8$^+$ T-cell responses (**Figure 3-figure supplement 1A**),

suggesting that *in vivo* selection and fixation of this Nef-G63E SIV was indeed Nef$_{62-70}$-specific

CD8$^+$ T cell-dependent. Replication kinetics of SIV carrying the Nef-G63E mutation was

comparable with wild-type (WT) on a cynomolgus macaque HSC-F CD4$^+$ T-cell line (Akari et

al., 1996), stating that the mutation does not critically impair viral replication (**Figure 3-figure**

supplement 1B), and plasma viral loads were comparable between Nef-G63E mutant-

selecting NAb inducers versus non-inducers (**Figure 3-figure supplement 1C**). These

somewhat contrasted conditional reversion of a more replication-impairing *nef* mutation

related with CD3 downregulation loss in SIV$_{mac239}$ infection (Joas et al., 2020). These results

indicate NAb induction following *in vivo* selection and fixation of the CD8$^+$ T-cell escape *nef*

mutation, Nef-G63E.

G63E mutation reduces aberrant Nef interaction-mediated drive of PI3K/mTORC2

We next focused on the functional phenotype of this highlighted Nef-G63E mutant SIV

in infected cells. An essence of host perturbation by Nef is its wide-spectrum molecular

downregulation, ultimately increasing viral replication (Schindler et al., 2004; Zhang et al.,

2009). To address possible amelioration, we compared downregulation of major targets CD3,

CD4, MHC-I, CXCR4 and BST-2 (Jia et al., 2009) in infected HSC-F cells. Nef-G63E mutation

did not confer notable change compared with wild-type (WT) [*P* = not significant (*n.s.*) for all

molecules, data not shown], implicating other non-canonical changes (**Figure 4A**). Literature

suggested ferritin negative influence on B cells proportionate to HIV-infected macrophage-

derived production (Swingler et al., 2008). Here, plasma ferritin showed no differences (**Figure**

4-figure supplement 1A) and viral loads were comparable between Nef-G63E-selecting NAb

inducers and non-inducers (**Figure 4-figure supplement 1B** and **1C**), arguing against gross

involvement of ferritin as well as other viral replication-changing Nef phenotypes at least in

this model.

Thus, we took interest in immunosignaling-modulating properties of Nef-G63E SIV.

Analysis of Akt, the predominant immune-intrinsic negative brake of B-cell maturation and

AFC/antibody responses *in vivo* (Omori et al., 2006; Limon et al., 2014; Fruman et al., 1999;

Ray et al., 2015; Sander et al., 2015), revealed that its serine 473 phosphorylation, non-

canonically known for Nef-mediated upregulation (Kumar et al., 2016), was significantly lower

on day 3 after Nef-G63E SIV infection at low multiplicity of infection (MOI) compared with WT

(**Figure 4B**), stronger than histogram deviation levels in PI3K gain-of-function mice with full

disease phenotype (Avery et al., 2018). The threonine 308-phosphorylated Akt (pAkt Thr308)

level was more unaffected (**Figure 4-figure supplement 1B**).

We further assessed properties of serine 473-phoyphorylated Akt (pAkt Ser473)

inhibition by Nef-G63E SIV, and found that external PI3K stimuli (serum starvation) (Kennedy

et al., 1997) accelerated phenotype appearance from day 3 to day 1 post-infection (**Figure**

4C, left). A similar trend was obtained by short-term PI3K stimulation with interferon-γ (Nguyen

et al., 2001), interleukin-2 (Marzec et al., 2008) and SIV Env (François & Klotman, 2003)

(**Figure 4-figure supplement 1C**). Env-stimulating high-MOI infection also accelerated

phenotype appearance, with enhanced 50% reduction (**Figure 4C**, right). These showed that

the Akt-inhibitory G63E mutant Nef phenotype is PI3K stimuli-dependent. Transcriptome

analysis signatured decrease in PI3K-pAkt-FoxO1-related genes in mutant SIV infection

(**Figure 4-figure supplement 1D**). As Akt is also a survival mediator this implicates that Nef-

G63E mutation may be Th-cytopathic, decreasing Th dysfunctional expansion in LCMV

(Recher et al., 2004) and HIV/SIV (Gray et al., 2011; Lindqvist et al., 2012; Petrovas et al.,

2012) infections. Here, reduced peripheral $CXCR3^-CXCR5^+PD-1^+$ memory follicular Th (Tfh)

(Locci et al., 2013) was in keeping (**Figure 4-figure supplement 1E** and **1F**).

We further explored molecular traits of this decreased Akt hyperactivation. We reasoned

that comparing endogenous Nef binding patterns would be adequate and analyzed SIV-

infected HSC-F cells with a flow cytometry-based proximity ligation assay (PLA) (Leuchowius

et al., 2009; Avin et al., 2017), by calculating a mean fluorescence intensity (MFI) deviation-

based binding index for each. We found that this Nef-G63E mutation causes significant

decrease in Nef binding to PI3K p85/p110α and downstream mTORC2 components mTOR

(Sarbassov et al., 2005) and GβL/mLST8 (Kim et al., 2003) in the $CD4^{lo}$-SIV Gag $p27^+$ infected

population (**Figure 4D**). Collectively, the Nef-G63E mutation results in decreased aberrant Nef

bridging and signal transduction of PI3K/mTORC2, explaining decreased substrate Akt

Ser473 phosphorylation by mTORC2. Thus, Nef-G63E SIV is a mutant virus decreased in

aberrant interaction/drive of B-cell-inhibitory PI3K/mTORC2 signaling. This molecular

signature was reciprocal to a recently found human inborn error of immunity (IEI), activated

PI3 kinase delta syndrome (APDS) (Angulo et al., 2013; Lucas et al., 2014; Avery et al., 2018).

Targeting of lymph node Env-specific B cells by Nef *in vivo*

Next, we analyzed *in vivo* targeting of virus-specific B cells by Nef in lymph nodes to explore the potential B-cell-intrinsic influence of the G63E-Nef phenotype. Previously suggested influence of soluble Nef itself (Qiao et al., 2006) and/or related host factors (Swingler et al., 2008) may derive generalized influence on B cells; however, SIV antigen-specific binding antibody responses were rather decreased in NAb inducers (**Figure 1-figure supplement 1**), viremia was comparable (**Figure 3-figure supplement 1C**) and ferritin levels were not altered (**Figure 4-figure supplement 1A**), arguing against these. Therefore, we surmised that some targeted Nef intrusion against Env-specific B cells may be occurring and that the decreased aberrant Nef-PI3K/mTORC2 drive may result in their enhanced lymph node maturation.

While reports (Xu et al., 2009) histologically proposed Nef B-cell transfer, quantitative traits, e.g., invasion frequency and influence on virus-specific B cells, have remained unvisited. Intracellular Nef staining is dim (particularly for B cell-acquired Nef), difficult to examine by conventional flow cytometry. Defining staining cutline requires single-cell images, overcoming confounding of high Nef false-staining via pre-permeation rupture and post-permeation confounding as judged by biologically discontinuous staining and/or batch-inflated signals. Collectively, at the expense of spatial information, we judged that sophisticating imaging cytometry would best visualize Nef-mediated B-cell perturbation *in vivo*. We analyzed lymph node B cells with Image Stream X MKII, with high-power (> 10-fold) antigen detection [e.g., molecules of equivalent soluble fluorochrome (MESF) 5 *vs.* MESF 80 in an average flow cytometer for FITC detection] ideal for detection/single-cell verification.

We designed a triple noise cancellation strategy to overcome the above. Firstly, amine reactive dye staining (Perfetto et al., 2006) gated out B cells being Nef-positive due to pre-experimental membrane damage (**Figure 5-figure supplement 1A**, left). Next, we deployed secondary quantitative parameters of Nef signals deriving from each pixel of the images. Definition of a gray-level cooccurrence matrix (GLCM) (Haralick et al., 1973) numerating

adjacent signal deviation as their frequencies on a transpose matrix enables calculation of a variety of feature values summarizing traits of the whole image. A biological assumption of Nef signal continuity in a true staining suggested that the sum of weighted square variance of GLCM, i.e., the {sum of [square of (value – average signal strength) × frequency of each value occurrence]} would separate natural versus artificial Nef signals. This value, Haralick variance, was computed for multiple directions and averaged, deriving Haralick variance mean (**Figure 5-figure supplement 1A**, X axis), proportionate with staining image unnaturalness. Finally, Nef signal intensity threshold (**Figure 5-figure supplement 1A**, Y axis) gated out overtly-stained cells void for true-false verification. These excluded B cells showing non-specific, strong-signal binary-clustered staining pixels for Nef deriving a large summated variance (X axis), and/or cells batch-stained for Nef (Y axis), likely originating from post-experimental membrane damage.

With this approach we acquired images of a Live/Dead$^-$-Nef signal Haralick variance mean^{lo}-Nef signal intensity threshold^{lo}-Nef$^+$CD19$^+$ B-cells, with fine-textured gradation of Nef^{int-lo} staining with continuity from membrane-proximal regions without sporadic staining speckles (**Figure 5A** and **Figure 5-figure supplement 1A**). Images resembled Nef-transferred B cells generated *in vitro* by a SIV-infected CD4+ T-cell coculture (**Figure 5-figure supplement 1B**). A linear discriminant analysis-based machine learning module, scoring segregation of typical void/valid images, showed this gating to provide the highest two-dimensional separation (**Figure 5-figure supplement 1A**). Combined with biotinylated Env/streptavidin staining, this reproducibly obtained images of Env-specific (Ch 12) CD19$^+$ (Ch 11) B cells without membrane ruptures (Ch 08) and showing fine-textured transferred Nef (Ch 02) (**Figure 5B**). Nef invasion upregulated pAkt Ser473 (Ch 03) to a range resembling *in vitro* analysis (**Figure 5C**), stating that Nef-driven aberrant PI3K/mTORC2 signaling does occur in Nef-invaded B cells *in vivo* and the current phenotypic change likely affects B-cell dynamics. Strikingly, lymph node Env-specific B cells showed significantly higher Nef-positive frequencies as compared with batch non-Env-specific B cells (**Figure 5D**). This indicated that infected cell-generated Nef preferentially targets adjacent Env-specific B cells, putatively through infected CD4$^+$ cell-to-B-

cell contact/transfer (Xu et al., 2009; Hashimoto et al., 2016) and that a Nef phenotypic change

likely dictates Env-specific B-cell maturation.

Enhanced Env-specific B-cell responses after PI3K-diminuting mutant selection

Finally, peripheral SIV Env-specific IgG$^+$ B-cell responses comprising plasmablasts

(PBs) and memory B cells (B$_{mem}$) were analyzed to assess *in vivo* B-cell quality in NAb

inducers with Nef-G63E mutant selection. After excluding lineage-specific cells [T/NK/pro-B

cells/monocytes/myeloid dendritic cells (DCs)/plasmacytoid DCs], we defined IgG$^+$ PBs as

showing: high replication (Ki-67$^+$), post-activation (HLA-DR$^+$), transcriptional switching for

terminal differentiation (IRF4^{hi}) and downregulated antigen surface binding (surface [s]Env^{lo}-

cytoplasmic [Cy]Env$^+$) (Nutt et al., 2015; Silveira et al, 2015). Env-specific PBs were defined

as CD3$^-$CD8$^-$CD10$^-$CD14$^-$CD16$^-$CD11c$^-$CD123$^-$CD19$^-$CyIgG$^+$Ki-67$^+$IRF4^{hi}HLA-DR$^+$ (total PB)

sEnv^{lo}CyEnv$^+$ (Env-specific) (**Figure 6-figure supplement 1**). NAb inducers showed

significantly higher Env-specific IgG$^+$ PB responses around week 30 p.i., after Nef-G63E

selection, compared to those in NAb non-inducers (**Figures 6A** and **6B**). At 1 year, this

difference became enhanced; Env-specific PBs were reduced in non-inducers, while this

reduction was ameliorated in the NAb inducers with Nef-G63E.

Concomitant determination of Env-specific IgD$^-$CD27$^+$IgG$^+$ B$_{mem}$ responses allowed pair-

wise analysis of Env-specific IgG$^+$ B$_{mem}$/PB as a projection of germinal center (GC) output as

an indicator of overall B-cell response quality (Zotos et al., 2010) (**Figure 6C**, left). In this two-

dimensional vector charting of Env-specific B$_{mem}$/PB responses, vector protrusion towards

upper right reasonably represents higher gross GC output of antibody-forming cells (AFCs)

(B$_{mem}$/PBs). In NAb non-inducers, all vectors converged on an empirically defined polygonal

attractor area D_n (gray area surrounded with dotted lines in **Figure 6C**, right) at year 1 p.i. and

beyond, describing that these NAb non-inducers failed to sustain Env-specific GC output. In

contrast, the vectors were consistently tracked outside D_n in the NAb inducers with Nef G63E

(**Figure 6C**). At the moment of NAb induction, they converged on another upper-right GC

output attractor area D_i (red area surrounded with dotted lines in **Figure 6C**, right), mutually

exclusive with D_n (P < 0.0001 by Fisher's exact test on NAb non-inducer/inducer vector

distribution frequency within D_n). These suggest more robust virus-specific IgG$^+$ B-cell

responses following Nef-G63E CD8$^+$ T-cell escape mutant selection, predisposing to NAb

induction. This enhanced cognate B-cell signature was also an inverted pattern of impaired

terminally-differentiated B-cell responses in APDS (Al Qureshah et al., 2021).

Taken together, in the current model, Nef$_{62-70}$-specific CD8$^+$ T-cell responses in

persistently SIV$_{mac239}$-infected macaques selected for an escape mutant, Nef-G63E SIV, which

decreased aberrant Nef binding and drive of B-cell-inhibitory PI3K/mTORC2. B-cell Nef

invasion *in vivo* occurred more preferentially in Env-specific B cells, suggesting diminution in

tonic Nef-mediated B-cell dysregulation after mutant selection. These predisposed to

enhanced Env-specific B-cell responses and subsequent SIV$_{mac239}$-specific NAb induction,

altogether, in a manner reciprocal to human APDS-mediated immune dysregulation.

Discussion

In the present study, we found NAb induction against an NAb-resistant SIV after selection of a CD8[+] T-cell escape *nef* mutant virus. The Nef-G63E mutation, positioned in cognate B cells, results in reduction of binding/drive of PI3K/mTORC2, being concordant with *in vivo* enhancement of Env-specific IgG[+] B cells up to NAb induction following mutant selection. Importantly, this manifested through a pAkt deviation level more vivid than wild-type versus germline *PIK3CD* gain-of-function mutation heterozygote mice showing full APDS phenotype, stating that the current Nef-G63E-associated B-cell/NAb phenotype likely occurs in a "reciprocal APDS-like" manner. This proposition was enhanced based on extension of immune cell-intrinsic Nef influence on cognate B cells (Fig. 5), in addition to infected T cells. This work is, to our knowledge, the first to interlink a PI3K/mTORC2-modulating viral signature and enhanced B-cell/NAb responses in a viral infection model.

A link between viral T-cell escape and consequent immune modulation has been previously partially explored, including enhanced mutant HIV-1 epitope peptide binding to inhibitory MHC-I receptor on DCs impairing T cells (Lichterfeld et al., 2007) and decreased CTL-mediated lymph node immunopathology rather driving LCMV-specific antibody production (Battegay et al., 1993). Our results now evidence a new pattern of NAb responses bivalently shaped through viral interactions with both humoral and cellular immunity in AIDS virus infection. SIV$_{mac239}$ infection of macaques possessing MHC-I alleles associated with Nef-G63E mutation can be one unique model to analyze virus-host interaction for B cell maturation leading to NAb induction. A *nef* polymorphism was extracted, partially appending to a virological impact of Nef (Kirchhoff et al., 1995; Gauduin et al., 2006; Stolp et al., 2012) in perturbing immunity whereas the current phenotype differed from such replication-related properties. The *nef* region encoding Nef-G63E is not conserved in HIV-1 (Schindler et al., 2004) and beyond this model study it remains to be addressed whether mutant HIV-1 with a similar phenotype can be obtained.

The major *in vivo* readout of this study is autologous neutralization of a highly resistant

SIV, differing from HIV-specific broadly neutralizing antibodies (bNAbs). Importantly, however, obtained signatures of enhanced IgG[+] Env-specific AFC B cells (class switch/terminal differentiation) and virus neutralization (hypermutation) directly evidence enhanced activity of activation-induced cytidine deaminase (AID), the canonical positive driver of B-cell fate, showing a strong conceptual continuity with bNAb regulation by AID.

Our results extracted an *in vivo* link between a decreased aberrant Nef-PI3K/mTORC2 interaction and major enhancement in B-cell responses. The relationship is indeed reciprocal to immunogenetic mechanisms of human PI3K gain-of-function mutations resulting in APDS with multiply impaired anti-viral B-cell responses. While the exact hierarchy of binding between Nef and PI3K/mTORC2 components (including PI3K isoforms) and its perturbation by Nef-G63E mutation remains much to be investigated, PI3Kp85 did reasonably show the strongest Nef-binding index in PLA assay (**Figure 4D**) as Nef and PI3Kp85 possess SH3-binding PxxP and SH3 domains, respectively (Rickles et al., 1994). Binding decrease in canonical partner PI3Kp85 may domino-trigger decrease with others, resulting in quadruple PI3K/mTORC2 binding decrease and fragile downstream Akt signaling.

In conclusion, we exemplify in a non-human primate AIDS model that NAb induction against a highly resistant SIV occurs after selection of a CD8[+] T-cell escape variant reduced in excess PI3K/mTORC2 drive. These results provide an understanding of how immunodeficiency virus-specific NAb responses can be shaped by both wings of adaptive immune pressure and depicts that immune cell-intrinsic PI3K/mTORC2 manipulation may be pivotal for harnessing antiviral NAb responses. Furthermore, search of other analogies between viral immune dysregulations and human IEIs may lead to identification of novel targets to modulate and harness immune responses in translational settings.

Limitations of study

A limitation of this study is the use of retrospective samples, posing constraints for detecting exact temporal changes in tonic viral B-cell perturbation. Related with this, sampling of lymph node cells in animals belonging to the subgroup of interest at the optimal moment was

362   unpredictable. Nevertheless, usage of such specific animal models does help to depict a

363   temporal cascade of *in vivo* events leading to NAb induction, assisting human immunology. As

364   future extension of this study extracting the NAb-involved molecular axis,

365   manipulation/reconstitution experiments shall be designed.

Acknowledgments

This work was supported by Japanese Initiative for Progress of Research on Infectious Diseases for Global Epidemic (J-PRIDE) (JP19fm0208017 to H.Y.), Japan Program for Infectious Diseases Research and Infrastructure (Interdisciplinary Cutting-edge Research) (JP22wm0325006 to H.Y.), Research Programs on HIV/AIDS, Emerging and Re-emerging Infectious Diseases, and Global Health Issues (JP20fk0410022 to H.Y.; JP18fk0410003, JP20fk0410011, JP20fk0108125, JP20jm0110012, JP21fk0410035, and JP21jk0210002 to T.M.) from Japan Agency for Medical Research and Development, the Ministry of Education, Culture, Sports, Science, and Technology in Japan ([JSPS] KAKENHI Grants 17H02185 and 18K07157 to H.Y.; 21H02745 to T.M.), the Takeda Science Foundation (to H.Y.) and the Imai Memorial Trust for AIDS Research (to H.Y.). We thank Fumiko Ono, Koji Hanari, Katsuhiko Komatsuzaki, Akio Hiyaoka, Hiromi Ogawa, Keiko Oto, Hirofumi Akari, Yasuhiro Yasutomi, Hiromi Sakawaki, Tomoyuki Miura and Yoshio Koyanagi for animal experiment assistance; Hiroyuki Kogure, Sayuri Seki and Julia R. Hirsiger for imaging cytometry technical assistance; Masako Nishizawa, Trang Thi Thu Hau, Shigeyoshi Harada, Takushi Nomura, Akiko Takeda, Taku Nakane, Nami Iwamoto, Taeko K. Naruse, Akinori Kimura and Mark S. de Souza for their help. H.Y. thanks Jun Abe, Shoi Shi, Bettina Stolp, Jens V. Stein, Makoto Yamagishi and Yusuke Yanagi for conceptual discussions.

Author Contributions

H.Y. conceived/designed/performed study and wrote manuscript; T.M. provided animal cohort, co-edited manuscript and co-directed study.

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Figure 1. NAb induction against NAb-resistant SIV$_{mac239}$.

(A) Study design.

(B) Plasma viral loads (SIV *gag* RNA copies/ml plasma) in NAb non-inducers (left) and

inducers (right). Viral loads have been previously partially reported (Yamamoto et al., 2007;

Nomura et al., 2012; Takahashi et al., 2013; Nakane et al., 2013; Iwamoto et al., 2014).

(C) Plasma SIV$_{mac239}$ 100% neutralizing end point titers by 10 TCID$_{50}$ killing assay on MT4-R5

cells. Points on the dotted line show marginally NAb-positive results ($< 1{:}2$). In some animals,

titers were comparable with our reported results using MT4 cells (Kawada et al., 2007).

Figure 2. Selection of a viral *nef* mutation (Nef-G63E) before NAb induction.

(A) Viral *nef* mutations in NAb inducers. A linear schema indicating Nef functional domains is

aligned above. Mutations at timepoints with NAb$^-$ (indicated by white wedge; mostly before 6

months p.i.), NAb$^{+/-}$ (green wedge) and NAb$^+$) (purple wedge; mostly after 1 year) are shown

in individual animals. Black and dark gray represent dominant and subdominant mutations (or

deletion in R-360) by direct sequencing, respectively. Red and pink indicate dominant and

subdominant G63E detection, respectively.

(B) Schema of SIV$_{mac239}$ Nef structure and Nef-G63E mutation orientation.

(C) Comparison of frequencies of macaques having Nef-G63E in plasma viruses between

NAb non-inducers and inducers. Compared by Fisher's exact test.

(D) Temporal relationship of Nef G63E frequencies in plasma virus and NAb induction. Black

boxes (left Y axis) show log$_2$ NAb titers; red triangles and green diamonds (right Y axis) show

percentage of G63E and G63R mutations detected by subcloning (15 clones/point on average),

respectively.

Figure 3. Nef-G63E is a CD8$^+$ T-cell escape mutation.

(A) Nef$_{62-70}$ QW9-specific CD8$^+$ T-cell frequencies and related MHC-I alleles in seven NAb

inducers selecting Nef-G63E. Mamu-B*039:01 and Mamu-B*004:01 are known to restrict

Nef_{62-70} QW9 epitope (Evans et al., 1999; Sette et al., 2012) and binding of Nef_{63-70} peptide to

Mamu-A1*032:02 was predicted.

(B) CD8[+] T-cell responses specific to wild-type Nef_{62-70} or mutant Nef_{62-70}.G63E or Nef_{62-}

$_{70}$.G63R peptides. See also Figure S3.

Figure 4. Nef-G63E mutation reduces PI3K/mTORC2 binding and pAkt drive.

(A) Representative surface expression level histograms of CD3, CD4, MHC class I, CXCR4

and BST-2 after wild-type (WT) or Nef-G63E mutant SIV infection at multiplicity of infection

(MOI) 0.1 on HSC-F cells.

(B) Left: representative histograms of relative pAkt serine (Ser) 473 levels in p27[+] cells after

WT or Nef-G63E mutant SIV infection at MOI 0.1 on HSC-F cells. Numbers show pAkt Ser473

mean fluorescence intensities (MFIs) for each. Right: Deviation of pAkt Ser473 MFIs in p27[+]

HSC-F cells compared with mean MFI of uninfected cells. Compared by unpaired t test.

(C) Left: Relative pAkt Ser473 levels (normalized to mean MFI of uninfected controls) in Nef[+]

HSC-F cells assessed for serum starvation (MOI 0.2, 1 day p.i.). Adjusted *P* values show

results of comparison via Sidak's post-hoc test of 2-way ANOVA (C, left).   Right: Deviation of

pAkt Ser473 MFIs in Nef[+] HSC-F cells assessed for high-MOI infection (MOI 5, 1-day p.i.).

Compared by unpaired t test.

(D) Proximity ligation assay (PLA) of Nef binding with mTOR, GβL/mLST8, PI3Kp85 and

PI3Kp110α. MFI-based binding index was calculated as (anti-Nef/anti-partner) - (isotype/anti-

partner) - (anti-Nef/isotype) + (isotype/isotype). Histograms for samples and isotype/isotype

are representatively shown. Differences in MFI binding indexes can be enhanced compared

with comparison of raw MFIs. Compared by paired t tests. Data represent one of two [(A); (B);

(C, right)] independent experiments in quadruplicate, four independent experiments performed

in triplicate [(C, left)] or four independent single-well comparison experiments pooled for

statistical analysis (D). Bars: mean ± SD (B, C right), mean ± SEM (C left).

Figure 5. Preferential targeting of lymph node Env-specific B cells by Nef *in vivo*.

(A) Representative gating of triple noise-cancellation *in vivo* Nef staining in B cells analyzed by ImageStreamX MKII. Pre-experimental damaged cells are first excluded with Live/Dead from focused/centroid/singlet image-acquired CD19$^+$ B cells (first lane right/R4 gated on "Focused"). Following Nef$^+$ gating (second lane left/R5 gated on R4), a second step of Nef noise cancellation (second lane middle/R7 on R5) comprises double-negative removal of post-experimental stochastic irregular staining building a disparate intracellular staining gradient (X axis, Nef signal pixel Haralick variance mean) and post-experimental batch overt cellular staining (Y axis, Nef signal pixel intensity threshold). This outputs a B-cell population with a fine-textured pericellular Nef^{int-lo} staining, biologically concordant with Nef membrane-anchoring. Probing of anti-Env BCR (αEnv) by recombinant SIV Env (second lane, right) is combined, resulting in a 2-D panel of intracellular Nef versus αEnv for noise-cancelled Nef$^+$ B cells (R7) plus Nef$^-$ B cells (R8) (third lane left/"R7+R8"). pAkt Ser473 expression (third lane right) and cellular morphology (B) was further analyzed.

(B) Typical images of Nef-transferred Env-specific B cells defined αEnv$^+$-intracellular Nef$^+$-Nef Haralick variance mean^{lo}-Nef Intensity threshold^{lo}-Live/Dead$^-$-CD19$^+$ cells ["Nef$^+$aEnv$^+$" population of lower left panel in (A), gated on "R7, R8"]. Note the pericellular pAkt Ser473 upregulation in these cells (Ch 03/yellow). Data on inguinal lymph node lymphocytes of macaque R10-007 at week 62 post-SIV$_{mac239}$ infection are shown in (A) and (B).

(C) Comparison of pAkt Ser473 median fluorescence (medFI) intensity levels in Nef$^-$ B cells (R8) versus noise-cancelled Nef$^+$ B cells (R7). Analyzed by paired t test.

(D) Comparison of Nef-positive cell frequencies in non-Env-specific (left) versus Env-specific (right) B cells in lymph nodes of persistently SIV-infected macaques (*n* = 6). Analyzed by paired t test.

Figure 6. Enhanced SIV Env-specific B-cell output up to NAb induction following Nef-G63E selection.

(A) Changes in SIV Env gp140-specific plasmablast (PB) frequencies in viremic NAb non-

inducers (left, n = 12) and Nef-G63E-selecting NAb inducers (right, n = 6). Available samples

of twelve viremic NAb non-inducers (including ten with $\geq$ 1-year survival) and six NAb inducers

were tracked. In the right panel, the frequencies in NAb non-inducers are shown in background

(gray) for comparison.

(B) Comparison of Env gp140-specific PB frequencies between NAb non-inducers and

inducers by Mann Whitney U tests. In the right, the frequencies at NAb induction were

compared with those at year 1 in NAb non-inducers with $\geq$ 1-year survival (n = 10). Bars: mean

$\pm$ SD.

(C) Left: vector chart of Env gp140-specific memory B cell (B_{mem}) and PB levels. Legends for

each animal correspond to the ones in (B). Right: NAb non-inducer vectors empirically define

a polygonal GC output attractor area D_n (gray area surrounded with dotted lines) on which

they converge by and beyond year 1 p.i. The NAb inducer vectors (shown up to the time of

NAb induction) remained outside of D_n. At the moment of NAb induction they converged on a

second GC output attractor area D_i (red area surrounded with dotted lines), mutually exclusive

with D_n (P < 0.0001 by Fisher's exact test on NAb inducer vector convergence frequency within

D_n). Legends for year 1 p.i. in the NAb non-inducers and moment of NAb induction in the NAb

inducers are specified.

Materials and Methods

Material availability

This study did not generate new unique reagents.

Data and code availability

Data were analyzed using existing computational packages. The accession number for the transcriptome data [wild-type SIV$_{mac239}$-infected HSC-F cell line (n = 3), Nef-G63E SIV$_{mac239}$-infected HSC-F cell line (n = 3) and uninfected HSC-F cell line (n = 3)] has been deposited under the accession number GenBank: GSE65806.

Rhesus macaques retrospectively utilized for samples

Seventy Burmese rhesus macaques (*macaca mulatta*) (57 males and 13 females) were retrospectively analyzed in this study. Experiments were previously carried out (Matano et al., 2004; Yamamoto et al., 2007; Iseda et al., 2016; Nomura et al., 2012; Ishii et al., 2012; Takahashi et al., 2013; Nakane et al., 2013; Shi et al., 2013; Iwamoto et al., 2014; Terahara et al., 2014) in the Tsukuba Primate Research Center, National Institutes of Biomedical Innovation, Health and Nutrition (NIBIOHN) with the help of the Corporation for Production and Research of Laboratory Primates and the Institute for Frontier Life and Medical Sciences, Kyoto University (IFLMS-KU) after approval by the Committee on the Ethics of Animal Experiments of NIBIOHN and IFLMS-KU under the guidelines for animal experiments at NIBIOHN, IFLMS-KU and National Institute of Infectious Diseases in accordance with the Guidelines for Proper Conduct of Animal Experiments established by Science Council of Japan (http://www.scj.go.jp/ja/info/kohyo/pdf/kohyo-20-k16-2e.pdf). The experiments were in accordance with the "Weatherall report for the use of non-human primates in research" recommendations (https://royalsociety.org/topics-policy/publications/2006/weatherall-report/). Animals were housed in adjoining individual primate cages allowing them to make sight and sound contact with one another for social interactions, where the temperature was kept at

25°C with light for 12 hours per day. Animals were fed with apples and commercial monkey

diet (Type CMK-2, Clea Japan, Inc.). Blood collection and virus challenge were performed

under ketamine anesthesia.

Cells and viruses

MT4-R5 cells were maintained in RPMI1640 (Invitrogen) supplemented with 10% fetal bovine

serum (FBS) (Clontech) and antibiotics. HSC-F cells (cynomolgus CD4$^+$ T-cell line) and

HSR5.4 cells (rhesus macaque CD4$^+$ T-cell line) were maintained in RPMI1640 (Invitrogen)

supplemented with 10% fetal bovine serum (FBS) (Clontech), human interleukin-2 (IL-2) (10

IU/ml, Roche), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (Invitrogen) and

2-mercaptoethanol (Gibco). Rhesus macaque primary CD20$^+$ B cells were magnetically

isolated from peripheral blood mononuclear cells (PBMCs) with anti-CD20 microbeads

(Miltenyi Biotec). SIV$_{mac239}$ molecular clone derivative was generated by mutagenesis PCR

(Agilent Technologies) using the primers listed. For virus preparation, COS-1 cells were

transfected with pBR$_{mac239}$ proviral DNA using FuGENE 6 (Promega). At 48 h post-transfection,

culture supernatants were harvested, centrifuged and filtered through a 0.45-μm pore-size

filter (Merck Millipore). To titrate infectivity, prepared viruses were serially diluted and infected

on HSC-F cells in 96-well plates (Falcon) in quadruplicate. At 10 days post-infection, the

endpoint was determined using SIV p27 antigen enzyme-linked immunosorbent assay

(ELISA) kit (ABL), and virus infectivity was calculated as the 50 percent tissue culture infective

dose (TCID$_{50}$) according to the Reed-Muench method.

Identification of SIV$_{mac239}$-NAb inducers

Burmese rhesus macaques previously challenged with the highly pathogenic molecular clone

virus SIV$_{mac239}$ (n = 70) were retrospectively examined approximately up to 2 years for their

plasma NAb profiles. Animals were challenged intravenously with 1,000 TCID$_{50}$ of SIV$_{mac239}$.

In the current study, virological and immunological profiles were compared between the newly

identified NAb inducers (n = 9) and representative NAb non-inducers (n = 19). These NAb

non-inducers and eight NAb inducers except R06-032 were previously partially reported for

their plasma viral loads (Yamamoto et al., 2007; Iseda et al., 2016; Nomura et al., 2012;

Takahashi et al., 2013; Nakane et al., 2013; Iwamoto et al., 2014) (**Figure 1B**). MHC-I

haplotypes and alleles were determined by reference strand-mediated conformation analysis,

PCR-SSP (PCR amplification utilizing sequence-specific priming) and cloning as described

(Naruse et al., 2010). MHC-I binding prediction was made on April 27, 2013 using the IEDB

analysis resource NetMHCpan tool (Hoof et al., 2009). Alleles of interest in the study have

been previously identified in macaques (Nomura et al., 2012; Evans et al., 1999; Sette et al.,

2012). Macaques R06-032, R03-015, R01-010 and R05-010 received a prime-boost

vaccination (Matano et al., 2004) composed of a DNA prime/intranasal Sendai virus vector

expressing SIV_{mac239} Gag (SeV-Gag). R03-015, R06-019, R06-038 and R10-001 received 300

mg of non-specific rhesus IgG at day 7 post-SIV_{mac239} challenge as an experimental control in

our previous reports (Yamamoto et al., 2007; Iseda et al., 2016; Nakane et al., 2013).

Plasma viral load quantitation

Plasma viral RNA samples were extracted with High Pure Viral RNA kit (Roche Diagnostics).

Serial fivefold sample dilutions were amplified in quadruplicate by reverse transcription and

nested PCR using SIV_{mac239} *gag*-specific primers to determine end point via the Reed-Muench

method as described previously (Matano et al., 2004; Iseda et al., 2016). The lower limit of

detection is approximately 400 viral RNA copies/ml plasma.

SIV_{mac239}-specific neutralization assay

NAbs were titrated as described (Yamamoto et al., 2007; Iseda et al., 2016). Serial twofold

dilutions of heat-inactivated plasma, or polyclonal IgG affinity-purified with Protein G

Sepharose 4 Fast Flow (GE Healthcare) from heat-inactivated and filtered plasma were mixed

with 10 $TCID_{50}$ of SIV_{mac239} at a 1:1 ratio (5 µl:5 µl) in quadruplicate. After 45-min incubation at

room temperature, the 10 µl mixtures were added into 5 x 10^4 MT4-R5 cells/well in 96-well

plates. Progeny virus production in day 12 culture supernatants was examined by SIV p27

ELISA (ABL) to determine 100% neutralizing endpoint. The lower limit of titration is 1:2.

Neutralization in three out of four wells at a dilution of 1:2 is defined as marginally NAb-positive

($< 1:2$). Results were comparable when the same assay was performed with macaque HSC-

F cells (Akari et al., 1996) as targets. NAb inducers R01-012, R02-004, R01-010, R02-007

and NAb non-inducer R01-011 were previously partially reported for their NAb titers measured

with the same method using MT4 cells as targets (Kawada et al., 2007), which derived

comparable results. For assessment of neutralizing activity in IgG, SIV_{mac239}-specific IgGs

purified from pools of plasma with SIV_{mac239}-specific NAb titers were obtained from each animal

as described (Yamamoto et al., 2007). After complement heat-inactivation at 56°C, 30 min and

0.45 μm filtration, IgGs were purified by Protein G Sepharose 4 Fast Flow (GE Healthcare)

and concentrated by Amicon Ultra 4, MW 50,000 (Millipore) to 30 mg/ml and similarly

examined for their 10 $TCID_{50}$ SIV_{mac239} killing titers on MT4-R5 cells.

SIV Env-specific IgG ELISA and immunoblotting

Plasma Env-specific IgG titers were measured as described (Nakane et al., 2013). SIV_{mac251}

Env gp120 (ImmunoDiagnostics) were coated on 96-well assay plates (BD) at 1000 ng/ml

(100 μl/well). Wells were prewashed with phosphate-buffered saline (PBS), blocked with 0.5%

bovine serum albumin (BSA)/PBS overnight and plasma samples were incubated at a 1:20

dilution (5 μl:95 μl) for 2 hours. Wells were washed with PBS and SIV Env-bound antibodies

were detected with a horseradish peroxidase (HRP)-conjugated goat anti-monkey IgG (H+L)

(Bethyl Laboratory) and SureBlue TMB 1-Component Microwell Peroxidase Substrate (KPL).

Absorbance at 450 nm was measured. Samples from week 0 pre-challenge and month 3,

month 6 and year 1 post-challenge were assessed in duplicate. Week 0 average values were

subtracted from corresponding later time point values for calibration. For immunoblotting, SIV

virion-specific IgGs in plasma were detected with a SIV_{mac239}-cross-reactive western blotting

system (ZeptoMetrix). In the NAb non-inducers, samples from those showing rapid

progression (euthanized due to AIDS progression within approximately 1 year) with low

plasma anti-SIV reactivity (Nakane et al., 2013) were not included.

Sequencing

Sequencing was performed as described (Matano et al., 2004; Iseda et al., 2016). Viral cDNA

fragments spanning from nt (nucleotide) 4829 to nt 7000, nt 6843 to nt 8831 and nt 8677 to nt

10196 in SIV_{mac239} (GenBank accession number MM33262) covering SIV *env* and *nef* were

amplified from plasma viral RNA by nested RT-PCR using Prime-Script one-step RT-PCR kit

v2 (TaKaRa) and KOD-Plus v2 (Toyobo). PCR products were either directly sequenced or

subcloned with a TOPO blunt-end cloning system (Invitrogen). Sequencing was performed

using dye terminator chemistry with an ABI 3730 DNA sequencer (Applied Biosystems). On

average, 15 clones were obtained per sample and 20 clones were assessed when *nef*

mutations of interest in early time points were subdominant.

SIV_{mac239}-specific CD8$^+$ T-cell responses

Virus-specific CD8$^+$ T-cell frequencies were measured as described (Matano et al., 2004;

Iseda et al., 2016). Peripheral blood mononuclear cells (PBMCs) were cocultured for 6 hours

with autologous herpesvirus papio-immortalized B lymphoblastoid cell lines (B-LCLs) pulsed

with Nef peptides (Sigma-Aldrich Japan) at 1 µM concentration or as indicated otherwise

under GolgiStop (monensin, BD) presence. Intracellular gamma interferon (IFN-γ) staining

was performed using Cytofix/Cytoperm kit (BD) and the following conjugated anti-human

monoclonal antibodies (mAbs): anti-CD4-FITC (M-T477, BD Pharmingen), anti-CD8-PerCP

(SK1, BD Biosciences), anti-CD3-APC (SP34-2, BD Pharmingen) and anti-IFN-γ-PE (4S.B3,

BioLegend). Specific CD8$^+$ T-cell frequencies were determined by subtracting nonspecific IFN-

γ$^+$ CD8$^+$ T-cell frequencies from those after peptide-specific stimulation; frequencies beneath

100 cells/million PBMCs were considered negative. Cells acquired by FACS Canto II (BD)

were analyzed by FACS Diva (BD) and FlowJo (Treestar). Approximately 1 x 10^5 PBMCs were

gated per test.

Nef-mediated signaling perturbation analysis

Virus supernatants obtained from COS-1 cells after transfection with wild-type or mutant

SIV_{mac239} molecular clones were used for infection of CD4$^+$ T-cell lines, cynomolgus macaque-

derived HSC-F and rhesus macaque-derived HSR5.4 (Akari et al., 1996). QuikChange II XL

site-directed mutagenesis kit (Agilent Technologies) was used to construct a mutant SIV_{mac239}

molecular clones possessing a *nef* mutation Nef G63E (G-to-A mutation at nt 9520) from the

wild-type SIV_{mac239} molecular clone (Kestler et al., 1991) (nt number from GenBank accession

number M33262). Cells (1 x 10^5 cells/well in U-bottomed 96-well culture plates [BD]) were

infected with wild-type or mutant SIV_{mac239} cultured in RPMI-1640 medium supplemented with

10% fetal bovine serum (FBS) for indicated periods at MOI 0.1 (intracellular signaling analysis)

or 0.001 (supernatant analysis). Culture supernatants were subjected to measurement of SIV

capsid p27 concentrations by ELISA. Harvested cells were fixed and permeabilized with

Cytofix/Cytoperm kit, washed twice and subjected to immunostaining. The following antibodies

were used; anti-SIV_{mac251} Gag p27 mAb (ABL) manually conjugated with Alexa 647 (Life

Technologies), anti-SIV_{mac251} Nef mAb (clone 17, epitope peptide corresponding to SIV_{mac239}

Nef 71-80 not including the residue G63: Thermo Scientific/Pierce) manually conjugated with

Alexa 647, anti-human CD3-APC-Cy7 (SP34-2, BD Pharmingen), anti-human CD4-PerCP

(L200, BD Pharmingen), anti-human HLA-ABC-FITC (G46-2.6, BD Pharmingen), anti-human

CXCR4-APC (12G5, BioLegend), BST2-Alexa647 (RS38E, BioLegend), Alexa 488-

conjugated anti-phospho-Akt (Ser473) (D9E, CST) and Alexa 488-conjugated anti-phospho-

Akt (Thr308) (C31E5E, CST). Cells acquired by FACS Canto II were analyzed by FACS Diva

and FlowJo. Approximately 5 x 10^4 cells were gated for each test.

Ferritin ELISA

Plasma ferritin levels in NAb inducers and control animals were analyzed by monkey ferritin

sandwich ELISA kit (LS Bio) according to the manufacturer's instructions.

PI3K stimulation assay

5 x 10^4 HSC-F cells were infected with wild-type or Nef G63E mutant SIV$_{mac239}$ at MOI 0.2 and cultured for 1 day in medium supplemented with 10% (normal), 1% (1/10 starvation) or 0.1% (1/100 starvation) FBS. For ligand stimulation (**Figure S4B**), cells at the end of 1-day culture were pulsed for 20 minutes with 40 ng/ml of recombinant human IFN-γ (Gibco/Thermo Fisher), 100 IU/ml of recombinant human IL-2 (Roche Diagnostics) or 10 μg/ml of SIV$_{mac251}$ Env gp130 (ImmunoDx). Cells were intracellularly stained using Cytofix/Cytoperm kit with anti-SIV$_{mac251}$ Nef mAb or anti-SIV$_{mac251}$ p27 mAb manually conjugated to Alexa 647 and PE-conjugated anti-human phospho-Akt (Ser473) (D9E, CST) or Alexa 488-conjugated anti-human phospho-Akt (Ser473). Cells acquired by FACS Canto II were analyzed by FACS Diva and FlowJo. Approximately 7 x 10^4 HSC-F cells were gated per test.

Transcriptome analysis

Total RNAs were extracted using RNeasy Plus Mini kit (Qiagen) from 2 x 10^6 HSC-F cells 1 day after infection with wild-type or Nef G63E mutant SIV$_{mac239}$ at MOI 5. Negative control RNA samples were extracted from 2 x 10^6 uninfected HSC-F cells after culture with the same condition. Three sets of experiments were performed. Total RNA samples were subjected to a quality control (QC) analysis using an Agilent 2100 Bioanalyzer. The obtained amounts of total RNAs were: 12.37 $\pm$ 0.39 (uninfected), 8.80 $\pm$ 0.44 (wild-type) and 9.02 $\pm$ 0.17 (Nef G63E) μg (P = 0.66 for wild-type versus Nef G63E by unpaired t test). In all samples, two bands of 18S and 28S rRNA were confirmed and the RNA integrity number (RIN) was 10. 500 ng of total RNA samples were processed with GeneChip WT Plus reagent (Affymetrix/Thermo Fisher Scientific) to produce 150 μl of fragmented and labeled cDNA samples. These were incubated with a Human Gene 2.0 ST Array (Affymetrix/Thermo Fisher Scientific) for 16 hours, 60 rpm at 45°C using GeneChip hybridization oven 645 (Affymetrix/Thermo Fisher Scientific). Results were scanned with GeneChip Scanner 3000 7G (Affymetrix/Thermo Fisher Scientific) and processed with Affymetrix Expression Console Software (Affymetrix/Thermo Fisher Scientific)

according to the manufacturer's instructions. Expression values were normalized by the RMA

method. Genes above mean background expression within the cognate sample and showing

significant difference between wild-type and Nef G63E SIV ($P < 0.05$ via unpaired t test for

$\log_2$-transformed values, 768 candidates) were determined. Akt-related genes exhibiting a

change of approximately 10% or more were representatively extracted and manually aligned

by the authors.

Peripheral CD4$^+$ T-cell surface staining

Cryopreserved/thawed PBMCs were stained for 30 minutes at 4°C with the following reagents

or conjugated anti-human mAbs: Live/Dead Aqua (Life Technologies), anti-CD4-PerCP (L200,

BD Pharmingen), anti-CD8-APC-Cy7 (RPA-T8, BD Pharmingen), anti-CD3-Alexa 700 (SP34-

2, BD Pharmingen), anti-CD95-PE-Cy7 (DX2, eBioscience), anti-CXCR5-PE (87.1,

eBioscience), anti-PD-1-Brilliant Violet 421 (EH12.2H7, BioLegend) and anti-CXCR3-Alexa

488 (G025H7, BioLegend). Cells acquired by FACS LSRII Fortessa (BD) were analyzed by

FACS Diva and FlowJo. Approximately 1.5×10^5 PBMCs were gated per test.

Proximity ligation assay (PLA)

A flow cytometry-based arrangement of proximity ligation assay (PLA) (Avin et al., 2017) was

performed to quantitatively assess Nef binding to candidate interacting molecules. 1×10^5

HSC-F cells were infected with SIV$_{mac239}$ at MOI 0.05 in U-bottomed plates and permeated

with Cytofix/Perm kit (BD Biosciences). After two washes, they were resuspended in 0.5%

BSA/PBS for prevention of experimental procedure-related loss and stained with mouse anti-

SIV$_{mac251}$ Nef mAb (clone 17, Thermo Scientific/Pierce) or mouse IgG1 isotype control mAb

(P3.6.2.8.1, Abcam) in combination with either of the following rabbit antibodies: anti-GβL

(86B8, CST), anti-mTOR (7C10, CST), polyclonal anti-human PI3K p85 (Merck Millipore), anti-

PI3 Kinase p110α (C73F8, CST) or rabbit IgG1 isotype control mAb (DA1E, CST). Antibody-

stained cells were subsequently probed with Duolink In Situ PLA Probe anti-mouse PLUS and

anti-mouse MINUS probes (Sigma/Merck). Next, they were detected for intermolecular binding

using Duolink flow PLA detection kit (Green) (Sigma/Merck) with a reaction time of 100

minutes for post-mouse/rabbit probe linking amplification. Finally, these PLA-subjected cells

were additionally stained with anti-SIV$_{mac251}$ Gag p27 mAb manually conjugated with Alexa

647 and anti-CD4-PerCP for 30 minutes for 20 minutes at 4°C. Cells acquired by FACS Canto

II were analyzed by FACS Diva and FlowJo. Approximately 1 x 10^5 cells were gated per test.

Binding index (Y axis) was calculated by deriving the deviation from the summation of:

{baseline P: (MFI of background reaction with mouse isotype control/rabbit isotype control) +

anti-Nef antibody-derived background Q: [(MFI of reaction with mouse anti-Nef/rabbit isotype

control) - (MFI of background reaction with mouse isotype control/rabbit isotype control)] +

anti-binding partner antibody-derived background R:[(MFI of reaction with mouse isotype

control/rabbit anti-binding partner molecule) - (MFI of background reaction with mouse isotype

control/rabbit isotype control)]}.

B-cell Nef invasion *in vitro* assay

For Nef invasion inhibition assay, 5 x 10^4 HSC-F cells were infected at MOI 1 for 48 hours.

These were then additionally cocultured with 7.5 x 10^4 Ramos B cells for 12 hours. Ramos B

cells were double-stained with Alexa 488-cojugated Nef and anti-CD19-PC-5.5 (J3-119,

Beckman Coulter), and interacting infected cell-invaded B cell doublet images acquired with

ImageStream X MKII (Amnis/Merck Millipore) were analyzed by IDEAS 6.3 (Amnis/Merck

Millipore/Luminex).

Quantitative *in vivo* imaging flow cytometry

Cryopreserved/thawed lymph node cells (LNCs) from several persistently SIV-infected

macaques used in previous experiments (Nakane et al., 2013; Takahashi et al., 2013; Iwamoto

et al., 2014) were seeded in V-bottomed 96-well plates (Nunc) and blocked with 25 μg/ml of

anti-human CD4 (clone L200: BD) in 100 μl volume for 15 minutes at 4°C. After three washes,

they were stained with 10 μg/ml of recombinant SIV$_{mac239}$ Env gp140-biotin (Immune

Technology) for 30 minutes at 4°C. Cells were then stained with anti-CD19-PC5.5 (J3-119,

Beckman Coulter), Live/Dead Aqua (Invitrogen) and streptavidin-APC-Cy7 (BioLegend) for 30

minutes at 4°C. After two washes, cells were processed with Cytofix/Cytoperm kit (BD) for 20

minutes at 4°C, washed twice, and next intracellularly stained with PE-conjugated anti-human

phospho-Akt (Ser473) (D9E, CST) and anti-SIV_{mac251} Nef mAb (clone 17, Thermo

Scientific/Pierce) manually conjugated with Alexa 488 or mouse IgG1 isotype control mAb

(P3.6.2.8.1, Abcam) for 30 minutes at 4°C. After final two washes, cells were suspended in

0.8% PFA/PBS. All centrifugations for washing (1,200 x g, 2 minutes) were performed at 4°C.

Cells were subjected to image acquisition with Image Stream X MKII imaging flow cytometer

(Amnis/Merck Millipore/Luminex) and analyzed with IDEAS 6.3 software (Amnis/Merck

Millipore/Luminex). Approximately 5 x 10^5 LNCs were acquired for analyses (**Figure 5**). A

custom-implemented linear discriminant analysis-based machine learning module (Luminex)

was utilized for verifying candidate Nef staining signal-related secondary parameters (**Figures**

5 and **S5**) for their efficacy of target population separation.

Peripheral SIV Env-specific B-cell responses

Peripheral SIV Env gp140-specific memory B cells (B_{mem}) and plasmablasts (PBs) were

measured by flow cytometry with procedures modified from previous reports and our

experience (Silveira et al., 2015; Hau et al., 2022). First, cryopreserved/thawed PBMCs

seeded in V-bottomed 96-well plates (Nunc) were blocked with 25 µg/ml of anti-human CD4

(clone L200: BD) in 100 µl volume for 15 minutes at 4°C. This concentration attains blockade

of promiscuous SIV Env binding to CD4 comparable to levels by magnetic depletion of CD3[+]

T cells (data not shown). After three washes, they were next stained with 10 µg/ml of

recombinant SIV_{mac239} Env gp140-biotin (Immune Technology) for 30 minutes at 4°C. They

were subsequently stained with the following anti-human antibodies and fluorochromes for 30

minutes at 4°C with combinations as follows; PB/B_{mem}: anti-CD3-APC-Cy7 (SP34-2, BD

Pharmingen), anti-CD8-APC-Cy7 (RPA-T8, BD Pharmingen), anti-CD14-APC-Cy7 (M5E2,

BioLegend), anti-CD16-APC-Cy7 (3G8, BioLegend), streptavidin-Brilliant Violet 421

(BioLegend), anti-CD19-PC5.5 (J3-119, Beckman Coulter) and Live/Dead Aqua (Invitrogen);

PB: anti-CD10-APC-Cy7 (HI10a, BioLegend), anti-CD11c-PE-Cy7 (3.9, BioLegend), anti-

CD123-PE-Cy7 (6H6, BioLegend) and anti-HLA-DR-PE-Texas Red (TU36, Invitrogen); B_{mem}:

anti-CD27-PE/Dazzle 594 (M-T271, BioLegend), anti-CD10-PE-Cy7 (HI10a, BioLegend), anti-

IgD-FITC (DaKo), anti-CD38-Alexa 647 (AT1, Santa Cruz Biotechnology), anti-IgG-Alexa 700

(G18-145, BD Pharmingen) and anti-CD138-PE (DL-101, eBioscience). After two washes,

B_{mem} samples were suspended in 0.8% PFA/PBS. For PB staining, surface-stained cells were

further processed with Cytofix/Cytoperm kit (BD) for 20 minutes at 4°C, washed twice, and

next intracellularly stained with 10 μg/ml of recombinant SIV_{mac239} Env gp140-biotin (Immune

Technology) for 30 minutes at 4°C. After three washes, they were next stained with anti-human

IgG-PE (G18-145, BD), anti-human/mouse IRF4-eFluor 660 (3E4, eBioscience), anti-human

Ki67-Alexa 700 (B56, BD Pharmingen) and streptavidin-Alexa 488 (BioLegend) for 30 minutes

at 4°C. After final two washes, cells were suspended in 0.8% PFA/PBS. Cells acquired with

FACS LSRII Fortessa were analyzed with FACS Diva and FlowJo. Approximately 1.5×10^5

PBMCs were acquired for B_{mem} and 4×10^5 PBMCs were acquired for PB analyses (**Figure

6**). All centrifugations for washing (1,200 x g, 2 minutes) were performed at 4°C. Env-specific

PB frequencies were 0 cells/million PBMCs for pre-challenge samples in all animals examined,

ruling out background confounding for high-sensitivity quantitation of the population.

Statistical Analysis

Analyses were performed via Prism 8 (GraphPad Software). $P < 0.05$ were considered

significant in two-tailed unpaired t tests, paired t tests, Mann Whitney U tests, Fisher's exact

tests, Wilcoxon signed-rank tests and two-way ANOVA with Sidak's post-hoc multiple

comparison tests. For analysis of Nef-invaded B-cell pAkt levels, median fluorescence

intensities (mFIs) were analyzed due to relatively small numbers of Nef^+ B cells. Analyses

involving pAkt Ser473 levels all derived comparable results between mFI and MFI for wild type

versus Nef G63E virus. Machine learning-based rating of Nef signal parameters was verified

with IDEAS 6.3 Machine Learning Module (Luminex).

References

Akari, H., Mori, K., Terao, K., Otani, I., Fukasawa, M., Mukai, R., and Yoshikawa, Y. (1996). In

vitro immortalization of Old World monkey T lymphocytes with Herpesvirus saimiri: its

susceptibility to infection with simian immunodeficiency viruses. *Virology* 218, 382–388.

Al Qureshah, F., Sagadiev, S., Thouvenel, C.D., Liu, S., Hua, Z., Hou, B., Acharya, M., James,

R.G., and Rawlings, D.J. (2021). Activated PI3Kδ signals compromise plasma cell survival via

limiting autophagy and increasing ER stress. *J Exp Med.* 218, e20211035.

Angulo, I., Vadas, O., Garçon, F., Banham-Hall, E., Plagnol, V., Leahy, T.R., Baxendale, H.,

Coulter, T., Curtis, J., Wu, C., et al. (2013). Phosphoinositide 3-kinase δ gene mutation

predisposes to respiratory infection and airway damage. *Science* 342, 866–871.

Avery DT, Kane A, Nguyen T, Lau A, Nguyen A, Lenthall H, Payne K, Shi W, Brigden H, French

E, et al. (2018). Germline-activating mutations in PIK3CD compromise B cell development and

function. *J Exp Med* 215, 2073–2095.

Avin, A., Levy, M., Porat, Z., and Abramson, J. (2017) Quantitative analysis of protein-protein

interactions and post-translational modifications in rare immune populations. *Nat. Commun.*

8, 1524.

Battegay, M., Moskophidis, D., Waldner, H., Bründler, M.A., Fung-Leung, W.P., Mak, T.W.,

Hengartner, H., and Zinkernagel, R.M. (1993). Impairment and delay of neutralizing antiviral

antibody responses by virus-specific cytotoxic T cells. *J. Immunol.* 151, 5408–5415.

Baumjohann, D., Preite, S., Reboldi, A., Ronchi, F., Ansel, K.M., Lanzavecchia, A., and

Sallusto, F. (2013). Persistent antigen and germinal center B cells sustain T follicular helper

cell responses and phenotype. *Immunity* 38, 596–605.

Bradley, T., Peppa, D., Pedroza-Pacheco, I., Li, D., Cain, D.W., Henao, R., Venkat, V., Hora,

B., Chen, Y., Vandergrift, N.A., et al. (2018). RAB11FIP5 Expression and Altered Natural Killer

Cell Function Are Associated with Induction of HIV Broadly Neutralizing Antibody Responses.

Cell 175, 387–399.e17.

Burns, D.P., Collignon, C., and Desrosiers, R.C. (1993). Simian immunodeficiency virus

mutants resistant to serum neutralization arise during persistent infection of rhesus monkeys.

J. Virol. 67, 4104–4113.

Cubas, R.A., Mudd, J.C., Savoye, A.L., Perreau, M., van Grevenynghe, J., Metcalf, T., Connick,

E., Meditz, A., Freeman, G.J., Abesada-Terk, G. Jr., et al. (2013). Inadequate T follicular cell

help impairs B cell immunity during HIV infection. *Nat. Med.* 19, 494–499.

Evans, D.T., O'Connor, D.H., Jing, P., Dzuris, J.L., Sidney, J., da Silva, J., Allen, T.M., Horton,

H., Venham, J.E., Rudersdorf, R.A., et al. (1999). Virus-specific cytotoxic T-lymphocyte

responses select for amino-acid variation in simian immunodeficiency virus Env and Nef. *Nat.*

Med. 5, 1270–1276.

Fallet, B., Narr, K., Ertuna, Y.I., Remy, M., Sommerstein, R., Cornille, K., Kreutzfeldt, M., Page,

N., Zimmer, G., Geier, F., et al. (2016). Interferon-driven deletion of antiviral B cells at the onset

of chronic infection. *Sci. Immunol.* 1, pii: eaah6817.

François, F., and Klotman, M.E. (2003). Phosphatidylinositol 3-kinase regulates human

immunodeficiency virus type 1 replication following viral entry in primary CD4+ T lymphocytes

and macrophages. *J. Virol.* 77, 2539–2549.

Fruman, D.A., Snapper, S.B., Yballe, C.M., Davidson, L., Yu, J.Y., Alt, F.W., and Cantley, L.C.

(1999). Impaired B cell development and proliferation in absence of phosphoinositide 3-kinase

p85alpha. *Science* 283, 393–397.

Gauduin, M.C., Yu, Y., Barabasz, A., Carville, A., Piatak, M., Lifson, J.D., Desrosiers, R.C.,

and Johnson, R.P. (2006). Induction of a virus-specific effector-memory CD4+ T cell response

by attenuated SIV infection. *J. Exp. Med.* 203, 2661–2672.

Geyer, M., Fackler, O.T., and Peterlin, B.M. (2001). Structure--function relationships in HIV-1

Nef. *EMBO Rep.* 2, 580–585.

Gitlin, A.D., Shulman, Z. and Nussenzweig, M.C. (2014). Clonal selection in the germinal

centre by regulated proliferation and hypermutation. *Nature* 509, 637–640.

Gray, E.S., Madiga, M.C., Hermanus, T., Moore, P.L., Wibmer, C.K., Tumba, N.L., Werner, L.,

Mlisana, K., Sibeko, S., Williamson, C., et al. (2011). The neutralization breadth of HIV-1

develops incrementally over four years and is associated with CD4+ T cell decline and high

viral load during acute infection. *J. Virol.* 85, 4828–4840.

Haigwood, N.L., Watson, A., Sutton, W.F., McClure, J., Lewis, A., Ranchalis, J., Travis, B.,

Voss, G., Letvin, N.L., Hu, S.L., et al. (1996). Passive immune globulin therapy in the

SIV/macaque model: early intervention can alter disease profile. *Immunol. Lett.* 51, 107–114.

Haralick, R.M., Shanmugam, K., and Dinstein, I. (1973) Textural features for image

classification. *IEEE Transactions on Systems, Man, and Cybernetics* SMC-3, 610–621.

Hashimoto, M., Bhuyan, F., Hiyoshi, M., Noyori, O., Nasser, H., Miyazaki, M., Saito, T., Kondoh,

Y., Osada, H., Kimura, S., et al. (2016). Potential role of the formation of tunneling nanotubes

in HIV-1 spread in macrophages. *J. Immunol.* 196, 1832–1841.

Hau, T.T.T., Nishizawa, M., Harada, S., Phan, M.H., Kanno, Y., Nomura, T., Matsuoka, S.,

Kawana-Tachikawa, A., Hall, W.W., Matano, T., et al. (2022). Association of envelope-specific

B-cell differentiation and viral selective pressure signatures in HIV-1 CRF01_AE infection.

AIDS 36, 1629–1641.

Havenar-Daughton, C., Carnathan, D.G., Torrents de la Peña, A., Pauthner, M., Briney, B.,

Reiss, S.M., Wood, J.S., Kaushik, K., van Gils, M.J., Rosales, S.L., et al. (2016). Direct Probing

of Germinal Center Responses Reveals Immunological Features and Bottlenecks for

Neutralizing Antibody Responses to HIV Env Trimer. *Cell Rep.* 17, 2195–2209.

Hoof, I., Peters, B., Sidney, J., Pedersen, L.E., Sette, A., Lund, O., Buus, S., and Nielsen, M..

(2009). NetMHCpan, a method for MHC class I binding prediction beyond humans.

Immunogenetics 61, 1–13.

Hunziker, L., Recher, M., Macpherson, A.J., Ciurea, A., Freigang, S., Hengartner, H., and

Zinkernagel, R.M. (2003). Hypergammaglobulinemia and autoantibody induction mechanisms

in viral infections. *Nat. Immunol.* 4, 343–349.

Iseda, S., Takahashi, N., Poplimont, H., Nomura, T., Seki, S., Nakane, T., Nakamura, M., Shi,

S., Ishii, H., Furukawa, S., et al. (2016). Biphasic CD8+ T-cell defense in simian

immunodeficiency virus control by acute-phase passive neutralizing antibody immunization. *J.*

Virol. 90, 6276–6290.

Ishii, H., Kawada, M., Tsukamoto, T., Yamamoto, H., Matsuoka, S., Shiino, T., Takeda, A.,

Inoue, M., Iida, A., Hara, H., et al. (2012). Impact of vaccination on cytotoxic T lymphocyte

immunodominance and cooperation against simian immunodeficiency virus replication in

rhesus macaques. *J. Virol.* 86, 738–745.

Iwamoto, N., Takahashi, N., Seki, S., Nomura, T., Yamamoto, H., Inoue, M., Shu, T., Naruse,

T.K., Kimura, A., and Matano, T. (2014). Control of simian immunodeficiency virus replication

by vaccine-induced Gag- and Vif-specific CD8$^+$ T cells. *J. Virol.* 88, 425–433.

Jia, B., Serra-Moreno, R., Neidermyer, W., Rahmberg, A., Mackey, J., Fofana, I.B., Johnson,

W.E., Westmoreland, S., and Evans, D.T. (2009). Species-specific activity of SIV Nef and HIV-

1 Vpu in overcoming restriction by tetherin/BST2. *PLoS Pathog.* 5, e1000429.

Joas, S., Sauermann, U., Roshani, B., Klippert, A., Daskalaki, M., Mätz-Rensing, K., Stolte-

Leeb, N., Heigele, A., Tharp, G.K., Gupta, P.M., et al. (2020). Nef-Mediated CD3-TCR

Downmodulation Dampens Acute Inflammation and Promotes SIV Immune Evasion. *Cell Rep.*

30, 2261–2274.e7.

Junt, T., Moseman, E.A., Iannacone, M., Massberg, S., Lang, P.A., Boes, M., Fink, K.,

Henrickson, S.E., Shayakhmetov, D.M., Di Paolo, N.C., et al. (2007). Subcapsular sinus

macrophages in lymph nodes clear lymph-borne viruses and present them to antiviral B cells.

Nature 450, 110–114.

Kawada, M., Tsukamoto, T., Yamamoto, H., Takeda, A., Igarashi, H., Watkins, D.I., and Matano,

T. (2007). Long-term control of simian immunodeficiency virus replication with central memory

CD4+ T-cell preservation after nonsterile protection by a cytotoxic T-lymphocyte-based

vaccine. *J. Virol.* 81, 5202–5211.

Kennedy, S.G., Wagner, A.J., Conzen, S.D., Jordán, J., Bellacosa, A., Tsichlis, P.N., and Hay,

N. (1997). The PI 3-kinase/Akt signaling pathway delivers an anti-apoptotic signal. *Genes Dev.*

11, 701–713.

Kestler, H.W. 3rd, Ringler, D.J., Mori, K., Panicali, D.L., Sehgal, P.K., Daniel, M.D., and

Desrosiers, R.C. (1991). Importance of the nef gene for maintenance of high virus loads and

for development of AIDS. *Cell* 65, 651–662.

Kim, D.H., Sarbassov, D.D., Ali, S.M., Latek, R.R., Guntur, K.V., Erdjument-Bromage, H.,

Tempst, P., and Sabatini, D.M. (2003). GbetaL, a positive regulator of the rapamycin-sensitive

pathway required for the nutrient-sensitive interaction between raptor and mTOR. *Mol. Cell* 11,

895–904.

Kirchhoff, F., Greenough, T.C., Brettler, D.B., Sullivan, L.L. and Desrosiers, R.C. (1995).

Absence of intact nef sequences in a long-term, nonprogressing survivor of HIV-1 infection. N.

Engl. J. Med. *332*, 228–232.

Kumar, A., Abbas, W., Colin, L., Khan, K.A., Bouchat, S., Varin, A., Larbi, A., Gatot, J.S.,

Kabeya, K., Vanhulle, C., et al. (2016). Tuning of AKT-pathway by Nef and its blockade by

protease inhibitors results in limited recovery in latently HIV infected T-cell line. *Sci. Rep.* 6,

24090.

Kumar, V., Scandella, E., Danuser, R., Onder, L., Nitschké, M., Fukui, Y., Halin, C., Ludewig,

B., and Stein, J.V. (2010). Global lymphoid tissue remodeling during a viral infection is

orchestrated by a B cell-lymphotoxin-dependent pathway. *Blood* 115, 4725–4733.

Leuchowius, K.J., Weibrecht, I., Landegren, U., Gedda, L., and Söderberg, O. (2009). Flow

cytometric in situ proximity ligation analyses of protein interactions and post-translational

modification of the epidermal growth factor receptor family. *Cytometry A* 75, 833–839.

Levesque, M.C., Moody, M.A., Hwang, K.K., Marshall, D.J., Whitesides, J.F., Amos, J.D.,

Gurley, T.C., Allgood, S., Haynes, B.B., Vandergrift, N.A., et al. (2009). Polyclonal B cell

differentiation and loss of gastrointestinal tract germinal centers in the earliest stages of HIV-

1 infection. *PLoS Med.* 6, e1000107.

Lichterfeld, M., Kavanagh, D.G., Williams, K.L., Moza, B., Mui, S.K., Miura, T., Sivamurthy, R.,

Allgaier, R., Pereyra, F., Trocha, A., et al. (2007). A viral CTL escape mutation leading to

immunoglobulin-like transcript 4-mediated functional inhibition of myelomonocytic cells. *J. Exp.*

Med. 204, 2813–2824.

Limon, J.J., So, L., Jellbauer, S., Chiu, H., Corado, J., Sykes, S.M., Raffatellu, M., and Fruman,

D.A. (2014). mTOR kinase inhibitors promote antibody class switching via mTORC2 inhibition.

Proc. Natl. Acad. Sci. USA 111, E5076–5085.

Lindqvist, M., van Lunzen, J., Soghoian, D.Z., Kuhl, B.D., Ranasinghe, S., Kranias, G.,

Flanders, M.D., Cutler, S., Yudanin, N., Muller, M.I., et al. (2012). Expansion of HIV-specific T

follicular helper cells in chronic HIV infection. *J. Clin. Invest.* 122, 3271–3280.

Locci, M., Havenar-Daughton, C., Landais, E., Wu, J., Kroenke, M.A., Arlehamn, C.L., Su, L.F.,

Cubas, R., Davis, M.M., Sette, A., et al. (2013). Human circulating PD-1+CXCR3-CXCR5+

memory Tfh cells are highly functional and correlate with broadly neutralizing HIV antibody

responses. *Immunity* 39, 758–769.

Lucas, C.L., Kuehn, H.S., Zhao, F., Niemela, J.E., Deenick, E.K., Palendira, U., Avery, D.T.,
Moens, L., Cannons, J.L., Biancalana, M., et al. (2014). Dominant-activating germline
mutations in the gene encoding the PI(3)K catalytic subunit p110δ result in T cell senescence
and human immunodeficiency. *Nat. Immunol.* 15*, 88–97.

Luo, W., Hawse, W., Conter, L., Trivedi, N., Weisel, F., Wikenheiser, D., Cattley, R.T., and
Shlomchik, M.J. (2019). The AKT kinase signaling network is rewired by PTEN to control
proximal BCR signaling in germinal center B cells. *Nat. Immunol.* 20, 736–746.

Marzec, M., Liu, X., Kasprzycka, M., Witkiewicz, A., Raghunath, P.N., El-Salem, M., Robertson,
E., Odum, N., and Wasik, M.A. (2008). IL-2- and IL-15-induced activation of the rapamycin-
sensitive mTORC1 pathway in malignant CD4+ T lymphocytes. *Blood* 111, 2181–2189.

1002

1003 Mason, R.D., Welles, H.C., Adams, C., Chakrabarti, B.K., Gorman, J., Zhou, T., Nguyen, R.,
1004 O'Dell, S., Lusvarghi, S., Bewley, C.A., et al. (2016). Targeted Isolation of Antibodies Directed
1005 against Major Sites of SIV Env Vulnerability. *PLoS Pathog.* 12, e1005537.

1006

1007 Matano, T., Kobayashi, M., Igarashi, H., Takeda, A., Nakamura, H., Kano, M., Sugimoto, C.,
1008 Mori, K., Iida, A., Hirata, T., et al. (2004). Cytotoxic T lymphocyte-based control of simian
1009 immunodeficiency virus replication in a preclinical AIDS vaccine trial. *J. Exp. Med.* 199, 1709–
1010 1718.

1011

1012 Mattapallil, J.J., Douek, D.C., Hill, B., Nishimura, Y., Martin, M., and Roederer, M. (2005).
1013 Massive infection and loss of memory CD4+ T cells in multiple tissues during acute SIV
1014 infection. *Nature* 434, 1093–1097.

1015

1016 Mikell, I., Sather, D.N., Kalams, S.A., Altfeld, M., Alter, G., and Stamatatos, L. (2011).
1017 Characteristics of the earliest cross-neutralizing antibody response to HIV-1. *PLoS Pathog.* 7,
1018 e1001251.

1019

1020 Moir, S., Malaspina, A., Ogwaro, K.M., Donoghue, E.T., Hallahan, C.W., Ehler, L.A., Liu, S.,
1021 Adelsberger, J., Lapointe, R., Hwu, P., et al. (2001). HIV-1 induces phenotypic and functional
1022 perturbations of B cells in chronically infected individuals. *Proc. Natl. Acad. Sci. USA* 98,
1023 10362–10367.

1024

1025 Moody, M.A., Pedroza-Pacheco, I., Vandergrift, N.A., Chui, C., Lloyd, K.E., Parks, R.,
1026 Soderberg, K.A., Ogbe, A.T., Cohen, M.S., Liao, H.X., Gao, F., et al. (2016). Immune
1027 perturbations in HIV-1-infected individuals who make broadly neutralizing antibodies. *Sci.*
1028 *Immunol.* 1, aag0851.

1029

1030 Moore, P.L., Gray, E.S., Wibmer, C.K., Bhiman, J.N., Nonyane, M., Sheward, D.J., Hermanus,
1031 T., Bajimaya, S., Tumba, N.L., Abrahams, M.R., et al. (2012). Evolution of an HIV glycan-
1032 dependent broadly neutralizing antibody epitope through immune escape. *Nat. Med.* 18,
1033 1688–1692.

1034

1035 Nakane, T., Nomura, T., Shi, S., Nakamura, M., Naruse, T.K., Kimura, A., Matano, T., and
1036 Yamamoto, H. (2013). Limited impact of passive non-neutralizing antibody immunization in
1037 acute SIV infection on viremia control in rhesus macaques. *PLoS ONE* 8, e73453.

1038

1039 Naruse, T.K., Chen, Z., Yanagida, R., Yamashita, T., Saito, Y., Mori, K., Akari, H., Yasutomi, Y.,
1040 Miyazawa, M., Matano, T., et al. (2010). Diversity of MHC class I genes in Burmese-origin
1041 rhesus macaques. *Immunogenetics* 62, 601–611.

1042

1043   Ng, C.T., Jaworski, J.P., Jayaraman, P., Sutton, W.F., Delio, P., Kuller, L., Anderson, D.,

1044   Landucci, G., Richardson, B.A., Burton, D.R., et al. (2010). Passive neutralizing antibody

1045   controls SHIV viremia and enhances B cell responses in infant macaques. *Nat. Med.* 16, 1117–

1046   1119.

1047

1048   Nguyen, H., Ramana, C.V., Bayes, J., and Stark, G.R. (2001). Roles of phosphatidylinositol 3-

1049   kinase in interferon-gamma-dependent phosphorylation of STAT1 on serine 727 and activation

1050   of gene expression. *J. Biol. Chem.* 276, 33361–33368.

1051

1052   Nishimura, Y., Gautam, R., Chun, T.W., Sadjadpour, R., Foulds, K.E., Shingai, M., Klein, F.,

1053   Gazumyan, A., Golijanin, J., Donaldson, M., et al. (2017). Early antibody therapy can induce

1054   long-lasting immunity to SHIV. *Nature* 543, 559–563.

1055

1056   Nomura, T., Yamamoto, H., Shiino, T., Takahashi, N., Nakane, T., Iwamoto, N., Ishii, H.,

1057   Tsukamoto, T., Kawada, M., Matsuoka, S., et al. (2012). Association of major histocompatibility

1058   complex class I haplotypes with disease progression after simian immunodeficiency virus

1059   challenge in Burmese rhesus macaques. *J. Virol.* 86, 6481–6490.

1060

1061   Nutt, S.L., Hodgkin, P.D., Tarlinton, D.M., and Corcoran, L.M. (2015). The generation of

1062   antibody-secreting plasma cells. *Nat. Rev. Immunol.* 15, 160–171.

1063

1064   Omori, S.A., Cato, M.H., Anzelon-Mills, A., Puri, K.D., Shapiro-Shelef, M., Calame, K., and

1065   Rickert, R.C. (2006). Regulation of class-switch recombination and plasma cell differentiation

1066   by phosphatidylinositol 3-kinase signaling. *Immunity* 25, 545–557.

1067

1068   Perfetto, S.P., Chattopadhyay, P.K., Lamoreaux, L., Nguyen, R., Ambrozak, D., Koup, R.A.,

1069   and Roederer, M. (2006). Amine reactive dyes: an effective tool to discriminate live and dead

1070   cells in polychromatic flow cytometry. *J. Immunol. Methods* 313, 199–208.

1071

1072 Petrovas, C., Yamamoto, T., Gerner, M.Y., Boswell, K.L., Wloka, K., Smith, E.C., Ambrozak,

1073 D.R., Sandler, N.G., Timmer, K.J., Sun, X., et al. (2012). CD4 T follicular helper cell dynamics

1074 during SIV infection. *J. Clin. Invest.* 122, 3281–3294.

1075

1076 Qiao, X., He, B., Chiu, A., Knowles, D.M., Chadburn, A., and Cerutti, A. (2006). Human

1077 immunodeficiency virus 1 Nef suppresses CD40-dependent immunoglobulin class switching

1078 in bystander B cells. *Nat. Immunol.* 7, 302–310.

1079

1080 Ray, J.P., Staron, M.M., Shyer, J.A., Ho, P.C., Marshall, H.D., Gray, S.M., Laidlaw, B.J., Araki,

1081 K., Ahmed, R., Kaech, S.M., et al. (2015). The interleukin-2-mTORc1 kinase axis defines the

1082 signaling, differentiation, and metabolism of T helper 1 and follicular B helper T cells. *Immunity*

1083 43, 690–702.

1084

1085 Recher, M., Lang, K.S., Hunziker, L., Freigang, S., Eschli, B., Harris, N.L., Navarini, A., Senn,

1086 B.M., Fink, K., Lötscher, M., et al. (2004). Deliberate removal of T cell help improves virus-

1087 neutralizing antibody production. *Nat. Immunol.* 5, 934–942.

1088

1089 Recher, M., Lang, K.S., Navarini, A., Hunziker, L., Lang, P.A., Fink, K., Freigang, S., Georgiev,

1090 P., Hangartner, L., Zellweger, R., et al. (2007). Extralymphatic virus sanctuaries as a

1091 consequence of potent T-cell activation. *Nat. Med.* 13, 1316–1323.

1092

1093 Rickles, R.J., Botfield, M.C., Weng, Z., Taylor, J.A., Green, O.M., Brugge, J.S., and Zoller, M.J.

1094 (1994). Identification of Src, Fyn, Lyn, PI3K and Abl SH3 domain ligands using phage display

1095 libraries. *EMBO J.* 13, 5598–5604.

1096

1097 Sammicheli, S., Kuka, M., Di Lucia, P., de Oya, N.J., De Giovanni, M., Fioravanti, J., Cristofani,

1098 C., Maganuco, C.G., Fallet, B., Ganzer, L., et al. (2016). Inflammatory monocytes hinder

1099 antiviral B cell responses. *Sci. Immunol.* 1, pii: eaah6789.

1100

1101 Sander, S., Chu, V.T., Yasuda, T., Franklin, A., Graf, R., Calado, D.P., Li, S., Imami, K., Selbach,

1102 M., Di Virgilio, M., et al. (2015). PI3 kinase and FOXO1 transcription factor activity differentially

1103 control B cells in the germinal center light and dark zones. *Immunity* 43, 1075–1086.

1104

1105 Sarbassov, D.D., Guertin, D.A., Ali, S.M., and Sabatini, D.M. (2005) Phosphorylation and

1106 regulation of Akt/PKB by the rictor-mTOR complex. *Science* 307, 1098–1101.

1107

1108 Schindler, M., Münch, J., Brenner, M., Stahl-Hennig, C., Skowronski, J., and Kirchhoff, F.

1109 (2004). Comprehensive analysis of nef functions selected in simian immunodeficiency virus-

1110 infected macaques. *J. Virol.* 78, 10588–10597.

1111

1112 Sette A, Sidney J, Southwood S, Moore C, Berry J, Dow C, Bradley K, Hoof I, Lewis MG,

1113 Hildebrand WH et al. (2012). A shared MHC supertype motif emerges by convergent evolution

1114 in macaques and mice, but is totally absent in human MHC molecules. *Immunogenetics* 64,

1115 421–434.

1116

1117 Shi, S., Seki, S., Matano, T., and Yamamoto, H. (2013). IL-21-producer CD4+ T cell kinetics

1118 during primary simian immunodeficiency virus infection. *Microbes Infect.* 15, 697–707.

1119

1120 Shulman, Z., Gitlin, A.D., Weinstein, J.S., Lainez, B., Esplugues, E., Flavell, R.A., Craft, J.E.,

1121 and Nussenzweig, M.C. (2014). Dynamic signaling by T follicular helper cells during germinal

1122 center B cell selection. *Science* 345, 1058–1062.

1123

1124 Silveira, E.L., Kasturi, S.P., Kovalenkov, Y., Rasheed, A.U., Yeiser, P., Jinnah, Z.S., Legere,

1125 T.H., Pulendran, B., Villinger, F., and Wrammert, J. (2015). Vaccine-induced plasmablast

1126 responses in rhesus macaques: phenotypic characterization and a source for generating

1127 antigen-specific monoclonal antibodies. *J. Immunol. Methods* 416, 69–83.

1128

1129 Sommerstein, R., Flatz, L., Remy, M.M., Malinge, P., Magistrelli, G., Fischer, N., Sahin, M.,

1130 Bergthaler, A., Igonet, S., Ter Meulen, J., et al. (2015). Arenavirus glycan shield promotes

1131 neutralizing antibody evasion and protracted infection. *PLoS Pathog.* 11, e1005276.

1132

1133 Srinivasan, L., Sasaki, Y., Calado, D.P., Zhang, B., Paik, J.H., DePinho, R.A., Kutok, J.L.,

1134 Kearney, J.F., Otipoby, K.L., and Rajewsky, K. (2009). PI3 kinase signals BCR-dependent

1135 mature B cell survival. *Cell* 139, 573–586.

1136

1137 Stolp, B., Imle, A., Coelho, F.M., Hons, M., Gorina, R., Lyck, R., Stein, J.V., and Fackler, O.T..

1138 (2012). HIV-1 Nef interferes with T-lymphocyte circulation through confined environments in

1139 vivo. *Proc. Natl. Acad. Sci. USA* 109, 18541–18546.

1140

1141 Swingler, S., Zhou, J., Swingler, C., Dauphin, A., Greenough, T., Jolicoeur, P., and Stevenson,

1142 M. (2008). Evidence for a pathogenic determinant in HIV-1 Nef involved in B cell dysfunction

1143 in HIV/AIDS. *Cell Host Microbe* 4, 63–76.

1144

1145 Takahashi, N., Nomura, T., Takahara, Y., Yamamoto, H., Shiino, T., Takeda, A., Inoue, M., Iida,

1146 A., Hara,H., Shu, T., et al. (2013). A novel protective MHC-I haplotype not associated with

1147 dominant Gag-specific CD8[+] T-cell responses in SIV_{mac239} infection of Burmese rhesus

1148 macaques. *PLoS ONE* 8, e54300.

1149

1150 Terahara, K., Ishii, H., Nomura, T., Takahashi, N., Takeda, A., Shiino, T., Tsunetsugu-Yokota,

1151 Y., and Matano, T. (2014). Vaccine-induced CD107a+ CD4+ T cells are resistant to depletion

1152 following AIDS virus infection. *J. Virol.* 88, 14232–14240.

1153

1154 Tomaras, G.D., Yates, N.L., Liu, P., Qin, L., Fouda, G.G., Chavez, L.L., Decamp, A.C., Parks,

1155 R.J., Ashley, V.C., Lucas, J.T., et al. (2008). Initial B-cell responses to transmitted human

1156 immunodeficiency virus type 1: virion-binding immunoglobulin M (IgM) and IgG antibodies

1157 followed by plasma anti-gp41 antibodies with ineffective control of initial viremia. *J. Virol.* 82,

1158 12449–12463.

1159

1160 Wang, J., Sohn, H., Sun, G., Milner, J.D. and Pierce, S.K. (2014). The autoinhibitory C-terminal

1161 SH2 domain of phospholipase C-γ2 stabilizes B cell receptor signalosome assembly. *Sci.*

1162 *Signal.* 7, ra89.

1163

1164 Xu, W., Santini, P.A., Sullivan, J.S., He, B., Shan, M., Ball, S.C., Dyer, W.B., Ketas, T.J.,

1165 Chadburn, A., Cohen-Gould, L., et al. (2009). HIV-1 evades virus-specific IgG2 and IgA

1166 responses by targeting systemic and intestinal B cells via long-range intercellular conduits.

1167 *Nat. Immunol.* 10, 1008–1017.

1168

1169 Yamamoto, H., Kawada, M., Takeda, A., Igarashi, H. and Matano, T. (2007). Post-infection

1170 immunodeficiency virus control by neutralizing antibodies. *PLoS ONE 2*, e540.

1171

1172 Zhang, F., Wilson, S.J., Landford, W.C., Virgen, B., Gregory, D., Johnson, M.C., Munch, J.,

1173 Kirchhoff, F., Bieniasz, P.D., and Hatziioannou, T. (2009). Nef proteins from simian

1174 immunodeficiency viruses are tetherin antagonists. *Cell Host Microbe 6*, 54–67.

1175

1176 Zotos, D., Coquet, J.M., Zhang, Y., Light, A., D'Costa, K., Kallies, A., Corcoran, L.M., Godfrey,

1177 D.I., Toellner, K.M., Smyth, M.J., et al. (2010). IL-21 regulates germinal center B cell

1178   differentiation and proliferation through a B cell-intrinsic mechanism. *J. Exp. Med.* 207, 365–

1179   378.

1180

     Supplemental Information Legends

     Fig. 1-Supplement 1. Profile of SIV$_{mac239}$-NAb non-inducer subgroup cohort.

     Summary of a SIV$_{mac239}$-NAb non-inducer subgroup cohort (*n* = 19), composed from previously

     partially characterized, MHC class I haplotype-balanced naïve animals. Symbols A+, B+, E+,

     J+ and D+ represent possession of MHC-I haplotypes *90-120-Ia*, *90-120-Ib*, *90-010-Ie*, *90-*

     088-Ij and *90-010-Id*, respectively. For the titers, - represents negative; +/- represents

     neutralization in three out of four wells in a quadruplicate test. Gray shadings indicate

     euthanasia due to AIDS progression. Mamu-A and Mamu-B alleles of interest in this study are

     listed. These NAb non-inducers were previously partially reported for their plasma viral loads

     and MHC-I alleles (Nomura et al., 2012; Takahashi et al., 2013; Nakane et al., 2013; Iwamoto

     et al., 2014). R01-011 was previously partially reported for NAb titers measured with the same

     method using MT4 cells as targets (Kawada et al., 2007), which derived comparable results.

     Fig. 1-Supplement 2. IgG class-switched NAbs in the inducers.

     Titration of SIV$_{mac239}$-neutralizing IgGs in the NAb inducers. SIV$_{mac239}$-specific IgGs, purified

     from pools of plasma with SIV$_{mac239}$-specific NAb titers, were obtained from each animal as

     described (Yamamoto et al., 2007) and examined for their neutralizing activity by 10 TCID$_{50}$

     SIV$_{mac239}$ killing on MT4-R5 cells. NAb-negative anti-SIV IgG and control IgG were prepared

     from a representative animal (R01-011) with NAb-negative plasma and pooled plasma of

     uninfected rhesus macaques, respectively.

     Fig.1-Supplement 3. Anti-SIV binding IgG profiles in NAb inducers and non-inducers.

     (A) Temporal changes in SIV$_{mac251}$ Env gp120-specific antibody titers in the NAb non-inducers

     without rapid progression [NAb-negative non-rapid progressors, NAb(-) non-RPs] (left, *n* = 11)

     and NAb inducers [NAb(+)] (right, *n* = 9). NAb non-inducers showing rapid progression

     (euthanized with AIDS onset within approximately 1 year) were not included except for

     macaque R10-001 (shown in the left). Samples for non-RPs R06-001 and R10-002 were

unavailable.

(B) Unpaired t test of log-transformed anti-SIV Env Ab titers in NAb(-) non-RPs and NAb

inducers. Bars represent geometric mean absorbance within each group.

(C) Lysed SIV$_{mac}$ virion linear antigen-binding at year 1 p.i. in NAb(-) non-RPs and NAb

inducers. Plasma anti-SIV$_{mac251}$ IgGs were detected using a commercial western blotting

system against the parental strain SIV$_{mac251}$ (ZeptoMetrix). White arrowheads indicate Env

gp120-specific bands. Experiments were performed twice with comparable results.

1216

Fig. 1-Supplement 4. Env sequence variation pattern is not a major characteristic of

NAb induction.

SIV$_{mac239}$ Env sequence variations before and after NAb induction. For each animal, the upper

time point shows pre-induction and the lower time point (in purple) shows post-induction (first

or second time point with NAb titer > 1:2). Dominant residue changes are shown as follows:

gp120 (gray), variable regions V1-V5 (blue), and gp41 (blue gray). Subdominant mutations

are uncolored. Unlabeled gray and black shadings show subdominant and dominant deletion

of residues. Parental SIV$_{mac239}$ sequence is listed for V1-V5. Residues mutated in multiple

animals within V1-V5 are shown in red.

1226

Fig. 3-Supplement 1. Selection of Nef-G63E mutant SIV is linked with Nef$_{62-70}$-specific

CD8⁺ T-cell response positivity.

(A) Nef$_{62-70}$-specific CD8+ T-cell responses around week 20 post-SIV challenge in seven

viremic NAb non-inducers possessing either of the MHC-I alleles Mamu-B*039:01, Mamu-

B*004:01 or Mamu-A1*032:02.

(B) Temporal supernatant SIV p27 concentrations after wild type (WT) or Nef-G63E mutant

SIV infection at multiplicity of infection (MOI) 0.001 on HSC-F cells. Data represent one of two

independent experiments performed in triplicate. Bars: mean ± SEM.

(C) Comparison of plasma viral loads in Nef-G63E-selecting NAb non-inducers (n = 7) versus

non-inducers at indicated time points post-SIV challenge. Bars represent geometric means.

log10-transformed values are compared by unpaired t tests.

1238

Fig. 4-Supplement 1. Decreased Akt hyperactivation properties of Nef-G63E mutant SIV.

(A) Plasma ferritin levels (ng/ml) in examined NAb non-inducers versus Nef-G63E mutant SIV-

selecting NAb inducers around year 1 post-infection. Bars represent mean in each group.

Compared by unpaired t test.

(B) Relative levels of pAkt Ser473 (left panel) and threonine (Thr) 308 (right panel) in SIV Nef+

HSR5.4 cells (MOI 0.1, 4 days p.i.).

(C) Relative pAkt Ser473 levels in Nef+ HSC-F cells without stimuli and transiently pulsed with

IFN-γ, IL-2 or SIV$_{mac}$ Env gp130 after infection (MOI 0.2, 1 day).

(D) Gene expression significantly differing between WT and Nef-G63E SIV infection (MOI 5,

1 day). Black bars show potentially Akt-related genes. Asterisks show dynamics concordant

with the Nef-G63E phenotype (PI3K-pAkt Ser473 downstream downregulation or inhibition

upregulation).

(E) Representative flow cytometric gating (NAb inducer R01-012, wk 30 p.i.) of peripheral

blood CXCR3$^-$CXCR5$^+$PD-1$^+$ memory follicular CD4+ T cells. Cells were gated as CD3$^+$CD8$^-$

CD4$^+$CD95$^+$CXCR5$^+$PD-1$^+$CXCR3$^-$ live singlet PBMCs. Available samples of six NAb inducers

showing peak Nef-G63E mutant selection around week 30 p.i. and thirteen viremic NAb non-

inducers were tracked.

(F) Comparison of peripheral CXCR3$^-$CXCR5$^+$PD-1$^+$ memory Tfh cell frequencies between

pre-SIV infection and around week 30 post-infection in NAb non-inducers developing disease

before week 60 (left: $n = 5$) and alive more than 70 weeks (middle: $n = 8$) and Nef-G63E-

detected NAb inducers alive more than 70 weeks (right: $n = 6$). Analyzed by Wilcoxon signed-

rank tests.

MFIs relative to those in uninfected controls (%) are shown. **: $P < 0.01$, ***: $P < 0.001$, n.s.:

not significant between WT and Nef-G63E mutant by unpaired t tests (B-C). Data represent

one of two (B, C) independent experiments performed in triplicate (C: unstimulated) or

quadruplicate (B, C: PI3K-pulsed). Bars: mean ± SEM.

1265

Fig. 5-Supplement 1. Machine learning-verified morphological gating of Nef-invaded B

cells *in vivo*.

(A) Representative gating strategy plus schematic of inguinal lymph node Nef$^+$CD19$^+$ B cells

of an SIV$_{mac239}$-infected rhesus macaque analyzed by ImageStreamX MK II imaging flow

cytometry (R10-012, week 160 p.i.). First-step incorporation of Live/Dead staining gates out

CD19$^+$ B cells staining positive for Nef potentially due to pre-experimental membrane

permeation (e.g., image #246896, #248952 and #71859). Second-step dual gating for

intracellular Nef signal pixel Haralick variance mean (H Variance mean, X axis), a textural

feature calculated from a gray level co-occurrence matrix (GLCM) for each image representing

adjacent pixel signal strength heterogeneity and Nef signal intensity threshold (Y axis) gates

out CD19$^+$ B cells showing potentially non-specific, strong intracellular binary clustered

staining pixels for Nef deriving a large variance (X axis), and/or cells stained overtly strong for

Nef and hence becoming void for feasible image verification (high threshold) (Y axis), both

likely originating stochastically from post-experimental rupturing procedures including

membrane permeation (e.g. image #18541). This two-step noise cancellation results in

acquisition of a Live/Dead$^-$-Nef signal Haralick variance mean^{lo}-Nef signal intensity threshold^{lo}-

Nef$^+$CD19$^+$ B-cell population (e.g. image #116318, #116250 and #132927) with a pericellular

Nef^{int-lo} staining, biologically concordant with Nef membrane anchoring, resembling Nef$^+$ cells

experimentally generated in vitro and showed the highest score in IDEAS6.3 machine learning

module scoring for segregation of typical void vs. valid images (upper right). This gating was

implemented for counting and analysis of Env-specific B-cell Nef invasion *in vivo* (Figure 5).

(B) Detection of contact-dependent Nef invasion from SIV-infected HSC-F CD4$^+$ T cells to

cocultured Ramos B cells *in vitro*. Two typical images of an intercellular conduit (SSC, purple)

with Nef (green) protrusions from infected HSC-F cells to CD19$^+$ (red) Ramos B cells from two

independent experiments are shown.

1291

Fig. 6-Supplement 1. Gating strategy of SIV$_{mac239}$ Env gp140-specific B-cell responses.

Representative gating (R02-004, week 32 p.i.) of SIV Env gp140-specific memory B cells

(B_{mem}) and plasmablasts (PBs). Two panels for PB gating are shown in lower resolution for

visibility. B_{mem} staining was performed separately and gatings are merged with PB panels

(second row, panels 2–4). For lineage-specific exclusion (first row, panel 4), CD10 was stained

independently for B_{mem} (see parenthesis).

A Surface downregulation

CD3 CD4 MHC Class I CXCR4 BST-2

WT
Nef-G63E
Uninfected

B MOI 0.1 Day 3

pAkt Ser473

P = 0.0009

WT
Nef-G63E
Uninfected

C Accaleration w/upstream PI3K stimuli Day 3→Day 1

Serum starvation

P = 0.99 *P* = 0.0017

WT
Nef-G63E

Starvation

%serum in culture

High MOI

P = 0.0011

WT Nef G63E

D Intracellular PI3K-pathway binding

mTORC2 PI3 Kinase

GβL/mLST8 mTOR PI3Kp85 PI3Kp110α

CD4^lop27+ singlet HSC-F

WT
Nef-G63E
WT (isotype)
Nef G63E (isotype)

P = 0.039 *P* = 0.028 *P* = 0.022 *P* = 0.043

NAb non-inducer cohort (n = 19)

MHC-I/ID	\multicolumn SIV$_{mac239}$ neutralization at 1:2 plasma dilution by 10TCID$_{50}$ killing assay (wks p.i.)									pVL at 6 mo	Nef G63E at 1 yr	Detected *Mamu* alleles
	wk11–20	wk21–30	wk31–40	wk41–50	wk51–60	wk61–70	wk71–80	wk81–90	wk91–104			
A+												
R06-037		wk25 -		wk46 +/-			wk72 -	wk87 -	wk95 -	6.4 x 10^2	- (pVL < 400)	
R07-001		wk25 -		wk42 -				wk88 +/-	wk96 -	< 400	- (pVL < 400)	*A1*032:02*
R07-004		wk25 -		wk46 -		wk64 -	wk74 -	wk86 -		4.5 x 10^3	+	*B*039:01*
R07-009		wk25 -		wk46 -		wk64 -	wk74 -			4.5 x 10^3	-	
B+												
R06-001		wk25 -		wk48 -	wk60 -		wk80 -	wk90 -		8.2 x 10^3	-	
R06-039		wk25 -		wk46 -	wk56 -					8.5 x 10^6	-	*A1*032:02 , B*004:01*
E+												
R01-011		wk22 -	wk40 -		wk52 -		wk71 -	wk85 -	wk104 -	1.3 x 10^5	-	*B*004:01*
R05-007		wk26 -			wk55 +/-		wk71 -	wk89 -	wk97 -	1.6 x 10^4	-	
R08-007		wk21 -		wk46 -		wk69 -		wk82 +/-		8.5 x 10^6	-	
R09-011	wk18 -		wk36 -		wk52 -					8.5 x 10^6	-	
R06-038		wk25 -			wk58 +/-		wk80 +/-		wk95 -	1.1 x 10^5	-	
J+												
R04-014	wk12 -	wk25 -	wk40 -							1.5 x 10^5	+ (wk40)	*B*039:01*
R06-022	wk12 -	wk23 -								1.4 x 10^7	- (wk23)	*B*039:01*
R10-001	wk14 -	wk30 -	wk37 -							1.9 x 10^7	- (wk37)	*B*039:01*
D+												
R05-006		wk25 -		wk48 -			wk80 -	wk90 -	wk98 -	5.4 x 10^2	-	*A1*032:02 , B*004:01*
R06-034		wk25 -		wk42 -		wk61 -	wk80 +/-	wk88 +/-	wk96 -	4.5 x 10^3	-	*A1*032:02 , B*004:01*
R08-005		wk25 -		wk48 -		wk68 -			wk100 -	9.0 x 10^2	-	*A1*032:02 , B*004:01*
Others												
R10-002	wk14 -				wk59 -				wk104 -	5.7 x 10^5	-	
R10-006	wk14 -	wk30 -			wk55 -					1.0 x 10^6	-	

Fig. 1-Supplement 1.

Group	Rhesus ID	mg/ml (IgG)	Collected/pooled (wks post-challenge)	10 $TCID_{50}$ SIV_{mac239} 100% killing titer
NAb (+)	R06-032	30	wk 76	1:16
	R01-012	30	wk 73–87	1:32
	R03-015	30	wk 112	1:8
	R02-004	30	wk 76	1:4
	R01-010	30	wk 77	1:16
	R06-019	30	wk 85	1:4
	R-360	30	wk 60–86	1:4
	R02-007	30	wk 42–70	1:32
	R05-010	30	wk 71	1:64
NAb (-)	R01-011	42	wk 104	< 1:2
uninfected	pooled	36	-	< 1:2

Fig. 1-Supplement 2.

Fig.1-Supplement 3.

Fig. 1-Supplement 4.

A

Nef$_{62-70}$ QW9-specific responses in NAb non-inducers

Nef$_{62-70}$-specific CD8+ T-cell frequencies (/million PBMCs)

R06-039 R01-011 R06-022 R10-001 R05-006 R06-034 R08-005

B

SIV p27 (ng/ml)

Days post-SIV infection on HSC-F cells

WT
Nef-G63E

C

3 mo	6 mo	1 yr	1.5 yr
P = 0.2495	*P* = 0.5078	*P* = 0.8928	*P* = 0.6488

Plasma viral loads (vRNA copies/ml)

NAb-   NAb+
Nef-G63E+

Fig. 3-Supplement 1.

Fig. 4-Supplement 1.

A

Intracellular pixel Nef signal
Intensity threshold

Signal intensity threshold (int/hi)

Signal intensity threshold (lo)

Live/Dead⁻CD19⁺
Nef⁺ B cells

IDEAS6.3 machine learning-based 2-D
segregation of typical valid *vs.* void
Live/Dead⁺Nef⁺CD19⁺ B-cell images

Good,bad

Void

Valid

Intracellular pixel Nef signal
strength distribution

2-D adjacent pixel signal heterogeneity (++)
→Haralick variance mean (int-hi)

2-D adjacent pixel signal heterogeneity (+/-)
→Haralick variance mean (lo)

Live/Dead⁺CD19⁺Nef⁺ B cells (void)

**Live/Dead⁻CD19⁺Nef⁺-(Intracellular Nef Haralick variance
mean^int/hi or Nef Intensity threshold^int/hi) B cells (void)**

**Live/Dead⁻CD19⁺Nef⁺-Intracellular Nef Haralick variance
mean^lo-Nef Intensity threshold^lo B cells (valid)**

B

***In vitro* CD19⁺Nef⁺ B cells**

DIC   Nef   CD19   SSC   Merged

Fig. 5-Supplement 1.

Fig. 6-Supplement 1.