SIV-specific neutralizing antibody induction following selection of a PI3K drive-

attenuated *nef* variant

Hiroyuki Yamamoto[*,1,2,3] and Tetsuro Matano[*,1,3,4]

[1]AIDS Research Center, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashi-

Murayama City, Tokyo 208-0011, Japan

[2]Department of Biomedicine, University Hospital Basel, Hebelstrasse 20, Basel 4031,

Switzerland

[3]Joint Research Center for Human Retrovirus Infection, Kumamoto University, 2-2-1 Honjo,

Kumamoto 860-0811, Japan

[4]The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku,

Tokyo 108-8639, Japan

*Correspondence:

Hiroyuki Yamamoto, AIDS Research Center, National Institute of Infectious Diseases, 4-7-1

Gakuen, Musashi-Murayama City, Tokyo 208-0011, Japan. Phone: +81-42-848-7154; Fax:

+81-42-848-7155. Email: h-yamato@niid.go.jp

Tetsuro Matano, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo

162-8640, Japan. Phone: +81-3-4582-2811; Fax: +81-3-5285-1165. Email:

tmatano@niid.go.jp

Running title: SIV neutralization post-PI3K diminution

Abstract

HIV and simian immunodeficiency virus (SIV) infections are known for impaired neutralizing

antibody (NAb) responses. While sequential virus-host B cell interaction appears to be basally

required for NAb induction, driver molecular signatures predisposing to NAb induction still

remain largely unknown. Here we describe SIV-specific NAb induction following a virus-host

interplay decreasing aberrant viral drive of phosphoinositide 3-kinase (PI3K). Screening of

seventy difficult-to-neutralize SIV_{mac239}-infected macaques found nine NAb-inducing animals,

with seven selecting for a specific CD8$^+$ T-cell escape mutation in viral *nef* before NAb

induction. This Nef-G63E mutation reduced excess Nef interaction-mediated drive of B-cell

maturation-limiting PI3K/mammalian target of rapamycin complex 2 (mTORC2). *In vivo*

imaging cytometry depicted preferential Nef perturbation of cognate Envelope-specific B cells,

suggestive of polarized contact-dependent Nef transfer and corroborating cognate B-cell

maturation post-mutant selection up to NAb induction. Results collectively exemplify a NAb

induction pattern extrinsically reciprocal to human PI3K gain-of-function antibody-

dysregulating disease, and indicate that harnessing the PI3K/mTORC2 axis may facilitate NAb

induction against difficult-to-neutralize viruses including HIV/SIV.

Keywords

neutralizing antibody; B cells; CD8$^+$ T cells; PI3 kinase; inborn errors of immunity; HIV/SIV

Introduction

Virus-specific neutralizing antibody (NAb) responses by B cells are induced by a tight cooperation of adaptive immune cells (Kumar et al., 2010; Shulman et al., 2014; Giltin et al., 2014; Wang et al., 2014) and often play a central role in clearance of acute viral infections (Junt et al., 2007). In contrast, persistence-prone viruses such as human immunodeficiency virus type 1 (HIV-1), simian immunodeficiency virus (SIV) and lymphocytic choriomeningitis virus (LCMV) variously equip themselves with B cell/antibody-inhibitory countermeasures (Moir et al., 2001; Mattapallil et al., 2005; Sommerstein et al., 2015; Sammicheli et al., 2016; Fallet et al., 2016; Mason et al., 2016) and NAb responses are impaired (Levesque et al.,2009; Mikell et al., 2011). These viruses successfully suppress elicitation of potent NAb responses particularly in acute infection (Hunziker et al., 2003; Tomaras et al., 2008) and establish viral persistence, posing considerable challenge for developing protective strategies. Particularly, HIV and SIV establish early a large *in vivo* body of infection with a distinct host genome-integrating retroviral life cycle (Whitney et al., 2014). In addition, lentiviruses HIV/SIV are unique in launching a matrix of host immune-perturbing interactions mainly by their six remarkably pleiotropic accessory viral proteins optimally fueling pathogenesis (Kirchhoff et al., 1995; Sheehy et al., 2002; Harris et al., 2003; Schindler et al., 2006; Neil et al., 2008; Zhang et al., 2009; Laguette et al., 2011; Yamada et al., 2018; Joas et al., 2018; Langer et al., 2019; Yan et al., 2019; Joas et al., 2020; Khan et al., 2020; Volcic et al., 2020; Reuschl et al., 2022). A body of evidence has depicted this over these 30 years, whereas its entity, including focal mechanisms of humoral immune perturbation in HIV/SIV infection, still remains elusive.

Contrasting this detrimental consequence of acute-phase endogenous NAb absence driven by adverse virus-host interactions in early HIV/SIV infection, we and others have previously described in *in vivo* experimental models that acute-phase passive NAb infusion can trigger an endogenous T-cell synergism resulting in robust SIV/SHIV (simian/human immunodeficiency virus) replication control (Haigwood et al., 1996; Yamamoto et al., 2007; Ng et al., 2010; Iseda et al., 2016; Nishimura et al., 2017). This shows that virus-specific NAbs

not only confer sterile protection but also can evoke T-cell-mediated non-sterile viral control,

suggesting the importance of inducing endogenous NAb responses during an optimal time

frame as well as humoral-cellular response synergisms. Therefore, identifying mechanisms

driving NAb induction against such viruses is an important step to eventually design NAb-

based HIV control strategies.

As an approach, analysis of *in vivo* immunological events linked with NAb induction

against difficult-to-neutralize SIVs in a non-human primate model can provide important

insights into this goal. Various *in vivo* signatures of HIV-specific NAb induction, such as

antibody-NAb coevolution (Moore et al., 2012), autoimmune-driven induction (Moody et al.,

2016) and natural killer cell-related host polymorphisms (Bradley et al., 2018) have been

reported to date. The variety collectively, and interestingly, indicates that pathways to NAb

induction against difficult-to-neutralize viruses including HIV/SIV are redundant, conversely

implicating as-yet-unknown mechanisms driving NAb induction. For example, the

neutralization resistance of certain SIV strains apparently is not explanatory by any of the

aforementioned, posing such models as attractive analytic targets of NAb induction

mechanisms.

In the present study, we examined virus-specific antibody responses in rhesus

macaques infected with a highly difficult-to-neutralize SIV strain, SIV_{mac239}. This virus is

pathogenic in rhesus macaques causing simian AIDS across a broad range of geographical

origin of macaques (Cumont et al., 2008), shows persistent viremia and hosts lack NAb

responses throughout infection (Kestler et al., 1991; Nomura et al., 2012). Yet, upon

performing a large-scale screening of SIV_{mac239}-infected Burmese rhesus macaques for up to

100 weeks, we noticed a subgroup inducing NAbs in the chronic phase. Interestingly, these

animals commonly selected for a specific CD8$^+$ cytotoxic T lymphocyte (CTL) escape mutation

in the viral Nef-coding gene before NAb induction. The mutant Nef, compared with wild-type

(WT) Nef, manifested a decrease in aberrant interaction with phosphoinositide 3-kinase

(PI3K)/mammalian target of rapamycin complex 2 (mTORC2), resulting in decreased

downstream hyperactivation of the canonical B-cell negative regulator Akt (Omori et al., 2006;

Limon et al., 2014). Nef was graphically identified via machine learning-assisted imaging

cytometry to preferentially target Env-specific B cells *in vivo*, and in those macaques inducing

NAbs, the NAb induction was linked with sustained Env-specific B-cell responses after or

during the mutant Nef selection. Results describe a B-cell functional boosting phenotypically

reciprocal to a recently found human PI3K gain-of-function and antibody-dysregulating inborn

error of immunity (IEI), activated PI3 kinase delta syndrome (APDS) (Angulo et al., 2013;

Lucas et al., 2014), and suggest that intervening PI3K/mTORC2 signaling can potentially

result in harnessing NAb induction against difficult-to-neutralize viruses.

Results

Identification of macaques inducing SIV$_{mac239}$-neutralizing antibodies

112   We performed a retrospective antibody profile screening in rhesus macaques infected

with NAb-resistant SIV$_{mac239}$ (n = 70) (**Figure 1A**) and identified a group of animals inducing

anti-SIV$_{mac239}$ NAb responses (n = 9), which were subjected to characterization. These NAb

inducers showed persistent viremia with no significant difference in viral loads compared with

a profile-clarified naïve, major histocompatibility complex class I (MHC-I) haplotype-balanced

subgroup of NAb non-inducers (n = 19) representatively utilized for comparison hereafter

(**Figure 1B** and **Figure 1-figure supplement 1**). Plasma SIV$_{mac239}$-NAb titers measured by 10

TCID$_{50}$ SIV$_{mac239}$ virus-killing assay showed an average maximum titer of 1:16, being induced

at an average of 48 weeks post-infection (p.i.) (**Figure 1C**). Two of the nine NAb inducers

showed detectable NAb responses before six months p.i., while the remaining seven induced

NAbs after six months. Anti-SIV$_{mac239}$ neutralizing activity was confirmed in immunoglobulin G

(IgG) purified from plasma of these NAb inducers (**Figure 1-figure supplement 2**). SIV Env-

binding IgGs developed from early infection in NAb inducers as well as non-rapid-progressing

NAb non-inducers, the latter differing from rapid progressors known to manifest serological

failure (Hirsch et al., 2004; Nakane et al., 2013) (**Figure 1-figure supplement 3A**); titers were

not higher but rather lower at year 1 p.i. in the NAb inducers (**Figure 1-figure supplement 3B**

and **3C**). This was consistent with other reports (Havenar-Daughton et al., 2016) and differing

with gross B-cell enhancement in Nef-deleted SIV infection with enhanced anti-SIV binding

antibody titers accompanying marginal neutralization (Adnan et al., 2016). NAb inducers and

non-inducers showed similar patterns of variations in viral *env* sequences (Burns et al., 1993),

mainly in variable regions 1, 2 and 4 (V1, V2 and V4) (**Figure 1-figure supplement 4**).

Selection of a viral *nef* mutation (Nef-G63E) precedes chronic-phase SIV$_{mac239}$-specific

NAb induction

136   Viral nonsynonymous polymorphisms outside *env* were next assessed. Strikingly, we

found selection of a viral genome mutation resulting in G (glycine)-to-E (glutamic acid)

substitution at residue 63 of Nef (Nef-G63E) in seven of the nine NAb inducers (**Figure 2A**).

This 63rd residue lies in the unstructured N-terminus of Nef, flanked by two α-helices

conserved in SIVmac/SIVsmm (**Figure 2B**). This region is generally polymorphic among HIV-

1/HIV-2/SIV, occasionally being deleted in laboratory and primary isolate HIV-1 strains (Geyer

et al., 2001). AlphaFold2-based structure prediction did not derive palpable change except for

low-probability disruption of the alpha helices (data not shown). The Nef G63E mutation was

not observed in two early NAb inducers (with detectable NAbs before 6 months p.i.) but

selected in all of the remaining seven inducing detectable NAb responses after 6 months. In

contrast, this mutation was found only in two of the nineteen NAb non-inducers, including one

rapid progressor (**Figure 1-figure supplement 1**) and selection was significantly enriched in

the NAb inducers as compared to the nineteen non-inducers (**Figure 2C**). Analysis of Nef-

G63E mutation frequencies in plasma viruses showed that Nef-G63E selection preceded or

at least paralleled NAb induction (**Figure 2D**).

Nef-G63E is a CD8$^+$ T-cell escape mutation

These seven NAb inducers selecting this mutation elicited CD8$^+$ T-cell responses

specific for a 9-mer peptide Nef$_{62-70}$ QW9 (QGQYMNTPW) (**Figure 3A**). This Nef$_{62-70}$ QW9

epitope is restricted by MHC-I molecules Mamu-B*004:01 and Mamu-B*039:01 (Evans et al.,

1999; Sette et al., 2012). Possession of these accounted for six cases of Nef-G63E selection

(**Figure 3A**), and the remaining one animal possessed Mamu-A1*032:02 predicted to bind to

Nef$_{63-70}$ GW8 (NetMHCpan). Ten of nineteen NAb non-inducers also had either of these alleles

(**Figure 1-figure supplement 1**). This did not significantly differ with the NAb inducer group

(P = 0.25 by Fisher's exact test, data not shown), indicating that NAb induction was not simply

linked with possession of these MHC-I genotypes but instead required furthermore specific

selection of the Nef-G63E mutation (**Figure 2C**). Nef-G63E was confirmed to be an escape

mutation from Nef$_{62-70}$-specific CD8$^+$ T-cell responses (**Figure 3B**). NAb non-inducers

possessing these alleles elicited little or no Nef$_{62-70}$-specific CD8$^+$ T-cell responses (**Figure 3-**

figure supplement 1A), suggesting that *in vivo* selection and fixation of this Nef-G63E SIV

was indeed Nef_{62-70}-specific $CD8^+$ T cell-dependent. Replication kinetics of SIV carrying the

Nef-G63E mutation was comparable with wild-type (WT) on a cynomolgus macaque HSC-F

$CD4^+$ T-cell line (Akari et al., 1996), excluding the possibility that the mutation critically impairs

viral replication (**Figure 3-figure supplement 1B**), and plasma viral loads were comparable

between Nef-G63E mutant-selecting NAb inducers versus non-inducers (**Figure 3-figure**

supplement 1C). These results indicate NAb induction following *in vivo* selection and fixation

of the $CD8^+$ T-cell escape *nef* mutation, Nef-G63E under viral persistence.

G63E mutation reduces aberrant Nef interaction-mediated drive of PI3K/mTORC2

175   We next focused on the functional phenotype of this highlighted Nef-G63E mutant SIV

in infected cells. An essence of host perturbation by Nef is its wide-spectrum molecular

downregulation, ultimately increasing viral replication (Schindler et al., 2004; Zhang et al.,

2009). To address possible amelioration, we compared downregulation of major targets CD3,

CD4, MHC-I, CXCR4 and BST-2 (Jia et al., 2009) in infected HSC-F cells. Nef-G63E mutation

did not confer notable change compared with wild-type (WT) [P = not significant (*n.s.*) for all

molecules, data not shown], implicating other non-canonical changes (**Figure 4A**). Literature

suggested ferritin to exert negative influence on B cells proportionate to HIV-infected

macrophage-derived production (Swingler et al., 2008). Here, plasma ferritin levels showed

no differences (**Figure 4-figure supplement 1A**) and viral loads were comparable between

Nef-G63E-selecting NAb inducers and non-inducers (**Figure 4-figure supplement 1B** and

1C), arguing against gross involvement of ferritin as well as other viral replication-changing

Nef phenotypes at least in this model.

188   Thus, we took interest in immunosignaling-modulating properties of Nef-G63E SIV.

189   Analysis of Akt, the predominant immune-intrinsic negative brake of B-cell maturation and

190   AFC/antibody responses *in vivo* (Omori et al., 2006; Limon et al., 2014; Fruman et al., 1999;

191   Ray et al., 2015; Sander et al., 2015), revealed that its serine 473 phosphorylation, non-

192   canonically known for Nef-mediated upregulation (Kumar et al., 2016), was significantly lower

on day 3 after Nef-G63E SIV infection at low multiplicity of infection (MOI) compared with WT (**Figure 4B**), stronger than histogram deviation levels in PI3K gain-of-function mice with full recapitulation of B-cell/antibody-dysregulating human APDS disease phenotype (Avery et al., 2018). The threonine 308-phosphorylated Akt (pAkt Thr308) level remained unaffected (**Figure 4-figure supplement 1B**).

We further assessed properties of serine 473-phoyphorylated Akt (pAkt Ser473) inhibition by Nef-G63E SIV, and found that external PI3K stimuli (serum starvation) (Kennedy et al., 1997) accelerated phenotype appearance from day 3 to day 1 post-infection (**Figure 4C**, left). A similar trend was obtained by short-term PI3K stimulation with interferon-γ (Nguyen et al., 2001), interleukin-2 (Marzec et al., 2008) and SIV Env (François & Klotman, 2003) (**Figure 4-figure supplement 1C**). A high-MOI SIV infection, comprising higher initial concentration of extracellular Env stimuli, also accelerated phenotype appearance from day 3 to day 1 post-infection, with stronger pAkt reduction (**Figure 4C**, right). These showed that the Akt-inhibitory G63E mutant Nef phenotype is PI3K stimuli-dependent. Transcriptome analysis signatured decrease in PI3K-pAkt-FoxO1-related genes in mutant SIV infection (**Figure 4-figure supplement 1D**). As Akt is also a survival mediator this implicated that Nef-G63E mutation may be Th-cytopathic, decreasing Th dysfunctional expansion (Baumjohann et al., 2013) as had been observed in LCMV strain WE (Recher et al., 2004) and chronic HIV/SIV (Gray et al., 2011; Lindqvist et al., 2012; Petrovas et al., 2012) infections. Here, reduced chronic-phase peripheral CXCR3$^-$CXCR5$^+$PD-1$^+$ memory follicular Th (Tfh), linked with antibody cross-reactivity in one cohort (Locci et al., 2013), was somewhat in line with this notion (**Figure 4-figure supplement 1E** and **1F**). We additionally examined another CTL escape mutant Nef-G63R, selected with marginal statistical significance primarily in early stage (**Figure 4-figure supplement 2A**). This mutant did not associate with NAb induction in one control animal (R06-034) even upon preferential selection (**Figure 4-figure supplement 2B**) and Nef-G63R mutant was not similarly decreased in pAkt drive (**Figure 4-figure supplement 2C**), implicating that the Nef-G63E mutation was more tightly linked with NAb induction in terms of an immunosignaling phenotype.

We further explored molecular traits of this decreased Akt hyperactivation. We reasoned

that comparing endogenous Nef binding patterns would be adequate and analyzed SIV-

infected HSC-F cells with a flow cytometry-based proximity ligation assay (PLA) (Leuchowius

et al., 2009; Avin et al., 2017), by calculating a mean fluorescence intensity (MFI) deviation-

based binding index for each. We found that this Nef-G63E mutation causes significant

decrease in Nef binding to PI3K p85/p110α and downstream mTORC2 components mTOR

(Sarbassov et al., 2005) and GβL/mLST8 (Kim et al., 2003) in the CD4^{lo}-SIV Gag p27^{+} infected

population (**Figure 4D**). Results were biochemically confirmed by a decrease in G63E-

mutated Nef binding to the mTORC2-intrinsic cofactor Sin1 in coimmunoprecipitation analysis

of infected HSC-F cells (**Figure 4E**). Collectively, the Nef-G63E mutation results in decreased

aberrant Nef bridging and signal transduction of PI3K/mTORC2, explaining decreased

substrate Akt Ser473 phosphorylation by mTORC2. These properties were in keeping with an

acceleration in CD4-downregulated/infected HSC-F cell apoptosis upon Nef-G63E SIV

infection (**Figure 4F**). Thus, Nef-G63E SIV is a mutant virus decreased in aberrant

interaction/drive of B-cell-inhibitory PI3K/mTORC2 signaling, manifesting a molecular

signature reciprocal to human APDS (Angulo et al., 2013; Lucas et al., 2014; Avery et al.,

2018).

Targeting of lymph node Env-specific B cells by Nef *in vivo*

Next, we analyzed *in vivo* targeting of virus-specific B cells by Nef in lymph nodes to

explore the potential B-cell-intrinsic influence of the G63E-Nef phenotype. Previously

suggested influence of soluble Nef itself (Qiao et al., 2006) and/or related host factors

(Swingler et al., 2008) may derive generalized influence on B cells; however, SIV antigen-

specific binding antibody responses were rather decreased in NAb inducers (**Figure 1-figure**

supplement 1), viremia was comparable (**Figure 3-figure supplement 1C**) and ferritin levels

were not altered (**Figure 4-figure supplement 1A**), differing from the aforementioned

literature stating positive correlation between the three. Therefore, we surmised that some

targeted Nef intrusion against Env-specific B cells may be occurring and that the decreased

aberrant Nef-PI3K/mTORC2 drive may result in their enhanced lymph node maturation.

While reports (Xu et al., 2009) histologically proposed Nef B-cell transfer, quantitative

traits, e.g., invasion frequency and influence on virus-specific B cells, have remained unvisited.

Intracellular Nef staining is dim (particularly for B cell-acquired Nef), difficult to examine by

conventional flow cytometry. Defining staining cutline requires single-cell images, overcoming

confounding of high Nef false-staining via pre-permeation rupture and post-permeation

confounding as judged by biologically discontinuous staining and/or batch-inflated signals.

Collectively, at the expense of spatial information, we judged that sophisticating imaging

cytometry would best visualize Nef-mediated B-cell perturbation *in vivo*. We analyzed lymph

node B cells with Image Stream X MKII, with high-power (> 10-fold) antigen detection [e.g.,

molecules of equivalent soluble fluorochrome (MESF) 5 *vs.* MESF 80 in an average flow

cytometer for FITC detection] ideal for detection/single-cell verification.

We designed a triple noise cancellation strategy to overcome the above. Firstly, amine

reactive dye staining (Perfetto et al., 2006) gated out B cells being Nef-positive due to pre-

experimental membrane damage (**Figure 5-figure supplement 1A**, left). Next, we deployed

secondary quantitative parameters of Nef signals deriving from each pixel of the images.

Definition of a gray-level cooccurrence matrix (GLCM) (Haralick et al., 1973) numerating

adjacent signal deviation as their frequencies on a transpose matrix enables calculation of a

variety of feature values summarizing traits of the whole image. A biological assumption of Nef

signal continuity in a true staining suggested that the sum of weighted square variance of

GLCM, i.e., the {sum of [square of (value – average signal strength) × frequency of each value

occurrence]} would separate natural versus artificial Nef signals. This value, Haralick variance,

was computed for multiple directions and averaged, deriving Haralick variance mean (**Figure

5-figure supplement 1A**, X axis), proportionate with staining image unnaturalness. Finally,

Nef signal intensity threshold (**Figure 5-figure supplement 1A**, Y axis) gated out overtly-

stained cells void for true-false verification. These excluded B cells showing non-specific,

strong-signal binary-clustered staining pixels for Nef deriving a large summated variance (X

axis), and/or cells batch-stained for Nef (Y axis), likely originating from post-experimental

membrane damage.

With this approach we acquired images of a Live/Dead$^-$-Nef signal Haralick variance

mean^{lo}-Nef signal intensity threshold^{lo}-Nef$^+$CD19$^+$ B-cells, with fine-textured gradation of Nef^{int-}

^{lo} staining with continuity from membrane-proximal regions without sporadic staining speckles

(**Figure 5A** and **Figure 5-figure supplement 1**). A linear discriminant analysis-based machine

learning module, scoring segregation of typical void/valid images, showed this gating to

provide the highest two-dimensional separation (**Figure 5-figure supplement 1**). Combined

with biotinylated Env/streptavidin staining, this reproducibly obtained images of Env-specific

(Ch 12) CD19$^+$ (Ch 11) B cells without membrane ruptures (Ch 08) and showing fine-textured

transferred Nef (Ch 02) (**Figure 5B**). Nef invasion upregulated pAkt Ser473 (Ch 03) to a range

resembling *in vitro* analysis (**Figure 5C**), stating that Nef-driven aberrant PI3K/mTORC2

signaling does occur in Nef-invaded B cells *in vivo* and the current phenotypic change likely

affects B-cell dynamics. Strikingly, lymph node Env-specific B cells showed significantly higher

Nef-positive frequencies as compared with batch non-Env-specific B cells (**Figure 5D**). This

indicated that infected cell-generated Nef preferentially targets adjacent Env-specific B cells,

putatively through infected CD4$^+$ cell-to-B-cell contact/transfer (Xu et al., 2009; Hashimoto et

al., 2016) and that a Nef phenotypic change likely dictates Env-specific B-cell maturation.

Cell contact-dependent B-cell invasion from infected cells *in vitro*

To understand cell-intrinsic properties enhancing B-cell Nef acquisition *in vivo*, we

addressed in an *in vitro* coculture reconstitution how infected cell-to-B-cell Nef invasion may

take place or be modulated. We performed imaging cytometry on a twelve-hour coculture of

SIV-disseminating HSC-F cells (reaching around 30% Nef-positivity at moment of coculture)

with Ramos B cells (**Figure 6A**, top left), independent of MHC-related interactions known to

enhance T-cell/B-cell contact (Wülfing et al., 1998). Machine learning-verified noise

cancellation of fragmented signal acquisition (**Figure 6A**, middle left) filtered reproducible

images of Nef-positive HSC-F cells directly adhering to Ramos B cells within the doublet

population (**Figure 6A**, top to middle right). Importantly, addition of a soluble macaque CD3

stimulation-competent antibody resulted in enhanced Nef+ T-cell adhesion to B cells (**Figure**

6A, bottom center), accompanying images of polarized Nef protrusion/injection from T cells to

B cells (second column, filled wedges) as well as trogocytosed B-cell membranes (third

column, open wedge), in keeping with dynamic B-cell Nef acquisition. Centrifugation strengths

upon cell collection and staining (190 x g versus 1,200 x g and 600 x g versus 1,200 x g,

respectively) shifted the tendency between enriched Nef+ HSC-F-B-cell doublet versus Nef+

B-cell singlet detection via CD3 stimulation, respectively (**Figure 6A**, lower right).

Substantiated by the acquired images, an enhancement of live singlet Ramos B-cell Nef

acquisition (Nef+ frequencies/total) was also observed by CD3 stimulation in conventional flow

cytometry (**Figure 6B**). We observed a similar enhancement by phytohemagglutinin (PHA)

stimulation whereas phorbol-12-myristate-13-acetate (PMA)/ionomycin did not enhance B-cell

Nef acquisition. In a similar 72-hour static infected cell coculture with primary CD20+ B cells,

we detected recapitulation of the decreased B-cell pAkt Ser473 phosphorylation upon G63E

Nef transfer compared to that of WT Nef (**Figure 6C**), implicating that the Nef-G63E mutation

can directly alleviate maturation inhibition in preferentially Nef-targeted Env-specific B cells.

Enhanced Env-specific B-cell responses after PI3K-diminuting mutant selection

Finally, peripheral SIV Env-specific IgG+ B-cell responses comprising plasmablasts

(PBs) and memory B cells (B_{mem}) were analyzed to assess *in vivo* B-cell quality in NAb

inducers with Nef-G63E mutant selection. After excluding lineage-specific cells [T/NK/pro-B

cells/monocytes/myeloid dendritic cells (DCs)/plasmacytoid DCs], we defined IgG+ PBs as

showing: high replication (Ki-67+), post-activation (HLA-DR+), transcriptional switching for

terminal differentiation (IRF4hi) and downregulated antigen surface binding (surface [s]Envlo-

cytoplasmic [Cy]Env+) (Nutt et al., 2015; Silveira et al, 2015). Env-specific PBs were defined

as CD3-CD8-CD10-CD14-CD16-CD11c-CD123-CD19-CyIgG+Ki-67+IRF4hiHLA-DR+ (total PB)

sEnvloCyEnv+ (Env-specific) (**Figure 7A** and **Figure 7-figure supplement 1**). NAb inducers

showed significantly higher Env-specific IgG+ PB responses around week 30 p.i., after Nef-

G63E selection, compared to those in NAb non-inducers (**Figures 7B** and **7C**). At 1 year, this

difference became enhanced; Env-specific PB frequencies were reduced in non-inducers, while this reduction was ameliorated in the NAb inducers with Nef-G63E.

Concomitant determination of Env-specific $IgD^-CD27^+IgG^+$ B_{mem} responses allowed pair-wise analysis of Env-specific IgG^+ B_{mem}/PB as a projection of germinal center (GC) output as an indicator of overall B-cell response quality (Zotos et al., 2010) (**Figure 7D**, left). In this two-dimensional vector charting of Env-specific B_{mem}/PB responses, vector protrusion towards upper right reasonably represents higher gross GC output of antibody-forming cells (AFCs) (B_{mem}/PBs). In NAb non-inducers, all vectors converged on an empirically defined polygonal attractor area D_n (gray area surrounded with dotted lines in **Figure 7D**, right) at year 1 p.i. and beyond, describing that these NAb non-inducers failed to sustain Env-specific GC output. In contrast, the vectors were consistently tracked outside D_n in the NAb inducers with Nef G63E (**Figure 7D**). At the moment of NAb induction, they converged on another upper-right GC output attractor area D_i (red area surrounded with dotted lines in **Figure 7D**, right), mutually exclusive with D_n ($P < 0.0001$ by Fisher's exact test on NAb non-inducer/inducer vector distribution frequency within D_n). These suggest more robust virus-specific IgG^+ B-cell responses following Nef-G63E $CD8^+$ T-cell escape mutant selection, predisposing to NAb induction. This enhanced cognate B-cell signature was also an inverted pattern of impaired terminally-differentiated B-cell responses in APDS (Al Qureshah et al., 2021).

Taken together, in the current model, Nef_{62-70}-specific $CD8^+$ T-cell responses in persistently SIV_{mac239}-infected macaques selected for an escape mutant, Nef-G63E SIV, which decreased aberrant Nef binding and drive of B-cell-inhibitory PI3K/mTORC2. B-cell Nef invasion *in vivo* occurred more preferentially in Env-specific B cells, suggesting diminution in tonic Nef-mediated B-cell dysregulation after mutant selection. These predisposed to enhanced Env-specific B-cell responses and subsequent SIV_{mac239}-specific NAb induction, altogether, in a manner reciprocal to human APDS-mediated immune dysregulation.

Discussion

In the present study, we found NAb induction against an NAb-resistant SIV after selection of a CD8$^+$ T-cell escape *nef* mutant virus. The Nef-G63E mutation, positioned in cognate B cells, results in reduction of binding/drive of PI3K/mTORC2, being concordant with *in vivo* enhancement of Env-specific IgG$^+$ B cells up to NAb induction following mutant selection. Importantly, this manifested through a pAkt deviation level more vivid than wild-type versus germline *PIK3CD* gain-of-function mutation heterozygote mice recapitulating full human APDS phenotype (Avery et al., 2018). Thus, the current Nef-G63E-associated B-cell/NAb phenotype likely occurs in a "reciprocal APDS-like" manner. This proposition was enhanced based on extension of immune cell-intrinsic Nef influence on cognate B cells (Fig. 5), in addition to infected T cells. This work is, to our knowledge, the first to interlink a PI3K/mTORC2-modulating viral signature and enhanced B-cell/NAb responses in a viral infection model.

A link between viral T-cell escape and consequent immune modulation has been previously partially explored, including enhanced mutant HIV-1 epitope peptide binding to inhibitory MHC-I receptor on DCs impairing T cells (Lichterfeld et al., 2007) and decreased CTL-mediated lymph node immunopathology rather driving LCMV-specific antibody production (Battegay et al., 1993). Our results now evidence a new pattern of NAb responses bivalently shaped through viral interactions with both humoral and cellular immunity in AIDS virus infection. SIV$_{mac239}$ infection of macaques possessing MHC-I alleles associated with Nef-G63E mutation can be one unique model to analyze virus-host interaction for B cell maturation leading to NAb induction. The time frame of Nef-G63E mutant selection versus moment of NAb induction was variable in the inducers, for which we did not obtain a clear explanation. This may be influenced by certain basal competitive balance between humoral versus cellular adaptive immunity, as observed in certain vaccination settings (Querec et al., 2009).

Identification of the current Nef-G63E candidate phenotype adds one aspect to the wealth of evidence documenting the multifaceted impact of Nef in HIV/SIV infection (Kirchhoff

et al., 1995; Gauduin et al., 2006; Stolp et al., 2012). It appeared unique that the current

phenotype differed from decreased replication-related properties or generalized amelioration

in immune impairment such as those documented in delta-Nef HIV/SIV infection (Kirchhoff et

al., 1995; Johnson et al., 1997; Gauduin et al., 2006; Fukazawa et al., 2012). The SIV Nef N-

terminal unstructured region comprising Nef-G63E is not conserved in HIV-1 (Schindler et al.,

2004) and beyond this model study it remains to be addressed whether a mutant HIV-1 with a

similar immunosignaling-affecting phenotype can be obtained and how much lentiviral

conservation exists for such interactions. Cognate B-cell maturation analysis on cohort basis

(Hau et al., 2022) may potentially assist this approach.

Obtained results on preferential Nef transfer to Env-specific B cells (**Figure 5**) implicate

involvement of polarized cognate immune cell perturbation *in vivo*, best depicted for

preferential infectivity of HIV-specific CD4$^+$ T cells (Douek et al., 2002) and HIV condensation

to the dendritic cell-T-cell interface (McDonald et al., 2003). As cognate MHC class II

interactions (Wülfing et al., 1998; Giltin et al., 2014) regularly occur between HIV/SIV-specific

CD4$^+$ T cells and Env-specific B cells in the germinal center, results may show that Env-

specific B-cell dysregulation stems from a negative consequence of exploited cognate T-

cell/B-cell orchestration. Enhanced T-cell attachment and transfer of Nef to B cells *in vitro* by

CD3 activation (**Figure 6**), mimicking peptide/MHC (pMHC) signaling, may facilitate recovery

from viral T-cell cytoskeletal impairment (Campbell et al., 2004; Jolly et al., 2004; Stolp et al.,

2009) antagonizing reduced motility (Stolp et al., 2012), resulting in *in vivo* "filtering" of

enhanced cognate CD4$^+$ T-cell interaction with cognate B cells. Combined with recent *in vivo*

detection of trogocytosed CD20-bearing follicular CD4$^+$ T cells positive for viral DNA and

showing increase during SIV infection (Samer et al., 2023), results suggest occurrence of

contact-dependent, polarized interactions facilitating Nef transfer to cognate B cells.

The major *in vivo* readout of this study is autologous neutralization of a highly difficult-

to-neutralize SIV, differing from HIV-specific broadly neutralizing antibodies (bNAbs).

Importantly, however, obtained signatures of enhanced IgG$^+$ Env-specific AFC B cells (class

switch/terminal differentiation) and virus neutralization (hypermutation) directly evidence

enhanced activity of activation-induced cytidine deaminase (AID), the canonical positive driver

of B-cell fate, showing a strong conceptual continuity with bNAb regulation by AID.

Our results extracted an *in vivo* link between a decreased aberrant Nef-PI3K/mTORC2

interaction and major enhancement in B-cell responses. The relationship is indeed reciprocal

to immunogenetic mechanisms of human PI3K gain-of-function mutations resulting in APDS

with multiply impaired anti-viral B-cell responses. APDS occurs from germline mutations in

leukocyte-intrinsic p110δ catalytic subunit-coding *PIK3CD* or the ubiquitous p85α regulatory

subunit-coding *PIK3R1*, both deriving class IA PI3K gain-of-function residue substitutions

(Dornan et al., 2017). Such mutations lead to immune cell-intrinsic perturbations with cognate

B-cell blockade, manifesting as impaired class-switched B-cell responses and susceptibility

against various infections (Angulo et al., 2013; Lucas et al., 2014; Coulter et al., 2017). In the

current work, we conversely document a decrease in virally induced tonic PI3K drive,

enhanced IgG class-switched cognate B-cell responses and NAb induction against a notably

difficult-to-neutralize SIV. While the exact hierarchy of binding between Nef and

PI3K/mTORC2 components (including PI3K isoforms) and its perturbation by Nef-G63E

mutation remains to be investigated, PI3Kp85 did reasonably show the strongest Nef-binding

index in PLA assay (**Figure 4D**) as Nef and PI3Kp85 possess SH3-binding PxxP and SH3

domains, respectively (Rickles et al., 1994). Binding decrease in canonical partner PI3Kp85

may domino-trigger decrease with others, resulting in quintuple PI3K/mTORC2 binding

decrease and fragile downstream Akt signaling. Such domino-triggering is also observed from

gain-of-function-mutated p85 to p110 evoking its activation (Dornan et al., 2017) similarly

highlighting the fine-tuned nature of the pathway.

A limitation of this study is the use of retrospective samples, posing constraints for

detecting exact temporal changes in tonic viral B-cell perturbation. Related with this, sampling

of lymph node cells in animals belonging to the subgroup of interest at the optimal moment

was unpredictable. Certain plasma and cellular samples were also unavailable in differing

experiments. While partially addressed in *in vitro* reconstitution, much remains to be

investigated for the observed polarity of Env-specific B-cell perturbation *in vivo*. Similarly,

influence of the Nef-G63E phenotype on CD4$^+$ T cells *in vivo* and its consequence on B-cell

modulation remains largely elusive, and in the current work we focused mainly on molecular

properties of the NAb-correlating Nef-G63E mutant strain. As HIV/SIV infection uniquely

shows a biphasic Tfh dysregulation of acute-phase destruction (Mattapallil et al., 2005;

Moukambi et al., 2015; Moukambi et al., 2019) followed by dysregulated chronic-phase hyper-

expansion (Lindqvist et al., 2012; Petrovas et al., 2012; Cubas et al., 2013; Xu et al., 2015), it

is of interest how the PI3K drive-ameliorated and proapoptotic mutant influenced this axis

post-*in vivo* selection, warranting further studies ahead. While influence on B cells appears to

be stronger by unexplained reasons (Asano et al., 2018), it is noteworthy that PI3K gain-of-

function also evokes basal dysregulation in peripheral CD4$^+$ T cells, locally recapitulating the

consequence of immunodeficiency virus replication and implicating importance of this

signaling axis in HIV/SIV infection. In any event, usage of specific animal models does help to

depict a temporal cascade of *in vivo* events leading to NAb induction, assisting human

immunology. As future extension of this study extracting the NAb-involved molecular axis,

manipulation/reconstitution experiments shall be designed.

In conclusion, we exemplify in a non-human primate AIDS model that NAb induction

against a highly resistant SIV occurs after selection of a CD8$^+$ T-cell escape variant reduced

in excess PI3K/mTORC2 drive. These results provide an understanding of how

immunodeficiency virus-specific NAb responses can be shaped by both wings of adaptive

immune pressure and depicts that immune cell-intrinsic PI3K/mTORC2 manipulation may be

pivotal for harnessing antiviral NAb responses. As human IEIs are increasingly becoming

discovered with various immune-perturbing phenotypes and inheritance patterns (Jauch et al.,

2023), search of other types of analogies between viral immune dysregulations and human

IEIs may lead to identification of novel targets to modulate and harness immune responses in

translational settings.

Acknowledgments

This work was supported by Japanese Initiative for Progress of Research on Infectious

Diseases for Global Epidemic (J-PRIDE) (JP19fm0208017 to H.Y.), Japan Program for

Infectious Diseases Research and Infrastructure (Interdisciplinary Cutting-edge Research)

(JP22wm0325006 to H.Y.), Research Programs on HIV/AIDS, Emerging and Re-emerging

Infectious Diseases, and Global Health Issues (JP20fk0410022 and JP24fk0410066 to H.Y.;

JP18fk0410003, JP20fk0410011, JP20fk0108125, JP20jm0110012, JP21fk0410035, and

JP21jk0210002 to T.M.) from Japan Agency for Medical Research and Development, the

Ministry of Education, Culture, Sports, Science, and Technology in Japan ([JSPS] KAKENHI

Grants 17H02185, 18K07157 and 24K21287 to H.Y.; 21H02745 to T.M.), the Takeda Science

Foundation (to H.Y.), the Imai Memorial Trust for AIDS Research (to H.Y.), and the Mitsui-

Sumitomo Insurance Welfare Foundation (to H.Y.). We thank Fumiko Ono, Koji Hanari,

Katsuhiko Komatsuzaki, Akio Hiyaoka, Hiromi Ogawa, Keiko Oto, Hirofumi Akari, Yasuhiro

Yasutomi, Hiromi Sakawaki, Tomoyuki Miura and Yoshio Koyanagi for animal experiment

assistance; Hiroyuki Kogure, Sayuri Seki and Julia R. Hirsiger for imaging cytometry technical

assistance; Masako Nishizawa, Trang Thi Thu Hau, Shigeyoshi Harada, Takushi Nomura,

Akiko Takeda, Taku Nakane, Nami Iwamoto, Taeko K. Naruse, Akinori Kimura and Mark S. de

Souza for their help. H.Y. thanks Jun Abe, Shoi Shi, Bettina Stolp, Jens V. Stein, Makoto

Yamagishi and Yusuke Yanagi for conceptual discussions, and Mike Recher for collaborative

support.

Author Contributions

H.Y. conceived/designed/performed study and wrote manuscript; T.M. provided animal cohort,

co-edited manuscript and co-directed study.

CRediT taxonomy:

Conceptualization: HY

Methodology: HY

Validation: HY

Formal analysis: HY

Investigation: HY

Resources: HY, TM

Data curation: HY

Writing—original draft: HY

Writing—review & editing: HY, TM

Visualization: HY

Supervision: HY, TM

Project administration: HY

Funding acquisition: HY, TM

Declaration of Interests

The authors declare no competing interests.

Figure Legends

Figure 1. NAb induction against NAb-resistant SIV$_{mac239}$.

(A) Study design.

(B) Plasma viral loads (SIV *gag* RNA copies/ml plasma) in NAb non-inducers (left) and inducers (right).

(C) Plasma SIV$_{mac239}$ 100% neutralizing end point titers by 10 TCID$_{50}$ killing assay on MT4-R5 cells. Points on the dotted line show marginally NAb-positive results ($< 1:2$). In some animals, titers were comparable with our reported results using MT4 cells.

Figure 2. Selection of a viral *nef* mutation (Nef-G63E) before NAb induction.

(A) Viral *nef* mutations in NAb inducers. A linear schema indicating Nef functional domains is aligned above. Mutations at timepoints with NAb$^-$ (indicated by white wedge; mostly before 6 months p.i.), NAb$^{+/-}$ (green wedge) and NAb$^+$) (purple wedge; mostly after 1 year) are shown in individual animals. Black and dark gray represent dominant and subdominant mutations (or deletion in R-360) by direct sequencing, respectively. Red and pink indicate dominant and subdominant G63E detection, respectively.

(B) Schema of SIV$_{mac239}$ Nef structure and Nef-G63E mutation orientation.

(C) Comparison of frequencies of macaques having Nef-G63E in plasma viruses between NAb non-inducers and inducers. Compared by Fisher's exact test.

(D) Temporal relationship of Nef G63E frequencies in plasma virus and NAb induction. Black boxes (left Y axis) show log$_2$ NAb titers; red triangles and blue diamonds (right Y axis) show percentage of G63E and G63R mutations detected by subcloning (15 clones/point on average), respectively.

Figure 3. Nef-G63E is a CD8$^+$ T-cell escape mutation.

(A) Nef$_{62-70}$ QW9-specific CD8$^+$ T-cell frequencies and related MHC-I alleles in seven NAb inducers selecting Nef-G63E. Mamu-B*039:01 and Mamu-B*004:01 are known to restrict

Nef$_{62-70}$ QW9 epitope and binding of Nef$_{63-70}$ peptide to Mamu-A1*032:02 was predicted.

(B) CD8$^+$ T-cell responses specific to wild-type Nef$_{62-70}$ or mutant Nef$_{62-70}$.G63E or Nef$_{62-}$

$_{70}$.G63R peptides.

Figure 4. Nef-G63E mutation reduces PI3K/mTORC2 binding and pAkt drive.

(A) Representative surface expression level histograms of CD3, CD4, MHC class I, CXCR4

and BST-2 in CD4^{lo}Nef$^+$ subpopulations after wild-type (WT) or Nef-G63E mutant SIV infection

at multiplicity of infection (MOI) 0.1 on HSC-F cells.

(B) Left: representative histograms of relative pAkt serine (Ser) 473 levels in p27$^+$

subpopulations after WT or Nef-G63E mutant SIV infection at MOI 0.1 on HSC-F cells.

Numbers show pAkt Ser473 mean fluorescence intensities (MFIs) for each. Right: Deviation

of pAkt Ser473 MFIs in p27$^+$ HSC-F cells compared with mean MFI of uninfected cells.

Compared by unpaired t test.

(C) Left: Relative pAkt Ser473 levels (normalized to mean MFI of uninfected controls) in Nef$^+$

HSC-F cells assessed for serum starvation (MOI 0.2, 1 day p.i.). Adjusted *P* values show

results of comparison via Sidak's post-hoc test of 2-way ANOVA (C, left). Right: Deviation of

pAkt Ser473 MFIs in Nef$^+$ HSC-F cells assessed for high-MOI infection (MOI 5, 1-day p.i.).

Compared by unpaired t test.

(D) Proximity ligation assay (PLA) of Nef binding with mTOR, GβL/mLST8, PI3Kp85 and

PI3Kp110α. MFI-based binding index was calculated as (anti-Nef/anti-partner) - (isotype/anti-

partner) - (anti-Nef/isotype) + (isotype/isotype). Histograms for samples and isotype/isotype

are representatively shown. Differences in MFI binding indexes can be enhanced compared

with comparison of raw MFIs. Compared by paired t tests.

(E) Sin1 co-immunoprecipitation analysis of WT versus Nef-G63E mutant SIV-infected HSC-

F cells (infected at MOI 0.05, day 3). Immunoblotting of Nef (37 kDa) in whole cell lysates

(lanes 4-6) and anti-Sin1 antibody immunoprecipitates (lanes 1-3) are shown.

(F) Cell death frequencies of SIV-infected cells measured by Annexin V positivity (% of CD4^{lo})

(infected at MOI 0.1, day 3). Compared by unpaired t test.

Data represent one of two [(A); (B); (C, right)] independent experiments in quadruplicate, four

independent experiments performed in triplicate [(C, left)], four independent single-well

comparison experiments pooled for statistical analysis (D), two experiments (E) or two

experiments performed with six wells/control (F). Bars: mean $\pm$ SD (B, C right), mean $\pm$ SEM

(C left, F).

Figure 5. Preferential targeting of lymph node Env-specific B cells by Nef *in vivo*.

(A) Representative gating of triple noise-cancellation *in vivo* Nef staining in B cells analyzed

by ImageStreamX MKII. Pre-experimental damaged cells are first excluded with Live/Dead

from focused/centroid/singlet image-acquired CD19$^+$ B cells (first lane right/R4 gated on

"Focused"). Following Nef$^+$ gating (second lane left/R5 gated on R4), a second step of Nef

noise cancellation (second lane middle/R7 on R5) comprises double-negative removal of post-

experimental stochastic irregular staining building a disparate intracellular staining gradient (X

axis, Nef signal pixel Haralick variance mean) and post-experimental batch overt cellular

staining (Y axis, Nef signal pixel intensity threshold). This outputs a B-cell population with a

fine-textured pericellular Nef^{int-lo} staining, biologically concordant with Nef membrane-

anchoring. Probing of anti-Env BCR (αEnv) by recombinant SIV Env (second lane, right) is

combined, resulting in a 2-D panel of intracellular Nef versus αEnv for noise-cancelled Nef$^+$ B

cells (R7) plus Nef$^-$ B cells (R8) (third lane left/"R7+R8"). pAkt Ser473 expression (third lane

right) and cellular morphology (B) was further analyzed. DN, double-negative.

(B) Typical images of Nef-transferred Env-specific B cells defined αEnv$^+$-intracellular Nef$^+$-Nef

Haralick variance mean^{lo}-Nef Intensity threshold^{lo}-Live/Dead$^-$-CD19$^+$ cells ["Nef$^+$aEnv$^+$"

population of lower left panel in (A), gated on "R7, R8"]. Note the pericellular pAkt Ser473

upregulation in these cells (Ch 03/yellow). Data on inguinal lymph node lymphocytes of

macaque R10-007 at week 62 post-SIV$_{mac239}$ infection are shown in (A) and (B).

(C) Comparison of pAkt Ser473 median fluorescence (medFI) intensity levels in Nef$^-$ B cells

(R8) versus noise-cancelled Nef$^+$ B cells (R7). Analyzed by paired t test.

(D) Comparison of Nef-positive cell frequencies in non-Env-specific (left) versus Env-specific

(right) B cells in lymph nodes of persistently SIV-infected macaques (n = 6). Analyzed by paired

t test.

Figure 6. Contact-dependent enhancement of B-cell Nef acquisition in reconstitution.

(A) Imaging flow cytometric analyses of stimuli-dependent enhancement of infected HSC-F

cell adhesion to non-permissive CD19$^+$ Ramos B cells with the indicated gating (left).

Cocultured doublet cells subjected to machine learning-verified filtration of fragmented signal

acquisition (via gating Nef pixel signal contrast^{lo}-Nef mean pixel per object^{hi} cells) deriving

typical images of infected HSC-F cell (Nef stain shown in pale blue pseudocolor) adhesion to

Ramos cells (CD19 stain shown in purple pseudocolor) are shown. Analyzed by unpaired t

test for doublet formation in the unstimulated versus anti-CD3 monoclonal antibody (clone FN-

18)-stimulated group.

(B) Flow cytometric analyses of Nef-acquiring live CD19$^+$ Ramos B cells upon coculture with

SIV-infected HSC-F cells with or without the indicated stimuli. Analyzed by one-way ANOVA

with Tukey's post-hoc multiple comparison tests.

(C) Representative flow cytometric plot (left), histogram (middle) and pAkt Ser473 signal

deviation in wild type versus G63E Nef-acquiring CD20$^+$ primary B cells (six wells/control).

Analyzed by unpaired t test. Data represent pooled data of two (A) or three [(B) and (C)]

independent experiments with indicated number of replicates showing similar results.

Figure 7. Enhanced SIV Env-specific B-cell output up to NAb induction following Nef-

G63E selection.

(A) Representative gating (R02-004, week 32 p.i.) of SIV Env gp140-specific memory B cells

(B$_{mem}$) and plasmablasts (PBs). Two panels for PB gating are shown in lower resolution for

visibility. B$_{mem}$ staining was performed separately and gating are merged with PB panels

(second row, panels 2–4). For lineage-specific exclusion (first row, panel 4), CD10 was stained

independently for B$_{mem}$ (see parenthesis).

(B) Changes in SIV Env gp140-specific plasmablast (PB) frequencies in viremic NAb non-

inducers (left, *n* = 12) and Nef-G63E-selecting NAb inducers (right, *n* = 6). Available samples

of twelve viremic NAb non-inducers (including ten with ≥ 1-year survival) and six NAb inducers

were tracked. In the right panel, the frequencies in NAb non-inducers are shown in background

(gray) for comparison.

(C) Comparison of Env gp140-specific PB frequencies between NAb non-inducers and

inducers by Mann Whitney U tests. In the right, the frequencies at NAb induction were

compared with those at year 1 in NAb non-inducers with ≥ 1-year survival (*n* = 10). Bars: mean

± SD.

(D) Left: vector chart of Env gp140-specific memory B cell (B_{mem}) and PB levels. Legends for

each animal correspond to the ones in (B). Right: NAb non-inducer vectors empirically define

a polygonal GC output attractor area D_n (gray area surrounded with dotted lines) on which

they converge by and beyond year 1 p.i. The NAb inducer vectors (shown up to the time of

NAb induction) remained outside of D_n. At the moment of NAb induction they converged on a

second GC output attractor area D_i (red area surrounded with dotted lines), mutually exclusive

with D_n ($P < 0.0001$ by Fisher's exact test on NAb inducer vector convergence frequency within

D_n). Legends for year 1 p.i. in the NAb non-inducers and moment of NAb induction in the NAb

inducers are specified.

Materials and Methods

Material availability

This study did not generate new unique reagents.

Data and code availability

Data were analyzed using existing computational packages. The accession number for the transcriptome data [wild-type SIV_{mac239}-infected HSC-F cell line (n = 3), Nef-G63E SIV_{mac239}-infected HSC-F cell line (n = 3) and uninfected HSC-F cell line (n = 3)] has been deposited under the accession number GenBank: GSE65806.

Rhesus macaques retrospectively utilized for samples

Seventy Burmese rhesus macaques (*macaca mulatta*) (57 males and 13 females) were retrospectively analyzed in this study. Experiments were previously carried out (Matano et al., 2004; Yamamoto et al., 2007; Iseda et al., 2016; Nomura et al., 2012; Ishii et al., 2012; Takahashi et al., 2013; Nakane et al., 2013; Shi et al., 2013; Iwamoto et al., 2014; Terahara et al., 2014) in the Tsukuba Primate Research Center, National Institutes of Biomedical Innovation, Health and Nutrition (NIBIOHN) with the help of the Corporation for Production and Research of Laboratory Primates and the Institute for Frontier Life and Medical Sciences, Kyoto University (IFLMS-KU) after approval by the Committee on the Ethics of Animal Experiments of NIBIOHN and IFLMS-KU under the guidelines for animal experiments at NIBIOHN, IFLMS-KU and National Institute of Infectious Diseases in accordance with the Guidelines for Proper Conduct of Animal Experiments established by Science Council of Japan (http://www.scj.go.jp/ja/info/kohyo/pdf/kohyo-20-k16-2e.pdf). The experiments were in accordance with the "Weatherall report for the use of non-human primates in research" recommendations (https://royalsociety.org/topics-policy/publications/2006/weatherall-report/). Animals were housed in adjoining individual primate cages allowing them to make sight and sound contact with one another for social interactions, where the temperature was kept at

25°C with light for 12 hours per day. Animals were fed with apples and commercial monkey

diet (Type CMK-2, Clea Japan, Inc.). Blood collection and virus challenge were performed

under ketamine anesthesia.

Cells and viruses

MT4-R5 cells were maintained in RPMI1640 (Invitrogen) supplemented with 10% fetal bovine

serum (FBS) (Clontech) and antibiotics. HSC-F cells (cynomolgus CD4$^+$ T-cell line) and

HSR5.4 cells (rhesus macaque CD4$^+$ T-cell line) were maintained in RPMI1640 (Invitrogen)

supplemented with 10% fetal bovine serum (FBS) (Clontech), human interleukin-2 (IL-2) (10

IU/ml, Roche), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (Invitrogen) and

2-mercaptoethanol (Gibco). SIV$_{mac239}$ molecular clone derivative was generated by

mutagenesis PCR (Agilent Technologies) using the primers listed. For virus preparation, COS-

1 cells were transfected with pBR$_{mac239}$ proviral DNA using FuGENE 6 (Promega). At 48 h post-

transfection, culture supernatants were harvested, centrifuged and filtered through a 0.45-μm

pore-size filter (Merck Millipore). To titrate infectivity, prepared viruses were serially diluted and

infected on HSC-F cells in 96-well plates (Falcon) in quadruplicate. At 10 days post-infection,

the endpoint was determined using SIV p27 antigen enzyme-linked immunosorbent assay

(ELISA) kit (ABL), and virus infectivity was calculated as the 50 percent tissue culture infective

dose (TCID$_{50}$) according to the Reed-Muench method.

Identification of SIV$_{mac239}$-NAb inducers

Burmese rhesus macaques previously challenged with the highly pathogenic molecular clone

virus SIV$_{mac239}$ (n = 70) were retrospectively examined approximately up to 2 years for their

plasma NAb profiles. Animals were challenged intravenously with 1,000 TCID$_{50}$ of SIV$_{mac239}$.

In the current study, virological and immunological profiles were compared between the newly

identified NAb inducers (n = 9) and representative NAb non-inducers (n = 19). Sex distribution

of the NAb inducers (eight males and one female) did not significantly differ with the total non-

inducers (49 males and 12 females, P = 0.99 by Fisher's exact test). These NAb non-inducers

and eight NAb inducers except R06-032 were previously partially reported for their plasma

viral loads (Yamamoto et al., 2007; Iseda et al., 2016; Nomura et al., 2012; Takahashi et al.,

2013; Nakane et al., 2013; Iwamoto et al., 2014) (**Figure 1B**). MHC-I haplotypes and alleles

were determined by reference strand-mediated conformation analysis, PCR-SSP (PCR

amplification utilizing sequence-specific priming) and cloning as described (Naruse et al.,

2010). MHC-I binding prediction was made on April 27, 2013 using the IEDB analysis resource

NetMHCpan tool (Hoof et al., 2009). Alleles of interest in the study have been previously

identified in macaques (Nomura et al., 2012; Evans et al., 1999; Sette et al., 2012). Macaques

R06-032, R03-015, R01-010 and R05-010 received a prime-boost vaccination (Matano et al.,

2004) composed of a DNA prime/intranasal Sendai virus vector expressing SIV_{mac239} Gag

(SeV-Gag). R03-015, R06-019, R06-038 and R10-001 received 300 mg of non-specific rhesus

IgG at day 7 post-SIV_{mac239} challenge as an experimental control in our previous reports

(Yamamoto et al., 2007; Iseda et al., 2016; Nakane et al., 2013).

Plasma viral load quantitation

Plasma viral RNA samples were extracted with High Pure Viral RNA kit (Roche Diagnostics).

Serial fivefold sample dilutions were amplified in quadruplicate by reverse transcription and

nested PCR using SIV_{mac239} *gag*-specific primers to determine end point via the Reed-Muench

method as described previously (Matano et al., 2004; Iseda et al., 2016). The lower limit of

detection is approximately 400 viral RNA copies/ml plasma. Viral loads have been previously

partially reported (Yamamoto et al., 2007; Nomura et al., 2012; Takahashi et al., 2013; Nakane

et al., 2013; Iwamoto et al., 2014).

SIV_{mac239}-specific neutralization assay

NAbs were titrated as described (Yamamoto et al., 2007; Iseda et al., 2016). Serial twofold

dilutions of heat-inactivated plasma, or polyclonal IgG affinity-purified with Protein G

Sepharose 4 Fast Flow (GE Healthcare) from heat-inactivated and filtered plasma were mixed

with 10 $TCID_{50}$ of SIV_{mac239} at a 1:1 ratio (5 µl:5 µl) in quadruplicate. After 45-min incubation at

room temperature, the 10 μl mixtures were added into 5 x 10^4 MT4-R5 cells/well in 96-well

plates. Progeny virus production in day 12 culture supernatants was examined by SIV p27

ELISA (ABL) to determine 100% neutralizing endpoint. The lower limit of titration is 1:2.

Neutralization in three out of four wells at a dilution of 1:2 is defined as marginally NAb-positive

(< 1:2). Results were comparable when the same assay was performed with macaque HSC-

F cells (Akari et al., 1996) as targets. NAb inducers R01-012, R02-004, R01-010, R02-007

and NAb non-inducer R01-011 were previously partially reported for their NAb titers measured

with the same method using MT4 cells as targets (Kawada et al., 2007), which derived

comparable results. For assessment of neutralizing activity in IgG, SIV$_{mac239}$-specific IgGs

purified from pools of plasma with SIV$_{mac239}$-specific NAb titers were obtained from each animal

as described (Yamamoto et al., 2007). After complement heat-inactivation at 56°C, 30 min and

0.45 μm filtration, IgGs were purified by Protein G Sepharose 4 Fast Flow (GE Healthcare)

and concentrated by Amicon Ultra 4, MW 50,000 (Millipore) to 30 mg/ml and similarly

examined for their 10 TCID$_{50}$ SIV$_{mac239}$ killing titers on MT4-R5 cells.

SIV Env-specific IgG ELISA and immunoblotting

Plasma Env-specific IgG titers were measured as described (Nakane et al., 2013). SIV$_{mac251}$

Env gp120 (ImmunoDiagnostics) were coated on 96-well assay plates (BD) at 1000 ng/ml

(100 μl/well). Wells were prewashed with phosphate-buffered saline (PBS), blocked with 0.5%

bovine serum albumin (BSA)/PBS overnight and plasma samples were incubated at a 1:20

dilution (5 μl:95 μl) for 2 hours. Wells were washed with PBS and SIV Env-bound antibodies

were detected with a horseradish peroxidase (HRP)-conjugated goat anti-monkey IgG (H+L)

(Bethyl Laboratory) and SureBlue TMB 1-Component Microwell Peroxidase Substrate (KPL).

Absorbance at 450 nm was measured. Samples from week 0 pre-challenge and month 3,

month 6 and year 1 post-challenge were assessed in duplicate. Week 0 average values were

subtracted from corresponding later time point values for calibration. For immunoblotting, SIV

virion-specific IgGs in plasma were detected with a SIV$_{mac239}$-cross-reactive western blotting

system (ZeptoMetrix). In the NAb non-inducers, samples from those close to rapid progression

(euthanized due to AIDS progression within approximately 1 year) known for low plasma anti-

SIV reactivity (Hirsch et al., 2004; Nakane et al., 2013) were not included.

Sequencing

Sequencing was performed as described (Matano et al., 2004; Iseda et al., 2016). Viral cDNA

fragments spanning from nt (nucleotide) 4829 to nt 7000, nt 6843 to nt 8831 and nt 8677 to nt

10196 in SIV_{mac239} (GenBank accession number MM33262) covering SIV *env* and *nef* were

amplified from plasma viral RNA by nested RT-PCR using Prime-Script one-step RT-PCR kit

v2 (TaKaRa) and KOD-Plus v2 (Toyobo). PCR products were either directly sequenced or

subcloned with a TOP10-transforming TOPO blunt-end cloning system (Invitrogen).

Sequencing was performed using dye terminator chemistry with an ABI 3730 DNA sequencer

(Applied Biosystems). On average, 15 clones were obtained per sample and 20 clones were

assessed when *nef* mutations of interest in early time points were subdominant.

SIV_{mac239}-specific CD8$^+$ T-cell responses

Virus-specific CD8$^+$ T-cell frequencies were measured as described (Matano et al., 2004;

Iseda et al., 2016). Peripheral blood mononuclear cells (PBMCs) were cocultured for 6 hours

with autologous herpesvirus papio-immortalized B lymphoblastoid cell lines (B-LCLs) pulsed

with Nef peptides (Sigma-Aldrich Japan) at 1 µM concentration or as indicated otherwise

under GolgiStop (monensin, BD) presence. Intracellular gamma interferon (IFN-γ) staining

was performed using Cytofix/Cytoperm kit (BD) and the following conjugated anti-human

monoclonal antibodies (mAbs): anti-CD4-FITC (M-T477, BD Pharmingen), anti-CD8-PerCP

(SK1, BD Biosciences), anti-CD3-APC (SP34-2, BD Pharmingen) and anti-IFN-γ-PE (4S.B3,

BioLegend). Specific CD8$^+$ T-cell frequencies were determined by subtracting nonspecific IFN-

γ^+ CD8$^+$ T-cell frequencies from those after peptide-specific stimulation; frequencies beneath

100 cells/million PBMCs were considered negative. Cells acquired by FACS Canto II (BD)

were analyzed by FACS Diva (BD) and FlowJo (Treestar). Approximately 1 x 10^5 PBMCs were

gated per test.

Nef-mediated signaling perturbation analysis

Virus supernatants obtained from COS-1 cells after transfection with wild-type or mutant

SIV_{mac239} molecular clones were used for infection of $CD4^+$ T-cell lines, cynomolgus macaque-

derived HSC-F and rhesus macaque-derived HSR5.4 (Akari et al., 1996). QuikChange II XL

site-directed mutagenesis kit (Agilent Technologies) was used to construct mutant SIV_{mac239}

molecular clones possessing *nef* mutations Nef-G63E (G-to-A mutation at nt 9520) and Nef-

G63R (G-to-A mutation at nt 9519) from the wild-type SIV_{mac239} molecular clone (Kestler et al.,

1991) (nt number from GenBank accession number M33262). Cells (1 x 10^5 cells/well in U-

bottomed 96-well culture plates [BD]) were infected with wild-type or mutant SIV_{mac239} cultured

in RPMI-1640 medium supplemented with 10% fetal bovine serum (FBS) for indicated periods

at MOI 0.1 (intracellular signaling analysis) or 0.001 (supernatant analysis). Culture

supernatants were subjected to measurement of SIV capsid p27 concentrations by ELISA.

Harvested cells were fixed and permeabilized with Cytofix/Cytoperm kit, washed twice and

subjected to immunostaining. The following antibodies were used; anti-SIV_{mac251} Gag p27 mAb

(ABL) manually conjugated with Alexa 647 (Life Technologies), anti-SIV_{mac251} Nef mAb (clone

17, epitope peptide corresponding to SIV_{mac239} Nef 71-80 not including the residue G63:

Thermo Fisher Scientific/Pierce) manually conjugated with Alexa 488 or Alexa 647, anti-

human CD3-APC-Cy7 (SP34-2, BD Pharmingen), anti-human CD4-PerCP (L200, BD

Pharmingen), anti-human HLA-ABC-FITC (G46-2.6, BD Pharmingen), anti-human CXCR4-

APC (12G5, BioLegend), BST2-Alexa 647 (RS38E, BioLegend), Alexa 488-conjugated anti-

phospho-Akt (Ser473) (D9E, CST) and Alexa 488-conjugated anti-phospho-Akt (Thr308)

(C31E5E, CST). Cells acquired by FACS Canto II were analyzed by FACS Diva and FlowJo.

Approximately 5 x 10^4 cells were gated for each test.

Ferritin ELISA

Plasma ferritin levels in NAb inducers and control animals were analyzed by monkey ferritin

sandwich ELISA kit (LS Bio) according to the manufacturer's instructions.

PI3K stimulation assay

5×10^4 HSC-F cells were infected with wild-type or Nef G63E mutant SIV_{mac239} at MOI 0.2 and

cultured for 1 day in medium supplemented with 10% (normal), 1% (1/10 starvation) or 0.1%

(1/100 starvation) FBS. For ligand stimulation (**Figure 4-figure supplement 1C**), cells at the

end of 1-day culture were pulsed for 20 minutes with 40 ng/ml of recombinant human IFN-γ

(Gibco/Thermo Fisher Scientific), 100 IU/ml of recombinant human IL-2 (Roche Diagnostics)

or 10 µg/ml of SIV_{mac251} Env gp130 (ImmunoDx). Cells were intracellularly stained using

Cytofix/Cytoperm kit with anti-SIV_{mac251} Nef mAb or anti-SIV_{mac251} p27 mAb manually

conjugated to Alexa 647 and PE-conjugated anti-human phospho-Akt (Ser473) (D9E, CST) or

Alexa 488-conjugated anti-human phospho-Akt (Ser473). Cells acquired by FACS Canto II

were analyzed by FACS Diva and FlowJo. Approximately 7×10^4 HSC-F cells were gated per

test.

Transcriptome analysis

Total RNAs were extracted using RNeasy Plus Mini kit (Qiagen) from 2×10^6 HSC-F cells 1

day after infection with wild-type or Nef G63E mutant SIV_{mac239} at MOI 5. Negative control RNA

samples were extracted from 2×10^6 uninfected HSC-F cells after culture with the same

condition. Three sets of experiments were performed. Total RNA samples were subjected to a

quality control (QC) analysis using an Agilent 2100 Bioanalyzer. The obtained amounts of total

RNAs were: 12.37 ± 0.39 (uninfected), 8.80 ± 0.44 (wild-type) and 9.02 ± 0.17 (Nef G63E) µg

($P = 0.66$ for wild-type versus Nef G63E by unpaired t test). In all samples, two bands of 18S

and 28S rRNA were confirmed and the RNA integrity number (RIN) was 10. 500 ng of total

RNA samples were processed with GeneChip WT Plus reagent (Affymetrix/Thermo Fisher

Scientific) to produce 150 µl of fragmented and labeled cDNA samples. These were incubated

with a Human Gene 2.0 ST Array (Affymetrix/Thermo Fisher Scientific) for 16 hours, 60 rpm

at 45°C using GeneChip hybridization oven 645 (Affymetrix/Thermo Fisher Scientific). Results

were scanned with GeneChip Scanner 3000 7G (Affymetrix/Thermo Fisher Scientific) and

processed with Affymetrix Expression Console Software (Affymetrix/Thermo Fisher Scientific)

according to the manufacturer's instructions. Expression values were normalized by the RMA

method. Genes above mean background expression within the cognate sample and showing

significant difference between wild-type and Nef G63E SIV ($P < 0.05$ via unpaired t test for

log_2-transformed values, 768 candidates) were determined. Akt-related genes exhibiting a

change of approximately 10% or more were representatively extracted and manually aligned

by the authors.

Peripheral CD4+ T-cell surface staining

Cryopreserved/thawed PBMCs were stained for 30 minutes at 4°C with the following reagents

or conjugated anti-human mAbs: Live/Dead Aqua (Life Technologies), anti-CD4-PerCP (L200,

BD Pharmingen), anti-CD8-APC-Cy7 (RPA-T8, BD Pharmingen), anti-CD3-Alexa 700 (SP34-

2, BD Pharmingen), anti-CD95-PE-Cy7 (DX2, eBioscience), anti-CXCR5-PE (87.1,

eBioscience), anti-PD-1-Brilliant Violet 421 (EH12.2H7, BioLegend) and anti-CXCR3-Alexa

488 (G025H7, BioLegend). Cells acquired by FACS LSRII Fortessa (BD) were analyzed by

FACS Diva and FlowJo. Approximately 1.5×10^5 PBMCs were gated per test.

Proximity ligation assay (PLA)

A flow cytometry-based arrangement of proximity ligation assay (PLA) (Avin et al., 2017) was

performed to quantitatively assess Nef binding to candidate interacting molecules. 1×10^5

HSC-F cells were infected with SIV_{mac239} at MOI 0.05 in U-bottomed plates and permeated

with Cytofix/Perm kit (BD Biosciences). After two washes, they were resuspended in 0.5%

BSA/PBS for prevention of experimental procedure-related loss and stained with mouse anti-

SIV_{mac251} Nef mAb (clone 17, Thermo Fisher Scientific/Pierce) or mouse IgG1 isotype control

mAb (P3.6.2.8.1, Abcam) in combination with either of the following rabbit antibodies: anti-

GβL (86B8, CST), anti-mTOR (7C10, CST), polyclonal anti-human PI3K p85 (Merck Millipore),

anti-PI3 Kinase p110α (C73F8, CST) or rabbit IgG1 isotype control mAb (DA1E, CST).

Antibody-stained cells were subsequently probed with Duolink In Situ PLA Probe anti-mouse

PLUS and anti-mouse MINUS probes (Sigma/Merck). Next, they were detected for

intermolecular binding using Duolink flow PLA detection kit (Green) (Sigma/Merck) with a

reaction time of 100 minutes for post-mouse/rabbit probe linking amplification. Finally, these

PLA-subjected cells were additionally stained with anti-SIV$_{mac251}$ Gag p27 mAb manually

conjugated with Alexa 647 and anti-CD4-PerCP for 30 minutes for 20 minutes at 4°C. Cells

acquired by FACS Canto II were analyzed by FACS Diva and FlowJo. Approximately 1 x 10^5

cells were gated per test. Binding index (Y axis) was calculated by deriving the deviation from

the summation of: {baseline P: (MFI of background reaction with mouse isotype control/rabbit

isotype control) + anti-Nef antibody-derived background Q: [(MFI of reaction with mouse anti-

Nef/rabbit isotype control) - (MFI of background reaction with mouse isotype control/rabbit

isotype control)] + anti-binding partner antibody-derived background R:[(MFI of reaction with

mouse isotype control/rabbit anti-binding partner molecule) - (MFI of background reaction with

mouse isotype control/rabbit isotype control)]}.

Co-immunoprecipitation analysis

1 x 10^5 HSC-F cells were infected with SIV$_{mac239}$ at MOI 0.05 in U-bottomed plates in

quadruplicate for each SIV strain for three days and pooled for each for acquiring cell pellets.

These pellets were lysed with Capturem IP & Co-IP Kit lysis buffer (Takara) and portions for

each were immunoprecipitated by anti-Sin1 mAb (D7G1A, CST) (1:50 dilution, 20 minutes,

room temperature). Spin column membrane-bound immunoprecipitates were obtained by

centrifugation with Capturem IP & Co-IP Kit (Clontech/Takara). Whole cell lysates and Sin1-

immunoprecipitated products were subjected to SDS-polyacrylamide gel electrophoresis

separation on a Mini Protean TGX 4-15% gel (BioRad) and transferred to a Immun-Blot PVDF

membrane (BioRad). Immunoblotting was performed by mouse anti-SIV$_{mac251}$ Nef mAb (clone

17, Thermo Fisher Scientific/Pierce) primary antibody probing (1:1000 dilution, 18 hours, 4°

C) and Mouse TrueBlot ULTRA anti-mouse Ig HRP (eB144, Rockland Immunochemicals)

secondary antibody incubation (1:1000 dilution, 30 minutes, room temperature). Nef-specific

bands in infected controls were visualized by enhanced chemiluminescence using

SuperSignal West Pico PLUS (Thermo Fisher Scientific). Bands auto-detectable by Image

Lab software (BioRad) were analyzed for signal intensities. Uncropped images are provided

as supplementary data.

Cell death assay

Infected cell apoptosis was measured by methods previously described (Jauch et al., 2023).

1 x 10^5 HSC-F cells were infected with SIV$_{mac239}$ at MOI 0.1 in U-bottomed plates (six

wells/control) for each SIV strain for three days. These cells were stained with CD4-FITC (M-

T477, BD Pharmingen) for 15 minutes at room temperature, washed once and next stained

with APC Annexin V (Biolegend) in Annexin V Binding Buffer (Biolegend) for 20 minutes at

room temperature. Stained reactions were then diluted fivefold with Annexin V Binding Buffer

and subjected to analysis. Cells acquired by FACS Lyric were analyzed by FACS Suite and

FlowJo. Approximately 1 x 10^4 HSC-F cells were gated per test.

Quantitative *in vivo* imaging flow cytometry

Cryopreserved/thawed lymph node cells (LNCs) from several persistently SIV-infected

macaques used in previous experiments (Nakane et al., 2013; Takahashi et al., 2013; Iwamoto

et al., 2014) were seeded in V-bottomed 96-well plates (Nunc) and blocked with 25 μg/ml of

anti-human CD4 (clone L200: BD) in 100 μl volume for 15 minutes at 4°C. After three washes,

they were stained with 10 μg/ml of recombinant SIV$_{mac239}$ Env gp140-biotin (Immune

Technology) for 30 minutes at 4°C. Cells were then stained with anti-CD19-PC5.5 (J3-119,

Beckman Coulter), Live/Dead Aqua (Invitrogen) and streptavidin-APC-Cy7 (BioLegend) for 30

minutes at 4°C. After two washes, cells were processed with Cytofix/Cytoperm kit (BD) for 20

minutes at 4°C, washed twice, and next intracellularly stained with PE-conjugated anti-human

phospho-Akt (Ser473) (D9E, CST) and anti-SIV$_{mac251}$ Nef mAb (clone 17, Thermo Fisher

Scientific/Pierce) manually conjugated with Alexa 488 or mouse IgG1 isotype control mAb

(P3.6.2.8.1, Abcam) for 30 minutes at 4°C. After final two washes, cells were suspended in

0.8% PFA/PBS. All centrifugations for washing (1,200 x g, 2 minutes) were performed at 4°C.

Cells were subjected to image acquisition with Image Stream X MKII imaging flow cytometer

(Amnis/Merck Millipore/Luminex) and analyzed with IDEAS 6.3 software (Amnis/Merck

Millipore/Luminex). Approximately 5 x 10^5 LNCs were acquired for analyses (**Figure 5**). A

custom-implemented linear discriminant analysis-based machine learning module (Luminex)

was utilized for verifying candidate Nef staining signal-related secondary parameters (**Figure**

5 and **Figure 5-figure supplement 1A**) for their efficacy of target population separation.

B-cell Nef invasion *in vitro* assay

For Nef invasion assay, 5 x 10^4 HSC-F cells were infected at MOI 1 for 48 hours. These were

then additionally cocultured with 7.5 x 10^4 Ramos B cells for 12 hours with or without addition

of macaque T cell-stimulating anti-CD3 antibody (FN-18, Abcam) at 1 μg/ml. Cells were

double-stained with Alexa 488-conjugated mouse anti-Nef (clone 17: Thermo

Scientific/Pierce) and anti-CD19-PC-5.5 (J3-119, Beckman Coulter). For graphical

visualization (**Figure 6A**), interacting Nef$^+$ infected cell-invaded B cell doublet images acquired

with ImageStream X MKII (Amnis/Merck Millipore) were analyzed by IDEAS 6.3 (Amnis/Merck

Millipore/Luminex) with the defined noise-cancelling gating (via gating Nef pixel signal

contrast^{lo}-Nef mean pixel per object^{hi} cells). For Nef invasion modulation verification on

conventional flow cytometry, the aforementioned culture condition was similarly used. Anti-

CD3 antibody (FN-18, Abcam) addition at 1 μg/ml and phorbol 12-myristate 13-acetate (PMA,

Sigma) plus ionomycin (Sigma) addition (10 ng/ml and 50 ng/ml, respectively) l were evaluated.

Cells were stained with Alexa 647-conjugated mouse anti-Nef (clone 17: Thermo

Scientific/Pierce), anti-CD19-PC-5.5 (J3-119, Beckman Coulter), anti-human phospho-Akt

(Ser473)-Alexa 488 (D9E, CST) and Live/Dead Near-IR (Invitrogen) for dead cell exclusion.

Dead cell frequencies were less than 3%. For viral phenotype evaluation, 1 x 10^5 HSC-F cells

(surface CD40L$^+$) were pre-infected at MOI 0.2 with wild type or Nef-G63E mutant SIV$_{mac239}$

six hours before coculture. 1.5 x 10^5 CD20$^+$ B cells, positively selected from fresh PBMCs of

uninfected rhesus macaques via anti-CD20 microbeads (Miltenyi Biotec), were cocultured with

these HSC-F cells at an E:T ratio of 2:3 for 3 days in the presence of 40 ng/ml carrier-free

recombinant human IL-4 (R&D Systems). Cells were surface-stained with anti-human CD20-

PerCP (2H7, BioLegend) and intracellularly stained using Cytofix/Cytoperm kit with anti-

SIV$_{mac251}$ Nef mAb manually conjugated to Alexa 488 and Alexa 647-conjugated anti-human

phospho-Akt (Ser473) (D9E, CST). For **Figure 6B** and **Figure 6C**, Cells acquired by FACS

Canto II were analyzed by FACS Diva and FlowJo. Approximately 5 x 10^4 CD20$^+$ primary B

cells and 3 x 10^4 CD19$^+$ Ramos B cells were analyzed.

Peripheral SIV Env-specific B-cell responses

Peripheral SIV Env gp140-specific memory B cells (B$_{mem}$) and plasmablasts (PBs) were

measured by flow cytometry with procedures modified from previous reports and our

experience (Silveira et al., 2015; Hau et al., 2022). First, cryopreserved/thawed PBMCs

seeded in V-bottomed 96-well plates (Nunc) were blocked with 25 μg/ml of anti-human CD4

(clone L200: BD) in 100 μl volume for 15 minutes at 4°C. This concentration attains blockade

of promiscuous SIV Env binding to CD4 comparable to levels by magnetic depletion of CD3$^+$

T cells (data not shown). After three washes, they were next stained with 10 μg/ml of

recombinant SIV$_{mac239}$ Env gp140-biotin (Immune Technology) for 30 minutes at 4°C. They

were subsequently stained with the following anti-human antibodies and fluorochromes for 30

minutes at 4°C with combinations as follows; PB/B$_{mem}$: anti-CD3-APC-Cy7 (SP34-2, BD

Pharmingen), anti-CD8-APC-Cy7 (RPA-T8, BD Pharmingen), anti-CD14-APC-Cy7 (M5E2,

BioLegend), anti-CD16-APC-Cy7 (3G8, BioLegend), streptavidin-Brilliant Violet 421

(BioLegend), anti-CD19-PC5.5 (J3-119, Beckman Coulter) and Live/Dead Aqua (Invitrogen);

PB: anti-CD10-APC-Cy7 (HI10a, BioLegend), anti-CD11c-PE-Cy7 (3.9, BioLegend), anti-

CD123-PE-Cy7 (6H6, BioLegend) and anti-HLA-DR-PE-Texas Red (TU36, Invitrogen); B_{mem}:

anti-CD27-PE/Dazzle 594 (M-T271, BioLegend), anti-CD10-PE-Cy7 (HI10a, BioLegend), anti-

IgD-FITC (DaKo), anti-CD38-Alexa 647 (AT1, Santa Cruz Biotechnology), anti-IgG-Alexa 700

(G18-145, BD Pharmingen) and anti-CD138-PE (DL-101, eBioscience). After two washes,

B_{mem} samples were suspended in 0.8% PFA/PBS. For PB staining, surface-stained cells were

further processed with Cytofix/Cytoperm kit (BD) for 20 minutes at 4°C, washed twice, and

next intracellularly stained with 10 μg/ml of recombinant SIV_{mac239} Env gp140-biotin (Immune

Technology) for 30 minutes at 4°C. After three washes, they were next stained with anti-human

IgG-PE (G18-145, BD), anti-human/mouse IRF4-eFluor 660 (3E4, eBioscience), anti-human

Ki67-Alexa 700 (B56, BD Pharmingen) and streptavidin-Alexa 488 (BioLegend) for 30 minutes

at 4°C. After final two washes, cells were suspended in 0.8% PFA/PBS. Cells acquired with

FACS LSRII Fortessa were analyzed with FACS Diva and FlowJo. Approximately 1.5 x 10^5

PBMCs were acquired for B_{mem} and 4 x 10^5 PBMCs were acquired for PB analyses (**Figure**

6). All centrifugations for washing (1,200 x g, 2 minutes) were performed at 4°C. Env-specific

PB frequencies were 0 cells/million PBMCs for pre-challenge samples in all animals examined,

ruling out background confounding for high-sensitivity quantitation of the population.

Statistical Analysis

Analyses were performed via Prism 8 (GraphPad Software). $P < 0.05$ were considered

significant in two-tailed unpaired t tests, paired t tests, Mann Whitney U tests, Fisher's exact

tests, Wilcoxon signed-rank tests, one-way ANOVA with Tukey's post-hoc multiple comparison

tests and two-way ANOVA with Sidak's post-hoc multiple comparison tests. For analysis of

Nef-invaded B-cell pAkt levels, median fluorescence intensities (mFIs) were analyzed due to

relatively small numbers of Nef[+] B cells. Analyses involving pAkt Ser473 levels all derived

comparable results between mFI and MFI for wild type versus Nef G63E virus. Machine

learning-based rating of Nef signal parameters was verified with IDEAS 6.3 Machine Learning

Module (Luminex).

References

Adnan, S., Reeves, R.K., Gillis, J., Wong, F.E., Yu, Y., Camp, J.V., Li, Q., Connole, M., Li, Y., Piatak, M. Jr., et al. (2016). Persistent low-level replication of SIVΔnef drives maturation of antibody and CD8 T cell responses to induce protective immunity against vaginal SIV infection. *PLoS Pathog*. 12, e1006104.

Akari, H., Mori, K., Terao, K., Otani, I., Fukasawa, M., Mukai, R., and Yoshikawa, Y. (1996). In vitro immortalization of Old World monkey T lymphocytes with Herpesvirus saimiri: its susceptibility to infection with simian immunodeficiency viruses. *Virology* 218, 382–388.

Al Qureshah, F., Sagadiev, S., Thouvenel, C.D., Liu, S., Hua, Z., Hou, B., Acharya, M., James, R.G., and Rawlings, D.J. (2021). Activated PI3Kδ signals compromise plasma cell survival via limiting autophagy and increasing ER stress. *J. Exp. Med.* 218, e20211035.

Angulo, I., Vadas, O., Garçon, F., Banham-Hall, E., Plagnol, V., Leahy, T.R., Baxendale, H., Coulter, T., Curtis, J., Wu, C., et al. (2013). Phosphoinositide 3-kinase δ gene mutation predisposes to respiratory infection and airway damage. *Science* 342, 866–871.

Asano, T., Okada, S., Tsumura, M., Yeh, T.W., Mitsui-Sekinaka, K., Tsujita, Y., Ichinose, Y., Shimada, A., Hashimoto, K., Wada, T., et al. (2018). Enhanced AKT phosphorylation of circulating B cells in patients with activated PI3Kδ syndrome. *Front. Immunol.* 9, 568.

Avery, D.T., Kane, A., Nguyen, T., Lau, A., Nguyen, A., Lenthall, H., Payne, K., Shi, W., Brigden, H., French, E., et al. (2018). Germline-activating mutations in PIK3CD compromise B cell development and function. *J. Exp. Med.* 215, 2073–2095.

1026   Avin, A., Levy, M., Porat, Z., and Abramson, J. (2017) Quantitative analysis of protein-protein

1027   interactions and post-translational modifications in rare immune populations. *Nat. Commun.*

1028   8, 1524.

1029

1030   Battegay, M., Moskophidis, D., Waldner, H., Bründler, M.A., Fung-Leung, W.P., Mak, T.W.,

1031   Hengartner, H., and Zinkernagel, R.M. (1993). Impairment and delay of neutralizing antiviral

1032   antibody responses by virus-specific cytotoxic T cells. *J. Immunol.* 151, 5408–5415.

1033

1034   Baumjohann, D., Preite, S., Reboldi, A., Ronchi, F., Ansel, K.M., Lanzavecchia, A., and

1035   Sallusto, F. (2013). Persistent antigen and germinal center B cells sustain T follicular helper

1036   cell responses and phenotype. *Immunity* 38, 596–605.

1037

1038   Bradley, T., Peppa, D., Pedroza-Pacheco, I., Li, D., Cain, D.W., Henao, R., Venkat, V., Hora,

1039   B., Chen, Y., Vandergrift, N.A., et al. (2018). RAB11FIP5 Expression and Altered Natural Killer

1040   Cell Function Are Associated with Induction of HIV Broadly Neutralizing Antibody Responses.

1041   *Cell* 175, 387–399.e17.

1042

1043   Burns, D.P., Collignon, C., and Desrosiers, R.C. (1993). Simian immunodeficiency virus

1044   mutants resistant to serum neutralization arise during persistent infection of rhesus monkeys.

1045   *J. Virol.* 67, 4104–4113.

1046

1047   Campbell, E.M., Nunez, R., and Hope, T.J. (2004). Disruption of the actin cytoskeleton can

1048   complement the ability of Nef to enhance human immunodeficiency virus type 1 infectivity. *J.*

1049   *Virol.* 78, 5745–5755.

1050

1051   Coulter, T.I., Chandra, A., Bacon, C.M., Babar, J., Curtis, J., Screaton, N., Goodlad, J.R.,

1052   Farmer, G., Steele, C.L., Leahy, T.R., et al. (2017). Clinical spectrum and features of activated

1053   phosphoinositide 3-kinase δ syndrome: A large patient cohort study. *J. Allergy Clin. Immunol.*

1054   139, 597–606.e4.

1055

1056   Cubas, R.A., Mudd, J.C., Savoye, A.L., Perreau, M., van Grevenynghe, J., Metcalf, T., Connick,

1057   E., Meditz, A., Freeman, G.J., Abesada-Terk, G. Jr., et al. (2013). Inadequate T follicular cell

1058   help impairs B cell immunity during HIV infection. *Nat. Med.* 19, 494–499.

1059

1060   Cumont, M.C., Diop, O., Vaslin, B., Elbim, C., Viollet, L., Monceaux, V., Lay, S., Silvestri, G.,

1061   Le Grand, R., Müller-Trutwin, M., et al. (2008). Early divergence in lymphoid tissue apoptosis

1062   between pathogenic and nonpathogenic simian immunodeficiency virus infections of

1063   nonhuman primates. *J. Virol.* 82, 1175–1184.

1064

1065   Dornan, G.L., Siempelkamp, B.D., Jenkins, M.L., Vadas, O., Lucas, C.L., and Burke, J.E.

1066   (2017). Conformational disruption of PI3Kδ regulation by immunodeficiency mutations in

1067   PIK3CD and PIK3R1. *Proc. Natl. Acad. Sci. USA* 114, 1982–1987.

1068

1069   Douek, D.C., Brenchley, J.M., Betts, M.R., Ambrozak, D.R., Hill, B.J., Okamoto, Y., Casazza,

1070   J.P., Kuruppu, J., Kunstman, K., Wolinsky, S., et al. (2002). HIV preferentially infects HIV-

1071   specific CD4[+] T cells. *Nature* 417, 95–98.

1072

1073   Evans, D.T., O'Connor, D.H., Jing, P., Dzuris, J.L., Sidney, J., da Silva, J., Allen, T.M., Horton,

1074   H., Venham, J.E., Rudersdorf, R.A., et al. (1999). Virus-specific cytotoxic T-lymphocyte

1075   responses select for amino-acid variation in simian immunodeficiency virus Env and Nef. *Nat.*

1076   *Med.* 5, 1270–1276.

1077

1078   Fallet, B., Narr, K., Ertuna, Y.I., Remy, M., Sommerstein, R., Cornille, K., Kreutzfeldt, M., Page,

1079   N., Zimmer, G., Geier, F., et al. (2016). Interferon-driven deletion of antiviral B cells at the onset

1080   of chronic infection. *Sci. Immunol.* 1, pii: eaah6817.

1081

1082   François, F., and Klotman, M.E. (2003). Phosphatidylinositol 3-kinase regulates human

1083   immunodeficiency virus type 1 replication following viral entry in primary CD4+ T lymphocytes

1084   and macrophages. *J. Virol.* 77, 2539–2549.

1085

1086   Fruman, D.A., Snapper, S.B., Yballe, C.M., Davidson, L., Yu, J.Y., Alt, F.W., and Cantley, L.C.

1087   (1999). Impaired B cell development and proliferation in absence of phosphoinositide 3-kinase

1088   p85alpha. *Science* 283, 393–397.

1089

1090   Fukazawa, Y., Park, H., Cameron, M.J., Lefebvre, F., Lum, R., Coombes, N., Mahyari, E.,

1091   Hagen, S.I., Bae, J.Y., Reyes, M.D. 3rd., et al. (2012). Lymph node T cell responses predict

1092   the efficacy of live attenuated SIV vaccines. *Nat. Med.* 18, 1673–1681.

1093

1094   Gauduin, M.C., Yu, Y., Barabasz, A., Carville, A., Piatak, M., Lifson, J.D., Desrosiers, R.C.,

1095   and Johnson, R.P. (2006). Induction of a virus-specific effector-memory CD4+ T cell response

1096   by attenuated SIV infection. *J. Exp. Med.* 203, 2661–2672.

1097

1098   Geyer, M., Fackler, O.T., and Peterlin, B.M. (2001). Structure--function relationships in HIV-1

1099   Nef. *EMBO Rep.* 2, 580–585.

1100

1101   Gitlin, A.D., Shulman, Z. and Nussenzweig, M.C. (2014). Clonal selection in the germinal

1102   centre by regulated proliferation and hypermutation. *Nature* 509, 637–640.

1103

1104   Gray, E.S., Madiga, M.C., Hermanus, T., Moore, P.L., Wibmer, C.K., Tumba, N.L., Werner, L.,

1105   Mlisana, K., Sibeko, S., Williamson, C., et al. (2011). The neutralization breadth of HIV-1

1106   develops incrementally over four years and is associated with CD4+ T cell decline and high

1107   viral load during acute infection. *J. Virol.* 85, 4828–4840.

1108

1109   Haigwood, N.L., Watson, A., Sutton, W.F., McClure, J., Lewis, A., Ranchalis, J., Travis, B.,

1110   Voss, G., Letvin, N.L., Hu, S.L., et al. (1996). Passive immune globulin therapy in the

1111   SIV/macaque model: early intervention can alter disease profile. *Immunol. Lett.* 51, 107–114.

1112

1113   Haralick, R.M., Shanmugam, K., and Dinstein, I. (1973) Textural features for image

1114   classification. *IEEE Transactions on Systems, Man, and Cybernetics* SMC-3, 610–621.

1115

1116   Harris, R.S., Bishop, K.N., Sheehy, A.M., Craig, H.M., Petersen-Mahrt, S.K., Watt, I.N.,

1117   Neuberger, M.S., and Malim, M.H. (2003). DNA deamination mediates innate immunity to

1118   retroviral infection. *Cell* 113, 803–809.

1119

1120   Hashimoto, M., Bhuyan, F., Hiyoshi, M., Noyori, O., Nasser, H., Miyazaki, M., Saito, T., Kondoh,

1121   Y., Osada, H., Kimura, S., et al. (2016). Potential role of the formation of tunneling nanotubes

1122   in HIV-1 spread in macrophages. *J. Immunol.* 196, 1832–1841.

1123

1124   Hau, T.T.T., Nishizawa, M., Harada, S., Phan, M.H., Kanno, Y., Nomura, T., Matsuoka, S.,

1125   Kawana-Tachikawa, A., Hall, W.W., Matano, T., et al. (2022). Association of envelope-specific

1126   B-cell differentiation and viral selective pressure signatures in HIV-1 CRF01_AE infection.

1127   *AIDS* 36, 1629–1641.

1128

1129   Havenar-Daughton, C., Carnathan, D.G., Torrents de la Peña, A., Pauthner, M., Briney, B.,

1130   Reiss, S.M., Wood, J.S., Kaushik, K., van Gils, M.J., Rosales, S.L., et al. (2016). Direct Probing

1131   of Germinal Center Responses Reveals Immunological Features and Bottlenecks for

1132   Neutralizing Antibody Responses to HIV Env Trimer. *Cell Rep.* 17, 2195–2209.

1133

1134   Hirsch, V.M., Santra, S., Goldstein, S., Plishka, R., Buckler-White, A., Seth, A., Ourmanov, I.,

1135   Brown, C.R., Engle, R., Montefiori, D., et al. (2004). Immune failure in the absence of profound

1136   CD4+ T-lymphocyte depletion in simian immunodeficiency virus-infected rapid progressor

1137   macaques. *J. Virol.* 78, 275–284.

1138

1139   Hoof, I., Peters, B., Sidney, J., Pedersen, L.E., Sette, A., Lund, O., Buus, S., and Nielsen, M..

1140   (2009). NetMHCpan, a method for MHC class I binding prediction beyond humans.

1141   *Immunogenetics* 61, 1–13.

1142

1143   Hunziker, L., Recher, M., Macpherson, A.J., Ciurea, A., Freigang, S., Hengartner, H., and

1144   Zinkernagel, R.M. (2003). Hypergammaglobulinemia and autoantibody induction mechanisms

1145   in viral infections. *Nat. Immunol.* 4, 343–349.

1146

1147   Iseda, S., Takahashi, N., Poplimont, H., Nomura, T., Seki, S., Nakane, T., Nakamura, M., Shi,

1148   S., Ishii, H., Furukawa, S., et al. (2016). Biphasic CD8+ T-cell defense in simian

1149   immunodeficiency virus control by acute-phase passive neutralizing antibody immunization. *J.*

1150   *Virol.* 90, 6276–6290.

1151

1152   Ishii, H., Kawada, M., Tsukamoto, T., Yamamoto, H., Matsuoka, S., Shiino, T., Takeda, A.,

1153   Inoue, M., Iida, A., Hara, H., et al. (2012). Impact of vaccination on cytotoxic T lymphocyte

1154   immunodominance and cooperation against simian immunodeficiency virus replication in

1155   rhesus macaques. *J. Virol.* 86, 738–745.

1156

1157   Iwamoto, N., Takahashi, N., Seki, S., Nomura, T., Yamamoto, H., Inoue, M., Shu, T., Naruse,

1158   T.K., Kimura, A., and Matano, T. (2014). Control of simian immunodeficiency virus replication

1159   by vaccine-induced Gag- and Vif-specific CD8[+] T cells. *J. Virol.* 88, 425–433.

1160

1161   Jauch, A.J., Bignucolo, O., Seki, S., Ghraichy, M., Delmonte, O.M., von Niederhäusern, V.,

1162   Higgins, R., Ghosh, A., Nishizawa, M., Tanaka, M., et al. (2023). Autoimmunity and

1163   immunodeficiency associated with monoallelic LIG4 mutations via haploinsufficiency. *J.*

1164   *Allergy Clin. Immunol.* 152, 500–516.

1165

1166   Jia, B., Serra-Moreno, R., Neidermyer, W., Rahmberg, A., Mackey, J., Fofana, I.B., Johnson,

1167   W.E., Westmoreland, S., and Evans, D.T. (2009). Species-specific activity of SIV Nef and HIV-

1168   1 Vpu in overcoming restriction by tetherin/BST2. *PLoS Pathog.* 5, e1000429.

1169

1170   Joas, S., Parrish, E.H., Gnanadurai, C.W., Lump, E., Stürzel, C.M., Parrish, N.F., Learn, G.H.,

1171   Sauermann, U., Neumann, B., Rensing, K.M., et al. (2018) Species-specific host factors rather

1172   than virus-intrinsic virulence determine primate lentiviral pathogenicity. *Nat. Commun.* 9, 1371.

1173

1174   Joas, S., Sauermann, U., Roshani, B., Klippert, A., Daskalaki, M., Mätz-Rensing, K., Stolte-

1175   Leeb, N., Heigele, A., Tharp, G.K., Gupta, P.M., et al. (2020). Nef-Mediated CD3-TCR

1176   Downmodulation Dampens Acute Inflammation and Promotes SIV Immune Evasion. *Cell Rep.*

1177   30, 2261–2274.e7.

1178

1179   Johnson, R.P., Glickman, R.L., Yang, J.Q., Kaur, A., Dion, J.T., Mulligan, M.J., and Desrosiers,

1180   R.C. (1997). Induction of vigorous cytotoxic T-lymphocyte responses by live attenuated simian

1181   immunodeficiency virus. *J. Virol.* 71, 7711–7718.

1182

1183   Jolly, C., Kashefi, K., Hollinshead, M., and Sattentau, Q.J. (2004). HIV-1 cell to cell transfer

1184   across an Env-induced, actin-dependent synapse. *J. Exp. Med.* 199, 283–293.

1185

1186   Junt, T., Moseman, E.A., Iannacone, M., Massberg, S., Lang, P.A., Boes, M., Fink, K.,

1187   Henrickson, S.E., Shayakhmetov, D.M., Di Paolo, N.C., et al. (2007). Subcapsular sinus

1188   macrophages in lymph nodes clear lymph-borne viruses and present them to antiviral B cells.

1189   *Nature* 450, 110–114.

1190

1191   Kawada, M., Tsukamoto, T., Yamamoto, H., Takeda, A., Igarashi, H., Watkins, D.I., and Matano,

1192   T. (2007). Long-term control of simian immunodeficiency virus replication with central memory

CD4+ T-cell preservation after nonsterile protection by a cytotoxic T-lymphocyte-based vaccine. *J. Virol.* 81, 5202–5211.

Kennedy, S.G., Wagner, A.J., Conzen, S.D., Jordán, J., Bellacosa, A., Tsichlis, P.N., and Hay, N. (1997). The PI 3-kinase/Akt signaling pathway delivers an anti-apoptotic signal. *Genes Dev.* 11, 701–713.

Kestler, H.W. 3rd, Ringler, D.J., Mori, K., Panicali, D.L., Sehgal, P.K., Daniel, M.D., and Desrosiers, R.C. (1991). Importance of the nef gene for maintenance of high virus loads and for development of AIDS. *Cell* 65, 651–662.

Khan, H., Sumner, R.P., Rasaiyaah, J., Tan, C.P., Rodriguez-Plata, M.T., Van Tulleken, C., Fink, D., Zuliani-Alvarez, L., Thorne, L., Stirling, D., et al. (2020). HIV-1 Vpr antagonizes innate immune activation by targeting karyopherin-mediated NF-κB/IRF3 nuclear transport. *Elife* 9, e60821.

Kim, D.H., Sarbassov, D.D., Ali, S.M., Latek, R.R., Guntur, K.V., Erdjument-Bromage, H., Tempst, P., and Sabatini, D.M. (2003). GbetaL, a positive regulator of the rapamycin-sensitive pathway required for the nutrient-sensitive interaction between raptor and mTOR. *Mol. Cell* 11, 895–904.

Kirchhoff, F., Greenough, T.C., Brettler, D.B., Sullivan, L.L. and Desrosiers, R.C. (1995). Absence of intact nef sequences in a long-term, nonprogressing survivor of HIV-1 infection. *N. Engl. J. Med. 332*, 228–232.

Kumar, A., Abbas, W., Colin, L., Khan, K.A., Bouchat, S., Varin, A., Larbi, A., Gatot, J.S., Kabeya, K., Vanhulle, C., et al. (2016). Tuning of AKT-pathway by Nef and its blockade by

protease inhibitors results in limited recovery in latently HIV infected T-cell line. *Sci. Rep.* 6, 24090.

Kumar, V., Scandella, E., Danuser, R., Onder, L., Nitschké, M., Fukui, Y., Halin, C., Ludewig, B., and Stein, J.V. (2010). Global lymphoid tissue remodeling during a viral infection is orchestrated by a B cell-lymphotoxin-dependent pathway. *Blood* 115, 4725–4733.

Laguette, N., Sobhian, B., Casartelli, N., Ringeard, M., Chable-Bessia, C., Ségéral, E., Yatim, A., Emiliani, S., Schwartz, O., and Benkirane, M. (2011). SAMHD1 is the dendritic- and myeloid-cell-specific HIV-1 restriction factor counteracted by Vpx. *Nature* 474, 654–657.

Langer, S., Hammer, C., Hopfensperger, K., Klein, L., Hotter, D., De Jesus, P.D., Herbert, K.M., Pache, L., Smith, N., van der Merwe, J.A., et al. (2019) HIV-1 Vpu is a potent transcriptional suppressor of NF-κB-elicited antiviral immune responses. *Elife* 8, e41930.

Leuchowius, K.J., Weibrecht, I., Landegren, U., Gedda, L., and Söderberg, O. (2009). Flow cytometric in situ proximity ligation analyses of protein interactions and post-translational modification of the epidermal growth factor receptor family. *Cytometry A* 75, 833–839.

Levesque, M.C., Moody, M.A., Hwang, K.K., Marshall, D.J., Whitesides, J.F., Amos, J.D., Gurley, T.C., Allgood, S., Haynes, B.B., Vandergrift, N.A., et al. (2009). Polyclonal B cell differentiation and loss of gastrointestinal tract germinal centers in the earliest stages of HIV-1 infection. *PLoS Med.* 6, e1000107.

Lichterfeld, M., Kavanagh, D.G., Williams, K.L., Moza, B., Mui, S.K., Miura, T., Sivamurthy, R., Allgaier, R., Pereyra, F., Trocha, A., et al. (2007). A viral CTL escape mutation leading to immunoglobulin-like transcript 4-mediated functional inhibition of myelomonocytic cells. *J. Exp. Med.* 204, 2813–2824.

1248

1249 Limon, J.J., So, L., Jellbauer, S., Chiu, H., Corado, J., Sykes, S.M., Raffatellu, M., and Fruman,

1250 D.A. (2014). mTOR kinase inhibitors promote antibody class switching via mTORC2 inhibition.

1251 *Proc. Natl. Acad. Sci. USA* 111, E5076–5085.

1252

1253 Lindqvist, M., van Lunzen, J., Soghoian, D.Z., Kuhl, B.D., Ranasinghe, S., Kranias, G.,

1254 Flanders, M.D., Cutler, S., Yudanin, N., Muller, M.I., et al. (2012). Expansion of HIV-specific T

1255 follicular helper cells in chronic HIV infection. *J. Clin. Invest.* 122, 3271–3280.

1256

1257 Locci, M., Havenar-Daughton, C., Landais, E., Wu, J., Kroenke, M.A., Arlehamn, C.L., Su, L.F.,

1258 Cubas, R., Davis, M.M., Sette, A., et al. (2013). Human circulating PD-1$^+$CXCR3$^-$CXCR5$^+$

1259 memory Tfh cells are highly functional and correlate with broadly neutralizing HIV antibody

1260 responses. *Immunity* 39, 758–769.

1261

1262 Lucas, C.L., Kuehn, H.S., Zhao, F., Niemela, J.E., Deenick, E.K., Palendira, U., Avery, D.T.,

1263 Moens, L., Cannons, J.L., Biancalana, M., et al. (2014). Dominant-activating germline

1264 mutations in the gene encoding the PI(3)K catalytic subunit p110δ result in T cell senescence

1265 and human immunodeficiency. *Nat. Immunol.* 15*,* 88–97.

1266

1267 Luo, W., Hawse, W., Conter, L., Trivedi, N., Weisel, F., Wikenheiser, D., Cattley, R.T., and

1268 Shlomchik, M.J. (2019). The AKT kinase signaling network is rewired by PTEN to control

1269 proximal BCR signaling in germinal center B cells. *Nat. Immunol.* 20, 736–746.

1270

1271 Marzec, M., Liu, X., Kasprzycka, M., Witkiewicz, A., Raghunath, P.N., El-Salem, M., Robertson,

1272 E., Odum, N., and Wasik, M.A. (2008). IL-2- and IL-15-induced activation of the rapamycin-

1273 sensitive mTORC1 pathway in malignant CD4+ T lymphocytes. *Blood* 111, 2181–2189.

1274

1275   Mason, R.D., Welles, H.C., Adams, C., Chakrabarti, B.K., Gorman, J., Zhou, T., Nguyen, R.,

1276   O'Dell, S., Lusvarghi, S., Bewley, C.A., et al. (2016). Targeted Isolation of Antibodies Directed

1277   against Major Sites of SIV Env Vulnerability. *PLoS Pathog.* 12, e1005537.

1278

1279   Matano, T., Kobayashi, M., Igarashi, H., Takeda, A., Nakamura, H., Kano, M., Sugimoto, C.,

1280   Mori, K., Iida, A., Hirata, T., et al. (2004). Cytotoxic T lymphocyte-based control of simian

1281   immunodeficiency virus replication in a preclinical AIDS vaccine trial. *J. Exp. Med.* 199, 1709–

1282   1718.

1283

1284   Mattapallil, J.J., Douek, D.C., Hill, B., Nishimura, Y., Martin, M., and Roederer, M. (2005).

1285   Massive infection and loss of memory CD4+ T cells in multiple tissues during acute SIV

1286   infection. *Nature* 434, 1093–1097.

1287

1288   McDonald, D., Wu, L., Bohks, S.M., KewalRamani, V.N., Unutmaz, D., and Hope, T.J. (2003).

1289   Recruitment of HIV and its receptors to dendritic cell-T cell junctions. *Science* 300, 1295–1297.

1290

1291   Mikell, I., Sather, D.N., Kalams, S.A., Altfeld, M., Alter, G., and Stamatatos, L. (2011).

1292   Characteristics of the earliest cross-neutralizing antibody response to HIV-1. *PLoS Pathog.* 7,

1293   e1001251.

1294

1295   Moir, S., Malaspina, A., Ogwaro, K.M., Donoghue, E.T., Hallahan, C.W., Ehler, L.A., Liu, S.,

1296   Adelsberger, J., Lapointe, R., Hwu, P., et al. (2001). HIV-1 induces phenotypic and functional

1297   perturbations of B cells in chronically infected individuals. *Proc. Natl. Acad. Sci. USA* 98,

1298   10362–10367.

1299

1300   Moody, M.A., Pedroza-Pacheco, I., Vandergrift, N.A., Chui, C., Lloyd, K.E., Parks, R.,

1301   Soderberg, K.A., Ogbe, A.T., Cohen, M.S., Liao, H.X., Gao, F., et al. (2016). Immune

1302   perturbations in HIV-1-infected individuals who make broadly neutralizing antibodies. *Sci.*

1303   *Immunol.* 1, aag0851.

1304

1305   Moore, P.L., Gray, E.S., Wibmer, C.K., Bhiman, J.N., Nonyane, M., Sheward, D.J., Hermanus,

1306   T., Bajimaya, S., Tumba, N.L., Abrahams, M.R., et al. (2012). Evolution of an HIV glycan-

1307   dependent broadly neutralizing antibody epitope through immune escape. *Nat. Med.* 18,

1308   1688–1692.

1309

1310   Moukambi, F., Rabezanahary, H., Rodrigues, V., Racine, G., Robitaille, L., Krust, B., Andreani,

1311   G., Soundaramourty, C., Silvestre, R., Laforge, M., et al. (2015). Early loss of splenic Tfh cells

1312   in SIV-infected rhesus macaques. *PLoS Pathog.* 11, e1005287.

1313

1314   Moukambi, F., Rabezanahary, H., Fortier, Y., Rodrigues, V., Clain, J., Benmadid-Laktout, G.,

1315   Zghidi-Abouzid, O., Soundaramourty, C., Laforge, M., and Estaquier, J. (2019) Mucosal T

1316   follicular helper cells in SIV-infected rhesus macaques: contributing role of IL-27. *Mucosal*

1317   *Immunol.* 12, 1038–1054.

1318

1319   Nakane, T., Nomura, T., Shi, S., Nakamura, M., Naruse, T.K., Kimura, A., Matano, T., and

1320   Yamamoto, H. (2013). Limited impact of passive non-neutralizing antibody immunization in

1321   acute SIV infection on viremia control in rhesus macaques. *PLoS ONE* 8, e73453.

1322

1323   Naruse, T.K., Chen, Z., Yanagida, R., Yamashita, T., Saito, Y., Mori, K., Akari, H., Yasutomi, Y.,

1324   Miyazawa, M., Matano, T., et al. (2010). Diversity of MHC class I genes in Burmese-origin

1325   rhesus macaques. *Immunogenetics* 62, 601–611.

1326

1327   Neil, S.J., Zang, T., and Bieniasz, P.D. (2008). Tetherin inhibits retrovirus release and is

1328   antagonized by HIV-1 Vpu. *Nature* 451, 425–430.

1329

1330 Ng, C.T., Jaworski, J.P., Jayaraman, P., Sutton, W.F., Delio, P., Kuller, L., Anderson, D.,

1331 Landucci, G., Richardson, B.A., Burton, D.R., et al. (2010). Passive neutralizing antibody

1332 controls SHIV viremia and enhances B cell responses in infant macaques. *Nat. Med.* 16, 1117–

1333 1119.

1334

1335 Nguyen, H., Ramana, C.V., Bayes, J., and Stark, G.R. (2001). Roles of phosphatidylinositol 3-

1336 kinase in interferon-gamma-dependent phosphorylation of STAT1 on serine 727 and activation

1337 of gene expression. *J. Biol. Chem.* 276, 33361–33368.

1338

1339 Nishimura, Y., Gautam, R., Chun, T.W., Sadjadpour, R., Foulds, K.E., Shingai, M., Klein, F.,

1340 Gazumyan, A., Golijanin, J., Donaldson, M., et al. (2017). Early antibody therapy can induce

1341 long-lasting immunity to SHIV. *Nature* 543, 559–563.

1342

1343 Nomura, T., Yamamoto, H., Shiino, T., Takahashi, N., Nakane, T., Iwamoto, N., Ishii, H.,

1344 Tsukamoto, T., Kawada, M., Matsuoka, S., et al. (2012). Association of major histocompatibility

1345 complex class I haplotypes with disease progression after simian immunodeficiency virus

1346 challenge in Burmese rhesus macaques. *J. Virol.* 86, 6481–6490.

1347

1348 Nutt, S.L., Hodgkin, P.D., Tarlinton, D.M., and Corcoran, L.M. (2015). The generation of

1349 antibody-secreting plasma cells. *Nat. Rev. Immunol.* 15, 160–171.

1350

1351 Omori, S.A., Cato, M.H., Anzelon-Mills, A., Puri, K.D., Shapiro-Shelef, M., Calame, K., and

1352 Rickert, R.C. (2006). Regulation of class-switch recombination and plasma cell differentiation

1353 by phosphatidylinositol 3-kinase signaling. *Immunity* 25, 545–557.

1354

1355 Perfetto, S.P., Chattopadhyay, P.K., Lamoreaux, L., Nguyen, R., Ambrozak, D., Koup, R.A.,

1356 and Roederer, M. (2006). Amine reactive dyes: an effective tool to discriminate live and dead

1357 cells in polychromatic flow cytometry. *J. Immunol. Methods* 313, 199–208.

1358

1359 Petrovas, C., Yamamoto, T., Gerner, M.Y., Boswell, K.L., Wloka, K., Smith, E.C., Ambrozak,

1360 D.R., Sandler, N.G., Timmer, K.J., Sun, X., et al. (2012). CD4 T follicular helper cell dynamics

1361 during SIV infection. *J. Clin. Invest.* 122, 3281–3294.

1362

1363 Qiao, X., He, B., Chiu, A., Knowles, D.M., Chadburn, A., and Cerutti, A. (2006). Human

1364 immunodeficiency virus 1 Nef suppresses CD40-dependent immunoglobulin class switching

1365 in bystander B cells. *Nat. Immunol.* 7, 302–310.

1366

1367 Querec, T.D., Akondy, R.S., Lee, E.K., Cao, W., Nakaya, H.I., Teuwen, D., Pirani, A., Gernert,

1368 K., Deng, J., Marzolf, B., et al. (2009). Systems biology approach predicts immunogenicity of

1369 the yellow fever vaccine in humans. *Nat. Immunol.* 10, 116–125.

1370

1371 Ray, J.P., Staron, M.M., Shyer, J.A., Ho, P.C., Marshall, H.D., Gray, S.M., Laidlaw, B.J., Araki,

1372 K., Ahmed, R., Kaech, S.M., et al. (2015). The interleukin-2-mTORc1 kinase axis defines the

1373 signaling, differentiation, and metabolism of T helper 1 and follicular B helper T cells. *Immunity*

1374 43, 690–702.

1375

1376 Recher, M., Lang, K.S., Hunziker, L., Freigang, S., Eschli, B., Harris, N.L., Navarini, A., Senn,

1377 B.M., Fink, K., Lötscher, M., et al. (2004). Deliberate removal of T cell help improves virus-

1378 neutralizing antibody production. *Nat. Immunol.* 5, 934–942.

1379

1380 Recher, M., Lang, K.S., Navarini, A., Hunziker, L., Lang, P.A., Fink, K., Freigang, S., Georgiev,

1381 P., Hangartner, L., Zellweger, R., et al. (2007). Extralymphatic virus sanctuaries as a

1382 consequence of potent T-cell activation. *Nat. Med.* 13, 1316–1323.

1383

1384 Reuschl, A.K., Mesner, D., Shivkumar, M., Whelan, M.V.X., Pallett, L.J., Guerra-Assunção,

1385 J.A., Madansein, R., Dullabh, K.J., Sigal, A., Thornhill, J.P., et al. (2022). HIV-1 Vpr drives a

tissue residency-like phenotype during selective infection of resting memory T cells. *Cell Rep.* 39, 110650.

Rickles, R.J., Botfield, M.C., Weng, Z., Taylor, J.A., Green, O.M., Brugge, J.S., and Zoller, M.J. (1994). Identification of Src, Fyn, Lyn, PI3K and Abl SH3 domain ligands using phage display libraries. *EMBO J.* 13, 5598–5604.

Samer, S., Chowdhury, A., Wiche Salinas, T.R., Estrada, P.M.D.R., Reuter, M., Tharp, G., Bosinger, S., Cervasi, B., Auger, J., Gill, K., et al. (2023). Lymph-Node-based CD3[+] CD20[+] cells emerge from membrane exchange between T follicular helper cells and B cells and increase their frequency following simian immunodeficiency virus infection. *J. Virol.* 97, e0176022.

Sammicheli, S., Kuka, M., Di Lucia, P., de Oya, N.J., De Giovanni, M., Fioravanti, J., Cristofani, C., Maganuco, C.G., Fallet, B., Ganzer, L., et al. (2016). Inflammatory monocytes hinder antiviral B cell responses. *Sci. Immunol.* 1, pii: eaah6789.

Sander, S., Chu, V.T., Yasuda, T., Franklin, A., Graf, R., Calado, D.P., Li, S., Imami, K., Selbach, M., Di Virgilio, M., et al. (2015). PI3 kinase and FOXO1 transcription factor activity differentially control B cells in the germinal center light and dark zones. *Immunity* 43, 1075–1086.

Sarbassov, D.D., Guertin, D.A., Ali, S.M., and Sabatini, D.M. (2005) Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. *Science* 307, 1098–1101.

Schindler, M., Münch, J., Brenner, M., Stahl-Hennig, C., Skowronski, J., and Kirchhoff, F. (2004). Comprehensive analysis of nef functions selected in simian immunodeficiency virus-infected macaques. *J. Virol.* 78, 10588–10597.

1414 Schindler, M., Münch, J., Kutsch, O., Li, H., Santiago, M.L., Bibollet-Ruche, F., Müller-Trutwin,

1415 M.C., Novembre, F.J., Peeters, M., Courgnaud, V., et al. (2006). Nef-mediated suppression of

1416 T cell activation was lost in a lentiviral lineage that gave rise to HIV-1. *Cell* 125, 1055–1067.

1417

1418 Sette, A., Sidney, J., Southwood, S., Moore, C., Berry, J., Dow, C., Bradley, K., Hoof, I., Lewis,

1419 M.G., Hildebrand, W.H., et al. (2012). A shared MHC supertype motif emerges by convergent

1420 evolution in macaques and mice, but is totally absent in human MHC molecules.

1421 *Immunogenetics* 64, 421–434.

1422

1423 Sheehy, A.M., Gaddis, N.C., Choi, J.D., and Malim, M.H. (2002). Isolation of a human gene

1424 that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* (2002) 418, 646–

1425 650.

1426

1427 Shi, S., Seki, S., Matano, T., and Yamamoto, H. (2013). IL-21-producer CD4$^+$ T cell kinetics

1428 during primary simian immunodeficiency virus infection. *Microbes Infect.* 15, 697–707.

1429

1430 Shulman, Z., Gitlin, A.D., Weinstein, J.S., Lainez, B., Esplugues, E., Flavell, R.A., Craft, J.E.,

1431 and Nussenzweig, M.C. (2014). Dynamic signaling by T follicular helper cells during germinal

1432 center B cell selection. *Science* 345, 1058–1062.

1433

1434 Silveira, E.L., Kasturi, S.P., Kovalenkov, Y., Rasheed, A.U., Yeiser, P., Jinnah, Z.S., Legere,

1435 T.H., Pulendran, B., Villinger, F., and Wrammert, J. (2015). Vaccine-induced plasmablast

1436 responses in rhesus macaques: phenotypic characterization and a source for generating

1437 antigen-specific monoclonal antibodies. *J. Immunol. Methods* 416, 69–83.

1438

1439 Sommerstein, R., Flatz, L., Remy, M.M., Malinge, P., Magistrelli, G., Fischer, N., Sahin, M.,

1440 Bergthaler, A., Igonet, S., Ter Meulen, J., et al. (2015). Arenavirus glycan shield promotes

1441 neutralizing antibody evasion and protracted infection. *PLoS Pathog.* 11, e1005276.

1442

1443   Srinivasan, L., Sasaki, Y., Calado, D.P., Zhang, B., Paik, J.H., DePinho, R.A., Kutok, J.L.,

1444   Kearney, J.F., Otipoby, K.L., and Rajewsky, K. (2009). PI3 kinase signals BCR-dependent

1445   mature B cell survival. *Cell* 139, 573–586.

1446

1447   Stolp, B., Reichman-Fried, M., Abraham, L., Pan, X., Giese, S.I., Hannemann, S., Goulimari,

1448   P., Raz, E., Grosse, R., and Fackler, O.T. (2009). HIV-1 Nef interferes with host cell motility by

1449   deregulation of Cofilin. *Cell Host Microbe* 6, 174–186.

1450

1451   Stolp, B., Imle, A., Coelho, F.M., Hons, M., Gorina, R., Lyck, R., Stein, J.V., and Fackler, O.T.

1452   (2012). HIV-1 Nef interferes with T-lymphocyte circulation through confined environments in

1453   vivo. *Proc. Natl. Acad. Sci. USA* 109, 18541–18546.

1454

1455   Swingler, S., Zhou, J., Swingler, C., Dauphin, A., Greenough, T., Jolicoeur, P., and Stevenson,

1456   M. (2008). Evidence for a pathogenic determinant in HIV-1 Nef involved in B cell dysfunction

1457   in HIV/AIDS. *Cell Host Microbe* 4, 63–76.

1458

1459   Takahashi, N., Nomura, T., Takahara, Y., Yamamoto, H., Shiino, T., Takeda, A., Inoue, M., Iida,

1460   A., Hara,H., Shu, T., et al. (2013). A novel protective MHC-I haplotype not associated with

1461   dominant Gag-specific CD8$^+$ T-cell responses in SIV$_{mac239}$ infection of Burmese rhesus

1462   macaques. *PLoS ONE* 8, e54300.

1463

1464   Terahara, K., Ishii, H., Nomura, T., Takahashi, N., Takeda, A., Shiino, T., Tsunetsugu-Yokota,

1465   Y., and Matano, T. (2014). Vaccine-induced CD107a$^+$ CD4$^+$ T cells are resistant to depletion

1466   following AIDS virus infection. *J. Virol.* 88, 14232–14240.

1467

1468   Tomaras, G.D., Yates, N.L., Liu, P., Qin, L., Fouda, G.G., Chavez, L.L., Decamp, A.C., Parks,

1469   R.J., Ashley, V.C., Lucas, J.T., et al. (2008). Initial B-cell responses to transmitted human

immunodeficiency virus type 1: virion-binding immunoglobulin M (IgM) and IgG antibodies followed by plasma anti-gp41 antibodies with ineffective control of initial viremia. *J. Virol.* 82, 12449–12463.

Volcic, M., Sparrer, K.M.J., Koepke, L., Hotter, D., Sauter, D., Stürzel, C.M., Scherer, M., Stamminger, T., Hofmann, T.G., Arhel, N.J., et al. (2020). Vpu modulates DNA repair to suppress innate sensing and hyper-integration of HIV-1. *Nat. Microbiol.* 5, 1247–1261.

Wang, J., Sohn, H., Sun, G., Milner, J.D. and Pierce, S.K. (2014). The autoinhibitory C-terminal SH2 domain of phospholipase C-γ2 stabilizes B cell receptor signalosome assembly. *Sci. Signal.* 7, ra89.

Whitney, J.B., Hill, A.L., Sanisetty, S., Penaloza-MacMaster, P., Liu, J., Shetty, M., Parenteau, L., Cabral, C., Shields, J., Blackmore, S., et al. (2014). Rapid seeding of the viral reservoir prior to SIV viraemia in rhesus monkeys. *Nature* 512, 74–77.

Wülfing, C., Sjaastad, M.D., and Davis, M.M. (1998). Visualizing the dynamics of T cell activation: intracellular adhesion molecule 1 migrates rapidly to the T cell/B cell interface and acts to sustain calcium levels. *Proc. Natl. Acad. Sci. USA* 95, 6302–6307.

Xu, H., Wang, X., Malam, N., Aye, P.P., Alvarez, X., Lackner, A.A., and Veazey, R.S. (2015). Persistent simian immunodeficiency virus infection drives differentiation, aberrant accumulation, and latent infection of germinal center follicular T helper cells. *J. Virol.* 90,1578–1587.

Xu, W., Santini, P.A., Sullivan, J.S., He, B., Shan, M., Ball, S.C., Dyer, W.B., Ketas, T.J., Chadburn, A., Cohen-Gould, L., et al. (2009). HIV-1 evades virus-specific IgG2 and IgA

responses by targeting systemic and intestinal B cells via long-range intercellular conduits. *Nat. Immunol.* 10, 1008–1017.

Yamada, E., Nakaoka, S., Klein, L., Reith, E., Langer, S., Hopfensperger, K., Iwami, S., Schreiber, G., Kirchhoff, F., Koyanagi, Y., et al. (2018) Human-specific adaptations in Vpu conferring anti-tetherin activity are critical for efficient early HIV-1 replication in vivo. *Cell Host Microbe* 23, 110–120.e7.

Yamamoto, H., Kawada, M., Takeda, A., Igarashi, H. and Matano, T. (2007). Post-infection immunodeficiency virus control by neutralizing antibodies. *PLoS ONE* 2, e540.

Yan, J., Shun, M.C., Zhang, Y., Hao, C., and Skowronski, J. (2019). HIV-1 Vpr counteracts HLTF-mediated restriction of HIV-1 infection in T cells. *Proc. Natl. Acad. Sci. USA.* 116, 9568–9577.

Zhang, F., Wilson, S.J., Landford, W.C., Virgen, B., Gregory, D., Johnson, M.C., Munch, J., Kirchhoff, F., Bieniasz, P.D., and Hatziioannou, T. (2009). Nef proteins from simian immunodeficiency viruses are tetherin antagonists. *Cell Host Microbe* 6, 54–67.

Zotos, D., Coquet, J.M., Zhang, Y., Light, A., D'Costa, K., Kallies, A., Corcoran, L.M., Godfrey, D.I., Toellner, K.M., Smyth, M.J., et al. (2010). IL-21 regulates germinal center B cell differentiation and proliferation through a B cell-intrinsic mechanism. *J. Exp. Med.* 207, 365–378.

 Supplemental Information Legends

 Fig. 1-Supplement 1. Profile of SIV$_{mac239}$-NAb non-inducer subgroup cohort.

 Summary of a SIV$_{mac239}$-NAb non-inducer subgroup cohort (*n* = 19), composed from previously

 partially characterized, MHC class I haplotype-balanced naïve animals. Symbols A+, B+, E+,

 J+ and D+ represent possession of MHC-I haplotypes *90-120-Ia*, *90-120-Ib*, *90-010-Ie*, *90-*

 088-Ij and *90-010-Id*, respectively. For the titers, - represents negative; +/- represents

 neutralization in three out of four wells in a quadruplicate test. Gray shadings indicate

 euthanasia due to AIDS progression. Mamu-A and Mamu-B alleles of interest in this study are

 listed. These NAb non-inducers were previously partially reported for their plasma viral loads

 and MHC-I alleles (Nomura et al., 2012; Takahashi et al., 2013; Nakane et al., 2013; Iwamoto

 et al., 2014). R01-011 was previously partially reported for NAb titers measured with the same

 method using MT4 cells as targets (Kawada et al., 2007), which derived comparable results.

 Fig. 1-Supplement 2. IgG class-switched NAbs in the inducers.

 Titration of SIV$_{mac239}$-neutralizing IgGs in the NAb inducers. SIV$_{mac239}$-specific IgGs, purified

 from pools of plasma with SIV$_{mac239}$-specific NAb titers, were obtained from each animal as

 described (Yamamoto et al., 2007) and examined for their neutralizing activity by 10 TCID$_{50}$

 SIV$_{mac239}$ killing on MT4-R5 cells. NAb-negative anti-SIV IgG and control IgG were prepared

 from a representative animal (R01-011) with NAb-negative plasma and pooled plasma of

 uninfected rhesus macaques, respectively.

 Fig.1-Supplement 3. Anti-SIV binding IgG profiles in NAb inducers and non-inducers.

 (A) Temporal changes in SIV$_{mac251}$ Env gp120-specific antibody titers in the NAb non-inducers

 without rapid progression [NAb-negative non-rapid progressors, NAb(-) non-RPs] (left, *n* = 11)

 and NAb inducers [NAb(+)] (right, *n* = 9). NAb non-inducers showing rapid progression

 (euthanized with AIDS onset within approximately 1 year) were not included except for

 macaque R10-001 (shown in the left for aligned comparison). Samples for non-RPs R06-001

and R10-002 were unavailable.

(B) Unpaired t test of log-transformed anti-SIV Env Ab titers in NAb(-) non-RPs and NAb

inducers. Bars represent geometric mean absorbance within each group.

(C) Lysed SIV_{mac} virion linear antigen-binding at year 1 p.i. in NAb(-) non-RPs (left, n = 12)

and NAb inducers (right, n = 9). Plasma anti-SIV_{mac251} IgGs were detected using a commercial

western blotting system against the parental strain SIV_{mac251} (ZeptoMetrix). White arrowheads

indicate Env gp120-specific bands. Experiments were performed twice with comparable

results.

1557

Fig. 1-Supplement 4. Env sequence variation pattern is not a major characteristic of

NAb induction.

SIV_{mac239} Env sequence variations before and after NAb induction. For each animal, the upper

time point shows pre-induction and the lower time point (in purple) shows post-induction (first

or second time point with NAb titer > 1:2). Dominant residue changes are shown as follows:

gp120 (gray), variable regions V1-V5 (blue), and gp41 (blue gray). Subdominant mutations

are uncolored. Unlabeled gray and black shadings show subdominant and dominant deletion

of residues. Parental SIV_{mac239} sequence is listed for V1-V5. Residues mutated in multiple

animals within V1-V5 are shown in red.

1567

Fig. 3-Supplement 1. Selection of Nef-G63E mutant SIV is linked with Nef$_{62-70}$-specific

CD8$^+$ T-cell response positivity.

(A) Nef$_{62-70}$-specific CD8+ T-cell responses around week 20 post-SIV challenge in seven

viremic NAb non-inducers possessing either of the MHC-I alleles Mamu-B*039:01, Mamu-

B*004:01 or Mamu-A1*032:02.

(B) Temporal supernatant SIV p27 concentrations after wild type (WT) or Nef-G63E mutant

SIV infection at multiplicity of infection (MOI) 0.001 on HSC-F cells. Data represent one of two

independent experiments performed in triplicate. Bars: mean ± SEM.

(C) Comparison of plasma viral loads in Nef-G63E-selecting NAb non-inducers (n = 7) versus

non-inducers at indicated time points post-SIV challenge. Bars represent geometric means.

log10-transformed values are compared by unpaired t tests.

1579

Fig. 4-Supplement 1. Decreased Akt hyperactivation properties of Nef-G63E mutant SIV.

(A) Plasma ferritin levels (ng/ml) in examined NAb non-inducers versus Nef-G63E mutant SIV-

selecting NAb inducers around year 1 post-infection. Bars represent mean in each group. Log-

transformed values compared by unpaired t test.

(B) Relative levels of pAkt Ser473 (left panel) and threonine (Thr) 308 (right panel) in SIV Nef$^+$

HSR5.4 cells (MOI 0.1, 4 days p.i.).

(C) Relative pAkt Ser473 levels in Nef$^+$ HSC-F cells without stimuli and transiently pulsed with

IFN-γ, IL-2 or SIV$_{mac}$ Env gp130 after infection (MOI 0.2, 1 day).

(D) Gene expression significantly differing between WT and Nef-G63E SIV infection (MOI 5,

1 day). Black bars show potentially Akt-related genes. Asterisks show dynamics concordant

with the Nef-G63E phenotype (PI3K-pAkt Ser473 downstream downregulation or inhibition

upregulation).

(E) Representative flow cytometric gating (NAb inducer R01-012, wk 30 p.i.) of peripheral

blood CXCR3$^-$CXCR5$^+$PD-1$^+$ memory follicular CD4+ T cells. Cells were gated as CD3$^+$CD8$^-$

CD4$^+$CD95$^+$CXCR5$^+$PD-1$^+$CXCR3$^-$ live singlet PBMCs. Available samples of six NAb inducers

showing peak Nef-G63E mutant selection around week 30 p.i. and thirteen viremic NAb non-

inducers were tracked.

(F) Comparison of peripheral CXCR3$^-$CXCR5$^+$PD-1$^+$ memory Tfh cell frequencies between

pre-SIV infection and around week 30 post-infection in examined NAb non-inducers

developing disease before week 60 (left: n = 5) and alive more than 70 weeks (middle: n = 8)

and Nef-G63E-detected NAb inducers alive more than 70 weeks (right: n = 6). Analyzed by

Wilcoxon signed-rank tests.

MFIs relative to those in uninfected controls (%) are shown. **: $P < 0.01$, ***: $P < 0.001$, n.s.:

not significant between WT and Nef-G63E mutant by unpaired t tests (B-C). Data represent

one of two (B, C) independent experiments performed in triplicate (C: unstimulated) or

quadruplicate (B, C: PI3K-pulsed). Bars: mean ± SEM.

1606

Fig. 4-Supplement 2. Nef-G63R mutant SIV properties.

(A) Comparison of frequencies of macaques selecting for Nef-G63R mutant in plasma

between NAb non-inducers and inducers. Compared by Fisher's exact test.

(B) Temporal relationship of Nef G63R frequencies in plasma virus in a NAb non-inducer

macaque R06-034 preferentially selecting for Nef-G63R but not Nef-G63E mutant. Black

boxes (left Y axis) show $\log_2$ NAb titers; red triangles and blue diamonds (right Y axis) show

percentage of G63E and G63R mutations detected by subcloning (15 clones/point on average),

respectively.

(C) Left: representative histograms of relative pAkt serine (Ser) 473 levels in p27$^+$

subpopulations after WT, Nef-G63E or Nef-G63R mutant SIV infection at MOI 0.1 on HSC-F

cells. Numbers show pAkt Ser473 mean fluorescence intensities (MFIs) for each. Right:

Deviation of pAkt Ser473 MFIs in p27$^+$ HSC-F cells compared with mean MFI of uninfected

cells. Analyzed by one-way ANOVA with Tukey's post-hoc multiple comparison tests. Data

represent one of two independent experiments performed in quadruplicate with similar results.

1621

Fig. 5-Supplement 1. Machine learning-verified morphological gating of Nef-invaded B

cells *in vivo*.

(A) Representative gating strategy plus schematic of inguinal lymph node Nef$^+$CD19$^+$ B cells

of an SIV$_{mac239}$-infected rhesus macaque analyzed by ImageStreamX MK II imaging flow

cytometry (R10-012, week 160 p.i.). First-step incorporation of Live/Dead staining gates out

CD19$^+$ B cells staining positive for Nef potentially due to pre-experimental membrane

permeation (e.g., image #246896, #248952 and #71859). Second-step dual gating for

intracellular Nef signal pixel Haralick variance mean (H Variance mean, X axis), a textural

feature calculated from a gray level co-occurrence matrix (GLCM) for each image representing

adjacent pixel signal strength heterogeneity and Nef signal intensity threshold (Y axis) gates

out CD19$^+$ B cells showing potentially non-specific, strong intracellular binary clustered

staining pixels for Nef deriving a large variance (X axis), and/or cells stained overtly strong for

Nef and hence becoming void for feasible image verification (high threshold) (Y axis), both

likely originating stochastically from post-experimental rupturing procedures including

membrane permeation (e.g. image #18541). This two-step noise cancellation results in

acquisition of a Live/Dead$^-$-Nef signal Haralick variance mean^{lo}-Nef signal intensity threshold^{lo}-

Nef$^+$CD19$^+$ B-cell population (e.g. image #116318, #116250 and #132927) with a pericellular

Nef^{int-lo} staining, biologically concordant with Nef membrane anchoring, resembling Nef$^+$ cells

experimentally generated *in vitro* and showed the highest score in IDEAS6.3 machine learning

module scoring for segregation of typical void vs. valid images (upper right). This gating was

implemented for counting and analysis of Env-specific B-cell Nef invasion *in vivo* (**Figure 5**).

(B) Detection of contact-dependent Nef invasion from SIV-infected HSC-F CD4$^+$ T cells to

cocultured Ramos B cells *in vitro*. One typical image of an intercellular conduit (SSC, purple)

with Nef (green) protrusions from infected HSC-F cells to CD19$^+$ (red) Ramos B cells from two

independent experiments are shown.

1647

1648 **Fig. 7-Supplement 1. Full gating strategy of SIV$_{mac239}$ Env gp140-specific B-cell**

1649 **responses.**

1650 Full gating (R02-004, week 32 p.i.) of SIV Env gp140-specific memory B cells (B$_{mem}$) and

1651 plasmablasts (PBs) shown in **Figure 7**.

A

SIVmac239 Nef

B

C Nef-G63E selection

P = 0.0009

NAb- NAb+
(2/19) (7/9)

Status: ▷ NAb (-) ▷ NAb (+/-) ▷ NAb (+)

D Temporal Nef-G63E mutant selection vs NAb induction

SIV 100% neutralizing titers (log₂)

Mutant SIV frequencies (%)

SIVmac239-NAb
Nef G63E
Nef G63R

Weeks post-SIV challenge

A

Surface downregulation

CD3 · CD4 · MHC Class I · CXCR4 · BST-2

WT · Nef-G63E · Uninfected

Gated on singlet CD4loNef+

B

MOI 0.1 Day 3

pAkt Ser473

1059 · 896 · 746

Gated on singlet p27+

WT · Nef-G63E · Uninfected

P = 0.0009

ΔMFI (+ mean of uninfected)

WT · Nef G63E

C

Accaleration w/upstream PI3K stimuli Day 3→Day 1

Serum starvation

P = 0.99 · P = 0.0017

pAkt relative MFIs in Nef+ cells (%)

WT · Nef-G63E

Starvation

% serum in culture

High MOI

P = 0.0011

ΔMFI (+ mean of uninfected)

WT · Nef G63E

D

Intracellular PI3K-pathway binding

CD4lop27+ singlet HSC-F

68.3

mTORC2 — GβL/mLST8 · mTOR

PI3 Kinase — PI3Kp85 · PI3Kp110α

WT · Nef-G63E · WT (isotype) · Nef G63E (isotype)

P = 0.039 · P = 0.028 · P = 0.022 · P = 0.043

Binding Index (ΔMFI)

WT · Nef G63E

E

IP: αSin1 · Lysate

WT · Nef-G63E · NC · WT · Nef-G63E · NC · Ladder · kDa

Nef ► · 40 · 37 · 30

IB: αNef

F

Cell death

CD4lo singlet HSC-F

20.9 · 29.0

Comp-APC-A :: AnnexinV APC-A

WT · Nef-G63E

P = 0.0041

Annexin V+ (% of CD4-low)

WT · Nef G63E

A

Gated on Lin [CD3/8/(10)14/16]^lo-Live singlet PBMCs

B

NAb non-inducers

NAb inducers (wk 30 Nef-G63E+)

- ■ R07-004
- ▲ R07-009
- ▼ R06-001
- ◆ R01-011
- ● R05-007
- ▣ R06-034
- △ R08-005
- ▽ R09-011
- ◇ R10-001
- ○ R06-022
- ✕ R10-002
- ✳ R10-006

- ■ R06-032
- ▲ R01-012
- ▼ R03-015
- ✕ R02-004
- ● R01-010
- ▣ R06-019

SIV Env gp140-specific IgG+ plasmablasts/million PBMCs

time post-SIV challenge

C

wk 30 — at NAb induction vs yr 1

P = 0.0131

P = 0.0030

SIV Env gp140-specific IgG+ plasmablasts/million PBMCs

NAb- / Nef-G63E+ NAb+

D

Temporal 2-D plot of Env gp140-specific Bmem vs PB

D_i = GC output attractor 2 (Nef-G63E+, at NAb induction)

D_n = GC output attractor 1 (NAb-, at yr 1)

gp140-specific IgG+ plasmablasts/million PBMCs

gp140-specific IgG+ memory B cells/million PBMCs

MHC-I/ID	SIV$_{mac239}$ neutralization at 1:2 plasma dilution by 10TCID$_{50}$ killing assay (wks p.i.)									pVL at 6 mo	Nef G63E at 1 yr	Detected *Mamu* alleles
	wk11–20	wk21–30	wk31–40	wk41–50	wk51–60	wk61–70	wk71–80	wk81–90	wk91–104			
A+												
R06-037		wk25 -		wk46 +/-			wk72 -	wk87 -	wk95 -	6.4×10^2	- (pVL < 400)	
R07-001		wk25 -		wk42 -				wk88 +/-	wk96 -	< 400	- (pVL < 400)	*A1*032:02*
R07-004		wk25 -		wk46 -		wk64 -	wk74 -	wk86 -		4.5×10^3	+	*B*039:01*
R07-009		wk25 -		wk46 -		wk64 -	wk74 -			4.5×10^3	-	
B+												
R06-001		wk25 -		wk48 -	wk60 -		wk80 -	wk90 -		8.2×10^3	-	
R06-039		wk25 -		wk46 -	wk56 -					8.5×10^6	-	*A1*032:02 , B*004:01*
E+												
R01-011		wk22 -	wk40 -		wk52 -		wk71 -	wk85 -	wk104 -	1.3×10^5	-	*B*004:01*
R05-007		wk26 -			wk55 +/-		wk71 -	wk89 -	wk97 -	1.6×10^4	-	
R08-007		wk21 -		wk46 -		wk69 -		wk82 +/-		8.5×10^6	-	
R09-011	wk18 -		wk36 -		wk52 -					8.5×10^6	-	
R06-038		wk25 -			wk58 +/-		wk80 +/-		wk95 -	1.1×10^5	-	
J+												
R04-014	wk12 -	wk25 -	wk40 -							1.5×10^5	+ (wk40)	*B*039:01*
R06-022	wk12 -	wk23 -								1.4×10^7	- (wk23)	*B*039:01*
R10-001	wk14 -	wk30 -	wk37 -							1.9×10^7	- (wk37)	*B*039:01*
D+												
R05-006		wk25 -		wk48 -			wk80 -	wk90 -	wk98 -	5.4×10^2	-	*A1*032:02 , B*004:01*
R06-034		wk25 -		wk42 -		wk61 -	wk80 +/-	wk88 +/-	wk96 -	4.5×10^3	-	*A1*032:02 , B*004:01*
R08-005		wk25 -		wk48 -		wk68 -			wk100 -	9.0×10^2	-	*A1*032:02 , B*004:01*
Others												
R10-002	wk14 -				wk59 -				wk104 -	5.7×10^5	-	
R10-006	wk14 -	wk30 -			wk55 -					1.0×10^6	-	

Group	Rhesus ID	mg/ml (IgG)	Collected/pooled (wks post-challenge)	10 TCID$_{50}$ SIV$_{mac239}$ 100% killing titer
NAb (+)	R06-032	30	wk 76	1:16
	R01-012	30	wk 73–87	1:32
	R03-015	30	wk 112	1:8
	R02-004	30	wk 76	1:4
	R01-010	30	wk 77	1:16
	R06-019	30	wk 85	1:4
	R-360	30	wk 60–86	1:4
	R02-007	30	wk 42–70	1:32
	R05-010	30	wk 71	1:64
NAb (-)	R01-011	42	wk 104	< 1:2
uninfected	pooled	36	-	< 1:2

A

NAb(-) non-RPs

AU (OD 450 nm Abs)

weeks post-SIV challenge

- ■ R06-037 △ R08-007
- ▲ R07-001 ▽ R06-038
- ▼ R07-004 ○ R05-006
- ◆ R07-009 ✕ R06-034
- ● R01-011 + R08-005
- ▢ R05-007 ◇ R10-001

NAb(+)

weeks post-SIV challenge

- ■ R02-007 △ R02-004
- ▲ R06-019 ▽ R03-015
- ▼ R01-012 ◇ R06-032
- ◆ R-360
- ● R01-010
- ▢ R05-010

B

Anti-Env Ab 1 yr

P = 0.044

AU (OD 450 nm Abs)

NAb- non-RPs   NAb+

C

NAb- non-RPs   NAb+

R06-037, R07-001, R07-004, R07-009, R06-001, R01-011, R05-007, R08-007, R06-038, R05-006, R06-034, R08-005   R02-007, R06-019, R01-010, R05-010, R02-004, R03-015, R06-032, R01-012, R-360

- gp160 (*env*)
- gp120 (*env*)
- p66 (*pol*)
- p55 (*gag*)
- gp41 (*env*)
- p31 (*pol*)
- p27 (*gag*)
- p17 (*gag*)
- Serum control

Top panel region labels:

gp120   V1 (113-151)   V2 (194-206)   V3 (311-344)

L V I K K T C N K S E T D R W G L T K S I T T T A S T T S T T A S A K V D W V N E T S S G A G E Q G N N T G N E S R C C N P K Y C R R P G N K T V L P V T I W S G L V F H S Q P I N D R P K Q A W C W K K

Position numbers (top): 24 57 98 99 103 110 120 130 132 135 136 138 140 150 190 201 211 24/253 254 303 320 330 340 345 349 351

Patient	wk	
R06-032	wk30	
	wk46	
R01-012	wk30	
	wk48	
R03-015	wk20	
	wk53	
R02-004	wk40	
	wk76	
R01-010	wk18	
	wk57	
R06-019	wk24	
	wk69	
K-360	wk12	
	wk60	
R02-007	wk12	
	wk24	
R05-010	wk12	
	wk20	

Bottom panel region labels:

V4 (402-432)   V5 (459-491)   gp41

D V Y K E L C K N N V F L N W V E D R N T A N Q K P K E Q H K R N Y Y P C N S T V T S L I A N I D W I D G N Q T N I T M S A E V A E L Y R D T G V F T S N K Q E K E R G G T I P L A V L I

Position numbers (bottom): 352 362 375 398 400 410 417 420 425 430 460 470 480 490 511 516 518 527 528 594 62/ 633 689 739 747 749 750 751 753 756 789 797 800 802 836 837 845 863

Patient	wk
R06-032	wk30 / wk46
R01-012	wk30 / wk48
R03-015	wk20 / wk53
R02-004	wk40 / wk76
R01-010	wk18 / wk57
R06-019	wk24 / wk69
K-360	wk12 / wk60
R02-007	wk12 / wk24
R05-010	wk12 / wk20

A

Nef$_{62-70}$ QW9-specific responses in NAb non-inducers (viremic/Nef-G63E non-selecting)

Y-axis: Nef$_{62-70}$-specific CD8+ T-cell frequencies (/million PBMCs)

X-axis: R06-039, R01-011, R06-022, R10-001, R05-006, R06-034, R08-005

B

Y-axis: SIV p27 (ng/ml)
X-axis: Days post-SIV infection on HSC-F cells

Legend: WT, Nef-G63E

C

Y-axis: Plasma viral loads (vRNA copies/ml)

3 mo	6 mo	1 yr	1.5 yr
P = 0.2495	*P* = 0.5078	*P* = 0.8928	*P* = 0.6488

NAb−   NAb+
          Nef-G63E+

A

R3

CD19-PC5.5 — Intensity_MC_Ch11

R4

Live/Dead-Aqua — Intensity_MC_Ch08

R4 — Intensity_MC_Ch11

R5

Intensity_MC_Ch02

R5

Intensity_MC_Ch11 — Intensity_MC_Ch02

Nef-AF488

Intracellular pixel Nef signal Intensity threshold

Signal intensity threshold (int/hi)

Signal intensity threshold (lo)

Live/Dead⁻CD19⁺ Nef⁺ B cells

R5

Intensity_Threshold(M02 Ch2 30)_ — H Variance Mean_M02_Ch02_1

R7

IDEAS6.3 machine learning-based 2-D segregation of typical valid vs. void Live/Dead⁺Nef⁺CD19⁺ B-cell images

Good,bad

Void

Valid

Intensity_Threshold(M02 Ch0 — H Variance Mean_M02_Ch02_

Intracellular pixel Nef signal strength distribution

2-D adjacent pixel signal heterogeneity (++)
→Haralick variance mean (int-hi)

2-D adjacent pixel signal heterogeneity (+/-)
→Haralick variance mean (lo)

Live/Dead⁺CD19⁺Nef⁺ B cells (void)

Ch01 | Ch02 | Ch08 | Ch11 | Ch02/Ch08/Ch11
246896
248952
71859
7 μm

Live/Dead⁻CD19⁺Nef⁺-(Intracellular Nef Haralick variance mean^int/hi or Nef Intensity threshold^int/hi) B cells (void)

Ch01 | Ch02 | Ch08 | Ch11 | Ch02/Ch08/Ch11
18541
7 μm

Live/Dead⁻CD19⁺Nef⁺-Intracellular Nef Haralick variance mean^lo-Nef Intensity threshold^lo B cells (valid)

Ch01 | Ch02 | Ch08 | Ch11 | Ch02/Ch08/Ch11
116318
116250
132927
7 μm

B

In vitro CD19⁺Nef⁺ B cells

DIC | Nef | CD19 | SSC | Merged

Ch01 | Ch02 | Ch05 | Ch06 | Ch02/Ch05
13088
10 μm

A

**Nef-G63R
selection**

P = 0.068

B

**Temporal Nef-G63E/R mutant selection
vs NAb induction**

R06-034

■ SIV$_{mac239}$-NAb
▼ Nef G63E
◆ Nef G63R

C

MOI 0.1 Day 3

pAkt Ser473

Gated on singlet p27$^+$

WT
Nef-G63E
Nef-G63R
Uninified

P = 0.70
P < 0.0001

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
gene (*Lentivirus*)	*env*, *nef*	GenBank	M33262.1	SIV_{mac239}
strain, strain background (*Lentivirus*)	SIV_{mac239}	PMID: 2032289		Obtained via COS-1 transfection of pBR_{mac239} and repropagation on MT4-R5 cells
strain, strain background (*Escherichia coli*)	One Shot™ TOP10 Chemically Competent *E. coli*	Invitrogen (Thermo Fisher Scientific)	Cat#:C404003	For subcloning sequencing
cell line (*Homo-sapiens*)	MT4-R5	PMID: 27122584		STR profiling (+)
cell line (*Chlorocebus aethiops*)	COS-1	ATCC	CRL-1650	STR profiling (+)
cell line (*Macaca fascicularis*)	HSC-F cells (CD4+ T-cell line)	PMID: 8610465	Gift from producer (Dr. Hirofumi Akari)	
cell line (*Macaca mulatta*)	HSR5.4 cells (CD4+ T-cell line)	PMID: 8610465	Gift from producer (Dr. Hirofumi Akari)	
cell line (*Homo-sapiens*)	Ramos B cell line	ATCC	CRL-1596	STR profiling (+)
biological sample (*Macaca mulatta*)	Primary simian PBMCs from SIV-infected macaques	Japan SLC	http://www.jslc.co.jp/service/	Cryopreserved from *Macaca mulatta*

biological sample (*Macaca mulatta*)	Primary simian lymph node cells from SIV-infected macaques	Japan SLC	http://www.jslc.co.jp/service/	Cryopreserved from *Macaca mulatta*
antibody	Mouse monoclonal anti-human CD4-FITC (M-T477)	BD Pharmingen	Cat#556615	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD8-PerCP (SK1)	BD Biosciences	Cat#347314	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD3-APC (SP34-2)	BD Pharmingen	Cat#557597	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human IFN-γ-PE (4S.B3)	BioLegend	Cat#502510	FACS (5 µl per test)
antibody	Mouse monoclonal anti-SIV_{mac251} Gag p27 mAb	ABL	Cat#4324	FACS (1:500 with conjugation)
antibody	Mouse monoclonal anti-SIV_{mac251} Nef mAb (clone 17)	Themo Fisher Scientific	Cat#MA1-71522	FACS, PLA, immuno-blotting and imaging cytometry (FACS and imaging cytometry, 1:500 with conjugation; PLA, 1:100; immuno-blotting, 1:1000)
antibody	Mouse IgG1 isotype control mAb (P3.6.2.8.1)	Themo Fisher Scientific	Cat#14-4714-85	PLA (1:100)

antibody	Rabbit monoclonal Alexa Fluor 488-conjugated anti-human phospho-Akt (Ser473) (D9E)	Cell Signaling Technology	Cat#4071	FACS (5 μl per test)
antibody	Rabbit monoclonal Alexa Fluor 488-conjugated anti-human phospho-Akt (Thr308) (C31E5E)	Cell Signaling Technology	Cat#2918	FACS (5 μl per test)
antibody	Rabbit monoclonal PE-conjugated anti-human phospho-Akt (Ser473) (D9E)	Cell Signaling Technology	Cat#5315	FACS and imaging cytometry (1:100)
antibody	Rabbit monoclonal Alexa Fluor 647-conjugated anti-human phospho-Akt (Ser473) (D9E)	Cell Signaling Technology	Cat#4075	FACS (5 μl per test)
antibody	Rabbit monoclonal Alexa Fluor 647-conjugated anti-human phospho-Akt (Thr308) (C31E5E)	Cell Signaling Technology	Cat#3375	FACS (5 μl per test)
antibody	Mouse monoclonal anti-human CD19-PC-5.5 (J3-119)	Beckman Coulter	Cat#66328	FACS and imaging cytometry (5 μl per test)
antibody	Mouse monoclonal anti-human CD4, purified (clone L200)	BD Biosciences	Cat#550625	FACS (25 μg/ml per test for blockade)
antibody	Mouse monoclonal anti-human CD3-	BD Pharmingen	Cat#557757	FACS (5 μl per test)

	APC-Cy7 (SP34-2)			
antibody	Mouse monoclonal anti-human CD8-APC-Cy7 (RPA-T8)	BD Pharmingen	Cat#557760	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD14-APC-Cy7 (M5E2)	BioLegend	Cat#301820	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD16-APC-Cy7 (3G8)	BioLegend	Cat#302018	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD10-APC-Cy7 (HI10a)	BioLegend	Cat#312212	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD11c-PE-Cy7 (3.9)	BioLegend	Cat#301608	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD123-PE-Cy7 (6H6)	BioLegend	Cat#306010	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human HLA-DR-PE-Texas Red (TU36)	Invitrogen (Thermo Fisher Scientific)	Cat# MHLDR17	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD27-PE/Dazzle 594 (M-T271)	BioLegend	Cat#356422	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD10-PE-Cy7 (HI10a)	BioLegend	Cat#312214	FACS (5 µl per test)

antibody	Rabbit polyclonal anti-human IgD-FITC F(ab')$_2$	DaKo	Cat#F0189	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD38-Alexa 647 (AT1)	Santa Cruz Biotechnology	Cat# sc-7325 AF647	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human IgG-Alexa 700 (G18-145)	BD Pharmingen	Cat#561296	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD138-PE (DL-101)	eBioscience (Thermo Fisher Scientific)	Cat#12-1389-42	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human IgG-PE (G18-145)	BD Biosciences	Cat#555787	FACS (5 µl per test)
antibody	Rat monoclonal anti-human/mouse IRF4-eFluor 660 (3E4)	eBioscience (Thermo Fisher Scientific)	Cat#50-9858-80	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human Ki67-Alexa 700 (B56)	BD Pharmingen	Cat#561277	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CD4-PerCP (L200)	BD Pharmingen	Cat#550631	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CD3-Alexa 700 (SP34-2)	BD Pharmingen	Cat#561805	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CD95-PE-Cy7 (DX2)	eBioscience (Thermo Fisher Scientific)	Cat#25-0959-42	FACS (5 µl per test)

antibody	Mouse monoclonal anti-human anti-CXCR5-PE (87.1)	eBioscience (Thermo Fisher Scientific)	Cat#12-1859-42	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human PD-1-Brilliant Violet 421 (EH12.2H7)	BioLegend	Cat#329920	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CXCR3-Alexa 488 (G025H7)	BioLegend	Cat#353710	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human HLA-ABC-FITC (G46-2.6)	BD Biosciences	Cat#557348	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CXCR4-APC (12G5)	BioLegend	Cat#306510	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human BST2-Alexa647 (RS38E)	BioLegend	Cat#348404	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD20 (2H7)	Biolegend	Cat#302324	FACS (5 µl per test)
antibody	Rabbit monoclonal anti-GβL (86B8)	Cell Signaling Technology	Cat#3274	PLA (1:100)
antibody	Rabbit monoclonal anti-mTOR (7C10)	Cell Signaling Technology	Cat#2983	PLA (1:100)
antibody	Rabbit polyclonal anti-human PI3K p85	Merck Millipore	Cat#ABS1856	PLA (1:100)

antibody	Rabbit monoclonal anti-PI3 Kinase p110α (C73F8)	Cell Signaling Technology	Cat#4249	PLA (1:100)
antibody	Goat polyclonal HRP-conjugated anti-monkey IgG (H+L)	Bethyl Laboratory	Cat# A140-102P	ELISA (1:1000)
antibody	Rabbit monoclonal anti-Sin1 (D7G1A)	Cell Signaling Technology	Cat#12860	Co-immuno-precipitation (1:50)
antibody	Mouse TrueBlot™ ULTRA anti-mouse Ig HRP (eB144)	Rockland Immuno-chemicals	Cat# 18-8817-30	Immuno-blotting (1:1000)
antibody	Mouse monoclonal anti-CD3 (FN-18)	Abcam	Cat#ab25109	Stimulation assay (1 µg/ml)
antibody	CD20 MicroBeads, non-human primate	Miltenyi Biotec	Cat#130-091-105	
recombinant DNA reagent	pBR$_{mac239}$	PMID: 2032289	GenBank M33262	coding full sequence of *nef*-open SIV$_{mac239}$
sequence-based reagent	9520GA-F	This paper	PCR primers	GGGACAG AAATACAA TCAGGAA CAGTATAT GAATACTC CATGGAG
sequence-based reagent	9520GA-R	This paper	PCR primers	CTCCATG GAGTATTC ATATACTG TTCCTGAT TGTATTTC TGTCCC
sequence-based reagent	9519GA-F	This paper	PCR primers	GAGGGAC AGAAATAC AATCAGA GACAGTA

				TATGAATA CTCCAT
sequence-based reagent	9519GA-R	This paper	PCR primers	ATGGAGT ATTCATAT ACTGTCTC TGATTGTA TTTCTGTC CCTC
peptide, recombinant protein	Peptide Nef$_{62-70}$	Sigma Aldrich	CTL epitope peptide	QGQYMNT PW
peptide, recombinant protein	Peptide Nef$_{62-70}$.G63E	Sigma Aldrich	Mutant CTL epitope peptide	QEQYMNT PW
peptide, recombinant protein	Peptide Nef$_{62-70}$.G63R	Sigma Aldrich	Mutant CTL epitope peptide	QRQYMNT PW
peptide, recombinant protein	GolgiStop	BD Biosciences	Cat#554724	
peptide, recombinant protein	recombinant SIV$_{mac251}$ Env gp130	ImmunoDx	Cat#5001	Ligand stimulation assay (10 µg/ml)
peptide, recombinant protein	recombinant human IL-2	Roche Diagnostics	Cat#1114752 8001	Ligand stimulation assay (100 IU/ml)
peptide, recombinant protein	recombinant human IFN-γ	Gibco (Thermo Fisher Scientific)	Cat#PHC403 3	Ligand stimulation assay (40 ng/ml)
peptide, recombinant protein	recombinant SIV$_{mac239}$ Env gp140 (biotin-conjugated)	Immune Technology	Cat#IT-001-140p	FACS and imaging cytometry (10µg/ml for probing)

peptide, recombinant protein	streptavidin-Brilliant Violet 421	BioLegend	Cat#405225	FACS (1:100)
peptide, recombinant protein	streptavidin-Alexa Fluor 488	BioLegend	Cat#405235	FACS (1:100)
peptide, recombinant protein	streptavidin-PE-Cy7	BioLegend	Cat#405206	Imaging cytometry (1:100)
peptide, recombinant protein	Recombinant Human IL-4 Protein (carrier-free)	R&D Systems	Cat# 204-IL-050/CF	Functional assay (40 ng/ml)
commercial assay or kit	High Pure Viral RNA kit	Roche Diagnostics	Cat#11858888 2001	
commercial assay or kit	SIV p27 ELISA	ABL	Cat#5436	
commercial assay or kit	SureBlue TMB 1-Component Microwell Peroxidase Substrate	KPL	Cat# 5120-0075	
commercial assay or kit	Prime-Script one-step RT-PCR kit v2	TaKaRa	Cat#RR055A	
commercial assay or kit	KOD-Plus v2	Toyobo	Cat#KOD-211	
commercial assay or kit	Zero Blunt TOPO PCR cloning kit for sequencing	Invitrogen (Thermo Fisher Scientific)	Cat#K287540	
commercial assay or kit	Cytofix/Cytoperm kit	BD Biosciences	Cat#554714	
commercial assay or kit	Monkey Ferritin (Sandwich	LS Bio	Cat#LS-F36490-1	

	ELISA) ELISA Kit			
commercial assay or kit	QuikChange II XL site-directed mutagenesis kit	Agilent Technologies	Cat#200522	
commercial assay or kit	RNeasy Plus Mini kit	Qiagen	Cat#74136	
commercial assay or kit	GeneChip WT Plus reagent	Affymetrix (Thermo Fisher Scientific)	Cat#902280	
commercial assay or kit	GeneChip Human Gene 2.0 ST Array	Affymetrix (Thermo Fisher Scientific)	Cat#902113	
commercial assay or kit	Alexa Fluor™ 488 Antibody Labeling Kit	Invitrogen (Thermo Fisher Scientific)	Cat#A20181	
commercial assay or kit	Alexa Fluor™ 647 Antibody Labeling Kit	Invitrogen (Thermo Fisher Scientific)	Cat# A20186	
commercial assay or kit	Duolink In Situ PLA Probe anti-mouse PLUS	Sigma (Merck)	Cat#DUO920 01	
commercial assay or kit	Duolink In Situ PLA Probe anti-rabbit MINUS	Sigma (Merck)	Cat#DUO920 05	
commercial assay or kit	Duolink flow PLA detection kit-Green	Sigma (Merck)	Cat#DUO940 02	
commercial assay or kit	FuGENE6	Promega	Cat#E2691	
commercial assay or kit	Capturem IP & Co-IP Kit	Clontech/Tak ara	Cat#635721	

software, algorithm	FACS Diva	BD Biosciences	http://www.bdbiosciences.com/jp/instruments/software/facsdiva/	
software, algorithm	FlowJo v10.7.1	BD/Treestar	https://www.flowjo.com/solutions/flowjo	
software, algorithm	Affymetrix Expression Console Software	Affymetrix/Thermo Fisher Scientific	https://www.thermofisher.com/jp/ja/home/life-science/microarray-analysis/microarray-analysis-instruments-software-services/microarray-analysis-software/affymetrix-transcriptome-analysis-console-software.html	
software, algorithm	IDEAS 6.3 (+Machine learning module)	Amnis/Merck Millipore/Luminex	https://www.luminexcorp.com/imaging-flow-cytometry/	
software, algorithm	Image Lab software	BioRad	https://www.bio-rad.com/en-jp/product/image-lab-software	
software, algorithm	Prism 8	GraphPad Software	https://www.graphpad.com	
other	LIVE/DEAD® Fixable Aqua stain	Invitrogen (Thermo Fisher Scientific)	Cat#L34957	FACS and imaging cytometry (1:1000)

other	APC Annexin V	Biolegend	Cat#640920	FACS (5 µl per 100 µl test)
other	Annexin V Binding Buffer	Biolegend	Cat#422201	Diluent
other	LIVE/DEAD® Fixable Near-IR stain	Invitrogen (Thermo Fisher Scientific)	Cat#L10119	FACS (1:1000)
other	phorbol-12-myristate-13-acetate (PMA)	Sigma/Merck	Cat#P1585	Stimulation assay (10 ng/ml)
other	ionomycin	Sigma/Merck	Cat#I3909	Stimulation assay (10 ng/ml)
other	4–15% Mini-PROTEAN® TGX™ Precast Protein Gels	BioRad	Cat#4561085	
other	SuperSignal West Pico	Thermo Fisher Scientific	Cat#34580	