SIV-specific neutralizing antibody induction following selection of a PI3K drive-

attenuated *nef* variant

Hiroyuki Yamamoto[*,1,2,3] and Tetsuro Matano[*,1,3,4]

[1]AIDS Research Center, National Institute of Infectious Diseases, 4-7-1 Gakuen, Musashi-

Murayama City, Tokyo 208-0011, Japan

[2]Department of Biomedicine, University Hospital Basel, Hebelstrasse 20, Basel 4031,

Switzerland

[3]Joint Research Center for Human Retrovirus Infection, Kumamoto University, 2-2-1 Honjo,

Kumamoto 860-0811, Japan

[4]The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku,

Tokyo 108-8639, Japan

*Correspondence:

Hiroyuki Yamamoto, AIDS Research Center, National Institute of Infectious Diseases, 4-7-1

Gakuen, Musashi-Murayama City, Tokyo 208-0011, Japan. Phone: +81-42-848-7154; Fax:

+81-42-848-7155. Email: h-yamato@niid.go.jp

Tetsuro Matano, National Institute of Infectious Diseases, 1-23-1 Toyama, Shinjuku-ku, Tokyo

162-8640, Japan. Phone: +81-3-4582-2811; Fax: +81-3-5285-1165. Email:

tmatano@niid.go.jp

Running title: SIV neutralization post-PI3K diminution

Abstract

HIV and simian immunodeficiency virus (SIV) infections are known for impaired neutralizing antibody (NAb) responses. While sequential virus-host B cell interaction appears to be basally required for NAb induction, driver molecular signatures predisposing to NAb induction still remain largely unknown. Here we describe SIV-specific NAb induction following a virus-host interplay decreasing aberrant viral drive of phosphoinositide 3-kinase (PI3K). Screening of seventy difficult-to-neutralize SIV$_{mac239}$-infected macaques found nine NAb-inducing animals, with seven selecting for a specific CD8$^+$ T-cell escape mutation in viral *nef* before NAb induction. This Nef-G63E mutation reduced excess Nef interaction-mediated drive of B-cell maturation-limiting PI3K/mammalian target of rapamycin complex 2 (mTORC2). *In vivo* imaging cytometry depicted preferential Nef perturbation of cognate Envelope-specific B cells, suggestive of polarized contact-dependent Nef transfer and corroborating cognate B-cell maturation post-mutant selection up to NAb induction. Results collectively exemplify a NAb induction pattern extrinsically reciprocal to human PI3K gain-of-function antibody-dysregulating disease, and indicate that harnessing the PI3K/mTORC2 axis may facilitate NAb induction against difficult-to-neutralize viruses including HIV/SIV.

Keywords

neutralizing antibody; B cells; CD8$^+$ T cells; PI3 kinase; inborn errors of immunity; HIV/SIV

Introduction

Virus-specific neutralizing antibody (NAb) responses by B cells are induced by an intricate cooperation of adaptive immune cells (Kumar et al., 2010; Shulman et al., 2014; Giltin et al., 2014; Wang et al., 2014) and often play a central role in clearance of acute viral infections (Junt et al., 2007). In contrast, persistence-prone viruses such as human immunodeficiency virus type 1 (HIV-1), simian immunodeficiency virus (SIV) and lymphocytic choriomeningitis virus (LCMV) variously equip themselves with B cell/antibody-inhibitory countermeasures (Moir et al., 2001; Mattapallil et al., 2005; Sommerstein et al., 2015; Sammicheli et al., 2016; Fallet et al., 2016; Mason et al., 2016), impairing NAb induction (Levesque et al.,2009; Mikell et al., 2011). These viruses successfully suppress elicitation of potent NAb responses, especially in acute infection (Hunziker et al., 2003; Tomaras et al., 2008), and establish viral persistence, posing considerable challenge for developing protective strategies. In particular, HIV and SIV establish early a large body of infection *in vivo* from early on with a distinct host genome-integrating retroviral life cycle (Whitney et al., 2014). In addition, these lentiviruses are unique in launching a matrix of host immune-perturbing interactions mainly by their six remarkably pleiotropic accessory viral proteins, which optimally fuels viral pathogenesis (Kirchhoff et al., 1995; Sheehy et al., 2002; Harris et al., 2003; Schindler et al., 2006; Neil et al., 2008; Zhang et al., 2009; Laguette et al., 2011; Yamada et al., 2018; Joas et al., 2018; Langer et al., 2019; Yan et al., 2019; Joas et al., 2020; Khan et al., 2020; Volcic et al., 2020; Reuschl et al., 2022). A body of evidence has depicted this in the last decades, its entity, including focal mechanisms of humoral immune perturbation in HIV/SIV infection, remains elusive to date.

Adverse virus-host interactions in HIV/SIV infection leads to a detrimental consequence of the absence of acute-phase endogenous NAb responses. Contrasting this, we and others have previously described in *in vivo* experimental models that passive NAb infusion in the acute phase can trigger an endogenous T-cell synergism resulting in robust control of SIV/SHIV (simian/human immunodeficiency virus) (Haigwood et al., 1996; Yamamoto et al.,

2007; Ng et al., 2010; Iseda et al., 2016; Nishimura et al., 2017). This indicates that virus-

specific NAbs not only confer sterile protection but also can evoke T-cell-mediated non-sterile

viral control, suggesting the importance of endogenous NAb responses supported by humoral-

cellular response synergisms, during an optimal time frame. Therefore, identifying

mechanisms driving NAb induction against such viruses is an important step to eventually

design NAb-based HIV control strategies.

One approach that can provide important insights into this goal is analysis of *in vivo*

immunological events linked with NAb induction against difficult-to-neutralize SIVs in a non-

human primate model. Various *in vivo* signatures of HIV-specific NAb induction, such as

antibody-NAb coevolution (Moore et al., 2012), autoimmune-driven induction (Moody et al.,

2016) and natural killer cell-related host polymorphisms (Bradley et al., 2018) have been

reported to date. The broad range of contributing factors collectively, and interestingly,

indicates that pathways to NAb induction against difficult-to-neutralize viruses including

HIV/SIV are redundant, and may potentially involve as-yet-unknown mechanisms driving NAb

induction. For example, the neutralization resistance of certain SIV strains does not appear to

be explained by any of the aforementioned, posing SIV models as attractive tools to analyze

NAb induction mechanisms.

In the present study, we examined virus-specific antibody responses in rhesus

macaques infected with a highly difficult-to-neutralize SIV strain, SIV_{mac239}. This virus is

pathogenic in rhesus macaques causing simian AIDS across a broad range of geographical

origin of macaques (Cumont et al., 2008). Macaques infected with SIV_{mac239} show persistent

viremia and generally lack NAb responses throughout infection (Kestler et al., 1991; Nomura

et al., 2012). In this study, a large-scale screening of SIV_{mac239}-infected Burmese rhesus

macaques for up to 100 weeks identified a subgroup inducing NAbs in the chronic phase.

Interestingly, before NAb induction, these animals commonly selected for a specific CD8$^+$

cytotoxic T lymphocyte (CTL) escape mutation in the viral Nef-coding gene. Compared with

wild-type (WT) Nef, this mutant Nef manifested a decrease in aberrant interaction with

phosphoinositide 3-kinase (PI3K)/mammalian target of rapamycin complex 2 (mTORC2),

resulting in decreased downstream hyperactivation of the canonical B-cell negative regulator

Akt (Omori et al., 2006; Limon et al., 2014). Machine learning-assisted imaging cytometry

revealed that Nef preferentially targets Env-specific B cells *in vivo*. Furthermore, the NAb

induction was linked with sustained Env-specific B-cell responses after or during the mutant

Nef selection. Thus, NAb induction in SIV_{mac239}-infected hosts conceivably involves a

functional boosting of B cells that is phenotypically reciprocal to a recently found human PI3K

gain-of-function and antibody-dysregulating inborn error of immunity (IEI), activated PI3 kinase

delta syndrome (APDS) (Angulo et al., 2013; Lucas et al., 2014). Our results suggest that

intervening PI3K/mTORC2 signaling can potentially result in harnessing NAb induction against

difficult-to-neutralize viruses.

Results

Identification of macaques inducing SIV$_{mac239}$-neutralizing antibodies

We performed a retrospective antibody profile screening in rhesus macaques infected

with NAb-resistant SIV$_{mac239}$ (n = 70) (**Figure 1A**) and identified a group of animals inducing

anti-SIV$_{mac239}$ NAb responses (n = 9), which were subjected to characterization. These NAb

inducers showed persistent viremia with no significant difference in viral loads compared with

a subgroup of NAb non-inducers (n = 19) that were previously profile-clarified, naïve and major

histocompatibility complex class I (MHC-I) haplotype-balanced (used for comparison

hereafter) (**Figure 1B** and **Figure 1-figure supplement 1**). Plasma SIV$_{mac239}$-NAb titers

measured by 10 TCID$_{50}$ SIV$_{mac239}$ virus-killing assay showed an average maximum titer of 1:16,

being induced at an average of 48 weeks post-infection (p.i.) (**Figure 1C**). Two of the nine

NAb inducers showed detectable NAb responses by 24 weeks p.i., while the remaining seven

induced NAbs after 30 weeks p.i. Anti-SIV$_{mac239}$ neutralizing activity was confirmed in

immunoglobulin G (IgG) purified from plasma of these NAb inducers (**Figure 1-figure**

supplement 2). SIV Env-binding IgGs developed from early infection in both NAb inducers

and non-rapid-progressing NAb non-inducers, the latter differing from rapid progressors

known to manifest serological failure (Hirsch et al., 2004; Nakane et al., 2013) (**Figure 1-**

figure supplement 3A). Titers of Env-binding IgG were not higher but rather lower at year 1

p.i. in the NAb inducers (**Figure 1-figure supplement 3B** and **3C**). This was consistent with

other reports (Havenar-Daughton et al., 2016) and differing with gross B-cell enhancement in

Nef-deleted SIV infection with enhanced anti-SIV binding antibody titers accompanying

marginal neutralization (Adnan et al., 2016). NAb inducers and non-inducers showed similar

patterns of variations in viral *env* sequences (Burns et al., 1993), mainly in variable regions 1,

2 and 4 (V1, V2 and V4) (**Figure 1-figure supplement 4**).

Selection of a viral *nef* mutation (Nef-G63E) precedes chronic-phase SIV$_{mac239}$-specific

NAb induction

To explore viral mutations linked to NAb induction, we assessed viral nonsynonymous

polymorphisms outside *env*. Strikingly, we found selection of a viral genome mutation resulting

in G (glycine)-to-E (glutamic acid) substitution at residue 63 of Nef (Nef-G63E) in seven of the

nine NAb inducers (**Figure 2A**). Two inducers that did not select Nef-G63E were the early

inducers detected for NAb positivity before 24 weeks p.i. This 63rd residue lies in the

unstructured N-terminus of Nef, flanked by two α-helices conserved in SIVmac/SIVsmm

(**Figure 2B**). This region is generally polymorphic among HIV-1/HIV-2/SIV, occasionally being

deleted in laboratory and primary isolate HIV-1 strains (Geyer et al., 2001). AlphaFold2-based

structure prediction did not derive palpable change except for low-probability disruption of the

alpha helices (data not shown). This mutation was found only in two of the nineteen NAb non-

inducers, including one rapid progressor (**Figure 1-figure supplement 1**) and selection was

significantly enriched in the NAb inducers as compared to the nineteen non-inducers (**Figure**

2C). Analysis of Nef-G63E mutation frequencies in plasma viruses showed that Nef-G63E

selection preceded or at least paralleled NAb induction (**Figure 2D**).

Nef-G63E is a CD8$^+$ T-cell escape mutation

To explore the mechanistic link between Nef-G63E selection and NAb induction, first we

assessed whether CD8$^+$ T-cell responses target this region. We found that these seven NAb

inducers selecting this mutation elicited CD8$^+$ T-cell responses specific for a 9-mer peptide

Nef$_{62–70}$ QW9 (QGQYMNTPW) (**Figure 3A**). This Nef$_{62–70}$ QW9 epitope is restricted by MHC-

I molecules Mamu-B*004:01 and Mamu-B*039:01 (Evans et al., 1999; Sette et al., 2012).

Possession of these accounted for six cases of Nef-G63E selection (**Figure 3A**), and the

remaining one animal possessed Mamu-A1*032:02 predicted to bind to Nef$_{63–70}$ GW8

(NetMHCpan). Ten of nineteen NAb non-inducers also had either of these alleles (**Figure 1-**

figure supplement 1). This did not significantly differ with the NAb inducer group (P = 0.25

by Fisher's exact test, data not shown), indicating that NAb induction was not simply linked

with possession of these MHC-I genotypes but instead required furthermore specific selection

of the Nef-G63E mutation (**Figure 2C**). Nef-G63E was confirmed to be an escape mutation

from Nef$_{62-70}$-specific CD8$^+$ T-cell responses (**Figure 3B**). NAb non-inducers possessing these

alleles elicited little or no Nef$_{62-70}$-specific CD8$^+$ T-cell responses (**Figure 3-figure supplement**

1A), suggesting that *in vivo* selection and fixation of this Nef-G63E SIV was indeed Nef$_{62-70}$-

specific CD8$^+$ T cell-dependent. Replication of SIV carrying the Nef-G63E mutation was

comparable with WT on a cynomolgus macaque HSC-F CD4$^+$ T-cell line (Akari et al., 1996),

excluding the possibility that the mutation critically impairs viral replication (**Figure 3-figure**

supplement 1B), and plasma viral loads were comparable between Nef-G63E mutant-

selecting NAb inducers versus non-inducers (**Figure 3-figure supplement 1C**). These results

indicate NAb induction following *in vivo* selection and fixation of the CD8$^+$ T-cell escape *nef*

mutation, Nef-G63E under viral persistence.

G63E mutation reduces aberrant Nef interaction-mediated drive of PI3K/mTORC2

We next focused on the functional phenotype of Nef-G63E mutant SIV in infected cells.

An essence of host perturbation by Nef is its wide-spectrum molecular downregulation,

ultimately facilitating viral replication (Schindler et al., 2004; Zhang et al., 2009). To evaluate

possible amelioration of this property, we compared downregulation of major targets CD3,

CD4, MHC-I, CXCR4 and BST-2 (Jia et al., 2009) in infected HSC-F cells. Nef-G63E mutation

did not confer notable change compared with WT [*P* = not significant (*n.s.*) for all molecules,

data not shown], implicating other non-canonical changes (**Figure 4A**). One report suggested

Nef to drive macrophage production of soluble ferritin and perturb B cells, with its plasma level

correlating with viremia (Swingler et al., 2008). Here, plasma ferritin levels showed no

differences (**Figure 4-figure supplement 1A**) and viral loads were comparable as

aforementioned between Nef-G63E-selecting NAb inducers and non-inducers, arguing

against gross involvement of ferritin as well as other viral replication-related Nef phenotypes

at least in this model.

Next, we elucidated immunosignaling-modulating properties of Nef-G63E SIV. Akt is

known as the predominant immune-intrinsic negative brake of B-cell maturation and

AFC/antibody responses *in vivo* (Omori et al., 2006; Limon et al., 2014; Fruman et al., 1999;

Ray et al., 2015; Sander et al., 2015). Thus, we analyzed Akt phosphorylation on day 3 after

Nef-G63E SIV infection at low multiplicity of infection (MOI). This analysis revealed that its

serine 473-phosphorylated form (pAkt Ser473), non-canonically known for Nef-mediated

upregulation (Kumar et al., 2016), was significantly lower in Nef-G63E mutant-infected cells

compared with WT (**Figure 4B**). The difference observed here was more pronounced than

histogram deviation levels in PI3K gain-of-function mice with full recapitulation of B-

cell/antibody-dysregulating human APDS disease phenotype (Avery et al., 2018). In contrast,

the threonine 308-phosphorylated Akt (pAkt Thr308) level remained unaffected (**Figure 4-**

figure supplement 1B).

Interestingly, external PI3K stimuli by serum starvation (Kennedy et al., 1997)

accelerated phenotype appearance from day 3 to day 1 post-infection (**Figure 4C**, left). A

similar trend was obtained by short-term PI3K stimulation with interferon-γ (Nguyen et al.,

2001), interleukin-2 (Marzec et al., 2008) and SIV Env (François & Klotman, 2003) (**Figure 4-**

figure supplement 1C). A high-MOI SIV infection, comprising higher initial concentration of

extracellular Env stimuli, also accelerated phenotype appearance from day 3 to day 1 post-

infection, with stronger pAkt reduction (**Figure 4C**, right). These data indicate that the Akt-

inhibitory G63E mutant Nef phenotype is PI3K stimuli-dependent. Transcriptome analysis

signatured decrease in PI3K-pAkt-FoxO1-related genes in mutant SIV infection (**Figure 4-**

figure supplement 1D). Given that Akt is also a survival mediator, Nef-G63E mutation may

be Th-cytopathic and decrease Th dysfunctional expansion (Baumjohann et al., 2013), as had

been observed in LCMV strain WE (Recher et al., 2004) and chronic HIV/SIV (Gray et al.,

2011; Lindqvist et al., 2012; Petrovas et al., 2012) infections. Here, reduced chronic-phase

peripheral CXCR3⁻CXCR5⁺PD-1⁺ memory follicular Th (Tfh), linked with antibody cross-

reactivity in one cohort (Locci et al., 2013), was somewhat in line with this notion (**Figure 4-**

figure supplement 1E and **1F**). We additionally examined another CTL escape mutant Nef-

G63R, selected with marginal statistical significance primarily in early stage (**Figure 4-figure**

supplement 2A). This mutant did not associate with NAb induction in one control animal (R06-

034) even upon preferential selection (**Figure 4-figure supplement 2B**) and Nef-G63R

mutant was not similarly decreased in pAkt drive (**Figure 4-figure supplement 2C**),

implicating that the Nef-G63E mutation was more tightly linked with NAb induction in terms of

an immunosignaling phenotype.

We further explored molecular traits of this decreased Akt hyperactivation. We reasoned

that comparing endogenous Nef binding patterns would be adequate and analyzed SIV-

infected HSC-F cells with a flow cytometry-based proximity ligation assay (PLA) (Leuchowius

et al., 2009; Avin et al., 2017). We found that this Nef-G63E mutation causes significant

decrease in Nef binding to PI3K p85/p110α and downstream mTORC2 components mTOR

(Sarbassov et al., 2005) and GβL/mLST8 (Kim et al., 2003) in the CD4^{lo}-SIV Gag p27^{+} infected

population (**Figure 4D**). Results were biochemically confirmed by a decrease in G63E-

mutated Nef binding to the mTORC2-intrinsic cofactor Sin1 in coimmunoprecipitation analysis

of infected HSC-F cells (**Figure 4E**). Collectively, the Nef-G63E mutation attenuates

PI3K/mTORC2 signaling driven by aberrant Nef bridging, explaining decreased Akt Ser473

phosphorylation by mTORC2. These properties were in keeping with more pronounced

apoptosis of CD4-downregulated/infected HSC-F cells upon Nef-G63E SIV infection (**Figure

4F**). Thus, Nef-G63E SIV is a mutant virus decreased in aberrant interaction/drive of B-cell-

inhibitory PI3K/mTORC2 signaling, manifesting a molecular signature reciprocal to human

APDS (Angulo et al., 2013; Lucas et al., 2014; Avery et al., 2018).

Targeting of lymph node Env-specific B cells by Nef *in vivo*

Next, we analyzed *in vivo* targeting of virus-specific B cells by Nef in lymph nodes to

explore the potential B-cell-intrinsic influence of the Nef-G63E phenotype. Previously

suggested influence of soluble Nef itself (Qiao et al., 2006) and/or related host factors

(Swingler et al., 2008) may derive generalized negative influence on B cells. However, SIV

antigen-specific binding antibody responses were rather decreased in NAb inducers (**Figure

1-figure supplement 1**), with comparable viral loads (**Figure 3-figure supplement 1C**) and

ferritin levels (**Figure 4-figure supplement 1A**), differing from the aforementioned literature

stating positive correlation between the three. We surmised that some targeted Nef intrusion

against Env-specific B cells may be occurring and that the decreased aberrant drive of Nef-

PI3K/mTORC2 may result in their enhanced maturation in lymph nodes.

While reports (Xu et al., 2009) histologically proposed Nef B-cell transfer, quantitative

traits, e.g., invasion frequency and influence on virus-specific B cells, have remained unvisited.

One of the reasons is that intracellular Nef staining is dim (particularly for B cell-acquired Nef)

and difficult to examine by conventional flow cytometry. To circumvent this issue, defining

staining cutline using single-cell images potentially overcomes confounding technical hurdles,

such as high Nef false-staining signals owing to pre-permeation rupture and post-permeation

processing that derives biologically discontinuous staining and/or batch-inflated signals. Thus,

we reasoned that at the expense of spatial information, sophisticating imaging cytometry

would best visualize Nef-mediated B-cell perturbation *in vivo*. We analyzed lymph node B cells

with Image Stream X MKII, with high-power (> 10-fold) antigen detection [e.g., molecules of

equivalent soluble fluorochrome (MESF) 5 *vs.* MESF 80 in an average flow cytometer for FITC

detection] ideal for detection/single-cell verification.

We designed a triple noise cancellation strategy to overcome the issues stated above.

Firstly, amine reactive dye staining (Perfetto et al., 2006) gated out B cells being Nef-positive

due to pre-experimental membrane damage (**Figure 5-figure supplement 1A**, left). Next, we

deployed secondary quantitative parameters of Nef signals deriving from each pixel of the

images. Generation of a gray-level cooccurrence matrix (GLCM) (Haralick et al., 1973), which

computes adjacent signal deviation as their frequencies on a transpose matrix, enables

calculation of a variety of feature values summarizing traits of the whole image. A biological

assumption of Nef signal continuity in a true staining suggested that the sum of weighted

square variance of GLCM, i.e., the {sum of [square of (value – average signal strength) ×

frequency of each value occurrence]} would separate natural versus artificial Nef signals. This

value, known as Haralick variance, was computed for multiple directions and averaged. The

resultant Haralick variance mean (**Figure 5-figure supplement 1A**, X axis) is proportionate

with unnaturalness of signals in the Nef channel. Finally, Nef signal intensity threshold (**Figure

5-figure supplement 1A**, Y axis) gated out overtly-stained cells void for true-false verification.

These measures excluded B cells with non-specific, strong-signal binary-clustered staining

pixels for Nef (deriving a large summated variance: X axis, and/or batch-stained for Nef: Y

axis; likely originating from post-experimental membrane damage).

Using this approach, we successfully acquired images of viable Nef+CD19+ B cells with

low Nef signal Haralick variance mean and low Nef signal intensity threshold, with fine-textured

gradation of low-to-intermediate Nef staining with continuity from membrane-proximal regions

and without sporadic staining speckles (**Figure 5A** and **Figure 5-figure supplement 1**).

Scoring segregation of typical void/valid images by a linear discriminant analysis-based

machine learning module showed that this gating provides the highest two-dimensional

separation (**Figure 5-figure supplement 1**). Combined with visualization of Env-binding B

cells, we reproducibly obtained images of Env-specific (Ch 12) CD19+ (Ch 11) B cells without

membrane ruptures (Ch 08) and showing fine-textured transferred Nef (Ch 02) (**Figure 5B**).

Nef invasion upregulated pAkt Ser473 (Ch 03) to a range resembling *in vitro* analysis (**Figure**

5C), demonstrating that Nef-driven aberrant PI3K/mTORC2 signaling does occur in Nef-

invaded B cells *in vivo*. Strikingly, lymph node Env-specific B cells showed significantly higher

Nef-positive frequencies as compared with batch non-Env-specific B cells (**Figure 5D**). This

indicated that infected cell-derived Nef preferentially targets adjacent Env-specific B cells,

putatively through its contact/transfer from infected CD4+ cells to B cells (Xu et al., 2009;

Hashimoto et al., 2016). Thus, the phenotypic change in Nef can directly dictate Env-specific

B-cell maturation.

Cell contact-dependent B-cell invasion from infected cells *in vitro*

To understand cell-intrinsic properties enhancing B-cell Nef acquisition *in vivo*, we

addressed in an *in vitro* coculture reconstitution how infected cell-to-B-cell Nef invasion takes

place and how it is modulated. We performed imaging cytometry on a twelve-hour coculture

of SIV-disseminating HSC-F cells (reaching around 30% Nef-positivity at moment of coculture)

with Ramos B cells, with or without modulators added throughout the coculture period (**Figure**

6A, top left). In this coculture, there is no MHC-related interaction known to enhance T-cell/B-

307 cell contact (Wülfing et al., 1998). Machine learning-verified noise cancellation of fragmented

308 signal acquisition (**Figure 6A**, middle left) filtered reproducible images of Nef-positive HSC-F

309 cells directly adhering to Ramos B cells within the doublet population (**Figure 6A**, top to middle

310 right). Importantly, addition of soluble, stimulation-competent antibodies against macaque

311 CD3 enhanced adhesion of Nef$^+$ T cells to B cells (**Figure 6A**, bottom center). In these

312 doublets, we readily detected images of polarized Nef protrusion/injection from T cells to B

313 cells (second column, filled wedges) as well as trogocytosed B-cell membranes (third column,

314 open wedge), suggesting dynamic Nef acquisition by B cells. Changing centrifugation

315 strengths upon coculture collection and staining (190 x g versus 1,200 x g and 600 x g versus

316 1,200 x g, respectively) shifted the tendency between enriched Nef$^+$ HSC-F-B-cell doublet

317 versus singlet detection via CD3 stimulation, respectively (**Figure 6A**, lower right).

318  Substantiated by the acquired images, an enhancement of live singlet Ramos B-cell Nef

319 acquisition (% Nef$^+$ in total B cells) was also observed by CD3 stimulation in conventional flow

320 cytometry (**Figure 6B**). There was a similar enhancement when we added phytohemagglutinin

321 (PHA) whereas phorbol-12-myristate-13-acetate (PMA)/ionomycin did not affect Nef

322 acquisition by B cells. In a similar 72-hour static coculture with primary CD20$^+$ B cells, we

323 detected recapitulation of the decreased B-cell pAkt Ser473 phosphorylation upon G63E Nef

324 transfer compared to that of WT Nef (**Figure 6C**), indicating that the Nef-G63E mutation can

325 directly alleviate maturation inhibition in preferentially Nef-targeted Env-specific B cells.

327 **Enhanced Env-specific B-cell responses after PI3K-diminuting mutant selection**

328  Finally, to assess *in vivo* B-cell quality in NAb inducers with Nef-G63E mutant selection,

329 we examined peripheral SIV Env-specific IgG$^+$ B-cell responses comprising plasmablasts

330 (PBs) and memory B cells (B$_{mem}$). After excluding lineage-specific cells [T/NK/pro-B

331 cells/monocytes/myeloid dendritic cells (DCs)/plasmacytoid DCs], we defined IgG$^+$ PBs as

332 showing: high replication (Ki-67$^+$), post-activation (HLA-DR$^+$), transcriptional switching for

333 terminal differentiation (IRF4^{hi}) and downregulated antigen surface binding (surface [s]Env^{lo}-

334 cytoplasmic [Cy]Env$^+$) (Nutt et al., 2015; Silveira et al, 2015) (**Figure 7A** and **Figure 7-figure**

supplement 1). NAb inducers showed significantly higher Env-specific IgG$^+$ PB responses

around week 30 p.i., after Nef-G63E selection, compared to those in NAb non-inducers

(**Figures 7B** and **7C**). At 1 year, this difference became enhanced as a result of less

pronounced decrease in Env-specific PB frequencies in the NAb inducers.

Simultaneous quantification of Env-specific PB and IgD$^-$CD27$^+$IgG$^+$ B$_{mem}$ responses

allowed for an assessment of overall B-cell response quality by pair-wise analysis of Env-

specific IgG$^+$ B$_{mem}$/PB as a projection of germinal center (GC) output (Zotos et al., 2010)

(**Figure 7D**, left). In this two-dimensional vector temporally plotting the frequency of B$_{mem}$ and

PBs, vector protrusion towards upper right represents higher gross GC output of antibody-

forming cells (AFCs). In NAb non-inducers, all vectors converged on an empirically defined

polygonal attractor area D_n (gray area surrounded with dotted lines in **Figure 7D**, right) at year

1 p.i. and beyond, describing that these NAb non-inducers failed to sustain Env-specific GC

output. In contrast, the vectors were consistently tracked outside D_n in the NAb inducers with

Nef-G63E (**Figure 7D**). At the time of NAb induction, they converged on another upper-right

GC output attractor area D_i (red area surrounded with dotted lines in **Figure 7D**, right) that is

mutually exclusive with D_n ($P < 0.0001$ by Fisher's exact test on NAb non-inducer/inducer

vector distribution frequency within D_n). These suggest more robust virus-specific IgG$^+$ B-cell

responses following Nef-G63E CD8$^+$ T-cell escape mutant selection, ultimately leading to NAb

induction. The enhanced signature of cognate B-cells was also an inverted pattern of impaired

terminal differentiation of B-cell responses in APDS (Al Qureshah et al., 2021).

Taken together, in the current model, Nef$_{62-70}$-specific CD8$^+$ T-cell responses in

persistently SIV$_{mac239}$-infected macaques selected for an escape mutant, Nef-G63E, which

displays attenuated aberrant Nef binding and ensuing drive of B-cell-inhibitory PI3K/mTORC2.

Nef invasion of B cells *in vivo* occurred more preferentially in Env-specific B cells, suggesting

diminution in Nef-mediated tonic dysregulation of B cells after mutant selection. These events

conceivably predisposed to enhanced Env-specific B-cell responses and subsequent

SIV$_{mac239}$-specific NAb induction, altogether, in a manner reciprocal to human APDS-mediated

immune dysregulation.

Discussion

In the present study, we found that in macaques infected with an NAb-resistant SIV, selection of a CD8[+] T-cell escape *nef* mutant virus, Nef-G63E, precedes NAb induction. As a result, Nef binding-mediated drive of PI3K/mTORC2 in Env-specific B cells becomes attenuated, which in turn unleashes maturation of antiviral B-cell responses *in vivo* to induce NAbs. Importantly, this manifested through a pAkt deviation level more pronounced than WT versus germline *PIK3CD* gain-of-function mutation heterozygote mice recapitulating full human APDS phenotype (Avery et al., 2018). Thus, the current Nef-G63E-associated B-cell/NAb phenotype likely occurs in a "reciprocal APDS-like" manner. This proposition was enhanced based on extension of immune cell-intrinsic Nef influence on cognate B cells (Fig. 5), in addition to infected T cells. This work is, to our knowledge, the first to interlink a PI3K/mTORC2-modulating viral signature and enhanced B-cell/NAb responses in a viral infection model.

A link between viral T-cell escape and consequent immune modulation has been previously explored to some extent. For example, enhanced binding of mutant HIV-1 epitope peptide to inhibitory MHC-I on DCs impairs T cells (Lichterfeld et al., 2007) and decreased CTL-mediated lymph node immunopathology can drive, and not inhibit, the production of LCMV-specific antibodies (Battegay et al., 1993). Our results now evidence a new pattern of NAb responses that are bivalently shaped through viral interactions with both humoral and cellular immunity in AIDS virus infection. SIV$_{mac239}$ infection of macaques possessing MHC-I alleles associated with Nef-G63E mutation can be one unique model to analyze virus-host interaction for B cell maturation leading to NAb induction. The interval between Nef-G63E mutant selection and NAb induction was variable in the NAb inducers, for which we did not obtain a clear explanation. This may be influenced by certain basal competitive balance between humoral versus cellular adaptive immunity, as observed in certain vaccination settings (Querec et al., 2009).

The current Nef-G63E phenotype identified here adds yet another aspect to the wealth

of evidence documenting the multifaceted impact of Nef in HIV/SIV infection (Kirchhoff et al., 1995; Gauduin et al., 2006; Stolp et al., 2012). Based on our data, the current phenotype differed from decreased replication-related properties or generalized amelioration in immune impairment such as those of delta-Nef HIV/SIV (Kirchhoff et al., 1995; Johnson et al., 1997; Gauduin et al., 2006; Fukazawa et al., 2012). The SIV Nef N-terminal unstructured region comprising Nef-G63E is not conserved in HIV-1 (Schindler et al., 2004) and beyond this model study it remains to be addressed whether a mutant HIV-1 with a similar immunosignaling-related phenotype can be obtained and how much lentiviral conservation exists for such interactions. Analysis of cognate B-cell maturation on cohort basis (Hau et al., 2022) may potentially assist this approach.

Preferential Nef transfer to Env-specific B cells (**Figure 5**) suggests potential involvement of polarized perturbation in cognate immune cells *in vivo*, which was best depicted for preferential infectivity of HIV-specific CD4$^+$ T cells (Douek et al., 2002) and HIV condensation to the interface of dendritic cell-T-cell contacts (McDonald et al., 2003). As cognate MHC class II interactions (Wülfing et al., 1998; Giltin et al., 2014) regularly occur between HIV/SIV-specific CD4$^+$ T cells and Env-specific B cells in GCs, dysregulation of Env-specific B cells may well result from hijacking of cognate T-cell/B-cell interactions by the virus. Enhanced T-cell attachment and transfer of Nef to B cells by TCR stimulation (**Figure 6**) may override constraints of infection-mediated CD4$^+$ T-cell cytoskeletal impairment (Campbell et al., 2004; Jolly et al., 2004; Stolp et al., 2009) particularly by antagonizing reduced motility (Stolp et al., 2012), resulting in *in vivo* "filtering" of enhanced cognate CD4$^+$ T-cell interaction with cognate B cells. Interestingly, Tfh cells that bear trogocytosed CD20 and are positive for viral DNA appear *in vivo* during SIV infection and increase when infection progresses (Samer et al., 2023). This is consistent with our data that suggest occurrence of contact-dependent, polarized interactions facilitating Nef transfer to cognate B cells.

The major *in vivo* readout of this study is autologous neutralization of a highly difficult-to-neutralize SIV strain, which is different from HIV-specific broadly neutralizing antibodies (bNAbs). Importantly, however, obtained signatures of enhanced IgG$^+$ Env-specific AFC and

B cells (class switch/terminal differentiation) and virus neutralization (hypermutation) in our

model are indicative of elevated activity of activation-induced cytidine deaminase (AID), the

canonical positive driver of B-cell fate. Thus, we reason that findings in this study have a strong

conceptual continuity with bNAb regulation by AID in HIV infection.

Our results extracted an *in vivo* link between a decrease in aberrant Nef-PI3K/mTORC2

interaction and major enhancement in B-cell responses. The relationship is reciprocal to

immunogenetic mechanisms of human PI3K gain-of-function mutations resulting in APDS with

multiply impaired anti-viral B-cell responses. APDS occurs from germline mutations in

leukocyte-intrinsic p110δ catalytic subunit-coding *PIK3CD* or the ubiquitous p85α regulatory

subunit-coding *PIK3R1*. In both cases, the mutations cause residue substitutions resulting in

class IA PI3K gain-of-function (Dornan et al., 2017). Hyperactivated PI3K interferes with

cognate B cells in an immune cell-intrinsic manner, leading to impaired B-cell class switching

and resultant susceptibility against various infections (Angulo et al., 2013; Lucas et al., 2014;

Coulter et al., 2017). In the current work, we conversely document a decrease in virally

induced tonic PI3K drive, enhanced IgG class-switched cognate B-cell responses and NAb

induction against a notably difficult-to-neutralize SIV. The exact hierarchy of binding between

Nef and PI3K/mTORC2 components (including PI3K isoforms) and its alteration by Nef-G63E

mutation remains to be investigated. Yet PI3Kp85 showed the strongest Nef-binding index in

PLA assay (**Figure 4D**) presumably because of the SH3-binding PxxP domain in Nef which

binds the SH3 domain of PI3Kp85 (Rickles et al., 1994). Decreased binding of Nef to its

canonical target PI3Kp85 may precipitate into attenuated interactions of Nef with quintuple

PI3K/mTORC2, resulting in fragile downstream Akt signaling. Such domino-triggering is also

observed in APDS patients, from gain-of-function mutant p85 to p110 (Dornan et al., 2017),

which similarly illustrates the fine-tuned nature of the PI3K/mTORC2/Akt signaling pathway.

A limitation of this study is the use of retrospective samples, posing constraints for

detecting exact temporal changes in tonic viral B-cell perturbation. Related with this, it was not

attainable to optimally time the sampling of lymph node cells in animals belonging to the

subgroup of interest. Certain plasma and cellular samples were also unavailable in differing

experiments. While partially addressed in *in vitro* reconstitution, much remains to be

investigated for the observed polarity of Env-specific B-cell perturbation *in vivo*. Similarly,

influence of the Nef-G63E phenotype on $CD4^+$ T cells *in vivo* and its consequence on B-cell

modulation remains largely elusive, and in the current work we focused mainly on molecular

properties of the NAb-correlating Nef-G63E mutant strain. HIV/SIV infection uniquely shows a

biphasic Tfh dysregulation of acute-phase destruction (Mattapallil et al., 2005; Moukambi et

al., 2015; Moukambi et al., 2019) followed by dysregulated chronic-phase hyper-expansion

(Lindqvist et al., 2012; Petrovas et al., 2012; Cubas et al., 2013; Xu et al., 2015). Further

studies are therefore warranted to uncover how the Nef mutant, which is proapoptotic and

displays ameliorated PI3K drive, influences Tfh responses at different phases of the infection.

Gain-of-function mutations in PI3K-encoding genes appear to affect B cells more strongly

(Asano et al., 2018). Nonetheless, it is noteworthy that they also evoke basal dysregulation in

peripheral $CD4^+$ T cells, which locally recapitulates dysregulation and death of HIV/SIV-

infected $CD4^+$ T cells, highlighting the importance of PI3K-mediated signaling in HIV/SIV

infection. In any event, we surmise that specific animal models like the current one help to

depict a temporal cascade of *in vivo* events leading to NAb induction, assisting human

immunology. As an extension of this study extracting the NAb-involved molecular axis,

manipulation/reconstitution experiments shall be designed.

In conclusion, we demonstrated in a non-human primate AIDS model that NAb induction

against a difficult-to-neutralize SIV strain occurs after selection of a $CD8^+$ T-cell escape variant

with a reduced ability to drive excess PI3K/mTORC2 signaling. These results collectively offer

an example of how NAb responses against immunodeficiency viruses can be shaped by both

wings of adaptive immune pressure. Given the key role for Nef-mediated perturbation of

PI3K/mTORC2 in NAb resistance, immune cell-intrinsic PI3K/mTORC2 manipulation may

offer a new possibility to harness antiviral NAb responses. As human IEIs are increasingly

becoming discovered with various immune-perturbing phenotypes and inheritance patterns

(Jauch et al., 2023), search of other types of analogies between viral immune dysregulations

and human IEIs may fuel the discovery of novel targets to modulate and harness immune

responses in translational settings.

Acknowledgments

This work was supported by Japanese Initiative for Progress of Research on Infectious Diseases for Global Epidemic (J-PRIDE) (JP19fm0208017 to H.Y.), Japan Program for Infectious Diseases Research and Infrastructure (Interdisciplinary Cutting-edge Research) (JP22wm0325006 to H.Y.), Research Programs on HIV/AIDS, Emerging and Re-emerging Infectious Diseases, and Global Health Issues (JP20fk0410022 and JP24fk0410066 to H.Y.; JP18fk0410003, JP20fk0410011, JP20fk0108125, JP20jm0110012, JP21fk0410035, and JP21jk0210002 to T.M.) from Japan Agency for Medical Research and Development, the Ministry of Education, Culture, Sports, Science, and Technology in Japan ([JSPS] KAKENHI Grants 17H02185, 18K07157 and 24K21287 to H.Y.; 21H02745 to T.M.), the Takeda Science Foundation (to H.Y.), the Imai Memorial Trust for AIDS Research (to H.Y.), and the Mitsui-Sumitomo Insurance Welfare Foundation (to H.Y.). We thank Fumiko Ono, Koji Hanari, Katsuhiko Komatsuzaki, Akio Hiyaoka, Hiromi Ogawa, Keiko Oto, Hirofumi Akari, Yasuhiro Yasutomi, Hiromi Sakawaki, Tomoyuki Miura and Yoshio Koyanagi for animal experiment assistance; Hiroyuki Kogure, Sayuri Seki and Julia R. Hirsiger for imaging cytometry technical assistance; Masako Nishizawa, Trang Thi Thu Hau, Shigeyoshi Harada, Takushi Nomura, Akiko Takeda, Taku Nakane, Nami Iwamoto, Taeko K. Naruse, Akinori Kimura and Mark S. de Souza for their help. H.Y. thanks Shoi Shi, Bettina Stolp, Jens V. Stein, Makoto Yamagishi and Yusuke Yanagi for conceptual discussions, and Jun Abe and Mike Recher for collaborative support.

Author Contributions

H.Y. conceived/designed/performed study and wrote manuscript; T.M. provided animal cohort, co-edited manuscript and co-directed study.

CRediT taxonomy:

Conceptualization: HY

Methodology: HY

Validation: HY

Formal analysis: HY

Investigation: HY

Resources: HY, TM

Data curation: HY

Writing—original draft: HY

Writing—review & editing: HY, TM

Visualization: HY

Supervision: HY, TM

Project administration: HY

Funding acquisition: HY, TM

Declaration of Interests

The authors declare no competing interests.

 Figure Legends

Figure 1. NAb induction against NAb-resistant SIV$_{mac239}$.

(A) Study design.

(B) Plasma viral loads (SIV *gag* RNA copies/ml plasma) in NAb non-inducers (left) and

inducers (right).

(C) Plasma SIV$_{mac239}$ 100% neutralizing end point titers by 10 TCID$_{50}$ killing assay on MT4-R5

cells. Points on the dotted line show marginally NAb-positive results ($< 1:2$). In some animals,

titers were comparable with our reported results using MT4 cells.

Figure 2. Selection of a viral *nef* mutation (Nef-G63E) before NAb induction.

(A) Viral *nef* mutations in NAb inducers. A linear schema indicating Nef functional domains is

aligned above. Mutations at timepoints with NAb$^-$ (indicated by white wedge; mostly before 6

months p.i.), NAb$^{+/-}$ (green wedge) and NAb$^+$) (purple wedge; mostly after 1 year) are shown

in individual animals. Black and dark gray represent dominant and subdominant mutations (or

deletion in R-360) by direct sequencing, respectively. Red and pink indicate dominant and

subdominant G63E detection, respectively.

(B) Schema of SIV$_{mac239}$ Nef structure and Nef-G63E mutation orientation.

(C) Comparison of frequencies of macaques having Nef-G63E in plasma viruses between

NAb non-inducers and inducers. Compared by Fisher's exact test.

(D) Temporal relationship of Nef-G63E frequencies in plasma virus and NAb induction. Black

boxes (left Y axis) show log$_2$ NAb titers; red triangles and blue diamonds (right Y axis) show

percentage of G63E and G63R mutations detected by subcloning (15 clones/point on average),

respectively.

Figure 3. Nef-G63E is a CD8$^+$ T-cell escape mutation.

(A) Nef$_{62-70}$ QW9-specific CD8$^+$ T-cell frequencies and related MHC-I alleles in seven NAb

inducers selecting Nef-G63E. Mamu-B*039:01 and Mamu-B*004:01 are known to restrict

Nef$_{62-70}$ QW9 epitope and binding of Nef$_{63-70}$ peptide to Mamu-A1*032:02 was predicted.

(B) CD8$^+$ T-cell responses specific to wild-type (WT) Nef$_{62-70}$ or mutant Nef$_{62-70}$.G63E or Nef$_{62-70}$.G63R peptides.

$_{70}$.G63R peptides.

Figure 4. Nef-G63E mutation reduces PI3K/mTORC2 binding and pAkt drive.

(A) Representative surface expression level histograms of CD3, CD4, MHC class I, CXCR4

and BST-2 in CD4^{lo}Nef$^+$ subpopulations after WT or Nef-G63E mutant SIV infection at

multiplicity of infection (MOI) 0.1 on HSC-F cells.

(B) Left: representative histograms of relative pAkt serine (Ser) 473 levels in p27$^+$

subpopulations after WT or Nef-G63E mutant SIV infection at MOI 0.1 on HSC-F cells.

Numbers show pAkt Ser473 mean fluorescence intensities (MFIs) for each. Right: Deviation

of pAkt Ser473 MFIs in p27$^+$ HSC-F cells compared with mean MFI of uninfected cells.

Compared by unpaired t test.

(C) Left: Relative pAkt Ser473 levels (normalized to mean MFI of uninfected controls) in Nef$^+$

HSC-F cells assessed for serum starvation (MOI 0.2, 1 day p.i.). Adjusted P values show

results of comparison via Sidak's post-hoc test of 2-way ANOVA (C, left).   Right: Deviation of

pAkt Ser473 MFIs in Nef$^+$ HSC-F cells assessed for high-MOI infection (MOI 5, 1-day p.i.).

Compared by unpaired t test.

(D) Proximity ligation assay (PLA) of Nef binding with mTOR, GβL/mLST8, PI3Kp85 and

PI3Kp110α. MFI-based binding index was calculated as (anti-Nef/anti-partner) - (isotype/anti-

partner) - (anti-Nef/isotype) + (isotype/isotype). Histograms for samples and isotype/isotype

are representatively shown. Differences in MFI binding indexes can be enhanced compared

with comparison of raw MFIs. Compared by paired t tests.

(E) Sin1 co-immunoprecipitation analysis of WT versus Nef-G63E mutant SIV-infected HSC-

F cells (infected at MOI 0.05, day 3). Immunoblotting of Nef (37 kDa) in whole cell lysates

(lanes 4-6) and anti-Sin1 antibody immunoprecipitates (lanes 1-3) are shown.

(F) Cell death frequencies of SIV-infected cells measured by Annexin V positivity (% of CD4^{lo})

(infected at MOI 0.1, day 3). Compared by unpaired t test.

Data represent one of two [(A); (B); (C, right)] independent experiments in quadruplicate, four

independent experiments performed in triplicate [(C, left)], four independent single-well

comparison experiments pooled for statistical analysis (D), two experiments (E) or two

experiments performed with six wells/control (F). Bars: mean $\pm$ SD (B, C right), mean $\pm$ SEM

(C left, F).

Figure 5. Preferential targeting of lymph node Env-specific B cells by Nef *in vivo*.

(A) Representative gating of triple noise-cancellation *in vivo* Nef staining in B cells analyzed

by ImageStreamX MKII. Pre-experimental damaged cells are first excluded with Live/Dead

from focused/centroid/singlet image-acquired CD19$^+$ B cells (first lane right/R4 gated on

"Focused"). Following Nef$^+$ gating (second lane left/R5 gated on R4), a second step of Nef

noise cancellation (second lane middle/R7 on R5) comprises double-negative removal of post-

experimental stochastic irregular staining building a disparate intracellular staining gradient (X

axis, Nef signal pixel Haralick variance mean) and post-experimental batch overt cellular

staining (Y axis, Nef signal pixel intensity threshold). This outputs a B-cell population with a

fine-textured pericellular Nef^{int-lo} staining, biologically concordant with Nef membrane-

anchoring. Probing of anti-Env BCR (αEnv) by recombinant SIV Env (second lane, right) is

combined, resulting in a 2-D panel of intracellular Nef versus αEnv for noise-cancelled Nef$^+$ B

cells (R7) plus Nef$^-$ B cells (R8) (third lane left/"R7+R8"). pAkt Ser473 expression (third lane

right) and cellular morphology (B) was further analyzed. DN, double-negative.

(B) Typical images of Nef-transferred Env-specific B cells defined αEnv$^+$-intracellular Nef$^+$-Nef

Haralick variance mean^{lo}-Nef Intensity threshold^{lo}-Live/Dead$^-$-CD19$^+$ cells ["Nef$^+$aEnv$^+$"

population of lower left panel in (A), gated on "R7, R8"]. Note the pericellular pAkt Ser473

upregulation in these cells (Ch 03/yellow). Data on inguinal lymph node lymphocytes of

macaque R10-007 at week 62 post-SIV$_{mac239}$ infection are shown in (A) and (B).

(C) Comparison of pAkt Ser473 median fluorescence (medFI) intensity levels in Nef$^-$ B cells

(R8) versus noise-cancelled Nef$^+$ B cells (R7). Analyzed by paired t test.

(D) Comparison of Nef-positive cell frequencies in non-Env-specific (left) versus Env-specific

(right) B cells in lymph nodes of persistently SIV-infected macaques (n = 6). Analyzed by paired

t test.

Figure 6. Contact-dependent enhancement of B-cell Nef acquisition in reconstitution.

(A) Imaging flow cytometric analyses of stimuli-dependent enhancement of infected HSC-F

cell adhesion to non-permissive CD19$^+$ Ramos B cells with the indicated gating (left).

Cocultured doublet cells subjected to machine learning-verified filtration of fragmented signal

acquisition (via gating Nef pixel signal contrast^{lo}-Nef mean pixel per object^{hi} cells) deriving

typical images of infected HSC-F cell (Nef stain shown in pale blue pseudocolor) adhesion to

Ramos cells (CD19 stain shown in purple pseudocolor) are shown. Analyzed by unpaired t

test for doublet formation in the unstimulated versus anti-CD3 monoclonal antibody (clone FN-

18)-stimulated group.

(B) Flow cytometric analyses of Nef-acquiring live CD19$^+$ Ramos B cells upon coculture with

SIV-infected HSC-F cells with or without the indicated stimuli. Analyzed by one-way ANOVA

with Tukey's post-hoc multiple comparison tests.

(C) Representative flow cytometric plot (left), histogram (middle) and pAkt Ser473 signal

deviation in WT versus G63E Nef-acquiring CD20$^+$ primary B cells (six wells/control). Analyzed

by unpaired t test. Data represent pooled data of two (A) or three [(B) and (C)] independent

experiments with indicated number of replicates showing similar results.

Figure 7. Enhanced SIV Env-specific B-cell output up to NAb induction following Nef-

G63E selection.

(A) Representative gating (R02-004, week 32 p.i.) of SIV Env gp140-specific memory B cells

(B$_{mem}$) and plasmablasts (PBs). Two panels for PB gating are shown in lower resolution for

visibility. B$_{mem}$ staining was performed separately and gating are merged with PB panels

(second row, panels 2–4). For lineage-specific exclusion (first row, panel 4), CD10 was stained

independently for B$_{mem}$ (see parenthesis).

(B) Changes in SIV Env gp140-specific plasmablast (PB) frequencies in viremic NAb non-

inducers (left, n = 12) and Nef-G63E-selecting NAb inducers (right, n = 6). Available samples

of twelve viremic NAb non-inducers (including ten with $\geq$ 1-year survival) and six NAb inducers

were tracked. In the right panel, the frequencies in NAb non-inducers are shown in background

(gray) for comparison.

(C) Comparison of Env gp140-specific PB frequencies between NAb non-inducers and

inducers by Mann Whitney U tests. In the right, the frequencies at NAb induction were

compared with those at year 1 in NAb non-inducers with $\geq$ 1-year survival (n = 10). Bars: mean

$\pm$ SD.

(D) Left: vector chart of Env gp140-specific memory B cell (B_{mem}) and PB levels. Legends for

each animal correspond to the ones in (B). Right: NAb non-inducer vectors empirically define

a polygonal GC output attractor area D_n (gray area surrounded with dotted lines) on which

they converge by and beyond year 1 p.i. The NAb inducer vectors (shown up to the time of

NAb induction) remained outside of D_n. At the moment of NAb induction they converged on a

second GC output attractor area D_i (red area surrounded with dotted lines), mutually exclusive

with D_n (P < 0.0001 by Fisher's exact test on NAb inducer vector convergence frequency within

D_n). Legends for year 1 p.i. in the NAb non-inducers and moment of NAb induction in the NAb

inducers are specified.

Materials and Methods

Material availability

This study did not generate new unique reagents.

Data and code availability

Data were analyzed using existing computational packages. The accession number for the transcriptome data [WT SIV$_{mac239}$-infected HSC-F cell line (n = 3), Nef-G63E SIV$_{mac239}$-infected HSC-F cell line (n = 3) and uninfected HSC-F cell line (n = 3)] has been deposited under the accession number GenBank: GSE65806.

Rhesus macaques retrospectively utilized for samples

Seventy Burmese rhesus macaques (*macaca mulatta*) (57 males and 13 females) were retrospectively analyzed in this study. Experiments were previously carried out (Matano et al., 2004; Yamamoto et al., 2007; Iseda et al., 2016; Nomura et al., 2012; Ishii et al., 2012; Takahashi et al., 2013; Nakane et al., 2013; Shi et al., 2013; Iwamoto et al., 2014; Terahara et al., 2014) in the Tsukuba Primate Research Center, National Institutes of Biomedical Innovation, Health and Nutrition (NIBIOHN) with the help of the Corporation for Production and Research of Laboratory Primates and the Institute for Frontier Life and Medical Sciences, Kyoto University (IFLMS-KU) after approval by the Committee on the Ethics of Animal Experiments of NIBIOHN and IFLMS-KU under the guidelines for animal experiments at NIBIOHN, IFLMS-KU and National Institute of Infectious Diseases in accordance with the Guidelines for Proper Conduct of Animal Experiments established by Science Council of Japan (http://www.scj.go.jp/ja/info/kohyo/pdf/kohyo-20-k16-2e.pdf). The experiments were in accordance with the "Weatherall report for the use of non-human primates in research" recommendations (https://royalsociety.org/topics-policy/publications/2006/weatherall-report/). Animals were housed in adjoining individual primate cages allowing them to make sight and sound contact with one another for social interactions, where the temperature was kept at

25°C with light for 12 hours per day. Animals were fed with apples and commercial monkey

diet (Type CMK-2, Clea Japan, Inc.). Blood collection and virus challenge were performed

under ketamine anesthesia.

Cells and viruses

MT4-R5 cells were maintained in RPMI1640 (Invitrogen) supplemented with 10% fetal bovine

serum (FBS) (Clontech) and antibiotics. HSC-F cells (cynomolgus CD4$^+$ T-cell line) and

HSR5.4 cells (rhesus macaque CD4$^+$ T-cell line) were maintained in RPMI1640 (Invitrogen)

supplemented with 10% fetal bovine serum (FBS) (Clontech), human interleukin-2 (IL-2) (10

IU/ml, Roche), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (Invitrogen) and

2-mercaptoethanol (Gibco). SIV$_{mac239}$ molecular clone derivative was generated by

mutagenesis PCR (Agilent Technologies) using the primers listed. For virus preparation, COS-

1 cells were transfected with pBR$_{mac239}$ proviral DNA using FuGENE 6 (Promega). At 48 h post-

transfection, culture supernatants were harvested, centrifuged and filtered through a 0.45-μm

pore-size filter (Merck Millipore). To titrate infectivity, prepared viruses were serially diluted and

infected on HSC-F cells in 96-well plates (Falcon) in quadruplicate. At 10 days post-infection,

the endpoint was determined using SIV p27 antigen enzyme-linked immunosorbent assay

(ELISA) kit (ABL), and virus infectivity was calculated as the 50 percent tissue culture infective

dose (TCID$_{50}$) according to the Reed-Muench method.

Identification of SIV$_{mac239}$-NAb inducers

Burmese rhesus macaques previously challenged with the highly pathogenic molecular clone

virus SIV$_{mac239}$ (n = 70) were retrospectively examined approximately up to 2 years for their

plasma NAb profiles. Animals were challenged intravenously with 1,000 TCID$_{50}$ of SIV$_{mac239}$.

In the current study, virological and immunological profiles were compared between the newly

identified NAb inducers (n = 9) and representative NAb non-inducers (n = 19). Sex distribution

of the NAb inducers (eight males and one female) did not significantly differ with the total non-

inducers (49 males and 12 females, P = 0.99 by Fisher's exact test). These NAb non-inducers

and eight NAb inducers except R06-032 were previously partially reported for their plasma

viral loads (Yamamoto et al., 2007; Iseda et al., 2016; Nomura et al., 2012; Takahashi et al.,

2013; Nakane et al., 2013; Iwamoto et al., 2014) (**Figure 1B**). MHC-I haplotypes and alleles

were determined by reference strand-mediated conformation analysis, PCR-SSP (PCR

amplification utilizing sequence-specific priming) and cloning as described (Naruse et al.,

2010). MHC-I binding prediction was made on April 27, 2013 using the IEDB analysis resource

NetMHCpan tool (Hoof et al., 2009). Alleles of interest in the study have been previously

identified in macaques (Nomura et al., 2012; Evans et al., 1999; Sette et al., 2012). Macaques

R06-032, R03-015, R01-010 and R05-010 received a prime-boost vaccination (Matano et al.,

2004) composed of a DNA prime/intranasal Sendai virus vector expressing SIV_{mac239} Gag

(SeV-Gag). R03-015, R06-019, R06-038 and R10-001 received 300 mg of non-specific rhesus

IgG at day 7 post-SIV_{mac239} challenge as an experimental control in our previous reports

(Yamamoto et al., 2007; Iseda et al., 2016; Nakane et al., 2013).

Plasma viral load quantitation

Plasma viral RNA samples were extracted with High Pure Viral RNA kit (Roche Diagnostics).

Serial fivefold sample dilutions were amplified in quadruplicate by reverse transcription and

nested PCR using SIV_{mac239} *gag*-specific primers to determine end point via the Reed-Muench

method as described previously (Matano et al., 2004; Iseda et al., 2016). The lower limit of

detection is approximately 400 viral RNA copies/ml plasma. Viral loads have been previously

partially reported (Yamamoto et al., 2007; Nomura et al., 2012; Takahashi et al., 2013; Nakane

et al., 2013; Iwamoto et al., 2014).

SIV_{mac239}-specific neutralization assay

NAbs were titrated as described (Yamamoto et al., 2007; Iseda et al., 2016). Serial twofold

dilutions of heat-inactivated plasma, or polyclonal IgG affinity-purified with Protein G

Sepharose 4 Fast Flow (GE Healthcare) from heat-inactivated and filtered plasma were mixed

with 10 $TCID_{50}$ of SIV_{mac239} at a 1:1 ratio (5 µl:5 µl) in quadruplicate. After 45-min incubation at

room temperature, the 10 μl mixtures were added into 5 x 10^4 MT4-R5 cells/well in 96-well

plates. Progeny virus production in day 12 culture supernatants was examined by SIV p27

ELISA (ABL) to determine 100% neutralizing endpoint. The lower limit of titration is 1:2.

Neutralization in three out of four wells at a dilution of 1:2 is defined as marginally NAb-positive

(< 1:2). Results were comparable when the same assay was performed with macaque HSC-

F cells (Akari et al., 1996) as targets. NAb inducers R01-012, R02-004, R01-010, R02-007

and NAb non-inducer R01-011 were previously partially reported for their NAb titers measured

with the same method using MT4 cells as targets (Kawada et al., 2007), which derived

comparable results. For assessment of neutralizing activity in IgG, SIV_{mac239}-specific IgGs

purified from pools of plasma with SIV_{mac239}-specific NAb titers were obtained from each animal

as described (Yamamoto et al., 2007). After complement heat-inactivation at 56°C, 30 min and

0.45 μm filtration, IgGs were purified by Protein G Sepharose 4 Fast Flow (GE Healthcare)

and concentrated by Amicon Ultra 4, MW 50,000 (Millipore) to 30 mg/ml and similarly

examined for their 10 $TCID_{50}$ SIV_{mac239} killing titers on MT4-R5 cells.

SIV Env-specific IgG ELISA and immunoblotting

Plasma Env-specific IgG titers were measured as described (Nakane et al., 2013). SIV_{mac251}

Env gp120 (ImmunoDiagnostics) were coated on 96-well assay plates (BD) at 1000 ng/ml

(100 μl/well). Wells were prewashed with phosphate-buffered saline (PBS), blocked with 0.5%

bovine serum albumin (BSA)/PBS overnight and plasma samples were incubated at a 1:20

dilution (5 μl:95 μl) for 2 hours. Wells were washed with PBS and SIV Env-bound antibodies

were detected with a horseradish peroxidase (HRP)-conjugated goat anti-monkey IgG (H+L)

(Bethyl Laboratory) and SureBlue TMB 1-Component Microwell Peroxidase Substrate (KPL).

Absorbance at 450 nm was measured. Samples from week 0 pre-challenge and month 3,

month 6 and year 1 post-challenge were assessed in duplicate. Week 0 average values were

subtracted from corresponding later time point values for calibration. For immunoblotting, SIV

virion-specific IgGs in plasma were detected with a SIV_{mac239}-cross-reactive western blotting

system (ZeptoMetrix). In the NAb non-inducers, samples from those close to rapid progression

(euthanized due to AIDS progression within approximately 1 year) known for low plasma anti-

SIV reactivity (Hirsch et al., 2004; Nakane et al., 2013) were not included.

Sequencing

Sequencing was performed as described (Matano et al., 2004; Iseda et al., 2016). Viral cDNA

fragments spanning from nt (nucleotide) 4829 to nt 7000, nt 6843 to nt 8831 and nt 8677 to nt

10196 in SIV$_{mac239}$ (GenBank accession number MM33262) covering SIV *env* and *nef* were

amplified from plasma viral RNA by nested RT-PCR using Prime-Script one-step RT-PCR kit

v2 (TaKaRa) and KOD-Plus v2 (Toyobo). PCR products were either directly sequenced or

subcloned with a TOP10-transforming TOPO blunt-end cloning system (Invitrogen).

Sequencing was performed using dye terminator chemistry with an ABI 3730 DNA sequencer

(Applied Biosystems). On average, 15 clones were obtained per sample and 20 clones were

assessed when *nef* mutations of interest in early time points were subdominant.

SIV$_{mac239}$-specific CD8$^+$ T-cell responses

Virus-specific CD8$^+$ T-cell frequencies were measured as described (Matano et al., 2004;

Iseda et al., 2016). Peripheral blood mononuclear cells (PBMCs) were cocultured for 6 hours

with autologous herpesvirus papio-immortalized B lymphoblastoid cell lines (B-LCLs) pulsed

with Nef peptides (Sigma-Aldrich Japan) at 1 µM concentration or as indicated otherwise

under GolgiStop (monensin, BD) presence. Intracellular gamma interferon (IFN-γ) staining

was performed using Cytofix/Cytoperm kit (BD) and the following conjugated anti-human

monoclonal antibodies (mAbs): anti-CD4-FITC (M-T477, BD Pharmingen), anti-CD8-PerCP

(SK1, BD Biosciences), anti-CD3-APC (SP34-2, BD Pharmingen) and anti-IFN-γ-PE (4S.B3,

BioLegend). Specific CD8$^+$ T-cell frequencies were determined by subtracting nonspecific IFN-

γ$^+$ CD8$^+$ T-cell frequencies from those after peptide-specific stimulation; frequencies beneath

100 cells/million PBMCs were considered negative. Cells acquired by FACS Canto II (BD)

were analyzed by FACS Diva (BD) and FlowJo (Treestar). Approximately 1 x 10^5 PBMCs were

gated per test.

Nef-mediated signaling perturbation analysis

Virus supernatants obtained from COS-1 cells after transfection with WT or mutant SIV$_{mac239}$

molecular clones were used for infection of CD4$^+$ T-cell lines, cynomolgus macaque-derived

HSC-F and rhesus macaque-derived HSR5.4 (Akari et al., 1996). QuikChange II XL site-

directed mutagenesis kit (Agilent Technologies) was used to construct mutant SIV$_{mac239}$

molecular clones possessing *nef* mutations Nef-G63E (G-to-A mutation at nt 9520) and Nef-

G63R (G-to-A mutation at nt 9519) from the WT SIV$_{mac239}$ molecular clone (Kestler et al., 1991)

(nt number from GenBank accession number M33262). Cells (1 x 10^5 cells/well in U-bottomed

96-well culture plates [BD]) were infected with WT or mutant SIV$_{mac239}$ cultured in RPMI-1640

medium supplemented with 10% fetal bovine serum (FBS) for indicated periods at MOI 0.1

(intracellular signaling analysis) or 0.001 (supernatant analysis). Culture supernatants were

subjected to measurement of SIV capsid p27 concentrations by ELISA. Harvested cells were

fixed and permeabilized with Cytofix/Cytoperm kit, washed twice and subjected to

immunostaining. The following antibodies were used; anti-SIV$_{mac251}$ Gag p27 mAb (ABL)

manually conjugated with Alexa 647 (Life Technologies), anti-SIV$_{mac251}$ Nef mAb (clone 17,

epitope peptide corresponding to SIV$_{mac239}$ Nef 71-80 not including the residue G63: Thermo

Fisher Scientific/Pierce) manually conjugated with Alexa 488 or Alexa 647, anti-human CD3-

APC-Cy7 (SP34-2, BD Pharmingen), anti-human CD4-PerCP (L200, BD Pharmingen), anti-

human HLA-ABC-FITC (G46-2.6, BD Pharmingen), anti-human CXCR4-APC (12G5,

BioLegend), BST2-Alexa 647 (RS38E, BioLegend), Alexa 488-conjugated anti-phospho-Akt

(Ser473) (D9E, CST) and Alexa 488-conjugated anti-phospho-Akt (Thr308) (C31E5E, CST).

Cells acquired by FACS Canto II were analyzed by FACS Diva and FlowJo. Approximately 5

x 10^4 cells were gated for each test.

Ferritin ELISA

Plasma ferritin levels in NAb inducers and control animals were analyzed by monkey ferritin

sandwich ELISA kit (LS Bio) according to the manufacturer's instructions.

PI3K stimulation assay

5×10^4 HSC-F cells were infected with WT or Nef-G63E mutant SIV$_{mac239}$ at MOI 0.2 and

cultured for 1 day in medium supplemented with 10% (normal), 1% (1/10 starvation) or 0.1%

(1/100 starvation) FBS. For ligand stimulation (**Figure 4-figure supplement 1C**), cells at the

end of 1-day culture were pulsed for 20 minutes with 40 ng/ml of recombinant human IFN-γ

(Gibco/Thermo Fisher Scientific), 100 IU/ml of recombinant human IL-2 (Roche Diagnostics)

or 10 μg/ml of SIV$_{mac251}$ Env gp130 (ImmunoDx). Cells were intracellularly stained using

Cytofix/Cytoperm kit with anti-SIV$_{mac251}$ Nef mAb or anti-SIV$_{mac251}$ p27 mAb manually

conjugated to Alexa 647 and PE-conjugated anti-human phospho-Akt (Ser473) (D9E, CST) or

Alexa 488-conjugated anti-human phospho-Akt (Ser473). Cells acquired by FACS Canto II

were analyzed by FACS Diva and FlowJo. Approximately 7×10^4 HSC-F cells were gated per

test.

Transcriptome analysis

Total RNAs were extracted using RNeasy Plus Mini kit (Qiagen) from 2×10^6 HSC-F cells 1

day after infection with WT or Nef-G63E mutant SIV$_{mac239}$ at MOI 5. Negative control RNA

samples were extracted from 2×10^6 uninfected HSC-F cells after culture with the same

condition. Three sets of experiments were performed. Total RNA samples were subjected to a

quality control (QC) analysis using an Agilent 2100 Bioanalyzer. The obtained amounts of total

RNAs were: 12.37 ± 0.39 (uninfected), 8.80 ± 0.44 (WT) and 9.02 ± 0.17 (Nef-G63E) μg ($P =$

0.66 for WT versus Nef-G63E by unpaired t test). In all samples, two bands of 18S and 28S

rRNA were confirmed and the RNA integrity number (RIN) was 10. 500 ng of total RNA

samples were processed with GeneChip WT Plus reagent (Affymetrix/Thermo Fisher

Scientific) to produce 150 μl of fragmented and labeled cDNA samples. These were incubated

with a Human Gene 2.0 ST Array (Affymetrix/Thermo Fisher Scientific) for 16 hours, 60 rpm

at 45°C using GeneChip hybridization oven 645 (Affymetrix/Thermo Fisher Scientific). Results

were scanned with GeneChip Scanner 3000 7G (Affymetrix/Thermo Fisher Scientific) and

processed with Affymetrix Expression Console Software (Affymetrix/Thermo Fisher Scientific)

according to the manufacturer's instructions. Expression values were normalized by the RMA

method. Genes above mean background expression within the cognate sample and showing

significant difference between WT and Nef-G63E SIV (P < 0.05 via unpaired t test for $\log_2$-

transformed values, 768 candidates) were determined. Akt-related genes exhibiting a change

of approximately 10% or more were representatively extracted and manually aligned by the

authors.

Peripheral CD4$^+$ T-cell surface staining

Cryopreserved/thawed PBMCs were stained for 30 minutes at 4°C with the following reagents

or conjugated anti-human mAbs: Live/Dead Aqua (Life Technologies), anti-CD4-PerCP (L200,

BD Pharmingen), anti-CD8-APC-Cy7 (RPA-T8, BD Pharmingen), anti-CD3-Alexa 700 (SP34-

2, BD Pharmingen), anti-CD95-PE-Cy7 (DX2, eBioscience), anti-CXCR5-PE (87.1,

eBioscience), anti-PD-1-Brilliant Violet 421 (EH12.2H7, BioLegend) and anti-CXCR3-Alexa

488 (G025H7, BioLegend). Cells acquired by FACS LSRII Fortessa (BD) were analyzed by

FACS Diva and FlowJo. Approximately 1.5 x 10^5 PBMCs were gated per test.

Proximity ligation assay (PLA)

A flow cytometry-based arrangement of proximity ligation assay (PLA) (Avin et al., 2017) was

performed to quantitatively assess Nef binding to candidate interacting molecules. 1 x 10^5

HSC-F cells were infected with SIV$_{mac239}$ at MOI 0.05 in U-bottomed plates and permeated

with Cytofix/Perm kit (BD Biosciences). After two washes, they were resuspended in 0.5%

BSA/PBS for prevention of experimental procedure-related loss and stained with mouse anti-

SIV$_{mac251}$ Nef mAb (clone 17, Thermo Fisher Scientific/Pierce) or mouse IgG1 isotype control

mAb (P3.6.2.8.1, Abcam) in combination with either of the following rabbit antibodies: anti-

GβL (86B8, CST), anti-mTOR (7C10, CST), polyclonal anti-human PI3K p85 (Merck Millipore),

anti-PI3 Kinase p110α (C73F8, CST) or rabbit IgG1 isotype control mAb (DA1E, CST).

Antibody-stained cells were subsequently probed with Duolink In Situ PLA Probe anti-mouse

PLUS and anti-mouse MINUS probes (Sigma/Merck). Next, they were detected for

intermolecular binding using Duolink flow PLA detection kit (Green) (Sigma/Merck) with a

reaction time of 100 minutes for post-mouse/rabbit probe linking amplification. Finally, these

PLA-subjected cells were additionally stained with anti-SIV$_{mac251}$ Gag p27 mAb manually

conjugated with Alexa 647 and anti-CD4-PerCP for 30 minutes for 20 minutes at 4°C. Cells

acquired by FACS Canto II were analyzed by FACS Diva and FlowJo. Approximately 1 x 10^5

cells were gated per test. Binding index (Y axis) was calculated by deriving the deviation from

the summation of: {baseline P: (MFI of background reaction with mouse isotype control/rabbit

isotype control) + anti-Nef antibody-derived background Q: [(MFI of reaction with mouse anti-

Nef/rabbit isotype control) - (MFI of background reaction with mouse isotype control/rabbit

isotype control)] + anti-binding partner antibody-derived background R:[(MFI of reaction with

mouse isotype control/rabbit anti-binding partner molecule) - (MFI of background reaction with

mouse isotype control/rabbit isotype control)]}.

Co-immunoprecipitation analysis

1 x 10^5 HSC-F cells were infected with SIV$_{mac239}$ at MOI 0.05 in U-bottomed plates in

quadruplicate for each SIV strain for three days and pooled for each for acquiring cell pellets.

These pellets were lysed with Capturem IP & Co-IP Kit lysis buffer (Takara) and portions for

each were immunoprecipitated by anti-Sin1 mAb (D7G1A, CST) (1:50 dilution, 20 minutes,

room temperature). Spin column membrane-bound immunoprecipitates were obtained by

centrifugation with Capturem IP & Co-IP Kit (Clontech/Takara). Whole cell lysates and Sin1-

immunoprecipitated products were subjected to SDS-polyacrylamide gel electrophoresis

separation on a Mini Protean TGX 4-15% gel (BioRad) and transferred to a Immun-Blot PVDF

membrane (BioRad). Immunoblotting was performed by mouse anti-SIV$_{mac251}$ Nef mAb (clone

17, Thermo Fisher Scientific/Pierce) primary antibody probing (1:1000 dilution, 18 hours, 4°

C) and Mouse TrueBlot ULTRA anti-mouse Ig HRP (eB144, Rockland Immunochemicals)

secondary antibody incubation (1:1000 dilution, 30 minutes, room temperature). Nef-specific

bands in infected controls were visualized by enhanced chemiluminescence using

SuperSignal West Pico PLUS (Thermo Fisher Scientific). Bands auto-detectable by Image

Lab software (BioRad) were analyzed for signal intensities. Uncropped images are provided

as supplementary data.

Cell death assay

Infected cell apoptosis was measured by methods previously described (Jauch et al., 2023).

1×10^5 HSC-F cells were infected with SIV_{mac239} at MOI 0.1 in U-bottomed plates (six

wells/control) for each SIV strain for three days. These cells were stained with CD4-FITC (M-

T477, BD Pharmingen) for 15 minutes at room temperature, washed once and next stained

with APC Annexin V (Biolegend) in Annexin V Binding Buffer (Biolegend) for 20 minutes at

room temperature. Stained reactions were then diluted fivefold with Annexin V Binding Buffer

and subjected to analysis. Cells acquired by FACS Lyric were analyzed by FACS Suite and

FlowJo. Approximately 1×10^4 HSC-F cells were gated per test.

Quantitative *in vivo* imaging flow cytometry

Cryopreserved/thawed lymph node cells (LNCs) from several persistently SIV-infected

macaques used in previous experiments (Nakane et al., 2013; Takahashi et al., 2013; Iwamoto

et al., 2014) were seeded in V-bottomed 96-well plates (Nunc) and blocked with 25 μg/ml of

anti-human CD4 (clone L200: BD) in 100 μl volume for 15 minutes at 4°C. After three washes,

they were stained with 10 μg/ml of recombinant SIV_{mac239} Env gp140-biotin (Immune

Technology) for 30 minutes at 4°C. Cells were then stained with anti-CD19-PC5.5 (J3-119,

Beckman Coulter), Live/Dead Aqua (Invitrogen) and streptavidin-APC-Cy7 (BioLegend) for 30

minutes at 4°C. After two washes, cells were processed with Cytofix/Cytoperm kit (BD) for 20

minutes at 4°C, washed twice, and next intracellularly stained with PE-conjugated anti-human

phospho-Akt (Ser473) (D9E, CST) and anti-SIV$_{mac251}$ Nef mAb (clone 17, Thermo Fisher

Scientific/Pierce) manually conjugated with Alexa 488 or mouse IgG1 isotype control mAb

(P3.6.2.8.1, Abcam) for 30 minutes at 4°C. After final two washes, cells were suspended in

0.8% PFA/PBS. All centrifugations for washing (1,200 x g, 2 minutes) were performed at 4°C.

Cells were subjected to image acquisition with Image Stream X MKII imaging flow cytometer

(Amnis/Merck Millipore/Luminex) and analyzed with IDEAS 6.3 software (Amnis/Merck

Millipore/Luminex). Approximately 5 x 10^5 LNCs were acquired for analyses (**Figure 5**). A

custom-implemented linear discriminant analysis-based machine learning module (Luminex)

was utilized for verifying candidate Nef staining signal-related secondary parameters (**Figure

5** and **Figure 5-figure supplement 1A**) for their efficacy of target population separation.

B-cell Nef invasion *in vitro* assay

For Nef invasion assay, 5 x 10^4 HSC-F cells were infected at MOI 1 for 48 hours. These were

then additionally cocultured with 7.5 x 10^4 Ramos B cells for 12 hours with or without addition

of macaque T cell-stimulating anti-CD3 antibody (FN-18, Abcam) at 1 µg/ml. Cells were

double-stained with Alexa 488-conjugated mouse anti-Nef (clone 17: Thermo

Scientific/Pierce) and anti-CD19-PC-5.5 (J3-119, Beckman Coulter). For graphical

visualization (**Figure 6A**), interacting Nef$^+$ infected cell-invaded B cell doublet images acquired

with ImageStream X MKII (Amnis/Merck Millipore) were analyzed by IDEAS 6.3 (Amnis/Merck

Millipore/Luminex) with the defined noise-cancelling gating (via gating Nef pixel signal

contrast^{lo}-Nef mean pixel per object^{hi} cells). For Nef invasion modulation verification on

conventional flow cytometry, the aforementioned culture condition was similarly used. Anti-

CD3 antibody (FN-18, Abcam) addition at 1 µg/ml and phorbol 12-myristate 13-acetate (PMA,

Sigma) plus ionomycin (Sigma) addition (10 ng/ml and 50 ng/ml, respectively) l were evaluated.

Cells were stained with Alexa 647-conjugated mouse anti-Nef (clone 17: Thermo

Scientific/Pierce), anti-CD19-PC-5.5 (J3-119, Beckman Coulter), anti-human phospho-Akt

(Ser473)-Alexa 488 (D9E, CST) and Live/Dead Near-IR (Invitrogen) for dead cell exclusion.

Dead cell frequencies were less than 3%. For viral phenotype evaluation, 1 x 10^5 HSC-F cells

(surface CD40L$^+$) were pre-infected at MOI 0.2 with WT or Nef-G63E mutant SIV$_{mac239}$ six

hours before coculture. 1.5 x 10^5 CD20$^+$ B cells, positively selected from fresh PBMCs of

uninfected rhesus macaques via anti-CD20 microbeads (Miltenyi Biotec), were cocultured with

these HSC-F cells at an E:T ratio of 2:3 for 3 days in the presence of 40 ng/ml carrier-free

recombinant human IL-4 (R&D Systems). Cells were surface-stained with anti-human CD20-

PerCP (2H7, BioLegend) and intracellularly stained using Cytofix/Cytoperm kit with anti-

SIV$_{mac251}$ Nef mAb manually conjugated to Alexa 488 and Alexa 647-conjugated anti-human

phospho-Akt (Ser473) (D9E, CST). For **Figure 6B** and **Figure 6C**, Cells acquired by FACS

Canto II were analyzed by FACS Diva and FlowJo. Approximately 5 x 10^4 CD20$^+$ primary B

cells and 3 x 10^4 CD19$^+$ Ramos B cells were analyzed.

Peripheral SIV Env-specific B-cell responses

Peripheral SIV Env gp140-specific memory B cells (B$_{mem}$) and plasmablasts (PBs) were

measured by flow cytometry with procedures modified from previous reports and our

experience (Silveira et al., 2015; Hau et al., 2022). First, cryopreserved/thawed PBMCs

seeded in V-bottomed 96-well plates (Nunc) were blocked with 25 μg/ml of anti-human CD4

(clone L200: BD) in 100 μl volume for 15 minutes at 4°C. This concentration attains blockade

of promiscuous SIV Env binding to CD4 comparable to levels by magnetic depletion of CD3$^+$

T cells (data not shown). After three washes, they were next stained with 10 μg/ml of

recombinant SIV$_{mac239}$ Env gp140-biotin (Immune Technology) for 30 minutes at 4°C. They

were subsequently stained with the following anti-human antibodies and fluorochromes for 30

minutes at 4°C with combinations as follows; PB/B$_{mem}$: anti-CD3-APC-Cy7 (SP34-2, BD

Pharmingen), anti-CD8-APC-Cy7 (RPA-T8, BD Pharmingen), anti-CD14-APC-Cy7 (M5E2,

BioLegend), anti-CD16-APC-Cy7 (3G8, BioLegend), streptavidin-Brilliant Violet 421

(BioLegend), anti-CD19-PC5.5 (J3-119, Beckman Coulter) and Live/Dead Aqua (Invitrogen);

PB: anti-CD10-APC-Cy7 (HI10a, BioLegend), anti-CD11c-PE-Cy7 (3.9, BioLegend), anti-

CD123-PE-Cy7 (6H6, BioLegend) and anti-HLA-DR-PE-Texas Red (TU36, Invitrogen); B_{mem}:

anti-CD27-PE/Dazzle 594 (M-T271, BioLegend), anti-CD10-PE-Cy7 (HI10a, BioLegend), anti-

IgD-FITC (DaKo), anti-CD38-Alexa 647 (AT1, Santa Cruz Biotechnology), anti-IgG-Alexa 700

(G18-145, BD Pharmingen) and anti-CD138-PE (DL-101, eBioscience). After two washes,

B_{mem} samples were suspended in 0.8% PFA/PBS. For PB staining, surface-stained cells were

further processed with Cytofix/Cytoperm kit (BD) for 20 minutes at 4°C, washed twice, and

next intracellularly stained with 10 μg/ml of recombinant SIV_{mac239} Env gp140-biotin (Immune

Technology) for 30 minutes at 4°C. After three washes, they were next stained with anti-human

IgG-PE (G18-145, BD), anti-human/mouse IRF4-eFluor 660 (3E4, eBioscience), anti-human

Ki67-Alexa 700 (B56, BD Pharmingen) and streptavidin-Alexa 488 (BioLegend) for 30 minutes

at 4°C. After final two washes, cells were suspended in 0.8% PFA/PBS. Cells acquired with

FACS LSRII Fortessa were analyzed with FACS Diva and FlowJo. Approximately 1.5 x 10^5

PBMCs were acquired for B_{mem} and 4 x 10^5 PBMCs were acquired for PB analyses (**Figure**

6). All centrifugations for washing (1,200 x g, 2 minutes) were performed at 4°C. Env-specific

PB frequencies were 0 cells/million PBMCs for pre-challenge samples in all animals examined,

ruling out background confounding for high-sensitivity quantitation of the population.

Statistical Analysis

Analyses were performed via Prism 8 (GraphPad Software). $P < 0.05$ were considered

significant in two-tailed unpaired t tests, paired t tests, Mann Whitney U tests, Fisher's exact

tests, Wilcoxon signed-rank tests, one-way ANOVA with Tukey's post-hoc multiple comparison

tests and two-way ANOVA with Sidak's post-hoc multiple comparison tests. For analysis of

Nef-invaded B-cell pAkt levels, median fluorescence intensities (mFIs) were analyzed due to

relatively small numbers of Nef^+ B cells. Analyses involving pAkt Ser473 levels all derived

comparable results between mFI and MFI for WT versus Nef-G63E virus. Machine learning-

based rating of Nef signal parameters was verified with IDEAS 6.3 Machine Learning Module

(Luminex).

References

1009

1010 Adnan, S., Reeves, R.K., Gillis, J., Wong, F.E., Yu, Y., Camp, J.V., Li, Q., Connole, M., Li, Y.,

1011 Piatak, M. Jr., et al. (2016). Persistent low-level replication of SIVΔnef drives maturation of

1012 antibody and CD8 T cell responses to induce protective immunity against vaginal SIV infection.

1013 *PLoS Pathog*. 12, e1006104.

1014

1015 Akari, H., Mori, K., Terao, K., Otani, I., Fukasawa, M., Mukai, R., and Yoshikawa, Y. (1996). In

1016 vitro immortalization of Old World monkey T lymphocytes with Herpesvirus saimiri: its

1017 susceptibility to infection with simian immunodeficiency viruses. *Virology* 218, 382–388.

1018

1019 Al Qureshah, F., Sagadiev, S., Thouvenel, C.D., Liu, S., Hua, Z., Hou, B., Acharya, M., James,

1020 R.G., and Rawlings, D.J. (2021). Activated PI3Kδ signals compromise plasma cell survival via

1021 limiting autophagy and increasing ER stress. *J. Exp. Med.* 218, e20211035.

1022

1023 Angulo, I., Vadas, O., Garçon, F., Banham-Hall, E., Plagnol, V., Leahy, T.R., Baxendale, H.,

1024 Coulter, T., Curtis, J., Wu, C., et al. (2013). Phosphoinositide 3-kinase δ gene mutation

1025 predisposes to respiratory infection and airway damage. *Science* 342, 866–871.

1026

1027 Asano, T., Okada, S., Tsumura, M., Yeh, T.W., Mitsui-Sekinaka, K., Tsujita, Y., Ichinose, Y.,

1028 Shimada, A., Hashimoto, K., Wada, T., et al. (2018). Enhanced AKT phosphorylation of

1029 circulating B cells in patients with activated PI3Kδ syndrome. *Front. Immunol.* 9, 568.

1030

1031 Avery, D.T., Kane, A., Nguyen, T., Lau, A., Nguyen, A., Lenthall, H., Payne, K., Shi, W., Brigden,

1032 H., French, E., et al. (2018). Germline-activating mutations in PIK3CD compromise B cell

1033 development and function. *J. Exp. Med.* 215, 2073–2095.

1034

1035   Avin, A., Levy, M., Porat, Z., and Abramson, J. (2017) Quantitative analysis of protein-protein

1036   interactions and post-translational modifications in rare immune populations. *Nat. Commun.*

1037   8, 1524.

1038

1039   Battegay, M., Moskophidis, D., Waldner, H., Bründler, M.A., Fung-Leung, W.P., Mak, T.W.,

1040   Hengartner, H., and Zinkernagel, R.M. (1993). Impairment and delay of neutralizing antiviral

1041   antibody responses by virus-specific cytotoxic T cells. *J. Immunol.* 151, 5408–5415.

1042

1043   Baumjohann, D., Preite, S., Reboldi, A., Ronchi, F., Ansel, K.M., Lanzavecchia, A., and

1044   Sallusto, F. (2013). Persistent antigen and germinal center B cells sustain T follicular helper

1045   cell responses and phenotype. *Immunity* 38, 596–605.

1046

1047   Bradley, T., Peppa, D., Pedroza-Pacheco, I., Li, D., Cain, D.W., Henao, R., Venkat, V., Hora,

1048   B., Chen, Y., Vandergrift, N.A., et al. (2018). RAB11FIP5 Expression and Altered Natural Killer

1049   Cell Function Are Associated with Induction of HIV Broadly Neutralizing Antibody Responses.

1050   *Cell* 175, 387–399.e17.

1051

1052   Burns, D.P., Collignon, C., and Desrosiers, R.C. (1993). Simian immunodeficiency virus

1053   mutants resistant to serum neutralization arise during persistent infection of rhesus monkeys.

1054   *J. Virol.* 67, 4104–4113.

1055

1056   Campbell, E.M., Nunez, R., and Hope, T.J. (2004). Disruption of the actin cytoskeleton can

1057   complement the ability of Nef to enhance human immunodeficiency virus type 1 infectivity. *J.*

1058   *Virol.* 78, 5745–5755.

1059

1060   Coulter, T.I., Chandra, A., Bacon, C.M., Babar, J., Curtis, J., Screaton, N., Goodlad, J.R.,

1061   Farmer, G., Steele, C.L., Leahy, T.R., et al. (2017). Clinical spectrum and features of activated

1062 phosphoinositide 3-kinase δ syndrome: A large patient cohort study. *J. Allergy Clin. Immunol.*

1063 139, 597–606.e4.

1064

1065 Cubas, R.A., Mudd, J.C., Savoye, A.L., Perreau, M., van Grevenynghe, J., Metcalf, T., Connick,

1066 E., Meditz, A., Freeman, G.J., Abesada-Terk, G. Jr., et al. (2013). Inadequate T follicular cell

1067 help impairs B cell immunity during HIV infection. *Nat. Med.* 19, 494–499.

1068

1069 Cumont, M.C., Diop, O., Vaslin, B., Elbim, C., Viollet, L., Monceaux, V., Lay, S., Silvestri, G.,

1070 Le Grand, R., Müller-Trutwin, M., et al. (2008). Early divergence in lymphoid tissue apoptosis

1071 between pathogenic and nonpathogenic simian immunodeficiency virus infections of

1072 nonhuman primates. *J. Virol.* 82, 1175–1184.

1073

1074 Dornan, G.L., Siempelkamp, B.D., Jenkins, M.L., Vadas, O., Lucas, C.L., and Burke, J.E.

1075 (2017). Conformational disruption of PI3Kδ regulation by immunodeficiency mutations in

1076 PIK3CD and PIK3R1. *Proc. Natl. Acad. Sci. USA* 114, 1982–1987.

1077

1078 Douek, D.C., Brenchley, J.M., Betts, M.R., Ambrozak, D.R., Hill, B.J., Okamoto, Y., Casazza,

1079 J.P., Kuruppu, J., Kunstman, K., Wolinsky, S., et al. (2002). HIV preferentially infects HIV-

1080 specific CD4[+] T cells. *Nature* 417, 95–98.

1081

1082 Evans, D.T., O'Connor, D.H., Jing, P., Dzuris, J.L., Sidney, J., da Silva, J., Allen, T.M., Horton,

1083 H., Venham, J.E., Rudersdorf, R.A., et al. (1999). Virus-specific cytotoxic T-lymphocyte

1084 responses select for amino-acid variation in simian immunodeficiency virus Env and Nef. *Nat.*

1085 *Med.* 5, 1270–1276.

1086

1087 Fallet, B., Narr, K., Ertuna, Y.I., Remy, M., Sommerstein, R., Cornille, K., Kreutzfeldt, M., Page,

1088 N., Zimmer, G., Geier, F., et al. (2016). Interferon-driven deletion of antiviral B cells at the onset

1089 of chronic infection. *Sci. Immunol.* 1, pii: eaah6817.

1090

1091   François, F., and Klotman, M.E. (2003). Phosphatidylinositol 3-kinase regulates human

1092   immunodeficiency virus type 1 replication following viral entry in primary CD4+ T lymphocytes

1093   and macrophages. *J. Virol.* 77, 2539–2549.

1094

1095   Fruman, D.A., Snapper, S.B., Yballe, C.M., Davidson, L., Yu, J.Y., Alt, F.W., and Cantley, L.C.

1096   (1999). Impaired B cell development and proliferation in absence of phosphoinositide 3-kinase

1097   p85alpha. *Science* 283, 393–397.

1098

1099   Fukazawa, Y., Park, H., Cameron, M.J., Lefebvre, F., Lum, R., Coombes, N., Mahyari, E.,

1100   Hagen, S.I., Bae, J.Y., Reyes, M.D. 3rd., et al. (2012). Lymph node T cell responses predict

1101   the efficacy of live attenuated SIV vaccines. *Nat. Med.* 18, 1673–1681.

1102

1103   Gauduin, M.C., Yu, Y., Barabasz, A., Carville, A., Piatak, M., Lifson, J.D., Desrosiers, R.C.,

1104   and Johnson, R.P. (2006). Induction of a virus-specific effector-memory CD4+ T cell response

1105   by attenuated SIV infection. *J. Exp. Med.* 203, 2661–2672.

1106

1107   Geyer, M., Fackler, O.T., and Peterlin, B.M. (2001). Structure--function relationships in HIV-1

1108   Nef. *EMBO Rep.* 2, 580–585.

1109

1110   Gitlin, A.D., Shulman, Z. and Nussenzweig, M.C. (2014). Clonal selection in the germinal

1111   centre by regulated proliferation and hypermutation. *Nature* 509, 637–640.

1112

1113   Gray, E.S., Madiga, M.C., Hermanus, T., Moore, P.L., Wibmer, C.K., Tumba, N.L., Werner, L.,

1114   Mlisana, K., Sibeko, S., Williamson, C., et al. (2011). The neutralization breadth of HIV-1

1115   develops incrementally over four years and is associated with CD4+ T cell decline and high

1116   viral load during acute infection. *J. Virol.* 85, 4828–4840.

1117

1118 Haigwood, N.L., Watson, A., Sutton, W.F., McClure, J., Lewis, A., Ranchalis, J., Travis, B.,

1119 Voss, G., Letvin, N.L., Hu, S.L., et al. (1996). Passive immune globulin therapy in the

1120 SIV/macaque model: early intervention can alter disease profile. *Immunol. Lett.* 51, 107–114.

1121

1122 Haralick, R.M., Shanmugam, K., and Dinstein, I. (1973) Textural features for image

1123 classification. *IEEE Transactions on Systems, Man, and Cybernetics* SMC-3, 610–621.

1124

1125 Harris, R.S., Bishop, K.N., Sheehy, A.M., Craig, H.M., Petersen-Mahrt, S.K., Watt, I.N.,

1126 Neuberger, M.S., and Malim, M.H. (2003). DNA deamination mediates innate immunity to

1127 retroviral infection. *Cell* 113, 803–809.

1128

1129 Hashimoto, M., Bhuyan, F., Hiyoshi, M., Noyori, O., Nasser, H., Miyazaki, M., Saito, T., Kondoh,

1130 Y., Osada, H., Kimura, S., et al. (2016). Potential role of the formation of tunneling nanotubes

1131 in HIV-1 spread in macrophages. *J. Immunol.* 196, 1832–1841.

1132

1133 Hau, T.T.T., Nishizawa, M., Harada, S., Phan, M.H., Kanno, Y., Nomura, T., Matsuoka, S.,

1134 Kawana-Tachikawa, A., Hall, W.W., Matano, T., et al. (2022). Association of envelope-specific

1135 B-cell differentiation and viral selective pressure signatures in HIV-1 CRF01_AE infection.

1136 *AIDS* 36, 1629–1641.

1137

1138 Havenar-Daughton, C., Carnathan, D.G., Torrents de la Peña, A., Pauthner, M., Briney, B.,

1139 Reiss, S.M., Wood, J.S., Kaushik, K., van Gils, M.J., Rosales, S.L., et al. (2016). Direct Probing

1140 of Germinal Center Responses Reveals Immunological Features and Bottlenecks for

1141 Neutralizing Antibody Responses to HIV Env Trimer. *Cell Rep.* 17, 2195–2209.

1142

1143 Hirsch, V.M., Santra, S., Goldstein, S., Plishka, R., Buckler-White, A., Seth, A., Ourmanov, I.,

1144 Brown, C.R., Engle, R., Montefiori, D., et al. (2004). Immune failure in the absence of profound

1145 CD4+ T-lymphocyte depletion in simian immunodeficiency virus-infected rapid progressor

1146   macaques. *J. Virol.* 78, 275–284.

1147

1148   Hoof, I., Peters, B., Sidney, J., Pedersen, L.E., Sette, A., Lund, O., Buus, S., and Nielsen, M..

1149   (2009). NetMHCpan, a method for MHC class I binding prediction beyond humans.

1150   *Immunogenetics* 61, 1–13.

1151

1152   Hunziker, L., Recher, M., Macpherson, A.J., Ciurea, A., Freigang, S., Hengartner, H., and

1153   Zinkernagel, R.M. (2003). Hypergammaglobulinemia and autoantibody induction mechanisms

1154   in viral infections. *Nat. Immunol.* 4, 343–349.

1155

1156   Iseda, S., Takahashi, N., Poplimont, H., Nomura, T., Seki, S., Nakane, T., Nakamura, M., Shi,

1157   S., Ishii, H., Furukawa, S., et al. (2016). Biphasic CD8+ T-cell defense in simian

1158   immunodeficiency virus control by acute-phase passive neutralizing antibody immunization. *J.*

1159   *Virol.* 90, 6276–6290.

1160

1161   Ishii, H., Kawada, M., Tsukamoto, T., Yamamoto, H., Matsuoka, S., Shiino, T., Takeda, A.,

1162   Inoue, M., Iida, A., Hara, H., et al. (2012). Impact of vaccination on cytotoxic T lymphocyte

1163   immunodominance and cooperation against simian immunodeficiency virus replication in

1164   rhesus macaques. *J. Virol.* 86, 738–745.

1165

1166   Iwamoto, N., Takahashi, N., Seki, S., Nomura, T., Yamamoto, H., Inoue, M., Shu, T., Naruse,

1167   T.K., Kimura, A., and Matano, T. (2014). Control of simian immunodeficiency virus replication

1168   by vaccine-induced Gag- and Vif-specific CD8$^+$ T cells. *J. Virol.* 88, 425–433.

1169

1170   Jauch, A.J., Bignucolo, O., Seki, S., Ghraichy, M., Delmonte, O.M., von Niederhäusern, V.,

1171   Higgins, R., Ghosh, A., Nishizawa, M., Tanaka, M., et al. (2023). Autoimmunity and

1172   immunodeficiency associated with monoallelic LIG4 mutations via haploinsufficiency. *J.*

1173   *Allergy Clin. Immunol.* 152, 500–516.

1174

1175   Jia, B., Serra-Moreno, R., Neidermyer, W., Rahmberg, A., Mackey, J., Fofana, I.B., Johnson,

1176   W.E., Westmoreland, S., and Evans, D.T. (2009). Species-specific activity of SIV Nef and HIV-

1177   1 Vpu in overcoming restriction by tetherin/BST2. *PLoS Pathog.* 5, e1000429.

1178

1179   Joas, S., Parrish, E.H., Gnanadurai, C.W., Lump, E., Stürzel, C.M., Parrish, N.F., Learn, G.H.,

1180   Sauermann, U., Neumann, B., Rensing, K.M., et al. (2018) Species-specific host factors rather

1181   than virus-intrinsic virulence determine primate lentiviral pathogenicity. *Nat. Commun.* 9, 1371.

1182

1183   Joas, S., Sauermann, U., Roshani, B., Klippert, A., Daskalaki, M., Mätz-Rensing, K., Stolte-

1184   Leeb, N., Heigele, A., Tharp, G.K., Gupta, P.M., et al. (2020). Nef-Mediated CD3-TCR

1185   Downmodulation Dampens Acute Inflammation and Promotes SIV Immune Evasion. *Cell Rep.*

1186   30, 2261–2274.e7.

1187

1188   Johnson, R.P., Glickman, R.L., Yang, J.Q., Kaur, A., Dion, J.T., Mulligan, M.J., and Desrosiers,

1189   R.C. (1997). Induction of vigorous cytotoxic T-lymphocyte responses by live attenuated simian

1190   immunodeficiency virus. *J. Virol.* 71, 7711–7718.

1191

1192   Jolly, C., Kashefi, K., Hollinshead, M., and Sattentau, Q.J. (2004). HIV-1 cell to cell transfer

1193   across an Env-induced, actin-dependent synapse. *J. Exp. Med.* 199, 283–293.

1194

1195   Junt, T., Moseman, E.A., Iannacone, M., Massberg, S., Lang, P.A., Boes, M., Fink, K.,

1196   Henrickson, S.E., Shayakhmetov, D.M., Di Paolo, N.C., et al. (2007). Subcapsular sinus

1197   macrophages in lymph nodes clear lymph-borne viruses and present them to antiviral B cells.

1198   *Nature* 450, 110–114.

1199

1200   Kawada, M., Tsukamoto, T., Yamamoto, H., Takeda, A., Igarashi, H., Watkins, D.I., and Matano,

1201   T. (2007). Long-term control of simian immunodeficiency virus replication with central memory

CD4+ T-cell preservation after nonsterile protection by a cytotoxic T-lymphocyte-based vaccine. *J. Virol.* 81, 5202–5211.

Kennedy, S.G., Wagner, A.J., Conzen, S.D., Jordán, J., Bellacosa, A., Tsichlis, P.N., and Hay, N. (1997). The PI 3-kinase/Akt signaling pathway delivers an anti-apoptotic signal. *Genes Dev.* 11, 701–713.

Kestler, H.W. 3rd, Ringler, D.J., Mori, K., Panicali, D.L., Sehgal, P.K., Daniel, M.D., and Desrosiers, R.C. (1991). Importance of the nef gene for maintenance of high virus loads and for development of AIDS. *Cell* 65, 651–662.

Khan, H., Sumner, R.P., Rasaiyaah, J., Tan, C.P., Rodriguez-Plata, M.T., Van Tulleken, C., Fink, D., Zuliani-Alvarez, L., Thorne, L., Stirling, D., et al. (2020). HIV-1 Vpr antagonizes innate immune activation by targeting karyopherin-mediated NF-κB/IRF3 nuclear transport. *Elife* 9, e60821.

Kim, D.H., Sarbassov, D.D., Ali, S.M., Latek, R.R., Guntur, K.V., Erdjument-Bromage, H., Tempst, P., and Sabatini, D.M. (2003). GbetaL, a positive regulator of the rapamycin-sensitive pathway required for the nutrient-sensitive interaction between raptor and mTOR. *Mol. Cell* 11, 895–904.

Kirchhoff, F., Greenough, T.C., Brettler, D.B., Sullivan, L.L. and Desrosiers, R.C. (1995). Absence of intact nef sequences in a long-term, nonprogressing survivor of HIV-1 infection. *N. Engl. J. Med. 332*, 228–232.

Kumar, A., Abbas, W., Colin, L., Khan, K.A., Bouchat, S., Varin, A., Larbi, A., Gatot, J.S., Kabeya, K., Vanhulle, C., et al. (2016). Tuning of AKT-pathway by Nef and its blockade by

1229   protease inhibitors results in limited recovery in latently HIV infected T-cell line. *Sci. Rep.* 6,

1230   24090.

1231

1232   Kumar, V., Scandella, E., Danuser, R., Onder, L., Nitschké, M., Fukui, Y., Halin, C., Ludewig,

1233   B., and Stein, J.V. (2010). Global lymphoid tissue remodeling during a viral infection is

1234   orchestrated by a B cell-lymphotoxin-dependent pathway. *Blood* 115, 4725–4733.

1235

1236   Laguette, N., Sobhian, B., Casartelli, N., Ringeard, M., Chable-Bessia, C., Ségéral, E., Yatim,

1237   A., Emiliani, S., Schwartz, O., and Benkirane, M. (2011). SAMHD1 is the dendritic- and

1238   myeloid-cell-specific HIV-1 restriction factor counteracted by Vpx. *Nature* 474, 654–657.

1239

1240   Langer, S., Hammer, C., Hopfensperger, K., Klein, L., Hotter, D., De Jesus, P.D., Herbert, K.M.,

1241   Pache, L., Smith, N., van der Merwe, J.A., et al. (2019) HIV-1 Vpu is a potent transcriptional

1242   suppressor of NF-κB-elicited antiviral immune responses. *Elife* 8, e41930.

1243

1244   Leuchowius, K.J., Weibrecht, I., Landegren, U., Gedda, L., and Söderberg, O. (2009). Flow

1245   cytometric in situ proximity ligation analyses of protein interactions and post-translational

1246   modification of the epidermal growth factor receptor family. *Cytometry A* 75, 833–839.

1247

1248   Levesque, M.C., Moody, M.A., Hwang, K.K., Marshall, D.J., Whitesides, J.F., Amos, J.D.,

1249   Gurley, T.C., Allgood, S., Haynes, B.B., Vandergrift, N.A., et al. (2009). Polyclonal B cell

1250   differentiation and loss of gastrointestinal tract germinal centers in the earliest stages of HIV-

1251   1 infection. *PLoS Med.* 6, e1000107.

1252

1253   Lichterfeld, M., Kavanagh, D.G., Williams, K.L., Moza, B., Mui, S.K., Miura, T., Sivamurthy, R.,

1254   Allgaier, R., Pereyra, F., Trocha, A., et al. (2007). A viral CTL escape mutation leading to

1255   immunoglobulin-like transcript 4-mediated functional inhibition of myelomonocytic cells. *J. Exp.*

1256   *Med.* 204, 2813–2824.

1257

1258   Limon, J.J., So, L., Jellbauer, S., Chiu, H., Corado, J., Sykes, S.M., Raffatellu, M., and Fruman,

1259   D.A. (2014). mTOR kinase inhibitors promote antibody class switching via mTORC2 inhibition.

1260   *Proc. Natl. Acad. Sci. USA* 111, E5076–5085.

1261

1262   Lindqvist, M., van Lunzen, J., Soghoian, D.Z., Kuhl, B.D., Ranasinghe, S., Kranias, G.,

1263   Flanders, M.D., Cutler, S., Yudanin, N., Muller, M.I., et al. (2012). Expansion of HIV-specific T

1264   follicular helper cells in chronic HIV infection. *J. Clin. Invest.* 122, 3271–3280.

1265

1266   Locci, M., Havenar-Daughton, C., Landais, E., Wu, J., Kroenke, M.A., Arlehamn, C.L., Su, L.F.,

1267   Cubas, R., Davis, M.M., Sette, A., et al. (2013). Human circulating PD-1^{+}CXCR3^{-}CXCR5^{+}

1268   memory Tfh cells are highly functional and correlate with broadly neutralizing HIV antibody

1269   responses. *Immunity* 39, 758–769.

1270

1271   Lucas, C.L., Kuehn, H.S., Zhao, F., Niemela, J.E., Deenick, E.K., Palendira, U., Avery, D.T.,

1272   Moens, L., Cannons, J.L., Biancalana, M., et al. (2014). Dominant-activating germline

1273   mutations in the gene encoding the PI(3)K catalytic subunit p110δ result in T cell senescence

1274   and human immunodeficiency. *Nat. Immunol.* 15*,* 88–97.

1275

1276   Luo, W., Hawse, W., Conter, L., Trivedi, N., Weisel, F., Wikenheiser, D., Cattley, R.T., and

1277   Shlomchik, M.J. (2019). The AKT kinase signaling network is rewired by PTEN to control

1278   proximal BCR signaling in germinal center B cells. *Nat. Immunol.* 20, 736–746.

1279

1280   Marzec, M., Liu, X., Kasprzycka, M., Witkiewicz, A., Raghunath, P.N., El-Salem, M., Robertson,

1281   E., Odum, N., and Wasik, M.A. (2008). IL-2- and IL-15-induced activation of the rapamycin-

1282   sensitive mTORC1 pathway in malignant CD4+ T lymphocytes. *Blood* 111, 2181–2189.

1283

1284   Mason, R.D., Welles, H.C., Adams, C., Chakrabarti, B.K., Gorman, J., Zhou, T., Nguyen, R.,

1285   O'Dell, S., Lusvarghi, S., Bewley, C.A., et al. (2016). Targeted Isolation of Antibodies Directed

1286   against Major Sites of SIV Env Vulnerability. *PLoS Pathog.* 12, e1005537.

1287

1288   Matano, T., Kobayashi, M., Igarashi, H., Takeda, A., Nakamura, H., Kano, M., Sugimoto, C.,

1289   Mori, K., Iida, A., Hirata, T., et al. (2004). Cytotoxic T lymphocyte-based control of simian

1290   immunodeficiency virus replication in a preclinical AIDS vaccine trial. *J. Exp. Med.* 199, 1709–

1291   1718.

1292

1293   Mattapallil, J.J., Douek, D.C., Hill, B., Nishimura, Y., Martin, M., and Roederer, M. (2005).

1294   Massive infection and loss of memory CD4+ T cells in multiple tissues during acute SIV

1295   infection. *Nature* 434, 1093–1097.

1296

1297   McDonald, D., Wu, L., Bohks, S.M., KewalRamani, V.N., Unutmaz, D., and Hope, T.J. (2003).

1298   Recruitment of HIV and its receptors to dendritic cell-T cell junctions. *Science* 300, 1295–1297.

1299

1300   Mikell, I., Sather, D.N., Kalams, S.A., Altfeld, M., Alter, G., and Stamatatos, L. (2011).

1301   Characteristics of the earliest cross-neutralizing antibody response to HIV-1. *PLoS Pathog.* 7,

1302   e1001251.

1303

1304   Moir, S., Malaspina, A., Ogwaro, K.M., Donoghue, E.T., Hallahan, C.W., Ehler, L.A., Liu, S.,

1305   Adelsberger, J., Lapointe, R., Hwu, P., et al. (2001). HIV-1 induces phenotypic and functional

1306   perturbations of B cells in chronically infected individuals. *Proc. Natl. Acad. Sci. USA* 98,

1307   10362–10367.

1308

1309   Moody, M.A., Pedroza-Pacheco, I., Vandergrift, N.A., Chui, C., Lloyd, K.E., Parks, R.,

1310   Soderberg, K.A., Ogbe, A.T., Cohen, M.S., Liao, H.X., Gao, F., et al. (2016). Immune

1311   perturbations in HIV-1-infected individuals who make broadly neutralizing antibodies. *Sci.*

1312   *Immunol.* 1, aag0851.

1313

1314   Moore, P.L., Gray, E.S., Wibmer, C.K., Bhiman, J.N., Nonyane, M., Sheward, D.J., Hermanus,

1315   T., Bajimaya, S., Tumba, N.L., Abrahams, M.R., et al. (2012). Evolution of an HIV glycan-

1316   dependent broadly neutralizing antibody epitope through immune escape. *Nat. Med.* 18,

1317   1688–1692.

1318

1319   Moukambi, F., Rabezanahary, H., Rodrigues, V., Racine, G., Robitaille, L., Krust, B., Andreani,

1320   G., Soundaramourty, C., Silvestre, R., Laforge, M., et al. (2015). Early loss of splenic Tfh cells

1321   in SIV-infected rhesus macaques. *PLoS Pathog.* 11, e1005287.

1322

1323   Moukambi, F., Rabezanahary, H., Fortier, Y., Rodrigues, V., Clain, J., Benmadid-Laktout, G.,

1324   Zghidi-Abouzid, O., Soundaramourty, C., Laforge, M., and Estaquier, J. (2019) Mucosal T

1325   follicular helper cells in SIV-infected rhesus macaques: contributing role of IL-27. *Mucosal*

1326   *Immunol.* 12, 1038–1054.

1327

1328   Nakane, T., Nomura, T., Shi, S., Nakamura, M., Naruse, T.K., Kimura, A., Matano, T., and

1329   Yamamoto, H. (2013). Limited impact of passive non-neutralizing antibody immunization in

1330   acute SIV infection on viremia control in rhesus macaques. *PLoS ONE* 8, e73453.

1331

1332   Naruse, T.K., Chen, Z., Yanagida, R., Yamashita, T., Saito, Y., Mori, K., Akari, H., Yasutomi, Y.,

1333   Miyazawa, M., Matano, T., et al. (2010). Diversity of MHC class I genes in Burmese-origin

1334   rhesus macaques. *Immunogenetics* 62, 601–611.

1335

1336   Neil, S.J., Zang, T., and Bieniasz, P.D. (2008). Tetherin inhibits retrovirus release and is

1337   antagonized by HIV-1 Vpu. *Nature* 451, 425–430.

1338

1339   Ng, C.T., Jaworski, J.P., Jayaraman, P., Sutton, W.F., Delio, P., Kuller, L., Anderson, D.,

1340   Landucci, G., Richardson, B.A., Burton, D.R., et al. (2010). Passive neutralizing antibody

1341   controls SHIV viremia and enhances B cell responses in infant macaques. *Nat. Med.* 16, 1117–

1342   1119.

1343

1344   Nguyen, H., Ramana, C.V., Bayes, J., and Stark, G.R. (2001). Roles of phosphatidylinositol 3-

1345   kinase in interferon-gamma-dependent phosphorylation of STAT1 on serine 727 and activation

1346   of gene expression. *J. Biol. Chem.* 276, 33361–33368.

1347

1348   Nishimura, Y., Gautam, R., Chun, T.W., Sadjadpour, R., Foulds, K.E., Shingai, M., Klein, F.,

1349   Gazumyan, A., Golijanin, J., Donaldson, M., et al. (2017). Early antibody therapy can induce

1350   long-lasting immunity to SHIV. *Nature* 543, 559–563.

1351

1352   Nomura, T., Yamamoto, H., Shiino, T., Takahashi, N., Nakane, T., Iwamoto, N., Ishii, H.,

1353   Tsukamoto, T., Kawada, M., Matsuoka, S., et al. (2012). Association of major histocompatibility

1354   complex class I haplotypes with disease progression after simian immunodeficiency virus

1355   challenge in Burmese rhesus macaques. *J. Virol.* 86, 6481–6490.

1356

1357   Nutt, S.L., Hodgkin, P.D., Tarlinton, D.M., and Corcoran, L.M. (2015). The generation of

1358   antibody-secreting plasma cells. *Nat. Rev. Immunol.* 15, 160–171.

1359

1360   Omori, S.A., Cato, M.H., Anzelon-Mills, A., Puri, K.D., Shapiro-Shelef, M., Calame, K., and

1361   Rickert, R.C. (2006). Regulation of class-switch recombination and plasma cell differentiation

1362   by phosphatidylinositol 3-kinase signaling. *Immunity* 25, 545–557.

1363

1364   Perfetto, S.P., Chattopadhyay, P.K., Lamoreaux, L., Nguyen, R., Ambrozak, D., Koup, R.A.,

1365   and Roederer, M. (2006). Amine reactive dyes: an effective tool to discriminate live and dead

1366   cells in polychromatic flow cytometry. *J. Immunol. Methods* 313, 199–208.

Petrovas, C., Yamamoto, T., Gerner, M.Y., Boswell, K.L., Wloka, K., Smith, E.C., Ambrozak, D.R., Sandler, N.G., Timmer, K.J., Sun, X., et al. (2012). CD4 T follicular helper cell dynamics during SIV infection. *J. Clin. Invest.* 122, 3281–3294.

Qiao, X., He, B., Chiu, A., Knowles, D.M., Chadburn, A., and Cerutti, A. (2006). Human immunodeficiency virus 1 Nef suppresses CD40-dependent immunoglobulin class switching in bystander B cells. *Nat. Immunol.* 7, 302–310.

Querec, T.D., Akondy, R.S., Lee, E.K., Cao, W., Nakaya, H.I., Teuwen, D., Pirani, A., Gernert, K., Deng, J., Marzolf, B., et al. (2009). Systems biology approach predicts immunogenicity of the yellow fever vaccine in humans. *Nat. Immunol.* 10, 116–125.

Ray, J.P., Staron, M.M., Shyer, J.A., Ho, P.C., Marshall, H.D., Gray, S.M., Laidlaw, B.J., Araki, K., Ahmed, R., Kaech, S.M., et al. (2015). The interleukin-2-mTORc1 kinase axis defines the signaling, differentiation, and metabolism of T helper 1 and follicular B helper T cells. *Immunity* 43, 690–702.

Recher, M., Lang, K.S., Hunziker, L., Freigang, S., Eschli, B., Harris, N.L., Navarini, A., Senn, B.M., Fink, K., Lötscher, M., et al. (2004). Deliberate removal of T cell help improves virus-neutralizing antibody production. *Nat. Immunol.* 5, 934–942.

Recher, M., Lang, K.S., Navarini, A., Hunziker, L., Lang, P.A., Fink, K., Freigang, S., Georgiev, P., Hangartner, L., Zellweger, R., et al. (2007). Extralymphatic virus sanctuaries as a consequence of potent T-cell activation. *Nat. Med.* 13, 1316–1323.

Reuschl, A.K., Mesner, D., Shivkumar, M., Whelan, M.V.X., Pallett, L.J., Guerra-Assunção, J.A., Madansein, R., Dullabh, K.J., Sigal, A., Thornhill, J.P., et al. (2022). HIV-1 Vpr drives a

tissue residency-like phenotype during selective infection of resting memory T cells. *Cell Rep.* 39, 110650.

Rickles, R.J., Botfield, M.C., Weng, Z., Taylor, J.A., Green, O.M., Brugge, J.S., and Zoller, M.J. (1994). Identification of Src, Fyn, Lyn, PI3K and Abl SH3 domain ligands using phage display libraries. *EMBO J.* 13, 5598–5604.

Samer, S., Chowdhury, A., Wiche Salinas, T.R., Estrada, P.M.D.R., Reuter, M., Tharp, G., Bosinger, S., Cervasi, B., Auger, J., Gill, K., et al. (2023). Lymph-Node-based CD3[+] CD20[+] cells emerge from membrane exchange between T follicular helper cells and B cells and increase their frequency following simian immunodeficiency virus infection. *J. Virol.* 97, e0176022.

Sammicheli, S., Kuka, M., Di Lucia, P., de Oya, N.J., De Giovanni, M., Fioravanti, J., Cristofani, C., Maganuco, C.G., Fallet, B., Ganzer, L., et al. (2016). Inflammatory monocytes hinder antiviral B cell responses. *Sci. Immunol.* 1, pii: eaah6789.

Sander, S., Chu, V.T., Yasuda, T., Franklin, A., Graf, R., Calado, D.P., Li, S., Imami, K., Selbach, M., Di Virgilio, M., et al. (2015). PI3 kinase and FOXO1 transcription factor activity differentially control B cells in the germinal center light and dark zones. *Immunity* 43, 1075–1086.

Sarbassov, D.D., Guertin, D.A., Ali, S.M., and Sabatini, D.M. (2005) Phosphorylation and regulation of Akt/PKB by the rictor-mTOR complex. *Science* 307, 1098–1101.

Schindler, M., Münch, J., Brenner, M., Stahl-Hennig, C., Skowronski, J., and Kirchhoff, F. (2004). Comprehensive analysis of nef functions selected in simian immunodeficiency virus-infected macaques. *J. Virol.* 78, 10588–10597.

1423 Schindler, M., Münch, J., Kutsch, O., Li, H., Santiago, M.L., Bibollet-Ruche, F., Müller-Trutwin,

1424 M.C., Novembre, F.J., Peeters, M., Courgnaud, V., et al. (2006). Nef-mediated suppression of

1425 T cell activation was lost in a lentiviral lineage that gave rise to HIV-1. *Cell* 125, 1055–1067.

1426

1427 Sette, A., Sidney, J., Southwood, S., Moore, C., Berry, J., Dow, C., Bradley, K., Hoof, I., Lewis,

1428 M.G., Hildebrand, W.H., et al. (2012). A shared MHC supertype motif emerges by convergent

1429 evolution in macaques and mice, but is totally absent in human MHC molecules.

1430 *Immunogenetics* 64, 421–434.

1431

1432 Sheehy, A.M., Gaddis, N.C., Choi, J.D., and Malim, M.H. (2002). Isolation of a human gene

1433 that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* (2002) 418, 646–

1434 650.

1435

1436 Shi, S., Seki, S., Matano, T., and Yamamoto, H. (2013). IL-21-producer CD4$^+$ T cell kinetics

1437 during primary simian immunodeficiency virus infection. *Microbes Infect.* 15, 697–707.

1438

1439 Shulman, Z., Gitlin, A.D., Weinstein, J.S., Lainez, B., Esplugues, E., Flavell, R.A., Craft, J.E.,

1440 and Nussenzweig, M.C. (2014). Dynamic signaling by T follicular helper cells during germinal

1441 center B cell selection. *Science* 345, 1058–1062.

1442

1443 Silveira, E.L., Kasturi, S.P., Kovalenkov, Y., Rasheed, A.U., Yeiser, P., Jinnah, Z.S., Legere,

1444 T.H., Pulendran, B., Villinger, F., and Wrammert, J. (2015). Vaccine-induced plasmablast

1445 responses in rhesus macaques: phenotypic characterization and a source for generating

1446 antigen-specific monoclonal antibodies. *J. Immunol. Methods* 416, 69–83.

1447

1448 Sommerstein, R., Flatz, L., Remy, M.M., Malinge, P., Magistrelli, G., Fischer, N., Sahin, M.,

1449 Bergthaler, A., Igonet, S., Ter Meulen, J., et al. (2015). Arenavirus glycan shield promotes

1450 neutralizing antibody evasion and protracted infection. *PLoS Pathog.* 11, e1005276.

1451

1452 Srinivasan, L., Sasaki, Y., Calado, D.P., Zhang, B., Paik, J.H., DePinho, R.A., Kutok, J.L.,

1453 Kearney, J.F., Otipoby, K.L., and Rajewsky, K. (2009). PI3 kinase signals BCR-dependent

1454 mature B cell survival. *Cell* 139, 573–586.

1455

1456 Stolp, B., Reichman-Fried, M., Abraham, L., Pan, X., Giese, S.I., Hannemann, S., Goulimari,

1457 P., Raz, E., Grosse, R., and Fackler, O.T. (2009). HIV-1 Nef interferes with host cell motility by

1458 deregulation of Cofilin. *Cell Host Microbe* 6, 174–186.

1459

1460 Stolp, B., Imle, A., Coelho, F.M., Hons, M., Gorina, R., Lyck, R., Stein, J.V., and Fackler, O.T.

1461 (2012). HIV-1 Nef interferes with T-lymphocyte circulation through confined environments in

1462 vivo. *Proc. Natl. Acad. Sci. USA* 109, 18541–18546.

1463

1464 Swingler, S., Zhou, J., Swingler, C., Dauphin, A., Greenough, T., Jolicoeur, P., and Stevenson,

1465 M. (2008). Evidence for a pathogenic determinant in HIV-1 Nef involved in B cell dysfunction

1466 in HIV/AIDS. *Cell Host Microbe* 4, 63–76.

1467

1468 Takahashi, N., Nomura, T., Takahara, Y., Yamamoto, H., Shiino, T., Takeda, A., Inoue, M., Iida,

1469 A., Hara,H., Shu, T., et al. (2013). A novel protective MHC-I haplotype not associated with

1470 dominant Gag-specific CD8$^+$ T-cell responses in SIV$_{mac239}$ infection of Burmese rhesus

1471 macaques. *PLoS ONE* 8, e54300.

1472

1473 Terahara, K., Ishii, H., Nomura, T., Takahashi, N., Takeda, A., Shiino, T., Tsunetsugu-Yokota,

1474 Y., and Matano, T. (2014). Vaccine-induced CD107a$^+$ CD4$^+$ T cells are resistant to depletion

1475 following AIDS virus infection. *J. Virol.* 88, 14232–14240.

1476

1477 Tomaras, G.D., Yates, N.L., Liu, P., Qin, L., Fouda, G.G., Chavez, L.L., Decamp, A.C., Parks,

1478 R.J., Ashley, V.C., Lucas, J.T., et al. (2008). Initial B-cell responses to transmitted human

immunodeficiency virus type 1: virion-binding immunoglobulin M (IgM) and IgG antibodies followed by plasma anti-gp41 antibodies with ineffective control of initial viremia. *J. Virol.* 82, 12449–12463.

Volcic, M., Sparrer, K.M.J., Koepke, L., Hotter, D., Sauter, D., Stürzel, C.M., Scherer, M., Stamminger, T., Hofmann, T.G., Arhel, N.J., et al. (2020). Vpu modulates DNA repair to suppress innate sensing and hyper-integration of HIV-1. *Nat. Microbiol.* 5, 1247–1261.

Wang, J., Sohn, H., Sun, G., Milner, J.D. and Pierce, S.K. (2014). The autoinhibitory C-terminal SH2 domain of phospholipase C-γ2 stabilizes B cell receptor signalosome assembly. *Sci. Signal.* 7, ra89.

Whitney, J.B., Hill, A.L., Sanisetty, S., Penaloza-MacMaster, P., Liu, J., Shetty, M., Parenteau, L., Cabral, C., Shields, J., Blackmore, S., et al. (2014). Rapid seeding of the viral reservoir prior to SIV viraemia in rhesus monkeys. *Nature* 512, 74–77.

Wülfing, C., Sjaastad, M.D., and Davis, M.M. (1998). Visualizing the dynamics of T cell activation: intracellular adhesion molecule 1 migrates rapidly to the T cell/B cell interface and acts to sustain calcium levels. *Proc. Natl. Acad. Sci. USA* 95, 6302–6307.

Xu, H., Wang, X., Malam, N., Aye, P.P., Alvarez, X., Lackner, A.A., and Veazey, R.S. (2015). Persistent simian immunodeficiency virus infection drives differentiation, aberrant accumulation, and latent infection of germinal center follicular T helper cells. *J. Virol.* 90,1578–1587.

Xu, W., Santini, P.A., Sullivan, J.S., He, B., Shan, M., Ball, S.C., Dyer, W.B., Ketas, T.J., Chadburn, A., Cohen-Gould, L., et al. (2009). HIV-1 evades virus-specific IgG2 and IgA

responses by targeting systemic and intestinal B cells via long-range intercellular conduits. *Nat. Immunol.* 10, 1008–1017.

Yamada, E., Nakaoka, S., Klein, L., Reith, E., Langer, S., Hopfensperger, K., Iwami, S., Schreiber, G., Kirchhoff, F., Koyanagi, Y., et al. (2018) Human-specific adaptations in Vpu conferring anti-tetherin activity are critical for efficient early HIV-1 replication in vivo. *Cell Host Microbe* 23, 110–120.e7.

Yamamoto, H., Kawada, M., Takeda, A., Igarashi, H. and Matano, T. (2007). Post-infection immunodeficiency virus control by neutralizing antibodies. *PLoS ONE 2*, e540.

Yan, J., Shun, M.C., Zhang, Y., Hao, C., and Skowronski, J. (2019). HIV-1 Vpr counteracts HLTF-mediated restriction of HIV-1 infection in T cells. *Proc. Natl. Acad. Sci. USA.* 116, 9568–9577.

Zhang, F., Wilson, S.J., Landford, W.C., Virgen, B., Gregory, D., Johnson, M.C., Munch, J., Kirchhoff, F., Bieniasz, P.D., and Hatziioannou, T. (2009). Nef proteins from simian immunodeficiency viruses are tetherin antagonists. *Cell Host Microbe 6*, 54–67.

Zotos, D., Coquet, J.M., Zhang, Y., Light, A., D'Costa, K., Kallies, A., Corcoran, L.M., Godfrey, D.I., Toellner, K.M., Smyth, M.J., et al. (2010). IL-21 regulates germinal center B cell differentiation and proliferation through a B cell-intrinsic mechanism. *J. Exp. Med.* 207, 365–378.

  Supplemental Information Legends

  Fig. 1-Supplement 1. Profile of SIV$_{mac239}$-NAb non-inducer subgroup cohort.

  Summary of a SIV$_{mac239}$-NAb non-inducer subgroup cohort (n = 19), composed from previously

  partially characterized, MHC class I haplotype-balanced naïve animals. Symbols A+, B+, E+,

  J+ and D+ represent possession of MHC-I haplotypes *90-120-Ia*, *90-120-Ib*, *90-010-Ie*, *90-*

  088-Ij and *90-010-Id*, respectively. For the titers, - represents negative; +/- represents

  neutralization in three out of four wells in a quadruplicate test. Gray shadings indicate

  euthanasia due to AIDS progression. Mamu-A and Mamu-B alleles of interest in this study are

  listed. These NAb non-inducers were previously partially reported for their plasma viral loads

  and MHC-I alleles (Nomura et al., 2012; Takahashi et al., 2013; Nakane et al., 2013; Iwamoto

  et al., 2014). R01-011 was previously partially reported for NAb titers measured with the same

  method using MT4 cells as targets (Kawada et al., 2007), which derived comparable results.

  Fig. 1-Supplement 2. IgG class-switched NAbs in the inducers.

  Titration of SIV$_{mac239}$-neutralizing IgGs in the NAb inducers. SIV$_{mac239}$-specific IgGs, purified

  from pools of plasma with SIV$_{mac239}$-specific NAb titers, were obtained from each animal as

  described (Yamamoto et al., 2007) and examined for their neutralizing activity by 10 TCID$_{50}$

  SIV$_{mac239}$ killing on MT4-R5 cells. NAb-negative anti-SIV IgG and control IgG were prepared

  from a representative animal (R01-011) with NAb-negative plasma and pooled plasma of

  uninfected rhesus macaques, respectively.

  Fig.1-Supplement 3. Anti-SIV binding IgG profiles in NAb inducers and non-inducers.

  (A) Temporal changes in SIV$_{mac251}$ Env gp120-specific antibody titers in the NAb non-inducers

  without rapid progression [NAb-negative non-rapid progressors, NAb(-) non-RPs] (left, n = 11)

  and NAb inducers [NAb(+)] (right, n = 9). NAb non-inducers showing rapid progression

  (euthanized with AIDS onset within approximately 1 year) were not included except for

  macaque R10-001 (shown in the left for aligned comparison). Samples for non-RPs R06-001

and R10-002 were unavailable.

(B) Unpaired t test of log-transformed anti-SIV Env Ab titers in NAb(-) non-RPs and NAb

inducers. Bars represent geometric mean absorbance within each group.

(C) Lysed SIV_{mac} virion linear antigen-binding at year 1 p.i. in NAb(-) non-RPs (left, n = 12)

and NAb inducers (right, n = 9). Plasma anti-SIV_{mac251} IgGs were detected using a commercial

western blotting system against the parental strain SIV_{mac251} (ZeptoMetrix). White arrowheads

indicate Env gp120-specific bands. Experiments were performed twice with comparable

results.

1566

Fig. 1-Supplement 4. Env sequence variation pattern is not a major characteristic of

NAb induction.

SIV_{mac239} Env sequence variations before and after NAb induction. For each animal, the upper

time point shows pre-induction and the lower time point (in purple) shows post-induction (first

or second time point with NAb titer > 1:2). Dominant residue changes are shown as follows:

gp120 (gray), variable regions V1-V5 (blue), and gp41 (blue gray). Subdominant mutations

are uncolored. Unlabeled gray and black shadings show subdominant and dominant deletion

of residues. Parental SIV_{mac239} sequence is listed for V1-V5. Residues mutated in multiple

animals within V1-V5 are shown in red.

1576

Fig. 3-Supplement 1. Selection of Nef-G63E mutant SIV is linked with Nef_{62-70}-specific

CD8$^+$ T-cell response positivity.

(A) Nef_{62-70}-specific CD8+ T-cell responses around week 20 post-SIV challenge in seven

viremic NAb non-inducers possessing either of the MHC-I alleles Mamu-B*039:01, Mamu-

B*004:01 or Mamu-A1*032:02.

(B) Temporal supernatant SIV p27 concentrations after wild type (WT) or Nef-G63E mutant

SIV infection at multiplicity of infection (MOI) 0.001 on HSC-F cells. Data represent one of two

independent experiments performed in triplicate. Bars: mean ± SEM.

(C) Comparison of plasma viral loads in Nef-G63E-selecting NAb non-inducers (n = 7) versus

non-inducers at indicated time points post-SIV challenge. Bars represent geometric means.

log10-transformed values are compared by unpaired t tests.

1588

Fig. 4-Supplement 1. Decreased Akt hyperactivation properties of Nef-G63E mutant SIV.

(A) Plasma ferritin levels (ng/ml) in examined NAb non-inducers versus Nef-G63E mutant SIV-

selecting NAb inducers around year 1 post-infection. Bars represent mean in each group. Log-

transformed values compared by unpaired t test.

(B) Relative levels of pAkt Ser473 (left panel) and threonine (Thr) 308 (right panel) in SIV Nef$^+$

HSR5.4 cells (MOI 0.1, 4 days p.i.).

(C) Relative pAkt Ser473 levels in Nef$^+$ HSC-F cells without stimuli and transiently pulsed with

IFN-γ, IL-2 or SIV$_{mac}$ Env gp130 after infection (MOI 0.2, 1 day).

(D) Gene expression significantly differing between WT and Nef-G63E SIV infection (MOI 5,

1 day). Black bars show potentially Akt-related genes. Asterisks show dynamics concordant

with the Nef-G63E phenotype (PI3K-pAkt Ser473 downstream downregulation or inhibition

upregulation).

(E) Representative flow cytometric gating (NAb inducer R01-012, wk 30 p.i.) of peripheral

blood CXCR3$^-$CXCR5$^+$PD-1$^+$ memory follicular CD4+ T cells. Cells were gated as CD3$^+$CD8$^-$

CD4$^+$CD95$^+$CXCR5$^+$PD-1$^+$CXCR3$^-$ live singlet PBMCs. Available samples of six NAb inducers

showing peak Nef-G63E mutant selection around week 30 p.i. and thirteen viremic NAb non-

inducers were tracked.

(F) Comparison of peripheral CXCR3$^-$CXCR5$^+$PD-1$^+$ memory Tfh cell frequencies between

pre-SIV infection and around week 30 post-infection in examined NAb non-inducers

developing disease before week 60 (left: $n = 5$) and alive more than 70 weeks (middle: $n = 8$)

and Nef-G63E-detected NAb inducers alive more than 70 weeks (right: $n = 6$). Analyzed by

Wilcoxon signed-rank tests.

MFIs relative to those in uninfected controls (%) are shown. **: $P < 0.01$, ***: $P < 0.001$, n.s.:

not significant between WT and Nef-G63E mutant by unpaired t tests (B-C). Data represent

one of two (B, C) independent experiments performed in triplicate (C: unstimulated) or

quadruplicate (B, C: PI3K-pulsed). Bars: mean ± SEM.

1615

Fig. 4-Supplement 2. Nef-G63R mutant SIV properties.

(A) Comparison of frequencies of macaques selecting for Nef-G63R mutant in plasma

between NAb non-inducers and inducers. Compared by Fisher's exact test.

(B) Temporal relationship of Nef G63R frequencies in plasma virus in a NAb non-inducer

macaque R06-034 preferentially selecting for Nef-G63R but not Nef-G63E mutant. Black

boxes (left Y axis) show $\log_2$ NAb titers; red triangles and blue diamonds (right Y axis) show

percentage of G63E and G63R mutations detected by subcloning (15 clones/point on average),

respectively.

(C) Left: representative histograms of relative pAkt serine (Ser) 473 levels in p27$^+$

subpopulations after WT, Nef-G63E or Nef-G63R mutant SIV infection at MOI 0.1 on HSC-F

cells. Numbers show pAkt Ser473 mean fluorescence intensities (MFIs) for each. Right:

Deviation of pAkt Ser473 MFIs in p27$^+$ HSC-F cells compared with mean MFI of uninfected

cells. Analyzed by one-way ANOVA with Tukey's post-hoc multiple comparison tests. Data

represent one of two independent experiments performed in quadruplicate with similar results.

1630

Fig. 5-Supplement 1. Machine learning-verified morphological gating of Nef-invaded B

cells *in vivo*.

(A) Representative gating strategy plus schematic of inguinal lymph node Nef$^+$CD19$^+$ B cells

of an SIV$_{mac239}$-infected rhesus macaque analyzed by ImageStreamX MK II imaging flow

cytometry (R10-012, week 160 p.i.). First-step incorporation of Live/Dead staining gates out

CD19$^+$ B cells staining positive for Nef potentially due to pre-experimental membrane

permeation (e.g., image #246896, #248952 and #71859). Second-step dual gating for

intracellular Nef signal pixel Haralick variance mean (H Variance mean, X axis), a textural

feature calculated from a gray level co-occurrence matrix (GLCM) for each image representing

adjacent pixel signal strength heterogeneity and Nef signal intensity threshold (Y axis) gates

out CD19$^+$ B cells showing potentially non-specific, strong intracellular binary clustered

staining pixels for Nef deriving a large variance (X axis), and/or cells stained overtly strong for

Nef and hence becoming void for feasible image verification (high threshold) (Y axis), both

likely originating stochastically from post-experimental rupturing procedures including

membrane permeation (e.g. image #18541). This two-step noise cancellation results in

acquisition of a Live/Dead$^-$-Nef signal Haralick variance mean^{lo}-Nef signal intensity threshold^{lo}-

Nef$^+$CD19$^+$ B-cell population (e.g. image #116318, #116250 and #132927) with a pericellular

Nef^{int-lo} staining, biologically concordant with Nef membrane anchoring, resembling Nef$^+$ cells

experimentally generated *in vitro* and showed the highest score in IDEAS6.3 machine learning

module scoring for segregation of typical void vs. valid images (upper right). This gating was

implemented for counting and analysis of Env-specific B-cell Nef invasion *in vivo* (**Figure 5**).

(B) Detection of contact-dependent Nef invasion from SIV-infected HSC-F CD4$^+$ T cells to

cocultured Ramos B cells *in vitro*. One typical image of an intercellular conduit (SSC, purple)

with Nef (green) protrusions from infected HSC-F cells to CD19$^+$ (red) Ramos B cells from two

independent experiments are shown.

1656

1657   **Fig. 7-Supplement 1. Full gating strategy of SIV$_{mac239}$ Env gp140-specific B-cell**

1658   **responses.**

1659   Full gating (R02-004, week 32 p.i.) of SIV Env gp140-specific memory B cells (B$_{mem}$) and

1660   plasmablasts (PBs) shown in **Figure 7**.

A Surface downregulation

CD3, CD4, MHC Class I, CXCR4, BST-2

WT / Nef-G63E / Uninfected

Gated on singlet CD4loNef+

B MOI 0.1 Day 3

pAkt Ser473

1059 / 896 / 746

Gated on singlet p27+

WT / Nef-G63E / Uninfected

P = 0.0009

C Accaleration w/upstream PI3K stimuli Day 3→Day 1

Serum starvation

P = 0.99, P = 0.0017

High MOI

P = 0.0011

D Intracellular PI3K-pathway binding

CD4lop27+ singlet HSC-F

mTORC2: GβL/mLST8, mTOR

PI3 Kinase: PI3Kp85, PI3Kp110α

P = 0.039, P = 0.028, P = 0.022, P = 0.043

E IP: αSin1 / Lysate

WT / Nef-G63E / NC / WT / Nef-G63E / NC / Ladder

IB: αNef

F Cell death

CD4lo singlet HSC-F

20.9 / 29.0

P = 0.0041

WT / Nef-G63E

Figure 6

A

Gated on Lin [CD3/8/(10)14/16]^{lo}-Live singlet PBMCs

B

NAb non-inducers

NAb inducers (wk 30 Nef-G63E+)

— R07-004
— R07-009
— R06-001
— R01-011
— R05-007
— R06-034
— R08-005
— R09-011
— R10-001
— R06-022
— R10-002
— R10-006

— R06-032
— R01-012
— R03-015
— R02-004
— R01-010
— R06-019

SIV Env gp140-specific IgG$^+$ plasmablasts/million PBMCs

time post-SIV challenge

C

wk 30 · at NAb induction vs yr 1

SIV Env gp140-specific IgG$^+$ plasmablasts/million PBMCs

P = 0.0131

P = 0.0030

NAb- · Nef-G63E+ NAb+

NAb- · Nef-G63E+ NAb+

D

Temporal 2-D plot of Env gp140-specific Bmem vs PB

gp140-specific IgG$^+$ plasmablasts/million PBMCs

gp140-specific IgG$^+$ memory B cells/million PBMCs

D_i = GC output attractor 2 (Nef-G63E+, at NAb induction)

D_n = GC output attractor 1 (NAb-, at yr 1)

MHC-I/ID	SIV$_{mac239}$ neutralization at 1:2 plasma dilution by 10TCID$_{50}$ killing assay (wks p.i.)									pVL at 6 mo	Nef G63E at 1 yr	Detected *Mamu* alleles
	wk11–20	wk21–30	wk31–40	wk41–50	wk51–60	wk61–70	wk71–80	wk81–90	wk91–104			
A+												
R06-037		wk25 -		wk46 +/-			wk72 -	wk87 -	wk95 -	6.4 x 10^2	- (pVL < 400)	
R07-001		wk25 -		wk42 -				wk88 +/-	wk96 -	< 400	- (pVL < 400)	*A1*032:02*
R07-004		wk25 -		wk46 -		wk64 -	wk74 -	wk86 -		4.5 x 10^3	+	*B*039:01*
R07-009		wk25 -		wk46 -		wk64 -	wk74 -			4.5 x 10^3	-	
B+												
R06-001		wk25 -		wk48 -	wk60 -		wk80 -	wk90 -		8.2 x 10^3	-	
R06-039		wk25 -		wk46 -	wk56 -					8.5 x 10^6	-	*A1*032:02 , B*004:01*
E+												
R01-011		wk22 -	wk40 -		wk52 -		wk71 -	wk85 -	wk104 -	1.3 x 10^5	-	*B*004:01*
R05-007		wk26 -			wk55 +/-		wk71 -	wk89 -	wk97 -	1.6 x 10^4	-	
R08-007		wk21 -		wk46 -		wk69 -		wk82 +/-		8.5 x 10^6	-	
R09-011	wk18 -		wk36 -		wk52 -					8.5 x 10^6	-	
R06-038		wk25 -			wk58 +/-		wk80 +/-		wk95 -	1.1 x 10^5	-	
J+												
R04-014	wk12 -	wk25 -	wk40 -							1.5 x 10^5	+ (wk40)	*B*039:01*
R06-022	wk12 -	wk23 -								1.4 x 10^7	- (wk23)	*B*039:01*
R10-001	wk14 -	wk30 -	wk37 -							1.9 x 10^7	- (wk37)	*B*039:01*
D+												
R05-006		wk25 -		wk48 -			wk80 -	wk90 -	wk98 -	5.4 x 10^2	-	*A1*032:02 , B*004:01*
R06-034		wk25 -		wk42 -		wk61 -	wk80 +/-	wk88 +/-	wk96 -	4.5 x 10^3	-	*A1*032:02 , B*004:01*
R08-005		wk25 -		wk48 -		wk68 -			wk100 -	9.0 x 10^2	-	*A1*032:02 , B*004:01*
Others												
R10-002	wk14 -				wk59 -				wk104 -	5.7 x 10^5	-	
R10-006	wk14 -	wk30 -			wk55 -					1.0 x 10^6	-	

Group	Rhesus ID	mg/ml (IgG)	Collected/pooled (wks post-challenge)	10 $TCID_{50}$ SIV_{mac239} 100% killing titer
NAb (+)	R06-032	30	wk 76	1:16
	R01-012	30	wk 73–87	1:32
	R03-015	30	wk 112	1:8
	R02-004	30	wk 76	1:4
	R01-010	30	wk 77	1:16
	R06-019	30	wk 85	1:4
	R-360	30	wk 60–86	1:4
	R02-007	30	wk 42–70	1:32
	R05-010	30	wk 71	1:64
NAb (-)	R01-011	42	wk 104	< 1:2
uninfected	pooled	36	-	< 1:2

A

B

C

Anti-Env Ab 1 yr

A

Nef₆₂₋₇₀ QW9-specific responses in NAb non-inducers (viremic/Nef-G63E non-selecting)

Y-axis: Nef₆₂₋₇₀-specific CD8⁺ T-cell frequencies (/million PBMCs)

X-axis labels: R06-039, R01-011, R06-022, R10-001, R05-006, R06-034, R08-005

B

Y-axis: SIV p27 (ng/ml)

X-axis: Days post-SIV infection on HSC-F cells

Legend: WT, Nef-G63E

C

Y-axis: Plasma viral loads (vRNA copies/ml)

3 mo	6 mo	1 yr	1.5 yr
P = 0.2495	*P* = 0.5078	*P* = 0.8928	*P* = 0.6488

X-axis labels (each panel): NAb−, NAb+ Nef-G63E+

A

Nef-G63R selection

P = 0.068

% detected in group

NAb-   NAb+

B

Temporal Nef-G63E/R mutant selection vs NAb induction

R06-034

SIV 100% neutralizing titer (log$_2$)

Mutant SIV frequency (%)

Weeks post-SIV challenge

■ SIV$_{mac239}$-NAb
▼ Nef G63E
◆ Nef G63R

C

MOI 0.1 Day 3

pAkt Ser473

730
618
473

730
724
473

% of Max

PE-A :: pAkt473

PE-A :: pAkt473

Gated on singlet p27$^+$

— **WT**
— **Nef-G63E**
— **Nef-G63R**
— **Uninfected**

P = 0.70
P < 0.0001

ΔMFI (+ mean of uninfected)

WT   Nef G63E   Nef G63R

A

R3

CD19-PC5.5
Intensity_MC_Ch11

R4

Live/Dead-Aqua
Intensity_MC_Ch08

R4

R5

Intensity_MC_Ch11

Intensity_MC_Ch02

R4

R5

Intensity_MC_Ch11

Intensity_MC_Ch02

Nef-AF488

Intracellular pixel Nef signal
Intensity threshold

Signal intensity threshold (int/hi)

Signal intensity threshold (lo)

Live/Dead$^-$CD19$^+$
Nef$^+$ B cells

R5

Intensity_Threshold(M02 Ch2 30)_

R7

H Variance Mean_M02_Ch02_1

IDEAS6.3 machine learning-based 2-D
segregation of typical valid vs. void
Live/Dead$^+$Nef$^+$CD19$^+$ B-cell images

Good,bad

Void

Valid

Intensity_Threshold(M02 Ch0

H Variance Mean_M02_Ch02_

Intracellular pixel Nef signal
strength distribution

2-D adjacent pixel signal heterogeneity (++)
→Haralick variance mean (int-hi)

2-D adjacent pixel signal heterogeneity (+/-)
→Haralick variance mean (lo)

Live/Dead$^+$CD19$^+$Nef$^+$ B cells (void)

Ch01 | Ch02 | Ch08 | Ch11 | Ch02/Ch08/Ch11
246896
248952
71859
····7 µm

**Live/Dead$^-$CD19$^+$Nef$^+$-(Intracellular Nef Haralick variance
mean$^{int/hi}$or Nef Intensity threshold$^{int/hi}$) B cells (void)**

Ch01 | Ch02 | Ch08 | Ch11 | Ch02/Ch08/Ch11
18541
····7 µm

**Live/Dead$^-$CD19$^+$Nef$^+$-Intracellular Nef Haralick variance
mean^{lo}-Nef Intensity threshold^{lo} B cells (valid)**

Ch01 | Ch02 | Ch08 | Ch11 | Ch02/Ch08/Ch11
116318
116250
132927
····7 µm

B

***In vitro* CD19$^+$Nef$^+$ B cells**

DIC | Nef | CD19 | SSC | Merged

Ch01 | Ch02 | Ch05 | Ch06 | Ch02/Ch05
13088
··· 10 µm

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
gene (*Lentivirus*)	*env*, *nef*	GenBank	M33262.1	SIV_{mac239}
strain, strain background (*Lentivirus*)	SIV_{mac239}	PMID: 2032289		Obtained via COS-1 transfection of pBR_{mac239} and repropagation on MT4-R5 cells
strain, strain background (*Escherichia coli*)	One Shot™ TOP10 Chemically Competent *E. coli*	Invitrogen (Thermo Fisher Scientific)	Cat#:C404003	For subcloning sequencing
cell line (*Homo-sapiens*)	MT4-R5	PMID: 27122584		STR profiling (+)
cell line (*Chlorocebus aethiops*)	COS-1	ATCC	CRL-1650	STR profiling (+)
cell line (*Macaca fascicularis*)	HSC-F cells (CD4+ T-cell line)	PMID: 8610465	Gift from producer (Dr. Hirofumi Akari)	
cell line (*Macaca mulatta*)	HSR5.4 cells (CD4+ T-cell line)	PMID: 8610465	Gift from producer (Dr. Hirofumi Akari)	
cell line (*Homo-sapiens*)	Ramos B cell line	ATCC	CRL-1596	STR profiling (+)
biological sample (*Macaca mulatta*)	Primary simian PBMCs from SIV-infected macaques	Japan SLC	http://www.jslc.co.jp/service/	Cryopreserved from *Macaca mulatta*

biological sample (*Macaca mulatta*)	Primary simian lymph node cells from SIV-infected macaques	Japan SLC	http://www.jslc.co.jp/service/	Cryopreserved from *Macaca mulatta*
antibody	Mouse monoclonal anti-human CD4-FITC (M-T477)	BD Pharmingen	Cat#556615	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD8-PerCP (SK1)	BD Biosciences	Cat#347314	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD3-APC (SP34-2)	BD Pharmingen	Cat#557597	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human IFN-γ-PE (4S.B3)	BioLegend	Cat#502510	FACS (5 µl per test)
antibody	Mouse monoclonal anti-SIV$_{mac251}$ Gag p27 mAb	ABL	Cat#4324	FACS (1:500 with conjugation)
antibody	Mouse monoclonal anti-SIV$_{mac251}$ Nef mAb (clone 17)	Themo Fisher Scientific	Cat#MA1-71522	FACS, PLA, immuno-blotting and imaging cytometry (FACS and imaging cytometry, 1:500 with conjugation; PLA, 1:100; immuno-blotting, 1:1000)
antibody	Mouse IgG1 isotype control mAb (P3.6.2.8.1)	Themo Fisher Scientific	Cat#14-4714-85	PLA (1:100)

antibody	Rabbit monoclonal Alexa Fluor 488-conjugated anti-human phospho-Akt (Ser473) (D9E)	Cell Signaling Technology	Cat#4071	FACS (5 μl per test)
antibody	Rabbit monoclonal Alexa Fluor 488-conjugated anti-human phospho-Akt (Thr308) (C31E5E)	Cell Signaling Technology	Cat#2918	FACS (5 μl per test)
antibody	Rabbit monoclonal PE-conjugated anti-human phospho-Akt (Ser473) (D9E)	Cell Signaling Technology	Cat#5315	FACS and imaging cytometry (1:100)
antibody	Rabbit monoclonal Alexa Fluor 647-conjugated anti-human phospho-Akt (Ser473) (D9E)	Cell Signaling Technology	Cat#4075	FACS (5 μl per test)
antibody	Rabbit monoclonal Alexa Fluor 647-conjugated anti-human phospho-Akt (Thr308) (C31E5E)	Cell Signaling Technology	Cat#3375	FACS (5 μl per test)
antibody	Mouse monoclonal anti-human CD19-PC-5.5 (J3-119)	Beckman Coulter	Cat#66328	FACS and imaging cytometry (5 μl per test)
antibody	Mouse monoclonal anti-human CD4, purified (clone L200)	BD Biosciences	Cat#550625	FACS (25 μg/ml per test for blockade)
antibody	Mouse monoclonal anti-human CD3-	BD Pharmingen	Cat#557757	FACS (5 μl per test)

	APC-Cy7 (SP34-2)			
antibody	Mouse monoclonal anti-human CD8-APC-Cy7 (RPA-T8)	BD Pharmingen	Cat#557760	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD14-APC-Cy7 (M5E2)	BioLegend	Cat#301820	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD16-APC-Cy7 (3G8)	BioLegend	Cat#302018	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD10-APC-Cy7 (HI10a)	BioLegend	Cat#312212	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD11c-PE-Cy7 (3.9)	BioLegend	Cat#301608	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD123-PE-Cy7 (6H6)	BioLegend	Cat#306010	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human HLA-DR-PE-Texas Red (TU36)	Invitrogen (Thermo Fisher Scientific)	Cat# MHLDR17	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD27-PE/Dazzle 594 (M-T271)	BioLegend	Cat#356422	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD10-PE-Cy7 (HI10a)	BioLegend	Cat#312214	FACS (5 µl per test)

antibody	Rabbit polyclonal anti-human IgD-FITC F(ab')$_2$	DaKo	Cat#F0189	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD38-Alexa 647 (AT1)	Santa Cruz Biotechnology	Cat# sc-7325 AF647	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human IgG-Alexa 700 (G18-145)	BD Pharmingen	Cat#561296	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD138-PE (DL-101)	eBioscience (Thermo Fisher Scientific)	Cat#12-1389-42	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human IgG-PE (G18-145)	BD Biosciences	Cat#555787	FACS (5 µl per test)
antibody	Rat monoclonal anti-human/mouse IRF4-eFluor 660 (3E4)	eBioscience (Thermo Fisher Scientific)	Cat#50-9858-80	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human Ki67-Alexa 700 (B56)	BD Pharmingen	Cat#561277	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CD4-PerCP (L200)	BD Pharmingen	Cat#550631	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CD3-Alexa 700 (SP34-2)	BD Pharmingen	Cat#561805	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CD95-PE-Cy7 (DX2)	eBioscience (Thermo Fisher Scientific)	Cat#25-0959-42	FACS (5 µl per test)

antibody	Mouse monoclonal anti-human anti-CXCR5-PE (87.1)	eBioscience (Thermo Fisher Scientific)	Cat#12-1859-42	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human PD-1-Brilliant Violet 421 (EH12.2H7)	BioLegend	Cat#329920	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CXCR3-Alexa 488 (G025H7)	BioLegend	Cat#353710	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human HLA-ABC-FITC (G46-2.6)	BD Biosciences	Cat#557348	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CXCR4-APC (12G5)	BioLegend	Cat#306510	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human BST2-Alexa647 (RS38E)	BioLegend	Cat#348404	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD20 (2H7)	Biolegend	Cat#302324	FACS (5 µl per test)
antibody	Rabbit monoclonal anti-GβL (86B8)	Cell Signaling Technology	Cat#3274	PLA (1:100)
antibody	Rabbit monoclonal anti-mTOR (7C10)	Cell Signaling Technology	Cat#2983	PLA (1:100)
antibody	Rabbit polyclonal anti-human PI3K p85	Merck Millipore	Cat#ABS1856	PLA (1:100)

antibody	Rabbit monoclonal anti-PI3 Kinase p110α (C73F8)	Cell Signaling Technology	Cat#4249	PLA (1:100)
antibody	Goat polyclonal HRP-conjugated anti-monkey IgG (H+L)	Bethyl Laboratory	Cat# A140-102P	ELISA (1:1000)
antibody	Rabbit monoclonal anti-Sin1 (D7G1A)	Cell Signaling Technology	Cat#12860	Co-immuno-precipitation (1:50)
antibody	Mouse TrueBlot™ ULTRA anti-mouse Ig HRP (eB144)	Rockland Immuno-chemicals	Cat# 18-8817-30	Immuno-blotting (1:1000)
antibody	Mouse monoclonal anti-CD3 (FN-18)	Abcam	Cat#ab25109	Stimulation assay (1 µg/ml)
antibody	CD20 MicroBeads, non-human primate	Miltenyi Biotec	Cat#130-091-105	
recombinant DNA reagent	pBR$_{mac239}$	PMID: 2032289	GenBank M33262	coding full sequence of *nef*-open SIV$_{mac239}$
sequence-based reagent	9520GA-F	This paper	PCR primers	GGGACAG AAATACAA TCAGGAA CAGTATAT GAATACTC CATGGAG
sequence-based reagent	9520GA-R	This paper	PCR primers	CTCCATG GAGTATTC ATATACTG TTCCTGAT TGTATTTC TGTCCC
sequence-based reagent	9519GA-F	This paper	PCR primers	GAGGGAC AGAAATAC AATCAGA GACAGTA

				TATGAATA CTCCAT
sequence-based reagent	9519GA-R	This paper	PCR primers	ATGGAGT ATTCATAT ACTGTCTC TGATTGTA TTTCTGTC CCTC
peptide, recombinant protein	Peptide Nef$_{62-70}$	Sigma Aldrich	CTL epitope peptide	QGQYMNT PW
peptide, recombinant protein	Peptide Nef$_{62-70}$.G63E	Sigma Aldrich	Mutant CTL epitope peptide	QEQYMNT PW
peptide, recombinant protein	Peptide Nef$_{62-70}$.G63R	Sigma Aldrich	Mutant CTL epitope peptide	QRQYMNT PW
peptide, recombinant protein	GolgiStop	BD Biosciences	Cat#554724	
peptide, recombinant protein	recombinant SIV$_{mac251}$ Env gp130	ImmunoDx	Cat#5001	Ligand stimulation assay (10 µg/ml)
peptide, recombinant protein	recombinant human IL-2	Roche Diagnostics	Cat#1114752 8001	Ligand stimulation assay (100 IU/ml)
peptide, recombinant protein	recombinant human IFN-γ	Gibco (Thermo Fisher Scientific)	Cat#PHC403 3	Ligand stimulation assay (40 ng/ml)
peptide, recombinant protein	recombinant SIV$_{mac239}$ Env gp140 (biotin-conjugated)	Immune Technology	Cat#IT-001-140p	FACS and imaging cytometry (10µg/ml for probing)

peptide, recombinant protein	streptavidin-Brilliant Violet 421	BioLegend	Cat#405225	FACS (1:100)
peptide, recombinant protein	streptavidin-Alexa Fluor 488	BioLegend	Cat#405235	FACS (1:100)
peptide, recombinant protein	streptavidin-PE-Cy7	BioLegend	Cat#405206	Imaging cytometry (1:100)
peptide, recombinant protein	Recombinant Human IL-4 Protein (carrier-free)	R&D Systems	Cat# 204-IL-050/CF	Functional assay (40 ng/ml)
commercial assay or kit	High Pure Viral RNA kit	Roche Diagnostics	Cat#1185888 2001	
commercial assay or kit	SIV p27 ELISA	ABL	Cat#5436	
commercial assay or kit	SureBlue TMB 1-Component Microwell Peroxidase Substrate	KPL	Cat# 5120-0075	
commercial assay or kit	Prime-Script one-step RT-PCR kit v2	TaKaRa	Cat#RR055A	
commercial assay or kit	KOD-Plus v2	Toyobo	Cat#KOD-211	
commercial assay or kit	Zero Blunt TOPO PCR cloning kit for sequencing	Invitrogen (Thermo Fisher Scientific)	Cat#K287540	
commercial assay or kit	Cytofix/Cytoperm kit	BD Biosciences	Cat#554714	
commercial assay or kit	Monkey Ferritin (Sandwich	LS Bio	Cat#LS-F36490-1	

	ELISA) ELISA Kit			
commercial assay or kit	QuikChange II XL site-directed mutagenesis kit	Agilent Technologies	Cat#200522	
commercial assay or kit	RNeasy Plus Mini kit	Qiagen	Cat#74136	
commercial assay or kit	GeneChip WT Plus reagent	Affymetrix (Thermo Fisher Scientific)	Cat#902280	
commercial assay or kit	GeneChip Human Gene 2.0 ST Array	Affymetrix (Thermo Fisher Scientific)	Cat#902113	
commercial assay or kit	Alexa Fluor™ 488 Antibody Labeling Kit	Invitrogen (Thermo Fisher Scientific)	Cat#A20181	
commercial assay or kit	Alexa Fluor™ 647 Antibody Labeling Kit	Invitrogen (Thermo Fisher Scientific)	Cat# A20186	
commercial assay or kit	Duolink In Situ PLA Probe anti-mouse PLUS	Sigma (Merck)	Cat#DUO92001	
commercial assay or kit	Duolink In Situ PLA Probe anti-rabbit MINUS	Sigma (Merck)	Cat#DUO92005	
commercial assay or kit	Duolink flow PLA detection kit-Green	Sigma (Merck)	Cat#DUO94002	
commercial assay or kit	FuGENE6	Promega	Cat#E2691	
commercial assay or kit	Capturem IP & Co-IP Kit	Clontech/Takara	Cat#635721	

software, algorithm	FACS Diva	BD Biosciences	http://www.bdbiosciences.com/jp/instruments/software/facsdiva/	
software, algorithm	FlowJo v10.7.1	BD/Treestar	https://www.flowjo.com/solutions/flowjo	
software, algorithm	Affymetrix Expression Console Software	Affymetrix/Thermo Fisher Scientific	https://www.thermofisher.com/jp/ja/home/life-science/microarray-analysis/microarray-analysis-instruments-software-services/microarray-analysis-software/affymetrix-transcriptome-analysis-console-software.html	
software, algorithm	IDEAS 6.3 (+Machine learning module)	Amnis/Merck Millipore/Luminex	https://www.luminexcorp.com/imaging-flow-cytometry/	
software, algorithm	Image Lab software	BioRad	https://www.bio-rad.com/en-jp/product/image-lab-software	
software, algorithm	Prism 8	GraphPad Software	https://www.graphpad.com	
other	LIVE/DEAD® Fixable Aqua stain	Invitrogen (Thermo Fisher Scientific)	Cat#L34957	FACS and imaging cytometry (1:1000)

other	APC Annexin V	Biolegend	Cat#640920	FACS (5 µl per 100 µl test)
other	Annexin V Binding Buffer	Biolegend	Cat#422201	Diluent
other	LIVE/DEAD® Fixable Near-IR stain	Invitrogen (Thermo Fisher Scientific)	Cat#L10119	FACS (1:1000)
other	phorbol-12-myristate-13-acetate (PMA)	Sigma/Merck	Cat#P1585	Stimulation assay (10 ng/ml)
other	ionomycin	Sigma/Merck	Cat#I3909	Stimulation assay (10 ng/ml)
other	4–15% Mini-PROTEAN® TGX™ Precast Protein Gels	BioRad	Cat#4561085	
other	SuperSignal West Pico	Thermo Fisher Scientific	Cat#34580	