AIDS virus-neutralizing antibody induction reciprocal to a PI3K gain-of-function

disease

Hiroyuki Yamamoto[*1,2] and Tetsuro Matano[*1,3,4]

[1]AIDS Research Center, National Institute of Infectious Diseases, 4-7-1 Gakuen,

Musashi-Murayama City, Tokyo 208-0011, Japan

[2]Department of Biomedicine, University Hospital Basel, Hebelstrasse 20, Basel 4031,

Switzerland

[3]The Institute of Medical Science, The University of Tokyo, 4-6-1 Shirokanedai, Minato-ku,

Tokyo 108-8639, Japan

[4]Joint Research Center for Human Retrovirus Infection, Kumamoto University, 2-2-1 Honjo,

Kumamoto 860-0811, Japan

*Correspondence:

Hiroyuki Yamamoto, AIDS Research Center, National Institute of Infectious Diseases, 4-7-1

Gakuen, Musashi-Murayama City, Tokyo 208-0011, Japan. Phone: +81-42-848-7154; Fax:

+81-42-848-7155. Email: h-yamato@niid.go.jp

Tetsuro Matano, AIDS Research Center, National Institute of Infectious Diseases, 1-23-1

Toyama, Shinjuku-ku, Tokyo 162-8640, Japan. Phone: +81-3-4582-2811; Fax:

+81-3-5285-1165. Email: tmatano@niid.go.jp

Running title: SIV neutralization post-PI3K diminution

Abstract

HIV and simian immunodeficiency virus (SIV) infections are known for impaired neutralizing antibody (NAb) responses. While sequential virus-host B cell interaction appears to be basally required for NAb induction, driver molecular signatures predisposing to NAb induction still remain largely unknown. Here we describe SIV-specific NAb induction following a virus-host interplay reciprocal to a congenital human antibody-dysregulating disease. Screening of seventy neutralization-resistant SIV$_{mac239}$-infected macaques found nine NAb-inducing animals, with seven selecting for a specific CD8$^+$ T-cell escape mutation in viral *nef* before NAb induction. This mutation reduced aberrant Nef interaction-mediated drive of B-cell maturation-limiting phosphoinositide 3-kinase (PI3K)/mammalian target of rapamycin complex 2 (mTORC2). *In vivo* imaging cytometry depicted preferential Nef perturbation of cognate Envelope-specific B cells, corroborating cognate B-cell maturation post-mutant selection up to NAb induction. Results exemplify a NAb induction pattern extrinsically reciprocal to human PI3K gain-of-function antibody dysregulation, shaped through two-front, sequential virus interaction with both wings of adaptive immunity.

Keywords

neutralizing antibody; B cells; CD8$^+$ T cells; PI3 kinase; inborn errors of immunity; HIV/SIV

Introduction

Virus-specific neutralizing antibody (NAb) responses by B cells are induced by a tight cooperation of adaptive immune cells (Kumar et al., 2010; Shulman et al., 2014; Giltin et al., 2014; Wang et al., 2014) and often play a central role in clearance of acute viral infections (Junt et al., 2007). In contrast, persistence-prone viruses such as human immunodeficiency virus type 1 (HIV-1), simian immunodeficiency virus (SIV) and lymphocytic choriomeningitis virus (LCMV) variously equip themselves with B cell/antibody-inhibitory countermeasures (Moir et al., 2001; Mattapallil et al., 2005; Sommerstein et al., 2015; Sammicheli et al., 2016; Fallet et al., 2016; Mason et al., 2016) and NAb responses are impaired (Levesque et al.,2009; Mikell et al., 2011). These viruses successfully suppress elicitation of potent NAb responses particularly in acute infection (Hunziker et al., 2003; Tomaras et al., 2008) and establish viral persistence, posing considerable challenge for developing protective strategies.

Reciprocal to this detrimental acute-phase NAb absence, we and others have previously described that acute-phase passive NAb infusion can trigger an endogenous T-cell synergism resulting in robust SIV/SHIV (simian/human immunodeficiency virus) replication control (Haigwood et al., 1996; Yamamoto et al., 2007; Ng et al., 2010; Iseda et al., 2016; Nishimura et al., 2017). This shows that virus-specific NAbs not only confer sterile protection but also can evoke T-cell-mediated non-sterile viral control, suggesting the importance of inducing endogenous NAb responses during an optimal time frame as well as humoral-cellular response synergisms. Therefore, identifying mechanisms driving NAb induction against such viruses is an important step to eventually design NAb-based HIV control strategies.

As an approach, analysis of *in vivo* immunological events linked with NAb induction against difficult-to-neutralize SIVs in a non-human primate model can provide important insights into this goal. HIV-specific NAb induction patterns and correlates, such as antibody-NAb coevolution (Moore et al., 2012), autoimmune-driven induction (Moody et al.,

2016) and natural killer cell-related host polymorphisms (Bradley et al., 2018) have been

variously reported to date; the wealth of knowledge on this topic collectively, and interestingly,

rather suggests that pathways to NAb induction against neutralization-resistant viruses

including HIV/SIV can be highly redundant and there may be as-yet-unknown *in vivo*

mechanisms that can robustly drive NAb induction. For example, the neutralization resistance

of certain SIV strains apparently is not explanatory by the aforementioned, posing such

models as attractive analytic targets of NAb induction mechanisms.

In the present study, we examined virus-specific antibody responses in rhesus

macaques infected with a highly neutralization-resistant SIV strain, SIV_{mac239}. This virus is

highly pathogenic, shows persistent viremia and hosts lack NAb responses throughout

infection (Kestler et al., 1991). Yet, upon performing a large-scale screening of

SIV_{mac239}-infected rhesus macaques for up to 100 weeks, we noticed a subgroup inducing

NAbs in the chronic phase. Interestingly, these animals commonly selected for a specific

$CD8^+$ cytotoxic T lymphocyte (CTL) escape mutation in the viral Nef-coding gene before NAb

induction. The mutant Nef, compared with wild-type (WT) Nef, manifested a decrease in

aberrant interaction with phosphoinositide 3-kinase (PI3K)/mammalian target of rapamycin

complex 2 (mTORC2), resulting in decreased downstream hyperactivation of the canonical

B-cell negative regulator Akt (Omori et al., 2006; Limon et al., 2014). Nef was graphically

identified via machine learning-comprising imaging cytometry to preferentially target

Env-specific B cells *in vivo*, and in those macaques inducing NAbs, the NAb induction was

linked with sustained Env-specific B-cell responses after the mutant Nef selection. Results

describe a B-cell functional boosting reciprocal to human PI3K gain-of-function antibody

dysregulation, and suggest that intervening PI3K/mTORC2 signaling can potentially result in

harnessing NAb induction against difficult-to-neutralize viruses.

Results

Identification of macaques inducing SIV$_{mac239}$-neutralizing antibodies

99   We performed a retrospective antibody profile screening in rhesus macaques infected

with NAb-resistant SIV$_{mac239}$ (n = 70) (**Figure 1A**) and identified a group of animals inducing

anti-SIV$_{mac239}$ NAb responses (n = 9), which were subjected to characterization. These NAb

inducers showed persistent viremia with no significant difference in viral loads compared with

a profile-clarified naïve, major histocompatibility complex class I (MHC-I) haplotype-balanced

subgroup of NAb non-inducers (n = 19) representatively utilized for comparison hereafter

(**Figure 1B** and **Figure 1-figure supplement 1**). Plasma SIV$_{mac239}$-NAb titers measured by

10 TCID$_{50}$ SIV$_{mac239}$ virus-killing assay showed an average maximum titer of 1:16, being

induced at an average of 48 weeks post-infection (p.i.) (**Figure 1C**). Two of the nine NAb

inducers showed detectable NAb responses before six months p.i., while the remaining

seven induced NAbs after six months. Anti-SIV$_{mac239}$ neutralizing activity was confirmed in

immunoglobulin G (IgG) purified from plasma of these NAb inducers (**Figure 1-figure**

supplement 2). SIV Env-binding IgGs developed from early infection in NAb inducers as well

as NAb non-inducers (**Figure 1-figure supplement 3**); titers were not higher but rather lower

at year 1 p.i. in the NAb inducers (**Figure 1-figure supplement 3B** and **3C**), consistent with

other reports (Havenar-Daughton et al., 2016). NAb inducers and non-inducers showed

similar patterns of variations in viral *env* sequences (Burns et al., 1993), mainly in variable

regions 1, 2 and 4 (V1, V2 and V4) (**Figure 1-figure supplement 4**).

Selection of a viral *nef* mutation (Nef-G63E) precedes chronic-phase SIV$_{mac239}$-specific

NAb induction

Viral nonsynonymous polymorphisms outside *env* were next assessed. Strikingly, we

found selection of a viral genome mutation resulting in G (glycine)-to-E (glutamic acid)

substitution at residue 63 of Nef (Nef-G63E) in seven of the nine NAb inducers (**Figure 2A**).

This 63rd residue is in the unstructured N-terminus of Nef, flanked by conserved α-helices

(**Figure 2B**). This region is conserved in HIV-2/SIV but not in HIV-1, occasionally being

deleted in laboratory HIV-1 strains (Geyer et al., 2001). AlphaFold2-based structure

prediction did not derive palpable change except for low-probability disruption of the alpha

helices (data not shown). The Nef G63E mutation was not observed in two early NAb

inducers (with detectable NAbs before 6 months p.i.) but selected in all of the remaining

seven inducing detectable NAb responses after 6 months. In contrast, this mutation was

found only in two of the nineteen NAb non-inducers, including one rapid progressor (**Figure**

1-figure supplement 1) and selection was significantly enriched in the NAb inducers (**Figure**

2C). Analysis of Nef-G63E mutation frequencies in plasma viruses showed that Nef-G63E

selection preceded or at least paralleled NAb induction (**Figure 2D**).

Nef-G63E is a CD8$^+$ T-cell escape mutation

These seven NAb inducers selecting this mutation elicited CD8$^+$ T-cell responses

specific for a 9-mer peptide Nef$_{62-70}$ QW9 (QGQYMNTPW) (**Figure 3A**). This Nef$_{62-70}$ QW9

epitope is restricted by MHC-I molecules Mamu-B*004:01 and Mamu-B*039:01 (Evans et al.,

1999; Sette et al., 2012). Possession of these accounted for six cases of Nef-G63E selection

(**Figure 3A**), and the remaining one animal possessed Mamu-A1*032:02 predicted to bind to

Nef$_{63-70}$ GW8 (NetMHCpan). Ten of nineteen NAb non-inducers also had either of these

alleles (**Figure 1-figure supplement 1**), indicating that NAb induction is not associated with

these MHC-I genotypes (P = 0.25 by Fisher's exact test, data not shown) but with the

Nef-G63E mutation itself (**Figure 2C**). Nef-G63E was confirmed to be an escape mutation

from Nef$_{62-70}$-specific CD8$^+$ T-cell responses (**Figure 3B**). NAb non-inducers possessing

these alleles elicited little or no Nef$_{62-70}$-specific CD8$^+$ T-cell responses (**Figure 3-figure**

supplement 1A), suggesting that *in vivo* selection and fixation of this Nef-G63E SIV was

indeed Nef$_{62-70}$-specific CD8$^+$ T cell-dependent. Replication kinetics of SIV carrying the

Nef-G63E mutation was comparable with wild-type (WT) on a cynomolgus macaque HSC-F

CD4$^+$ T-cell line (Akari et al., 1996), stating that the mutation does not critically impair viral

replication (**Figure 3-figure supplement 1B**), and plasma viral loads were comparable

between Nef-G63E mutant-selecting NAb inducers versus non-inducers (**Figure 3-figure**

supplement 1C). These somewhat contrasted conditional reversion of a more

replication-impairing *nef* mutation related with CD3 downregulation loss in SIV_{mac239} infection

(Joas et al., 2020). These results indicate NAb induction following *in vivo* selection and

fixation of the CD8[+] T-cell escape *nef* mutation, Nef-G63E.

G63E mutation reduces aberrant Nef interaction-mediated drive of PI3K/mTORC2

We next focused on the functional phenotype of this highlighted Nef-G63E mutant SIV

in infected cells. An essence of host perturbation by Nef is its wide-spectrum molecular

downregulation, ultimately increasing viral replication (Schindler et al., 2004; Zhang et al.,

2009). To address possible amelioration, we compared downregulation of major targets CD3,

CD4, MHC-I, CXCR4 and BST-2 (Jia et al., 2009) in infected HSC-F cells. Nef-G63E mutation

did not confer notable change compared with wild-type (WT) [*P* = not significant (*n.s.*) for all

molecules, data not shown], implicating other non-canonical changes (**Figure 4A**). Literature

suggested ferritin negative influence on B cells proportionate to HIV-infected

macrophage-derived production (Swingler et al., 2008). Here, plasma ferritin showed no

differences (**Figure 4-figure supplement 1A**) and viral loads were comparable between

Nef-G63E-selecting NAb inducers and non-inducers (**Figure 4-figure supplement 1B** and

1C), arguing against gross involvement of ferritin as well as other viral replication-changing

Nef phenotypes at least in this model.

Thus, we took interest in immunosignaling-modulating properties of Nef-G63E SIV.

Analysis of Akt, the predominant immune-intrinsic negative brake of B-cell maturation and

AFC/antibody responses *in vivo* (Omori et al., 2006; Limon et al., 2014; Fruman et al., 1999;

Ray et al., 2015; Sander et al., 2015), revealed that its serine 473 phosphorylation,

non-canonically known for Nef-mediated upregulation (Kumar et al., 2016), was significantly

lower on day 3 after Nef-G63E SIV infection at low multiplicity of infection (MOI) compared

with WT (**Figure 4B**), stronger than histogram deviation levels in PI3K gain-of-function mice

with full disease phenotype (Avery et al., 2018). The threonine 308-phosphorylated Akt (pAkt

Thr308) level was more unaffected (**Figure 4-figure supplement 1B**).

We further assessed properties of serine 473-phoyphorylated Akt (pAkt Ser473)

inhibition by Nef-G63E SIV, and found that external PI3K stimuli (serum starvation) (Kennedy

et al., 1997) accelerated phenotype appearance from day 3 to day 1 post-infection (**Figure**

4C, left). A similar trend was obtained by short-term PI3K stimulation with interferon-γ

(Nguyen et al., 2001), interleukin-2 (Marzec et al., 2008) and SIV Env (François & Klotman,

2003) (**Figure 4-figure supplement 1C**). Env-stimulating high-MOI infection also

accelerated phenotype appearance, with enhanced 50% reduction (**Figure 4C**, right). These

showed that the Akt-inhibitory G63E mutant Nef phenotype is PI3K stimuli-dependent.

Transcriptome analysis signatured decrease in PI3K-pAkt-FoxO1-related genes in mutant

SIV infection (**Figure 4-figure supplement 1D**). As Akt is also a survival mediator this

implicates that Nef-G63E mutation may be Th-cytopathic, decreasing Th dysfunctional

expansion in LCMV (Recher et al., 2004) and HIV/SIV (Gray et al., 2011; Lindqvist et al.,

2012; Petrovas et al., 2012) infections. Here, reduced peripheral $CXCR3^-CXCR5^+PD-1^+$

memory follicular Th (Tfh) (Locci et al., 2013) was in keeping (**Figure 4-figure supplement**

1E and **1F**).

We further explored molecular traits of this decreased Akt hyperactivation. We

reasoned that comparing endogenous Nef binding patterns would be adequate and analyzed

SIV-infected HSC-F cells with a flow cytometry-based proximity ligation assay (PLA)

(Leuchowius et al., 2009; Avin et al., 2017), by calculating a mean fluorescence intensity

(MFI) deviation-based binding index for each. We found that this Nef-G63E mutation causes

significant decrease in Nef binding to PI3K p85/p110α and downstream mTORC2

components mTOR (Sarbassov et al., 2005) and GβL/mLST8 (Kim et al., 2003) in the

$CD4^{lo}$-SIV Gag $p27^+$ infected population (**Figure 4D**). Collectively, the Nef-G63E mutation

results in decreased aberrant Nef bridging and signal transduction of PI3K/mTORC2,

explaining decreased substrate Akt Ser473 phosphorylation by mTORC2. Thus, Nef-G63E

SIV is a mutant virus decreased in aberrant interaction/drive of B-cell-inhibitory

PI3K/mTORC2 signaling. This molecular signature was reciprocal to a recently found human

inborn error of immunity (IEI), activated PI3 kinase delta syndrome (APDS) (Angulo et al.,

2013; Lucas et al., 2014; Avery et al., 2018).

Targeting of lymph node Env-specific B cells by Nef *in vivo*

Next, we analyzed *in vivo* targeting of virus-specific B cells by Nef in lymph nodes to

explore the potential B-cell-intrinsic influence of the G63E-Nef phenotype. Previously

suggested influence of soluble Nef itself (Qiao et al., 2006) and/or related host factors

(Swingler et al., 2008) may derive generalized influence on B cells; however, SIV

antigen-specific binding antibody responses were rather decreased in NAb inducers (**Figure**

1-figure supplement 1), viremia was comparable (**Figure 3-figure supplement 1C**) and

ferritin levels were not altered (**Figure 4-figure supplement 1A**), arguing against these.

Therefore, we surmised that some targeted Nef intrusion against Env-specific B cells may be

occurring and that the decreased aberrant Nef-PI3K/mTORC2 drive may result in their

enhanced lymph node maturation.

While reports (Xu et al., 2009) histologically proposed Nef B-cell transfer, quantitative

traits, e.g., invasion frequency and influence on virus-specific B cells, have remained

unvisited. Intracellular Nef staining is dim (particularly for B cell-acquired Nef), difficult to

examine by conventional flow cytometry. Defining staining cutline requires single-cell images,

overcoming confounding of high Nef false-staining via pre-permeation rupture and

post-permeation confounding as judged by biologically discontinuous staining and/or

batch-inflated signals. Collectively, at the expense of spatial information, we judged that

sophisticating imaging cytometry would best visualize Nef-mediated B-cell perturbation *in*

vivo. We analyzed lymph node B cells with Image Stream X MKII, with high-power (> 10-fold)

antigen detection [e.g., molecules of equivalent soluble fluorochrome (MESF) 5 *vs.* MESF 80

in an average flow cytometer for FITC detection] ideal for detection/single-cell verification.

We designed a triple noise cancellation strategy to overcome the above. Firstly, amine

reactive dye staining (Perfetto et al., 2006) gated out B cells being Nef-positive due to

pre-experimental membrane damage (**Figure 5-figure supplement 1A**, left). Next, we

deployed secondary quantitative parameters of Nef signals deriving from each pixel of the images. Definition of a gray-level cooccurrence matrix (GLCM) (Haralick et al., 1973) numerating adjacent signal deviation as their frequencies on a transpose matrix enables calculation of a variety of feature values summarizing traits of the whole image. A biological assumption of Nef signal continuity in a true staining suggested that the sum of weighted square variance of GLCM, i.e., the {sum of [square of (value − average signal strength) × frequency of each value occurrence]} would separate natural versus artificial Nef signals. This value, Haralick variance, was computed for multiple directions and averaged, deriving Haralick variance mean (**Figure 5-figure supplement 1A**, X axis), proportionate with staining image unnaturalness. Finally, Nef signal intensity threshold (**Figure 5-figure supplement 1A**, Y axis) gated out overtly-stained cells void for true-false verification. These excluded B cells showing non-specific, strong-signal binary-clustered staining pixels for Nef deriving a large summated variance (X axis), and/or cells batch-stained for Nef (Y axis), likely originating from post-experimental membrane damage.

   With this approach we acquired images of a Live/Dead$^-$-Nef signal Haralick variance mean^{lo}-Nef signal intensity threshold^{lo}-Nef$^+$CD19$^+$ B-cells, with fine-textured gradation of Nef^{int-lo} staining with continuity from membrane-proximal regions without sporadic staining speckles (**Figure 5A** and **Figure 5-figure supplement 1A**). Images resembled Nef-transferred B cells generated *in vitro* by a SIV-infected CD4+ T-cell coculture (**Figure 5-figure supplement 1B**). A linear discriminant analysis-based machine learning module, scoring segregation of typical void/valid images, showed this gating to provide the highest two-dimensional separation (**Figure 5-figure supplement 1A**). Combined with biotinylated Env/streptavidin staining, this reproducibly obtained images of Env-specific (Ch 12) CD19$^+$ (Ch 11) B cells without membrane ruptures (Ch 08) and showing fine-textured transferred Nef (Ch 02) (**Figure 5B**). Nef invasion upregulated pAkt Ser473 (Ch 03) to a range resembling *in vitro* analysis (**Figure 5C**), stating that Nef-driven aberrant PI3K/mTORC2 signaling does occur in Nef-invaded B cells *in vivo* and the current phenotypic change likely affects B-cell dynamics. Strikingly, lymph node Env-specific B cells showed significantly higher Nef-positive

frequencies as compared with batch non-Env-specific B cells (**Figure 5D**). This indicated that

infected cell-generated Nef preferentially targets adjacent Env-specific B cells, putatively

through infected CD4$^+$ cell-to-B-cell contact/transfer (Xu et al., 2009; Hashimoto et al., 2016)

and that a Nef phenotypic change likely dictates Env-specific B-cell maturation.

Enhanced Env-specific B-cell responses after PI3K-diminuting mutant selection

Finally, peripheral SIV Env-specific IgG$^+$ B-cell responses comprising plasmablasts

(PBs) and memory B cells (B$_{mem}$) were analyzed to assess *in vivo* B-cell quality in NAb

inducers with Nef-G63E mutant selection. After excluding lineage-specific cells [T/NK/pro-B

cells/monocytes/myeloid dendritic cells (DCs)/plasmacytoid DCs], we defined IgG$^+$ PBs as

showing: high replication (Ki-67$^+$), post-activation (HLA-DR$^+$), transcriptional switching for

terminal differentiation (IRF4^{hi}) and downregulated antigen surface binding (surface

[s]Env^{lo}-cytoplasmic [Cy]Env$^+$) (Nutt et al., 2015; Silveira et al, 2015). Env-specific PBs were

defined as CD3$^-$CD8$^-$CD10$^-$CD14$^-$CD16$^-$CD11c$^-$CD123$^-$CD19$^-$CyIgG$^+$Ki-67$^+$IRF4^{hi}HLA-DR$^+$

(total PB) sEnv^{lo}CyEnv$^+$ (Env-specific) (**Figure 6-figure supplement 1**). NAb inducers

showed significantly higher Env-specific IgG$^+$ PB responses around week 30 p.i., after

Nef-G63E selection, compared to those in NAb non-inducers (**Figures 6A** and **6B**). At 1 year,

this difference became enhanced; Env-specific PBs were reduced in non-inducers, while this

reduction was ameliorated in the NAb inducers with Nef-G63E.

Concomitant determination of Env-specific IgD$^-$CD27$^+$IgG$^+$ B$_{mem}$ responses allowed

pair-wise analysis of Env-specific IgG$^+$ B$_{mem}$/PB as a projection of germinal center (GC)

output as an indicator of overall B-cell response quality (Zotos et al., 2010) (**Figure 6C**, left).

In this two-dimensional vector charting of Env-specific B$_{mem}$/PB responses, vector protrusion

towards upper right reasonably represents higher gross GC output of antibody-forming cells

(AFCs) (B$_{mem}$/PBs). In NAb non-inducers, all vectors converged on an empirically defined

polygonal attractor area D_n (gray area surrounded with dotted lines in **Figure 6C**, right) at

year 1 p.i. and beyond, describing that these NAb non-inducers failed to sustain Env-specific

GC output. In contrast, the vectors were consistently tracked outside D_n in the NAb inducers

with Nef G63E (**Figure 6C**). At the moment of NAb induction, they converged on another

upper-right GC output attractor area D_i (red area surrounded with dotted lines in **Figure 6C**,

right), mutually exclusive with D_n (P < 0.0001 by Fisher's exact test on NAb

non-inducer/inducer vector distribution frequency within D_n). These suggest more robust

virus-specific IgG$^+$ B-cell responses following Nef-G63E CD8$^+$ T-cell escape mutant selection,

predisposing to NAb induction. This enhanced cognate B-cell signature was also an inverted

pattern of impaired terminally-differentiated B-cell responses in APDS (Al Qureshah et al.,

2021).

Taken together, in the current model, Nef$_{62-70}$-specific CD8$^+$ T-cell responses in

persistently SIV$_{mac239}$-infected macaques selected for an escape mutant, Nef-G63E SIV,

which decreased aberrant Nef binding and drive of B-cell-inhibitory PI3K/mTORC2. B-cell Nef

invasion *in vivo* occurred more preferentially in Env-specific B cells, suggesting diminution in

tonic Nef-mediated B-cell dysregulation after mutant selection. These predisposed to

enhanced Env-specific B-cell responses and subsequent SIV$_{mac239}$-specific NAb induction,

altogether, in a manner reciprocal to human APDS-mediated immune dysregulation.

Discussion

In the present study, we found NAb induction against an NAb-resistant SIV after selection of a CD8[+] T-cell escape *nef* mutant virus. The Nef-G63E mutation, positioned in cognate B cells, results in reduction of binding/drive of PI3K/mTORC2, being concordant with *in vivo* enhancement of Env-specific IgG[+] B cells up to NAb induction following mutant selection. Importantly, this manifested through a pAkt deviation level more vivid than wild-type versus germline *PIK3CD* gain-of-function mutation heterozygote mice showing full APDS phenotype, stating that the current Nef-G63E-associated B-cell/NAb phenotype likely occurs in a "reciprocal APDS-like" manner. This proposition was enhanced based on extension of immune cell-intrinsic Nef influence on cognate B cells (Fig. 5), in addition to infected T cells. This work is, to our knowledge, the first to interlink a PI3K/mTORC2-modulating viral signature and enhanced B-cell/NAb responses in a viral infection model.

A link between viral T-cell escape and consequent immune modulation has been previously partially explored, including enhanced mutant HIV-1 epitope peptide binding to inhibitory MHC-I receptor on DCs impairing T cells (Lichterfeld et al., 2007) and decreased CTL-mediated lymph node immunopathology rather driving LCMV-specific antibody production (Battegay et al., 1993). Our results now evidence a new pattern of NAb responses bivalently shaped through viral interactions with both humoral and cellular immunity in AIDS virus infection. SIV$_{mac239}$ infection of macaques possessing MHC-I alleles associated with Nef-G63E mutation can be one unique model to analyze virus-host interaction for B cell maturation leading to NAb induction. A *nef* polymorphism was extracted, partially appending to a virological impact of Nef (Kirchhoff et al., 1995; Gauduin et al., 2006; Stolp et al., 2012) in perturbing immunity whereas the current phenotype differed from such replication-related properties. The *nef* region encoding Nef-G63E is not conserved in HIV-1 (Schindler et al., 2004) and beyond this model study it remains to be addressed whether mutant HIV-1 with a similar phenotype can be obtained.

The major *in vivo* readout of this study is autologous neutralization of a highly resistant

SIV, differing from HIV-specific broadly neutralizing antibodies (bNAbs). Importantly, however,

obtained signatures of enhanced IgG[+] Env-specific AFC B cells (class switch/terminal

differentiation) and virus neutralization (hypermutation) directly evidence enhanced activity of

activation-induced cytidine deaminase (AID), the canonical positive driver of B-cell fate,

showing a strong conceptual continuity with bNAb regulation by AID.

Our results extracted an *in vivo* link between a decreased aberrant Nef-PI3K/mTORC2

interaction and major enhancement in B-cell responses. The relationship is indeed reciprocal

to immunogenetic mechanisms of human PI3K gain-of-function mutations resulting in APDS

with multiply impaired anti-viral B-cell responses. While the exact hierarchy of binding

between Nef and PI3K/mTORC2 components (including PI3K isoforms) and its perturbation

by Nef-G63E mutation remains much to be investigated, PI3Kp85 did reasonably show the

strongest Nef-binding index in PLA assay (**Figure 4D**) as Nef and PI3Kp85 possess

SH3-binding PxxP and SH3 domains, respectively (Rickles et al., 1994). Binding decrease in

canonical partner PI3Kp85 may domino-trigger decrease with others, resulting in quadruple

PI3K/mTORC2 binding decrease and fragile downstream Akt signaling.

In conclusion, we exemplify in a non-human primate AIDS model that NAb induction

against a highly resistant SIV occurs after selection of a CD8[+] T-cell escape variant reduced

in excess PI3K/mTORC2 drive. These results provide an understanding of how

immunodeficiency virus-specific NAb responses can be shaped by both wings of adaptive

immune pressure and depicts that immune cell-intrinsic PI3K/mTORC2 manipulation may be

pivotal for harnessing antiviral NAb responses. Furthermore, search of other analogies

between viral immune dysregulations and human IEIs may lead to identification of novel

targets to modulate and harness immune responses in translational settings.

Limitations of study

A limitation of this study is the use of retrospective samples, posing constraints for detecting

exact temporal changes in tonic viral B-cell perturbation. Related with this, sampling of lymph

node cells in animals belonging to the subgroup of interest at the optimal moment was

unpredictable. Nevertheless, usage of such specific animal models does help to depict a

temporal cascade of *in vivo* events leading to NAb induction, assisting human immunology.

As future extension of this study extracting the NAb-involved molecular axis,

manipulation/reconstitution experiments shall be designed.

Acknowledgments

This work was supported by Japanese Initiative for Progress of Research on Infectious

Diseases for Global Epidemic (J-PRIDE) (JP19fm0208017 to H.Y.), Japan Program for

Infectious Diseases Research and Infrastructure (Interdisciplinary Cutting-edge Research)

(JP22wm0325006 to H.Y.), Research Programs on HIV/AIDS, Emerging and Re-emerging

Infectious Diseases, and Global Health Issues (JP20fk0410022 to H.Y.; JP18fk0410003,

JP20fk0410011, JP20fk0108125, JP20jm0110012, JP21fk0410035, and JP21jk0210002 to

T.M.) from Japan Agency for Medical Research and Development, the Ministry of Education,

Culture, Sports, Science, and Technology in Japan ([JSPS] KAKENHI Grants 17H02185 and

18K07157 to H.Y.; 21H02745 to T.M.), the Takeda Science Foundation (to H.Y.) and the Imai

Memorial Trust for AIDS Research (to H.Y.). We thank Fumiko Ono, Koji Hanari, Katsuhiko

Komatsuzaki, Akio Hiyaoka, Hiromi Ogawa, Keiko Oto, Hirofumi Akari, Yasuhiro Yasutomi,

Hiromi Sakawaki, Tomoyuki Miura and Yoshio Koyanagi for animal experiment assistance;

Hiroyuki Kogure, Sayuri Seki and Julia R. Hirsiger for imaging cytometry technical assistance;

Masako Nishizawa, Trang Thi Thu Hau, Shigeyoshi Harada, Takushi Nomura, Akiko Takeda,

Taku Nakane, Nami Iwamoto, Taeko K. Naruse, Akinori Kimura and Mark S. de Souza for

their help. H.Y. thanks Jun Abe, Shoi Shi, Bettina Stolp, Jens V. Stein, Makoto Yamagishi and

Yusuke Yanagi for conceptual discussions.

Author Contributions

H.Y. conceived/designed/performed study and wrote manuscript; T.M. provided animal cohort,

co-edited manuscript and co-directed study.

Declaration of Interests

The authors declare no competing interests.

 Figure Legends

 Figure 1. NAb induction against NAb-resistant SIV$_{mac239}$.

 (A) Study design.

 (B) Plasma viral loads (SIV *gag* RNA copies/ml plasma) in NAb non-inducers (left) and

 inducers (right). Viral loads have been previously partially reported (Yamamoto et al., 2007;

 Nomura et al., 2012; Takahashi et al., 2013; Nakane et al., 2013; Iwamoto et al., 2014).

 (C) Plasma SIV$_{mac239}$ 100% neutralizing end point titers by 10 TCID$_{50}$ killing assay on MT4-R5

 cells. Points on the dotted line show marginally NAb-positive results (< 1:2). In some animals,

 titers were comparable with our reported results using MT4 cells (Kawada et al., 2007).

 Figure 2. Selection of a viral *nef* mutation (Nef-G63E) before NAb induction.

 (A) Viral *nef* mutations in NAb inducers. A linear schema indicating Nef functional domains is

 aligned above. Mutations at timepoints with NAb$^-$ (indicated by white wedge; mostly before 6

 months p.i.), NAb$^{+/-}$ (green wedge) and NAb$^+$) (purple wedge; mostly after 1 year) are shown

 in individual animals. Black and dark gray represent dominant and subdominant mutations (or

 deletion in R-360) by direct sequencing, respectively. Red and pink indicate dominant and

 subdominant G63E detection, respectively.

 (B) Schema of SIV$_{mac239}$ Nef structure and Nef-G63E mutation orientation.

 (C) Comparison of frequencies of macaques having Nef-G63E in plasma viruses between

 NAb non-inducers and inducers. Compared by Fisher's exact test.

 (D) Temporal relationship of Nef G63E frequencies in plasma virus and NAb induction. Black

 boxes (left Y axis) show log$_2$ NAb titers; red triangles and green diamonds (right Y axis) show

 percentage of G63E and G63R mutations detected by subcloning (15 clones/point on

 average), respectively.

 Figure 3. Nef-G63E is a CD8$^+$ T-cell escape mutation.

 (A) Nef$_{62-70}$ QW9-specific CD8$^+$ T-cell frequencies and related MHC-I alleles in seven NAb

inducers selecting Nef-G63E. Mamu-B*039:01 and Mamu-B*004:01 are known to restrict

Nef$_{62-70}$ QW9 epitope (Evans et al., 1999; Sette et al., 2012) and binding of Nef$_{63-70}$ peptide to

Mamu-A1*032:02 was predicted.

(B) CD8$^+$ T-cell responses specific to wild-type Nef$_{62-70}$ or mutant Nef$_{62-70}$.G63E or

Nef$_{62-70}$.G63R peptides. See also Figure S3.

Figure 4. Nef-G63E mutation reduces PI3K/mTORC2 binding and pAkt drive.

(A) Representative surface expression level histograms of CD3, CD4, MHC class I, CXCR4

and BST-2 in CD4^{lo}Nef$^+$ subpopulations after wild-type (WT) or Nef-G63E mutant SIV

infection at multiplicity of infection (MOI) 0.1 on HSC-F cells.

(B) Left: representative histograms of relative pAkt serine (Ser) 473 levels in p27$^+$

subpopulations after WT or Nef-G63E mutant SIV infection at MOI 0.1 on HSC-F cells.

Numbers show pAkt Ser473 mean fluorescence intensities (MFIs) for each. Right: Deviation

of pAkt Ser473 MFIs in p27$^+$ HSC-F cells compared with mean MFI of uninfected cells.

Compared by unpaired t test.

(C) Left: Relative pAkt Ser473 levels (normalized to mean MFI of uninfected controls) in Nef$^+$

HSC-F cells assessed for serum starvation (MOI 0.2, 1 day p.i.). Adjusted *P* values show

results of comparison via Sidak's post-hoc test of 2-way ANOVA (C, left).   Right: Deviation of

pAkt Ser473 MFIs in Nef$^+$ HSC-F cells assessed for high-MOI infection (MOI 5, 1-day p.i.).

Compared by unpaired t test.

(D) Proximity ligation assay (PLA) of Nef binding with mTOR, GβL/mLST8, PI3Kp85 and

PI3Kp110α. MFI-based binding index was calculated as (anti-Nef/anti-partner) -

(isotype/anti-partner) - (anti-Nef/isotype) + (isotype/isotype). Histograms for samples and

isotype/isotype are representatively shown. Differences in MFI binding indexes can be

enhanced compared with comparison of raw MFIs. Compared by paired t tests. Data

represent one of two [(A); (B); (C, right)] independent experiments in quadruplicate, four

independent experiments performed in triplicate [(C, left)] or four independent single-well

comparison experiments pooled for statistical analysis (D). Bars: mean ± SD (B, C right),

mean ± SEM (C left).

Figure 5. Preferential targeting of lymph node Env-specific B cells by Nef *in vivo*.

(A) Representative gating of triple noise-cancellation *in vivo* Nef staining in B cells analyzed

by ImageStreamX MKII. Pre-experimental damaged cells are first excluded with Live/Dead

from focused/centroid/singlet image-acquired CD19$^+$ B cells (first lane right/R4 gated on

"Focused"). Following Nef$^+$ gating (second lane left/R5 gated on R4), a second step of Nef

noise cancellation (second lane middle/R7 on R5) comprises double-negative removal of

post-experimental stochastic irregular staining building a disparate intracellular staining

gradient (X axis, Nef signal pixel Haralick variance mean) and post-experimental batch overt

cellular staining (Y axis, Nef signal pixel intensity threshold). This outputs a B-cell population

with a fine-textured pericellular Nef^{int-lo} staining, biologically concordant with Nef

membrane-anchoring. Probing of anti-Env BCR (αEnv) by recombinant SIV Env (second lane,

right) is combined, resulting in a 2-D panel of intracellular Nef versus αEnv for

noise-cancelled Nef$^+$ B cells (R7) plus Nef$^-$ B cells (R8) (third lane left/"R7+R8"). pAkt Ser473

expression (third lane right) and cellular morphology (B) was further analyzed. DN,

double-negative.

(B) Typical images of Nef-transferred Env-specific B cells defined αEnv$^+$-intracellular

Nef$^+$-Nef Haralick variance mean^{lo}-Nef Intensity threshold^{lo}-Live/Dead$^-$-CD19$^+$ cells

["Nef$^+$aEnv$^+$" population of lower left panel in (A), gated on "R7, R8"]. Note the pericellular

pAkt Ser473 upregulation in these cells (Ch 03/yellow). Data on inguinal lymph node

lymphocytes of macaque R10-007 at week 62 post-SIV$_{mac239}$ infection are shown in (A) and

(B).

(C) Comparison of pAkt Ser473 median fluorescence (medFI) intensity levels in Nef$^-$ B cells

(R8) versus noise-cancelled Nef$^+$ B cells (R7). Analyzed by paired t test.

(D) Comparison of Nef-positive cell frequencies in non-Env-specific (left) versus Env-specific

(right) B cells in lymph nodes of persistently SIV-infected macaques (*n* = 6). Analyzed by

paired t test.

Figure 6. Enhanced SIV Env-specific B-cell output up to NAb induction following Nef-G63E selection.

(A) Changes in SIV Env gp140-specific plasmablast (PB) frequencies in viremic NAb non-inducers (left, n = 12) and Nef-G63E-selecting NAb inducers (right, n = 6). Available samples of twelve viremic NAb non-inducers (including ten with $\geq$ 1-year survival) and six NAb inducers were tracked. In the right panel, the frequencies in NAb non-inducers are shown in background (gray) for comparison.

(B) Comparison of Env gp140-specific PB frequencies between NAb non-inducers and inducers by Mann Whitney U tests. In the right, the frequencies at NAb induction were compared with those at year 1 in NAb non-inducers with $\geq$ 1-year survival (n = 10). Bars: mean $\pm$ SD.

(C) Left: vector chart of Env gp140-specific memory B cell (B_{mem}) and PB levels. Legends for each animal correspond to the ones in (B). Right: NAb non-inducer vectors empirically define a polygonal GC output attractor area D_n (gray area surrounded with dotted lines) on which they converge by and beyond year 1 p.i. The NAb inducer vectors (shown up to the time of NAb induction) remained outside of D_n. At the moment of NAb induction they converged on a second GC output attractor area D_i (red area surrounded with dotted lines), mutually exclusive with D_n (P < 0.0001 by Fisher's exact test on NAb inducer vector convergence frequency within D_n). Legends for year 1 p.i. in the NAb non-inducers and moment of NAb induction in the NAb inducers are specified.

Materials and Methods

Material availability

This study did not generate new unique reagents.

Data and code availability

Data were analyzed using existing computational packages. The accession number for the transcriptome data [wild-type SIV_{mac239}-infected HSC-F cell line (n = 3), Nef-G63E SIV_{mac239}-infected HSC-F cell line (n = 3) and uninfected HSC-F cell line (n = 3)] has been deposited under the accession number GenBank: GSE65806.

Rhesus macaques retrospectively utilized for samples

Seventy Burmese rhesus macaques (*macaca mulatta*) (57 males and 13 females) were retrospectively analyzed in this study. Experiments were previously carried out (Matano et al., 2004; Yamamoto et al., 2007; Iseda et al., 2016; Nomura et al., 2012; Ishii et al., 2012; Takahashi et al., 2013; Nakane et al., 2013; Shi et al., 2013; Iwamoto et al., 2014; Terahara et al., 2014) in the Tsukuba Primate Research Center, National Institutes of Biomedical Innovation, Health and Nutrition (NIBIOHN) with the help of the Corporation for Production and Research of Laboratory Primates and the Institute for Frontier Life and Medical Sciences, Kyoto University (IFLMS-KU) after approval by the Committee on the Ethics of Animal Experiments of NIBIOHN and IFLMS-KU under the guidelines for animal experiments at NIBIOHN, IFLMS-KU and National Institute of Infectious Diseases in accordance with the Guidelines for Proper Conduct of Animal Experiments established by Science Council of Japan (http://www.scj.go.jp/ja/info/kohyo/pdf/kohyo-20-k16-2e.pdf). The experiments were in accordance with the "Weatherall report for the use of non-human primates in research" recommendations (https://royalsociety.org/topics-policy/publications/2006/weatherall-report/). Animals were housed in adjoining individual primate cages allowing them to make sight and sound contact with one another for social interactions, where the temperature was kept at

25^{o}C with light for 12 hours per day. Animals were fed with apples and commercial monkey

diet (Type CMK-2, Clea Japan, Inc.). Blood collection and virus challenge were performed

under ketamine anesthesia.

Cells and viruses

MT4-R5 cells were maintained in RPMI1640 (Invitrogen) supplemented with 10% fetal bovine

serum (FBS) (Clontech) and antibiotics. HSC-F cells (cynomolgus $CD4^+$ T-cell line) and

HSR5.4 cells (rhesus macaque $CD4^+$ T-cell line) were maintained in RPMI1640 (Invitrogen)

supplemented with 10% fetal bovine serum (FBS) (Clontech), human interleukin-2 (IL-2) (10

IU/ml, Roche), 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES) (Invitrogen) and

2-mercaptoethanol (Gibco). SIV_{mac239} molecular clone derivative was generated by

mutagenesis PCR (Agilent Technologies) using the primers listed. For virus preparation,

COS-1 cells were transfected with pBR_{mac239} proviral DNA using FuGENE 6 (Promega). At 48

h post-transfection, culture supernatants were harvested, centrifuged and filtered through a

0.45-µm pore-size filter (Merck Millipore). To titrate infectivity, prepared viruses were serially

diluted and infected on HSC-F cells in 96-well plates (Falcon) in quadruplicate. At 10 days

post-infection, the endpoint was determined using SIV p27 antigen enzyme-linked

immunosorbent assay (ELISA) kit (ABL), and virus infectivity was calculated as the 50

percent tissue culture infective dose ($TCID_{50}$) according to the Reed-Muench method.

Identification of SIV_{mac239}-NAb inducers

Burmese rhesus macaques previously challenged with the highly pathogenic molecular clone

virus SIV_{mac239} (n = 70) were retrospectively examined approximately up to 2 years for their

plasma NAb profiles. Animals were challenged intravenously with 1,000 $TCID_{50}$ of SIV_{mac239}.

In the current study, virological and immunological profiles were compared between the newly

identified NAb inducers (n = 9) and representative NAb non-inducers (n = 19). These NAb

non-inducers and eight NAb inducers except R06-032 were previously partially reported for

their plasma viral loads (Yamamoto et al., 2007; Iseda et al., 2016; Nomura et al., 2012;

Takahashi et al., 2013; Nakane et al., 2013; Iwamoto et al., 2014) (**Figure 1B**). MHC-I

haplotypes and alleles were determined by reference strand-mediated conformation analysis,

PCR-SSP (PCR amplification utilizing sequence-specific priming) and cloning as described

(Naruse et al., 2010). MHC-I binding prediction was made on April 27, 2013 using the IEDB

analysis resource NetMHCpan tool (Hoof et al., 2009). Alleles of interest in the study have

been previously identified in macaques (Nomura et al., 2012; Evans et al., 1999; Sette et al.,

2012). Macaques R06-032, R03-015, R01-010 and R05-010 received a prime-boost

vaccination (Matano et al., 2004) composed of a DNA prime/intranasal Sendai virus vector

expressing SIV_{mac239} Gag (SeV-Gag). R03-015, R06-019, R06-038 and R10-001 received

300 mg of non-specific rhesus IgG at day 7 post-SIV_{mac239} challenge as an experimental

control in our previous reports (Yamamoto et al., 2007; Iseda et al., 2016; Nakane et al.,

2013).

Plasma viral load quantitation

Plasma viral RNA samples were extracted with High Pure Viral RNA kit (Roche Diagnostics).

Serial fivefold sample dilutions were amplified in quadruplicate by reverse transcription and

nested PCR using SIV_{mac239} *gag*-specific primers to determine end point via the

Reed-Muench method as described previously (Matano et al., 2004; Iseda et al., 2016). The

lower limit of detection is approximately 400 viral RNA copies/ml plasma.

SIV_{mac239}-specific neutralization assay

NAbs were titrated as described (Yamamoto et al., 2007; Iseda et al., 2016). Serial twofold

dilutions of heat-inactivated plasma, or polyclonal IgG affinity-purified with Protein G

Sepharose 4 Fast Flow (GE Healthcare) from heat-inactivated and filtered plasma were

mixed with 10 $TCID_{50}$ of SIV_{mac239} at a 1:1 ratio (5 µl:5 µl) in quadruplicate. After 45-min

incubation at room temperature, the 10 µl mixtures were added into 5×10^4 MT4-R5 cells/well

in 96-well plates. Progeny virus production in day 12 culture supernatants was examined by

SIV p27 ELISA (ABL) to determine 100% neutralizing endpoint. The lower limit of titration is

1:2. Neutralization in three out of four wells at a dilution of 1:2 is defined as marginally

NAb-positive ($<$ 1:2). Results were comparable when the same assay was performed with

macaque HSC-F cells (Akari et al., 1996) as targets. NAb inducers R01-012, R02-004,

R01-010, R02-007 and NAb non-inducer R01-011 were previously partially reported for their

NAb titers measured with the same method using MT4 cells as targets (Kawada et al., 2007),

which derived comparable results. For assessment of neutralizing activity in IgG,

SIV_{mac239}-specific IgGs purified from pools of plasma with SIV_{mac239}-specific NAb titers were

obtained from each animal as described (Yamamoto et al., 2007). After complement

heat-inactivation at 56°C, 30 min and 0.45 μm filtration, IgGs were purified by Protein G

Sepharose 4 Fast Flow (GE Healthcare) and concentrated by Amicon Ultra 4, MW 50,000

(Millipore) to 30 mg/ml and similarly examined for their 10 $TCID_{50}$ SIV_{mac239} killing titers on

MT4-R5 cells.

SIV Env-specific IgG ELISA and immunoblotting

Plasma Env-specific IgG titers were measured as described (Nakane et al., 2013). SIV_{mac251}

Env gp120 (ImmunoDiagnostics) were coated on 96-well assay plates (BD) at 1000 ng/ml

(100 μl/well). Wells were prewashed with phosphate-buffered saline (PBS), blocked with

0.5% bovine serum albumin (BSA)/PBS overnight and plasma samples were incubated at a

1:20 dilution (5 μl:95 μl) for 2 hours. Wells were washed with PBS and SIV Env-bound

antibodies were detected with a horseradish peroxidase (HRP)-conjugated goat anti-monkey

IgG (H+L) (Bethyl Laboratory) and SureBlue TMB 1-Component Microwell Peroxidase

Substrate (KPL). Absorbance at 450 nm was measured. Samples from week 0 pre-challenge

and month 3, month 6 and year 1 post-challenge were assessed in duplicate. Week 0

average values were subtracted from corresponding later time point values for calibration.

For immunoblotting, SIV virion-specific IgGs in plasma were detected with a

SIV_{mac239}-cross-reactive western blotting system (ZeptoMetrix). In the NAb non-inducers,

samples from those showing rapid progression (euthanized due to AIDS progression within

approximately 1 year) with low plasma anti-SIV reactivity (Nakane et al., 2013) were not

included.

Sequencing

Sequencing was performed as described (Matano et al., 2004; Iseda et al., 2016). Viral cDNA

fragments spanning from nt (nucleotide) 4829 to nt 7000, nt 6843 to nt 8831 and nt 8677 to nt

10196 in SIV_{mac239} (GenBank accession number MM33262) covering SIV *env* and *nef* were

amplified from plasma viral RNA by nested RT-PCR using Prime-Script one-step RT-PCR kit

v2 (TaKaRa) and KOD-Plus v2 (Toyobo). PCR products were either directly sequenced or

subcloned with a TOP10-transforming TOPO blunt-end cloning system (Invitrogen).

Sequencing was performed using dye terminator chemistry with an ABI 3730 DNA sequencer

(Applied Biosystems). On average, 15 clones were obtained per sample and 20 clones were

assessed when *nef* mutations of interest in early time points were subdominant.

SIV_{mac239}-specific CD8$^+$ T-cell responses

Virus-specific CD8$^+$ T-cell frequencies were measured as described (Matano et al., 2004;

Iseda et al., 2016). Peripheral blood mononuclear cells (PBMCs) were cocultured for 6 hours

with autologous herpesvirus papio-immortalized B lymphoblastoid cell lines (B-LCLs) pulsed

with Nef peptides (Sigma-Aldrich Japan) at 1 µM concentration or as indicated otherwise

under GolgiStop (monensin, BD) presence. Intracellular gamma interferon (IFN-γ) staining

was performed using Cytofix/Cytoperm kit (BD) and the following conjugated anti-human

monoclonal antibodies (mAbs): anti-CD4-FITC (M-T477, BD Pharmingen), anti-CD8-PerCP

(SK1, BD Biosciences), anti-CD3-APC (SP34-2, BD Pharmingen) and anti-IFN-γ-PE (4S.B3,

BioLegend). Specific CD8$^+$ T-cell frequencies were determined by subtracting nonspecific

IFN-γ$^+$ CD8$^+$ T-cell frequencies from those after peptide-specific stimulation; frequencies

beneath 100 cells/million PBMCs were considered negative. Cells acquired by FACS Canto II

(BD) were analyzed by FACS Diva (BD) and FlowJo (Treestar). Approximately 1 x 10^5

PBMCs were gated per test.

Nef-mediated signaling perturbation analysis

Virus supernatants obtained from COS-1 cells after transfection with wild-type or mutant

SIV_{mac239} molecular clones were used for infection of $CD4^+$ T-cell lines, cynomolgus

macaque-derived HSC-F and rhesus macaque-derived HSR5.4 (Akari et al., 1996).

QuikChange II XL site-directed mutagenesis kit (Agilent Technologies) was used to construct

a mutant SIV_{mac239} molecular clones possessing a *nef* mutation Nef G63E (G-to-A mutation at

nt 9520) from the wild-type SIV_{mac239} molecular clone (Kestler et al., 1991) (nt number from

GenBank accession number M33262). Cells (1 x 10^5 cells/well in U-bottomed 96-well culture

plates [BD]) were infected with wild-type or mutant SIV_{mac239} cultured in RPMI-1640 medium

supplemented with 10% fetal bovine serum (FBS) for indicated periods at MOI 0.1

(intracellular signaling analysis) or 0.001 (supernatant analysis). Culture supernatants were

subjected to measurement of SIV capsid p27 concentrations by ELISA. Harvested cells were

fixed and permeabilized with Cytofix/Cytoperm kit, washed twice and subjected to

immunostaining. The following antibodies were used; anti-SIV_{mac251} Gag p27 mAb (ABL)

manually conjugated with Alexa 647 (Life Technologies), anti-SIV_{mac251} Nef mAb (clone 17,

epitope peptide corresponding to SIV_{mac239} Nef 71-80 not including the residue G63: Thermo

Scientific/Pierce) manually conjugated with Alexa 488 or Alexa 647, anti-human

CD3-APC-Cy7 (SP34-2, BD Pharmingen), anti-human CD4-PerCP (L200, BD Pharmingen),

anti-human HLA-ABC-FITC (G46-2.6, BD Pharmingen), anti-human CXCR4-APC (12G5,

BioLegend), BST2-Alexa 647 (RS38E, BioLegend), Alexa 488-conjugated anti-phospho-Akt

(Ser473) (D9E, CST) and Alexa 488-conjugated anti-phospho-Akt (Thr308) (C31E5E, CST).

Cells acquired by FACS Canto II were analyzed by FACS Diva and FlowJo. Approximately 5 x

10^4 cells were gated for each test.

Ferritin ELISA

Plasma ferritin levels in NAb inducers and control animals were analyzed by monkey ferritin

sandwich ELISA kit (LS Bio) according to the manufacturer's instructions.

PI3K stimulation assay

5×10^4 HSC-F cells were infected with wild-type or Nef G63E mutant SIV_{mac239} at MOI 0.2 and

cultured for 1 day in medium supplemented with 10% (normal), 1% (1/10 starvation) or 0.1%

(1/100 starvation) FBS. For ligand stimulation (**Figure 4-figure supplement 1C**), cells at the

end of 1-day culture were pulsed for 20 minutes with 40 ng/ml of recombinant human IFN-γ

(Gibco/Thermo Fisher), 100 IU/ml of recombinant human IL-2 (Roche Diagnostics) or 10

µg/ml of SIV_{mac251} Env gp130 (ImmunoDx). Cells were intracellularly stained using

Cytofix/Cytoperm kit with anti-SIV_{mac251} Nef mAb or anti-SIV_{mac251} p27 mAb manually

conjugated to Alexa 647 and PE-conjugated anti-human phospho-Akt (Ser473) (D9E, CST)

or Alexa 488-conjugated anti-human phospho-Akt (Ser473). Cells acquired by FACS Canto II

were analyzed by FACS Diva and FlowJo. Approximately 7×10^4 HSC-F cells were gated per

test.

Transcriptome analysis

Total RNAs were extracted using RNeasy Plus Mini kit (Qiagen) from 2×10^6 HSC-F cells 1

day after infection with wild-type or Nef G63E mutant SIV_{mac239} at MOI 5. Negative control

RNA samples were extracted from 2×10^6 uninfected HSC-F cells after culture with the same

condition. Three sets of experiments were performed. Total RNA samples were subjected to a

quality control (QC) analysis using an Agilent 2100 Bioanalyzer. The obtained amounts of

total RNAs were: 12.37 ± 0.39 (uninfected), 8.80 ± 0.44 (wild-type) and 9.02 ± 0.17 (Nef

G63E) µg ($P = 0.66$ for wild-type versus Nef G63E by unpaired t test). In all samples, two

bands of 18S and 28S rRNA were confirmed and the RNA integrity number (RIN) was 10. 500

ng of total RNA samples were processed with GeneChip WT Plus reagent

(Affymetrix/Thermo Fisher Scientific) to produce 150 µl of fragmented and labeled cDNA

samples. These were incubated with a Human Gene 2.0 ST Array (Affymetrix/Thermo Fisher

Scientific) for 16 hours, 60 rpm at 45°C using GeneChip hybridization oven 645

(Affymetrix/Thermo Fisher Scientific). Results were scanned with GeneChip Scanner 3000

7G (Affymetrix/Thermo Fisher Scientific) and processed with Affymetrix Expression Console

Software (Affymetrix/Thermo Fisher Scientific) according to the manufacturer's instructions.

Expression values were normalized by the RMA method. Genes above mean background

expression within the cognate sample and showing significant difference between wild-type

and Nef G63E SIV ($P < 0.05$ via unpaired t test for $\log_2$-transformed values, 768 candidates)

were determined. Akt-related genes exhibiting a change of approximately 10% or more were

representatively extracted and manually aligned by the authors.

Peripheral CD4+ T-cell surface staining

Cryopreserved/thawed PBMCs were stained for 30 minutes at 4°C with the following

reagents or conjugated anti-human mAbs: Live/Dead Aqua (Life Technologies),

anti-CD4-PerCP (L200, BD Pharmingen), anti-CD8-APC-Cy7 (RPA-T8, BD Pharmingen),

anti-CD3-Alexa 700 (SP34-2, BD Pharmingen), anti-CD95-PE-Cy7 (DX2, eBioscience),

anti-CXCR5-PE (87.1, eBioscience), anti-PD-1-Brilliant Violet 421 (EH12.2H7, BioLegend)

and anti-CXCR3-Alexa 488 (G025H7, BioLegend). Cells acquired by FACS LSRII Fortessa

(BD) were analyzed by FACS Diva and FlowJo. Approximately 1.5×10^5 PBMCs were gated

per test.

Proximity ligation assay (PLA)

A flow cytometry-based arrangement of proximity ligation assay (PLA) (Avin et al., 2017) was

performed to quantitatively assess Nef binding to candidate interacting molecules. 1×10^5

HSC-F cells were infected with SIV$_{mac239}$ at MOI 0.05 in U-bottomed plates and permeated

with Cytofix/Perm kit (BD Biosciences). After two washes, they were resuspended in 0.5%

BSA/PBS for prevention of experimental procedure-related loss and stained with mouse

anti-SIV$_{mac251}$ Nef mAb (clone 17, Thermo Scientific/Pierce) or mouse IgG1 isotype control

mAb (P3.6.2.8.1, Abcam) in combination with either of the following rabbit antibodies:

anti-GβL (86B8, CST), anti-mTOR (7C10, CST), polyclonal anti-human PI3K p85 (Merck

Millipore), anti-PI3 Kinase p110α (C73F8, CST) or rabbit IgG1 isotype control mAb (DA1E,

CST). Antibody-stained cells were subsequently probed with Duolink In Situ PLA Probe

anti-mouse PLUS and anti-mouse MINUS probes (Sigma/Merck). Next, they were detected

for intermolecular binding using Duolink flow PLA detection kit (Green) (Sigma/Merck) with a

reaction time of 100 minutes for post-mouse/rabbit probe linking amplification. Finally, these

PLA-subjected cells were additionally stained with anti-SIV$_{mac251}$ Gag p27 mAb manually

conjugated with Alexa 647 and anti-CD4-PerCP for 30 minutes for 20 minutes at 4°C. Cells

acquired by FACS Canto II were analyzed by FACS Diva and FlowJo. Approximately 1 x 10^5

cells were gated per test. Binding index (Y axis) was calculated by deriving the deviation from

the summation of: {baseline P: (MFI of background reaction with mouse isotype control/rabbit

isotype control) + anti-Nef antibody-derived background Q: [(MFI of reaction with mouse

anti-Nef/rabbit isotype control) - (MFI of background reaction with mouse isotype

control/rabbit isotype control)] + anti-binding partner antibody-derived background R:[(MFI of

reaction with mouse isotype control/rabbit anti-binding partner molecule) - (MFI of

background reaction with mouse isotype control/rabbit isotype control)]}.

B-cell Nef invasion *in vitro* assay

For Nef invasion inhibition assay, 5 x 10^4 HSC-F cells were infected at MOI 1 for 48 hours.

These were then additionally cocultured with 7.5 x 10^4 Ramos B cells for 12 hours. Ramos B

cells were double-stained with Alexa 488-cojugated Nef and anti-CD19-PC-5.5 (J3-119,

Beckman Coulter), and interacting infected cell-invaded B cell doublet images acquired with

ImageStream X MKII (Amnis/Merck Millipore) were analyzed by IDEAS 6.3 (Amnis/Merck

Millipore/Luminex).

Quantitative *in vivo* imaging flow cytometry

Cryopreserved/thawed lymph node cells (LNCs) from several persistently SIV-infected

macaques used in previous experiments (Nakane et al., 2013; Takahashi et al., 2013;

Iwamoto et al., 2014) were seeded in V-bottomed 96-well plates (Nunc) and blocked with 25

μg/ml of anti-human CD4 (clone L200: BD) in 100 μl volume for 15 minutes at 4°C. After three

washes, they were stained with 10 μg/ml of recombinant SIV_{mac239} Env gp140-biotin (Immune

Technology) for 30 minutes at 4°C. Cells were then stained with anti-CD19-PC5.5 (J3-119,

Beckman Coulter), Live/Dead Aqua (Invitrogen) and streptavidin-APC-Cy7 (BioLegend) for

30 minutes at 4°C. After two washes, cells were processed with Cytofix/Cytoperm kit (BD) for

20 minutes at 4°C, washed twice, and next intracellularly stained with PE-conjugated

anti-human phospho-Akt (Ser473) (D9E, CST) and anti-SIV_{mac251} Nef mAb (clone 17, Thermo

Scientific/Pierce) manually conjugated with Alexa 488 or mouse IgG1 isotype control mAb

(P3.6.2.8.1, Abcam) for 30 minutes at 4°C. After final two washes, cells were suspended in

0.8% PFA/PBS. All centrifugations for washing (1,200 x g, 2 minutes) were performed at 4°C.

Cells were subjected to image acquisition with Image Stream X MKII imaging flow cytometer

(Amnis/Merck Millipore/Luminex) and analyzed with IDEAS 6.3 software (Amnis/Merck

Millipore/Luminex). Approximately 5 x 10^5 LNCs were acquired for analyses (**Figure 5**). A

custom-implemented linear discriminant analysis-based machine learning module (Luminex)

was utilized for verifying candidate Nef staining signal-related secondary parameters (**Figure

5** and **Figure 5-figure supplement 1A**) for their efficacy of target population separation.

Peripheral SIV Env-specific B-cell responses

Peripheral SIV Env gp140-specific memory B cells (B_{mem}) and plasmablasts (PBs) were

measured by flow cytometry with procedures modified from previous reports and our

experience (Silveira et al., 2015; Hau et al., 2022). First, cryopreserved/thawed PBMCs

seeded in V-bottomed 96-well plates (Nunc) were blocked with 25 μg/ml of anti-human CD4

(clone L200: BD) in 100 μl volume for 15 minutes at 4°C. This concentration attains blockade

of promiscuous SIV Env binding to CD4 comparable to levels by magnetic depletion of CD3[+]

T cells (data not shown). After three washes, they were next stained with 10 μg/ml of

recombinant SIV_{mac239} Env gp140-biotin (Immune Technology) for 30 minutes at 4°C. They

were subsequently stained with the following anti-human antibodies and fluorochromes for 30

minutes at 4°C with combinations as follows; PB/B_{mem}: anti-CD3-APC-Cy7 (SP34-2, BD

Pharmingen), anti-CD8-APC-Cy7 (RPA-T8, BD Pharmingen), anti-CD14-APC-Cy7 (M5E2,

BioLegend), anti-CD16-APC-Cy7 (3G8, BioLegend), streptavidin-Brilliant Violet 421

(BioLegend), anti-CD19-PC5.5 (J3-119, Beckman Coulter) and Live/Dead Aqua (Invitrogen);

PB: anti-CD10-APC-Cy7 (HI10a, BioLegend), anti-CD11c-PE-Cy7 (3.9, BioLegend),

anti-CD123-PE-Cy7 (6H6, BioLegend) and anti-HLA-DR-PE-Texas Red (TU36, Invitrogen);

B_{mem}: anti-CD27-PE/Dazzle 594 (M-T271, BioLegend), anti-CD10-PE-Cy7 (HI10a,

BioLegend), anti-IgD-FITC (DaKo), anti-CD38-Alexa 647 (AT1, Santa Cruz Biotechnology),

anti-IgG-Alexa 700 (G18-145, BD Pharmingen) and anti-CD138-PE (DL-101, eBioscience).

After two washes, B_{mem} samples were suspended in 0.8% PFA/PBS. For PB staining,

surface-stained cells were further processed with Cytofix/Cytoperm kit (BD) for 20 minutes at

4°C, washed twice, and next intracellularly stained with 10 μg/ml of recombinant SIV_{mac239}

Env gp140-biotin (Immune Technology) for 30 minutes at 4°C. After three washes, they were

next stained with anti-human IgG-PE (G18-145, BD), anti-human/mouse IRF4-eFluor 660

(3E4, eBioscience), anti-human Ki67-Alexa 700 (B56, BD Pharmingen) and

streptavidin-Alexa 488 (BioLegend) for 30 minutes at 4°C. After final two washes, cells were

suspended in 0.8% PFA/PBS. Cells acquired with FACS LSRII Fortessa were analyzed with

FACS Diva and FlowJo. Approximately 1.5×10^5 PBMCs were acquired for B_{mem} and 4×10^5

PBMCs were acquired for PB analyses (**Figure 6**). All centrifugations for washing (1,200 x g,

2 minutes) were performed at 4°C. Env-specific PB frequencies were 0 cells/million PBMCs

for pre-challenge samples in all animals examined, ruling out background confounding for

high-sensitivity quantitation of the population.

Statistical Analysis

Analyses were performed via Prism 8 (GraphPad Software). $P < 0.05$ were considered

significant in two-tailed unpaired t tests, paired t tests, Mann Whitney U tests, Fisher's exact

tests, Wilcoxon signed-rank tests and two-way ANOVA with Sidak's post-hoc multiple

comparison tests. For analysis of Nef-invaded B-cell pAkt levels, median fluorescence

intensities (mFIs) were analyzed due to relatively small numbers of Nef[+] B cells. Analyses

involving pAkt Ser473 levels all derived comparable results between mFI and MFI for wild

type versus Nef G63E virus. Machine learning-based rating of Nef signal parameters was

verified with IDEAS 6.3 Machine Learning Module (Luminex).

References

Akari, H., Mori, K., Terao, K., Otani, I., Fukasawa, M., Mukai, R., and Yoshikawa, Y. (1996). In

vitro immortalization of Old World monkey T lymphocytes with Herpesvirus saimiri: its

susceptibility to infection with simian immunodeficiency viruses. *Virology* 218, 382–388.

Al Qureshah, F., Sagadiev, S., Thouvenel, C.D., Liu, S., Hua, Z., Hou, B., Acharya, M., James,

R.G., and Rawlings, D.J. (2021). Activated PI3Kδ signals compromise plasma cell survival

via limiting autophagy and increasing ER stress. *J Exp Med.* 218, e20211035.

Angulo, I., Vadas, O., Garçon, F., Banham-Hall, E., Plagnol, V., Leahy, T.R., Baxendale, H.,

Coulter, T., Curtis, J., Wu, C., et al. (2013). Phosphoinositide 3-kinase δ gene mutation

predisposes to respiratory infection and airway damage. *Science* 342, 866–871.

Avery DT, Kane A, Nguyen T, Lau A, Nguyen A, Lenthall H, Payne K, Shi W, Brigden H,

French E, et al. (2018). Germline-activating mutations in PIK3CD compromise B cell

development and function. *J Exp Med* 215, 2073–2095.

Avin, A., Levy, M., Porat, Z., and Abramson, J. (2017) Quantitative analysis of protein-protein

interactions and post-translational modifications in rare immune populations. *Nat. Commun.* 8,

1524.

Battegay, M., Moskophidis, D., Waldner, H., Bründler, M.A., Fung-Leung, W.P., Mak, T.W.,

Hengartner, H., and Zinkernagel, R.M. (1993). Impairment and delay of neutralizing antiviral

antibody responses by virus-specific cytotoxic T cells. *J. Immunol.* 151, 5408–5415.

Baumjohann, D., Preite, S., Reboldi, A., Ronchi, F., Ansel, K.M., Lanzavecchia, A., and

Sallusto, F. (2013). Persistent antigen and germinal center B cells sustain T follicular helper

cell responses and phenotype. *Immunity* 38, 596–605.

Bradley, T., Peppa, D., Pedroza-Pacheco, I., Li, D., Cain, D.W., Henao, R., Venkat, V., Hora,

B., Chen, Y., Vandergrift, N.A., et al. (2018). RAB11FIP5 Expression and Altered Natural Killer

Cell Function Are Associated with Induction of HIV Broadly Neutralizing Antibody Responses.

Cell 175, 387–399.e17.

Burns, D.P., Collignon, C., and Desrosiers, R.C. (1993). Simian immunodeficiency virus

mutants resistant to serum neutralization arise during persistent infection of rhesus monkeys.

J. Virol. 67, 4104–4113.

Cubas, R.A., Mudd, J.C., Savoye, A.L., Perreau, M., van Grevenynghe, J., Metcalf, T.,

Connick, E., Meditz, A., Freeman, G.J., Abesada-Terk, G. Jr., et al. (2013). Inadequate T

follicular cell help impairs B cell immunity during HIV infection. *Nat. Med.* 19, 494–499.

Evans, D.T., O'Connor, D.H., Jing, P., Dzuris, J.L., Sidney, J., da Silva, J., Allen, T.M., Horton,

H., Venham, J.E., Rudersdorf, R.A., et al. (1999). Virus-specific cytotoxic T-lymphocyte

responses select for amino-acid variation in simian immunodeficiency virus Env and Nef. *Nat.*

Med. 5, 1270–1276.

Fallet, B., Narr, K., Ertuna, Y.I., Remy, M., Sommerstein, R., Cornille, K., Kreutzfeldt, M.,

Page, N., Zimmer, G., Geier, F., et al. (2016). Interferon-driven deletion of antiviral B cells at

the onset of chronic infection. *Sci. Immunol.* 1, pii: eaah6817.

François, F., and Klotman, M.E. (2003). Phosphatidylinositol 3-kinase regulates human

immunodeficiency virus type 1 replication following viral entry in primary CD4+ T lymphocytes

and macrophages. *J. Virol.* 77, 2539–2549.

Fruman, D.A., Snapper, S.B., Yballe, C.M., Davidson, L., Yu, J.Y., Alt, F.W., and Cantley, L.C.

(1999). Impaired B cell development and proliferation in absence of phosphoinositide

3-kinase p85alpha. *Science* 283, 393–397.

Gauduin, M.C., Yu, Y., Barabasz, A., Carville, A., Piatak, M., Lifson, J.D., Desrosiers, R.C.,

and Johnson, R.P. (2006). Induction of a virus-specific effector-memory CD4+ T cell response

by attenuated SIV infection. *J. Exp. Med.* 203, 2661–2672.

Geyer, M., Fackler, O.T., and Peterlin, B.M. (2001). Structure--function relationships in HIV-1

Nef. *EMBO Rep.* 2, 580–585.

Gitlin, A.D., Shulman, Z. and Nussenzweig, M.C. (2014). Clonal selection in the germinal

centre by regulated proliferation and hypermutation. *Nature* 509, 637–640.

Gray, E.S., Madiga, M.C., Hermanus, T., Moore, P.L., Wibmer, C.K., Tumba, N.L., Werner, L.,

Mlisana, K., Sibeko, S., Williamson, C., et al. (2011). The neutralization breadth of HIV-1

develops incrementally over four years and is associated with CD4+ T cell decline and high

viral load during acute infection. *J. Virol.* 85, 4828–4840.

Haigwood, N.L., Watson, A., Sutton, W.F., McClure, J., Lewis, A., Ranchalis, J., Travis, B.,

Voss, G., Letvin, N.L., Hu, S.L., et al. (1996). Passive immune globulin therapy in the

SIV/macaque model: early intervention can alter disease profile. *Immunol. Lett.* 51, 107–114.

Haralick, R.M., Shanmugam, K., and Dinstein, I. (1973) Textural features for image

classification. *IEEE Transactions on Systems, Man, and Cybernetics* SMC-3, 610–621.

Hashimoto, M., Bhuyan, F., Hiyoshi, M., Noyori, O., Nasser, H., Miyazaki, M., Saito, T.,

Kondoh, Y., Osada, H., Kimura, S., et al. (2016). Potential role of the formation of tunneling

nanotubes in HIV-1 spread in macrophages. *J. Immunol.* 196, 1832–1841.

Hau, T.T.T., Nishizawa, M., Harada, S., Phan, M.H., Kanno, Y., Nomura, T., Matsuoka, S.,

Kawana-Tachikawa, A., Hall, W.W., Matano, T., et al. (2022). Association of envelope-specific

B-cell differentiation and viral selective pressure signatures in HIV-1 CRF01_AE infection.

AIDS 36, 1629–1641.

Havenar-Daughton, C., Carnathan, D.G., Torrents de la Peña, A., Pauthner, M., Briney, B.,

Reiss, S.M., Wood, J.S., Kaushik, K., van Gils, M.J., Rosales, S.L., et al. (2016). Direct

Probing of Germinal Center Responses Reveals Immunological Features and Bottlenecks for

Neutralizing Antibody Responses to HIV Env Trimer. *Cell Rep.* 17, 2195–2209.

Hoof, I., Peters, B., Sidney, J., Pedersen, L.E., Sette, A., Lund, O., Buus, S., and Nielsen, M..

(2009). NetMHCpan, a method for MHC class I binding prediction beyond humans.

Immunogenetics 61, 1–13.

Hunziker, L., Recher, M., Macpherson, A.J., Ciurea, A., Freigang, S., Hengartner, H., and

Zinkernagel, R.M. (2003). Hypergammaglobulinemia and autoantibody induction

mechanisms in viral infections. *Nat. Immunol.* 4, 343–349.

Iseda, S., Takahashi, N., Poplimont, H., Nomura, T., Seki, S., Nakane, T., Nakamura, M., Shi,

S., Ishii, H., Furukawa, S., et al. (2016). Biphasic CD8+ T-cell defense in simian

immunodeficiency virus control by acute-phase passive neutralizing antibody immunization. *J.*

Virol. 90, 6276–6290.

Ishii, H., Kawada, M., Tsukamoto, T., Yamamoto, H., Matsuoka, S., Shiino, T., Takeda, A.,

Inoue, M., Iida, A., Hara, H., et al. (2012). Impact of vaccination on cytotoxic T lymphocyte

immunodominance and cooperation against simian immunodeficiency virus replication in

rhesus macaques. *J. Virol.* 86, 738–745.

Iwamoto, N., Takahashi, N., Seki, S., Nomura, T., Yamamoto, H., Inoue, M., Shu, T., Naruse,

T.K., Kimura, A., and Matano, T. (2014). Control of simian immunodeficiency virus replication

by vaccine-induced Gag- and Vif-specific CD8$^+$ T cells. *J. Virol.* 88, 425–433.

Jia, B., Serra-Moreno, R., Neidermyer, W., Rahmberg, A., Mackey, J., Fofana, I.B., Johnson,

W.E., Westmoreland, S., and Evans, D.T. (2009). Species-specific activity of SIV Nef and

HIV-1 Vpu in overcoming restriction by tetherin/BST2. *PLoS Pathog.* 5, e1000429.

Joas, S., Sauermann, U., Roshani, B., Klippert, A., Daskalaki, M., Mätz-Rensing, K.,

Stolte-Leeb, N., Heigele, A., Tharp, G.K., Gupta, P.M., et al. (2020). Nef-Mediated CD3-TCR

Downmodulation Dampens Acute Inflammation and Promotes SIV Immune Evasion. *Cell*

Rep. 30, 2261–2274.e7.

Junt, T., Moseman, E.A., Iannacone, M., Massberg, S., Lang, P.A., Boes, M., Fink, K.,

Henrickson, S.E., Shayakhmetov, D.M., Di Paolo, N.C., et al. (2007). Subcapsular sinus

macrophages in lymph nodes clear lymph-borne viruses and present them to antiviral B cells.

Nature 450, 110–114.

Kawada, M., Tsukamoto, T., Yamamoto, H., Takeda, A., Igarashi, H., Watkins, D.I., and

Matano, T. (2007). Long-term control of simian immunodeficiency virus replication with

central memory CD4+ T-cell preservation after nonsterile protection by a cytotoxic

T-lymphocyte-based vaccine. *J. Virol.* 81, 5202–5211.

Kennedy, S.G., Wagner, A.J., Conzen, S.D., Jordán, J., Bellacosa, A., Tsichlis, P.N., and Hay,

N. (1997). The PI 3-kinase/Akt signaling pathway delivers an anti-apoptotic signal. *Genes*

Dev. 11, 701–713.

Kestler, H.W. 3rd, Ringler, D.J., Mori, K., Panicali, D.L., Sehgal, P.K., Daniel, M.D., and

Desrosiers, R.C. (1991). Importance of the nef gene for maintenance of high virus loads and

for development of AIDS. *Cell* 65, 651–662.

Kim, D.H., Sarbassov, D.D., Ali, S.M., Latek, R.R., Guntur, K.V., Erdjument-Bromage, H.,

Tempst, P., and Sabatini, D.M. (2003). GbetaL, a positive regulator of the rapamycin-sensitive

pathway required for the nutrient-sensitive interaction between raptor and mTOR. *Mol. Cell*

11, 895–904.

Kirchhoff, F., Greenough, T.C., Brettler, D.B., Sullivan, L.L. and Desrosiers, R.C. (1995).

Absence of intact nef sequences in a long-term, nonprogressing survivor of HIV-1 infection. N.

Engl. J. Med. *332*, 228–232.

Kumar, A., Abbas, W., Colin, L., Khan, K.A., Bouchat, S., Varin, A., Larbi, A., Gatot, J.S.,

Kabeya, K., Vanhulle, C., et al. (2016). Tuning of AKT-pathway by Nef and its blockade by

protease inhibitors results in limited recovery in latently HIV infected T-cell line. *Sci. Rep.* 6,

24090.

Kumar, V., Scandella, E., Danuser, R., Onder, L., Nitschké, M., Fukui, Y., Halin, C., Ludewig,

B., and Stein, J.V. (2010). Global lymphoid tissue remodeling during a viral infection is

orchestrated by a B cell-lymphotoxin-dependent pathway. *Blood* 115, 4725–4733.

Leuchowius, K.J., Weibrecht, I., Landegren, U., Gedda, L., and Söderberg, O. (2009). Flow

cytometric in situ proximity ligation analyses of protein interactions and post-translational

modification of the epidermal growth factor receptor family. *Cytometry A* 75, 833–839.

Levesque, M.C., Moody, M.A., Hwang, K.K., Marshall, D.J., Whitesides, J.F., Amos, J.D.,

Gurley, T.C., Allgood, S., Haynes, B.B., Vandergrift, N.A., et al. (2009). Polyclonal B cell

differentiation and loss of gastrointestinal tract germinal centers in the earliest stages of HIV-1

infection. *PLoS Med.* 6, e1000107.

Lichterfeld, M., Kavanagh, D.G., Williams, K.L., Moza, B., Mui, S.K., Miura, T., Sivamurthy, R.,

Allgaier, R., Pereyra, F., Trocha, A., et al. (2007). A viral CTL escape mutation leading to

immunoglobulin-like transcript 4-mediated functional inhibition of myelomonocytic cells. *J.*

Exp. Med. 204, 2813–2824.

Limon, J.J., So, L., Jellbauer, S., Chiu, H., Corado, J., Sykes, S.M., Raffatellu, M., and

Fruman, D.A. (2014). mTOR kinase inhibitors promote antibody class switching via mTORC2

inhibition. *Proc. Natl. Acad. Sci. USA* 111, E5076–5085.

Lindqvist, M., van Lunzen, J., Soghoian, D.Z., Kuhl, B.D., Ranasinghe, S., Kranias, G.,

Flanders, M.D., Cutler, S., Yudanin, N., Muller, M.I., et al. (2012). Expansion of HIV-specific T

follicular helper cells in chronic HIV infection. *J. Clin. Invest.* 122, 3271–3280.

Locci, M., Havenar-Daughton, C., Landais, E., Wu, J., Kroenke, M.A., Arlehamn, C.L., Su,

L.F., Cubas, R., Davis, M.M., Sette, A., et al. (2013). Human circulating

PD-1+CXCR3-CXCR5+ memory Tfh cells are highly functional and correlate with broadly

neutralizing HIV antibody responses. *Immunity* 39, 758–769.

Lucas, C.L., Kuehn, H.S., Zhao, F., Niemela, J.E., Deenick, E.K., Palendira, U., Avery, D.T.,

Moens, L., Cannons, J.L., Biancalana, M., et al. (2014). Dominant-activating germline

mutations in the gene encoding the PI(3)K catalytic subunit p110δ result in T cell senescence

and human immunodeficiency. *Nat. Immunol.* 15*,* 88–97.

1003

1004   Luo, W., Hawse, W., Conter, L., Trivedi, N., Weisel, F., Wikenheiser, D., Cattley, R.T., and

1005   Shlomchik, M.J. (2019). The AKT kinase signaling network is rewired by PTEN to control

1006   proximal BCR signaling in germinal center B cells. *Nat. Immunol.* 20, 736–746.

1007

1008   Marzec, M., Liu, X., Kasprzycka, M., Witkiewicz, A., Raghunath, P.N., El-Salem, M.,

1009   Robertson, E., Odum, N., and Wasik, M.A. (2008). IL-2- and IL-15-induced activation of the

1010   rapamycin-sensitive mTORC1 pathway in malignant CD4+ T lymphocytes. *Blood* 111,

1011   2181–2189.

1012

1013   Mason, R.D., Welles, H.C., Adams, C., Chakrabarti, B.K., Gorman, J., Zhou, T., Nguyen, R.,

1014   O'Dell, S., Lusvarghi, S., Bewley, C.A., et al. (2016). Targeted Isolation of Antibodies Directed

1015   against Major Sites of SIV Env Vulnerability. *PLoS Pathog.* 12, e1005537.

1016

1017   Matano, T., Kobayashi, M., Igarashi, H., Takeda, A., Nakamura, H., Kano, M., Sugimoto, C.,

1018   Mori, K., Iida, A., Hirata, T., et al. (2004). Cytotoxic T lymphocyte-based control of simian

1019   immunodeficiency virus replication in a preclinical AIDS vaccine trial. *J. Exp. Med.* 199,

1020   1709–1718.

1021

1022   Mattapallil, J.J., Douek, D.C., Hill, B., Nishimura, Y., Martin, M., and Roederer, M. (2005).

1023   Massive infection and loss of memory CD4+ T cells in multiple tissues during acute SIV

1024   infection. *Nature* 434, 1093–1097.

1025

1026 Mikell, I., Sather, D.N., Kalams, S.A., Altfeld, M., Alter, G., and Stamatatos, L. (2011).

1027 Characteristics of the earliest cross-neutralizing antibody response to HIV-1. *PLoS Pathog.* 7,

1028 e1001251.

1029

1030 Moir, S., Malaspina, A., Ogwaro, K.M., Donoghue, E.T., Hallahan, C.W., Ehler, L.A., Liu, S.,

1031 Adelsberger, J., Lapointe, R., Hwu, P., et al. (2001). HIV-1 induces phenotypic and functional

1032 perturbations of B cells in chronically infected individuals. *Proc. Natl. Acad. Sci. USA* 98,

1033 10362–10367.

1034

1035 Moody, M.A., Pedroza-Pacheco, I., Vandergrift, N.A., Chui, C., Lloyd, K.E., Parks, R.,

1036 Soderberg, K.A., Ogbe, A.T., Cohen, M.S., Liao, H.X., Gao, F., et al. (2016). Immune

1037 perturbations in HIV-1-infected individuals who make broadly neutralizing antibodies. *Sci.*

1038 *Immunol.* 1, aag0851.

1039

1040 Moore, P.L., Gray, E.S., Wibmer, C.K., Bhiman, J.N., Nonyane, M., Sheward, D.J., Hermanus,

1041 T., Bajimaya, S., Tumba, N.L., Abrahams, M.R., et al. (2012). Evolution of an HIV

1042 glycan-dependent broadly neutralizing antibody epitope through immune escape. *Nat. Med.*

1043 18, 1688–1692.

1044

1045 Nakane, T., Nomura, T., Shi, S., Nakamura, M., Naruse, T.K., Kimura, A., Matano, T., and

1046 Yamamoto, H. (2013). Limited impact of passive non-neutralizing antibody immunization in

1047 acute SIV infection on viremia control in rhesus macaques. *PLoS ONE* 8, e73453.

1048

1049 Naruse, T.K., Chen, Z., Yanagida, R., Yamashita, T., Saito, Y., Mori, K., Akari, H., Yasutomi, Y.,

1050 Miyazawa, M., Matano, T., et al. (2010). Diversity of MHC class I genes in Burmese-origin

1051 rhesus macaques. *Immunogenetics* 62, 601–611.

1052

1053   Ng, C.T., Jaworski, J.P., Jayaraman, P., Sutton, W.F., Delio, P., Kuller, L., Anderson, D.,

1054   Landucci, G., Richardson, B.A., Burton, D.R., et al. (2010). Passive neutralizing antibody

1055   controls SHIV viremia and enhances B cell responses in infant macaques. *Nat. Med.* 16,

1056   1117–1119.

1057

1058   Nguyen, H., Ramana, C.V., Bayes, J., and Stark, G.R. (2001). Roles of phosphatidylinositol

1059   3-kinase in interferon-gamma-dependent phosphorylation of STAT1 on serine 727 and

1060   activation of gene expression. *J. Biol. Chem.* 276, 33361–33368.

1061

1062   Nishimura, Y., Gautam, R., Chun, T.W., Sadjadpour, R., Foulds, K.E., Shingai, M., Klein, F.,

1063   Gazumyan, A., Golijanin, J., Donaldson, M., et al. (2017). Early antibody therapy can induce

1064   long-lasting immunity to SHIV. *Nature* 543, 559–563.

1065

1066   Nomura, T., Yamamoto, H., Shiino, T., Takahashi, N., Nakane, T., Iwamoto, N., Ishii, H.,

1067   Tsukamoto, T., Kawada, M., Matsuoka, S., et al. (2012). Association of major

1068   histocompatibility complex class I haplotypes with disease progression after simian

1069   immunodeficiency virus challenge in Burmese rhesus macaques. *J. Virol.* 86, 6481–6490.

1070

1071   Nutt, S.L., Hodgkin, P.D., Tarlinton, D.M., and Corcoran, L.M. (2015). The generation of

1072   antibody-secreting plasma cells. *Nat. Rev. Immunol.* 15, 160–171.

1073

1074   Omori, S.A., Cato, M.H., Anzelon-Mills, A., Puri, K.D., Shapiro-Shelef, M., Calame, K., and

1075   Rickert, R.C. (2006). Regulation of class-switch recombination and plasma cell differentiation

1076   by phosphatidylinositol 3-kinase signaling. *Immunity* 25, 545–557.

1077

1078   Perfetto, S.P., Chattopadhyay, P.K., Lamoreaux, L., Nguyen, R., Ambrozak, D., Koup, R.A.,

1079   and Roederer, M. (2006). Amine reactive dyes: an effective tool to discriminate live and dead

1080   cells in polychromatic flow cytometry. *J. Immunol. Methods* 313, 199–208.

1081

1082 Petrovas, C., Yamamoto, T., Gerner, M.Y., Boswell, K.L., Wloka, K., Smith, E.C., Ambrozak,

1083 D.R., Sandler, N.G., Timmer, K.J., Sun, X., et al. (2012). CD4 T follicular helper cell dynamics

1084 during SIV infection. *J. Clin. Invest.* 122, 3281–3294.

1085

1086 Qiao, X., He, B., Chiu, A., Knowles, D.M., Chadburn, A., and Cerutti, A. (2006). Human

1087 immunodeficiency virus 1 Nef suppresses CD40-dependent immunoglobulin class switching

1088 in bystander B cells. *Nat. Immunol.* 7, 302–310.

1089

1090 Ray, J.P., Staron, M.M., Shyer, J.A., Ho, P.C., Marshall, H.D., Gray, S.M., Laidlaw, B.J., Araki,

1091 K., Ahmed, R., Kaech, S.M., et al. (2015). The interleukin-2-mTORc1 kinase axis defines the

1092 signaling, differentiation, and metabolism of T helper 1 and follicular B helper T cells.

1093 *Immunity* 43, 690–702.

1094

1095 Recher, M., Lang, K.S., Hunziker, L., Freigang, S., Eschli, B., Harris, N.L., Navarini, A., Senn,

1096 B.M., Fink, K., Lötscher, M., et al. (2004). Deliberate removal of T cell help improves

1097 virus-neutralizing antibody production. *Nat. Immunol.* 5, 934–942.

1098

1099 Recher, M., Lang, K.S., Navarini, A., Hunziker, L., Lang, P.A., Fink, K., Freigang, S., Georgiev,

1100 P., Hangartner, L., Zellweger, R., et al. (2007). Extralymphatic virus sanctuaries as a

1101 consequence of potent T-cell activation. *Nat. Med.* 13, 1316–1323.

1102

1103 Rickles, R.J., Botfield, M.C., Weng, Z., Taylor, J.A., Green, O.M., Brugge, J.S., and Zoller, M.J.

1104 (1994). Identification of Src, Fyn, Lyn, PI3K and Abl SH3 domain ligands using phage display

1105 libraries. *EMBO J.* 13, 5598–5604.

1106

1107 Sammicheli, S., Kuka, M., Di Lucia, P., de Oya, N.J., De Giovanni, M., Fioravanti, J.,

1108 Cristofani, C., Maganuco, C.G., Fallet, B., Ganzer, L., et al. (2016). Inflammatory monocytes

1109 hinder antiviral B cell responses. *Sci. Immunol.* 1, pii: eaah6789.

1110

1111 Sander, S., Chu, V.T., Yasuda, T., Franklin, A., Graf, R., Calado, D.P., Li, S., Imami, K.,

1112 Selbach, M., Di Virgilio, M., et al. (2015). PI3 kinase and FOXO1 transcription factor activity

1113 differentially control B cells in the germinal center light and dark zones. *Immunity* 43,

1114 1075–1086.

1115

1116 Sarbassov, D.D., Guertin, D.A., Ali, S.M., and Sabatini, D.M. (2005) Phosphorylation and

1117 regulation of Akt/PKB by the rictor-mTOR complex. *Science* 307, 1098–1101.

1118

1119 Schindler, M., Münch, J., Brenner, M., Stahl-Hennig, C., Skowronski, J., and Kirchhoff, F.

1120 (2004). Comprehensive analysis of nef functions selected in simian immunodeficiency

1121 virus-infected macaques. *J. Virol.* 78, 10588–10597.

1122

1123 Sette A, Sidney J, Southwood S, Moore C, Berry J, Dow C, Bradley K, Hoof I, Lewis MG,

1124 Hildebrand WH et al. (2012). A shared MHC supertype motif emerges by convergent

1125 evolution in macaques and mice, but is totally absent in human MHC molecules.

1126 *Immunogenetics* 64, 421–434.

1127

1128 Shi, S., Seki, S., Matano, T., and Yamamoto, H. (2013). IL-21-producer CD4+ T cell kinetics

1129 during primary simian immunodeficiency virus infection. *Microbes Infect.* 15, 697–707.

1130

1131 Shulman, Z., Gitlin, A.D., Weinstein, J.S., Lainez, B., Esplugues, E., Flavell, R.A., Craft, J.E.,

1132 and Nussenzweig, M.C. (2014). Dynamic signaling by T follicular helper cells during germinal

1133 center B cell selection. *Science* 345, 1058–1062.

1134

1135   Silveira, E.L., Kasturi, S.P., Kovalenkov, Y., Rasheed, A.U., Yeiser, P., Jinnah, Z.S., Legere,

1136   T.H., Pulendran, B., Villinger, F., and Wrammert, J. (2015). Vaccine-induced plasmablast

1137   responses in rhesus macaques: phenotypic characterization and a source for generating

1138   antigen-specific monoclonal antibodies. *J. Immunol. Methods* 416, 69–83.

1139

1140   Sommerstein, R., Flatz, L., Remy, M.M., Malinge, P., Magistrelli, G., Fischer, N., Sahin, M.,

1141   Bergthaler, A., Igonet, S., Ter Meulen, J., et al. (2015). Arenavirus glycan shield promotes

1142   neutralizing antibody evasion and protracted infection. *PLoS Pathog.* 11, e1005276.

1143

1144   Srinivasan, L., Sasaki, Y., Calado, D.P., Zhang, B., Paik, J.H., DePinho, R.A., Kutok, J.L.,

1145   Kearney, J.F., Otipoby, K.L., and Rajewsky, K. (2009). PI3 kinase signals BCR-dependent

1146   mature B cell survival. *Cell* 139, 573–586.

1147

1148   Stolp, B., Imle, A., Coelho, F.M., Hons, M., Gorina, R., Lyck, R., Stein, J.V., and Fackler, O.T..

1149   (2012). HIV-1 Nef interferes with T-lymphocyte circulation through confined environments in

1150   vivo. *Proc. Natl. Acad. Sci. USA* 109, 18541–18546.

1151

1152   Swingler, S., Zhou, J., Swingler, C., Dauphin, A., Greenough, T., Jolicoeur, P., and Stevenson,

1153   M. (2008). Evidence for a pathogenic determinant in HIV-1 Nef involved in B cell dysfunction

1154   in HIV/AIDS. *Cell Host Microbe* 4, 63–76.

1155

1156   Takahashi, N., Nomura, T., Takahara, Y., Yamamoto, H., Shiino, T., Takeda, A., Inoue, M., Iida,

1157   A., Hara,H., Shu, T., et al. (2013). A novel protective MHC-I haplotype not associated with

1158   dominant Gag-specific CD8[+] T-cell responses in SIV$_{mac239}$ infection of Burmese rhesus

1159   macaques. *PLoS ONE* 8, e54300.

1160

1161 Terahara, K., Ishii, H., Nomura, T., Takahashi, N., Takeda, A., Shiino, T., Tsunetsugu-Yokota,

1162 Y., and Matano, T. (2014). Vaccine-induced CD107a+ CD4+ T cells are resistant to depletion

1163 following AIDS virus infection. *J. Virol.* 88, 14232–14240.

1164

1165 Tomaras, G.D., Yates, N.L., Liu, P., Qin, L., Fouda, G.G., Chavez, L.L., Decamp, A.C., Parks,

1166 R.J., Ashley, V.C., Lucas, J.T., et al. (2008). Initial B-cell responses to transmitted human

1167 immunodeficiency virus type 1: virion-binding immunoglobulin M (IgM) and IgG antibodies

1168 followed by plasma anti-gp41 antibodies with ineffective control of initial viremia. *J. Virol.* 82,

1169 12449–12463.

1170

1171 Wang, J., Sohn, H., Sun, G., Milner, J.D. and Pierce, S.K. (2014). The autoinhibitory

1172 C-terminal SH2 domain of phospholipase C-γ2 stabilizes B cell receptor signalosome

1173 assembly. *Sci. Signal.* 7, ra89.

1174

1175 Xu, W., Santini, P.A., Sullivan, J.S., He, B., Shan, M., Ball, S.C., Dyer, W.B., Ketas, T.J.,

1176 Chadburn, A., Cohen-Gould, L., et al. (2009). HIV-1 evades virus-specific IgG2 and IgA

1177 responses by targeting systemic and intestinal B cells via long-range intercellular conduits.

1178 *Nat. Immunol.* 10, 1008–1017.

1179

1180 Yamamoto, H., Kawada, M., Takeda, A., Igarashi, H. and Matano, T. (2007). Post-infection

1181 immunodeficiency virus control by neutralizing antibodies. *PLoS ONE 2*, e540.

1182

1183 Zhang, F., Wilson, S.J., Landford, W.C., Virgen, B., Gregory, D., Johnson, M.C., Munch, J.,

1184 Kirchhoff, F., Bieniasz, P.D., and Hatziioannou, T. (2009). Nef proteins from simian

1185 immunodeficiency viruses are tetherin antagonists. *Cell Host Microbe 6*, 54–67.

1186

1187 Zotos, D., Coquet, J.M., Zhang, Y., Light, A., D'Costa, K., Kallies, A., Corcoran, L.M., Godfrey,

1188 D.I., Toellner, K.M., Smyth, M.J., et al. (2010). IL-21 regulates germinal center B cell

1189   differentiation and proliferation through a B cell-intrinsic mechanism. *J. Exp. Med.* 207,

1190   365–378.

1191

 Supplemental Information Legends

 Fig. 1-Supplement 1. Profile of SIV$_{mac239}$-NAb non-inducer subgroup cohort.

 Summary of a SIV$_{mac239}$-NAb non-inducer subgroup cohort (n = 19), composed from

 previously partially characterized, MHC class I haplotype-balanced naïve animals. Symbols

 A+, B+, E+, J+ and D+ represent possession of MHC-I haplotypes *90-120-Ia*, *90-120-Ib*,

 90-010-Ie, *90-088-Ij* and *90-010-Id*, respectively. For the titers, - represents negative; +/-

 represents neutralization in three out of four wells in a quadruplicate test. Gray shadings

 indicate euthanasia due to AIDS progression. Mamu-A and Mamu-B alleles of interest in this

 study are listed. These NAb non-inducers were previously partially reported for their plasma

 viral loads and MHC-I alleles (Nomura et al., 2012; Takahashi et al., 2013; Nakane et al.,

 2013; Iwamoto et al., 2014). R01-011 was previously partially reported for NAb titers

 measured with the same method using MT4 cells as targets (Kawada et al., 2007), which

 derived comparable results.

 Fig. 1-Supplement 2. IgG class-switched NAbs in the inducers.

 Titration of SIV$_{mac239}$-neutralizing IgGs in the NAb inducers. SIV$_{mac239}$-specific IgGs, purified

 from pools of plasma with SIV$_{mac239}$-specific NAb titers, were obtained from each animal as

 described (Yamamoto et al., 2007) and examined for their neutralizing activity by 10 TCID$_{50}$

 SIV$_{mac239}$ killing on MT4-R5 cells. NAb-negative anti-SIV IgG and control IgG were prepared

 from a representative animal (R01-011) with NAb-negative plasma and pooled plasma of

 uninfected rhesus macaques, respectively.

 Fig.1-Supplement 3. Anti-SIV binding IgG profiles in NAb inducers and non-inducers.

 (A) Temporal changes in SIV$_{mac251}$ Env gp120-specific antibody titers in the NAb non-inducers

 without rapid progression [NAb-negative non-rapid progressors, NAb(-) non-RPs] (left, n =

 11) and NAb inducers [NAb(+)] (right, n = 9). NAb non-inducers showing rapid progression

 (euthanized with AIDS onset within approximately 1 year) were not included except for

macaque R10-001 (shown in the left). Samples for non-RPs R06-001 and R10-002 were

unavailable.

(B) Unpaired t test of log-transformed anti-SIV Env Ab titers in NAb(-) non-RPs and NAb

inducers. Bars represent geometric mean absorbance within each group.

(C) Lysed SIV$_{mac}$ virion linear antigen-binding at year 1 p.i. in NAb(-) non-RPs and NAb

inducers. Plasma anti-SIV$_{mac251}$ IgGs were detected using a commercial western blotting

system against the parental strain SIV$_{mac251}$ (ZeptoMetrix). White arrowheads indicate Env

gp120-specific bands. Experiments were performed twice with comparable results.

1228

Fig. 1-Supplement 4. Env sequence variation pattern is not a major characteristic of

NAb induction.

SIV$_{mac239}$ Env sequence variations before and after NAb induction. For each animal, the

upper time point shows pre-induction and the lower time point (in purple) shows

post-induction (first or second time point with NAb titer > 1:2). Dominant residue changes are

shown as follows: gp120 (gray), variable regions V1-V5 (blue), and gp41 (blue gray).

Subdominant mutations are uncolored. Unlabeled gray and black shadings show

subdominant and dominant deletion of residues. Parental SIV$_{mac239}$ sequence is listed for

V1-V5. Residues mutated in multiple animals within V1-V5 are shown in red.

1238

Fig. 3-Supplement 1. Selection of Nef-G63E mutant SIV is linked with Nef$_{62-70}$-specific

CD8$^+$ T-cell response positivity.

(A) Nef$_{62-70}$-specific CD8+ T-cell responses around week 20 post-SIV challenge in seven

viremic NAb non-inducers possessing either of the MHC-I alleles Mamu-B*039:01,

Mamu-B*004:01 or Mamu-A1*032:02.

(B) Temporal supernatant SIV p27 concentrations after wild type (WT) or Nef-G63E mutant

SIV infection at multiplicity of infection (MOI) 0.001 on HSC-F cells. Data represent one of two

independent experiments performed in triplicate. Bars: mean ± SEM.

(C) Comparison of plasma viral loads in Nef-G63E-selecting NAb non-inducers (n = 7) versus

non-inducers at indicated time points post-SIV challenge. Bars represent geometric means.

log10-transformed values are compared by unpaired t tests.

1250

Fig. 4-Supplement 1. Decreased Akt hyperactivation properties of Nef-G63E mutant

SIV.

(A) Plasma ferritin levels (ng/ml) in examined NAb non-inducers versus Nef-G63E mutant

SIV-selecting NAb inducers around year 1 post-infection. Bars represent mean in each group.

Compared by unpaired t test.

(B) Relative levels of pAkt Ser473 (left panel) and threonine (Thr) 308 (right panel) in SIV

Nef+ HSR5.4 cells (MOI 0.1, 4 days p.i.).

(C) Relative pAkt Ser473 levels in Nef+ HSC-F cells without stimuli and transiently pulsed

with IFN-γ, IL-2 or SIV$_{mac}$ Env gp130 after infection (MOI 0.2, 1 day).

(D) Gene expression significantly differing between WT and Nef-G63E SIV infection (MOI 5, 1

day). Black bars show potentially Akt-related genes. Asterisks show dynamics concordant

with the Nef-G63E phenotype (PI3K-pAkt Ser473 downstream downregulation or inhibition

upregulation).

(E) Representative flow cytometric gating (NAb inducer R01-012, wk 30 p.i.) of peripheral

blood CXCR3$^-$CXCR5$^+$PD-1$^+$ memory follicular CD4+ T cells. Cells were gated as

CD3$^+$CD8$^-$CD4$^+$CD95$^+$CXCR5$^+$PD-1$^+$CXCR3$^-$ live singlet PBMCs. Available samples of six

NAb inducers showing peak Nef-G63E mutant selection around week 30 p.i. and thirteen

viremic NAb non-inducers were tracked.

(F) Comparison of peripheral CXCR3$^-$CXCR5$^+$PD-1$^+$ memory Tfh cell frequencies between

pre-SIV infection and around week 30 post-infection in NAb non-inducers developing disease

before week 60 (left: n = 5) and alive more than 70 weeks (middle: n = 8) and

Nef-G63E-detected NAb inducers alive more than 70 weeks (right: n = 6). Analyzed by

Wilcoxon signed-rank tests.

MFIs relative to those in uninfected controls (%) are shown. **: $P < 0.01$, ***: $P < 0.001$, n.s.:

not significant between WT and Nef-G63E mutant by unpaired t tests (B-C). Data represent

one of two (B, C) independent experiments performed in triplicate (C: unstimulated) or

quadruplicate (B, C: PI3K-pulsed). Bars: mean ± SEM.

1278

Fig. 5-Supplement 1. Machine learning-verified morphological gating of Nef-invaded B

cells *in vivo*.

(A) Representative gating strategy plus schematic of inguinal lymph node Nef$^+$CD19$^+$ B cells

of an SIV$_{mac239}$-infected rhesus macaque analyzed by ImageStreamX MK II imaging flow

cytometry (R10-012, week 160 p.i.). First-step incorporation of Live/Dead staining gates out

CD19$^+$ B cells staining positive for Nef potentially due to pre-experimental membrane

permeation (e.g., image #246896, #248952 and #71859). Second-step dual gating for

intracellular Nef signal pixel Haralick variance mean (H Variance mean, X axis), a textural

feature calculated from a gray level co-occurrence matrix (GLCM) for each image

representing adjacent pixel signal strength heterogeneity and Nef signal intensity threshold

(Y axis) gates out CD19$^+$ B cells showing potentially non-specific, strong intracellular binary

clustered staining pixels for Nef deriving a large variance (X axis), and/or cells stained overtly

strong for Nef and hence becoming void for feasible image verification (high threshold) (Y

axis), both likely originating stochastically from post-experimental rupturing procedures

including membrane permeation (e.g. image #18541). This two-step noise cancellation

results in acquisition of a Live/Dead$^-$-Nef signal Haralick variance mean^{lo}-Nef signal intensity

threshold^{lo}-Nef$^+$CD19$^+$ B-cell population (e.g. image #116318, #116250 and #132927) with a

pericellular Nef^{int-lo} staining, biologically concordant with Nef membrane anchoring,

resembling Nef$^+$ cells experimentally generated *in vitro* and showed the highest score in

IDEAS6.3 machine learning module scoring for segregation of typical void vs. valid images

(upper right). This gating was implemented for counting and analysis of Env-specific B-cell

Nef invasion *in vivo* (Figure 5).

(B) Detection of contact-dependent Nef invasion from SIV-infected HSC-F CD4$^+$ T cells to

cocultured Ramos B cells *in vitro*. Two typical images of an intercellular conduit (SSC, purple)

with Nef (green) protrusions from infected HSC-F cells to CD19$^+$ (red) Ramos B cells from

two independent experiments are shown.

1305

Fig. 6-Supplement 1. Gating strategy of SIV$_{mac239}$ Env gp140-specific B-cell responses.

Representative gating (R02-004, week 32 p.i.) of SIV Env gp140-specific memory B cells

(B$_{mem}$) and plasmablasts (PBs). Two panels for PB gating are shown in lower resolution for

visibility. B$_{mem}$ staining was performed separately and gatings are merged with PB panels

(second row, panels 2–4). For lineage-specific exclusion (first row, panel 4), CD10 was

stained independently for B$_{mem}$ (see parenthesis).

A

N-myristoylation (G2)

env

N-anchor 89 Core

104PxxP107

LTR129

SIVmac239 Nef

AP-2-binding

PPT

G63E

R06-032

R01-012

R03-015

R02-004

R01-010

R06-019

R-360

R02-007

R05-010

Status: ▷ NAb (-)   ▷ NAb (+/-)   ▷ NAb (+)

B

α-helix

N-myristoylation

N-anchor
(1–89)

G63E

PxxP motif

Core
(90–)

Acidic
cluster

D89

Core domain loop

C

Nef-G63E
selection

P = 0.0009

% detected in group

NAb-   NAb+

D

Temporal Nef-G63E mutant selection vs NAb induction

SIV 100% neutralizing titers (log₂)

Mutant SIV frequencies (%)

R06-032   R01-012   R03-015   R02-004

R01-010   R06-019   R-360

■ SIVmac239-NAb
▼ Nef G63E
◆ Nef G63R

Weeks post-SIV challenge

A

B

A

Surface downregulation

| CD3 | CD4 | MHC Class I | CXCR4 | BST-2 |

% of Max

<APC-Cy7-A>: CD3 <PerCP-A>: CD4 <FITC-A>: HLA-ABC <APC-A>: CXCR4 <APC-A>: BST2-A647

— WT
— Nef-G63E
— Uninfected

Gated on singlet CD4^{lo}Nef$^+$

B

MOI 0.1 Day 3

pAkt Ser473

% of Max

1059
896
746

Comp-FITC-A :: pAkt473

Gated on singlet p27$^+$

— WT
— Nef-G63E
— Uninfected

$P = 0.0009$

ΔMFI (+ mean of uninfected)

WT   Nef G63E

C

Accaleration w/upstream PI3K stimuli Day 3→Day 1

Serum starvation

pAkt relative MFIs in Nef$^+$ cells (%)

$P = 0.99$ $P = 0.0017$

■ WT
▲ Nef-G63E

10   1   0.1
Starvation
%serum in culture

High MOI

$P = 0.0011$

ΔMFI (+ mean of uninfected)

WT   Nef G63E

D

Intracellular PI3K-pathway binding

CD4^{lo}p27$^+$
singlet HSC-F

<APC-A>: p27-A647
68.3
<PerCP-A>: CD4

— WT
— Nef-G63E
— WT (isotype)
---- Nef G63E (isotype)

| mTORC2 | | PI3 Kinase | |
| GβL/mLST8 | mTOR | PI3Kp85 | PI3Kp110α |

% of Max

<FITC-A>: Interaction

$P = 0.039$ $P = 0.028$ $P = 0.022$ $P = 0.043$

Binding Index (ΔMFI)

WT Nef G63E   WT Nef G63E   WT Nef G63E   WT Nef G63E

A

NAb non-inducers

NAb inducers (wk 30 Nef-G63E+)

B

wk 30

at NAb induction vs yr 1

P = 0.0131

P = 0.0030

C

Temporal 2-D plot of Env gp140-specific Bmem vs PB

D_i = GC output attractor 2 (Nef-G63E+, at NAb induction)

D_n = GC output attractor 1 (NAb-, at yr 1)

NAb non-inducer cohort (_n_ = 19)

MHC-I/ID	SIVmac239 neutralization at 1:2 plasma dilution by 10TCID$_{50}$ killing assay (wks p.i.)									pVL at 6 mo	Nef G63E at 1 yr	Detected _Mamu_ alleles
	wk11–20	wk21–30	wk31–40	wk41–50	wk51–60	wk61–70	wk71–80	wk81–90	wk91–104			
_A+_												
R06-037		wk25 -		wk46 +/-			wk72 -	wk87 -	wk95 -	6.4 x 10^2	- (pVL < 400)	
R07-001		wk25 -		wk42 -				wk88 +/-	wk96 -	< 400	- (pVL < 400)	_A1*032:02_
R07-004		wk25 -		wk46 -		wk64 -	wk74 -	wk86 -		4.5 x 10^3	+	_B*039:01_
R07-009		wk25 -		wk46 -		wk64 -	wk74 -			4.5 x 10^3	-	
_B+_												
R06-001		wk25 -		wk48 -	wk60 -		wk80 -	wk90 -		8.2 x 10^3	-	
R06-039		wk25 -		wk46 -	wk56 -					8.5 x 10^6	-	_A1*032:02 , B*004:01_
_E+_												
R01-011		wk22 -	wk40 -		wk52 -		wk71 -	wk85 -	wk104 -	1.3 x 10^5	-	_B*004:01_
R05-007		wk26 -			wk55 +/-		wk71 -	wk89 -	wk97 -	1.6 x 10^4	-	
R08-007		wk21 -		wk46 -		wk69 -		wk82 +/-		8.5 x 10^6	-	
R09-011	wk18 -		wk36 -		wk52 -					8.5 x 10^6	-	
R06-038		wk25 -			wk58 +/-		wk80 +/-		wk95 -	1.1 x 10^5	-	
_J+_												
R04-014	wk12 -	wk25 -	wk40 -							1.5 x 10^5	+ (wk40)	_B*039:01_
R06-022	wk12 -	wk23 -								1.4 x 10^7	- (wk23)	_B*039:01_
R10-001	wk14 -	wk30 -	wk37 -							1.9 x 10^7	- (wk37)	_B*039:01_
_D+_												
R05-006		wk25 -		wk48 -			wk80 -	wk90 -	wk98 -	5.4 x 10^2	-	_A1*032:02 , B*004:01_
R06-034		wk25 -		wk42 -		wk61 -	wk80 +/-	wk88 +/-	wk96 -	4.5 x 10^3	-	_A1*032:02 , B*004:01_
R08-005		wk25 -		wk48 -		wk68 -			wk100 -	9.0 x 10^2	-	_A1*032:02 , B*004:01_
Others												
R10-002	wk14 -				wk59 -				wk104 -	5.7 x 10^5	-	
R10-006	wk14 -	wk30 -			wk55 -					1.0 x 10^6	-	

Group	Rhesus ID	mg/ml (IgG)	Collected/pooled (wks post-challenge)	10 TCID$_{50}$ SIV$_{mac239}$ 100% killing titer
NAb (+)	R06-032	30	wk 76	1:16
	R01-012	30	wk 73–87	1:32
	R03-015	30	wk 112	1:8
	R02-004	30	wk 76	1:4
	R01-010	30	wk 77	1:16
	R06-019	30	wk 85	1:4
	R-360	30	wk 60–86	1:4
	R02-007	30	wk 42–70	1:32
	R05-010	30	wk 71	1:64
NAb (-)	R01-011	42	wk 104	< 1:2
uninfected	pooled	36	-	< 1:2

A

NAb(-) non-RPs

R06-037 R08-007
R07-001 R06-038
R07-004 R05-006
R07-009 R06-034
R01-011 R08-005
R05-007 R10-001

NAb(+)

R02-007 R02-004
R06-019 R03-015
R01-012 R06-032
R-360
R01-010
R05-010

B

Anti-Env Ab 1 yr

P = 0.044

AU (OD 450 nm Abs)

NAb- non-RPs   NAb+

C

gp120   **V1 (113-151)**   **V2 (194-206)**   **V3 (311-344)**

L V I K K T C N K S E T D R W G L T K S I T T A S T T S T T A S A K V D W V N E T S S G A C E Q G N N T G N E S R C C W P K Y C R R P G N K T V L P V T I W S G L V F H S Q P I N D R P K Q A W C W K K

24 87 98 99 103 110   120   130 132 135 136 138 140   150 190   201   211 24 / 253 254 303   320   330   340   345 349 351

R06-032
- wk30
- wk46

R01-012
- wk30
- wk48

R03-015
- wk20
- wk53

R02-004
- wk40
- wk76

R01-010
- wk18
- wk57

R06-019
- wk24
- wk69

R-360
- wk12
- wk60

R02-007
- wk12
- wk24

R05-010
- wk12
- wk20

V4 (402-432)   **V5 (459-491)**   **gp41**

D Y K E L C K W N W F L W V E D R W T A W Q K P K E Q H K R N Y V P C N S T V T S L I A N I D W I D G N Q T N I T W S A E V A E L Y R D T G V F T S N K Q E K E R G G T I P L A V L I

352 362 375 398 400   410   417 420   425   430   460   470   480   490 511 516 518 52 / 528 594 62 / 633 689   739 74 / 749 750 751 753 756 789 79 / 800 802 836 83 / 845 863

R06-032
- wk30
- wk46

R01-012
- wk30
- wk48

R03-015
- wk20
- wk53

R02-004
- wk40
- wk76

R01-010
- wk18
- wk57

R06-019
- wk24
- wk69

R-360
- wk12
- wk60

R02-007
- wk12
- wk24

R05-010
- wk12
- wk20

A

Nef$_{62-70}$ QW9-specific responses in NAb non-inducers

Nef$_{62-70}$-specific CD8+ T-cell frequencies (/million PBMCs)

1000

R06-039 R01-011 R06-022 R10-001 R05-006 R06-034 R08-005

B

SIV p27 (ng/ml)

0 2 4 6 8 10 12 14

Days post-SIV infection on HSC-F cells

■ WT
▲ Nef-G63E

C

Plasma viral loads (vRNA copies/ml)

3 mo	6 mo	1 yr	1.5 yr
P = 0.2495	*P* = 0.5078	*P* = 0.8928	*P* = 0.6488

10^8
10^7
10^6
10^5
10^4
10^3

NAb- NAb+
Nef-G63E+

A

Intracellular pixel Nef signal Intensity threshold

Signal intensity threshold (int/hi)

Signal intensity threshold (lo)

Live/Dead⁻CD19⁺ Nef⁺ B cells

R5

IDEAS6.3 machine learning-based 2-D segregation of typical valid vs. void Live/Dead⁺Nef⁺CD19⁺ B-cell images

Good,bad

Void

Valid

Intracellular pixel Nef signal strength distribution

2-D adjacent pixel signal heterogeneity (++)
→Haralick variance mean (int-hi)

2-D adjacent pixel signal heterogeneity (+/-)
→Haralick variance mean (lo)

Live/Dead⁺CD19⁺Nef⁺ B cells (void)

Live/Dead⁻CD19⁺Nef⁺-(Intracellular Nef Haralick variance mean$^{int/hi}$or Nef Intensity threshold$^{int/hi}$) B cells (void)

Live/Dead⁻CD19⁺Nef⁺-Intracellular Nef Haralick variance mean^{lo}-Nef Intensity threshold^{lo} B cells (valid)

B

In vitro CD19⁺Nef⁺ B cells

DIC | Nef | CD19 | SSC | Merged

Key Resources Table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
gene (*Lentivirus*)	*env, nef*	GenBank	M33262.1	SIV$_{mac239}$
strain, strain background (*Lentivirus*)	SIV$_{mac239}$	PMID: 2032289		Obtained via COS-1 transfection of pBR$_{mac239}$ and repropagation on MT4-R5 cells
strain, strain background (*Escherichia coli*)	One Shot™ TOP10 Chemically Competent *E. coli*	Invitrogen (Thermo Fisher Scientific)	Cat#:C404003	For subcloning sequencing
cell line (*Homo-sapiens*)	MT4-R5	PMID: 27122584		
cell line (*Chlorocebus aethiops*)	COS-1	ATCC	CRL-1650	
cell line (*Macaca fascicularis*)	HSC-F cells (CD4+ T-cell line)	PMID: 8610465		
cell line (*Macaca mulatta*)	HSR5.4 cells (CD4+ T-cell line)	PMID: 8610465		
cell line (*Homo-sapiens*)	Ramos B cell line	ATCC	CRL-1596	
biological sample (*Macaca mulatta*)	Primary simian PBMCs from SIV-infected macaques	Japan SLC	http://www.jslc.co.jp/service/	Cryopreserved from *Macaca mulatta*
biological sample (*Macaca mulatta*)	Primary simian lymph node cells from SIV-infected	Japan SLC	http://www.jslc.co.jp/service/	Cryopreserved from *Macaca mulatta*

	macaques			
antibody	Mouse monoclonal anti-human CD4-FITC (M-T477)	BD Pharmingen	Cat#556615	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD8-PerCP (SK1)	BD Biosciences	Cat#347314	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD3-APC (SP34-2)	BD Pharmingen	Cat#557597	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human IFN-γ-PE (4S.B3)	BioLegend	Cat#502510	FACS (5 µl per test)
antibody	Mouse monoclonal anti-SIV_{mac251} Gag p27 mAb	ABL	Cat#4324	FACS (1:500 with conjugation)
antibody	Mouse monoclonal anti-SIV_{mac251} Nef mAb (clone 17)	Themo Fisher Scientific	Cat#MA1-71522	FACS, PLA and imaging cytometry (FACS and imaging cytometry, 1:500 with conjugation; PLA, 1:100)
antibody	Mouse IgG1 isotype control mAb (P3.6.2.8.1)	Themo Fisher Scientific	Cat#14-4714-85	PLA (1:100)
antibody	Rabbit monoclonal Alexa Fluor 488-conjugated anti-human phospho-Akt (Ser473) (D9E)	Cell Signaling Technology	Cat#4071	FACS (5 µl per test)

antibody	Rabbit monoclonal Alexa Fluor 488-conjugated anti-human phospho-Akt (Thr308) (C31E5E)	Cell Signaling Technology	Cat#2918	FACS (5 µl per test)
antibody	Rabbit monoclonal PE-conjugated anti-human phospho-Akt (Ser473) (D9E)	Cell Signaling Technology	Cat#5315	FACS and imaging cytometry (1:100)
antibody	Rabbit monoclonal Alexa Fluor 647-conjugated anti-human phospho-Akt (Ser473) (D9E)	Cell Signaling Technology	Cat#4075	FACS (5 µl per test)
antibody	Rabbit monoclonal Alexa Fluor 647-conjugated anti-human phospho-Akt (Thr308) (C31E5E)	Cell Signaling Technology	Cat#3375	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD19-PC-5.5 (J3-119)	Beckman Coulter	Cat#66328	FACS and imaging cytometry (5 µl per test)
antibody	Mouse monoclonal anti-human CD4, purified (clone L200)	BD Biosciences	Cat#550625	FACS (25 µg/ml per test for blockade)
antibody	Mouse monoclonal anti-human CD3-APC-Cy7 (SP34-2)	BD Pharmingen	Cat#557757	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD8-APC-Cy7 (RPA-T8)	BD Pharmingen	Cat#557760	FACS (5 µl per test)

antibody	Mouse monoclonal anti-human CD14-APC-Cy7 (M5E2)	BioLegend	Cat#301820	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD16-APC-Cy7 (3G8)	BioLegend	Cat#302018	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD10-APC-Cy7 (HI10a)	BioLegend	Cat#312212	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD11c-PE-Cy7 (3.9)	BioLegend	Cat#301608	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD123-PE-Cy7 (6H6)	BioLegend	Cat#306010	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human HLA-DR-PE-Texas Red (TU36)	Invitrogen (Thermo Fisher Scientific)	Cat# MHLDR17	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD27-PE/Dazzle 594 (M-T271)	BioLegend	Cat#356422	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD10-PE-Cy7 (HI10a)	BioLegend	Cat#312214	FACS (5 µl per test)
antibody	Rabbit polyclonal anti-human IgD-FITC F(ab')$_2$	DaKo	Cat#F0189	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD38-Alexa 647 (AT1)	Santa Cruz Biotechnology	Cat# sc-7325 AF647	FACS (5 µl per test)

antibody	Mouse monoclonal anti-human IgG-Alexa 700 (G18-145)	BD Pharmingen	Cat#561296	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CD138-PE (DL-101)	eBioscience (Thermo Fisher Scientific)	Cat#12-1389-42	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human IgG-PE (G18-145)	BD Biosciences	Cat#555787	FACS (5 µl per test)
antibody	Rat monoclonal anti-human/mouse IRF4-eFluor 660 (3E4)	eBioscience (Thermo Fisher Scientific)	Cat#50-9858-80	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human Ki67-Alexa 700 (B56)	BD Pharmingen	Cat#561277	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CD4-PerCP (L200)	BD Pharmingen	Cat#550631	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CD3-Alexa 700 (SP34-2)	BD Pharmingen	Cat#561805	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CD95-PE-Cy7 (DX2)	eBioscience (Thermo Fisher Scientific)	Cat#25-0959-42	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human anti-CXCR5-PE (87.1)	eBioscience (Thermo Fisher Scientific)	Cat#12-1859-42	FACS (5 µl per test)

antibody	Mouse monoclonal anti-human PD-1-Brilliant Violet 421 (EH12.2H7)	BioLegend	Cat#329920	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CXCR3-Alexa 488 (G025H7)	BioLegend	Cat#353710	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human HLA-ABC-FITC (G46-2.6)	BD Biosciences	Cat#557348	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human CXCR4-APC (12G5)	BioLegend	Cat#306510	FACS (5 µl per test)
antibody	Mouse monoclonal anti-human BST2-Alexa647 (RS38E)	BioLegend	Cat#348404	FACS (5 µl per test)
antibody	Rabbit monoclonal anti-GβL (86B8)	Cell Signaling Technology	Cat#3274	PLA (1:100)
antibody	Rabbit monoclonal anti-mTOR (7C10)	Cell Signaling Technology	Cat#2983	PLA (1:100)
antibody	Rabbit polyclonal anti-human PI3K p85	Merck Millipore	Cat#ABS1856	PLA (1:100)
antibody	Rabbit monoclonal anti-PI3 Kinase p110α (C73F8)	Cell Signaling Technology	Cat#4249	PLA (1:100)
antibody	Goat polyclonal HRP-conjugated anti-monkey IgG (H+L)	Bethyl Laboratory	Cat# A140-102P	ELISA (1:1000)

recombinant DNA reagent	pBR$_{mac239}$	PMID: 2032289	GenBank M33262	coding full sequence of *nef*-open SIV$_{mac239}$
sequence-based reagent	9520GA-F	This paper	PCR primers	GGGACAG AAATACAA TCAGGAA CAGTATAT GAATACTC CATGGAG
sequence-based reagent	9520GA-R	This paper	PCR primers	CTCCATG GAGTATTC ATATACTG TTCCTGAT TGTATTTC TGTCCC
peptide, recombinant protein	Peptide Nef$_{62-70}$	Sigma Aldrich	CTL epitope peptide	QGQYMNT PW
peptide, recombinant protein	Peptide Nef$_{62-70}$.G63E	Sigma Aldrich	Mutant CTL epitope peptide	QEQYMNT PW
peptide, recombinant protein	Peptide Nef$_{62-70}$.G63R	Sigma Aldrich	Mutant CTL epitope peptide	QRQYMNT PW
peptide, recombinant protein	GolgiStop	BD Biosciences	Cat#554724	
peptide, recombinant protein	recombinant SIV$_{mac251}$ Env gp130	ImmunoDx	Cat#5001	Ligand stimulation assay (10 µg/ml)
peptide, recombinant protein	recombinant human IL-2	Roche Diagnostics	Cat#1114752 8001	Ligand stimulation assay (100 IU/ml)
peptide, recombinant protein	recombinant human IFN-γ	Gibco (Thermo Fisher Scientific)	Cat#PHC403 3	Ligand stimulation assay (40 ng/ml)

peptide, recombinant protein	recombinant SIV$_{mac239}$ Env gp140 (biotin-conjugated)	Immune Technology	Cat#IT-001-140p	FACS and imaging cytometry (10µg/ml for probing)
peptide, recombinant protein	streptavidin-Brilliant Violet 421	BioLegend	Cat#405225	FACS (1:100)
peptide, recombinant protein	streptavidin-Alexa Fluor 488	BioLegend	Cat#405235	FACS (1:100)
peptide, recombinant protein	streptavidin-PE-Cy7	BioLegend	Cat#405206	Imaging cytometry (1:100)
commercial assay or kit	High Pure Viral RNA kit	Roche Diagnostics	Cat#11858882001	
commercial assay or kit	SIV p27 ELISA	ABL	Cat#5436	
commercial assay or kit	SureBlue TMB 1-Component Microwell Peroxidase Substrate	KPL	Cat# 5120-0075	
commercial assay or kit	SIV Western Blot Assay	ZeptoMetrix	Cat#0801500	
commercial assay or kit	Prime-Script one-step RT-PCR kit v2	TaKaRa	Cat#RR055A	
commercial assay or kit	KOD-Plus v2	Toyobo	Cat#KOD-211	
commercial assay or kit	Zero Blunt TOPO PCR cloning kit for sequencing	Invitrogen (Thermo Fisher Scientific)	Cat#K287540	

commercial assay or kit	Cytofix/Cytoperm kit	BD Biosciences	Cat#554714	
commercial assay or kit	Monkey Ferritin (Sandwich ELISA) ELISA Kit	LS Bio	Cat#LS-F36490-1	
commercial assay or kit	QuikChange II XL site-directed mutagenesis kit	Agilent Technologies	Cat#200522	
commercial assay or kit	RNeasy Plus Mini kit	Qiagen	Cat#74136	
commercial assay or kit	GeneChip WT Plus reagent	Affymetrix (Thermo Fisher Scientific)	Cat#902280	
commercial assay or kit	GeneChip Human Gene 2.0 ST Array	Affymetrix (Thermo Fisher Scientific)	Cat#902113	
commercial assay or kit	Alexa Fluor™ 488 Antibody Labeling Kit	Invitrogen (Thermo Fisher Scientific)	Cat#A20181	
commercial assay or kit	Alexa Fluor™ 647 Antibody Labeling Kit	Invitrogen (Thermo Fisher Scientific)	Cat# A20186	
commercial assay or kit	Duolink In Situ PLA Probe anti-mouse PLUS	Sigma (Merck)	Cat#DUO92001	
commercial assay or kit	Duolink In Situ PLA Probe anti-rabbit MINUS	Sigma (Merck)	Cat#DUO92005	
commercial assay or kit	Duolink flow PLA detection kit-Green	Sigma (Merck)	Cat#DUO94002	
commercial assay or kit	FuGENE6	Promega	Cat#E2691	

software, algorithm	FACS Diva	BD Biosciences	http://www.bdbiosciences.com/jp/instruments/software/facsdiva/	
software, algorithm	FlowJo v10.7.1	BD/Treestar	https://www.flowjo.com/solutions/flowjo	
software, algorithm	Affymetrix Expression Console Software	Affymetrix/Thermo Fisher Scientific	https://www.thermofisher.com/jp/ja/home/life-science/microarray-analysis/microarray-analysis-instruments-software-services/microarray-analysis-software/affymetrix-transcriptome-analysis-console-software.html	
software, algorithm	IDEAS 6.3 (+Machine learning module)	Amnis/Merck Millipore/Luminex	https://www.luminexcorp.com/imaging-flow-cytometry/	
software, algorithm	Prism 8	GraphPad Software	https://www.graphpad.com	
other	Live/Dead Aqua	Invitrogen (Thermo Fisher Scientific)	Cat#L34966	FACS and imaging cytometry (1:1000)