Characterization of Tryptophan Oxidation Affecting D1 Degradation by FtsH in the Photosystem II Quality Control of Chloroplasts

Yusuke Kato[1,2]*, Hiroshi Kuroda[3]*, Shin-Ichiro Ozawa[1,3]*, Keisuke Saito[4], Vivek Dogra[5,6], Martin Scholz[7], Guoxian Zhang[1], Catherine de Vitry[8], Hiroshi Ishikita[4], Chanhong Kim[5], Michael Hippler[1,7], Yuichiro Takahashi[3], and Wataru Sakamoto[1]**

[1] Institute of Plant Science and Resources (IPSR), Okayama University, Kurashiki, Okayama 710-0046, Japan

[2] Faculty of Agriculture, Setsunan University, Hirakata, Osaka 573-0101, Japan

[3] Research Institute for Interdisciplinary Science, Okayama University, Okayama, Okayama 700-8530, Japan

[4] Research Center for Advanced Science and Technology, The University of Tokyo, Meguro-ku, Tokyo 153-8904, Japan

[5] Shanghai Center for Plant Stress Biology, Center for Excellence in Molecular Plant Sciences, Chinese Academy of Sciences, Shanghai 200032, China

[6] Biotechnology Division, CSIR-Institute of Himalayan Bioresource Technology, Palampur 176061, India

[7] Institute of Plant Biology and Biotechnology, University of Münster, Münster 48143, Germany

[8] Institut de Biologie Physico-Chimique, Unité Mixte de Recherche 7141, Centre National de la Recherche Scientifique//Sorbonne Université Pierre et Marie Curie, Paris 75005, France.

*These authors contributed equally to this work.

**Corresponding author: Wataru Sakamoto
E-mail: saka@okayama-u.ac.jp

Author contributions

W.S. conceived the entire work along with C.K.; W.S. designed the experiments with Y.K., Y.T.; mass-spectrometric analysis was conducted by V.D. and C.K. in *Arabidopsis*, and by M.S., Y.K., S.O. and M.H. in *Chlamydomonas*; all *Chlamydomonas* mutants were generated by H.K. and were characterized by Y.K., G.Z., and S.O.; co-immunoprecipitation was performed by S.O.; the mutants in the *ftsh1* background was generated by Y.K. under the supervision of C.dV.; molecular dynamics simulation was performed by K.S. and H.I.; all authors analyzed data, and W.S. and Y.K. wrote the manuscript on behalf of all authors.

Competing Interest Statement: The authors have no conflict of interest, financial or otherwise, in relation to this study

Classification: Plant Science

Key words: *Chlamydomonas*, Oxidative post-translational modification, Photoinhibition, Photosynthesis, Protease

Running Title: D1 degradation and tryptophan oxidation

This PDF file includes: Main text
                                 Table 1
                                 Figures 1 to 6
                                 Supplementary Figures S1 to S8

Supplemental data not included in this PDF:
                                 Supplementary Methods
                                 Supplementary Tables 1 to 2
                                 Supplementary Movie 1

Significance Statement: In photosynthetic organisms, maintenance of photosynthetic light reaction is manifested by so called Photosystem II (PSII) repair system, where the reaction center protein D1 is targeted to photo-oxidative damage and rapidly degraded by the processive protease FtsH. While this system is well known to cope with photoinhibition, the actual oxidation within the D1 polypeptide and its association to degradation remained elusive. Here, we characterized oxidative modification of tryptophan (Trp) residues in the PSII core, and hypothesize that the oxidation of N-terminal Trp is one of the key oxidations in the PSII repair, likely enhancing D1's accessibility to FtsH.

Light reaction of photosynthesis is one of the most important reactions for sustaining our
environment. Photosystem II (PSII) is the initial site of photosynthetic electron transfer by water
oxidation. Light in excess, however, causes the simultaneous production of singlet oxygen, a
potent reactive oxygen species (ROS), leading to photo-oxidative damage in PSII. To maintain
photosynthetic activity, the PSII reaction center protein D1, which is the primary target of
unavoidable photo-oxidative damage, is efficiently degraded by FtsH protease. In PSII subunits,
photo-oxidative modifications of several amino acids such as Trp have been indeed documented,
whereas the linkage between such modifications and D1 degradation remains elusive. Here, we
show that an oxidative post-translational modification of Trp residue at the N-terminal tail of D1 is
correlated with D1 degradation by FtsH during high-light stress. We revealed that *Arabidopsis*
mutant lacking FtsH2 had increased levels of oxidative Trp residues in D1, among which an N-
terminal Trp-14 was distinctively localized in the stromal side. Further characterization of Trp-14
using chloroplast transformation in *Chlamydomonas* indicated that substitution of D1 Trp-14 to
Phe, mimicking Trp oxidation enhanced FtsH-mediated D1 degradation under high light,
although the substitution did not affect protein stability and PSII activity. Molecular dynamics
simulation of PSII implies that both Trp-14 oxidation and Phe substitution cause fluctuation of D1
N-terminal tail. Furthermore, Trp-14 to Phe modification appeared to have an additive effect in
the interaction between FtsH and PSII core in vivo. Together, our results suggest that the Trp
oxidation at its N-terminus of D1 may be one of the key oxidations in the PSII repair, leading to
processive degradation by FtsH.

Introduction

Light energy is essential for photosynthesis, which sustains our environment on Earth by generating oxygen and chemical energy for carbon fixation. The first step of photosynthetic electron transfer in the thylakoid membrane occurs at Photosystem II (PSII), where light energy absorbed by P_{680} chlorophyll molecules drives water oxidation, and electron is transferred to plastoquinone. PSII core complex is formed by two reaction center proteins D1 (PsbA) and D2 (PsbD), along with intrinsic antenna CP43 (PsbC) and CP47 (PsbB). Despite its orchestrated coordination, however, light energy is known to cause photooxidative damage in PSII, especially reaction center protein D1, with singlet oxygen. Photo-damaged D1 needs to be degraded to replace it with a newly synthesized one, known as the PSII repair (Lindahl, 2000; Bailey et al., 2002; Silva et al., 2003; Kato et al., 2009). When photooxidative damage in PSII exceeds the capacity of PSII repair, it ultimately leads to the status called 'photoinhibition' (Aro et al., 1993; Murata et al., 2007). PSII repair proceeds with the following steps, i) oxidative damage to D1 protein in the PSII complex, ii) migration of photo-damaged PSII laterally from grana stacks to non-appressed regions of thylakoid membranes, iii) detachment of CP43 from photo-damaged PSII allows for access of protease degrading damaged D1, and iv) concomitant D1 synthesis and reassembly of PSII into grana thylakoid. Substantial efforts to understand the mechanisms of PSII repair suggest that reversible phosphorylation of PSII core subunits is involved in fine-tuning the photo-damaged D1 turnover (Baena-González et al., 2002; Fristedt et al., 2009; Kato and Sakamoto, 2014). On the other hand, other critical steps, CP43 disassembly and the recognition of photooxidative damage in PSII for selective D1 degradation, remain to be elucidated.

Our previous studies along with those from other groups have shown that the fundamental D1 degradation in PSII repair is performed by FtsH, a membrane-bound ATP-dependent zinc metalloprotease that degrades membrane proteins in a processive manner, although several Deg proteases seem to facilitate the effective degradation by creating additional recognition sites for FtsH (Lindahl, 2000; Bailey et al., 2002; Sakamoto, 2003; Silva et al., 2003; Kato et al., 2009; Kato and Sakamoto, 2014). Their family proteins are universally conserved in prokaryotes and eukaryotic organelles (Hanna Janska, Malgorzata Kwasniaka, 2013). Photosynthetic organisms have hetero-hexameric FtsH complex in the thylakoid membrane, which is composed of type A and type B subunits (Kato and Sakamoto, 2018; Yi et al., 2022). In the thylakoid membrane of *Arabidopsis thaliana*, for example, FtsH1 or FtsH5 (type A) and FtsH2 or FtsH8 (type B) form functional FtsH complex (Yu et al., 2004; Yu et al., 2005; Zaltsman et al., 2005). *Arabidopsis* mutants lacking FtsH5 or FtsH2 are, though viable, highly vulnerable to PSII

photodamage under high light and are named *yellow variegated1* (*var1*) and *var2* from the
characteristic variegated phenotype (Chen et al., 2000; Takechi et al., 2000; Sakamoto et al.,
2002). This variegated phenotype implies that FtsH is required for proper protein quality control
during proplastid-to-chloroplast differentiation in seed plants (Miura et al., 2007; Sakamoto et al.,
2009). Similarly, the FtsH complex in the thylakoid membranes of *Chlamydomonas reinhardtii*
consists of FtsH1 (type A) and FtsH2 (type B) (Malnoe et al., 2014). *Chlamydomonas* studies
revealed that FtsH is also involved in the degradation of cytochrome b_6f complex and light-
harvesting antenna of photosystem I (PSI) (Malnoe et al., 2014; Bujaldon et al., 2017).
Furthermore, recent studies suggest that increased turnover of FtsH by itself, which is
compensated by upregulated gene expression, is crucial for their function under high-light stress
(Wang et al., 2017; Kato et al., 2018).

In *Escherichia coli*, FtsH was shown to recognize either N- or C-terminal tail of
membrane protein substrates to dislocate their substrates into the protease chamber (Ito and
Akiyama, 2005). Similarly, the fact that both D1's N-terminal end and the catalytic site of FtsH are
exposed to the stromal side implies D1 to be recognized by FtsH from its N-terminus (Umena et
al., 2011). Supporting this, processive degradation of D1 by FtsH was shown to be attenuated by
the loss of N-terminal tail of D1 (Komenda et al., 2007; Michoux et al., 2016). While these
observations suggest involvement of the N-terminal region in D1, whether D1 undergoes
oxidative modification at its N-terminal region has not been investigated.

What is the consequence of photoinhibition leading to photo-oxidative damage in PSII?
Light energy frequently leads to the generation of reactive oxygen species (ROS) such as singlet
oxygen at around PSII (Ohnishi et al., 2005; Tyystjärvi, 2008; Yamamoto et al., 2008), which may
cause oxidative post-translational modification (OPTM) of subunit proteins (Li and Kim, 2022). It
is noteworthy that light-dependent oxidation of amino acids, either in free forms or as peptide
residues, has been reported, including thiol-containing (Cys and Met) and aromatic (Tyr, Phe,
Trp) amino acids. For example, Cys and Met are prone to oxidation, whereas the oxidized Cys
and Met can be reduced enzymatically. In contrast to these reversible OPTMs, OPTM of Trp is
irreversible (Rinalducci et al., 2008; Ehrenshaft et al., 2015). Thus, the only way to remove
irreversible oxidized residues is proteolytic degradation, implying that Trp oxidation might trigger
D1 degradation, either directly or indirectly in the PSII repair. As summarized in Fig. 1a, oxidation
of Trp side chain results in the formation of oxindolylalanine (OIA), N-formylkyrnurenine (NFK),
and kynurenine (KYN). ROS, such as singlet oxygen, attacks and opens the pyrrole ring of Trp,
and forms a di-oxidized Trp derivative, NFK. Indeed, Trp residues in photosynthetic protein
components were shown to be oxidized both in vitro and in vivo (Anderson et al., 2002; Dreaden
et al., 2011; Dreaden Kasson et al., 2012). However, although oxidative modification of D1 and

other subunits has been documented previously, how these molecules are recognized and undergo degradation remains elusive. In this study, we investigated whether Trp oxidation in PSII core proteins influences D1 degradation mediated by FtsH. Our integrative approaches to address this question, by mass-spectrometry, site-directed mutagenesis, D1 degradation assay, and simulation model suggest that an N-terminal Trp oxidation is likely to be a key OPTM to trigger D1 degradation in the PSII repair.

Results

Increased OPTM of Trp residues in *Arabidopsis var2* mutant

Previous studies using isolated spinach thylakoid membranes and *Arabidopsis* seedlings revealed several Trp residues oxidized in PSII core proteins, as summarized (Fig.1b and Supplementary Table S1)(Rinalducci et al., 2008; Dreaden et al., 2011; Dreaden Kasson et al., 2012). Trp-oxidized derivatives, OIA, NFK, and KYN, give the shifts of peptide mass to + 16, + 32, and + 4 Da, respectively (Fig.1a, Fig. S1 and S2). In this study, we attempted to assess if the oxidation of certain Trp residues is associated with D1 degradation in the PSII repair cycle. To investigate this, comprehensive detection of Trp oxidation within protein extracts in *Arabidopsis* has been established using label-free quantitative mass-spectrometry, as previously reported (Dogra et al., 2019). First, we characterized Trp oxidation from total proteins of *Arabidopsis* wild-type seedlings grown in continuous light (100 μmol photons $m^{-2}s^{-1}$). Consistent with previous results, two Trp residues in D1, namely Trp-14 and Trp-317, were shown to be oxidized (Fig. S1 and S2). The total sequence coverage obtained by our mass spectrometry for D1 was 26%. Further mass spectrometry in extracts *Arabidopsis* mutants *var2* lacking FtsH2, which is shown to impair D1 degradation and exhibit substantial accumulation of ROS (Kato et al., 2009), revealed the accumulation of oxidized Trp in the PSII complex. The oxidation levels in Trp-14 and Trp-314 increased 1.8-fold and 1.4-fold in *var2* compared to the wild type, respectively (Fig. 1c). These results prompted us to characterize the role of these Trp oxidations in the PSII repair further.

OPTM of Trp residues in *Chlamydomonas* PSII core proteins

To validate whether Trp oxidation detected in *Arabidopsis* seedings is also detectable in other model organisms, we investigated extracts from *Chlamydomonas* thylakoid membranes (Table1 and Supplementary Table S2). Mass spectrometry following trypsin digestion demonstrated that several Trp residues in PSII were oxidized; among the core subunits, D1 had oxidation of four

residues, D2 had three, CP43 had five, and CP47 had three (Table1 and Supplementary Table
S2). The position of oxidized Trp residues in their amino acid sequence was shown in Fig. 1b
and Fig. S3. Together with previous studies, Trp-14 and Trp-317 in D1, Trp-21 and Trp-328 in D2,
Trp-353 and Trp-375 in CP43, and Trp-275 and Trp-302 in CP47 were commonly identified
among at least two organisms. Spatial arrangement of these Trp residues that were commonly
oxidized in *Arabidopsis* and *Chlamydomonas* was compared within the structure of PSII complex,
as shown in Fig.1d and e (oxidized Trp residues were assigned in the PSII dimer, and its 3D
image is shown in Supplementary Movie 1 online). Also, oxidized Trp residues were assigned in
the PSII structure from *Thermosynecoccocus vulcanus* and were shown in Fig. S4. Intriguingly,
most of these oxidized Trp residues are positioned at the luminal side of the PSII core complex,
which appeared to surround the Mn_4O_5Ca cluster in the PSII structure model. In contrast, two
oxidized Trp residues, Trp-14 in D1 and Trp-21 in D2 close to the N-terminus of the polypeptides,
are located on the stromal side. The fact that the oxidized Trp residues are predominantly
observed around the Mn_4O_5Ca cluster may reflect photoinhibition of PSII electron donor side and
concomitant ROS generation. In contrast, stromal Trp oxidation is novel and localized at the N-
terminal alpha-helix, which might suggest its effects in processive D1 degradation.
142
Site-directed mutagenesis of Trp residues undergoing OPTM in D1
To test whether any change in the oxidized Trp residues is associated with D1 degradation, we
performed site-directed mutagenesis using chloroplast transformation in *Chlamydomonas*, to
substitute the corresponding Trp for other amino acids in D1. Based on the mass-spectrometric
results, we focused on Trp-14 and Trp-317, each of which was replaced by Ala (non-polar and
hydrophobic) or Phe (aromatic and hydrophobic), respectively. The vectors harboring
spectinomycin/streptomycin-resistant *aadA* cassette and the mutated *psbA* gene were
transformed into Δ*psbA* mutant Fud7 (Fig. 2a). Transformants were selected on mixotrophic Tris-
acetate-phosphate (TAP) plates containing spectinomycin, and their homoplasmicity was
subsequently confirmed by PCR using specific primers and sequencing. All transformants grew
like the control strain on mixotrophic TAP plates (Fig. 2b). However, the transformants in which
Trp-14 or Trp-317 was substituted to Ala, (W14A, W317A, and W14A/W317A) showed
significantly impaired growth on photoautotrophic high salt minimal (HSM) plates. Ala substitution
at both Trp-14 and Trp-317 led to decreased photosynthetic activities due to reduced
accumulation of D1 and other PSII core proteins (Fig. 2b - d), indicating its defect in stability and/or
the translation of D1 protein.
In contrast, Phe substitution at the same sites had little effect on their growth under
growth light (30 μmol photons $m^{-2}s^{-1}$). These transformants (W14F, W317F, and W14F/W317F)

accumulated PSII core proteins whose amounts were comparable to the control levels. They
showed normal photosynthetic activities as evidenced by comparable electron transport rates
through the PSII complex and oxygen-evolving activity (Fig. 2c and e). We next examined their
photoautotrophic growth under high light (320 µmol photons $m^{-2}s^{-1}$). Under this condition,
however, W14F exhibited significantly impaired growth, and W317F grew slightly slower than
control cells. Double mutant W14F/W317F synergistically increased high-light sensitivity but the
growth defect appeared to be similar to W14F, suggesting that Phe substitution at Trp-14, but not
at Trp-317 had profound effects in the PSII repair cycle (Fig. 2b and Fig. S5).
169
Site-directed mutagenesis of Trp residues in CP43
We next examined if Trp modification in CP43 influences in high-light sensitivity. As an important
step in the PSII repair, PSII complex is partially disassembled by CP43 detachment, and this
process likely allows FtsH to access photo-damaged D1. Therefore, Trp oxidation in CP43 may
play a role in PSII disassembly and D1 degradation concomitantly. To test this, we substituted
Trp-353 and Trp-375 for either Ala or Phe as carried out in D1. Transformants were generated
by cotransformation of Fud7, using the vector harboring wild-type *psbA* gene and the vector
harboring the mutated *psbC* gene. Consequently, we obtained four single mutants (W353A,
W353F, W375A, and W375F) and two double mutants (W353A/W375A and W353A/W375F)
(Fig. S6). Mixotrophic growth on TAP plates was comparable among all transformants and
control cells (Fig. S6). All the transformants except for W353A/W375A grew normally on the
phototrophic condition under growth light condition. Supporting normal growth, immunoblot
analysis showed normal accumulation of PSII core proteins, D1 and CP43, in all lines except for
the double mutant W353A/W375A. On the other hand, D1 and CP43 were severely reduced in
W353A/W375A, indicating that Ala substitution in these residues resulted in highly unstable or
impaired PSII complex formation (Fig. S6). Further analysis of these mutants under high light
showed that the transformants did not increase high-light sensitivity on their growth unlike D1
W14F mutant. At least in our site-directed mutagenesis, Trp oxidation in CP43 appeared to have
little impact on the PSII repair.
189
Substitution of Trp-14 with Phe accelerates D1 degradation
To evaluate whether Trp substitution in D1 affects PSII damage or repair, we next measured the
maximum quantum yield of PSII (Fv/Fm) and subsequently monitored D1 levels under growth or
high-light conditions. Trp-substituted lines grown in TAP medium under growth light were pre-
incubated in the presence or absence of chloramphenicol (CAM), an inhibitor of chloroplast
protein synthesis. CAM blocks the PSII repair at the step of D1 synthesis and allows us to

evaluate photodamage and D1 degradation. Cells incubated under growth light or high light were
subjected to chlorophyll fluorescence measurement and immunoblot analysis. Under growth light
condition and in the absence of CAM, both PSII activity (Fv/Fm values) and D1 levels were
comparable among all Trp-substituted lines and the control (Fig. 3a). This result was consistent
with their photoautotrophic growth under growth light. When CAM was added, D1 levels
decreased only slightly during incubation (90 min) in the control. D1 degradation rate was
comparable in all Trp-substituted lines and control (Fig. 3c), indicating that all Trp-substituted D1
proteins formed stable and functional PSII complex under growth light.

Under high-light condition, however, Fv/Fm values in Trp-substituted lines significantly
decreased, compared to that observed in the control even in the absence of CAM (Fig. 3b).
These vulnerabilities to high light were consistent with their impaired growth under high light (Fig.
2). To our surprise, D1 levels in W14F and W14F/W317F concomitantly decreased during high-
light incubation (Fig. 3b). In contrast, D1 levels in W317F were similar to those in control cells.
When the PSII repair engages properly, high-light irradiation does not alter D1 levels because
turnover of photo-damaged D1 encounters rapid D1 synthesis. Given decreased D1 under high
light, W14F was likely to cause faster D1 degradation. To confirm this possibility, D1 degradation
in the presence of CAM was measured. PSII activity in all Trp-substituted lines fell at similar rates
compared with control cells in the presence of the CAM (Fig. 3d), indicating the light-induced
damage was at the similar level among all Trp-substituted lines and the control. In contrast, our
time course experiment indicated that W14F and W14F/W317F decreased D1 faster than the
control and W317F (Fig. 3d); the D1 level in W14F and W14F/W317F decreased approximately
60% and 50% of the initial level, respectively, whereas those in control cells and W317F remained
80% (Fig. 3d).

Based on these D1 degradation assays, we assumed that D1 degradation by
proteolysis was enhanced by W14F substitution, despite the fact that PSII suffered from
photodamage equally among other lines and the control during high-light irradiation. To exclude
the possibility that W14F decelerates D1 synthesis rather than accelerating degradation, we
analyzed protein synthesis in Trp-substituted lines by in vivo pulse labeling in the presence of
cycloheximide, which prevents the synthesis of the nuclear-encoded proteins. As shown in Fig.
S7, D1 synthesis was shown to proceed comparably in all lines. Collectively, our findings
demonstrated that Trp-14 substitution to Phe enhanced D1 degradation, whereas it affected
neither the light-induced damage in PSII, D1 synthesis, nor PSII stability.

229 Enhanced D1 degradation due to the substitution of Trp-14 is mitigated in the *ftsH* mutant

To address whether the increased D1 degradation in W14F (and W14F/W317F) involved

proteolysis by FtsH, these Trp substitutions were introduced into an *ftsH* mutant deficient in

thylakoid FtsH activity. In the thylakoid membrane of *Chlamydomonas*, a hetero-oligomeric FtsH

complex composed of FtsH1 (type-A) and FtsH2 (type-B) exists, and the *ftsh1-1* mutant,

expressing an inactive FtsH1 due to the amino-acid substitution in the ATP-binding domain, has

been reported (Malnoe et al., 2014). We performed mating W14F and W14F/W317F

transformants (mt+) with *ftsh1-1* (mt-), and the resulting mutants, W14F *ftsh1* and W14F/W317F

ftsh1, were subjected to D1 degradation assay. The results indicated that as expected, the

enhanced D1 degradation observed in W14F and W14F/W317F cells was partially mitigated

under *ftsh1-1* background (Fig. 4), when CAM was added. These results suggested that FtsH

plays a key role in the increased D1 degradation in W14F and W14F W317F.

Molecular dynamics simulation suggests W14F mimicking Trp-14 oxidation

Although our site-directed mutagenesis in Trp-14 showed its effect in D1 degradation, how Trp

oxidation can be structurally correlated with Trp to Phe mutagenesis should be taken into

consideration. To investigate this, we employed molecular dynamics (MD) simulation, a powerful

tool to simulate movements of amino acids in a protein complex, using the crystal structure of

PSII complex from *Thermosynechococcus vulcanus* (Sakashita et al., 2017b; Sakashita et al.,

2017a; Kawashima et al., 2018). D1 Trp-14 is located in the first α-helix at the N-terminus and

hydrogen-bonded with PsbI Ser-25 (Fig. 5a). It is deduced that this hydrogen bond restricts the

conformational change around D1 Trp-14 and limit the fluctuation of D1 N-terminus. The

simulation indicated that the hydrogen bond disappeared (Fig. 5b, c) and the structural fluctuation

of D1 Trp-14 was increased as compared with WT (Fig. 5d) when Trp-14 is oxidized to NFK or is

replaced with Phe residue. The increased fluctuation of the side chain also influences the Cβ-Cβ

distance between D1 Trp-14 and PsbI Ser-25; the two Cβ atoms became farther away from each

other when D1 Trp-14 is oxidized to NFK (Fig. 5e). Of note, the amino acid substitution on Trp-

14 to Phe showed similar trends as those observed when D1-Trp-14 is oxidized to NFK. These

results suggest that the structural change of Trp-14 affect the local movement. The increased

fluctuation of the first α-helix of D1 would give a chance to recognize the photo-damaged D1 by

FtsH protease.

Augmented interaction between D1 and FtsH by substituting Trp-14/317

Presented experimental results collectively raise the possibility that oxidation of Trp-14 is one of

the key OPTMs for D1 degradation by FtsH. We raised a possibility that W14F mimics Trp-14

oxidation and shows increased FtsH association with D1. Since quantitative interaction of the

protein and the protease remains to be elucidated, we performed differential pull-down assay. To
emphasize the effect of the substituted amino acid residues and minimize potential oxidation of
other amino acid residues, we decreased light intensity during cell culture and removed oxygen
molecule from the buffer solution during the assay. *Chlamydomonas* cells were grown under dim
light with gently shaking and were harvested at the mid-log phase. Subsequently, we isolated
thylakoid membrane from the gently disrupted cells and performed co-immunoprecipitation in
anoxic aqueous solution. Quantification of D1 and D2 levels, normalized by FtsH in the co-
immunoprecipitated sample showed that the relative D1 and D2 protein amounts were
statistically higher in W14F/W317F than the control (Fig. 6). We concluded that W14F increased
affinity between FtsH and the reaction center proteins, which leads to enhanced D1 degradation.
275
276

277   **Discussion**
278

Recent progress in mass-spectrometry has advanced our understanding of holistic OPTM in
photosynthetic protein complexes. Along this line, we investigated Trp oxidation in PSII in this
study, and attempted to address whether any modification in amino acid residues was correlated
with the PSII repair. PSII is one of the major sites for ROS generation due to photoinhibition, and
oxidized amino acid residues in PSII core proteins have been reported previously (Kale et al.,
2017)(Frankel et al., 2012). In general, Met and Cys are sensitive amino acid residues for ROS-
mediated oxidation (Rinalducci et al., 2008)(Ehrenshaft et al., 2015). However, those OPTM can
be converted back in reduced forms by methionine sulfoxide reductase and disulfide reductase,
respectively. In contrast, Trp oxidation is irreversible, and its replacement requires whole protein
degradation and de novo synthesis, implicating Trp suitable for flagging photo-oxidative
damaged proteins that undergo degradation in the PSII repair.
290   Previous studies have reported OPTM of several Trp residues in PSII core proteins in
vitro (Dreaden Kasson et al., 2012)(Dreaden et al., 2011)(Anderson et al., 2002). In addition,
Dogra et al. confirmed several oxidized Trp residues in PSII core proteins of *Arabidopsis* in vivo
(Dogra et al., 2019). Our mass-spectrometry in *Chlamydomonas* further indicated that oxidation
in some of the Trp residues was detected among algae and land plants. It is noteworthy that a
majority of Trp residues are located on the luminal side (Table 1) in the vicinity of the Mn_4O_5Ca
cluster, consistent with the previous observations that amino acid residues around the Mn_4O_5Ca
cluster are oxidized at the early stage of photoinhibition (Kale et al., 2017)(Frankel et al., 2012).
However, OPTM is not limited to the luminal side but also found in the stromal side, which
accounts for photoinhibition at the electron acceptor side (Vass, 2012). Our data indicated the

presence of commonly oxidized Trp residues as represented by Trp-14 and Trp-317 in D1 protein. These are in fact Trp residues highly conserved among photosynthetic organisms (Fig.1). Technically, detection of light-dependent OPTM was considered to be difficult; an extensive OPTM leads to fully inactive PSII, whereas D1 degradation in the PSII repair may diminish OPTM under optimal light conditions. We therefore assumed that OPTM associated with PSII repair might accumulate significantly in the mutant that is defective in FtsH. Supporting this, our quantitative mass spectrometry indicated that *var2* lacking FtsH2 in *Arabidopsis* had increased levels of oxidation in Trp residues, particularly in Trp-14 and Trp-317. Although indirect, these results strongly suggested an interconnection between Trp oxidation and D1 degradation.

OPTM of Trp residues causes irreversible modification in D1 protein and is likely to mark a substrate of D1 degradation. In the PSII repair, a series of events including migration of photo-damaged PSII to non-appressed regions of thylakoid membranes, release of CP43 from the PSII, and recognition of photo-damaged D1 for selective D1 degradation, are essential. In this scenario, FtsH interacts with a partially disassembled PSII complex lacking CP43 protein, called RC47 (Kato and Sakamoto, 2009; Järvi et al., 2015). Close access to the photo-damaged D1, followed by the recognition of its N-terminal region, is concomitantly necessary for FtsH to proceed with processive D1 degradation. Therefore, the OPTM would be involved in the CP43 disassembly or the recognition of damaged D1 protein. Krynická et al. indicate that the accessibility to PSII core proteins drives selective protein degradation by FtsH in the cyanobacterium *Synechocystis* PCC 6803 (Krynická et al., 2015). This observation suggests that D1 protein in the RC47 complex is promptly degraded even if D1 did not suffer from photodamage. However, all site-directed mutants mimicking Try oxidation by Trp to Phe substitution have stable and functional PSII complexes under growth light, suggesting that the OPTM would not induce the disassembly of CP43. We also tested whether Trp oxidation in CP43 affects PSII repair by site-directed mutagenesis. Similarly to the case in D1, none of Trp substitutions at the site of OPTM in CP43 (Trp-353 and Trp-375) affected the D1 degradation. Additionally, the CP43 transformants did not show the increased photosensitivity (Fig. S6). Our results somewhat appear to contradict the previous report in cyanobacterium that the mutants in which Trp-353 (Trp-352 in *Synechocystis* 6803) was substituted to Leu, Cyc, or Ala increased photo-sensitivity under high-light conditions (Anderson et al., 2002). This might be due to the use of extremely high-light irradiation (5,000 µmol photons m^{-2} s^{-1}), under which severe photo-damage in PSII complex was rendered. We consider that irreversible Trp oxidation in CP43, if to be repaired, may require a rapid turnover rate comparable to D1 degradation, which is not the case. Although further study is necessary to elucidate the disassembly mechanisms of CP43 during the PSII repair cycle, Trp oxidation in D1, rather than CP43 disassembly, might be important for the recognition of FtsH.

335    To examine its effect on D1 degradation, we performed site-directed mutagenesis of

336 the corresponding Trp residues using *Chlamydomonas* chloroplast transformation. While Trp to

337 Ala substitution in these sites (W14A or W317A) appeared to compromise PSII complex

338 formation, Trp to Phe substitution (W14F and W317) gave us a hint in the critical role of OPTM.

339 We showed that W14F, but not W317F, caused higher photo-sensitivity with the rapid decrease

340 of D1 under high-light irradiation (Fig. 2). Given that W14F affected neither D1 synthesis (Fig.

341 S7), stability of PSII complex formation, nor PSII activity under non-photoinhibitory conditions, it

342 was concluded that the mutation results in enhanced D1 degradation. In our D1 degradation

343 assay of wild type, generally, D1 turnover is too fast to detect unless inhibitor of chloroplast protein

344 synthesis (CAM) is added. In sharp contrast, W14F proceeds with rapid D1 degradation even

345 without CAM. Reportedly, numerous amino acid substitutions have been introduced in D1, which

346 may or may not compromise PSII activity. To our knowledge, however, mutations that accelerate

347 D1 degradation have not been found except for W14F in this study. We thus consider that Trp-

348 14 is particularly important, at least for FtsH to recognize photodamaged D1 as described below.

349    Because FtsH-mediated D1 degradation is crucial for the PSII repair cycle, recognition

350 of photodamaged D1 by FtsH protease is a critical step. Following observations suggest that Trp-

351 14 oxidation is one of the key OPTMs for degrading photodamaged D1. First, enhanced D1

352 degradation in W14F well fits the notion that PTM in the N-terminus of D1 is important to execute

353 processive degradation by FtsH, as proposed previously. For example, the lack of an N-terminal

354 helix attenuates proper D1 degradation (Komenda et al., 2007)(Michoux et al., 2016). The

355 excision of N-terminal Met by organellar Met aminopeptidase and prokaryotic-like peptide

356 deformylase was shown to be required for FtsH-mediated D1 degradation (Adam et al., 2011).

357 Phosphorylation of the D1 N-terminus was also shown to affect proteolysis and contribute to the

358 fine‐tuned D1 degradation pathway (Koivuniemi et al., 1995)(Rintamäki et al., 1996)(Kato and

359 Sakamoto, 2014). Together with these, it is possible that Trp-14 oxidation is likely to play a role

360 in 'photodamage-dependent' degradation, although we cannot rule out the possibility that other

361 OPTMs may have additive effects. Second, our MD simulation strongly suggests that W14F is

362 similar to Trp-14 oxidation W14* (modified as NFK) in terms of allowing a regional conformational

363 change around the N-terminal helix, thereby increasing fluctuation of the side chain. This

364 fluctuation appears to be manifested by losing hydrogen bonding with Ser-25 of PsbI, a short

365 peptide localized close to D1 and CP43 in the PSII core complex. Although further studies are

366 needed, our simulation is consistent with our notion that Trp-14 is a target of OPTM that alters

367 subtle but critical structural change at the N-terminus of D1.

368    Based on these observations, we propose a working model of 'photodamaged D1

369 recognition' in which Trp oxidation plays a role in processive degradation by FtsH (Fig. 6c). As a

consequence of photoinhibition, ROS is produced around PSII and leads to OPTM of numerous

residues. Among these, Trp-14 and Trp-317 are prone to oxidation likely due to their relative

positions in PSII. While oxidation takes place in both, Trp-14 causes a conformational change at

the N-terminus, which triggers enhanced access of FtsH for subsequent processive degradation.

Supporting this, we observed augmented association between D1 and FtsH in W14F/W317F

(Fig. 6). It is unlikely, however, that Trp-14 oxidation alone is sufficient to drive degradation of

photodamaged D1, because a stepwise dissociation of PSII core complexes is prerequisite.

OPTM of Trp residues has been observed in various proteins (Kasson and Barry, 2012).

For example, Trp oxidation has been identified in ATP synthase alpha subunit in mitochondria,

one of the target proteins for oxidative stress in the mitochondrial inner membrane (Rexroth et

al., 2012). In this case, the oxidation is not random but selectively targets specific Trp. The

oxidized ATP synthase might be degraded by mitochondrial FtsH homologs, m-AAA and i-AAA

proteases, which have an essential role in the quality control of aberrant proteins in mitochondrial

membranes. Furthermore, a previous study in chloroplasts suggests the Trp oxidation in the

stress responses related to singlet oxygen; Dogra et al. (2019) report specific oxidation of Trp in

EXECUTER1 (EX1), a sensor protein of singlet oxygen in plastid signaling. The oxidation of

specific Trp residue is required for ROS signaling mediated by light-dependent EX1 degradation

by FtsH (Dogra et al., 2019). Together with our results, these reports imply a general mechanism

between oxidized modification of target protein and substrate recognition by FtsH. Future

proteomic approaches for investigating OPTM will reveal the general substrate recognition

mechanisms by FtsH in the thylakoid membranes.

Methods

Detection of Trp oxidation in *Arabidopsis*

Chloroplasts were isolated from 3-week-old plants of WT and *var2* (SAIL_253_A03) grown under

continuous light (80 μmol photons $m^{-2}s^{-1}$ at 20 ± 2°C) conditions. The collected rosette leaves

were homogenated in chloroplast isolation buffer [50 mM Hepes-KOH pH 8, 5 mM $MgCl_2$, 5 mM

EDTA pH 8, 5 mM EGTA pH 8, 10 mM $NaHCO_3$, and 0.33 M D-sorbitol, supplemented with

SIGMAFAST^{TM} Protease Inhibitor (1 tablet per 100 ml)]. The homogenate was filtered through

four layers of Miracloth and centrifuged at 400 × *g* for 8 min at 4°C. The pellets were suspended

in isolation buffer and loaded onto a two-step Percoll gradient (40:80%) solution to separate intact

and broken chloroplasts. The intact chloroplasts enriched between the two Percoll steps were

carefully collected and washed twice with HS buffer (50 mM Hepes-KOH pH 8, and 0.33 M D-

sorbitol). Chloroplasts corresponding to equal amounts of chlorophyll were lysed, and the

proteins extracted using 6 M guanidine hydrochloride buffer (guanidine hydrochloride dissolved

in 100 mM Tris, pH 8.5). The lysed samples were sonicated in an ice bath for 1 min with a pulse

of 3 s 'on' and 5 s 'off', followed by heating at 95 °C for 5 min, and then centrifugation at 21 000

g for 30 min at 4 °C. Total protein content was estimated using a PierceTM BCA protein assay kit

(ThermoFisher Scientific).

Mass spectrometric analysis for protein identification and PTM analysis was done according

to our previous study (Dogra et al., 2019). For MS analysis, equal amounts of total protein (2 µg

µl^{-1}) from three independent biological samples were denatured using 10 mM DTT at 56 °C for

30 min followed by alkylation in 50 mM iodoacetamide at room temperature for 40 min in the dark.

Reduced-alkylated proteins were then desalted a Nanosep membrane (Pall Corporation,

MWCO 10K) in 200 µL of 100 mM NH_4HCO_3 buffer, followed by digestion in buffer containing 40

ng/µl trypsin in 100 mM NH_4HCO_3 (corresponding to the enzyme-to-protein ratio of 1:50) at 37 °C

for 20 h. The digested peptides were dried and resuspended in 0.1% (v/v) formic acid solution.

Digested peptides were separated using nanoAcquity Ultra Performance LC (Waters, Milford,

MA, USA) and analyzed by using Q Exactive Mass Spectrometer (Thermo Fisher Scientific, San

Jose, CA, USA) as described in our previous study (Dogra et al., 2019). The mass spectra were

submitted to the Mascot Server (version 2.5.1, Matrix Science, London, UK) for peptide

identification and scanned against the *Arabidopsis* protein sequences (downloaded from TAIR;

http://www.arabidopsis.org/). Database searches were carried out with peptide mass tolerance

of 20 ppm, fragment mass tolerance of 0.02 Da, and a maximum of two missed cleavages.

Carbamidomethylation of Cys was set as a fixed modification, while oxidations of Met and Trp

were defined as variable modifications. The significance threshold for search results was set at

a *P*-value of 0.05 and an Ions score cut-off of 15. For quantification, raw MS data files were

processed and analyzed using MaxQuant software (version 1.5.8.3) with a label-free quantitation

(LFQ) algorithm. Parent ion and MS2 spectra were searched against the *Arabidopsis* protein

sequences. The precursor ion tolerance was set at 7 ppm with an allowed fragment mass

deviation of 20 ppm. Carbamidomethylation of Cys was set as a fixed modification, while

oxidations of Met and Trp were defined as variable modifications. Peptides with a minimum of six

amino acids and a maximum of two missed cleavages were allowed. False discovery rate (FDR)

was set to 0.01 for both peptide and protein identification. The absolute intensity values were

used to calculate the abundance of oxidized peptides. Label-free quantitation of oxidized

peptides using mass spectrometry were performed according to previously described method

(Luber et al., 2010; Schwanhäusser et al., 2011; Duan et al., 2019).

Strains and generation of chloroplast transformants in *Chlamydomonas*

The *psbA* deletion mutant of the green alga *Chlamydomonas reinhardtii*, Fud7 (Bennoun et al., 1986) was used for chloroplast transformation in this study. The vector, which lacks large portion of *psbA* gene (Takahashi et al., 1996), was used for plasmid construction. Details in plasmid construction performed in this study were described as separate Supplementary Materials and Methods. To obtain *psbA* mutants, each *psbA* transformation vector was biolistically delivered into chloroplast of the Fud7 mutant using a particle gun (IDERA GIE-III, TANAKA Co. Ltd., Sapporo, Japan). Chloroplast transformants were selected by at least four rounds of single colony purification on TAP agar plates containing spectinomycin (150 µg mL^{-1}) as described previously (Takahashi et al., 1996). The CP43 mutants were generated according to Kuroda et al. (2021) (Kuroda et al., 2021). The psbD gene in the Fud7 was disrupted and we obtained Fud7-ΔpsbD mutant as a recipient for CP43 mutagenesis. The DNA delivery methods are the same with *psbA* mutagenesis experiment.

Detection of Trp oxidation in *Chlamydomonas*

Cultivation of the algae was carried out under constant light (50 µmol photons m^{-2}s^{-1} or 500 µmol photons m^{-2}s^{-1}) in TAP medium for 24h. Cells were harvested by centrifugation (2500 x g for 5 min at room temperature), frozen in liquid nitrogen and stored at -80°C until further use. For protein extraction, lysis buffer (100 mM Tris/HCl pH 8.5, 2% (w/v) SDS, 1mM PMSF, 1 mM benzamidine) was added to frozen cell pellets and incubated for 10 min at 65°C and 1000 rpm in a Thermomixer (Eppendorf, Germany). The lysate was cleared by centrifugation (18,000 x g for 10 min at 25°C) and the protein content of the supernatant was determined using the Pierce BCA protein assay kit (Thermo Fisher Scientific). Reduction, alkylation and tryptic digestion (50 µg of protein per sample) was performed in centrifugal filters (Amicon Ultra-0.5, 30 kDa cut-off, Merck Millipore) according to the FASP protocol (Wiśniewski et al., 2009). Peptides (5 µg per sample) were desalted using self-packed C18-StageTips as previously described (Kulak et al., 2014), followed by vacuum centrifugation until dry. Prior to LC-MS/MS analysis peptide samples were resuspended in 2% (v/v) acetonitrile/0.05% (v/v) trifluoroacetic acid at a concentration at a concentration of 1 µg/µl. LC-MS/MS analysis was carried out using an Ultimate 3000 nanoLC (Thermo Fisher Scientific) coupled to an Q Exactive Plus mass spectrometer (Thermo Fisher Scientific) via a nanospray interface. Samples (1 µl) were loaded on a trap column (C18, Acclaim PepMap 100, 300 µM × 5 mm, 5-µm particle size, 100-Å pore size; Thermo Scientific) at a flow rate of 10 µl/min for 3 min using 2% (v/v) acetonitrile/0.05% (v/v) trifluoroacetic acid in ultrapure water. Subsequently, peptide separation was performed on a reversed phase column (C18, Acclaim Pepmap C18, 75 µm x 50 cm, 2 µm particle size, 100 Å pore size, Thermo Fisher

Scientific) at a flow rate of 250 nl/min using the eluents 0.1% (v/v) formic acid in ultrapure water

(A) and 80% (v/v) acetonitrile/0.1 % (v/v) formic acid in ultrapure water (B). The following gradient

was applied: 2.5-5% B over 10 min, 5-22% B over 90 min, 22-30% B over 70 min, 30-99%B over

10 min, 99 % B for 20 min.

MS full scans (m/z 350-1600) were acquired in positive ion mode at a resolution of 70,000

(FWHM, at m/z 200) with internal lock mass calibration on m/z 445.120025. The AGC target was

set to 3e6 and the maximum injection time to 50 ms. For MS2, the 12 most intense ions with

charge states 2-4 were fragmented by higher-energy c-trap dissociation (HCD) at 27%

normalized collision energy. AGC target value was set to 5e4, minimum AGC target to 5.5e2,

maximum injection time to 55 ms and precursor isolation window to 1.5 m/z.

Peptide and protein identification were carried out in Proteome Discoverer 2.4 (Thermo Fisher

Scientific) using the MSFragger node (MSFragger 3.0)(Kong et al., 2017) with default parameters

for closed searches (precursor mass tolerance: 50 ppm, precursor true tolerance: 20 ppm,

fragment mass tolerance: 20 ppm, maximum missed cleavages: 1). Spectra were searched

against a concatenated sequence database containing nucleus-encoded proteins

(www.phytozome.org, assembly version 5.0, annotation version 5.6), supplemented with

proteins encoded in the chloroplast (NCBI BK000554.2) and mitochondria (NCBI NC_001638.1),

as well as common contaminants (cRAP, www.thegpm.org/crap/). Carbamidomethylation was

set as static modification. The following variable modifications were defined: N-acetylation of

protein N-termini, oxidation of methionine, and various products of tryptophan oxidation

(kynurenine (+3.995 Da), hydroxytryptophan (+15.995 Da), hydroxykynurenine (+19.990 Da), N-

formylkynurenine (+31.990 Da), dihydroxy-N-formylkynurenine (+63.980 Da). Peptide-spectrum-

matches (PSMs) were filtered using the Percolator node to satisfy a false discovery rate (FDR)

of 0.01. Subsequently, identifications were filtered to achieve a peptide and protein level FDR of

0.01.

Growth test

Cells were grown in TAP liquid medium without shaking at 23-24°C under the light-dark

synchronized condition (10 hours light at 50 µmol photons $m^{-2}s^{-1}$ or less and 14 hours darkness).

Subsequently the cells were harvested by centrifugation at 2000 × g for 10 min at 25°C and were

suspended in TP (Tris Phosphate) medium for washing. After finishing the washing process, the

cell concentration was adjusted at 25 ng Chl $µL^{-1}$ with TP medium. The liquid culture was spotted

on solid medium at 100 ng Chlorophylls/spot. When we evaluate the cellular growth rate in the

liquid culture, the cells grown under 30 µmol photons $m^{-2}s^{-1}$ in TAP medium were suspended in

the TP medium at 0.1 of OD750 and were incubated under 30 or 350 µmol photons $m^{-2}s^{-1}$.

Measurement of photosynthetic activity

Chlorophyll fluorescence induction kinetics of *Chlamydomonas* transformants were measured

using a pulse amplitude-modulated fluorometer (Dual-PAM-100; Heinz Walz GmbH). Before

measurements, cultured cells were maintained in the dark for 5 min to oxidize the plastoquinone

pool fully. Initial fluorescence yield of PSII (F_O) and maximal fluorescence yield of PSII (F_M) were

measured. Maximal PSII quantum yield (Fv/Fm) was determined as Fv/Fm = $(F_M–F_O)/F_M$. Light-

induced oxygen-evolving activity of cells was measured using a Clark-type O_2 electrode

(Oxytherm OXYT1; Hansatech Instruments). Briefly, cells were grown in TAP culture under 5

µmol photons $m^{-2} s^{-1}$ to reach 5-10 µg Chl mL^{-1}. O_2-evolving activity of cells (5 µg Chl mL^{-1}) in the

presence of 0.3 mM 2,6-dichloro-1,4-benzoquinone was measured using a Clarke-type O_2

electrode with an actinic light at 7,800 µmol photons $m^{-2} s^{-1}$ at 25°C as described (Kuroda et al.,

2014).

Immunoblotting

Total proteins were solubilized in SDS-PAGE sample (125 mM Tris-HCl, pH 6.8, 2% [w/v] SDS,

100 mM dithiothreitol, 10% [v/v] Glycerol, 0.05% [w/v] BPB) buffer at 96°C for 1 min, and then

were loaded based on equal chlorophyll. The proteins were electrophoretically transferred onto

polyvinylidene difluoride membrane (Atto Corp.) after SDS-PAGE. The membranes were

incubated with specific polyclonal antibodies: anti-D1 (raised against N-termimus, dilution 1:5,000)

(Kato et al., 2012), anti-D2 (AS06 146, Agrisera; dilution, 1:5,000), anti-CP43 (AS11 1787,

Agrisera; dilution, 1:5,000), anti-PsaA (a gift from Kevin Redding, Arizona State University, dilution

1:5000), and anti-Lhca1, dilution 1:5000)(Ozawa et al., 2018). The signals were visualized by

using a Luminata Forte Western HRP Substrate (Merck Millipore) with Molecular Imager

ChemiDoc XRS+ imaging system (Bio Rad Laboratories, Inc., USA). Signal intensities were

quantified using NIH Image.

D1 degradation assay

Cells were grown in TAP liquid medium at 22°C under continuous light-condition (30 µmol

photons $m^{-2}s^{-1}$). Cultured cells were harvested by centrifugation at 600 × g for 5 min. The cell

pellets were resuspended in a new TAP liquid medium as a final concentration of 0.5 µg Chl mL

$^{-1}$. Then, the cells were preincubated in the presence or absence of chloramphenicol (100 µg

mL^{-1}) in the dark for 30 min. Subsequently, the cells were incubated under high-light or growth

light conditions (350 or 30 µmol photons $m^{-2}s^{-1}$) with stirring. Cells in 400 µl culture were collected

at each time points (30, 60, 90 min) by centrifugation, and the resulting cell pellet was
resuspended in 100 µl of SDS-PAGE sample buffer.
546

Pulse labeling of chloroplastic proteins
Cells grown in pre-culture medium (TAP media with less sulfur) were harvested by centrifugation
at 600 × g for 5 min and were washed by TAP media containing no sulfur. After centrifugation,
the cells were resuspended to 25 µg Chl mL^{-1} in TAP media containing no sulfur and incubated
for 2h. Subsequently, sulfur-starved cells were labeled with 5 µ Ci mL^{-1} [^{35}S]Na$_2$SO$_4$ (American
Radiolabeled Chemicals) in the light at 50 µmol photons m^{-2}s^{-1} in the presence of 10 µg mL^{-1}
cycloheximide. At each time point (1, 2, 4 min), cell samples were collected and immediately
frozen in liquid nitrogen.
555

Thylakoid membrane isolation and the following co-immunoprecipitation in anoxic
aqueous solution
Cells grown in TAP medium under 5 µmol photons m^{-2} s^{-1} were harvested by centrifugation at
2,000 × g for 10 min at 25°C. All buffers were incubated at 25°C for 60 minutes in the presence
of 100 mM glucose, 40 U/mL glucose oxidase, and 50 U/mL catalase to remove oxygen before
chilling. Cells were suspended in suspension buffer (10 mM HEPES-KOH pH 8.0), broken by
double passage through an airbrush at a pressure of 0.2 MPa (0.2 mm aperture airbrush). The
broken materials were suspended in high sucrose concentration solution (1.8 M sucrose, 10 mM
HEPES-KOH pH 8.0), and then a low sucrose concentration solution (1.0 M sucrose, 10 mM
HEPES-KOH pH 8.0) and suspension buffer were layered in this order. Thylakoid membrane
was floated at the interface between high sucrose concentration solution and low sucrose
concentration solution after centrifugation (at 20,000 × g, for 60 min, at 25°C). The recovered
thylakoid membrane was suspended in the suspension buffer.
The anti-VAR2 antibody (Sakamoto, 2003) was conjugated with magnetic beads
(Magnosphere^{TM}, MS 160/Tosyl, JSR life sciences, Japan) by the presence of the fully chemically
synthesized polymer (Blockmaster^{TM} CE210, JSR life sciences, Japan) according to the
instruction manual. The conjugated and blocked magnetic beads were suspended in the
suspension buffer (10 mM HEPES-KOH pH 8.0) after washing TBS-T. Prior to the incubation
with solubilized thylakoid membrane, the magnetic beads were resuspended in the suspension
buffer of which oxygen was removed enzymatically by incubating at 25°C for 60 minutes in the
presence of 100 mM glucose, 40 U/mL glucose oxidase, and 50 U/mL catalase.
Thylakoid membrane was solubilized sequentially; thylakoid membrane (1.0 mg
Chlorophyll/mL) was incubated with 1.0% (w/v) glyco-diosgenin (GDN) and subsequently n-

dodecyl-α-maltoside was added at 1.0% (w/v), and finally the mixture was diluted at twice volume

with suspension buffer. The solubilized material was incubated with FtsH conjugated magnetic

beads for 60 minutes at 4°C after removal of debris by centrifugation (at 20,000 × g, for 1 min, at

4°C). The beads were washed 6 times with suspension buffer containing 0.02% (w/v) GDN and

were incubated with elution buffer (125 mM Tris-HCl pH 6.8, 2% (w/v) Lithium Dodecyl sulfate,

0.1% (w/v) Sodium Dodecyl sulfate, and 25% (w/v) glycerol) for 60 minutes on ice. The eluted

sample was directly loaded on individual sample slot on SDS-PAGE to separate polypeptides.

All buffers except elution buffer were incubated at 25°C for 60 minutes in the presence of 100

mM glucose, 40 U/mL glucose oxidase, and 50 U/mL catalase to remove oxygen before chilling.

Molecular dynamics simulations of D1 N-term in PSII complex

The MD simulations for PSII were performed using the X-ray crystal structure determined at 1.9-

Å resolution (PDB: 3ARC)(Umena et al., 2011) and based on the same procedure described

previously (Sakashita et al., 2017b)(Sakashita et al., 2017a)(Kawashima et al., 2018), except for

the following points. To investigate the structural fluctuation of the N terminal region of the D1

subunit, we restructured the N-terminal region between D1-Met1 and D1-Ser10 that was lacking

in the crystal structure, using MOE program (2018). After structural optimization with positional

restraints on heavy atoms of the PSII assembly, the system was heated from 0.001 to 300 K over

5.0 ps, with a 0.05-fs time step. The positional restraints on heavy atoms were gradually released

over 16.5 ns. After an equilibrating MD run for 40 ns, a production run was conducted over 495

ns with an MD time step of 1.5 fs. The SHAKE algorithm was used for hydrogen constraints

(Ryckaert et al., 1977). The structure of the D1-W14F mutant was modeled from the crystal

structure of WT. The MD simulations were based on the AMBER-ff14SB force field for protein

residues and lipids (Maier et al., 2015). The water molecules were described by TIP3P model

(Jorgensen et al., 1983). For NFK, we employed the generalized Amber force field (GAFF)

parameter set (Wang et al., 2004). The atomic partial charges of NFK were determined by fitting

the electrostatic potential by using the RESP procedure (Bayly et al., 1993) (for calculated

charges, see Fig. S8). The electronic wave functions were calculated after geometry optimization

with the density functional theory of the B3LYP/6-31G** level by using JAGUAR (2013). MD

Simulations were conducted using the MD engine NAMD (Phillips et al., 2005). The atomic

fluctuation was calculated as the root mean square fluctuation (RMSF) of heavy atoms from the

averaged structure of PSII over the whole MD trajectory.

Acknowledgements

We would like to thank Rie Hijiya and Tsuneaki Takami for their technical assistance.

Funding

This work was supported by KAKENHI grants (23H04959 from the Ministry of Education, Culture, Sports, Science and Technology to W.S.; 21H02508 from the Japan Society for the Promotion of Science to W.S.; 18K06290 from the Japan Society for the Promotion of Science to Y.K.; 16H06560 from the Ministry of Education, Culture, Sports, Science and Technology to K.S.; 18H01186 from the Japan Society for the Promotion of Science to K.S.) and by the Oohara Foundation (to W.S.); CNRS and Sorbonne Université (basic support to UMR7141) and by the Initiative d'Excellence program (grant DYNAMO, Agence Nationale de la Recherche ANR-11-LABX-0011-01).

Additional information

Supplementary Table S1. Modification reported in Trp in PSII core proteins.

Supplementary Table S2. Trp oxidation in Chlamydomonas PSII core proteins.

Supplementary Figure S1. $^1O_2$-induced oxidative modifications at Trp14 of D1.

Supplementary Figure S2. $^1O_2$-induced oxidative modifications at Trp317 of D1.

Supplementary Figure S3. The positions of oxidized Trp residues in the identified peptide of PSII core complex by the MS-MS analysis.

Supplementary Fig.ure S4. Structural positions of oxidized Trp residues in PSII core proteins.

Supplementary Figure S5. High-light sensitive phenotype in the *Chlamydomonas* D1 transformants in which Trp-14 and Trp317 were mutated.

Supplementary Figure S6. Characterization of *Chlamydomonas* CP43 transformants in which Trp-353 and Trp-375 were mutated.

Supplementary Figure S7. Protein synthesis in the transformants studied by in vivo protein labeling.

Supplementary Figure S8. Atomic partial charges of NFK.

Supplementary Movie 1. Oxidized Trp residues assigned in the PSII dimer.

Supplementary Materials and Methods.

Correspondence and requests for materials should be addressed to W.S.

References

Adam Z, Frottin F, Espagne C, Meinnel T, Giglione C (2011) Interplay between N-terminal
methionine excision and FtsH protease is essential for normal chloroplast development
and function in Arabidopsis. Plant Cell **23**: 3745–3760

Anderson LB, Maderia M, Ouellette AJA, Putnam-Evans C, Higgins L, Krick T, MacCoss
MJ, Lim H, Yates JR, Barry BA (2002) Posttranslational modifications in the CP43
subunit of photosystem II. Proc Natl Acad Sci U S A **99**: 14676–81

Aro EM, Virgin I, Andersson B (1993) Photoinhibition of Photosystem II. Inactivation, protein
damage and turnover. BBA - Bioenerg **1143**: 113–134

Baena-González E, Aro EM, Styring S, Andersson B, Hundal T, Diner B, Wikstrom M,
Fromme P, Junge W (2002) Biogenesis, assembly and turnover of photosystem II units.
Philos Trans R Soc B Biol Sci **357**: 1451–1460

Bailey S, Thompson E, Nixon PJ, Horton P, Mullineaux CW, Robinson C, Mann NH
(2002) A critical role for the Var2 FtsH homologue of Arabidopsis thaliana in the
photosystem II repair cycle in vivo. J Biol Chem **277**: 2006–2011

Bayly CI, Cieplak P, Cornell WD, Kollman PA (1993) A well-behaved electrostatic potential
based method using charge restraints for deriving atomic charges: The RESP model. J
Phys Chem **97**: 10269–10280

Bennoun P, Spierer-Herz M, Erickson J, Girard-Bascou J, Pierre Y, Delosme M, Rochaix
JD (1986) Characterization of photosystem II mutants of Chlamydomonas reinhardii
lacking the psbA gene. Plant Mol Biol **6**: 151–160

Bujaldon S, Kodama N, Rappaport F, Subramanyam R, de Vitry C, Takahashi Y,
Wollman FA (2017) Functional accumulation of antenna proteins in chlorophyll b-less
mutants of Chlamydomonas reinhardtii. Mol Plant **10**: 115–130

Chen M, Choi Y, Voytas DF, Rodermel S (2000) Mutations in the Arabidopsis VAR2 locus
cause leaf variegation due to the loss of a chloroplast FtsH protease. Plant J **22**: 303–313

Dogra V, Li M, Singh S, Li M, Kim C (2019) Oxidative post-translational modification of
EXECUTER1 is required for singlet oxygen sensing in plastids. Nat Commun **10**: 2834

Dreaden Kasson TM, Rexroth S, Barry BA (2012) Light-Induced Oxidative Stress, N-
Formylkynurenine, and Oxygenic Photosynthesis. PLoS One **7**: e42220

Dreaden TM, Chen J, Rexroth S, Barry BA (2011) N-formylkynurenine as a marker of high
light stress in photosynthesis. J Biol Chem **286**: 22632–22641

Duan J, Lee KP, Dogra V, Zhang S, Liu K, Caceres-Moreno C, Lv S, Xing W, Kato Y,
Sakamoto W, et al (2019) Impaired PSII Proteostasis Promotes Retrograde Signaling via
Salicylic Acid. Plant Physiol **180**: 2182–2197

Ehrenshaft M, Deterding LJ, Mason RP (2015) Tripping up Trp: Modification of protein
tryptophan residues by reactive oxygen species, modes of detection, and biological
consequences. Free Radic Biol Med **89**: 220–228

Frankel LK, Sallans L, Limbach PA, Bricker TM (2012) Identification of Oxidized Amino Acid
Residues in the Vicinity of the Mn 4 CaO 5 Cluster of Photosystem II: Implications for the
Identification of Oxygen Channels within the Photosystem. Biochemistry **51**: 6371–6377
Fristedt R, Willig A, Granath P, Crèvecoeur M, Rochaix JD, Vener A V. (2009)
Phosphorylation of photosystem II controls functional macroscopic folding of
photosynthetic membranes in Arabidopsis. Plant Cell **21**: 3950–3964
Hanna Janska, Malgorzata Kwasniaka JS (2013) Protein quality control in organelles -
AAA/FtsH story. Biochim Biophys Acta - Mol Cell Res **1833**: 381–387
Ito K, Akiyama Y (2005) Cellular functions, mechanism of action, and regulation of FtsH
protease. Annu Rev Microbiol **59**: 211–231
Järvi S, Suorsa M, Aro EM (2015) Photosystem II repair in plant chloroplasts--Regulation,
assisting proteins and shared components with photosystem II biogenesis. Biochim
Biophys Acta **1847**: 900–909
Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML (1983) Comparison of
simple potential functions for simulating liquid water. J Chem Phys **79**: 926–935
Kale R, Hebert AE, Frankel LK, Sallans L, Bricker TM, Pospíšil P (2017) Amino acid
oxidation of the D1 and D2 proteins by oxygen radicals during photoinhibition of
Photosystem II. Proc Natl Acad Sci **114**: 2988–2993
Kasson TMD, Barry BA (2012) Reactive oxygen and oxidative stress: N-formyl kynurenine in
photosystem II and non-photosynthetic proteins. Photosynth Res **114**: 97–110
Kato Y, Hyodo K, Sakamoto W (2018) The Photosystem II Repair Cycle Requires FtsH
Turnover through the EngA GTPase. Plant Physiol **178**: 596–611
Kato Y, Miura E, Ido K, Ifuku K, Sakamoto W (2009) The variegated mutants lacking
chloroplastic FtsHs are defective in D1 degradation and accumulate reactive oxygen
species. Plant Physiol **151**: 1790–1801
Kato Y, Sakamoto W (2014) Phosphorylation of photosystem II core proteins prevents
undesirable cleavage of D1 and contributes to the fine-tuned repair of photosystem II.
Plant J **79**: 312–321
Kato Y, Sakamoto W (2018) FtsH protease in the thylakoid membrane: Physiological
functions and the regulation of protease activity. Front Plant Sci **9**: 1–8
Kato Y, Sakamoto W (2009) Protein quality control in chloroplasts: A current model of D1
protein degradation in the photosystem II repair cycle. J Biochem **146**: 463–469
Kato Y, Sun X, Zhang L, Sakamoto W (2012) Cooperative D1 degradation in the
photosystem II repair mediated by chloroplastic proteases in Arabidopsis. Plant Physiol
159: 1428–1439
Kawashima K, Takaoka T, Kimura H, Saito K, Ishikita H (2018) O2 evolution and recovery
of the water-oxidizing enzyme. Nat Commun **9**: 1247
Koivuniemi A, Aro EM, Andersson B (1995) Degradation of the D1- and D2-proteins of
photosystem II in higher plants is regulated by reversible phosphorylation. Biochemistry
34: 16022–16029

Komenda J, Tichý M, Prášil O, Knoppová J, Kuviková S, de Vries R, Nixon PJ (2007) The Exposed N-Terminal Tail of the D1 Subunit Is Required for Rapid D1 Degradation during Photosystem II Repair in Synechocystis sp PCC 6803. Plant Cell **19**: 2839–2854

Kong AT, Leprevost F V., Avtonomov DM, Mellacheruvu D, Nesvizhskii AI (2017) MSFragger: ultrafast and comprehensive peptide identification in mass spectrometry–based proteomics. Nat Methods **14**: 513–520

Krynická V, Shao S, Nixon PJ, Komenda J (2015) Accessibility controls selective degradation of photosystem II subunits by FtsH protease. Nat Plants **1**: 15168

Kulak NA, Pichler G, Paron I, Nagaraj N, Mann M (2014) Minimal, encapsulated proteomic-sample processing applied to copy-number estimation in eukaryotic cells. Nat Methods **11**: 319–324

Kuroda H, Kawashima K, Ueda K, Ikeda T, Saito K, Ninomiya R, Hida C, Takahashi Y, Ishikita H (2021) Proton transfer pathway from the oxygen-evolving complex in photosystem II substantiated by extensive mutagenesis. Biochim Biophys Acta - Bioenerg **1862**: 148329

Kuroda H, Kodama N, Sun X-Y, Ozawa S, Takahashi Y (2014) Requirement for Asn298 on D1 Protein for Oxygen Evolution: Analyses by Exhaustive Amino Acid Substitution in the Green Alga Chlamydomonas reinhardtii. Plant Cell Physiol **55**: 1266–1275

Li M, Kim C (2022) Chloroplast ROS and stress signaling. Plant Commun **3**: 100264

Lindahl M (2000) The thylakoid FtsH protease plays a role in the light-induced turnover of the photosystem II D1 protein. Plant Cell **12**: 419–432

Luber CA, Cox J, Lauterbach H, Fancke B, Selbach M, Tschopp J, Akira S, Wiegand M, Hochrein H, O'Keeffe M, et al (2010) Quantitative Proteomics Reveals Subset-Specific Viral Recognition in Dendritic Cells. Immunity **32**: 279–289

Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C (2015) ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. J Chem Theory Comput **11**: 3696–3713

Malnoe A, Wang F, Girard-Bascou J, Wollman F-A, de Vitry C (2014) Thylakoid FtsH protease contributes to photosystem II and cytochrome b6f remodeling in Chlamydomonas reinhardtii under stress conditions. Plant Cell **26**: 373–390

Michoux F, Ahmad N, Wei Z-Y, Belgio E, Ruban A V., Nixon PJ (2016) Testing the Role of the N-Terminal Tail of D1 in the Maintenance of Photosystem II in Tobacco Chloroplasts. Front Plant Sci **7**: 1–9

Miura E, Kato Y, Matsushima R, Albrecht V, Laalami S, Sakamoto W (2007) The Balance between Protein Synthesis and Degradation in Chloroplasts Determines Leaf Variegation in Arabidopsis yellow variegated Mutants. Plant Cell **19**: 1313–1328

Murata N, Takahashi S, Nishiyama Y, Allakhverdiev SI (2007) Photoinhibition of photosystem II under environmental stress. Biochim Biophys Acta - Bioenerg **1767**: 414–421

Ohnishi N, Allakhverdiev SI, Takahashi S, Higashi S, Watanabe M, Nishiyama Y, Murata

N (2005) Two-Step Mechanism of Photodamage to Photosystem II: Step 1 Occurs at the Oxygen-Evolving Complex and Step 2 Occurs at the Photochemical Reaction Center. Biochemistry **44**: 8494–8499

Ozawa S-I, Bald T, Onishi T, Xue H, Matsumura T, Kubo R, Takahashi H, Hippler M, Takahashi Y (2018) Configuration of Ten Light-Harvesting Chlorophyll a / b Complex I Subunits in Chlamydomonas reinhardtii Photosystem I. Plant Physiol **178**: 583–595

Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, Chipot C, Skeel RD, Kalé L, Schulten K (2005) Scalable molecular dynamics with NAMD. J Comput Chem **26**: 1781–1802

Rexroth S, Poetsch A, Rögner M, Hamann A, Werner A, Osiewacz HD, Schäfer ER, Seelert H, Dencher NA (2012) Reactive oxygen species target specific tryptophan site in the mitochondrial ATP synthase. Biochim Biophys Acta - Bioenerg **1817**: 381–387

Rinalducci S, Murgiano L, Zolla L (2008) Redox proteomics: basic principles and future perspectives for the detection of protein oxidation in plants. J Exp Bot **59**: 3781–3801

Rintamäki E, Kettunen R, Aro E-M (1996) Differential D1 Dephosphorylation in Functional and Photodamaged Photosystem II Centers. J Biol Chem **271**: 14870–14875

Ryckaert JP, Ciccotti G, Berendsen HJC (1977) Numerical integration of the cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes. J Comput Phys **23**: 327–341

Sakamoto W (2003) Coordinated regulation and complex formation of YELLOW VARIEGATED1 and YELLOW VARIEGATED2, chloroplastic FtsH metalloproteases involved in the repair cycle of photosystem II in Arabidopsis thylakoid membranes. Plant Cell **15**: 2843–2855

Sakamoto W, Tamura T, Hanba-Tomita Y, Sodmergen, Murata M (2002) The VAR1 locus of Arabidopis encodes a choloroplastic FtsH and is responsible for leaf variegation in the mutant alleles. Genes to Cells **7**: 769–780

Sakamoto W, Uno Y, Zhang Q, Miura E, Kato Y, Sodmergen (2009) Arrested differentiation of proplastids into chloroplasts in variegated leaves characterized by plastid ultrastructure and nucleoid morphology. Plant Cell Physiol **50**: 2069–2083

Sakashita N, Watanabe HC, Ikeda T, Ishikita H (2017a) Structurally conserved channels in cyanobacterial and plant photosystem II. Photosynth Res **133**: 75–85

Sakashita N, Watanabe HC, Ikeda T, Saito K, Ishikita H (2017b) Origins of Water Molecules in the Photosystem II Crystal Structure. Biochemistry **56**: 3049–3057

Schwanhäusser B, Busse D, Li N, Dittmar G, Schuchhardt J, Wolf J, Chen W, Selbach M (2011) Global quantification of mammalian gene expression control. Nature **473**: 337–342

Silva P, Thompson E, Bailey S, Kruse O, Mullineaux CW, Robinson C, Mann NH, Nixon PJ (2003) FtsH is involved in the early stages of repair of photosystem II in Synechocystis sp PCC 6803. Plant Cell **15**: 2152–2164

Takahashi Y, Utsumi K, Yamamoto Y, Hatano A, Satoh K (1996) Genetic Engineering of the Processing Site of D1 Precursor Protein of Photosystem II Reaction Center in

Chlamydomonas reinhardtii. Plant Cell Physiol **37**: 161–168

Takechi K, Sodmergen, Murata M, Motoyoshi F, Sakamoto W (2000) The YELLOW

VARIEGATED (VAR2) locus encodes a homologue of FtsH, an ATP-dependent protease

in Arabidopsis. Plant Cell Physiol **41**: 1334–46

Tyystjärvi E (2008) Photoinhibition of Photosystem II and photodamage of the oxygen

evolving manganese cluster. Coord Chem Rev **252**: 361–376

Umena Y, Kawakami K, Shen J-R, Kamiya N (2011) Crystal structure of oxygen-evolving

photosystem II at a resolution of 1.9 Å. Nature **473**: 55–60

Vass I (2012) Molecular mechanisms of photodamage in the Photosystem II complex. Biochim

Biophys Acta - Bioenerg **1817**: 209–217

Wang F, Qi Y, Malnoë A, Choquet Y, Wollman FA, de Vitry C (2017) The high light

response and redox control of thylakoid FtsH protease in Chlamydomonas reinhardtii. Mol

Plant **10**: 99–114

Wang J, Wolf RM, Caldwell JW, Kollman PA, Case DA (2004) Development and testing of a

general amber force field. J Comput Chem **25**: 1157–1174

Wiśniewski JR, Zougman A, Mann M (2009) Combination of FASP and StageTip-Based

Fractionation Allows In-Depth Analysis of the Hippocampal Membrane Proteome. J

Proteome Res **8**: 5674–5678

Yamamoto Y, Aminaka R, Yoshioka M, Khatoon M, Komayama K, Takenaka D,

Yamashita A, Nijo N, Inagawa K, Morita N, et al (2008) Quality control of photosystem

II: impact of light and heat stresses. Photosynth Res **98**: 589–608

Yi L, Liu B, Nixon PJ, Yu J, Chen F (2022) Recent Advances in Understanding the Structural

and Functional Evolution of FtsH Proteases. Front Plant Sci **13**: 1–16

Yu F, Park S, Rodermel SR (2004) The Arabidopsis FtsH metalloprotease gene family:

Interchangeability of subunits in chloroplast oligomeric complexes. Plant J **37**: 864–876

Yu F, Park S, Rodermel SR (2005) Functional redundancy of AtFtsH metalloproteases in

thylakoid membrane complexes. Plant Physiol **138**: 1957–1966

Zaltsman A, Ori N, Adam Z (2005) Two Types of FtsH Protease Subunits Are Required for

Chloroplast Biogenesis and Photosystem II Repair in Arabidopsis. Plant Cell **17**: 2782–

2790

Molecular Operating Environment (MOE), 2018.01; Chemical Computing Group ULC., 1010

Sherbooke St. West, Suite #910, Montreal, QC, Canada, H3A 2R7.

Jaguar, version 8.0 Schrödinger, LLC, New York.

Table 1. Trp oxidation in Chlamydomonas PSII core proteins

Accession	Protein	Sequence	Range	Modified Trp residue	Oxidation status
DAA00922.1_20	D1	ENSSL(W*)AR	9-16	Trp14	OIA, NFK, KYN
		FC^{cam}E(W*)ITSTENR	17-27	Trp20	OIA, NFK, KYN
		E(W*)WELSFR	130-136	Trp131	OIA, NFK, KYN
		VLNT(W*)ADIINR	313-323	Trp317	OIA, NFK, KYN
DAA00964.1_63	D2	T(W*)FDDADDWLR	13-23	Trp14	OIA, NFK, KYN
		TWFDDADD(W*)LR	13-23	Trp21	OIA, NFK, KYN
		T(W*)FDDADD(W*)LR	13-23	Trp14, Trp21	OIA, NFK, KYN
		A(W*)MAAQDQPHER	327-338	Trp328	OIA, NFK, KYN
		A(W*)M^{ox}AAQDQPHER	327-338	Trp328	OIA, NFK, KYN
DAA00966.1_65	CP43	DQETTGFA(W*)WSGNAR	15-29	Trp23	OIA, NFK, KYN
		DQETTGFAW(W*)SGNAR	15-29	Trp24	OIA, NFK, KYN
		DQETTGFA(W*)(W*)SGNAR	15-29	Trp23, Trp24	OIA, NFK, KYN
		AM^{ox}YFGGVYDT(W*)APGGGDVR	167-185	Trp177	OIA, NFK, KYN
		GP(W*)LEPLR	351-358	Trp353	OIA, NFK, KYN
		NDIQP(W*)QER	370-378	Trp375	OIA, NFK, KYN
DAA00933.1_31	CP47	YQ(W*)DQGFFQQEIQK	273-286	Trp275	OIA, NFK, KYN
		VQASLAEGASLSDA(W*)SR	288-304	Trp302	OIA, NFK, KYN
		TGAM^{ox}NSGDGIAVG(W*)LGHASFK	327-347	Trp340	OIA, NFK, KYN

Ccam, Cyc carbamidomethylation; W*, Trp oxidative modifications; Mox, Met oxidation.

Figure 1. The Oxidized Trp residues in PSII complex.

a, Trp-oxidation pathway. OIA: oxindolylalanine, NFK: N-formylkynurenine, KYN: Kynurenine. **b,** Multiple

alignment of D1 protein from *Arabidopsis*, spinach, *Chlamydomonas*, and *Thermosynechococcus vulcanus*,

showing oxidized Trp residues. Orange color boxes indicate the identified peptide by the MS-MS analysis. Oxidized Trp residues are highlighted in red. **c,** Oxidation levels of three oxidative variants of Trp in Trp14 and Trp317 containing peptides in *var2* and WT obtained by label-free MS analysis. The abundance of oxidized variants of Trp14 and Trp317 were calculated using the intensity values. Asterisks indicate statistically significant differences between the mean values (* < 0.05, ** < 0.01; Student's t-test). **d-e,** Structural positions of oxidized Trp residues in PSII core proteins. The side chain of oxidized Trp residues are shown with red-colored space-filling model and indicated with arrows. The P680 special chlorophyll pair is indicated with dark-green colored ball-stick model in panel e. PSII dimer (panel d) and monomer (panel e) from *Chlamydomonas reinhardtii* (PDB ID is 6KAC) is shown in cartoon model without cofactors Top view from stromal side or luminal side (**d**) and the side view from the dimer interface (**e**) are shown respectively. The color code of each subunit is, Green, D1; Dark blue, D2; Purple, CP43, Cyan, CP47. Protein structure graphics were generated with PyMOL ver. 2.4.0 software.

Figure 2. High-light sensitive phenotype in the *Chlamydomonas* D1 transformants in which Trp-14 and Trp317 were mutated.

a, Schematic drawing of the transforming vector carrying *psbA*, its flanking regions of the chloroplast DNA, and the selectable *aadA* marker cassette. E and X represent restriction sites of *EcoR*I and *Xho*I, respectively. Green boxes represent exons 1–5 of *psbA*. Fud7 is the *psbA* deletion mutant of *Chlamydomonas*. **b,** Phototrophic growth of Trp-substituted transformants on HSM medium and mixotrophic growth on TAP medium. GL, growth light (30 µmol photons m^{-2}s^{-1}): HL, high light (320 µmol photons m^{-2}s^{-1}). **c,** Chlorophyll fluorescence induction kinetics in Trp-substituted transformants. SP, saturating pulse. AL, actinic light. **d,** Protein accumulation in the transformants. Thylakoid proteins of cells grown in TAP medium under growth-light condition were separated by SDS-PAGE and analyzed by immunoblotting with antibodies against PSII subunits (D1, D2, and CP43), PSI subunits (PsaA), and light-harvesting complex of PSI (Lhca1). **e,** oxygen-evolving activity of the transformants.

Figure 3. D1 degradation assay in W14F and W317F transformants demonstrating enhanced D1 degradation under high-light stress.

The transformants were incubated under high-light (320 µmol photons m^{-2}s^{-1}) or growth-light (30 µmol photons m^{-2}s^{-1}) conditions in the absence or presence of inhibitor of chloroplast protein synthesis, CAM, and subjected to D1 degradation assay. **a,** growth-light in the absence of CAM; **b,** high-light in the absence of CAM; **c,** growth-light in the presence of CAM; **d,** high-light in the presence of CAM. Immunoblot results of D1 in the transformants are shown at the top of each panel. A representative immunoblot using anti-D1 is depicted. Quantified D1 levels using NIH Image program are shown in the middle. Values are means ± SD. Asterisks indicate statistically significant differences between the mean values (* < 0.05, ** < 0.01; Student's t-test). Time course analysis of maximal photochemical efficiency of PSII, Fv/Fm, are shown at the bottom.

Figure 4. D1 degradation assay in W14F and W14F/W317F transformants in the *ftsh* mutant background.

Rate of D1 degradation in the W14F *ftsH* and W14F/W317F *ftsH* was investigated as shown in Fig. 3. Cultured cells were incubated under high-light conditions (320 μmol photons m^{-2}s^{-1}) in the absence (**a**) or presence (**b**) of CAM. Signals of immunoblots were quantified using NIH Image program. Values are means ± SD (n = 4). A representative immunoblot using anti-D1 is depicted. Values are means ± SD. Asterisks indicate statistically significant differences between the mean values (* < 0.05, ** < 0.01; Student's t-test).

Figure 5. Snapshots and structural fluctuation of D1-Trp14 in molecular dynamics simulations of PSII.

a, The interaction between D1 Trp-14 and PsbI Ser-25. Dash line indicates the hydrogen bond between the side chains. **b,** Position change of side-chain when D1 Trp-14 is oxidized to N-formylkynurenine. **c,** Position change of side-chain when D1 Trp-14 is substituted to Phe. **d,** The fluctuation of atoms at D1 Trp-14 in the MD simulation. **e,** Cβ-Cβ distance between side chains of D1 Trp-14 and PsbI Ser-25. The Cβ atoms are indicated as red arrowheads in **a**.

Figure 6. Augmented affinity of FtsH with D1 by W14F/W317F.

a, Coimmunoprecipitation was performed with anti-FtsH antibody using the thylakoid membrane isolated from control or D1-W14F/W317F. The polypeptides of thylakoid membrane or coimmunoprecipitated samples were separated by SDS-PAGE and detected by immunoblotting with anti-D1, anti-D2, and anti-FtsH antibody. **b**, The immunoblotting signals are quantified and the ratio of D1 or D2 to FtsH are calculated. The averaged value and standard error for three biological replicates are shown. Significant difference was calculated by t-test and 0.1 or 0.05 probability confidence were indicated respectively.**c**, A proposed model of photodamaged D1 recognition, in which Trp oxidation plays a role in recruiting FtsH. FtsH heterocomplexes (blue) and PSII core proteins (green) along with oxygen evolving protein complex (gray) in the thylakoid membrane are schematically shown. Trp-oxidized residues (red) are localized at both luminal and stromal sides. Trp-14 located at the N-terminus alpha helix enhances association of FtsH, whose catalytic site faces stroma.

Supplementary Figure S1 $^1O_2$-induced oxidative modifications at Trp14 of D1. a Trp-oxidation pathway. b Mass spectra of Trp14 carrying peptide [9]ESESL(W)GR[16] of D1 protein in *var2*. This oxidation led to the formation of oxindolylalanine (OIA), N-formylkynurenine (NFK), and kynurenine (KYN) with +16, +32, and +4 mass shifts, respectively.

Supplementary Figure S2. $^1O_2$-induced oxidative modifications at Trp317 of D1. a Trp-oxidation pathway. b Mass spectra of Trp317 carrying peptide [313]VINT(W)ADIINR[323] of D1 protein in *var2*. This oxidation led to the formation of oxindolylalanine (OIA), N-formylkyrnurenine (NFK), and kynurenine (KYN) with +16, +32, and +4 mass shifts, respectively.

Supplementary Figure S3. (Continued to next page)

c

Supplementary Figure S3. Positions of oxidized Trp residues in the identified peptide of PSII core complex by the MS-MS analysis.

The oxidized Trp residues in D2 (a), CP43(b), and CP47 (c) were highlighted. Orange color boxes indicate the identified peptide by the MS-MS analysis. Oxidized Trp residues are highlighted in red.

Supplementary Fig. S4. Structural positions of oxidized Trp residues in PSII core proteins.

The structure is from *Thermosynecoccocus vulcanus* (PDB id is 3WU2) and shown in cartoon model without cofactors by PyMOL ver. 2.4.0. The four PSII core subunits are colored in green (D1), marine blue (D2), pink (CP43), and cyan (CP47). The oxidized Trp residues are indicated by red. Side view (**a**) and top view (**b**) of the structure are respectively shown. Green, D1; Dark blue, D2; Purple, CP43, Cyan, CP47.

Supplementary Figure S5. High-light sensitive growth phenotype in the *Chlamydomonas* D1 transformants in which Trp-14 and Trp317 were mutated. Phototrophic growth of Trp-substituted transformants on HSM medium. GL, growth light (30 μmol photons $m^{-2}s^{-1}$): HL, high light (320 μmol photons $m^{-2}s^{-1}$).

Supplementary Figure S6. Characterization of *Chlamydomonas* CP43 transformants in which Trp-353 and Trp-375 were mutated.

a, Phototrophic growth on HSM medium and mixotrophic growth on TAP medium at growth light (GL) at 30 µmol m^{-2}s^{-1} or high light, (HL) at 320 µmol m^{-2}s^{-1}. **b**, Protein accumulation in the transformants. Thylakoid proteins of cells grown in TAP medium under growth light condition were separated by SDS-PAGE and analyzed by immunoblotting with antibodies against PSII subunits (D1 and CP43).

Supplementary Figure S7. Protein synthesis in the transformants studied by in vivo protein labeling.
Cells were radio-labeled in vivo with ^{35}S, in the presence of cycloheximide for 1, 2, and 4 min. Total proteins were separated by SDS-PAGE. The bands corresponding to D1 is indicated by arrowheads. Quantified newly synthesyzed D1 levels using the Image J program are shown in bottom panels. To normalize values from four independent experiments, the ratio of control at 4 min was adjusted as 1, and the relative ratios are indicated. Values are means ± SD.

Supplementary Figure S8. Atomic partial charges of NFK for MD simulations.

Red values represent atomic partial charges calculated by using the RESP procedure.

Supplementary Movie 1. Oxidized Trp residues assigned in the PSII dimer (Movie is available online).