1

Ulk4 promotes Shh signaling by regulating Stk36 ciliary localization and Gli2 phosphorylation

8
9
10

Mengmeng Zhou[1], Yuhong Han[1], and Jin Jiang[1,2,*]

[1]Department of Molecular Biology, [2]Department of Pharmacology, University of Texas Southwestern Medical Center, Dallas, TX 75390, USA

*Corresponding authors: jin.jiang@utsouthwestern.edu

Running title: Ulk4 promotes Gli phosphorylation and Hh signaling

Summary

The Hedgehog (Hh) family of secreted proteins governs embryonic development and adult tissue homeostasis through the Gli family of transcription factors. Gli is thought to be activated at the tip of primary cilium, but the underlying mechanism has remained poorly understood. Here, we show that Unc-51-like kinase 4 (Ulk4), a pseudokinase and a member of the Ulk kinase family, acts in conjunction with another Ulk family member Stk36 to promote Gli2 phosphorylation and Hh pathway activation. Ulk4 interacts with Stk36 through its N-terminal region containing the pseudokinase domain and Gli2 via its regulatory domain to bridge the kinase and substrate. Although dispensable for Hh-induced Stk36 kinase activation, Ulk4 is essential for Stk36 ciliary tip localization, Gli2 phosphorylation and activation. In response to Hh, both Ulk4 and Stk36 colocalize with Gli2 at ciliary tip, and Ulk4 and Stk36 depend on each other for their ciliary tip accumulation. We further show that ciliary localization of Ulk4 depends on Stk36 kinase activity and phosphorylation of Ulk4 on Thr1023, and that ciliary tip accumulation of Ulk4 is essential for its function in the Hh pathway. Taken together, our results suggest that Ulk4 regulates Hh signaling by promoting Stk36-mediated Gli2 phosphorylation and activation at ciliary tip.

Introduction

Pseudokinases, which comprise ~10% of the mammalian kinome, play essential

biological functions yet the mechanism of action for many pseudokinases has remained

unknown (Jacobsen and Murphy 2017). One such pseudokinase is Unc-51-like kinase 4

(Ulk4), which is a member of Ulk kinase family found in most eukaryotes (Luo et al.

2022). The Ulk family kinases, which include Ulk1 (Atg1 in yeast) and Ulk2 involved in

autophagy initiation, contain a conserved kinase domain in their N-terminal regions and

a variable regulatory domain in their C-terminal regions (Luo et al. 2022). However,

unlike other Ulk family members, Ulk4 contains a pseudokinase domain at its N-

terminus and is predicted to be catalytically inactive (Preuss et al. 2020). Ulk4 plays a

key role in neurogenesis in mice by regulating neural stem cell pool and neurite

branching morphogenesis (Lang et al. 2014; Liu et al. 2016a). In addition, Ulk4

regulates the integrity of white matter by promoting myelination (Liu et al. 2018b).

Recent studies have implicated Ulk4 as a risk factor for neuropsychiatric disorders

including schizophrenia (Lang et al. 2014; Lang et al. 2016; Liu et al. 2018a; Luo et al.

2022). However, the signaling pathways in which Ulk4 is involved and the substrates it

regulates have remained unknown. A recent proteome study has revealed that Ulk4

interacts with Stk36 in additional to many other proteins (Preuss et al. 2020). Stk36 is

also a member of the Ulk family and mammalian homolog of *Drosophila* Fused (Fu), a

Ser/Thr kinase implicated in Hedgehog (Hh) signaling (Murone et al. 2000; Wilson et al.

2009; Han et al. 2019). A recent study identified Ulk4 as a genetic modifier of the

holoprosencephaly phenotype caused by impaired Shh pathway activity (Mecklenburg

et al. 2021). In addition, this study showed that Ulk4 was localized to the primary cilia

and that Ulk4 knockdown affected Shh induced *Gli-luc* reporter gene expression and

Smoothened (Smo) ciliary localization (Mecklenburg et al. 2021). These observations

have raised an interesting possibility that Ulk4 may participate in Hh signal transduction.

The Hh family of secreted proteins governs embryonic development and adult tissue

homeostasis and deregulation of Hh signaling activity has been implicated in a myriad

of human disorders including birth defect and cancer (Villavicencio et al. 2000; Jiang

and Hui 2008; Briscoe and Therond 2013; Jiang 2021). Hh exerts its biological influence

via an evolutionarily conserved signaling cascade that culminates in the activation of Gli

family of Zn-finger transcription factors. The core Hh signal reception system consists of

two multi-span transmembrane proteins Patched1 (Ptch1) and Smoothened (Smo that

transduce vertebrate Hh signal at primary cilia (Goetz and Anderson 2010). In quiescent

cells, Ptch1 localizes to primary cilia where Smo is barely detectable (Corbit et al. 2005;

Rohatgi et al. 2007); the full-length Gli (Gli^F) undergoes ubiquitin/proteasome-mediated

proteolytic processing to generate a truncated transcriptional repressor form (Gli^R)

(Wang et al. 2000; Jiang 2002; Tempe et al. 2006; Wang and Li 2006; Chen and Jiang

2013). Hh inhibits Ptch1 activity and promotes its ciliary exit, allowing Smo and Gli

proteins to accumulate in the cilia to transduce the Hh signal (Corbit et al. 2005;

Haycraft et al. 2005; Rohatgi et al. 2007; Chen et al. 2009; Tukachinsky et al. 2010;

Chen et al. 2011). Activated Smo blocks the proteolytic processing of Gli into Gli^R and

converts Gli^F to an active form (Gli^A) that enters the nucleus to turn on the expression of

Hh target genes (Wang et al. 2000; Kim et al. 2009; Tukachinsky et al. 2010; Han et al.

2019; Arveseth et al. 2021).

Although it is generally thought that Smo converts Gli^F into Gli^A at ciliary tip (Goetz

and Anderson 2010), the underlying mechanism has remained an enigma. Our recent

studies have demonstrated that two Ulk family kinases Ulk3 and Stk36 play a role in

converting Gli^F into Gli^A by directly phosphorylating Gli2 at multiple sites in mammalian

cells (Han et al. 2019; Zhou et al. 2022; Zhou and Jiang 2022). Here, we provide

evidence that Ulk4 acts in conjunction with Stk36 to promote Gli2 phosphorylation and

Hh pathway activation. We show that Ulk4 interacts with Stk36 through its N-terminal

region containing the pseudokinase domain and with Gli2 via its C-terminal regulatory

domain to bring Stk36 and Gli2 into proximity. We find that Ulk4 is not required for Stk36

kinase activation by Shh but is essential for Stk36 ciliary localization and Gli2

phosphorylation. We show that, in response to Shh stimulation, both Ulk4 and Stk36

colocalize with Gli2 at the tips of primary cilia, and that Ulk4 and Stk36 depend on each

other for their ciliary tip accumulation. We further show that ciliary tip localization of Ulk4

depends on Stk36-mediated phosphorylation of Ulk4 on Thr1023, and that ciliary tip

localization of Ulk4 is essential for its function in Hh signal transduction.

107
108
109   **Results**
110
111
Ulk4 acts in conjunction with Stk36 to promote Gli2 phosphorylation and Hh

pathway activation

To determine whether Ulk4 is required for Hh pathway activation, we depleted Ulk4 by

RNA interference (RNAi) in NIH3T3 cells, a well-established system for studying Hh

signaling. We found that Ulk4 depletion significantly decreased Shh-induced expression

of *Gli1* and *Ptch1* determined by RT-qPCR, which was rescued by the expression of

human Ulk4 (hUlk4; Figure 1A-B; Figure 1-figure supplement 1). The requirement of

Ulk4 in Shh-induced expression of *Gli1* and *Ptch1* was confirmed in MEF cells (Figure

1-figure supplement 2).

Our previous study revealed that Stk36 and Ulk3, which are Ulk family kinases,

act in parallel to promote Gli2 phosphorylation and Hh pathway activation (Han et al.

2019). To determine the relationship between Ulk4 and Ulk3/Stk36, we carried out

double RNAi experiments. We found that double knockdown (KD) of Ulk4 and Ulk3

further decreased Shh target gene expression compared with single KD of either Ulk4

or Ulk3 whereas double KD of Ulk4 and Stk36 did not have an additive effect (Figure

1C-D; Figure 1-figure supplement 2). These results indicated that Ulk4 acts in the same

pathway with Stk36 but in parallel with Ulk3 to promote Hh signaling.

Our previous study demonstrated that Ulk3 and Stk36 phosphorylate Gli2 at

multiple sites including S230/232 to promote Gli2 activation (Han et al. 2019; Zhou et al.

2022). We found that Ulk4 KD reduced Gli2 phosphorylation on S230/232 stimulated by

Shh in NIH3T3 cells, which was determined by western blot analysis using a phospho-

specific antibody that recognizes pS230/232 (Han et al. 2019; Han and Jiang 2021),

and that expression of hUlk4 restored Gli2 phosphorylation on S230/232 in Ulk4

depleted cells (Figure 1E-E'). Consistent with Ulk4 acting in a liner pathway with Stk36

but in parallel with Ulk3, double KD of Ulk4 and Stk36 decreased S230/232

phosphorylation similarly to single KD of either Ulk4 or Stk36 whereas double KD of

Ulk3 and Ulk4 further decreased Shh-induced S230/232 phosphorylation compared with

single KD of Ulk3 or Ulk4 (Figure 1F-G'). Taken together, these results demonstrated

that Ulk4 and Stk36 act in a linear pathway but in parallel with Ulk3 to participate in Hh

transduction by facilitating Gli2 phosphorylation and activation (Figure 1H).

143
144
145   **Ulk4 forms a complex with Stk36 and Gli2**
146
147
A previous study using proximity-dependent biotin identification (BioID) identified Stk36

as a proximity interactor (Preuss et al. 2020). To verify that Stk36 formed a complex

with Ulk4, we carried out coimmunoprecipitation (Co-IP) experiments and found that a

C-terminal HA-tagged Ulk4 (hUlk4-HA) interacted with a N-terminal CFP-tagged Stk36

(CFP-Stk36) in HEK293T cells (Figure 2A). To explore whether Ulk4 interacted with Gli2

and Sufu, a Gli binding partner and Hh pathway inhibitor (Dunaeva et al. 2003; Cooper

et al. 2005; Han et al. 2015), we co-expressed hUlk4-HA, with a Myc-tagged Gli2 (Myc-

Gli2) and a Flag-tagged Sufu (Fg-Sufu) in HEK293T cells. Co-IP experiments showed

that both Myc-Gli2 and Fg-Sufu were pulled down by Ulk4-HA (Figure 2B), suggesting

that Ulk4 could form a complex with Stk36 and Gli2-Sufu.

Ulk4 contains an N-terminal pseudokinase domain and the C-terminal regulatory

domain with armadillo (Arm) repeats (Figure 2C) (Preuss et al. 2020). We generated

two truncated forms of Ulk4: hUlk4N (aa 1-450) that contains the pseudokinase domain

and hUlk4C (451-1275) that contains the C-terminal regulatory domain. Co-IP

experiments showed that hUlk4N-HA interacted with CFP-Stk36 but not with Myc-

Gli2/Fg-Sufu whereas hUlk4C-HA formed a complex with Myc-Gli2/Fg-Sufu but not with

CFP-Stk36 (Figure 2D-E), suggesting that Ulk4 interacted with Stk36 and Gli2-Sufu

through its N-terminal pseudokinase domain and C-terminal regulatory domain,

respectively. Consistent with our observation, previous BioID experiments using BirA*-

Ulk4 pseudokinase domain also identified Stk36 as a proximity binding partner of the

Ulk4 pseudokinase domain (Preuss et al. 2020).

Ulk4 is dispensable for Shh-induced Stk36 kinase activation
171
172
Pseudokinases could regulate the enzymatic activity of interacting kinases, as has been

shown for LKB1 kinase activation by its binding to the pseudokinases STRADα and

STARADβ (Baas et al. 2003). To determine whether Ulk4 regulates Stk36 enzymatic

activity, we carried out an *in vitro* kinase assay using GST-Gli2N, which contains the

Ulk3/Stk36 phosphorylation site S230, as substrate and either Fg-Stk36 or Fg-

Stk36/hUlk4-HA complex immunopurified from HEK293T cells as kinase. GST-Gli2N

containing S230A substitution was used as a negative control. As shown in Figure 3A,

Stk36 and Stk36/Ulk4 phosphorylated GST-Gli2N at similar levels and the

phosphorylation was abolished by the S230A mutation. This result suggests binding of

Ulk4 to Stk36 does not modulate Stk36 kinase activity.

We next determined whether Ulk4 is required for Stk36 kinase activation by Shh.

In *Drosophila*, Hh activates Fu by promoting its kinase domain dimerization and trans-

autophosphorylation on multiple S/T residues in its activation loop (AL) including T151,

T154, T158 and S159 (Shi et al. 2011; Zhang et al. 2011; Zhou and Kalderon 2011).

Interestingly, the AL phosphorylation sites as well as the neighboring residues are

highly conserved between Fu and Stk36 (Figure 3B) (Shi et al. 2011), raising the

possibility that Stk36 could be activated by Shh through a similar mechanism. Indeed,

we found that Shh induced phosphorylation of Stk36 on T158/S159 in NIH-3T3 cells as

determined by a phospho-specific antibody that recognized pT158/pS159 (Figure. 3C)

(Shi et al. 2011). To determine whether Shh-induced T158/S159 phosphorylation

depends on Stk36 kinase activity, we introduced either wild type Stk36 (CFP-Stk36^{WT})

or its kinase dead form (CFP-Stk36^{KR}) into NIH3T3 cells with endogenous Stk36 and

Ulk3 deleted (*Ulk3^{KO} Stk36^{KO}*) by CRISPR (Han et al. 2019). As shown in Figure 3D,

Shh induced phosphorylation of CFP-Stk36^{WT} but not CFP-Stk36^{KR} on T158/S159,

suggesting that Shh promotes the phosphorylation of Stk36 AL via its kinase activity,

likely through trans-autophosphorylation as are the cases for other kinases. Hence,

phosphorylation of T158/S159 can be used as a readout for Stk36 kinase activation.

To determine whether AL phosphorylation promotes Stk36 kinase activation, we

converted both S151 and T154 to E (EE) to mimic phosphorylation or to A (AA) to block

phosphorylation on these sites (Figure 3B). We found that the phospho-mimetic

mutation (Stk36^{EE}) increased whereas phospho-deficient mutation (Stk36^{AA}) abolished

Stk36 kinase activity as determined by both Stk36 T158/S159 phosphorylation and Gli2

S230/S232 phosphorylation (Figure3 E-H), which is similar to what we have observed

previously for Fu (Shi et al. 2011), suggesting that Hh stimulates kinase activation of

Stk36 and Fu through a similar mechanism.

To determine whether Ulk4 is required for Shh-induced Stk36 kinase activation,

we employed RNAi to deplete Ulk4 from NIH3T3 cells that express CFP-Stk36. As

shown in Figure 3I, depletion of Ulk4 did not affect Shh-induced phosphorylation of

CFP-Stk36 on T158/S159, suggesting that Ulk4 is not required for Stk36 kinase activation in response to Shh stimulation.

Ulk4 mediates the interaction between Stk36 and Gli2

Another possible mechanism for Ulk4 to influence Gli2 phosphorylation by Stk36 is to function as a scaffold that brings Stk36 and Gli2 into the same protein complex. Indeed, we found that Ulk4 depletion diminished the association between exogenously expressed Stk36 and Gli2 in HEK293T cells as determined by Co-IP experiments (Figure 4A). By domain mapping, we found that the kinase domain of Stk36 (Stk36N: aa1-260) but not its regulatory domain (Stk36C: aa261-1351) interacted with Ulk4 (Figure 4B-C). Consistent with Ulk4 mediating the interaction between Stk36 and Gli2, we found that Stk36N formed a complex with Gli2 more effectively than Stk36C (Figure 4D).

We next determined whether Ulk4 mediated Stk36/Gli2 interaction in NIH3T3 cells and whether this was regulated by Shh. NIH3T3 cells were infected with lentiviruses expressing low levels of a Flag-tagged Stk36 (Fg-Stk36) and Myc-Gli2 and treated with or without Shh. Co-IP experiments showed that Fg-Stk36 and Myc-Gli2 formed a complex regardless of Shh treatment (Figure 4E; lands 1-2). The interaction between Fg-Stk36 and Myc-Gli2 was diminished by Ulk4 RNAi (Figure 4E; lands 3-4). Reintroducing full-length hUlk4 (WT) but not hUlk4N or hUlk4C restored Fg-Stk36/Myc-Gli2 association in Ulk4-depleted cells (Figure 4E; lands 5-10). Taken together, these results suggest that Ulk4 functions as a molecular scaffold to bring Stk36 and Gli2 into

236 the same complex with its N-terminal region binding to Stk36 and its regulatory domain

237 interacting with Gli2 (Figure 4F).

239 Shh stimulates ciliary tip accumulation of both Ulk4 and Stk36

In response to Shh, Gli2 and Sufu accumulated at the tip of primary cilium in mammal

cells (Haycraft et al. 2005; Chen et al. 2009; Tukachinsky et al. 2010), leading to the

general view that Gli2 is converted to Gli2^A at the ciliary tip (Goetz and Anderson 2010;

Jiang 2021). Indeed, our previous study showed that Shh-stimulated Gli2

phosphorylation by Stk36/Ulk3 depended on ciliary localization of Gli2 (Han et al. 2019).

These observations imply that Stk36/Ulk3 may translocate to the primary cilium to

phosphorylate Gli2 in response to Shh. To test this hypothesis, we treated NIH3T3 cells

stably expressing an N-terminal Myc-tagged Stk36 (Myc-Stk36) with or without Shh and

examined ciliary localization of Stk36 by immunostaining with an anti-Myc antibody.

Ciliary localization of endogenous Gli2 was examined by immunostaining with an anti-

Gli2 antibody. As shown in Figure 5A-B and Figure 5-figure supplement 1A1-B2, Myc-

Stk36 (green signals) and Gli2 (blue signals) were barely detectable in primary cilia in

quiescent cells; however, Shh treatment induced ciliary tip localization of both Myc-

Stk36 and Gli2.

To determine whether Shh induced ciliary localization of Ulk4, NIH3T3 cells

stably expressing hUlk4-HA was treated with or without Shh, followed by

immunostaining with anti-HA and anti-Gli2 antibodies. Like Stk36, hUlk4-HA was barely

detectable in primary cilia in the absence of Shh but accumulated to the tips of primary

cilia and colocalized with Gli2 after Shh stimulation (Figure 5C-D; Figure 5-figure

supplement 1C1-D2). We also examined NIH3T3 cells simultaneously expressing Myc-

Stk36 and hUlk4-HA and found that Shh induced the ciliary tip co-localization of Ulk4

and Stk36 (Figure 5E-F; Figure 5-figure supplement 1E1-F2). We conformed these

results in MEF cells (Figure 5-figure supplement 1G1-J2). These observations suggest

that Shh induces ciliary tip accumulation of Stk36-Ulk4-Gli2 protein complexes to

promote Gli2 phosphorylation and activation.

Ulk4 and Stk36 depend on each other for their ciliary localization.
268
As pseudokinases could regulate the subcellular localization of their interacting kinases,

we asked whether Ulk4 is required for ciliary localization of Stk36 in response to Shh.

NIH3T3 cells stabling expressing Myc-Stk36 were depleted of Ulk4 by shRNA and

treated with and without Shh. As shown in Figure 6A-B, Ulk4 RNAi diminished Shh-

induced ciliary tip localization of Myc-Stk36, suggesting that Ulk4 is required for ciliary

localization of Stk36 in response to Shh.

We next asked whether Ulk4 ciliary localization depends on Stk36. We first

determined the ciliary location of Ulk4 in NIH3T3 cells with both Stk36 and Ulk3

knocked out (DKO) by the CRISPR-Cas9 system (Han et al. 2019). As shown in Figure

6C-D, Shh promoted ciliary tip enrichment of Ulk4-HA in the control cells but not in the

Stk36 and Ulk3 DKO cells. Adding back wild type Stk36 (Stk36^{WT}) but not its kinase

inactive form (Stk36^{AA}) restored Shh-induced Ulk4-HA ciliary tip accumulation in DKO

cells (Figure 6E-F). By contrast, neither wild type Ulk3 (Ulk3^{WT}) nor its kinase dead form

(Ulk3^{KR}) could rescue ciliary tip accumulation of Ulk4-HA in DKO cells (Figure 6E-F).

These results suggest that ciliary tip localization of Ulk4 depends on Stk36 kinase

activity.

Ulk4 ciliary localization is regulated by its C-terminal phosphorylation

Many pseudokinases are also the substrates of their interacting kinases (Kung and Jura

2019). To determine whether Ulk4 is a Stk36 substrate and whether Stk36-mediated

phosphorylation of Ulk4 promotes its ciliary tip localization, we sought out to identify the

relevant phosphorylation sites. We generated several C-terminally HA-tagged truncated

forms of hUlk4 to narrow down the region essential for ciliary tip localization of Ulk4

(Figure 7A). These truncated forms were expressed in NIH3T3 cells in which

endogenous Ulk4 depleted by RNAi. Deleting the entire regulatory domain abolished

Shh-induced ciliary tip localization of the Ulk4 variant (hUlk4$_{1-450}$; Figure 7B-C). Deleting

the Stk36 binding region from Ulk4 (hUlk4$_{451-1275}$) also abolished its ciliary tip

localization in response to Shh, suggesting that binding to Stk36 is essential for Ulk4

ciliary localization. We generated two small deletions from the C-terminus of hUlk4

(hUlk4$_{1-1000}$ and hUlk4$_{1-1200}$; Figure 7A). Whereas hUlk4$_{1-1200}$ still localized to the ciliary

tip in response to Shh, hUlk4$_{1-1000}$ failed to do so (Figure 7B-C), suggesting that the

region between aa1000-1200 is essential for Shh-induced hUlk4 ciliary tip localization.

Within this region, T1021 and T1088/T1090 fall into the Fu/Ulk3/Stk36 phosphorylation

consensus site: S/T [X$_5$] E/D (Figure 7D) (Han et al. 2019; Zhou and Jiang 2022).

Furthermore, phosphorylation on T1021 can prime CK1-mediated phosphorylation on

T1023 as T1023 falls into the CK1 phosphorylation consensus site: D/E/$_{(p)}$S/$_{(P)}$T[X$_{1-3}$]

S/T (Knippschild et al. 2005).

To determine whether phosphorylation on these sites may regulate Ulk4 ciliary

localization, we generated hUlk4 variants carrying T1021A/1023A (hUlk4$^{T1021A/1023A}$) or

T1088A/T1090A (hUlk4$^{T1088A/T1090A}$) substitutions. These hUlk4 variants were expressed

in NIH3T3 cells with endogenous Ulk4 depleted by shRNA. We found that

hUlk4$^{T1088A/T1090A}$ but not hUlk4$^{T1021A/1023A}$ accumulated at ciliary tip in response to Shh

stimulation (Figure 7E-F), suggesting phosphorylation on T1021/1023 is essential for

Shh-induced ciliary tip localization of Ulk4.

To confirm T1021 phosphorylation by Stk36, we carried out an *in vitro* kinase

assay using immunopurified Fg-Stk36 as kinase and GST fusion protein containing

hUlk4 sequence from aa1017 to aa1036 (GST-hUlk4C) as substrate. As shown in

Figure 7G, immunopurified wild type Stk36 (Fg-Stk36^{WT}) but not its kinase inactive

variant (Fg-Stk36^{AA}) phosphorylated GST-hUlk4C. Mutating both T1021 and T1023 to A

(T1021A /T1023A) abolished the phosphorylation; however, mutating T1021A reduced

but not eliminated the phosphorylation, suggesting that in the absence of T1021

phosphorylation, T1023 could also be phosphorylated by Stk36. Interestingly, T1023 but

not T1021 is conserved in mouse Ulk4 (mUlk4) and mUlk4 contains an Ala at T1021

(Figure 7H), implying that phosphorylation of T1021 might be dispensable for Ulk4

function. Indeed, we found that mutating T1021 to A in hUlk4 (hUlk4^{T1021A}) did not affect

its ciliary tip accumulation in response to Shh (Figure 7E-F) as well as its activity in the

Hh pathway (see below).

326
327
328   **Ciliary localization of Ulk4 is essential for its function in Shh signaling**
329

To determine the importance of ciliary localization of Ulk4 for its function in Shh signal

transduction, we compared the Hh pathway related activity of ciliary localization

deficient hUlk4$^{T1021A/1023A}$ with that of ciliary localization competent hUlk4^{T1021A}

(equivalent to mUlk4^{WT}) and hUlk4^{WT} in NIH3T3 cells with endogenous Ulk4 depleted by

shRNA. We found that both hUlk4^{T1021A} and hUlk4^{WT} rescued Shh-induced ciliary tip

localization of coexpressed Myc-Stk36 (Figure 8A-B), phosphorylation of Myc-Gli2 at

S230/S232 (Figure 8C-C'), and the expression of Shh target genes *Ptch1* and *Gli1*

(Figure 8D-E). By contrast, hUlk4$^{T1021A/1023A}$ failed to rescue Shh-stimulated Stk36 ciliary

tip localization, Gli2 phosphorylation, as well as *Ptch1* and *Gli1* expression in Ulk4

depleted cells (Figure 8A-E). These results suggest that ciliary localization of Ulk4 is

essential for its ability to promote Stk36 ciliary localization, Gli2 phosphorylation, and Hh

pathway activation (Figure 8F).

Discussion
344
The Gli family of transcription factors mediates the transcriptional output of the Hh

signaling pathway in species ranging from *Drosophila* to mammals but how Gli is

activated by Hh signaling is still poorly understood, especially in vertebrates where Gli

activation is thought to occur at the tip of primary cilium. We recently found that the

Fu/Ulk family kinases directly phosphorylate Ci and Gli2 on multiple sites to promote

their activation (Han et al. 2019; Zhou et al. 2022; Zhou and Jiang 2022). In mammalian

cells, the Ulk family kinases Ulk3 and Stk36 (mammalian homolog of Fu) act in parallel

to promote Gli2 phosphorylation and activation in response to Shh (Han et al. 2019). In

addition, Gli2 phosphorylation depends on its ciliary localization (Han et al. 2019). In this

study, we found that Stk36 accumulated to the tip of primary cilium and colocalized with

Gli2 in response to Shh stimulation, consistent with Gli2 being converted into Gli2^A at

the ciliary tip. In addition, we identified another Ulk family member Ulk4, which is a

pseudokinase, as a mediator of Shh-induced Gli2 phosphorylation. We found that Ulk4

and Stk36 act in the same pathway but in parallel with Ulk3 to promote Gli2

phosphorylation and activation in NIH3T3 and MEF cells. The positive role of Ulk4 in

Shh signaling could explain why elevated expression of Ulk4 in certain genetic

background could suppress the heart outflow tract defects and holoprosencephaly

caused by partially reduced Shh signaling activity in LRP2-deficient mice (Mecklenburg

et al. 2021). However, it remains to be determined whether Ulk4 participates in Hh

signaling *in vivo* by loss-of-function study. Because knockout of Stk36 in mice did not

affect Hh signaling during development, likely due to functional compensation by other

kinases such as Ulk3 (Chen et al. 2005; Merchant et al. 2005), we do not expect that

loss-of-function of Ulk4 will affect Hh signaling during development. Therefore, double

knockout of Ulk4 (or Stk36) and Ulk3 will be necessary to address the physiological role

of the Fu/Ulk family members in mammalian Hh signaling. In addition, it will also be

important to determine whether Ulk4/Stk36 and Ulk3 could be differentially required for

Hh signal transduction in different cell types/tissues or in different contexts, e.g.,

development vs. tissue homeostasis or physiological vs pathological settings.

To determine the mechanism by which Ulk4 assists Stk36 to phosphorylate Gli2,

we explored whether Ulk4 regulated 1) the enzymatic activity of Stk36; 2) Shh-induced

Stk36 kinase activation; 3) the association of Stk36 and its substrate; and 4) ciliary

localization Stk36. In an *in vitro* kinase assay, we found that immunopurified Stk36

phosphorylated GST-Gli2N equally well regardless of whether it interacted with Ulk4 or

not, suggesting that Ulk4 does not regulate Stk36 kinase activity. Rather, we found that

Ulk4 is required for Stk36 to bind to its substrate Gli2. Ulk4 interacts with Stk36 through

its N-terminal region that contains the pseudokinase domain and with Gli2 via its C-

terminal regulatory domain. Depletion of Ulk4 diminished the physical association

between Stk36 and Gli2. As a further support of the notion that Gli2 is brought to Stk36

through Ulk4, we found that the kinase domain of Stk36, which binds Ulk4, also

interacted with Gli2 whereas its C-terminal region interacted with neither Ulk4 nor Gli2.

Of note, despite being a pseudokinase, Ulk4 contains an unusual ATP binding site that

might regulate its ability to bind other proteins such as Stk36 (Preuss et al. 2020).

Therefore, it would be interesting to determine in the future whether ATP binding to Ulk4

could regulate Shh signaling or other processes that Ulk4 is involved.

389   Although Stk36/Ulk4/Gli2 complex formation appears to be constitutive

regardless of the presence or absence of Shh, Stk36 kinase activation depends on Shh

stimulation. Like Fu, Stk36 is phosphorylated on multiple sites in the activation loop of

its kinase domain in response to Shh likely through trans-autophosphorylation, and

activation loop phosphorylation is essential for its kinase activation. Using a phospho-

specific antibody that recognized pT158/pS159 in the kinase activation loop of

Fu/Stk36, we found that Ulk4 depletion did not affect Shh-induced phosphorylation of

Stk36 on T158/S159, a readout for Stk36 kinase activation, suggesting that Ulk4 is

dispensable for Shh signaling to activate Stk36. In *Drosophila*, the kinesin-like protein

Costal2 (Cos2) mediates the activation of Fu kinase by Hh. Therefore, it would be

interesting to determine whether mammalian homologs Cos2, Kif7 and/or Kif27,

participates in Stk36 kinase activation in response to Shh stimulation.

Hh signaling depends on primary cilia in vertebrates. Upon Shh stimulation, key

pathway components such as Smo and Gli proteins accumulate in primary cilia, and

Gli2 is accumulated to the ciliary tip and thought to be converted into its activator form

(Gli2^A) in this ciliary compartment. Consistent with this notion, we found that Shh

induced ciliary accumulation of both Ulk4 and Stk36 that colocalized with Gli2 at the

ciliary tip. Furthermore, we found that Ulk4 is required for ciliary tip accumulation of

Stk36. On the other hand, ciliary tip accumulation of Ulk4 depends on Stk36 kinase

activity and phosphorylation of its C-terminal region on T1023. Preventing Ulk4

phosphorylation on T1023 abolished its ciliary tip accumulation and its ability to promote

Stk36 ciliary tip accumulation, Gli2 phosphorylation and Hh pathway activation. These

results suggest that ciliary tip accumulation of Ulk4 is critical for its function in Hh signal

transduction, likely by increasing the local concentration of Stk36 and Gli2 to promote

effective phosphorylation of Gli2. It would also be important to examine the localization

of endogenous Ulk4 and Stk36 in the future as exogenously expressed epitope-tagged

protein might cause artifacts. In addition, the precise mechanism and dynamics of Ulk4

ciliary targeting remain to be determined in the future. e.g., it would be important to

determine how phosphorylation of T1023 promotes ciliary accumulation of Ulk4. A

recent study identified several motor proteins of the kinesin family (KIF1B, KIF3A,

KIF3B, and KAP3) as the interaction partners of Ulk4 (Preuss et al. 2020). KIF3A,

KIF3B and KAP3 are components of kinesin-2 motor that act in conjunction with

intraflagellar transport (IFT) complexes to transport cargoes into primary cilia (Goetz

and Anderson 2010). It would be interesting to determine whether Ulk4 T1023

phosphorylation affects its interaction with KIF3A, KIF3B, KAP3, or components in the

IFT complexes. Of note, a previous study suggested that Ulk4 regulated the ciliary

localization of Smo although the underlying mechanism was not explored (Mecklenburg

et al. 2021). Therefore, it remains possible that Ulk4 could regulate mammalian Hh

signaling at multiple steps similar to what Fu does in the *Drosophila* Hh pathway (Claret

et al. 2007; Liu et al. 2007).

Both Ulk4 and Stk36 have functions outside the Hh pathway. Stk36 is required

for motile ciliogenesis, and its loss of function affects the construction of central pair

apparatus of "9+2" motile cilia in both zebrafish and mice (Wilson et al. 2009; Nozawa et

al. 2013). Ulk4 malfunction also affects motile ciliogenesis of ependymal cells in mutant

mice, leading to hydrocephalus (Vogel et al. 2012; Liu et al. 2016b). A recent study

demonstrated that Ulk4 and Stk36 physically interact to mediate the assembly of motile

flagella in the flagellated protist *Leishmania Mexicana* (McCoy et al. 2023). However,

the biochemical mechanism by which Ulk4/Stk36 regulates motile ciliogenesis has

remained unknown. Ulk4 could regulate Stk36 ciliary localization and/or kinase-

substrate interaction like what we have discovered for their role in Shh signaling. The

relevant substrates for Stk36 in motile ciliogenesis remain to be determined. Knowing

the phosphorylation consensus sites for the Fu/Ulk family kinase may facilitate the

identification of Stk36 substrates involved in the assembly of motile cilia and flagella.

Initially identified Fu/Ulk3/Stk36 phosphorylation sites in Ci/Gli and Cos2 fall into the

consensus: $S/T(X)_5D/E$, where the acidic residue at +6 position is essential (Han et al.

2019; Zhou and Jiang 2022). However, subsequent studies suggested that the spacing

between the phospho-acceptor site and the acid residue appears to be flexible. For

example, a Ulk3/Stk36 phosphorylation site in the C-terminal region of Gli2 has a

spacing of six residues between the phospho-acceptor site and the acid residue (Zhou

et al. 2022). Here we found that a Stk36 phosphorylation site in the C-terminal region of

Ulk4 contains only four residues between the phospho-acceptor site and the acid

residue. Based on these findings, we propose an expanded consensus site for the

Fu/Ulk family kinases as $S/T(X)_{4-6}D/E$. It would be interesting to determine whether this

phosphorylation consensus sequence could be extended to other members of the Ulk

kinase family such as Ulk1/Atg1 and Ulk2, which are the initiating kinases for autophagy

(Zachari and Ganley 2017).

Ulk4 has been implicated in several human diseases including hypertension and

psychiatric disorder such as schizophrenia (Luo et al. 2022). Ulk4 also plays a role in

neurogenesis in mice by regulating neural stem cell pool and neurite branching

morphogenesis (Lang et al. 2014; Lang et al. 2016; Liu et al. 2016a). In addition, Ulk4

regulates the integrity of white matter by promoting myelination (Liu et al. 2018b). How

Ulk4 participates in these diverse processes has remained unclear. A recent study

revealed that Ulk4 binds several other kinases and phosphatases such as DAPK3,

LMTK2, PTPN14 as well as proteins implicated in the regulation of neuronal

differentiation and axonal regeneration including CAMSAP family proteins CAMSAP1/3

(Pongrakhananon et al. 2018; Preuss et al. 2020). Given our findings regarding how

Ulk4 regulates Hh signaling, it is tempting to speculate that Ulk4 may also function as a

scaffold to bridge kinases or other enzymes to their substrates and/or regulate the

subcellular localization of the interacting proteins in other biological processes.

Materials and methods

DNA constructs

Myc-Gli2 and GST-Gli2 were described previously (Han et al. 2019). All GST-fusion (the GST-tagged human Ulk4C^{WT}, Ulk4C^{T1021A}, Ulk4C$^{T1021A/1023A}$) proteins were subcloned into the *pGEX 4T-1* vector using EcoRI and XhoI sites. N-terminally CFP-tagged mouse Ulk3^{WT}, Ulk3^{KR}, N-terminally 6XMyc-tagged human Stk36, N-terminally 3XFg-tagged human Stk36^{WT}, Stk36^{AA} and the C-terminally 3XHA-tagged Human Ulk4^{WT}, Ulk4$^{1-451}$, Ulk4$^{452-1275}$, Ulk4$^{1-1000}$, Ulk4$^{1-1200}$, Ulk4$^{T1021/1023A}$, Ulk4$^{T1088/1090A}$, Ulk4^{T1021A} were subcloned into the *FUGW* vector with the EcoRI and BamHI sites. N-terminally 3XFg-tagged human Stk36^{WT}, Stk36^{EE}, Stk36^{AA}, Stk36$^{1-260}$, Stk36$^{261-1315}$, and C-terminally 3XHA-tagged Human Ulk4^{WT}, Ulk4$^{1-451}$, Ulk4$^{452-1275}$ were subcloned in the *PLVX* vectors using the EcoRI and MluI sites. The Ulk4$^{T1021/1023A}$, Ulk4$^{T1088/1090A}$, Ulk4^{T1021A}, Stk36^{EE}, Stk36^{AA}, were generated by PCR-mediated site-directed mutagenesis.

Cell culture, transfection, and lentiviral production

NIH3T3 cells were cultured in Dulbecco's modified Eagle's medium (DMEM, Sigma) containing10% bovine calf serum (Gibco) and 1% penicillin/streptomycin (Sigma). The transfection was carried out by using the GenJet Plus *in vitro* DNA transfection kit according to the manufacturer's instruction (SignaGen). HEK293T cells were cultured in DMEM (Sigma) supplemented with 10% fetal bovine serum (Gibco) and 1% penicillin/streptomycin (Sigma). PolyJet *in vitro* DNA transfection kit (SignaGen) was

used to do the transfection according to the manufacturer's instruction. Mouse

embryonic fibroblast cells (MEFs) were cultured in DMEM (Sigma) containing 10% FBS

(Gibco), and 1% streptomycin / penicillin (Sigma) according to the standard protocol.

For Shh-N treatment, cells were cultured at 50%-confluent densities in plate or chamber

slides for one day and starved in serum-free medium (0.5% BCS for NIH3T3 cells or

0.5% FBS for MEF cells) for 12 hours to allow ciliation. The recombinant human Shh N-

terminal fragment (1 ng/ml; R&D Systems, #8908-SH-005) was then added to the same

serum-free medium for 12 hours or overnight.

Immunostaining, immunoprecipitation, and western blotting analysis

The protocol for Ulk4-HA/Myc-Stk36 immunostaining was carried out as previously

described (Han et al. 2019). Immunofluorescence imaging was captured using a Zeiss

LSM700 laser scanning microscope. For ciliary Myc-Stk36/Ulk4-HA quantification, the

cilia were labelled with the acetylated-Tubulin and the Myc-Stk36/Ulk4-HA fluorescence

intensities were measured in the Myc-Stk36/Ulk4-HA image channel by ImageJ. Twenty

cells were randomly selected and calculated for each group. For immunoprecipitation

assay, after transfection for 48 hours, cells were washed twice with PBS and then lysed

on ice for 10 min with lysis buffer containing 1M Tris pH8.0, 5M NaCl, 1MNaF, 0.1M

Na3VO4, 1% CA630, 10% glycerol, and 0.5M EDTA (pH 8.0). Cell lysates were

incubated with protein A-Sepharose beads (Thermo scientific) for 1 hr at 4°C to

eliminate nonspecific binding proteins. After removal of the protein-A beads by

centrifugation, the cleared lysates were incubated with Myc (HA or Flag) antibody for 4

h or overnight. Protein complexes were collected by incubation with protein A-

Sepharose beads for 1 hr at 4°C, followed by centrifugation. Immunoprecipitates were

washed three times for 5 min each with lysis buffer and were separated on SDS-PAGE.

Western blot was carried out using standard protocol. Western blot quantification was

carried out by employing fluorogenic secondary antibody (IRDye 680LT Goat anti-

Mouse:926-68020 or Goat anti-Rabbit: 926-68021) and the blot was scanned and

quantified with an Odyssey CLx imaging system (LI-COR). Antibodies used in this study

were as follows: Mouse anti-Myc (Santa Cruz Biotechnology; 9E10), Mouse anti-Flag

(Sigma; M2), Mouse anti-HA (Santa Cruz Biotechnology; F7), Mouse anti-acetylated

Tubulin (Sigma; T7451; staining:1:1000), Rabbit anti-Myc (Abcam; ab9106; staining

1:2000), Rabbit anti-HA (AbCam; ab9110; staining 1:500), Goat anti-Gli2 (R&D system;

AF3635; staining 1:1000).

***In vitro* kinase assay**

In vitro kinase assay was performed by incubating 25 µL of reaction mixtures containing

150 mM Tris-HCl (pH 7.5), 0.2 mM Mg2+/ATP, 1 µg of purified GST-Ulk4/Gli2 fusion

proteins, together with appropriate amount of kinase for 30mins at 30°C. The reaction

was terminated by adding 2× SDS loading buffer. The resultant samples were load on

SDS-PAGE and subjected to pIMAGO phospho-protein detection kit with fluor-680

(Sigma) detection kit and LI-COR Odyssey platform for western blot analyses of

phosphorylated proteins (Millipore Sigma). Fg-Stk36^{WT} and Fg-Stk36^{AA} were

immunopurified from HEK293 cells transfected with the Fg-Stk36^{WT} or Fg-Stk36^{AA}

construct.

Viral Infection and shRNA

For lentivirus generation, *pLKO.1* vectors expressing shRNA or *FUGW* vectors

expressing transgenes were co-transfected with package vectors (PSPAX2 and

PMDG2) into HEK293T cells using PolyJet (SignaGen). The supernatants were

collected after 48 h and 72 h transfection, filtered through a 0.45 mM filter and

centrifuged at 20,000 g for 2 h. The sediment containing the viruses was re-suspended,

aliquoted in small volume of culture medium and stored at -80°C for future use. For cell

infection, lentiviruses were added to 70% confluent wild type NIH3T3 or MEF cells

together with Polybrene (Sigma, Cat. No. H9268) for overnight to make the stable

NIH3T3 or MEF cell lines expressing transgene or shRNA. The Mouse Ulk4 shRNA

lentiviral vector was purchased from Sigma (TRCN0000328268 for Ulk4 knockdown in

NIH3T3 cells and MEFs;TRCN0000002203 for Ulk4 knockdown in HEK293 cells). The

Non-Target shRNA Plasmid DNA (sigma; SHC016) was used as control.

RNA interference

For RNAi experiments in NIH3T3 or MEF cells, the siRNAs were purchased from

Sigma. The sequences for Ulk3 silencing were: 5'-GUGUACAAGGCCUACGCCA-3'

(Cat# SASI_ Mm02 _ 00444409) and 5'-CUGAGAAGGUGGCCCGUGU-3' (Cat#

SASI_Mm02_00444410). The sequences for Stk36 silencing were: 5'-

GGUAUACUGGCUUCAGAAA-3' (Cat# SASI_ Mm02 _00345637) and 5'-

GCCUUAUGUGCUUUGCUGU-3' (Cat# SASI_Mm01_00167751). The target sequence

of Mouse Ulk4 shRNA is: CTGCGAAGATTATCGAGAATG. The target sequence of

Human Ulk4 shRNA is: CCACTAGGTCACTCTTTCAGA. The

sequences for negative control were: 5'-UUCUCCGAACGUGUCACG U-3'. The RNAi

MAX reagent (Invitrogen Cat. No. 13778150) was used for the transfection of siRNA.

The knockdown efficiency was validated via real-time PCR.

Quantitative RT-PCR

Total RNA was extracted from 1×10^6 cells using the RNeasy Plus Mini Kit (Qiagen),

and cDNA was synthesized with High-Capacity cDNA Reverse Transcription Kit

(Applied Biosystems; 01279158) and qPCR was performed using Fast SYBR Green

Master Mix (Applied Biosystems; 2608088) and a Bio-Rad CFX96 real-time PCR

system (BIO-RAD; HSP9601). The RT-qPCR was performed in triplicate for each of

three independent biological replicates. Quantification of mRNA levels was calculated

using the comparative CT method. Primers: Gli1, 5'-GTGCACGTTTGAAGGCTGTC-3'

and 5'-GAGTGGGTCCGATTCTGGTG-3'; Ptch1, 5'-GAAGCCACAGAAAACCCTGTC-3'

and 5'-GCCGCAAGCCTTCTCTACG-3'; Ulk3, 5'- ACGAAACATCTCTCACTTG-3' and

5'- TGCTGGGCAAAGCCAAAGTC-3'; Ulk4, 5'- ATGCAGAGTGTGATTGCGTTG-3' and

5'- GAGTGGCAGTTTCTGTGAACA-3'; Stk36, 5'- CGCATCCTACACCGAGATATGA-3'

and 5'- AAATCCAAAGTCACAGAGCTTGA-3';GAPDH, 5'-

GTGGTGAAGCAGGCATCTGA-3' and 5'-GCCATGTAGGCCATGAGGTC-3'.

Acknowledgements

We thank Bing Wang, Xu Li, and Yong Cho for technical assistance, Pancheng Xie and

Yi Liu for reagents. J. Jiang is a Eugene McDermott Endowed Scholar in Biomedical

Science at the University of Texas Southwestern.

583
584 **Declaration of interests**
585
586 The authors declare no competing interests.

Reference

Arveseth CD, Happ JT, Hedeen DS, Zhu JF, Capener JL, Klatt Shaw D, Deshpande I, Liang J, Xu J, Stubben SL et al. 2021. Smoothened transduces Hedgehog signals via activity-dependent sequestration of PKA catalytic subunits. *PLoS biology* **19**: e3001191.

Baas AF, Boudeau J, Sapkota GP, Smit L, Medema R, Morrice NA, Alessi DR, Clevers HC. 2003. Activation of the tumour suppressor kinase LKB1 by the STE20-like pseudokinase STRAD. *Embo Journal* **22**: 3062-3072.

Briscoe J, Therond PP. 2013. The mechanisms of Hedgehog signalling and its roles in development and disease. *Nat Rev Mol Cell Biol* **14**: 418-431.

Chen MH, Gao N, Kawakami T, Chuang PT. 2005. Mice deficient in the fused homolog do not exhibit phenotypes indicative of perturbed hedgehog signaling during embryonic development. *Mol Cell Biol* **25**: 7042-7053.

Chen MH, Wilson CW, Li YJ, Law KK, Lu CS, Gacayan R, Zhang X, Hui CC, Chuang PT. 2009. Cilium-independent regulation of Gli protein function by Sufu in Hedgehog signaling is evolutionarily conserved. *Genes Dev* **23**: 1910-1928.

Chen Y, Jiang J. 2013. Decoding the phosphorylation code in Hedgehog signal transduction. *Cell Res* **23**: 186-200.

Chen Y, Sasai N, Ma G, Yue T, Jia J, Briscoe J, Jiang J. 2011. Sonic Hedgehog dependent phosphorylation by CK1alpha and GRK2 is required for ciliary accumulation and activation of smoothened. *PLoS biology* **9**: e1001083.

Claret S, Sanial M, Plessis A. 2007. Evidence for a novel feedback loop in the Hedgehog pathway involving Smoothened and Fused. *Curr Biol* **17**: 1326-1333.

Cooper AF, Yu KP, Brueckner M, Brailey LL, Johnson L, McGrath JM, Bale AE. 2005. Cardiac and CNS defects in a mouse with targeted disruption of suppressor of fused. *Development* **132**: 4407-4417.

Corbit KC, Aanstad P, Singla V, Norman AR, Stainier DY, Reiter JF. 2005. Vertebrate Smoothened functions at the primary cilium. *Nature* **437**: 1018-1021.

Dunaeva M, Michelson P, Kogerman P, Toftgard R. 2003. Characterization of the physical interaction of Gli proteins with SUFU proteins. *J Biol Chem* **278**: 5116-5122.

Goetz SC, Anderson KV. 2010. The primary cilium: a signalling centre during vertebrate development. *Nat Rev Genet* **11**: 331-344.

Han Y, Jiang J. 2021. Expression and purification of fused kinase from insect cells for in vitro kinase assay. *STAR Protoc* **2**: 100376.

Han Y, Shi Q, Jiang J. 2015. Multisite interaction with Sufu regulates Ci/Gli activity through distinct mechanisms in Hh signal transduction. *Proc Natl Acad Sci U S A* **112**: 6383-6388.

Han Y, Wang B, Cho YS, Zhu J, Wu J, Chen Y, Jiang J. 2019. Phosphorylation of Ci/Gli by Fused Family Kinases Promotes Hedgehog Signaling. *Dev Cell* **50**: 610-626 e614.

Haycraft CJ, Banizs B, Aydin-Son Y, Zhang Q, Michaud EJ, Yoder BK. 2005. Gli2 and gli3 localize to cilia and require the intraflagellar transport protein polaris for processing and function. *PLoS Genet* **1**: e53.

Jacobsen AV, Murphy JM. 2017. The secret life of kinases: insights into non-catalytic signalling functions from pseudokinases. *Biochemical Society transactions* **45**: 665-681.

Jiang J. 2002. Degrading Ci: who is Cul-pable? *Genes & Dev* **16**: 2315-2321.

Jiang J. 2021. Hedgehog signaling mechanism and role in cancer. *Semin Cancer Biol*.

Jiang J, Hui CC. 2008. Hedgehog signaling in development and cancer. *Dev Cell* **15**: 801-812.

Kim J, Kato M, Beachy PA. 2009. Gli2 trafficking links Hedgehog-dependent activation of
Smoothened in the primary cilium to transcriptional activation in the nucleus. *Proc Natl*
Acad Sci U S A **106**: 21666-21671.

Knippschild U, Gocht A, Wolff S, Huber N, Lohler J, Stoter M. 2005. The casein kinase 1 family:
participation in multiple cellular processes in eukaryotes. *Cellular signalling* **17**: 675-689.

Kung JE, Jura N. 2019. Prospects for pharmacological targeting of pseudokinases. *Nat Rev Drug*
Discov **18**: 501-526.

Lang B, Pu J, Hunter I, Liu M, Martin-Granados C, Reilly TJ, Gao GD, Guan ZL, Li WD, Shi YY et al.
2014. Recurrent deletions of ULK4 in schizophrenia: a gene crucial for neuritogenesis
and neuronal motility. *J Cell Sci* **127**: 630-640.

Lang B, Zhang L, Jiang G, Hu L, Lan W, Zhao L, Hunter I, Pruski M, Song NN, Huang Y et al. 2016.
Control of cortex development by ULK4, a rare risk gene for mental disorders including
schizophrenia. *Scientific reports* **6**: 31126.

Liu M, Fitzgibbon M, Wang Y, Reilly J, Qian X, O'Brien T, Clapcote S, Shen S, Roche M. 2018a.
Ulk4 regulates GABAergic signaling and anxiety-related behavior. *Transl Psychiatry* **8**: 43.

Liu M, Guan Z, Shen Q, Flinter F, Dominguez L, Ahn JW, Collier DA, O'Brien T, Shen S. 2016a.
Ulk4 Regulates Neural Stem Cell Pool. *Stem Cells* **34**: 2318-2331.

Liu M, Guan Z, Shen Q, Lalor P, Fitzgerald U, O'Brien T, Dockery P, Shen S. 2016b. Ulk4 Is
Essential for Ciliogenesis and CSF Flow. *J Neurosci* **36**: 7589-7600.

Liu M, Xu P, Guan Z, Qian X, Dockery P, Fitzgerald U, O'Brien T, Shen S. 2018b. Ulk4 deficiency
leads to hypomyelination in mice. *Glia* **66**: 175-190.

Liu Y, Cao X, Jiang J, Jia J. 2007. Fused-Costal2 protein complex regulates Hedgehog-induced
Smo phosphorylation and cell-surface accumulation. *Genes Dev* **21**: 1949-1963.

Luo S, Zheng N, Lang B. 2022. ULK4 in Neurodevelopmental and Neuropsychiatric Disorders.
Front Cell Dev Biol **10**: 873706.

McCoy CJ, Paupelin-Vaucelle H, Gorilak P, Beneke T, Varga V, Gluenz E. 2023. ULK4 and
Fused/STK36 interact to mediate assembly of a motile flagellum. *Mol Biol Cell*:
mbcE22060222.

Mecklenburg N, Kowalczyk I, Witte F, Gorne J, Laier A, Mamo TM, Gonschior H, Lehmann M,
Richter M, Sporbert A et al. 2021. Identification of disease-relevant modulators of the
SHH pathway in the developing brain. *Development* **148**.

Merchant M, Evangelista M, Luoh SM, Frantz GD, Chalasani S, Carano RA, van Hoy M, Ramirez J,
Ogasawara AK, McFarland LM et al. 2005. Loss of the serine/threonine kinase fused
results in postnatal growth defects and lethality due to progressive hydrocephalus. *Mol*
Cell Biol **25**: 7054-7068.

Murone M, Luoh SM, Stone D, Li W, Gurney A, Armanini M, Grey C, Rosenthal A, de Sauvage FJ.
2000. Gli regulation by the opposing activities of fused and suppressor of fused. *Nat Cell*
Biol **2**: 310-312.

Nozawa YI, Yao E, Lin C, Yang JH, Wilson CW, Gacayan R, Chuang PT. 2013. Fused (Stk36) is a
ciliary protein required for central pair assembly and motile cilia orientation in the
mammalian oviduct. *Dev Dyn* **242**: 1307-1319.

Pongrakhananon V, Saito H, Hiver S, Abe T, Shioi G, Meng W, Takeichi M. 2018. CAMSAP3
maintains neuronal polarity through regulation of microtubule stability. *Proceedings of*
the National Academy of Sciences **115**: 9750-9755.
Preuss F, Chatterjee D, Mathea S, Shrestha S, St-Germain J, Saha M, Kannan N, Raught B,
Rottapel R, Knapp S. 2020. Nucleotide Binding, Evolutionary Insights, and Interaction
Partners of the Pseudokinase Unc-51-like Kinase 4. *Structure* **28**: 1184-1196 e1186.
Rohatgi R, Milenkovic L, Scott MP. 2007. Patched1 regulates hedgehog signaling at the primary
cilium. *Science* **317**: 372-376.
Shi Q, Li S, Jia J, Jiang J. 2011. The Hedgehog-induced Smoothened conformational switch
assembles a signaling complex that activates Fused by promoting its dimerization and
phosphorylation. *Development* **138**: 4219-4231.
Tempe D, Casas M, Karaz S, Blanchet-Tournier MF, Concordet JP. 2006. Multisite protein kinase
A and glycogen synthase kinase 3beta phosphorylation leads to Gli3 ubiquitination by
SCFbetaTrCP. *Mol Cell Biol* **26**: 4316-4326.
Tukachinsky H, Lopez LV, Salic A. 2010. A mechanism for vertebrate Hedgehog signaling:
recruitment to cilia and dissociation of SuFu-Gli protein complexes. *Journal Of Cell*
Biology **191**: 415-428.
Villavicencio EH, Walterhouse DO, Iannaccone PM. 2000. The sonic hedgehog-patched-gli
pathway in human development and disease. *Am J Hum Genet* **67**: 1047-1054.
Vogel P, Read RW, Hansen GM, Payne BJ, Small D, Sands AT, Zambrowicz BP. 2012. Congenital
hydrocephalus in genetically engineered mice. *Vet Pathol* **49**: 166-181.
Wang B, Fallon JF, Beachy PA. 2000. Hedgehog-regulated processing of Gli3 produces an
anterior/posterior repressor gradient in the developing vertebrate limb. *Cell* **100**: 423-
434.
Wang B, Li Y. 2006. Evidence for the direct involvement of {beta}TrCP in Gli3 protein processing.
Proc Natl Acad Sci U S A **103**: 33-38.
Wilson CW, Nguyen CT, Chen MH, Yang JH, Gacayan R, Huang J, Chen JN, Chuang PT. 2009.
Fused has evolved divergent roles in vertebrate Hedgehog signalling and motile
ciliogenesis. *Nature* **459**: 98-102.
Zachari M, Ganley IG. 2017. The mammalian ULK1 complex and autophagy initiation. *Essays*
Biochem **61**: 585-596.
Zhang Y, Mao F, Lu Y, Wu W, Zhang L, Zhao Y. 2011. Transduction of the Hedgehog signal
through the dimerization of Fused and the nuclear translocation of Cubitus interruptus.
Cell Res **21**: 1436-1451.
Zhou M, Han Y, Wang B, Cho YS, Jiang J. 2022. Dose-dependent phosphorylation and activation
of Hh pathway transcription factors. *Life Sci Alliance* **5**.
Zhou M, Jiang J. 2022. Gli Phosphorylation Code in Hedgehog Signal Transduction. *Front Cell*
Dev Biol **10**: 846927.
Zhou Q, Kalderon D. 2011. Hedgehog Activates Fused through Phosphorylation to Elicit a Full
Spectrum of Pathway Responses. *Dev Cell* **20**: 802-814.
713
714
715

Figure legend

Figure 1. Ulk4 acts in conjunction with Stk36 to promote Shh-mediated Gli activation

(A-B) Relative *Gli1* (A) and *Ptch1* (B) mRNA levels measured by RT-qPCR in NIH3T3 cells expressing the indicated shRNA and a human Ulk4 (hUlk4) lentiviral construct and treated with or without Shh-N. Of note, hUlk4 is insensitive to RNAi, which targets mouse Ulk4.

(C-D) Relative *Ptch1* mRNA levels measured by RT-qPCR in NIH3T3 cells treated with the indicated RNAi and with or without Shh-N.

(E-E') Western blot analysis (E) and quantification (E') of Myc-Gli2 phosphorylation in NIH3T3 cells expressing the indicated shRNA and hUlk4 lentiviral construct and treated with or without Shh-N.

(F-G') Western blot analysis (F and G) and quantification (F' and G') of Myc-Gli2 phosphorylation in NIH3T3 cells with the indicated RNAi in the presence or absence of Shh-N treatment.

(H) Schematic diagram showing the functional relationship among Ulk4, Stk36 and Ulk3 in Shh-induced Gli activation.

The cells (A-G) were firstly starved in serum-free medium for 12 hours then cultured in the same medium with or without Shh-N fragment for another 12 hours before they are subjected into RNA preparation (A-D) or Western blot analysis (E-G). Data in (A-D) are mean ± SD from three independent experiments. Data in (E', F', G') are mean ± SD from two independent experiments. ***$p < 0.001$ (Student's t test). NS: not significant.

Figure 2. Ulk4 forms a complex with Stk36 and Gli2

(A) Ulk4 forms a complex with Stk36. HEK293T cells were transfected with hUlk4-HA

and CFP-Stk36 constructs, followed by Co-IP and western blot analyses with the

indicated antibodies.

(B) Ulk4 forms a complex with Gli2 and Sufu. HEK293T cells were transfected with

hUlk4-HA, Myc-Gli2 and Fg-Sufu constructs, followed by Co-IP and western blot

analyses with the indicated antibodies.

(C) Ulk4 domain structure and deletion mutants used for Co-IP experiments.

(D-E) Ulk4 interacted with Stk36 through its N-terminal domain (D) and with Myc-

Gli2/Fg-Sufu through its C-terminal region (E). HEK293T cells were transfected with the

indicated hUlk4, Fg-Stk36, Myc-Gli2 and Fg-Sufu constructs, followed by Co-IP and

western blot analyses.

Figure 3. Ulk4 is dispensable for Stk36 kinase activation.

(A) *In vitro* kinase assay using the immunopurified Fg-Stk36 or Fg-Stk36/hUlk4-HA as

kinase and GST-Gli2N^{WT} or GST-Gli2N^{S230A} as substrate. pIMAGO was used to detect

GST-Gli2N phosphorylation. The levels of Fg-Stk36/hUlk4-HA and GST/GST-Gli2N

were analyzed by western blot and coomassie blue (CB) staining, respectively.

(B) The schematic diagram showing the sequence alignment of the activation segments

of *Drosophila* Fu, mouse Stk36 (mStk36) and human Stk36 (hStk36). The conserved

Ser/Thr residues in the activation loop are color-coded in green. The residues used as

antigen for developing pT158/pS159 are colored in yellow. Amino acid substitutions for

Stk36 variants are indicated.

(C) Shh induced Stk36 phosphorylation on T158/S159. NIH3T3 cells expressing CFP-

Stk36 lentiviral construct were treated with or without Shh-N, followed by

IP with an GFP antibody and western blot analysis using an antibody against the

phosphorylated T158/S159 (pT158/pS159).

(D) Shh induced Stk36 phosphorylation on T158/S159 depending on Stk36 kinase

activity. Ulk3 and Stk36 double knockout NIH3T3 cells expressing CFP-Stk36^{WT} or

CFP-Stk36^{KR} (kinase dead) lentiviral construct were treated with or without Shh-N,

followed by IP with an GFP antibody and western blot analysis with the pT158/pS159

antibody.

(E-F) Western blot analysis (E) and quantification (F) of T158/S159 phosphorylation in

Fg-Stk36^{WT}, Fg-Stk36^{EE} or Fg-Stk36^{AA}. HEK293T cells were transfected with the

indicated Fg-Stk36 constructs, followed by IP with anti-Flag antibody and western blot

analysis with the indicated antibodies.

(G-H) Western blot analysis (G) and quantification (H) of Myc-Gli2 phosphorylation on

S230/S232 by the wild type (WT) or indicated Stk36 variants in HEK293T cells.

(I) Ulk4 is not required for Shh-induced Stk36 kinase activation. NIH3T3 cells

expressing the indicated shRNA and CFP-Stk36 lentiviral construct were treated with or

without Shh-N, followed by IP with an GFP antibody and western blot analysis with the

pT158/pS159 antibody (right). The knockdown efficiency of Ulk4 was determined by RT-

qPCR (left).

Data are mean ± SD from three independent experiments. **p < 0.01, and ***p < 0.001

(Student's t test).

Figure 4. Ulk4 mediates the interaction between Stk36 and Gli2.

789
(A) Depletion of Ulk4 diminished the association between Stk36 and Gli2. HEK293T

cells were transfected with the indicated shRNA and Fg-Stk36 and Myc-Gli2 constructs,

followed by CoIP and western blot analysis with the indicated antibodies.

(B) Domain structure and deletion mutants of human Stk36 used for the Co-IP

experiments.

(C-D) Stk36 binds to Ulk4 and Gli2 through its N-terminal kinase domain. HEK293T

cells were transfected with Ulk4-HA (C) or Myc-Gli2 (D) and the indicated Fg-Stk36

constructs, followed by Co-IP and western blot analyses with the indicated antibodies.

Black triangles indicate the heavy chain of IgG and asterisks indicate the protein bands

produced by individual Stk36 constructs.

(E) Both N- and C-terminal domains of Ulk4 are required for mediating Stk36/Gli2

association. NIH3T3 cell lines stably expressing control (Ctrl) or Ulk4 shRNA were

infected with lentiviruses expressing the indicated hUlk4 constructs (diagrams shown on

F, right), Fg-Stk36 and Myc-Gli2, and treated with or without the Shh-N, followed by

CoIP and western blot analysis with the indicated antibodies.

(F) Schematic diagram showing the interaction relationship between Ulk4 and Stk36,

Gli2.

Figure 5. Shh stimulates ciliary tip accumulation of both Ulk4 and Stk36.
809

(A-D) Representative images of immunostaining (A, C) and quantification (B, D) of

ciliary tip localized Myc-Stk36 (green in A) or Ulk4-HA (green in C) and Gli2 (blue in A

and C) in NIH3T3 cells infected with the Myc-Stk36 or Ulk4-HA lentivirus and treated

with or without Shh-N. Primary cilia were marked by acetylated tubulin (Ac-tub) staining

(red in A and C).

(E-F) Representative images of immunostaining (E) and quantification (F) of ciliary tip

localized Myc-Stk36 (green) and Ulk4-HA (blue) in NIH3T3 cells co-infected with Myc-

Stk36 and Ulk4-HA lentiviruses and treated with or without Shh-N. Primary cilia were

marked by Ac-Tub staining (red).

The cells are starved in serum-free medium for 12 hours to allow ciliation and cultured

in the same medium with or without Shh-N fragment for another 12 hours before they

are subjected into immunostaining assay. The intensity of ciliary-localized Myc-Stk36,

Ulk4-HA and Gli2 was measured by Image J. 20 cells were randomly selected and

counted for each group. Data are mean ± SD from three independent experiments. ***p

< 0.001 (Student's t test). Scale bars are 2 μM.

Figure 6. Stk36 and Ulk4 depend on each other for their ciliary tip accumulation

(A-B) Representative images of immunostaining (A) and quantification (B) of ciliary tip

localized Myc-Stk36 in NIH3T3 cells infected with the indicated shRNA and Myc-Stk36

lentivirus in the presence or absence of Shh-N. Primary cilia were marked by Ac-Tub

staining (red); Myc-Stk36 was marked by Myc staining (green); nuclei were marked by

DAPI (Blue).

(C-D) Representative images of immunostaining (C) and quantification (D) of ciliary tip

localized Ulk4-HA (green) in wild type (Ctrl) or Ulk3 and Stk36 double knockout (DKO)

NIH3T3 cells treated with or without Shh-N. Primary cilia were marked by Ac-Tub

staining (red) and nuclei by DAPI (Blue).

(E-F) Representative images of immunostaining (E) and quantification (F) of ciliary-

localized Ulk4-HA (green) in Shh-N treated control or DKO NIH3T3 cells infected with or

without lentiviruses expressing the indicated Ulk3 or Stk36 constructs.

The cells are starved in serum-free medium for 12 hours to allow ciliation and cultured

in the same medium with or without Shh-N fragment for another 12 hours before they

are subjected into immunostaining assay. The intensity of ciliary-localized Myc-Stk36 or

Ulk4-HA was measured by Image J. 20 cells were randomly selected and counted for

each group. Data are mean ± SD from three independent experiments. ***$p < 0.001$

(Student's t test). Scale bars are 2 μM.

Figure 7. Ulk4 ciliary tip accumulation is promoted by its phosphorylation by

Stk36

(A) Diagrams of hUlk4 deletion constructs.

(B-C) Representative images of immunostaining (B) and quantification (C) of ciliary tip

localized full-length or truncated hUlk4-HA in Ulk4 knockdown cells treated with Shh-N.

(D) Stk36 and CK1 phosphorylation sites in the C-terminal region of hUlk4.

(E-F) Representative images of immunostaining (E) and quantification (F) of ciliary tip

localized wild type (WT) or mutant hUlk4-HA baring the indicated amino acid

substitutions in Ulk4 knockdown NIH3T3 cells treated with Shh-N.

(G) *In vitro* kinase assay using the immunopurified Fg-Stk36^{WT} or Fg-Stk36^{AA} as kinase

and the indicated GST-Ulk4C fusion proteins as substrates. Phosphorylation was

detected by the pIMAGO system.

(H) Schematic diagram showing the sequence alignment of C-terminal phosphorylation

sites of mouse Ulk4 (mUlk4) and human Ulk4 (hUlk4). The Thr residues phosphorylated

by Stk36 are color-coded in green.

The cells are starved in serum-free medium for 12 hours to allow ciliation and cultured

in the same medium with or without Shh-N fragment for another 12 hours before they

are subjected into immunostaining assay. The intensity of ciliary-localized WT and

mutant Ulk4-HA was measured by Image J. 20 cells were randomly selected and

counted for each group. Data are mean ± SD from three independent experiments. ***p

< 0.001 (Student's t test). Scale bars are 2 μM.

867
868   **Figure 8. Ciliary tip localization of Ulk4 is required for Shh signal transduction**
869
(A-B) Representative images of immunostaining (A) and quantification (B) of ciliary tip

localized Myc-Stk36 and hUlk4 in Ulk4-depleted NIH3T3 cells expressing the indicated

hUlk4 constructs and treated with Shh.

(C-C') Western blot analysis (C) and quantification (C') of Myc-Gli2 phosphorylation on

S230/S232 in NIH3T3 cells expressing the indicated shRNA and hUlk4 lentiviral

constructs and treated with or without Shh-N. Data are mean ± SD from two

independent experiments. ***p < 0.001 (Student's t test).

(D-E) Relative *Gli1* (D) and *Ptch1* (E) mRNA levels in NIH3T3 cells expressing the

indicated shRNA and hUlk4 lentiviral constructs treated with or without Shh-N.

(F) Model for how Ulk4 participates in Shh signaling. See text for details.

The cells are starved in serum-free medium for 12 hours to allow ciliation and cultured

in the same medium with or without Shh-N fragment for another 12 hours before they

are subjected into RNA preparation, Western blot and immunostaining assay. The

intensity of ciliary-localized WT and mutant Ulk4-HA was measured by Image J. 20

cells were randomly selected and counted for each group. Data are mean ± SD from

three independent experiments. ***p < 0.001 (Student's t test). Scale bars are 2 μM.

Supplementary Figure legends

Figure 1- figure supplement 1. The knockdown efficiency of Ulk4, Ulk3 and Stk36

in NIH3T3 cells.

(A-C) Relative Ulk4 (A), Ulk3 (B), and Stk36 (C) mRNA levels measured by RT-qPCR in

NIH3T3 cells treated with the indicated shRNA or siRNA.

Data are mean ± SD from three independent experiments. ***p< 0.001 (Student's t test).

Figure 1- figure supplement 2. Ulk4 acts in conjunction with Stk36 to regulate

Shh-induced expression of *Gli1* and *Ptch1* in MEFs.

(A-B) Relative *Gli1* (A) and *Ptch1* (B) mRNA levels measured by RT-qPCR in MEF cells

expressing the indicated shRNA and hUlk4 lentiviral constructs with or without Shh-N

treatment.

(C-D) Relative *Gli1* mRNA levels measured by RT-qPCR in MEF cells with the indicated

RNAi knockdown in the presence or absence of Shh-N treatment.

(E) Relative Ulk4, Ulk3 and Stk36 mRNA levels measured by RT-qPCR in MEF cells

treated with the indicated shRNA or siRNA.

The cells were firstly starved in serum-free medium for 12 hours then cultured in the

same medium with or without Shh-N fragment for another 12 hours before they are

subjected into RNA preparation. Data are mean ± SD from three independent

experiments. ***$p < 0.001$ (Student's t test).

909
910
Figure 5- figure supplement1. Shh stimulates ciliary tip accumulation of both Ulk4

and Stk36

(A1-D2) Representative images of immunostaining ciliary tip localized Myc-Stk36 (green

in A1-B2) or Ulk4-HA (green in C1-D2) and Gli2 (blue) in NIH3T3 cells infected with the

Myc-Stk36 or Ulk4-HA lentivirus and treated with or without Shh-N.

(E1-F2) Representative images of immunostaining of ciliary tip localized Myc-Stk36

(blue) and Ulk4-HA (green) in NIH3T3 cells co-infected with Myc-Stk36 and Ulk4-HA

lentiviruses and treated with or without Shh-N.

(G1-J2) Representative images of immunostaining of ciliary tip localized Myc-Stk36

(green in G1-H2) and Gli2 (blue in G1-H2) or Ulk4-HA (green in I1-J2) and Myc-Stk36

(blue in I1-J2) in MEFs infected with either Myc-Stk36 (G1-H2) or both Myc-Stk36 and

Ulk4-HA lentiviruses and treated with or without Shh-N.

Primary cilia were marked by acetylated tubulin (Ac-tub) staining (red in A1-J2). Insets

show the enlarged images of primary cilia with individual channels shown separately.

Scale bars are 2 μM.

A

B

C

D

E

F

A

B

C

D

E

F