Article

THE EFFECT OF SELECTED COFFEE CONSTITUENTS ON

MITOCHON-DRIAL DYSFUNCTION IN AN EARLY ALZHEIMER

DISEASE CELL MODEL

Lukas Babylon[a], Micha T. Limbeck[a], Gunter P. Eckert[a*]

[a] Justus-Liebig-University – Laboratory for Nutrition in Prevention and Therapy,

Institute of Nutritional Sciences, Biomedical Research Center Seltersberg (BFS),

Schubertstr. 81, 35392 Giessen, Germany

* Corresponding author: Gunter P. Eckert - eckert@uni-giessen.de

THE EFFECT OF SELECTED COFFEE CONSTITUENTS ON MITOCHONDRIAL DYSFUNCTION IN AN EARLY ALZHEIMER DISEASE CELL MODEL

Alzheimer disease (AD) is an emerging medical problem worldwide without any cure yet. By 2050, more than 152 million people will be affected. AD is characterized by mitochondrial dys-function (MD) and increased amyloid beta (Aβ) levels. Coffee is one of the most commonly consumed beverages. It has many bioactive and neuroprotective ingredients of which caffeine (Cof), kahweohl (KW) and cafestol (CF) shows a variety of pharmacological properties such as anti-inflammatory and neuroprotective effects. Effects of Cof, KW, and CF were tested in a cel-lular model of AD on MD and Aβ. SH-SY5Y-APP695 cells were incubated with 50µM Cof, 1µM CF and 1µM KW for 24h. The energetic metabolite ATP was determined using a luciferase-catalyzed bioluminescence assay. The activity of mitochondrial respiration chain complexes was assessed by high-resolution respirometry using a Clarke electrode. Expression levels genes were deter-mined using quantitative real-time polymerase chain reaction (qRT-PCR). The levels of amyloid β-protein ($A\beta_{1-40}$) were measured using homogeneous time-resolved fluorescence (HTRF). ROS levels, cAMP levels, and peroxidase activity were determined using a fluorescence assay. The combination of Cof, KW and CF significantly increased ATP levels. The combination had neither a significant effect on MMP, on activity of respiration chain complexes, nor on $A\beta_{1-40}$ levels. cAMP levels were slightly increased after incubation with the combination, but not the peroxi-dase activity. Pyruvate levels and the lactate-pyruvate-ration but not lactate levels were signifi-cantly enhanced. No effect was seen on the expression level of lactate dehydrogenase and py-ruvate dehydrogenase kinase. In some experiments we have tested the single substances. They showed significant results especially in ATP, lactate and pyruvate values compared to the con-trol. The combinations have a lesser effect on mitochondrial dysfunction in cells and none on Aβ production. Whereas ATP levels and pyruvate levels were significantly increased. This suggests a change in glycolysis in neuronal cells harbouring human genes relevant for AD.

Keywords: Alzheimer disease; mitochondria; mitochondria dysfunction; caffeine; cafestol; kahweol; amyloid beta;

1. Introduction

Alzheimer's disease (AD) is a neurodegenerative disease belonging to the class of dementias. It manifests itself through a progressive loss of function of neurons of the central nervous system [1]. Dementia affects 5-7% of people over 60 in developed countries [2], of which Alzheimer's disease, discovered by German physician Alois Alzheimer in 1906, is the most common manifestation, accounting for 60% [1]. Clinical symptoms include memory loss as well as language and orientation problems [2]. One possible cause of AD is a manifestation of mitochondrial dysfunction (MD) as an early event of AD [3,4]. Almost all mitochondrial functions are impaired in AD [5,6]. First signs of MD are a reduction in key enzymes of in the oxidative metabolism [7,8] and glucose consumption [9]. The limited function of the electron transport chain (ETC) is the reason for the decrease activity of complexes IV and I. These complications lead to a lower membrane potential (MMP) and to reduced ATP levels [10,11]. A further possible trigger of the emergence of AD is the overexpression of amyloid-beta (Aβ) [12]. An imbalance between production and removal leads to accumulation and aggregation in the brain, this leads to inflammatory responses, production of reactive oxygen species (ROS) and loss of neurons, this leads to MD, dementia and Alzheimer disease [13]. An important antioxidant enzyme involved in AD is peroxidase, which catalyzes the oxidation of organic and non-organic substances using H_2O_2. In this process, it is supposed to protect against the harmful effects of physiological accumulation of ROS. A change in peroxidase activity could induce effects on ROS levels. Another impairment in AD is cyclic adenosine monophosphate (cAMP), which is one of the nucleotides and serves as a second messenger in the body for signaling metabolic pathways and hormone effects. [14,15]. The signaling cascade consisting of cAMP/PKA/CREB is considered to be generally important for the processes and

functioning in learning and memory [16]. Performance deficits may be mediated by

deficits in the signaling cascade. One trigger of this impairment could be Aβ toxicity

[17]. Up to now, there is no cure for it, which is why prevention and therapy are in the

focus of attention. A possibility for this offer might be coffee.

Today, there is a high consumption of coffee worldwide, which in Germany is

150 l per capita per year [18]. Due to a high diversity of bioactive ingredients, the

beverage has been the subject of research for many years regarding its influence on

human health. Caffeine has been the focus of attention due to its stimulant effects. In

addition, many other secondary plant compounds with bioactive effects are now known

in coffee. These potentially health-promoting effects are based on cardioprotective,

hepatoprotective and neuroprotective properties [19]. Roasted coffee contains a variety

of bioactive ingredients. Besides caffeine, coffee contains other diterpenes in the form

of fatty acid esters, with cafestol and kahweol making up the largest proportion [20].

Depending on the coffee variety, the composition of cafestol and kahweol differs.

Arabica coffee contains cafestol and kahweol, whereas robusta contains cafestol and

small amounts of kahweol [21]. The effects of caffeine are due to mechanisms of action

on nerve cells in the brain, where the pathological changes during AD also occur [22].

Coffee and caffeine intake is associated with a lower risk of developing dementia and

AD through a wide variety of mechanisms [23–25]. The data situation on kahweol and

cafestol is limited, but there are some evidence that both substances also have a

neuroprotective effect [26–28]. In this study, we focused on the combination of

caffeine, kahweol and cafestol in small doses on an early AD cell model.

2. Materials and Methods

2.1 Cell culture

Human neuroblastoma SH-SY5Y cells were cultured at 37 °C under an atmosphere of 5% CO_2 in DMEM supplemented with 10% (v/v) heat-inactivated fetal calf serum, 60µg/mL streptomycin, 60units/mL penicillin, 0.3mg/mL hygromycin, MEM non-essential amino acids, and 1mM sodium pyruvate 1%. SH-SY5Y cells were stably transfected with DNA constructs harboring human wild-type APP_{695} (APP_{695}) (for details; please refer to [29]). Cells were passaged every 3 days and were used for experiments when they reached 70-80% confluence.

2.2 Cell treatment

Cells were incubated with 1µM kahweol (KW), 1µM cafestol (Caf) and 50µM caffeine (Cof) (KCC) for 24h. The solution medium DMSO served as control.

2.3 ATP Measurement

A bioluminescence assay was used to determine the ATP levels, which is based on the production of light from ATP and luciferin in the presence of luciferase. The test was performed using the ATPlite Luminescence Assay System (PerkinElmer, Rodgau, Germany) according to the previously published protocol [30].

2.4 MMP Measurement

Mitochondrial membrane potential (MMP) was measured using the fluorescence dye rhodamine-123 (R123). Cells were incubated for 15 min with 0.4 µM R123 and centrifuged at 750 x g for 5 min before being washed with Hank's Balanced Salt

Solution (HBSS) buffer supplemented with Mg^{2+}, $Ca^{2+,}$ and HEPES. The Cells were

suspended with fresh HBSS before they were evaluated by measuring the R123

fluorescence. The excitation wavelength was set to 490 nm and the emission wavelength

to 535 nm.

2.5 Cellular Respiration

Respiration in SH-SY5Y$_{695}$ cells was assessed using an Oxygraph-2k (Oroboros,

Innsbruck, Austria) and DatLab 7.0.0.2. The cells were treated according to a complex

protocol developed by Dr Erich Gnaiger [31]. They were incubated with different

substrates, inhibitors and uncouplers. First, cells were washed with PBS (containing

potassium chloride 26.6 mM, potassium phosphate monobasic 14.705 mM, sodium

chloride 1379.31 mM and sodium phosphate dibasic 80.59 mM) and scraped into

mitochondrial respiration medium (MiRO5) developed by Oroboros [31]. Afterward,

they were centrifuged, resuspended in MiRO5, and diluted to 10^6 cells/mL. After 2mL

of cell suspension was added to each chamber and endogenous respiration was

stabilized, the cells were treated with digitonin ($10\mu g/10^6$ cells) to permeabilize the

membrane, leaving the outer and inner mitochondrial membrane intact. OXPHOS was

measured by adding the complex I and II substrates malate (2mM), glutamate (10mM)

and ADP (2mM), followed by succinate (10 mM). Gradual addition of carbonyl

cyanide-4- before it is evaluated by measuring the R123 fluorescence

(trifluoromethoxy) phenylhydrazone showed the maximum capacity of the electron

transfer system. To measure complex II activity, rotenone (0.1mM), a complex I

inhibitor, was injected. After that, oligomycin (2 µg/mL) was added to measure the leak

respiration. Inhibition of complex III by the addition of antimycin A (2.5 µM)

determined residual oxygen consumption, which was subtracted from all respiratory

states. The activity of complex IV was measured by adding N, N, N', N'-tetramethyl-p-

phenylenediamine (0.5 mM) and ascorbate (2 mM). To measure the sodium

autoxidation rate, azide ($\geq$100mM) was added. Afterward, complex IV respiration was

corrected for autoxidation. DMSO served as control.

2.6 Citrate Synthase Activity

Cell samples from respirometry measurements were frozen and stored at -80°C for the

determination of citrate synthase activity. Samples were thawed while the reaction mix

(0.1 mM 5,5′-dithiol-bis-(2-nitrobenzoic acid) (DTNB), 0.5 mM oxaloacetate, 50 μM

EDTA, 0.31 mM acetyl coenzyme A, 5 mM triethanolamine hydrochloride and 0.1 M

Tris-HCl) was mixed and heated at 30°C for 5 min. Afterward, 40 µl of samples were

submitted in triplets and mixed with 110µl of the reaction mix. The absorption was

measured at 412nm.

2.7 Aβ_{1-40} Determination

After 24 h incubation, the Aβ_{1-40} levels were determined in SH-SY5Y-APP$_{695}$ cells

using HTRF Amyloid-Beta 1-40 kit (Cisbio, Codolet, France). The protocol was the

same as recently described [32]. Aβ concentrations were normalized against the protein

content.

2.8. Peroxidase Activity

Peroxidase activation was measured using an AmplexTM Red Peroxidase Kit (Thermo

Fisher Scientific, Waltham, MA, USA) according to the manufacturer's instructions.

Cells were seeded in 96-well plates and incubated for 24 h.

2.9 Protein Quantification

Protein content was determined using a Pierce™ Protein Assay Kit (Thermo Fisher

Scientific, Waltham, MA, USA) according to the manufacturer's instructions. Bovine

serum albumin was used as a standard.

2.10. ROS Measurement

Cellular ROS production was determined using a DCFDA/H2DCFDA Cellular ROS

Assay Kit (ab113851; Abcam, Cambridge, UK). Cells were incubated for 24h with our

substances and then the manufacturer's instructions were followed.

2.11 Pyruvate and Lactate Content

Frozen cells, which were previously harvested and incubated for 24h, were thawed at

room temperature. Pyruvate and lactate concentrations were assessed using a pyruvate

assay kit (MAK071, Sigma Aldrich, Darmstadt, Germany) and a lactate assay kit

(MAK064, Sigma Aldrich, Darmstadt, Germany) according to the manufacturer's

instructions.

2.12 Real-Time qRT-PCR

Total RNA was isolated using the RNeasy Mini Kit (Qiagen, Hilden, Germany)

according to the manufacturer's guidelines. RNA was quantified using a Nanodrop™

2000c spectrometer (Thermo Fisher Scientific, Waltham, MA, USA). To remove

residual genomic DNA, samples were treated with a TURBO DNA-free™ kit according

to the manufacturer's instructions (Thermo Fisher Scientific, Waltham, MA, USA).

Complementary DNA was synthesized from 1 µg total RNA using an iScript cDNA

Synthesis Kit (Bio-Rad, Munich, Germany). qRT-PCR was conducted using a CfX 96

Connect™ system (Bio-Rad, Munich, Germany). Primers were provided from Biomol

(Hamburg, Germany). All primers are listed in Table 1. The cDNA aliquots were

diluted 1:10 with RNase-free water (Qiagen, Hilden, Germany), and all samples were

analyzed in triplicate. PCR cycling conditions were as follows: initial denaturation for 3

min at 95 °C, followed by 45 cycles at 95 °C for 10 s, 58 °C for 30 s (or 56 °C for 45 s,

depending on the primer), and 72 °C for 29 s. Expression was analyzed with –(2ΔΔCq)

using Bio-Rad CfX manager software. The normalization factor was calculated based

on the geometric mean of the levels of multiple control genes of *ß-actin (ACTB),*

glyceraldehyde-3-phosphate dehydrogenase (GAPDH), and *phosphoglycerate kinase 1*

(PGK1) according to the MIQE guidelines [33]. No-template control served as an assay

control to exclude impurities.

Table 1. Oligonucleotide primer sequences, product sizes, and primer concentrations
for quantitative real-time PCR.

Primer	Sequence	Manufacturer	Product size	Concentration (nM)
ß-Actin (ACTB) NM_001101.2	5′-ggacttcgagcaagagatgg-3′ 5′-agcactgtgttggcgtacag-3′	Biomol, Hamburg, Germany	234	200
Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) NM_002046.2	5′-gagtcaacggatttggtcgt-3′ 5′-ttgattttggagggatctcg-3′	Biomol, Hamburg, Germany	238	200
Phosphoglycerate kinase 1 (PGK1) NM_000291.2	5′-ctgtgggggtatttgaatgg-3′ 5′-cttccaggagctccaaa-3′	Biomol, Hamburg, Germany	198	200
Pyruvate dehydrogenase kinase, isozyme 1 (PDK1) NM_002610	5′-atacggatcagaaaccgaca-3′ 5′-cagacgcctagcattttcat-3′	Biomol, Hamburg, Germany	291	100
Human lactate dehydrogenase A like 6B (LDHA) NM_033195	5′-ggtgtccctttgaaggatct-3′ 5′-tgcagtcacttctttgtgga-3′	Biomol, Hamburg, Germany	87	400

206
207

2.13 cAMP level

Cellular peroxidase activity was determined using a cAMP Direct Immunoassays Kit

(Abcam, Cambridge, UK). Cells were incubated for 24h with our substances and then

the manufacturer's instructions were followed.

2.14 Statistics

Unless otherwise stated, values are presented as mean ± standard error of the mean

(SEM). Statistical analyses were performed by applying one-way ANOVA with

Tukey´s multiple comparison post-hock test and student's unpaired t-test (Prism 9.1

GraphPad Software, San Diego, CA, USA). Statistical significance was defined for *p*

values * p < 0.05, ** p < 0.01, *** p < 0.001 and **** p < 0.0001.

3. Results

3.1 Mitochondrial functions

To find the right concentrations for our combination we tested different concentrations

of Cof, CF and KW (SupFigure 1), as well as combinations of the three mentioned

substances, for the best combination, for the effects on the ATP levels (SupFigure 2).

The concentration of kahweol 1μM, cafestol 1μM and caffeine 50μM (KCC) contain all

of our three substances and turned out to be the best. There the highest ATP values for a

combination were achieved. We tested KCC on the mitochondrial functions, we

incubated SH-SY5Y-APP695 cells for 24h with all three of the substances combined.

Respiration under O2 consumption through the respiratory chain builds up the

mitochondrial membrane potential, which is used by the ATP synthase to produce ATP.

We measured the ATP level, MMP level, respiration and citrate synthase activity (Fig.

1). The KCC combination had a significant increasing effect (p = 0.0151) on the ATP

level compared to the control (Fig. 1A). Also the single substances CF1 and KW1 (p =

0.0001) had a significant increasing effect compared to the control. Whereas the

incubation with KCC had no effect on the MMP level, respiration or citrate synthase

activity. Caffeine alone does not increase or decrease ATP levels. Only the combination

with one or both diterpenes shows a significant increase at 50 μM (SupFigure 2). We

chose the concentration of 50μM caffeine to be closer to the doses that are realistic

when consuming coffee without damaging the cells or causing a loss of ATP.

Figure 1: ATP level, MMP level, respiration and citrate synthase activity of SH-SY5Y-APP$_{695}$ cells incubated for 24h with KCC. (A) ATP level of SH-SY5Y-APP$_{695}$ cells incubated with KCC. Cells treated with cell culture medium served as control (100%). N = 8. B) MMP level of 2x10^5 SH-SY5Y-APP$_{695}$ cells incubated with N = 16. (C) Respiration of SH-SY5Y-APP$_{695}$ cells incubated with KCC compared to the control. SH-SY5Y-APP$_{695}$ cells adjusted to international units (IU) of citrate synthase activity. N = 18. (G) Citrate synthase activity of SH-SY5Y-APP$_{695}$ cells incubated with KCC compared to control. N = 18. Significance was determined by Student's unpaired t-test. Data are displayed as the mean ± SEM. *P > 0.05, ****p> 0.0001. KCC = kahweol 1µM, cafestol 1µM and caffeine 50µM. Concentrations are given in µM

3.2 Aβ₁₋₄₀ Production

To measure the effect of KCC on the amyloid beta (Aβ) level SH-SY5Y-APP695 cells

were incubated for 24h. Here we found no effect of KCC on the Aβ1-40 level compared

to the control (Fig. 2).

Figure 2: Effect of KCC on the Aβ$_{1-40}$ level of SH-SY5Y-APP$_{695}$ cells incubated for

24h. N = 10. Aβ$_{1-40}$ levels were adjusted to the protein content. Significance was

determined by Student's unpaired t-test. Data are displayed as the mean ± SEM. KCC =

kahweol 1µM, cafestol 1µM and caffeine 50µM.

3.3 Peroxidase

In order to investigated the effect of KCC or the single substances on the peroxidase

activity in SH-SY5Y-APP695 cells we incubated them for 24h. Here we found no

differences between incubations and the control with regard to the peroxidase activity

(Fig. 3).

Peroxidase activity

Figure 3: Effect of KCC and the single substances on the peroxidase activity of SH-

SY5Y-APP$_{695}$ cells incubated for 24h. N = 14. Significance was determined by

Student's unpaired t-test. Data are displayed as the mean ± SEM. KCC = kahweol 1µM,

cafestol 1µM and caffeine 50µM. Concentrations are given in µM.

3.4 ROS measurement

In the next step, we examined the ROS level in SH-SY5Y-APP695 cells after the

incubation with KCC or the single substances for 24h. Here, we found no different

between both treatments (Fig. 4).

Figure 4: ROS measurement of SH-SY5Y-APP$_{695}$ cells after incubation with KCC for

24h. N = 12. Significance was determined by Student's unpaired t-test. Data are

displayed as the mean ± SEM. KCC = kahweol 1µM, cafestol 1µM and caffeine 50µM.

Concentrations are given in µM.

3.5 cAMP

To test the effect of KCC on the cAMP level, we incubated SH-SY5Y-APP$_{695}$ cells for

24h. Here we had an increase, but not a significant, of cAMP level after the incubation

with KCC compared to control (Fig. 5).

Figure 5: cAMP level of SH-SY5Y-APP$_{695}$ cells after incubation with KCC for 24h

compared to the control. N = 8. Significance was determined by Student's unpaired t-

test. Data are displayed as the mean ± SEM. KCC = kahweol 1µM, cafestol 1µM and

caffeine 50µM.

3.6 Lactate and pyruvate level

To examine the effect of KCC on the lactate and pyruvate level we incubated SH-

SY5Y-APP$_{695}$ cells for 24h. There were no different between KCC compared to the

control for the lactate level (Fig. 6A). However, the incubation with KCC significant

increased the pyruvate value compared to the control (p = 0.0027) (Fig. 6B). Likewise,

the lactate/pyruvate ratio (Fig. 6C) was significantly reduced by KCC compared to the

control (p = 0.0141). To investigate whether the increased pyruvate levels were caused

by a single substance. We determined the values of the individual substances. This

showed that the pyruvate values were significantly increased by KW (p = 0.0062, Fig

6B). All the individual substances (Fig 6C) also significantly reduced the

lactate/pyruvate ratio (Cof p = 0.0097, CF p = 0.0321, KW p = 0.0008).

A) Lactate level **B) Pyruvate level** **C) Lactate/pyruvate ratio**

Figure 6: Effect of the on lactate and pyruvate level after 24h of incubation with caffeine (Cof), cafestol (CF) and kahweol (KW) or the combination KCC. (A) Lactate level of SH-SY5Y-APP$_{695}$ cells after the incubation compared to the control. (B) Pyruvate level of SH-SY5Y-APP$_{695}$ cells after the incubation with KCC compared to the control. (C) Lactate to pyruvate ratio. N = 8. Levels were adjusted to the protein content. Concentrations are given in µM. Significance was determined by one-way ANOVA. * $p < 0.05$ and ** $p < 0.01$, ***$p > 0.001$. Data are displayed as ±SEM. Concentrations are given in µM.

3.7 qPCR

To investigate the molecular basis of altered lactate and pyruvate levels, the gene expression of pyruvate dehydrogenase kinase 1 (PDK1) and lactate dehydrogenase A (LDHA) were examined after 24 h incubation using qRT-PCR. KCC had no significant effect on LDHA or PDK1 gene expression compared to control (Fig.7).

Figure 7: Effect on the incubation with KCC on the gene expression after 24h of
incubation. (A) Gene expression of lactate dehydrogenase A (LDHA) of SH-SY5Y$_{695}$
cells after the incubation with KCC compared to the control. (B) Gene expression of
pyruvate dehydrogenase kinase 1 (PDK1) of SH-SY5Y$_{695}$ cells after the incubation with
KCC compared to the control. N = 8. The normalization factor was calculated based on
the geometric mean of the levels of multiple control genes of *ß-actin (ACTB),*
glyceraldehyde-3-phosphate dehydrogenase (GAPDH), and *phosphoglycerate kinase 1*
(PGK1) according to the MIQE guidelines [33]. Significance was determined by
Student's unpaired t-test. Data are displayed as the mean ± SEM. KCC = kahweol 1µM,
cafestol 1µM and caffeine 50µM.

4. Discussion

In the present work, we aimed to investigate different bioactive substances present in
coffee. For this purpose, we selected three molecules that occur in larger concentrations
in coffee - namely, caffeine, kahweol and cafestol. We tested various concentrations and
combinations of these three compounds, of which 1µM kahweol, 1µM cafestol and
50µM caffeine (KCC) together were most promising. The compounds were tested in a
SH-SY5Y-APP$_{695}$ cells, an established model for early AD.

First, we investigated the effects of the single coffee compounds and its

combination (KCC) on the mitochondrial dysfunction present occurring in SH-SY5Y-

APP_{695} cells. The incubation with KCC significantly increased the ATP levels

compared to the control (Fig. 1A). In addition, KW1 and CF1 increase the ATP levels

significant. Caffeine alone does not increase or decrease ATP levels. Only the

combination with one or both diterpenes shows a significant increase at Cof 50µM

(SupFigure 2). In a transgenic mouse model of AD that oral treatment with caffeine

intake of 0.6mg/d raised the ATP level significantly [34]. According to our result,

kahweol was able to significantly rise ATP levels in SH-SY5Y cells a model [28]. It

seems a combination of both compounds has equally positive effects on ATP levels.

Due to the lack of knowledge it is unclear to what extent CF affects ATP levels. We

found no indication that increased ATP levels were a result of enhanced MMP in

consequence of an accelerated complex activity through the respiratory chain. In studies

on SH-SY5Y cells with KW, no change in MMP or oxygen consumption could be

measured by single administration, but KW was able to attenuate the effect by the

stressor H_2O_2 [26,28]. This suggests that KW in combination with the other investigated

substances has a positive effect on the maintenance of the MMP and the complex

activity of the respiratory chain. In addition, the increased ATP level were not a result

of a boosted mitochondrial mass (Fig. 1D), which leads to the assumption that these

come from a different source. Another parameter related to mitochondrial quality and

functionality are reactive oxygen species (ROS) that are produced in excess due to an

insufficient proton transfer through the mitochondrial respiration chain. Due to their

ability to interact with cellular components such mitochondrial membrane lipids or

DNA, they can thus cumulatively cause oxidative damage that shortens lifespan and

leads to impaired mitochondrial function [35]. However, in our study the tested

compounds and their combination showed no antioxidant activity. Studies have shown

that the addition of caffeine can lower ROS levels and thus have anti-inflammatory

effects. Overall, caffeine is thought to reduce oxidative stress. Caffeine has been shown

to inhibit the activity of NF-$_k$B and increase that of superoxide dismutase [36]. CF is

also thought to have a suppressive function of NF-$_k$B [37]. Peroxidase, which catalyses

the oxidation of organic and non-organic substances using H_2O_2, is considered an

important antioxidant enzyme. In this process, it is supposed to protect against the

harmful effects of physiological accumulation of ROS. Caffeine has been shown to

increase the activity of antioxidant enzymes in the liver of mice or to increase overall

antioxidant [38,39]. Glutathione peroxidase as well as superoxide dismutase, both

antioxidant enzymes, benefit from increased activity after coffee addition in human

experiments [40]. We examined the peroxidase activity that represents an important

detoxification mechanism and found no significant differences by incubation with the

single substances or its combination. The pathological appearance of AD is

characterized by a deposition of oligomeric Aβ in the brain. For this purpose, we

measured Aβ$_{1-40}$ levels in our experiments (Fig. 2). There were no significant changes

due to the incubations compared to the control. However, there is evidence that there is

a caffeine-induced protective effect on the internalization of APP processing. Thus,

Aβ$_{1-40}$ concentration was significantly decreased due to the addition of caffeine [41]. In

animal experiments with mice, it was shown that both a single and long-term

administration of caffeine significantly reduced plasma Aβ levels. Increased cognitive

performance and decreased Aβ deposition in the brain were also observed [42]. Studies

in humans showed that moderate coffee consumption was associated with lower Aβ

levels. This potential neuroprotective effect is consistent with other studies that suggest

coffee consumers are at lower risk for AD [43–45]. It is possible that a longer

incubation period with the three components would have had a greater effect on Aβ

production.

Cyclic adenosine monophosphate (cAMP) belongs to the nucleotides and serves

as a second messenger in the body. Its tasks are signal transmission in the context of

metabolic pathways and hormonal effects, in which a signal cascade is induced by

cAMP. Its main mode of action is to activate the cAMP-dependent protein kinase A

enzyme family. One consequence of protein kinase A (PKA) being activated in this way

is that it influences gene transcription by means of phosyphorylation of the cAMP

response element binding protein (CREB) [14,15]. CREB affects neuron growth, neuron

differentiation, synaptic plasticity and neurogenesis. In diseases such as Parkinson's

disease, Huntington's disease and AD, an impairment of CREB has been observed [46].

The cAMP signaling cascades thus altered have the ability to modulate long-term

memory. This is based on the interaction of phosphorylated CREB with target genes.

Thus, it is postulated that cAMP has a beneficial effect on cognitive performance in the

context of short-term and long-term memory [47]. Cognitive performance deficits

occurring in AD is mediated, among other things, by the cAMP/PKA/CREB signaling

cascade. Aβ toxicity may represent one trigger of this impairment [17]. Furthermore,

pathological Aβ concentration shows to cause a rapid and sustained decrease in PKA

activity and inhibition of CREB phosphorylation [48]. Similarly, oxidative stress

inhibits PKA activity and phosphorylation of CREB have. Administration of

antioxidants has been shown to protect neurons from Aβ-induced inactivation by PKA

[16]. In our study, Figure 5 shows an increase in cAMP levels by incubation with KCC

compared with the control, although not a significant one. Studies have shown that

caffeine has an increasing effect on cAMP concentration [49,50]. Furthermore, caffeine

has shown a beneficial effect on PKA activity in animal models of AD [48]. Overall,

many benefits lie in increasing the cAMP signalling cascade, which includes activation

of CREB, sirtuin 1, and Nrf2. In the Nrf2 signalling pathway, potential modes of action

of caffeine, cafestol, and kahweol overlap. Studies suggest that CF and KW may

condition health-promoting effects via Nrf2, but unlike caffeine, without exerting

influence on cAMP levels. There is no known cAMP modulation by CF or KW [51–

53].

Glucose is the primary source of energy for brain activity. It is suspected that

due to this, an impairment of glucose metabolism is associated with neurodegenerative

diseases [54]. Lactate is the end product of anaerobic glycolysis catalysed by lactate

dehydrogenase (LDH). Lactate is considered as an important bioenergetic metabolite

formed in the absence of oxygen by fermentation or in the presence of oxygen by

aerobic glycolysis [55]. Lactate is the link between the glycolytic and aerobic metabolic

pathways [56]. It has been observed that when ATP production is impaired, lactate

production for short- and long-term energy provision is stimulated. Thus, increased

ATP levels may decrease cellular lactate demand [57]. In our work, despite increased

ATP levels, there was no decrease in lactate values (Fig. 6A). Pyruvate is the end

product of glycolysis and is considered a key molecule for numerous metabolic

pathways. These include mitochondrial ATP generation and several biosynthetic

pathways that cross the citrate cycle [58]. In general, neurons have an increased

pyruvate demand due to a high energy turnover [59,60]. The significantly increased

pyruvate concentration may indicates a relevant influence on the metabolism of

pyruvate [61]. In our study, incubation with the combination of compounds significantly

increased pyruvate levels (Fig. 6B). This effect might relay on KW, which as a single

substance significantly increased the pyruvate levels, whereas the other single

substances could only partly raise the levels. Other studies have shown that increased

pyruvate levels can protect neurons from damage. In this case, the damage triggered by

a stressor was annulated by an increased pyruvate concentration and at the same time

pyruvate specifically protects the mitochondria from oxidative stress [62,63]. With

regard to the lactate/pyruvate ratio (L/P), the results obtained indicate that there is a

lowered ratio in favour of pyruvate. An increase in L/P is accompanied by an increase

in $NADH/NAD^+$. In our experiments, there should be a decreased ratio of $NADH/NAD^+$

corresponding to L/P [56]. A shift in this ratio could have an impact on glycolysis, as

there is inhibition of glycolysis when the $NADH/NAD^+$ ratio is increased [56,64]. This

may suggest that increased glycolysis may be present in the experiments performed here

due to a decreased $NADH/NAD^+$ ratio. Gene expression analysis of lactate

dehydrogenase A (LDHA) and pyruvate dehydrogenase kinase 1 (PDK1) performed in

our study (Fig. 7) showed no significant differences from control. There was a slight

decrease in LDA and a slight increase in PDK1 expression. In an AD model with

Drosophila melanogaster, downregulation of LDHA was shown to stimulate

neuroprotective effects. Consistent with this, overexpression of LDHA showed a

shortened lifespan as well as brain dysfunction. Based on this, LDHA can be considered

a modulator of aging processes [65–67]. PDK1 inhibits pyruvate dehydrogenase (PDH)

via reversible phosphorylation. PDH catalyses the oxidative decarboxylation of

pyruvate to acetyl-CoA and CO_2 in the mitochondrial matrix [63]. Shift in PDK activity

may result in changes in mitochondrial function as well as neurological activity. With

age, there is a decrease in PDH via increased PDK activity, which shifts energy

production from aerobic to anaerobic glycolysis [68]. Overall, synergistically, the slight

decrease in LDHA in combination with the slightly increased PDK1 could explain the

increased pyruvate concentration. Thus, the increased ATP levels could be the result of

increased glycolysis. This would go hand in hand with the observations made of the

increased pyruvate concentration. Thus, an increased glucose metabolism up to the

generation of pyruvate can be assumed. If an increased metabolism of glucose persists,

need to be addressed by future studies.

5. Conclusion

In the present study, we observed the effect of a combination of different coffee

components, kahweol, caffestol, and caffeine, on early AD symptomatology in a cell

model of early AD. ATP levels were increase independent from mitochondrial function.

Despite the lack of change in gene expression and the lack of change in mitochondrial

complex activity, we suspect a change within glycolysis due to the change in pyruvate

level. Further investigations have to address if coffee components affect glycolysis.

Supplementary Materials: The following supporting information can be downloaded at:

Author Contributions; investigation, L.B, M.T.L.; writing—original draft preparation, L.B.;

supervision, G.P.E. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding

Data Availability Statement: The data presented in this study are available on request from the

corresponding author.

Conflicts of Interest: The authors declare no conflict of interest

References

1.   Lane CA, Hardy J, Schott JM (2018) Alzheimer's disease. Eur J Neurol 25:59–

70. https://doi.org/10.1111/ene.13439

2.   Jessen F (ed) (2018) Handbuch Alzheimer-Krankheit: Grundlagen - Diagnostik -

Therapie - Versorgung - Prävention. De Gruyter, Berlin, Boston

3.   Castellani R, Hirai K, Aliev G et al. (2002) Role of mitochondrial dysfunction in

Alzheimer's disease. J Neurosci Res 70:357–360. https://doi.org/10.1002/jnr.10389

4.   Swerdlow RH, Burns JM, Khan SM (2010) The Alzheimer's disease

mitochondrial cascade hypothesis. J Alzheimers Dis 20 Suppl 2:S265-79.

https://doi.org/10.3233/JAD-2010-100339

5.   Müller WE, Eckert A, Kurz C et al. (2010) Mitochondrial dysfunction: common

final pathway in brain aging and Alzheimer's disease--therapeutic aspects. Mol

Neurobiol 41:159–171. https://doi.org/10.1007/s12035-010-8141-5

6.   Wang X, Su B, Zheng L et al. (2009) The role of abnormal mitochondrial

dynamics in the pathogenesis of Alzheimer's disease. J Neurochem 109 Suppl 1:153–

159. https://doi.org/10.1111/j.1471-4159.2009.05867.x.

7.   Parker WD, Parks J, Filley CM et al. (1994) Electron transport chain defects in

Alzheimer's disease brain. Neurology 44:1090–1096.

https://doi.org/10.1212/wnl.44.6.1090

8.   Chandrasekaran K, Giordano T, Brady DR et al. (1994) Impairment in

mitochondrial cytochrome oxidase gene expression in Alzheimer disease. Molecular

Brain Research 24:336–340. https://doi.org/10.1016/0169-328X(94)90147-3

9.   Landau SM, Harvey D, Madison CM et al. (2011) Associations between

cognitive, functional, and FDG-PET measures of decline in AD and MCI. Neurobiol

Aging 32:1207–1218. https://doi.org/10.1016/j.neurobiolaging.2009.07.002

10. Wang X, Su B, Lee H et al. (2009) Impaired balance of mitochondrial fission

and fusion in Alzheimer's disease. J Neurosci 29:9090–9103.

https://doi.org/10.1523/JNEUROSCI.1357-09.2009

11. Rhein V, Song X, Wiesner A et al. (2009) Amyloid-beta and tau synergistically

impair the oxidative phosphorylation system in triple transgenic Alzheimer's disease

mice. Proc Natl Acad Sci U S A 106:20057–20062.

https://doi.org/10.1073/pnas.0905529106

12. Gallardo G, Holtzman DM (2019) Amyloid-β and Tau at the Crossroads of

Alzheimer's Disease. Adv Exp Med Biol 1184:187–203. https://doi.org/10.1007/978-

981-32-9358-8_16

13. Lemere CA, Masliah E (2010) Can Alzheimer disease be prevented by amyloid-

beta immunotherapy? Nat Rev Neurol 6:108–119.

https://doi.org/10.1038/nrneurol.2009.219

14. Serezani CH, Ballinger MN, Aronoff DM et al. (2008) Cyclic AMP: master

regulator of innate immune cell function. Am J Respir Cell Mol Biol 39:127–132.

https://doi.org/10.1165/rcmb.2008-0091TR

15. Yan K, Gao L-N, Cui Y-L et al. (2016) The cyclic AMP signaling pathway:

Exploring targets for successful drug discovery (Review). Mol Med Rep 13:3715–3723.

https://doi.org/10.3892/mmr.2016.5005

16. Du H, Guo L, Wu X et al. (2014) Cyclophilin D deficiency rescues Aβ-impaired

PKA/CREB signaling and alleviates synaptic degeneration. Biochim Biophys Acta

1842:2517–2527. https://doi.org/10.1016/j.bbadis.2013.03.004

17.   Tong L, Thornton PL, Balazs R et al. (2001) Beta -amyloid-(1-42) impairs

activity-dependent cAMP-response element-binding protein signaling in neurons at

concentrations in which cell survival Is not compromised. J Biol Chem 276:17301–

17306. https://doi.org/10.1074/jbc.M010450200

18.   Chu Y-F (ed) (2012) Coffee: Emerging health effects and disease prevention.

IFT Press series. IFT Press Wiley-Blackwell A John Wiley & Sons Ltd. Publication,

Ames, Iowa

19.   Sc Y, Muralidhara (2016) Beneficial Role of Coffee and Caffeine in

Neurodegenerative Diseases: A Minireview. AIMS Public Health 3:407–422.

https://doi.org/10.3934/publichealth.2016.2.407

20.   Hu GL, Wang X, Zhang L et al. (2019) The sources and mechanisms of

bioactive ingredients in coffee. Food Funct 10:3113–3126.

https://doi.org/10.1039/c9fo00288j

21.   Speer K, Kölling-Speer I (2006) The lipid fraction of the coffee bean. Braz J

Plant Physiol 18:201–216. https://doi.org/10.1590/S1677-04202006000100014

22.   Camandola S, Plick N, Mattson MP (2019) Impact of Coffee and Cacao Purine

Metabolites on Neuroplasticity and Neurodegenerative Disease. Neurochem Res

44:214–227. https://doi.org/10.1007/s11064-018-2492-0

23.   Patil H, Lavie CJ, O'Keefe JH (2011) Cuppa joe: friend or foe? Effects of

chronic coffee consumption on cardiovascular and brain health. Mo Med 108:431–438

24.   Kim JW, Byun MS, Yi D et al. (2019) Coffee intake and decreased amyloid

pathology in human brain. Transl Psychiatry 9:270. https://doi.org/10.1038/s41398-019-

0604-5

25.   Wasim S, Kukkar V, Awad VM et al. (2020) Neuroprotective and

Neurodegenerative Aspects of Coffee and Its Active Ingredients in View of Scientific

Literature. Cureus 12:e9578. https://doi.org/10.7759/cureus.9578

26.   Oliveira MR de, Souza ICC de, Fürstenau CR (2020) Mitochondrial Protection

Promoted by the Coffee Diterpene Kahweol in Methylglyoxal-Treated Human

Neuroblastoma SH-SY5Y Cells. Neurotox Res 37:100–110.

https://doi.org/10.1007/s12640-019-00107-w

27.   Lee KJ, Jeong HG (2007) Protective effects of kahweol and cafestol against

hydrogen peroxide-induced oxidative stress and DNA damage. Toxicol Lett 173:80–87.

https://doi.org/10.1016/j.toxlet.2007.06.008

28.   Fürstenau CR, Souza ICC de, Oliveira MR de (2019) The effects of kahweol, a
560   diterpene present in coffee, on the mitochondria of the human neuroblastoma
561   SH-SY5Y cells exposed to hydrogen peroxide. Toxicol In Vitro 61:104601.
562   https://doi.org/10.1016/j.tiv.2019.104601

29.   Grewal R, Reutzel M, Dilberger B et al. (2020) Purified oleocanthal and

ligstroside protect against mitochondrial dysfunction in models of early Alzheimer's

disease and brain ageing. Exp Neurol 328:113248.

https://doi.org/10.1016/j.expneurol.2020.113248.

30.   Hagl S, Grewal R, Ciobanu I et al. (2015) Rice bran extract compensates

mitochondrial dysfunction in a cellular model of early Alzheimer's disease. J

Alzheimers Dis 43:927–938. https://doi.org/10.3233/JAD-132084

31.   Stadlmann S, Renner K, Pollheimer J et al. (2006) Preserved Coupling of

Oxidative Phosphorylation But Decreased Mitochondrial Respiratory Capacity in IL-

1β-Treated Human Peritoneal Mesothelial Cells. CBB 44:179–186.

https://doi.org/10.1385/cbb:44:2:179

32.   Babylon L, Grewal R, Stahr P-L et al. (2021) Hesperetin Nanocrystals Improve

Mitochondrial Function in a Cell Model of Early Alzheimer Disease. Antioxidants

(Basel) 10. https://doi.org/10.3390/antiox10071003

33.   Bustin SA, Benes V, Garson JA et al. (2009) The MIQE guidelines: minimum

information for publication of quantitative real-time PCR experiments. Clin Chem

55:611–622. https://doi.org/10.1373/clinchem.2008.112797

34.   Dragicevic N, Delic V, Cao C et al. (2012) Caffeine increases mitochondrial

function and blocks melatonin signaling to mitochondria in Alzheimer's mice and cells.

Neuropharmacology 63:1368–1379. https://doi.org/10.1016/j.neuropharm.2012.08.018

35.   Giorgi C, Marchi S, Simoes ICM et al. (2018) Mitochondria and Reactive

Oxygen Species in Aging and Age-Related Diseases. Int Rev Cell Mol Biol 340:209–

344. https://doi.org/10.1016/bs.ircmb.2018.05.006

36.   Londzin P, Zamora M, Kąkol B et al. (2021) Potential of Caffeine in

Alzheimer's Disease-A Review of Experimental Studies. Nutrients 13.

https://doi.org/10.3390/nu13020537

37.   Shen T, Lee J, Lee E et al. (2010) Cafestol, a coffee-specific diterpene, is a

novel extracellular signal-regulated kinase inhibitor with AP-1-targeted inhibition of

prostaglandin E2 production in lipopolysaccharide-activated macrophages. Biol Pharm

Bull 33:128–132. https://doi.org/10.1248/bpb.33.128

38.   Mukhopadhyay S, Mondal A, Poddar MK (2003) Chronic administration of

caffeine: effect on the activities of hepatic antioxidant enzymes of Ehrlich ascites

tumor-bearing mice. Indian J Exp Biol 41:283–289

39.   Lee C (2000) Antioxidant ability of caffeine and its metabolites based on the

study of oxygen radical absorbing capacity and inhibition of LDL peroxidation. Clinica

Chimica Acta 295:141–154. https://doi.org/10.1016/S0009-8981(00)00201-1

40.   Bichler J, Cavin C, Simic T et al. (2007) Coffee consumption protects human

lymphocytes against oxidative and 3-amino-1-methyl-5H-pyrido4,3-bindole acetate

(Trp-P-2) induced DNA-damage: results of an experimental study with human

volunteers. Food Chem Toxicol 45:1428–1436.

https://doi.org/10.1016/j.fct.2007.02.001

41.   Li S, Geiger NH, Soliman ML et al. (2015) Caffeine, Through Adenosine A3

Receptor-Mediated Actions, Suppresses Amyloid-β Protein Precursor Internalization

and Amyloid-β Generation. J Alzheimers Dis 47:73–83. https://doi.org/10.3233/JAD-

142223

42.   Cao C, Cirrito JR, Lin X et al. (2009) Caffeine suppresses amyloid-beta levels in

plasma and brain of Alzheimer's disease transgenic mice. J Alzheimers Dis 17:681–697.

https://doi.org/10.3233/JAD-2009-1071

43.   Butt MS, Sultan MT (2011) Coffee and its consumption: benefits and risks. Crit

Rev Food Sci Nutr 51:363–373. https://doi.org/10.1080/10408390903586412

44.   Godos J, Pluchinotta FR, Marventano S et al. (2014) Coffee components and

cardiovascular risk: beneficial and detrimental effects. Int J Food Sci Nutr 65:925–936.

https://doi.org/10.3109/09637486.2014.940287

45. Socała K, Szopa A, Serefko A et al. (2020) Neuroprotective Effects of Coffee

Bioactive Compounds: A Review. Int J Mol Sci 22.

https://doi.org/10.3390/ijms22010107

46. Sharma VK, Singh TG (2020) CREB: A Multifaceted Target for Alzheimer's

Disease. Curr Alzheimer Res 17:1280–1293.

https://doi.org/10.2174/1567205018666210218152253

47. Chen Y, Huang X, Zhang Y et al. (2012) Alzheimer's β-secretase (BACE1)

regulates the cAMP/PKA/CREB pathway independently of β-amyloid. J Neurosci

32:11390–11395. https://doi.org/10.1523/JNEUROSCI.0757-12.2012

48. Teich AF, Nicholls RE, Puzzo D et al. (2015) Synaptic therapy in Alzheimer's

disease: a CREB-centric approach. Neurotherapeutics 12:29–41.

https://doi.org/10.1007/s13311-014-0327-5

49. Echeverri D, Montes FR, Cabrera M et al. (2010) Caffeine's Vascular

Mechanisms of Action. Int J Vasc Med 2010:834060.

https://doi.org/10.1155/2010/834060

50. Orbán C, Vásárhelyi Z, Bajnok A et al. (2018) Effects of caffeine and

phosphodiesterase inhibitors on activation of neonatal T lymphocytes. Immunobiology

223:627–633. https://doi.org/10.1016/j.imbio.2018.07.008

51. Sanders O, Rajagopal L (2020) Phosphodiesterase Inhibitors for Alzheimer's

Disease: A Systematic Review of Clinical Trials and Epidemiology with a Mechanistic

Rationale. J Alzheimers Dis Rep 4:185–215. https://doi.org/10.3233/ADR-200191

52. Seo H-Y, Lee S-H, Lee J-H et al. (2020) Kahweol activates the Nrf2/HO-1

pathway by decreasing Keap1 expression independently of p62 and autophagy

pathways. PLoS One 15:e0240478. https://doi.org/10.1371/journal.pone.0240478

53.   Hao W-R, Sung L-C, Chen C-C et al. (2019) Cafestol Activates Nuclear Factor

Erythroid-2 Related Factor 2 and Inhibits Urotensin II-Induced Cardiomyocyte

Hypertrophy. Am J Chin Med 47:337–350.

https://doi.org/10.1142/S0192415X19500162

54.   Hipkiss AR (2019) Aging, Alzheimer's Disease and Dysfunctional Glycolysis;

Similar Effects of Too Much and Too Little. Aging Dis 10:1328–1331.

https://doi.org/10.14336/AD.2019.0611

55.   Tauffenberger A, Fiumelli H, Almustafa S et al. (2019) Lactate and pyruvate

promote oxidative stress resistance through hormetic ROS signaling. Cell Death Dis

10:653. https://doi.org/10.1038/s41419-019-1877-6

56.   Mintun MA, Vlassenko AG, Rundle MM et al. (2004) Increased lactate/pyruvate

ratio augments blood flow in physiologically activated human brain. Proc Natl Acad Sci

U S A 101:659–664. https://doi.org/10.1073/pnas.0307457100

57.   Brooks GA (2018) The Science and Translation of Lactate Shuttle Theory. Cell

Metab 27:757–785. https://doi.org/10.1016/j.cmet.2018.03.008

58.   Zangari J, Petrelli F, Maillot B et al. (2020) The Multifaceted Pyruvate

Metabolism: Role of the Mitochondrial Pyruvate Carrier. Biomolecules 10.

https://doi.org/10.3390/biom10071068

59.   Hassel B (2001) Pyruvate carboxylation in neurons. J Neurosci Res 66:755–762.

https://doi.org/10.1002/jnr.10044

60.   Katoh-Semba R, Keino H, Kashiwamata S (1988) A possible contribution by

glial cells to neuronal energy production: enzyme-histochemical studies in the

developing rat cerebellum. Cell Tissue Res 252:133–139.

https://doi.org/10.1007/BF00213834

61.   Schormann N, Hayden KL, Lee P et al. (2019) An overview of structure,

function, and regulation of pyruvate kinases. Protein Sci 28:1771–1784.

https://doi.org/10.1002/pro.3691

62.   Desagher S, Glowinski J, Prémont J (1997) Pyruvate Protects Neurons against

Hydrogen Peroxide-Induced Toxicity. J Neurosci 17:9060–9067.

https://doi.org/10.1523/JNEUROSCI.17-23-09060.1997

63.   Wang X, Shen X, Yan Y et al. (2021) Pyruvate dehydrogenase kinases (PDKs):

an overview toward clinical applications. Biosci Rep 41.

https://doi.org/10.1042/BSR20204402

64.   Wu J, Jin Z, Zheng H et al. (2016) Sources and implications of NADH/NAD(+)

redox imbalance in diabetes and its complications. Diabetes Metab Syndr Obes 9:145–

153. https://doi.org/10.2147/DMSO.S106087

65.   Long DM, Frame AK, Reardon PN et al. (2020) Lactate dehydrogenase

expression modulates longevity and neurodegeneration in Drosophila melanogaster.

Aging (Albany NY) 12:10041–10058. https://doi.org/10.18632/aging.103373

66.   Ross JM, Öberg J, Brené S et al. (2010) High brain lactate is a hallmark of aging

and caused by a shift in the lactate dehydrogenase A/B ratio. Proc Natl Acad Sci U S A

107:20087–20092. https://doi.org/10.1073/pnas.1008189107

67.   Rossi A, Rigotto G, Valente G et al. (2020) Defective Mitochondrial Pyruvate

Flux Affects Cell Bioenergetics in Alzheimer's Disease-Related Models. Cell Rep

30:2332-2348.e10. https://doi.org/10.1016/j.celrep.2020.01.060

68.   Kumar Jha M, Jeon S, Suk K (2012) Pyruvate Dehydrogenase Kinases in the

Nervous System: Their Principal Functions in Neuronal-glial Metabolic Interaction and

Neuro-metabolic Disorders. Current Neuropharmacology 10:393–403.

https://doi.org/10.2174/157015912804499528

References

[1]   Lane CA, Hardy J, Schott JM. Alzheimer's disease. Eur J Neurol. 2018;25:59–70.

[2]   Jessen F, editor. Handbuch Alzheimer-Krankheit: Grundlagen - Diagnostik -

Therapie - Versorgung - Prävention. Berlin, Boston: De Gruyter; 2018.

[3]   Castellani R, Hirai K, Aliev G, et al. Role of mitochondrial dysfunction in

Alzheimer's disease. J Neurosci Res. 2002;70:357–360.

[4]   Swerdlow RH, Burns JM, Khan SM. The Alzheimer's disease mitochondrial

cascade hypothesis. J Alzheimers Dis. 2010;20 Suppl 2:S265-79.

[5]   Müller WE, Eckert A, Kurz C, et al. Mitochondrial dysfunction: common final

pathway in brain aging and Alzheimer's disease--therapeutic aspects. Mol

Neurobiol. 2010;41:159–171.

[6]   Wang X, Su B, Zheng L, et al. The role of abnormal mitochondrial dynamics in the

pathogenesis of Alzheimer's disease. J Neurochem. 2009;109 Suppl 1:153–159.

[7]   Parker WD, Parks J, Filley CM, et al. Electron transport chain defects in

Alzheimer's disease brain. Neurology. 1994;44:1090–1096.

[8]   Chandrasekaran K, Giordano T, Brady DR, et al. Impairment in mitochondrial

cytochrome oxidase gene expression in Alzheimer disease. Molecular Brain

Research. 1994;24:336–340.

[9]   Landau SM, Harvey D, Madison CM, et al. Associations between cognitive,

functional, and FDG-PET measures of decline in AD and MCI. Neurobiol Aging.

2011;32:1207–1218.

[10] Wang X, Su B, Lee H, et al. Impaired balance of mitochondrial fission and fusion

in Alzheimer's disease. J Neurosci. 2009;29:9090–9103.

[11] Rhein V, Song X, Wiesner A, et al. Amyloid-beta and tau synergistically impair the

oxidative phosphorylation system in triple transgenic Alzheimer's disease mice.

Proc Natl Acad Sci U S A. 2009;106:20057–20062.

[12] Gallardo G, Holtzman DM. Amyloid-β and Tau at the Crossroads of Alzheimer's

Disease. Adv Exp Med Biol. 2019;1184:187–203.

[13] Lemere CA, Masliah E. Can Alzheimer disease be prevented by amyloid-beta

immunotherapy? Nat Rev Neurol. 2010;6:108–119.

[14] Serezani CH, Ballinger MN, Aronoff DM, et al. Cyclic AMP: master regulator of

innate immune cell function. Am J Respir Cell Mol Biol. 2008;39:127–132.

[15] Yan K, Gao L-N, Cui Y-L, et al. The cyclic AMP signaling pathway: Exploring

targets for successful drug discovery (Review). Mol Med Rep. 2016;13:3715–

3723.

[16] Du H, Guo L, Wu X, et al. Cyclophilin D deficiency rescues Aβ-impaired

PKA/CREB signaling and alleviates synaptic degeneration. Biochim Biophys Acta.

2014;1842:2517–2527.

[17] Tong L, Thornton PL, Balazs R, et al. Beta -amyloid-(1-42) impairs activity-

dependent cAMP-response element-binding protein signaling in neurons at

concentrations in which cell survival Is not compromised. J Biol Chem.

2001;276:17301–17306.

[18] Chu Y-F, editor. Coffee: Emerging health effects and disease prevention. Ames,

Iowa: IFT Press Wiley-Blackwell A John Wiley & Sons Ltd. Publication; 2012.

(IFT Press series).

[19] Sc Y, Muralidhara. Beneficial Role of Coffee and Caffeine in Neurodegenerative

Diseases: A Minireview. AIMS Public Health. 2016;3:407–422.

[20] Hu GL, Wang X, Zhang L, et al. The sources and mechanisms of bioactive

ingredients in coffee. Food Funct. 2019;10:3113–3126.

[21] Speer K, Kölling-Speer I. The lipid fraction of the coffee bean. Braz. J. Plant

Physiol. 2006;18:201–216.

[22] Camandola S, Plick N, Mattson MP. Impact of Coffee and Cacao Purine

Metabolites on Neuroplasticity and Neurodegenerative Disease. Neurochem Res.

2019;44:214–227.

[23] Patil H, Lavie CJ, O'Keefe JH. Cuppa joe: friend or foe? Effects of chronic coffee

consumption on cardiovascular and brain health. Mo Med. 2011;108:431–438.

[24] Kim JW, Byun MS, Yi D, et al. Coffee intake and decreased amyloid pathology in

human brain. Transl Psychiatry. 2019;9:270.

[25] Wasim S, Kukkar V, Awad VM, et al. Neuroprotective and Neurodegenerative

Aspects of Coffee and Its Active Ingredients in View of Scientific Literature.

Cureus. 2020;12:e9578.

[26] Oliveira MR de, Souza ICC de, Fürstenau CR. Mitochondrial Protection Promoted

by the Coffee Diterpene Kahweol in Methylglyoxal-Treated Human

Neuroblastoma SH-SY5Y Cells. Neurotox Res. 2020;37:100–110.

[27] Lee KJ, Jeong HG. Protective effects of kahweol and cafestol against hydrogen

peroxide-induced oxidative stress and DNA damage. Toxicol Lett. 2007;173:80–

87.

[28] Fürstenau CR, Souza ICC de, Oliveira MR de. The effects of kahweol, a diterpene

present in coffee, on the mitochondria of the human neuroblastoma SH-SY5Y cells

exposed to hydrogen peroxide. Toxicol In Vitro. 2019;61:104601.

[29] Grewal R, Reutzel M, Dilberger B, et al. Purified oleocanthal and ligstroside

protect against mitochondrial dysfunction in models of early Alzheimer's disease

and brain ageing. Exp Neurol. 2020;328:113248.

[30] Hagl S, Grewal R, Ciobanu I, et al. Rice bran extract compensates mitochondrial

dysfunction in a cellular model of early Alzheimer's disease. J Alzheimers Dis.

2015;43:927–938.

[31] Stadlmann S, Renner K, Pollheimer J, et al. Preserved Coupling of Oxidative

Phosphorylation But Decreased Mitochondrial Respiratory Capacity in IL-1β-

Treated Human Peritoneal Mesothelial Cells. CBB. 2006;44:179–186.

[32] Babylon L, Grewal R, Stahr P-L, et al. Hesperetin Nanocrystals Improve

Mitochondrial Function in a Cell Model of Early Alzheimer Disease. Antioxidants

(Basel). 2021;10. DOI: 10.3390/antiox10071003.

[33] Bustin SA, Benes V, Garson JA, et al. The MIQE guidelines: minimum

information for publication of quantitative real-time PCR experiments. Clin Chem.

2009;55:611–622.

[34] Dragicevic N, Delic V, Cao C, et al. Caffeine increases mitochondrial function and

blocks melatonin signaling to mitochondria in Alzheimer's mice and cells.

Neuropharmacology. 2012;63:1368–1379.

[35] Giorgi C, Marchi S, Simoes ICM, et al. Mitochondria and Reactive Oxygen

Species in Aging and Age-Related Diseases. Int Rev Cell Mol Biol. 2018;340:209–

344.

[36] Londzin P, Zamora M, Kąkol B, et al. Potential of Caffeine in Alzheimer's

Disease-A Review of Experimental Studies. Nutrients. 2021;13. DOI:

10.3390/nu13020537.

[37] Shen T, Lee J, Lee E, et al. Cafestol, a coffee-specific diterpene, is a novel

extracellular signal-regulated kinase inhibitor with AP-1-targeted inhibition of

prostaglandin E2 production in lipopolysaccharide-activated macrophages. Biol

Pharm Bull. 2010;33:128–132.

[38] Mukhopadhyay S, Mondal A, Poddar MK. Chronic administration of caffeine:

effect on the activities of hepatic antioxidant enzymes of Ehrlich ascites tumor-

bearing mice. Indian J Exp Biol. 2003;41:283–289.

[39] Lee C. Antioxidant ability of caffeine and its metabolites based on the study of

oxygen radical absorbing capacity and inhibition of LDL peroxidation. Clinica

Chimica Acta. 2000;295:141–154.

[40] Bichler J, Cavin C, Simic T, et al. Coffee consumption protects human

lymphocytes against oxidative and 3-amino-1-methyl-5H-pyrido4,3-bindole acetate

(Trp-P-2) induced DNA-damage: results of an experimental study with human

volunteers. Food Chem Toxicol. 2007;45:1428–1436.

[41] Li S, Geiger NH, Soliman ML, et al. Caffeine, Through Adenosine A3 Receptor-

Mediated Actions, Suppresses Amyloid-β Protein Precursor Internalization and

Amyloid-β Generation. J Alzheimers Dis. 2015;47:73–83.

[42] Cao C, Cirrito JR, Lin X, et al. Caffeine suppresses amyloid-beta levels in plasma

802   and brain of Alzheimer's disease transgenic mice. J Alzheimers Dis. 2009;17:681–

803   697.

804   [43] Butt MS, Sultan MT. Coffee and its consumption: benefits and risks. Crit Rev Food

805   Sci Nutr. 2011;51:363–373.

806   [44] Godos J, Pluchinotta FR, Marventano S, et al. Coffee components and

807   cardiovascular risk: beneficial and detrimental effects. Int J Food Sci Nutr.

808   2014;65:925–936.

809   [45] Socała K, Szopa A, Serefko A, et al. Neuroprotective Effects of Coffee Bioactive

810   Compounds: A Review. Int J Mol Sci. 2020;22. DOI: 10.3390/ijms22010107.

811   [46] Sharma VK, Singh TG. CREB: A Multifaceted Target for Alzheimer's Disease.

812   Curr Alzheimer Res. 2020;17:1280–1293.

813   [47] Chen Y, Huang X, Zhang Y, et al. Alzheimer's β-secretase (BACE1) regulates the

814   cAMP/PKA/CREB pathway independently of β-amyloid. J Neurosci.

815   2012;32:11390–11395.

816   [48] Teich AF, Nicholls RE, Puzzo D, et al. Synaptic therapy in Alzheimer's disease: a

817   CREB-centric approach. Neurotherapeutics. 2015;12:29–41.

818   [49] Echeverri D, Montes FR, Cabrera M, et al. Caffeine's Vascular Mechanisms of

819   Action. Int J Vasc Med. 2010;2010:834060.

820   [50] Orbán C, Vásárhelyi Z, Bajnok A, et al. Effects of caffeine and phosphodiesterase

821   inhibitors on activation of neonatal T lymphocytes. Immunobiology.

822   2018;223:627–633.

823   [51] Sanders O, Rajagopal L. Phosphodiesterase Inhibitors for Alzheimer's Disease: A

824   Systematic Review of Clinical Trials and Epidemiology with a Mechanistic

825   Rationale. J Alzheimers Dis Rep. 2020;4:185–215.

[52] Seo H-Y, Lee S-H, Lee J-H, et al. Kahweol activates the Nrf2/HO-1 pathway by

decreasing Keap1 expression independently of p62 and autophagy pathways. PLoS

One. 2020;15:e0240478.

[53] Hao W-R, Sung L-C, Chen C-C, et al. Cafestol Activates Nuclear Factor Erythroid-

2 Related Factor 2 and Inhibits Urotensin II-Induced Cardiomyocyte Hypertrophy.

Am J Chin Med. 2019;47:337–350.

[54] Hipkiss AR. Aging, Alzheimer's Disease and Dysfunctional Glycolysis; Similar

Effects of Too Much and Too Little. Aging Dis. 2019;10:1328–1331.

[55] Tauffenberger A, Fiumelli H, Almustafa S, et al. Lactate and pyruvate promote

oxidative stress resistance through hormetic ROS signaling. Cell Death Dis.

2019;10:653.

[56] Mintun MA, Vlassenko AG, Rundle MM, et al. Increased lactate/pyruvate ratio

augments blood flow in physiologically activated human brain. Proc Natl Acad Sci

U S A. 2004;101:659–664.

[57] Brooks GA. The Science and Translation of Lactate Shuttle Theory. Cell Metab.

2018;27:757–785.

[58] Zangari J, Petrelli F, Maillot B, et al. The Multifaceted Pyruvate Metabolism: Role

of the Mitochondrial Pyruvate Carrier. Biomolecules. 2020;10. DOI:

10.3390/biom10071068.

[59] Hassel B. Pyruvate carboxylation in neurons. J Neurosci Res. 2001;66:755–762.

[60] Katoh-Semba R, Keino H, Kashiwamata S. A possible contribution by glial cells to

neuronal energy production: enzyme-histochemical studies in the developing rat

cerebellum. Cell Tissue Res. 1988;252:133–139.

[61] Schormann N, Hayden KL, Lee P, et al. An overview of structure, function, and

regulation of pyruvate kinases. Protein Sci. 2019;28:1771–1784.

[62] Desagher S, Glowinski J, Prémont J. Pyruvate Protects Neurons against Hydrogen

Peroxide-Induced Toxicity. J. Neurosci. 1997;17:9060–9067.

[63] Wang X, Shen X, Yan Y, et al. Pyruvate dehydrogenase kinases (PDKs): an

overview toward clinical applications. Biosci Rep. 2021;41. DOI:

10.1042/BSR20204402.

[64] Wu J, Jin Z, Zheng H, et al. Sources and implications of NADH/NAD(+) redox

imbalance in diabetes and its complications. Diabetes Metab Syndr Obes.

2016;9:145–153.

[65] Long DM, Frame AK, Reardon PN, et al. Lactate dehydrogenase expression

modulates longevity and neurodegeneration in Drosophila melanogaster. Aging

(Albany NY). 2020;12:10041–10058.

[66] Ross JM, Öberg J, Brené S, et al. High brain lactate is a hallmark of aging and

caused by a shift in the lactate dehydrogenase A/B ratio. Proc Natl Acad Sci U S A.

2010;107:20087–20092.

[67] Rossi A, Rigotto G, Valente G, et al. Defective Mitochondrial Pyruvate Flux

Affects Cell Bioenergetics in Alzheimer's Disease-Related Models. Cell Rep.

2020;30:2332-2348.e10.

[68] Kumar Jha M, Jeon S, Suk K. Pyruvate Dehydrogenase Kinases in the Nervous

System: Their Principal Functions in Neuronal-glial Metabolic Interaction and

Neuro-metabolic Disorders. Current Neuropharmacology. 2012;10:393–403.

871