Immune cells employ intermittent integrin-mediated traction forces for 3D migration

Tina Czerwinski,[a] Lars Bischof,[a] David Böhringer,[a] Sibel Kara,[b] Ernst Wittmann,[a] Alexander Winterl,[a] Richard Gerum,[a] Gina Nusser,[b] Manuel Wiesinger,[d,e,f] Silvia Budday,[c] Anja Lux,[b] Caroline Voskens,[d,e,f] Ben Fabry,[a] and Christoph Mark[a,*]

[a]Friedrich-Alexander University Erlangen-Nürnberg, Department of Physics, Erlangen, Germany
[b]Friedrich-Alexander University Erlangen-Nürnberg, Department of Genetics, Erlangen, Germany
[c]Friedrich-Alexander University Erlangen-Nürnberg, Department of Mechanical Engineering, Erlangen, Germany
[d]Universitätsklinikum Erlangen, Department of Dermatology, Erlangen, Germany
[e]Friedrich-Alexander University Erlangen-Nürnberg, Deutsches Zentrum Immuntherapie (DZI), Erlangen, Germany
[f]Comprehensive Cancer Center Erlangen-European Metropolitan Area of Nürnberg (CCC-ER-EMN), Erlangen, Germany

*Correspondence should be addressed to C.M. (christoph.mark@fau.de)

To reach targets outside the bloodstream, immune cells can extravasate and migrate through connective tissue. During tissue infiltration, immune cells migrate in an amoeboid fashion, characterized by weak matrix adhesions and low traction forces, that allows them to achieve high migration speeds of up to 10 μm/min. How immune cells reconcile amoeboid migration with the need to overcome steric hindrance in dense matrices is currently not understood. Here we show that when confronted with steric hindrance, immune cells can switch from their default amoeboid migration mode to a highly contractile, mesenchymal-like migration mode. We use time-lapse confocal reflection microscopy to obtain simultaneous measurements of migration speed, directional persistence, and cell contractility in 3D biopolymer networks. We find that NK92 (natural killer) cells are highly mechanoresponsive and exert substantial acto-myosin driven, integrin-mediated contractile forces of up to 100 nN on the extracellular matrix during short contractile phases. This burst-like contractile behavior is also found in primary B, T, NK cells, neutrophils, and monocytes, and is specifically used by the cells to avoid getting stuck in narrow pores of the surrounding matrix. Our results demonstrate that steric hindrance guides the rapid regulation of integrin-mediated adhesion to the ECM in a large number of immune cell subtypes.

Introduction

Immune cells such as monocytes, neutrophilic granulocytes, as well as B cells, T cells, and natural killer (NK) cells can migrate through tissues up to 100 times faster than mesenchymal cells[1,2]. Immune cells typically use an amoeboid migration mode with limited or no proteolytic activity and limited or no specific (e.g. integrin-mediated) adhesion to the extracellular matrix (ECM)[3–5]. Therefore, these cells are not expected to generate substantial traction forces, implying that they do not form mature focal adhesion contacts and thus are not able to significantly pull on and rearrange ECM fibers[4,6]. However, since migration in three-dimensional (3D) environments requires the cell to overcome the resistive forces of the matrix[7–9], some mechanism of force transmission across the plasma membrane to the ECM must exist but has not yet been characterized in these cells.

The involvement of integrins in leukocyte migration in 3D matrices has been studied previously, and the general conclusion is that integrins are required during extravasation, but are not essential during tissue infiltration: when integrin function is blocked in T cells or when integrins are depleted in mouse dendritic cells, neutrophils, and B cells, migration is only possible in confined 3D environments but not on two-dimensional (2D) surfaces[2,4].

More specifically, integrin depletion in mouse leukocytes does not affect migration speed in 3D collagen gels, indicating that these cells do not require integrins for interstitial migration, whereas they do for migration on 2D substrates[2]. Dendritic cells can switch from integrin-mediated to integrin-independent movement without any change in migration speed or directional persistence[10]. Furthermore, integrin-mediated force coupling is not required for migration in constrained environments. The currently established notion is that integrins in immune cells act as a low-affinity frictional interface between cortical actin flow and the substrate, thereby facilitating cell locomotion[11–13].

In this study, we use fast time-lapse 3D traction force microscopy (Fig. 1) in combination with high-throughput 3D migration assays to revisit this well-established notion, which, as our data on large traction forces generated by immune cells suggest, is not wrong but also not complete. Consistent with prevailing theories, our data confirm that immune cells are able to migrate within 3D networks of randomly oriented collagen fibers most of the time without

Figure 1: Traction force microscopy of immune cell migration in 3D collagen gels. a: Confocal image series of a ex-vivo expanded NK cell (magenta, Calcein stain) migrating in a 1.2 mg/ml collagen gel (imaged with confocal reflection microscopy). The dashed red lines indicate the position of exemplary collagen fibers at t=0s, to illustrate the movement of the fibers due to forces applied by the NK cell. The dashed blue line indicates the outline of the NK cell at t=0s. Scale bar: 10μm. **b:** Temporal evolution of the contractility of a NK cell over the course of 10min. A typical contractile burst is visible during minutes 4-7. **c:** Measured 3D deformation field of the collagen fiber network that surrounds the NK cell at t=5min (corresponding to the red line in (b)) and at t=10min (corresponding to the dashed red line in (b)). Scale bar = 20μm. **d:** Same as in (c), but showing the reconstructed force field that is obtained from the measured deformation fields.

significant tractional forces, but that this amoeboid migration mode is frequently interrupted by short contractile bursts of up to ~60 nN in NK92 cells. We argue that this switch to a highly contractile mesenchymal-like migration mode allows immune cells to overcome the steric hindrance imposed by narrow pores in the ECM and to avoid getting stuck in the matrix (Supplementary Videos 1,2). We provide evidence for this hypothesis in two ways: first, we find a significant temporal correlation between cell contractility and cell speed as well as directional persistence of individual motile immune cells. Second, we find that by preventing strong integrin adhesion to the matrix, NK92 cells can still migrate in an amoeboid migration mode with unchanged migration speed, but are forced to take more frequent turns to avoid small pores and obstacles, and become stuck more frequently (reduced motile fraction).

Results

Contractility and mechanosensitivity of NK92 cells

To test the ability of immune cells to exert traction forces on the extracellular matrix, we employ high-speed 3D traction force microscopy (TFM). Here, we quantify the 3D matrix deformations surrounding individual NK92 cells that are

Figure 2: Force generation of NK92 cells in reconstituted collagen gels. a: Confocal reflection image through a 3D reconstituted collagen gel with a collagen concentration of 0.6 mg/ml (shear modulus: 101 Pa; left), 1.2 mg/ml (286 Pa; center), and 2.4 mg/ml (1155 Pa; right). Collagen fibers are visible in black. Scale bar: 20 μm. **b:** 99th percentile of the matrix deformations generated by NK92 cells for different collagen concentrations (0.6 mg/ml & 1.2 mg/ml: n=51 in 4 independent experiments; 2.4 mg/ml: n=46 in 3 independent experiments). **c:** Maximum cell contractility during a 24 min observation period. **d:** Fraction of contractile phases (time periods when cell contractility > 5 nN per cell). **e:** Fraction of motile NK92 cells for different collagen concentrations. A cell is defined as motile if the 5 min bounding-box around its migration path has a mean diagonal length of 6.5μm or greater. Error bars denote 1 SEM.

embedded in reconstituted collagen gels over a time course of 24 min at a time interval of 1 min. As expected, the maximum deformation that the cells exert onto the collagen matrix within the measurement time decreases with increasing collagen concentration and thus with matrix stiffness (Fig. 2a,b). However, since collagen is a highly non-linear material, deformation is not strictly proportional to contractility. This nonlinearity thus prohibits the use of deformations as a proxy for cell contractility. To test how the contractility of the NK92 cells changes in response to a varying matrix stiffness, we reconstruct the cell contractility from the measured deformation fields. Importantly, we employ a material model that specifically includes non-affine deformations and accurately captures the non-linear effects of strain stiffening and buckling of collagen fibers[8,14]. As immune cells are expected to migrate without substantial force generation most of the time, we focus on the maximum contractility that we obtain within the 24 min measurement period of each cell. We find that cell contractility increases with increasing matrix stiffness, reaching ~60 nN at the highest collagen concentration (2.4 mg/ml, corresponding to 1155 Pa shear modulus; Fig. 2c; Supplementary Fig. 1). Given the small size of NK92 cells, this contractility is surprisingly high - comparable to mesenchymal cancer cells that generate forces of ~100 nN[8,15].

The increase in cell contractility with increasing collagen concentration further reveals that NK92 cells are able to sense mechanical changes of their environment and

upregulate force generation in denser collagen gels. However, denser collagen gels not only attain an increased gel stiffness but also a decreased pore size (Supplementary Fig. 1). To confirm that NK92 cells indeed adapt cell contractility specifically in response to matrix stiffness and not to pore size, we repeat the experiment in a different collagen batch with approximately the same pore size but lower stiffness. Indeed, we find approximately 2-fold lower contractile forces in the 4-fold softer batch (Supplementary Fig. 2). This degree of mechano-responsiveness of cell contractility is reminiscent of mesenchymal cells such as fibroblasts or invasive cancer cells[16,17].

Figure 3: Dynamic variations of contractility during the migration of NK92 cells. a: Exemplary migration path of a NK92 cell migrating within a 3D collagen gel for 10min. Cell outlines are shown in red, darker patches indicate a longer retention time at that spot, lighter patches indicate fast changes of cell morphology. Scale bar: 10µm. **b:** Reconstructed cell contractility (blue), cell speed (orange), and directional persistence (green) for the exemplary NK92 cell shown in (a). **c:** Cross-correlation function between cell contractility and cell speed. Positive x-values indicate that speed follows contractility in time, negative x-values indicate that contractility follows cell speed. The correlation function is based on data from n=52 cells and considers all time steps with a positive contractility. The red rectangle indicates the right-shifted peak of the correlation function. Error intervals indicate 1sem obtained by bootstrapping. **d:** Same as in (c), but showing the cross-correlation function between contractility and directional persistence.

In contrast to fibroblasts and cancer cells, however, NK92 cells are not always contractile but instead display short contractile bursts. We find that NK92 cells spend about 20-25% of the time in these contractile phases (> 5nN) when cultured in low- and medium-concentration collagen gels (Fig. 2d). In high-concentration collagen gels with narrower pores, the time spent in contractile phases drops to ~10% (Fig. 2d). To explain this decrease, we consider the motile phases of NK92 cells. We draw a bounding-box around a cell's migration path during each 5 min period and determine its diagonal length (a measure of the exploration volume). A cell is classified as motile if the diagonal is greater than 6.5 µm on average over the 24 min observation time.

In low- and medium-concentration collagen gels, we find a significant fraction of motile cells, ranging between 10-30% (Fig. 2e). In high-concentration collagen gels, however, almost all cells remain non-motile (Fig. 2e). This behavior appears to follow the product of maximum contractility and fraction of contractile phases (Fig. 2c-e). Accordingly, successful migration appears to require force bursts with sufficient magnitude and frequency. The dramatic decrease in the motile fraction in the high-concentration collagen gel suggests that narrow pores cause excessive steric hindrance, which cannot be overcome by increased magnitude or frequency of traction forces.

Contractile phases correlate with increased cell speed and directional persistence

In high-concentration collagen gels, NK92 cells upregulate traction forces but cannot remain motile due to abundant narrow pores[18]. This leads to the question of whether

traction forces can enhance motility in medium (and low) concentration collagen gels where pore size is not a limiting factor. Therefore, we investigate whether individual bursts of contractility lead to a higher cell speed and a directionally more persistent migration path (Fig. 3a,b). To this end, we track the position of individual cells during the 3D traction force microscopy measurement and extract the momentary cell speed and turning angle between subsequent time steps (see methods; Fig. 3a,b). As we are interested in the systematic movement of the cell as a whole, we employ Bayesian filtering to suppress artifactual movements of the cell due to cell shape variations and imaging noise (see methods, Supplementary Figure 3)[19,20].

To identify common patterns in the interplay between cell contractility and cell motility of many individual NK92 cells, we compute cross-correlation functions between contractility and cell speed as well as directional persistence (Fig. 3c,d). Importantly, we find that, on average, cell speed and directional persistence increase with cell contractility, with a small time lag of 2 min, suggesting that NK92 cell motility benefits from contractile forces. This interplay of cell speed, directional persistence, and cell contractility closely resembles the gliding motion during mesenchymal migration of, for example, breast cancer cells, but on a faster time scale (cross-correlation functions fall to zero within 10-

15 min for NK92 cells, compared to 1 h for MDA-MB-231 cells[8]). This striking similarity in migration dynamics suggests that NK92 cells are able to switch to a mesenchymal-like migration mode to maintain their motility in challenging microenvironments.

Inhibition of acto-myosin-driven forces decreases the motile fraction of NK92 cells

If NK92 cells use traction forces to overcome steric hindrance in 3D collagen gels, we expect that inhibition of cellular force generation will result in a lower motile fraction as more cells get stuck in the narrow pores of the gel. To test this hypothesis, we downregulate acto-myosin contraction in NK92 cells by treating the cells with Rho kinase inhibitor or with blebbistatin. The aim is to reduce but not to completely inhibit cellular force generation so that the cells retain their ability for dynamic shape changes and migration. As expected, cellular force generation is significantly reduced, both in magnitude (Fig. 4a) and frequency of force peaks (Fig. 4b). In agreement with our hypothesis, we find that the fraction of motile NK92 cells is substantially reduced by ~50% when acto-myosin contraction is reduced (Fig. 4c). Importantly, the treatment does not affect cell viability (Supplementary Figure 4).

To confirm this finding, we perform high-throughput 3D migration assays in which NK92 cells are seeded into a collagen matrix and then imaged and tracked at low magnification for 5 minutes in multiple field-of-views. This technique allows us to acquire data on 1000-3500 individual cells per condition and experiment. We perform this additional assay for two reasons: first, there may be a selection bias toward elongated, motile cells in single cell 3D TFM measurements, as the experimenter must manually search for and select cells for imaging. Second, the high-throughput 3D migration assay allows us to perform paired experiments for all conditions and obtain data on a much larger population of cells.

Figure 4: Inhibition of acto-myosin-driven forces in NK92 cells decreases motile fraction, but not motility. a: Cell contractility, evaluated by taking the mean over the maximum contractility of each cell during a 24 min observation period (to capture the contractility during the short contractile bursts that the NK92 cells exhibit), for the control and for treatment with 10µM Y-27632 Rock-inhibitor and 3 µM blebbistatin in a 1.2 mg/ml collagen gel (Ctrl n=50, blebbistatin n=44, Rock inhibitor n=44 cells) in 3 independent experiments. **b:** Contractile fraction of the NK92 cells, calculated as the average fraction of time steps that the cells reach a contractility >5 nN. **c:** Fraction of motile NK92 cells. A cell is motile if the 5 min bounding-box around its migration path has a mean diagonal length of 6.5µm or greater. **d:** Same as in (c), but measured in the high-throughput 3D migration assay (n=3 independent experiments, n = 1000-3500 cells per measurement and condition, one tailed paired t-test, p=6.86*10^-5 for Rock inhibitor, p=0.0051 for blebbistatin). **e:** Mean speed of motile NK92 cells as measured in the high-throughput 3D migration assay, paired for all three conditions. Individual paired measurements are indicated by the black dots and lines (n=4 independent experiments; 1000-3500 cells are measured per measurement and condition). **f:** Same as in (e), but showing the mean directional persistence of motile NK92 cells. **g:** Density plot of cell speed and persistence under control conditions. Each point represents the mean cell speed and persistence of a 5 min trajectory of a single NK92 cell. The color shading represents the Gaussian kernel density estimate (n=4 independent experiments). Red dashed circle approximately indicates the motile fraction of the cell population that migrates at least 6.5 µm from its point of origin within 5 min. **h:** Same as in (g), but for NK92 cells treated with 10 µM Y-27632 Rock-inhibitor. **i:** Same as in (g), but for NK92 cells treated with 3 µM blebbistatin. **j:** Exemplary time-lapse minimum intensity projection sequence of a NK92 cell that remains stuck in the collagen matrix from 0 min to 50 min. Note that the cell changes its morphology frequently, but is unable to move. After 45 min, the cell abruptly becomes migratory again. Red points indicate the cell trajectory. The red contour shows the cell morphology and the pink arrows show the constriction of the cell due to the collagen pore (scale bar: 10 µm).

The high-throughput migration assay shows a lower motile fraction in all conditions compared to the 3D TFM experiment (Fig. 4d), confirming a selection bias towards motile cells in the single-cell 3D TFM experiment. However, the 50% decrease in the motile fraction after downregulation of acto-myosin contraction remains and is consistently

observed in all independent replicates of the experiment (Fig. 4d, g-i). Interestingly, for the remaining motile fraction, we find only a small ~10% decrease in cell speed after treatment (Fig. 4e), and no change in directional persistence (Fig. 4f). Thus, the combination of single-cell 3D TFM and high-throughput 3D migration assay shows that down-regulation of acto-myosin contraction in NK92 cells drastically reduces the number of motile cells in the population, while the remaining motile cells retain most of their motility. Taken together, this result confirms our notion that large cell tractions are critical for overcoming the steric hindrance when encountering narrow pores, but are not critical for amoeboid motion in less challenging

Figure 5: Traction forces of primary immune cells. a: Cell contractility, evaluated by taking the mean over the maximum contractility of each cell during the 24min observation period (to capture the contractility during the short contractile bursts) for different primary immune cells 1h (blue) and 24h (gray) after seeding the cells into the 3D collagen gels (B-cells: 'B', n=43/44; Monocytes: 'Mono', n=39/44; Neutrophils: 'Neutro', n=47/0; NK cells: 'NK', n=41/43; T cells: 'T', n=48/43 in 4 independent experiments). Each black dot represents the maximum contractility of an individual cell. **b:** Contractile fraction of different primary immune cells, calculated as the average fraction of time steps that the cells reach a contractility >5nN. **c:** Fraction of motile primary immune cells as measured in the high-throughput migration assay. **d:** Change in the mean fraction of time steps with a contractility >5nN (called "contractile fraction") within the first 24h after seeding the cells into the collagen gel, as a function of the change in the motile fraction of the cell population (as measured in the migration assay). The red dashed line indicates the identity line. **e:** Cell contractility of primary NK cells after incubation with IL-2 (gray) and without (control; blue), for different incubation periods prior to seeding the cells into the collagen gel. Each black dot represents the maximum contractility of an individual cell. **f:** Contractile fraction of primary NK cells after incubation with IL-2 (gray) and without (control; blue), for different incubation periods prior to seeding the cells into the collagen gel. Error bars denote 1 SEM in (a)-(c) and (e)-(f), and the 1-sigma credible interval obtained by bootstrapping in (d).

environments. This view is further supported by our observation that cells that become stuck are still undergoing rapid morphological changes and continue their migration path once they have succeeded to overcome the obstacle (Fig. 7j, Supplementary Videos 3,4).

Contractile forces of primary immune cells

The observation of short contractile phases is not limited to the cultured NK92 cell line. Rather, we find that most immune cell types are capable of generating appreciable traction forces. In particular, we find that without expansion, primary human B cells, monocytes, NK cells, T cells, and neutrophils all exhibit contractile phases on the order of ~5 nN (with occasional peaks around 20 nN) in which they spend 5-10% of the time when placed in a 1.2 mg/ml collagen matrix with a low stiffness of 85 Pa (Fig. 5a, b; Supplementary Fig. 2). We further confirm that traction forces facilitate migration of primary cells, as cell speed and directional persistence increase after force upregulation (Supplementary Figure 5). For all cell types, the fraction of motile cells remains lower compared to NK92 cells and is subject to significant donor-to-donor variation (Fig. 5c).

Over a time course of 24 h, B cells increase the frequency of contractile bursts and also increase their motile fraction, whereas monocytes decrease the frequency of contractile bursts and also decrease their motile fraction (Fig. 5b, c). We find a positive correlation between the 24 h change in burst frequency and the change in motile fractions amongst different cell types (Fig. 5d), highlighting the importance of contractile force bursts in the 3D migration process for a diverse set of primary immune cells.

Migration and contractile forces of ex-vivo expanded NK cells

Ex-vivo expanded NK cells are currently explored for use in adoptive immunotherapy against blood-borne cancers[21,22]. In this type of therapy, patient-derived primary peripheral blood mononuclear cells (PBMCs) are expanded into a population of highly activated NK-cells. Cells are activated by adding IL-2 and culturing them in the presence of IL-15-secreting feeder cells. IL-2 activation is known to promote cell motility[23], and we reasoned that this effect might be supported by an increased contractility. We confirm that the motile fraction of primary NK cells without IL-2 decreases over a 72 h time period, but remains high when stimulated with IL-2 (Supplementary Figure 6). Similarly, the magnitude and frequency of contractile bursts increase within 1 h after IL-2 addition and remain stable for the following 72 h, whereas contractile forces tend towards zero over this time period in non-stimulated cells (Fig. 5e,f). These data suggest that the enhanced migration of NK cells after IL-2 activation is indeed facilitated by increased cellular force generation.

To relate these findings to a clinically relevant cell product, we next investigated NK cells that are expanded for 14 days in the presence of high concentrations of IL-2 and IL-15-

secreting feeder cells. In general, we find that expanded NK-cells are similarly motile and contractile as IL-2 stimulated primary cells and NK92 cells. However, we find some notable qualitative differences. Specifically, the maximum contractile force and the contractile fraction of expanded NK cells show a bi-phasic response to collagen concentration, reaching the highest values at an intermediate (1.2 mg/ml) collagen concentration (Fig. 6a-d). The motile fraction of ex-vivo expanded NK cells is roughly proportional to the product of the magnitude and frequency of force bursts (c.f. Fig. 2c-e), similar to our findings in NK92 cells. Notably, expanded NK cells retain a higher motile fraction of ~30% in the high-concentration collagen gel, likely due to their smaller size compared to NK92 cells (Supplementary Figure 7).

In ex-vivo expanded NK cells, we find an even more pronounced relationship between traction forces and migration compared to NK92 cells: phases of elevated contractility are followed by phases of elevated speed and persistence lasting for ~10 min (Fig. 6e-h). Phases of elevated cell speed, by contrast, are followed by phases of reduced contractility lasting for ~15 min (Fig. 6e-h). Hence, the cell preferentially upregulates traction forces when it becomes slower or less persistent.

Figure 6: Dynamic variations of contractility during the migration of ex-vivo expanded NK cells. a: 99th percentile of the matrix deformations generated by ex-vivo expanded NK cells for different collagen concentrations (0.6mg/ml: n=27; 1.2mg/ml: n=16; 2.4mg/ml: n=16 in 2 independent experiments) **b:** Maximum cell contractility during a 24 min observation period. **c:** Fraction of contractile phases (time periods when cell contractility > 5 nN). **d:** Fraction of motile NK92 cells for different collagen concentrations. A cell is defined as motile if the 5 min bounding-box around its migration path has a mean diagonal length of 13μm or greater. **e:** Exemplary migration path of an ex-vivo expanded NK cell migrating within a 3D collagen gel for 10min. Cell outlines are shown in red, darker patches indicate a longer retention time at that spot, lighter patches indicate a fast cell movement. Scale bar: 10μm. **f:** Cross-correlation function between cell contractility and cell speed. Positive x-values indicate that speed follows contractility in time, negative x-values indicate that contractility follows cell speed. The correlation function is based on data from n=64 cells and considers all time steps with a positive contractility. The solid red rectangle indicates the right-shifted peak of the correlation function. The dashed red rectangle marks the negative correlation between contractility following cell speed. Error intervals indicate 1sem obtained by bootstrapping. **g:** Reconstructed cell contractility (blue), cell speed (orange), and directional persistence (green) for the exemplary NK cell shown in (e). **h:** Same as in (f), but showing the cross-correlation function between contractility and directional persistence.

3D NK cell motility depends on cell-matrix adhesion

The reduced motile fraction of NK cells after force inhibition, and the correlations of contractility versus cell speed, provide strong but indirect evidence for our hypothesis that immune cells use traction forces to overcome steric hindrance during 3D migration. Traction forces require strong adhesion to the matrix[24,25]. If NK cells use adhesion-mediated traction forces only when confronted with steric hindrance, we predict that any down-regulation of cell-matrix adhesion will result in a reduced motile fraction (more cells get stuck) and a reduced directional persistence (cells take more turns to avoid obstacles). However, we do not expect cell speed to be significantly affected, as motile cells should still be able to migrate in a non-adhesive, amoeboid fashion as long as they do not encounter narrow pores within the matrix.

We approach the manipulation of cell-matrix adhesion by manipulating both, matrix adhesiveness and the ability of the cells to form adhesions. First, we add the adhesion-inhibiting surfactant pluronic to polymerized collagen gels to reduce direct contacts between the cell membrane and individual collagen fibers, forcing the NK cells into a non-adhesive migration mode[26]. For NK92 cells, pluronic treatment reduces the fraction of motile cells by ~25% and the directional persistence of the remaining motile cells also by ~25% (Fig. 7a, Supplementary Figure 4). Consistent with our hypothesis, the remaining motile cells maintain their full migratory speed (Fig. 7a). Second, we seed NK92 cells into non-adhesive carbomer gels (Fig. 7b). These gels are composed of cross-linked polymers and provide a soft elastic microenvironment containing dispersed stiff inclusions[15,27,28] (Supplementary Figure 8; Supplementary Video 5). The relative changes in NK92 cell migration in carbomer gels confirm the pluronic measurements, but show a much more pronounced decrease in the motile fraction and directional

persistence of motile cells by >75% (Fig. 7b). Strikingly, the migration speed of the remaining motile cells remains unchanged (Fig. 7b).

We further test how NK cell migration changes when cells are treated with the calcium chelator EDTA. EDTA binds calcium and magnesium ions that are required for the proper function of several adhesion proteins, including integrins[29]. For EDTA-treated NK92 cells in collagen gels, we again find that the fraction of motile cells is reduced by ~25%, the directional persistence of the remaining motile cells is reduced to a lesser extent, and the migration speed of the remaining motile cells is not affected (Fig. 7c). While the effects of pluronic, carbomer, and EDTA differ on a quantitative level, they all reflect the same qualitative effect of reduced cell-matrix adhesion: fewer cells remain motile and the remaining motile cells have to make more turns within their microenvironment.

For ex-vivo expanded NK cells, the response to manipulation of cell-matrix adhesions is more nuanced. Again, pluronic treatment results in a substantial reduction of the motile fraction by >50%, but the remaining motile cells retain not only their full migration speed but also their full directional persistence (Fig. 7d). In carbomer gels and after EDTA treatment, ex-vivo expanded NK cells respond similar to NK92 cells, with a large decrease in motile fraction, a less pronounced decrease in directional persistence, and a smaller decrease in cell speed (Fig. 7e, f). The difference in migratory changes after inhibition of cell-matrix adhesions between NK92 cells and ex-vivo expanded NK cells may be attributed to the smaller size of ex-vivo expanded NK cells with respect to pore size and spacing between obstacles in the matrix (Supplementary Fig. 1,5,6).

Pluronic, carbomer, and EDTA broadly inhibit cell adhesion to the matrix without specificity for particular adhesion molecules. Since the dynamic force regulation of NK cells is reminiscent of mesenchymal cells, we further hypothesize that force transmission to the ECM in NK cells is integrin-mediated. To test this hypothesis, we treat NK92 cells with antibodies that specifically block or activate β_1 and β_2 integrin subunits. Integrin β_1 (CD29) is expressed on many cultured immune cell lines, including THP1 monocytic cells, Daudi B cells, the NK92 cell line, and Jurkat T cells (Fig. 7g). Integrin β_2 (CD18) shows a 5-fold higher expression level in NK92 cells compared to integrin β_1. The expression of β_2 is at least 8-fold higher in NK92 cells compared to the other tested immune cell lines (Fig. 7h). Therefore, we expect that NK92 cells mainly rely on integrin β_2 for adhesion[30], and that blocking integrin β_2 will affect the motile fraction more than blocking integrin β_1. Indeed, 3D migration assays confirm that blocking integrin β_1 decreases the motile fraction of NK92 cells by ~10%, whereas blocking integrin β_2 results in a decrease of ~30% (Fig. 7i, Supplementary Figure 4). As in the experiments where adhesion is blocked in a nonspecific manner, we again find that directional persistence is

Figure 7: Inhibiting cell-matrix adhesion suppresses NK cell motility. a: Relative change of cell speed, persistence, and motile fraction of NK92 cells, measured in high-throughput 3D migration assays (n=3 independent experiments) after inhibiting cell-matrix adhesion externally by the addition of pluronic to collagen gels (hindering cells from making contact with collagen fibers; orange; p=0.02 for the motile fraction, p=0.94 for cell speed, p=0.08 for persistence). **b:** Same as in (a), but for NK92 cells in non-adhesive carbomer gels (n=3; green; p<0.001 for the motile fraction, p=0.90 for cell speed, p=0.001 for persistence). **c:** Same as in (a), but for EDTA-treated NK92 cells in collagen (n=3; blue; p=0.04 for the motile fraction, p=0.56 for cell speed, p=0.38 for persistence), limiting cell-matrix adhesion internally by removing calcium ions that integrins require to maintain adhesion to the matrix. **d:** Same as in (a), but for expanded, primary NK cells (n=2, p=0.24 for the motile fraction, p=0.19 for cell speed, p=0.69 for persistence). **e:** Same as in (b), but for ex-vivo expanded NK cells (n=8, p=0.09 for the motile fraction, p=0.13 for cell speed, p=0.03 for persistence). **f:** Same as in (c), but for expanded, primary NK cells (n=7, p=0.03 for the motile fraction, p=0.11 for cell speed, p=0.03 for persistence). **g:** Delta mean fluorescence intensity of CD29 (β_1 integrin) of immune cell lines (THP1, Daudi, NK92 and Jurkat cells), measured by flow cytometry (n=3). **h:** Same as in (g), but for CD18 (β_2 integrin) (n=3). **i:** Same as in (a), but for β_1 (blue; p=0.01 for the motile fraction, p=0.46 for cell speed, p=0.10 for persistence) and β_2 (orange; p=0.05 for the motile fraction, p=0.26 for cell speed, p=0.02 for persistence) blocking antibodies treated NK92 cells in collagen (n = 3). **j:** Same as in (i), but for β_1 (blue) activating antibody treated cells in collagen (n = 3; p=0.37 for the motile fraction, p=0.37 for cell speed, p=0.59 for persistence). Statistical significance is tested using a paired ratio-t test.

significantly lower after integrin blocking. Notably, we find that cell speed is slightly increased after blocking β-integrins, likely because any type of integrin-mediated adhesion slows cells during their default amoeboid migration.

Treatment of NK92 cells with an activating integrin β_1 antibody leaves cell motility unchanged across all parameters (Fig. 7j). While we see a small increase in directional persistence and motile fraction, these changes

are not statistically significant. This additional finding shows that NK cell motility is not a linear function of integrin activation, but that integrin activity is likely tightly regulated to achieve optimal motility in challenging microenvironments[13]. In summary, manipulation of NK cell adhesion to the ECM demonstrates that NK cells generally achieve rapid migration in 3D biopolymer networks independent of adhesion to the ECM, but are critically dependent on adhesion to maintain motility and directional persistence in the presence of steric hindrance.

Disruption of the microtubule network increases traction forces and motile fraction

Recent reports have demonstrated that the migration of T cells and their engagement with target cells can be improved by treatment with nocodazole [31,32]. Nocodazole inhibits the polymerization of tubulin and thus destabilizes the microtubule network in cells. Microtubule destabilization has two effects: first, the nuclear deformability is increased[31], and second, acto-myosin contraction is increased[32,33]. The current understanding is that enhanced immune cell migration under nocodazole treatment is achieved by an increased nuclear deformability as it eases the transition of the cell through narrow pores[2,31], and by increasing cortical contraction and thus facilitating a fast, contact-guided, amoeboid migration mode in "2.5D" nano-structured surfaces[32]. Here, we provide further context on the effect of nocodazole by investigating cellular force generation and migration of NK92 cells under treatment in 3D collagen gels.

We find that nocodazole treatment substantially increases traction forces of NK92 cells by more than 100% (Fig. 8a). Moreover, treated cells employ traction forces greater than 5 nN more often, in more than 30% of all measured time steps, compared to 15% for untreated cells (Fig. 8b). In accordance with our hypothesis that traction forces help cells to overcome steric hindrance, we find that nocodazole treatment of NK92 cells significantly increases the fraction of motile cells (Fig. 8c). Strikingly, we again see that cell speed of the motile cells is not affected by treatment (Fig. 8d), indicating once more that traction forces do not make cells faster, but instead prevent them from getting stuck. Directional persistence is significantly increased after nocodazole treatment (Fig. 8e), further underlining that higher traction forces alleviate the need to take turns to avoid obstacles within the ECM.

Finally, we investigate the effect of an initial nocodazole treatment of NK92 cells on their long-term migration behavior. We find that the initial, strong increase in the motile fraction and in persistence has a short life time of < 6 h (Fig. 8f). Importantly, after 18 h measurement time, the majority of all treated cells have become non-motile, whereas the motile fraction of the control cells remains stable (Fig. 8f). The short-term positive effect of nocodazole on NK cell migration thus reverts on longer time scales, leaving most cells in a non-motile state.

Figure 8: Disruption of microtubule network in NK92 cells increases traction forces and motile fraction. a: Maximum cell contractility of NK92 cells during a 24 min observation period for the control and for treatment with 10 μM nocodazole (control: 'Ctrl' n=46; nocodazole: 'Noco' n=49 cells) in 1.2 mg/ml collagen in 4 independent experiments. **b:** Contractile fraction of the NK92 cells, calculated as the average fraction of time steps that the cells reach a contractility >5 nN. **c:** Fraction of motile NK92 cells measured in the high-throughput 3D migration assay, paired for both conditions. Individual paired measurements are indicated by the black dots and lines (n=4 independent experiments; 3000-4000 cells are measured per measurement and condition, one tailed paired t-test, p=0.001). **d:** Same as in (c), but showing the mean speed of motile NK92 cells (one tailed paired t-test, p=0.409). **e:** Same as in (d), but showing the mean directional persistence of motile NK92 cells (one tailed paired t-test, p=0.005). **f:** Density plot of cell speed and persistence for NK92 cells under control conditions (top row) or for NK92 cells treated with 10 μM nocodazole (bottom row), for different measurement times (0 - 18h) after seeding untreated and treated NK92 cells in collagen. Each point represents the mean cell speed and persistence of a 5 min trajectory of a single NK92 cell. The color shading represents the Gaussian kernel density estimate (n=1 independent experiments, 1500-2000 cells are measured per condition for each measurement time).

Discussion

We have shown that immune cells are capable of exerting significant traction forces on the ECM, comparable to the forces exerted by motile mesenchymal cells, taking into account the smaller size of immune cells. While we confirm the prevailing understanding that immune cells migrate in an amoeboid or friction-mediated manner without significant traction forces[4,12], immune cells are able to switch to a highly contractile migratory mode, achieving non-negligible contractility in 10-20% (NK92 cells) and 10-40% (expanded NK cells) of the total measurement time. In particular, NK92 cells show a pronounced mechanosensitivity similar to mesenchymal cells and significantly increase their contractility with increasing matrix stiffness.

Both the migration speed and the directional persistence of individual NK cells increase dynamically during and after short contractile bursts. This suggests that the switch from an amoeboid to a contractile migration mode is triggered

when the cell encounters increased steric hindrance – possibly similar to the integrin-driven extravasation of immune cells. This synchronization of cell contractility, persistence, and cell speed is reminiscent of mesenchymal migration of motile cancer cells, which also rely on traction forces to overcome steric hindrance in 3D fiber networks[8,9].

In support of the hypothesis that immune cells use traction forces to overcome steric hindrance, we have further shown that inhibition of the acto-myosin contraction in NK92 cells dramatically reduces both contractility and the fraction of motile cells, but has little effect on the migration speed and directional persistence of the remaining motile cells. These results suggest that unless a cell experiences steric hindrance (for example, by migrating through a narrow pore), it retains its default exploratory behavior and amoeboid migration mode. However, with limited acto-myosin contraction, cells are more likely to get stuck, resulting in a significantly reduced motile fraction.

The combination of amoeboid migration and occasional contractile bursts is not specific to NK cells. Non-activated primary B, T, and NK cells, as well as monocytes and neutrophils, are all able to exert traction forces on the ECM, although less frequently and not as strong as NK92 cells or expanded ex-vivo NK cells. Interestingly, cellular force generation appears to be closely linked to the adaptation process that immune cells undergo after being seeded into a 3D collagen matrix. Immune cell types that adapt well and increase their motile fraction over 24 hours after seeding also increase the frequency of force bursts, and vice versa. Upon activation with IL-2, primary NK cells increase both the frequency and magnitude of force bursts. Thus, traction force generation may be a key determinant of immune cell infiltration and retention in mechanically challenging environments.

Because large traction forces require a sufficiently strong coupling of the cell's contractile machinery to the extracellular matrix, it follows that immune cell migration in 3D biopolymer networks depends on cell-matrix adhesion. We have used several methods to inhibit cell adhesion to the ECM. In a non-adhesive environment, either by making collagen fibers non-adhesive using pluronic or by seeding cells in non-adhesive carbomer gels, we find that the speed of NK92 cells is unchanged, but more cells get stuck, and the remaining motile cells take more turns within the matrix. Treating cells with EDTA to weaken adhesion in a non-specific manner confirms these findings. These effects are qualitatively replicated in ex-vivo expanded NK cells. In addition, we have shown that the migration of NK92 cells in 3D fiber networks depends on integrin-mediated adhesion to the matrix. Specifically, we find that blocking integrin β1 and β2 again results in a substantial decrease in directional persistence and motile fraction, while cell speed is not reduced. Importantly, the decrease in motile fraction is greater when integrin β2 is blocked compared to β1,

consistent with the expression level of the receptors in NK92 cells.

Finally, we have shown that nocodazole-induced depolymerization of microtubules in NK92 cells results in a substantial, short-term increase in traction forces, motile fraction, and directional persistence, while leaving cell speed of the motile cells unchanged. These results confirm a previously reported increase of the *overall* cell speed under treatment[31,32], but provide a more nuanced interpretation: not the cell speed of the motile cells changes under treatment, but the fraction of motile cells increases. Hence, nocodazole treatment does not make cells faster but prevents them from getting stuck. Our results suggest that apart from the previously reported increased nuclear deformation[2,31], traction forces play a crucial role in the nocodazole-induced enhancement of migration. These in-vitro experiments have implications for current cancer therapies: taxane-based chemotherapies stabilize microtubules to interfere with cell division, but they may further impair tissue- and tumor-infiltration by immune cells[32]. Different chemotherapy agents, such as vinblastine, instead interfere with cell division by destabilizing microtubules, and may thus further enhance immune cell migration[31], at least on short time-scales. Functional biophysical assays that quantify migration behavior and mechanical interactions between cells and the ECM thus may help to better understand side effects and synergetic effects of existing anti-cancer therapies on immune cell behavior.

In conclusion, this work shows that immune cells are able to switch to a mesenchymal-like migration mode during 3D migration in biopolymer networks. This mesenchymal-like migration mode is characterized by integrin-mediated adhesion to the matrix and large, intermittent traction forces that the cells use to overcome steric hindrance.

References

1. Springer, T. A. Traffic signals for lymphocyte recirculation and leukocyte emigration: The multistep paradigm. *Cell* **76**, 301–314 (1994).

2. Lämmermann, T. *et al.* Rapid leukocyte migration by integrin-independent flowing and squeezing. *Nature* **453**, 51–55 (2008).

3. Friedl, P., Noble, P. B. & Zänker, K. S. T lymphocyte locomotion in a three-dimensional collagen matrix. Expression and function of cell adhesion molecules. *J. Immunol.* **154**, 4973–4985 (1995).

4. Friedl, P., Entschladen, F., Conrad, C., Niggemann, B. & Zänker, K. S. CD4+ T lymphocytes migrating in three-dimensional collagen lattices lack focal adhesions and utilize β1 integrin-independent strategies for polarization, interaction with collagen fibers and locomotion. *Eur. J. Immunol.* **28**, 2331–2343 (1998).

5. Wolf, K., Müller, R., Borgmann, S., Bröcker, Eva.-B. & Friedl, P. Amoeboid shape change and contact guidance: T-lymphocyte crawling through fibrillar collagen is independent of matrix remodeling by MMPs and other proteases. *Blood* **102**, 3262–3269 (2003).

6. Friedl, P., Borgmann, S. & Bröcker, E.-B. Amoeboid leukocyte crawling through extracellular matrix: lessons from the *Dictyostelium* paradigm of cell movement. *J. Leukoc. Biol.* **70**, 491–509 (2001).

7. Mierke, C. T., Frey, B., Fellner, M., Herrmann, M. & Fabry, B. Integrin α5β1 facilitates cancer cell invasion through enhanced contractile forces. *J. Cell Sci.* **124**, 369–383 (2011).

8. Steinwachs, J. *et al.* Three-dimensional force microscopy of cells in biopolymer networks. *Nat. Methods* **13**, 171–176 (2016).

9. Cóndor, M. *et al.* Breast Cancer Cells Adapt Contractile Forces to Overcome Steric Hindrance. *Biophys. J.* **116**, 1305–1312 (2019).

10. Renkawitz, J. *et al.* Adaptive force transmission in amoeboid cell migration. *Nat. Cell Biol.* **11**, 1438–1443 (2009).

11. Hons, M. *et al.* Chemokines and integrins independently tune actin flow and substrate friction during intranodal migration of T cells. *Nat. Immunol.* **19**, 606–616 (2018).

12. Yamada, K. M. & Sixt, M. Mechanisms of 3D cell migration. *Nat. Rev. Mol. Cell Biol.* **20**, 738–752 (2019).

13. Fowell, D. J. & Kim, M. The spatio-temporal control of effector T cell migration. *Nat. Rev. Immunol.* **21**, 582–596 (2021).

14. Böhringer, D. *et al. Dynamic traction force measurements of migrating immune cells in 3D matrices.* http://biorxiv.org/lookup/doi/10.1101/2022.11.16.516758 (2022) doi:10.1101/2022.11.16.516758.

15. Mark, C. *et al.* Cryopreservation impairs 3-D migration and cytotoxicity of natural killer cells. *Nat. Commun.* **11**, 5224 (2020).

16. Discher, D. E., Janmey, P. & Wang, Y. Tissue Cells Feel and Respond to the Stiffness of Their Substrate. *Science* **310**, 1139–1143 (2005).

17. Paszek, M. J. *et al.* Tensional homeostasis and the malignant phenotype. *Cancer Cell* **8**, 241–254 (2005).

18. Wolf, K. *et al.* Physical limits of cell migration: Control by ECM space and nuclear deformation and tuning by proteolysis and traction force. *J. Cell Biol.* **201**, 1069–1084 (2013).

19. Metzner, C. *et al.* Superstatistical analysis and modelling of heterogeneous random walks. *Nat. Commun.* **6**, 7516 (2015).

20. Mark, C. *et al.* Bayesian model selection for complex dynamic systems. *Nat. Commun.* **9**, 1803 (2018).

21. Ruggeri, L. *et al.* Effectiveness of Donor Natural Killer Cell Alloreactivity in Mismatched Hematopoietic Transplants. *Science* **295**, 2097–2100 (2002).

22. Laskowski, T. J., Biederstädt, A. & Rezvani, K. Natural killer cells in antitumour adoptive cell immunotherapy. *Nat. Rev. Cancer* **22**, 557–575 (2022).

23. Olofsson, P. E. *et al.* Distinct Migration and Contact Dynamics of Resting and IL-2-Activated Human Natural Killer Cells. *Front. Immunol.* **5**, (2014).

24. Balaban, N. Q. *et al.* Force and focal adhesion assembly: a close relationship studied using elastic micropatterned substrates. *Nat. Cell Biol.* **3**, 466–472 (2001).

25. Riveline, D. *et al.* Focal Contacts as Mechanosensors: Externally Applied Local Mechanical Force Induces Growth of Focal Contacts by an mDia1-dependent and ROCK-independent Mechanism✪. *J. Cell Biol.* **153**, (2001).

26. Tharmalingam, T., Ghebeh, H., Wuerz, T. & Butler, M. Pluronic Enhances the Robustness and Reduces the Cell Attachment of Mammalian Cells. *Mol. Biotechnol.* **39**, 167–177 (2008).

27. Kim, J.-Y., Song, J.-Y., Lee, E.-J. & Park, S.-K. Rheological properties and microstructures of Carbopol gel network system. *Colloid Polym. Sci.* **281**, 614–623 (2003).

28. Bhattacharjee, T. & Angelini, T. E. 3D T cell motility in jammed microgels. *J. Phys. Appl. Phys.* **52**, 024006 (2019).

29. Reversat, A. *et al.* Cellular locomotion using environmental topography. *Nature* **582**, 582–585 (2020).

30. Garnotel, R. *et al.* Human Blood Monocytes Interact with Type I Collagen Through αxβ2 Integrin (CD11c-CD18, gp150-95). *J. Immunol.* **164**, 5928–5934 (2000).

31. Zhao, R. *et al.* Targeting the Microtubule-Network Rescues CTL Killing Efficiency in Dense 3D Matrices. *Front. Immunol.* **12**, 729820 (2021).

32. Tabdanov, E. D. *et al.* Engineering T cells to enhance 3D migration through structurally and mechanically complex tumor microenvironments. *Nat. Commun.* **12**, 2815 (2021).

33. Rape, A., Guo, W. & Wang, Y. Microtubule depolymerization induces traction force increase through two distinct pathways. *J. Cell Sci.* **124**, 4233–4240 (2011).

34. Imai, C., Iwamoto, S. & Campana, D. Genetic modification of primary natural killer cells overcomes inhibitory signals and induces specific killing of leukemic cells. *Blood* **106**, 376–383 (2005).

35. Voskens, C. J. *et al.* Ex-vivo expanded human NK cells express activating receptors that mediate cytotoxicity of allogeneic and autologous cancer cell lines by direct recognition and antibody directed cellular cytotoxicity. *J. Exp. Clin. Cancer Res.* **29**, 134 (2010).

36. Lang, N. R. *et al.* Estimating the 3D Pore Size Distribution of Biopolymer Networks from Directionally Biased Data. *Biophys. J.* **105**, 1967–1975 (2013).

37. Krauss, P., Metzner, C., Lange, J., Lang, N. & Fabry, B. Parameter-Free Binarization and Skeletonization of Fiber Networks from Confocal Image Stacks. *PLoS ONE* **7**, e36575 (2012).

38. Ronneberger, O., Fischer, P. & Brox, T. U-Net: Convolutional Networks for Biomedical Image Segmentation. Preprint at http://arxiv.org/abs/1505.04597 (2015).

39. Cóndor, M., Steinwachs, J., Mark, C., García-Aznar, J. M. & Fabry, B. Traction Force Microscopy in 3-Dimensional Extracellular Matrix Networks. *Curr. Protoc. Cell Biol.* **75**, (2017).

40. Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Comput. Sci. Eng.* **9**, 90–95 (2007).

41. Gerum, R. pylustrator: code generation for reproducible figures for publication. *J. Open Source Softw.* **5**, 1989 (2020).

Acknowledgements

This work was supported by the National Institutes of Health Grant HL120839. We acknowledge the support by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) project number 460333672 – CRC 1540 Exploring Brain Mechanics (subprojects A01, B01, and C05) to LB, SB and BF. We acknowledge the support by the DFG (FA 336-12/1, TRR-SFB 225) project 326998133 (subprojects A01, B09, and C02) to DB, SB and BF. We thank Dario Campana, Andreas Baur, and Felix Rico for providing cell lines. This paper was typeset with the bioRxiv word template by @Chrelli: www.github.com/chrelli/bioRxiv-word-template

Author contributions

C.M., B.F., A.L., C.V. and T.C. designed the study. S.K., G.N., and L.B. performed primary immune cell isolation and FACS analysis. M.W. and C.V. isolated PBMCs from peripheral blood and performed ex-vivo NK cell expansion. T.C. and R.G. developed, T.C performed and analyzed short and long-term 3-D migration assays in collagen and carbomer. L.B. and D.B. performed, L.B., D.B., and R.G. analyzed 3-D contraction force microscopy assays. S.B. and D.B. performed collagen and carbomer rheology measurements. E.W. and A.W. developed the neural network-based cell segmentation. T.C., S.K., D.B., L.B. and C.M. performed data analysis. T.C. and C.M. generated the figures. T.C., B.F. and C.M. wrote the manuscript.

Competing interest statement

The authors declare no competing interests.

Materials and Methods

NK92, Jurkat, Daudi and THP1 cell lines

NK-92 cells (purchased from ATCC CRL-2407) are cultured for three weeks prior to measurements in in Alpha-MEM medium (Stemcell Technologies) with 15% fetal calf serum (Sigma), 15% horse serum (Gibco), 500 IU/ml human interleukin-2 (IL-2) cytokine (Proleukin S, Aldesleukin, Novartis Pharma, cat. no. 02238131) and 1% penicillin-streptomycin solution (10.000 Units/ml Penicillin, 10.000 µg/ml Streptomycin, Gibco). Jurkat cells (gift from Prof. A. Baur, Department of Molecular Dermatology and Extracellular Vesicle Analysis, Erlangen, Germany), THP1 cells (gift from F. Rico, Aix-Marseille Université, CNRS, Inserm, LAI, Turing center for living systems), and Daudi cells (gift from A. Lux, Department of Genetics, Erlangen, Germany) are cultured for three weeks prior to measurements in RPMI 1640 medium (Gibco) with 10% fetal calf serum, 1% Natrium Pyruvate (Gibco), 1% 100x MEM NEAA (Gibco), 0.3% 1M HEPES (Gibco) and 1% penicillin-streptomycin solution (hereafter called R10 medium).

Human primary immune cell isolation and storage

Human peripheral blood is collected from healthy donors (Department of Transfusion Medicine and Hemostaseology, University Clinics Erlangen, Germany), with approved consent of all participants and the ethics committee (22-320-Bp) of the university. Immune cells are isolated by density gradient centrifugation using Pancoll centrifugation media (PAN-Biotech, Aidenbach, Germany). Monocytes, NK cells, B cells, and T cells are

extracted from the peripheral blood mononuclear cell (PBMC) layer (second layer) with negative magnetic-activated cell sorting (MACS) using the following Biolegend kits: cat. no. 480060 for monocytes, cat. no. 480054 for NK cells, cat. no. 480082 for B cells, cat. no. 480022 for T cells. Granulocytes (mostly neutrophils) are extracted from the bottom layer, whereby the erythrocytes are eliminated by water lysis for 30 s, followed by the addition of 10 vol% of 10x PBS to stop lysis. After cell isolation, monocytes, NK cells, B cells and T cells are stored in R10 medium at 4°C for 24 h, whereas neutrophils are used immediately.

Flow cytometry analysis

A purity of at least 85% for all isolated primary immune cell subsets (NK, T cells, B cells, monocytes and neutrophils) is confirmed by flow cytometry. Primary immune cells are identified by antibodies against CD45 (Biolegend, cat. no. 368518), a common marker of leukocytes, in combination with a strong forwards and side scatter (for neutrophils) or a second cell-type specific surface marker (CD3 (Biolegend, cat. no. 300408) for T cells, CD19 (Biolegend, cat. no. 302216) for B cells, CD56 (Biolegend, cat. no. 304608) for NK cells and CD33 (Biolegend, cat. no. 303304) for monocytes). Dead cells and cell fragments are identified based on DAPI staining (dilution 1:2,500 of a 5 mg/ml stock solution in PBS with 2% FCS and 0.05% NaN3, AppliChem, Darmstadt, cat. no. 28718-90-3) and excluded from the analysis. Flow cytometry is further used to determine the expression profile of the integrins CD29 (PE anti-human CD29 antibody; Biolegend, cat. no. 303003) and CD18 (PE anti-human CD18 antibody; Biolegend, cat. no. 302107) of the cultured immune cell lines (Fig. 7e - f). All measurements are performed on a FACS Canto II (BD Biosciences), and data are analyzed using FACSDiva and FlowJo Software.

NK cell IL-2 activation

After 24 h storage at 4°C, primary NK cells are incubated at 37°C in R10 medium with or without 200 IU/ml human IL-2 cytokine for 1h, 24h, 48h or 72h (Fig. 5e - f, Supplementary Figure 6) in a tissue-culture treated 24-well plate (Corning).

Ex-vivo expansion of NK cells

NK cells were ex-vivo expanded (Figs. 6,7) from PBMCs as previously published[15,34,35]. PBMCs are isolated from healthy donors after oral and written informed consent (Department of Transfusion Medicine, University Hospital Erlangen, Germany; IRB approval number 147_13B and 22-314-Bp), In brief, PBMCs are cultured in the presence of irradiated K562-mbIL15-41BBL feeder cells (gift from Prof. D. Campana, Department of Pediatrics, University Hospital of Singapore) for 14 days in RPMI 1640 medium supplemented with 10% fetal bovine serum, 20 µg/ml gentamycin, 1% L-glutamine and 200 IU/ml human IL-2 cytokine.

Collagen gel preparation

The collagen solution is prepared at a temperature of 4°C from a 2:1 mixture of rat tail collagen (Collagen R, 2 mg/ml, Matrix Bioscience) and bovine skin collagen (Collagen G, 4 mg/ml, Matrix Bioscience). It is important that all tubes and pipette tips that come in contact with the collagen solution are also cooled to 4°C. We then add 10% (vol/vol) sodium bicarbonate (23 mg/ml) and 10% (vol/vol) 10× cRPMI (Gibco), and dilute the solution to a collagen concentration of 0.6 mg/ml, 1.2 mg/ml or 2.4 mg/ml with a dilution medium containing 1 volume part NaHCO3, 1 part 10 × cRPMI and 8 parts H2O adjusted to pH 9 using NaOH. Immediately

after dilution, the cells are added to the collagen solution, and the cell-collagen mixture is pipetted into 35 mm dishes (ThermoFisher Scientific, Nunc) for traction force measurements, or 6-24 well plates (Corning) for migration measurements. The sample is quickly transferred to the incubator to initiate polymerization. After polymerization at 37 °C, 5% CO2 and 95% RH for 1 hour, 1-3 ml cell culture medium or PBS (as described below) is added. Time-lapse imaging of embedded cells is either started directly after this 1h polymerization period (Fig. 5), or 24 h later (for primary immune cells except neutrophils) to ensure that the cells have adapted to the collagen gel and have recovered their full migratory potential.

Two different batches of collagen are used in this work. The rheology of the collagen gels varies between these batches (Supplementary Fig. 1 - 2). Batch 1 results in stiffer collagen gels (storage modulus 286 Pa) and is used for Figures 2, 4, 7a-b, 7g-h and Supplementary Figure 9, while batch 2 (shear modulus 84 Pa) is used for Figures 3, 5, 6, 7c-d and Supplementary Figure 6. Rheological measurements are conducted as previously described in Böhringer et al.[14] In brief, collagens hydrogels (85µl) are polymerized for 30 minutes at 37°C between the plates of a shear rheometer (Discovery HR-3, TA Instruments, Milford, 20mm cone-plate geometry). Subsequently, a frequency sweep (0.2-2Hz) is performed at a low amplitude of 1%, and the stress-strain relationship is measured (0-100% strain) with a constant strain rate of 1%/s. Pore diameters for both batches are 4.4 µm as determined from confocal reflection image stacks (volume of 160x160x200 µm with a voxel sizes of 0.314x0.314x0.642 µm) as described in Refs.[14,36,37]

Manipulation of cell-matrix adhesion

Cell adhesion on collagen fibers (Fig. 7a - d) is prevented either by adding pluronic surfactant (1 vol% in PBS, Sigma-Aldrich, cat. no. 9003-11-6), or by calcium ion chelation with 5 mM ethylenediaminetetraacetic acid (EDTA, Roth, cat. no. 8040.3) in PBS. As a control, PBS is used.

Inhibition of Rho-kinase on NK92 cells (Fig. 4) is performed with Y-27632 Rock-inhibitor (Sigma-Aldrich, cat. no. 129830-38-2). Y-27632 Rock-inhibitor is dissolved in DMSO (Sigma-Aldrich, cat. no. 67-68-5) at a stock concentration of 1mM. Y-27632 Rock-inhibitor is diluted in cell culture medium to a final concentration of 10 µM. Inhibition of myosin in NK92 cells (Fig. 4) is performed with blebbistatin (Sigma-Aldrich,cat. no. 856925-71-8). Blebbistatin is dissolved in DMSO at a stock concentration of 10 mM and mixed with cell culture medium to a final concentration of 3 µM. As control, DMSO is mixed in cell culture medium to a final concentration of 0.075 vol%.

Cells are mixed in collagen gel as described above. 2 ml of the PBS-drug solution (pluronic, EDTA, Y-27632, blebbistatin) or 2 ml of PBS with or without DMSO as control, are added one hour after initiating collagen polymerisation to the cell samples, and incubated for 1 h before measurements.

For integrin blocking or integrin activation with antibodies (Fig. 7g - h, Supplementary Fig. 8), we incubate 5-6*10⁵ NK92 cells prior to mixing them into the collagen for one hour in serum-free cell culture medium at 37°C with or without the following antibodies: 10 µg/ml anti-integrin activating β1 (CD29) antibody (Biolegend, cat. no. 303002) and the corresponding IgG1 isotype (Biolegend, cat. no. 400101) as negative control; anti-integrin blocking β1 antibody (P5D2, Abcam, cat. no. ab24693) or anti-integrin

Cell type	Cell number	Collagen volume	Collagen concentration	Type of well plate	Figure number
Non-activated primary immune cells	75.000	450 µl	0.3, 0.6, 1.2, 2.4 mg/ml	24	5c-d
IL-2 activated NK cells	150.000	750 µl	1.2 mg/ml	12	5e-f, and S4
Ex-vivo expanded NK cells	300.000	1-1.5 ml	1.2 mg/ml	6	7c-d
NK92 cells	300.000	1-1.5 ml	1.2 mg/ml	6	4d-j; 7a-b; 7g-h; 8; S6,

Table 1: Experimental set-up of the 3D migration assays with different cell types and different collagen concentrations and volumes due to the use of different tissue treated plates shown in several figures.

blocking β2 (CD18) antibody (Merck, cat. no. CBL158), and the corresponding IgG1 isotype (rndsystems, cat. no. MAB002). After incubation with the antibodies, we mix the NK92 cells in collagen gel as described above.

Manipulation of microtubules

Destabilization of microtubules in NK92 cells (Fig. 8) is performed with nocodazole (Biomol, Cay13857-10). Nocodazole is dissolved in DMSO at a stock concentration of 10 mM. Serum-free cell culture medium is either mixed with nocodazole to a final concentration of 10 μM or mixed with DMSO to a final concentration of 0.1% (DMSO control).

300.000 NK92 cells are incubated for 10 min with serum-free cell culture medium with or without nocodazole. Subsequently, the cells are washed with serum-free cell culture medium and mixed in collagen gel as described above. After one hour of polymerisation, 1ml of the serum-free cell culture medium solution with or without nocodazole is added to the gel. This results in two conditions: cells that are never or continuously exposed to nocodazole.

Carbomer hydrogel preparation

Carbomer hydrogel (Fig. 7a - d, Supplementary Figure 8) is prepared by mixing 9 mg carbomer powder (Ashland 980 carbomer, Covington, USA) with 1 ml R10 medium. The pH is titrated to a value of 7.4 with 10 M NaOH. The carbomer solution is incubated at 37°C and 5% CO2 for at least one hour for equilibration. The migration assay is started directly after mixing 300.000 cells in 2 ml of 9 mg/ml carbomer in each well of a tissue-culture treated 6-well plate (Corning).

High-throughput 3D migration assay

After gel preparation (Table 1) and polymerisation, the well plate is transferred to a motorized microscope (Leica DMi6000B, equipped with a 10x 0.25 NA Leica objective and an Infinity III CCD camera, Lumenera) that provides an incubation chamber (Tokai HIT model: WSKMX). For 3D time-lapse imaging, we perform z-scans (10 μm apart) through the 1 mm thick gel every 15 s for a duration of 2.5 min for primary immune cells or for a duration of 5 min for ex-vivo expanded NK cells and NK92 cells. For each condition, we repeat this scanning procedure for 10 field-of-views in sequence.

For analysis, we detect individual cells using a convolutional neural network that is trained on 4 - 16 manually labeled minimum/maximum intensity projections with approximately 100 cells in each projection. Additionally, we do data augmentation by image flipping, transposing, and cropping. The network is based on the U-Net architecture[38]. The labeling accuracy of the network ranges from 82% for monocytes to 96% for T cells, 86% for NK92 cells and 94% [15] for ex-vivo expanded NK cells (F1 score). We then connect the x,y positions of detected immune cells between subsequent images to obtain migration trajectories. An immune cell is classified as motile if it moves away from its starting point by ≥ 6.5 μm within 2.5 min for primary immune cells, 13 μm within 5 min for ex-vivo expanded NK cells and 6.5 μm within 5 min for NK92 cells. Cell speed is determined as the diagonal of the bounding box of each cell trajectory, divided by the measurement time of 2.5 min or 5 min. Directional persistence is determined as the average cosine of the turning angles between consecutive cell movements. Zero persistence corresponds to random motion, whereas a persistence of unity corresponds to ballistic motion.

Single cell 3-D traction force microscopy

3D traction force microscopy (Fig. 2b-e, 3, 4a-c, 5a,b, 5d-f, 6) is conducted as described in Böhringer et al.[14] and Condor et al.[39]. In brief, we pipette 3 ml collagen (0.6 - 2.4 mg/ml) solution with 200,000 cells into a 35 mm Nunc cell culture treated Petri dish (ThermoFisher Scientific). After incubation for one hour to ensure the complete polymerization of the gel, 2 ml of cell culture medium are added, and the first measurements start immediately. An additional waiting time of 24 hours before imaging ensures that non-activated primary immune cells have properly adapted to the collagen gel. We image z-stacks around single cells with a cubic volume of $(123\mu m)^3$ using confocal reflection microscopy (Leica DM 6000 CFS with HCX APO L 20x/1,00 W objective). A combination of galvanometric stage and resonance scanner allows for time-lapse imaging of 4 single cells simultaneously with a temporal resolution of 60 seconds. In each independent experiment, up to 12 individual cells were recorded for 24

minutes experiencing constant atmospheric conditions in a climate chamber (37°C, 5% CO2.). We select elongated cells for traction force measurements specifically, as these are assumed to be motile. From the recorded 3D reflection-channel image stacks, we extract the cell-induced deformations of the collagen fibers using 3D particle image velocimetry as implemented by the SAENOPY Python software package[14]. Based on these 3D deformation fields, SAENOPY computes the contractility of each cell at each measured time step. We exclude deformation fields for which the sum of absolute z-deformations exceeds 45% of the sum of the absolute deformations in all directions, as these deformations fields are perturbed by the acceleration of the galvo stage or by a wobbling motion of the collagen. We obtain the cell contractility and force polarity of individual cells from up to 4 independent experiments each.

The 3D traction force microscopy experiments shown in Fig. 2b,c, Fig. 3b, Fig. 4a,b, and Supplementary Figures 1 and 2 have previously been reported in the work by Böhringer et al.[14]

Bayesian filtering of cell trajectories

Bayesian filtering of the individual NK cell trajectories is employed as described in Mark et al.[20], but adapted to the fast-moving immune cells. In particular, we use the probabilistic programming framework *bayesloop* to infer the temporal evolution of the filtered cell speed and the filtered directional persistence from a series of (x_t, y_t) -coordinates that are obtained from the TFM experiments (we ignore the movement of the cells in z-direction as cell movement is more pronounced in the x/y-plane due to the free surface of the collagen gel). First, we model the measured cell speed v_t by a Poisson random variable with a time-varying rate parameter λ_t:

$$v_t \sim Poisson(\lambda_t) \quad \text{with} \quad v_t = \frac{\sqrt{\Delta x_t^2 + \Delta y_t^2}}{\Delta t}$$

where Δx_t and Δy_t denote the measured cell movement from time step $(t-1)$ to time step t, and Δt denotes the time interval between two time steps (1min in the TFM experiments). The rate parameter λ_t is allowed to vary slowly over time according to a Gaussian random walk, such that the difference of two subsequent rate parameter values follows a Normal distribution with zero mean and the standard deviation σ:

$$(\lambda_t - \lambda_{t-1}) \sim Normal(\mu = 0, \sigma)$$

The hyper-parameter σ thus describes how fast the rate λ_t changes over longer time scales. The rate parameter λ_t represents the filtered cell speed that we subsequently use to compute cross-correlation functions. The complete temporal evolution of the filtered cell speed λ_t as well as the hyper-parameter σ are then inferred from the series of measured cell positions using the grid-based approach detailed in Mark et al.[20] (Supplementary Figure 3).

Next, we model the measured turning angle ϕ_t (the change in direction between two cell movements) by a zero-centered von Mises distribution:

$$\phi_t \sim vonMises(\mu = 0, \kappa_t) \quad \text{with}$$
$$\phi_t = atan2(v_{x,t} - v_{x,(t-1)}, v_{y,t} - v_{y,(t-1)})$$

Here, $atan2$ denotes the multi-valued inverse tangent function (c.f. Mark et al.[20]), and κ_t denotes the time-varying concentration parameter of the von Mises distribution. We use κ_t as the filtered directional persistence, as a higher concentration of turning angles around zero describes a more persistent cell movement. For $\kappa = 0$, the model describes non-persistent Brownian random motion. For the asymptotic limit $\kappa \to \infty$, the model describes a ballistic movement. As for the filtered cell speed λ_t, the filtered directional persistence κ_t is allowed to vary slowly over time according to a Gaussian random walk:

$$(\kappa_t - \kappa_{t-1}) \sim Normal(\mu = 0, \sigma)$$

Again, the hyper-parameter σ describes how fast the directional persistence κ_t changes over longer time scales. The complete temporal evolution of the filtered cell speed κ_t as well as the hyper-parameter σ are then inferred from the series of measured cell positions using the grid-based approach detailed in Mark et al.[20] (Supplementary Figure 3).

Cross-correlation functions

To extract common patterns in the dynamic regulation of cell contractility c_t in relation to cell speed λ_t and directional persistence κ_t, we compute cross-correlation functions $\rho(\Delta t)$ across n cell trajectories by first creating

two vectors V and W that contain the values (for example of contractility) from all cells pooled together, but shifted in time by the lag-time Δt:

$$V(c, \Delta t) = (c_{t=0,k=0}, c_{t=1,k=0}, \ldots, c_{t=(T-\Delta t),k=0},$$
$$c_{t=0,k=1}, \ldots c_{t=(T-\Delta t),k=1}, \ldots c_{t=(T-\Delta t),k=n})$$

$$W(c, \Delta t) = (c_{t=\Delta t,k=0}, c_{t=\Delta t+1,k=0}, \ldots, c_{t=T,k=0},$$
$$c_{t=\Delta t,k=1}, \ldots c_{t=T,k=1}, \ldots c_{t=T,k=n})$$

where the index k denotes the cell, and the index t denoted the time step. Note that we further mask out all time steps for which we have obtained a negative contractility estimate to focus on the dynamics of contractile cellular forces. For positive lag-times $\Delta t \geq 0$, we then compute the cross-correlation function between contractility and cell speed at lag-time Δt (i.e. contractility following cell speed with a lag of Δt):

$$\rho(c, \Delta t) = SpearmanR(W(c, \Delta t), V(\lambda, \Delta t))$$

where *SpearmanR* denotes the Spearman's rank correlation coefficient. For negative lag-times $\Delta t < 0$, we adjust the indexing as follows:

$$\rho(c, \Delta t) = SpearmanR(W(\lambda, -\Delta t), V(c, -\Delta t))$$

Error bars for cross-correlation functions are determined via bootstrapping, by selecting n cell trajectories at random with replacement from the original set of n cell trajectories for 1,000 times, resulting in 1,000 bootstrapped correlation values. We subsequently compute the 1-sigma credible interval by taking the 31.73th percentile and the 68.27th percentile as the lower and the upper error bar boundary, respectively.

Code availability

The 3D traction force microscopy method used in this work is implemented in the Python package *SAENOPY*[14]. The software is open source (under the MIT License) and is hosted on GitHub (https://github.com/rgerum/saenopy). The Bayesian filtering of cell trajectories is implemented in the Python package *bayesloop*[20]. The software is open source (under the MIT License) and is hosted on GitHub (https://github.com/christophmark/bayesloop). The detection and tracking of immune cells in the high-throughput migration assays is implemented in custom Python scripts that are available from the corresponding author upon request. The figures in this study have been created using the Python packages matplotlib[40] and Pylustrator[41].

Data availability

Data supporting the findings of this study are available from the corresponding author upon reasonable request.

Supplementary Information

Supplementary Figure 1: Collagen Rheology and pore size for collagen batch 1. Rheology and pore size measurements of collagen hydrogels at different concentrations of 0.6 mg/ml (blue), 1.2 mg/ml (orange) and 2.4 mg/ml (green) for collagen batch 1 (see Batch C in Böhringer et al.[14]). **a:** Storage modulus scales for small frequency (mean value at 0.02 Hz) for different collagen concentration (0.6 mg/ml: n=3, 1.2 mg/ml: n=4, 2.4 mg/ml: n=4). Stiffness increases with increasing collagen concentration. **b:** Mean pore size for different collagen concentrations averaged from 8 different regions (80x80x100\,μm) within an image stack (160x160x200\,μm) per concentration (n=1). The pore size decreases with increasing collagen concentration. **c:** Stress-strain relationship of collagen gels for different collagen concentration. Solid lines represent the mean, shaded areas represent the standard deviation (0.6 mg/ml: n=3, 1.2 mg/ml: n=4, 2.4 mg/ml: n=4).

Supplementary Figure 2: Mechanoresponsiveness of NK92 cells. 1.2 mg/ml collagen hydrogels for two different batches of collagen batch 1 (gray) and 2 (blue) (see Batch A and C in Böhringer et al.[14]). **a:** Storage modulus scales for small frequency (mean value at 0.02 Hz) for batch 1 (n=4) and batch 2 (n=8). The stiffness of batch 1 is more than three times higher compared to batch 2. **b:** Mean pore size measured from an image stack of collagen 1.2mg/ml gels of batch 1 and batch 2. Pore size is averaged from 8 different regions (80x80x100μm) within an individual image stack (160x160x200μm). For both batches, the pore size is 4.4 μm. **c:** Contractility of NK92 cells in batch 1 (n=51) and batch 2 (n=50). The contractility of NK92 cells is twice as high in batch 1 as in batch 2.

a

Supplementary Figure 3: Bayesian filtering of cell speed and persistence. a: Exemplary migration path (black) of an NK92 cells embedded in a 1.2 mg/ml collagen gel, measured by confocal microscopy over a time course of 20 min, at an interval of 1 min. The cell trajectory starts at the green circle, and ends at the red circle. Scale bar: 10µm. **b:** The momentary cell speed (dashed black line), and the filtered cell speed (solid black line) as obtained from the Bayesian filtering method. The red shading indicates the probability density of the inferred, filtered cell speed. **c:** The cosine of the turning angle as a momentary measure of directional persistence (dashed black line), and the filtered directional persistence (solid black line) as obtained from the Bayesian filtering method. The blue shading indicates the probability density of the inferred, filtered persistence.

Supplementary Figure 4: Effect of cell viability on the motile fraction in high-throughput migration assays. To test whether the variations in cell viability due to cell culture conditions and various cell treatments systematically affect the measured motile fraction of cells in high-throughput migration assays, we repeat the migration assays shown in the main text and extend the measurement protocol by a live/dead staining using propidium iodide (Sigma-Aldrich, cat. no. P4864, incubation with 1:1000 dilution of 1 mg/ml for 20 min) at the end of the time-lapse imaging period. By overlaying the detected cell trajectories with the fluorescent images, we determine the number of non-motile dead cells in addition to the number of live motile, and live non-motile cells. This additional metric allows us to compute a corrected motile fraction that is defined as the number of motile cells divided by the number of live non-motile cells. This corrected motile fraction is always equal of greater than the "naïve" motile fraction that does not account for cell viability. **a:** Corrected motile fraction that accounts for cell viability plotted against the naïve motile fraction for n=19 individual experiments using NK92 cells. We do not find any systematic bias of the naïve motile fraction, indicating that variations in cell viability do not systematically affect the outcomes of the high-throughput migration assay with NK92 cells. **b:** Naïve motile fraction (blue) and corrected motile fraction (orange) for different cell treatments (n=1 for each condition; motile fraction of cells treated with Nocodazole is measured 8h after treatment). **c:** Relative change of the motile fraction between control conditions and treatment conditions for the different treatments shown in (b), using the naïve motile fraction (blue) and the corrected motile fraction (orange).

Supplementary Figure 5: Dynamic regulation of force generation and cell motility in primary immune cells. Left column shows the cross-correlation function between cell contractility and cell speed. Positive x-values indicate that speed follows contractility, negative x-values indicate that contractility follows cell speed. Right column shows the cross-correlation function between cell contractility and directional persistence. Error bars indicate 1 sem obtained by bootstrapping. In general, positive values on the right-hand side of the plots and negative values on the left-hand side of the plots indicate that the cells increase contractility after getting slower or less persistent, and before getting faster or more persistent. **a:** Combined cross-correlation functions computed from all primary cells (n=116). **b:** Cross-correlation functions for B cells (n=20). **c:** Cross-correlation functions for Monocytes (n=28). **d:** Cross-correlation functions for NK cells (n=16). **e:** Cross-correlation functions for T cells (n=16). **f:** Cross-correlation functions for B cells (n=36).

Supplementary Figure 6: Migration of primary NK cells after IL-2 activation measured in a high-throughput 3D migration assay. Motile fraction, migration speed and migration persistence of primary NK cells incubated with (orange) and without (blue, ctrl) IL-2 from n=2 independent donors for different incubation times (1h - 72h) in 1.2 mg/ml collagen. Each bar represents mean +- se from multiple fields of views (9-20) for each subject. **a:** Motile fraction decreases over time without the addition of IL-2, whereas the motile fraction remains stable over time with the addition of IL-2. **b:** Cell speed of the motile cells increases slightly over time with and without the addition of IL-2, but falls back to the approximate initial state after 72 hours. **c:** Directional persistence of motile cells remains the same over time, with or without the addition of IL-2.

Supplementary Figure 7: Size of NK92 cells vs. ex-vivo expanded NK cells Cell size distribution of motile and non-motile ex-vivo expanded NK cells (blue, n = 24.453) and NK92 cells (orange, n = 6.757) using masks generated by the neural network. Ex-vivo expanded NK cells have an average cell diameter of about 7.6 µm, while NK-92 cells have about 9.9 µm.

Supplementary Figure 8: Carbomer rheology and structure. **a:** Shear modulus of Carbomer as a function of angular frequency, measured with a rotational rheometer. **b:** Shear modulus of Carbomer as a function of the oscillation strain, showing the fluidization of Carbomer at high strains. **c:** Patches of low diffusivity (dark) and high diffusivity (light) in Carbomer. Diffusivity was quantified by recording the fluorescent intensity of 1µm-sized beads that are suspended in Carbomer and move randomly due to thermal diffusion (c.f. Supplementary Video 5). Black patches correspond to the 5th percentile of all intensity values that are cumulated over time, and indicate stiff inclusions within the Carbomer matrix. Scale bar is 10µm.

Supplementary Figure 9: Antibody-based manipulation of cell adhesion measured in a high-throughput 3D migration assay. Motile fraction, migration speed and migration persistence of NK92 cells incubated with blocking (blue or orange) or activating integrin antibody (AB, blue), with the corresponding IgG1 isotype as negative control (white) or without any antibody (gray, control) from n=3-4 independent samples in 1.2 mg/ml collagen. Each bar represents mean +- se from 10 fields of views for each subject. Each circle represents individual values of each sample. **a:** Motile fraction, **b:** cell speed and **c:** directional persistence of NK92 cells treated with β1 blocking AB (n=3) **d-f:** Same as in A - C, but for NK92 cells treated with β1 activating AB (n=3) **g-i:** Same as in A - C, but for NK92 cells treated with β2 blocking AB (n=4).

Immune cells employ integrin-mediated traction forces to overcome steric hindrance in 3D biopolymer networks

Supplementary Video 1: Confocal reflection movie of an ex-vivo expanded NK cell (cyan blue, calcein stain) migrating in a 1.2 mg/ml collagen gel (white, confocal reflection channel), imaged for 15 min at an interval of 15 sec. During the first 4 minutes, the NK cell migrates in an amoeboid way without visible deformations of the ECM. After 5 minutes, the cell changes its migration mode by pulling on the collagen fibers. Scale bar: 20µm.

Supplementary Video 2: Confocal reflection movie of two ex-vivo expanded NK cells (cyan blue, calcein stain) migrating in a 1.2 mg/ml collagen gel (white, confocal reflection channel), imaged for 15 min at an interval of 15 sec. The cell in the upper right quadrant of the field of view exerts strong pulling forces that result in large deformations of the collagen network. At t=13min, the cell succeeds in pulling its uropod out of the narrow pore and continue its migration path. Scale bar: 20µm.

Supplementary Video 3: Long-term time-lapse imaging of NK92 cells migrating in a 1 mm thick 1.2 mg/ml collagen gel for 24 h at an interval of 15 sec (1 sec of the video corresponds to 8 min measurement time). The movie shows the minimum intensity projection of recorded image stacks with a total height of 1 mm at a z-interval of 10 µm.

Supplementary Video 4: Confocal reflection movie of an NK92 cell (green, calcein staining) embedded in a 1.2mg/ml collagen fiber network (red, confocal reflection), imaged for 48 min at an interval of 1 min. Arrows indicate the deformations of the extracellular matrix. This video has been published before as part of Ref.[14]

Supplementary Video 5: Confocal images of 0.1 µm fluorescent beads (FluoSpheres carboxylate-modified microspheres 0.1 µm, orange (540/560), ThermoFisher, LOT: 2201623) mixed in 1.5 ml carbomer at an interval of 15 sec. Permanently dark regions in the video indicate stiff inclusions within the carbomer gel, as the beads are unable to diffuse into these regions (some beads become stuck at the boundary of these inclusions). Scale bar: 10 µm.