A core Nrf2 gene set defined through comprehensive transcriptomic analysis predicts drug resistance and poor multi-cancer prognosis

George Luo[1], Harshita Kumar[2], Kristin Aldridge[3], Stevie Rieger[4], Ethan Jiang[5], Ernest R. Chan[6], Ahmed Soliman[7], Haider Mahdi[8,9,10], and John J. Letterio[11,12,13]

Department of Pathology, Case Western Reserve University School of Medicine, Cleveland, Ohio
Department of Engineering, Case Western Reserve University, Cleveland, Ohio
Triterpenoid Therapeutics Inc., Tunbridge Vermont
Department of Biochemistry, Case Western Reserve University, Cleveland, Ohio
Booth School of Business, University of Chicago, Chicago, Illinois
Institute for Computational Biology, Case Western Reserve University, Cleveland, Ohio
Department of Pediatrics, SUNY Downstate Hospital, Brooklyn, New York
Magee Women's Research Institute, University of Pittsburgh Medical Center, Pittsburgh, PA
Magee Women's Hospital, University of Pittsburgh Medical Center, Pittsburgh, PA,
Hillman Cancer Center, University of Pittsburgh, Pittsburgh, PA,
The Angie Fowler Adolescent and Young Adult Cancer Institute, University Hospitals Rainbow Babies & Children's Hospital, Cleveland, Ohio
The Case Comprehensive Cancer Center, Cleveland, Ohio
Department of Pediatrics, Case Western Reserve University, Cleveland, Ohio

Corresponding Authors:
George Luo, Case Western Reserve University School of Medicine, 2103 Cornell Rd., Wolstein Research Bldg. Rm 3501, Cleveland, Ohio 44106; gxl263@case.edu

John J. Letterio, M.D., The Angie Fowler Adolescent and Young Adult Cancer Institute, University Hospitals Rainbow Babies & Children's Hospital, 11100 Euclid Avenue, Cleveland, Ohio 44106; jjl26@case.edu

Running Title: 'Nrf2 activation leads to selective drug resistance and poor cancer prognosis'

Key Words: Nrf2, Drug Resistance, Cancer Prognosis, Transcriptomics,

ABSTRACT

The Nrf2-KEAP1 pathway plays an important role in the cellular response to oxidative stress and confers protection in diseases associated with chronic inflammation. However, chronic activation of the Nrf2 pathway may contribute to metabolic changes and disease progression in cancer. We investigated the activation of Nrf2 in human cancers and fibroblast cells through KEAP1 inhibition and cancer associated KEAP1/Nrf2 mutations. We define a core set of 14 upregulated Nrf2 target genes from seven RNA-Sequencing databases that we generated and analyzed. Additionally, we validated this universal Nrf2 target gene set through analyses of published databases. An Nrf2 activity score based on expression of these core target genes correlates with resistance to drugs such as PX-12 and necrosulfonamide but not to paclitaxel or bardoxolone methyl. We validated these findings and found Nrf2 activation also led to radioresistance in cancer cell lines. Finally, our Nrf2 score is prognostic for survival for a variety of cancers, an observation validated in additional independent cohorts. These analyses define a core Nrf2 gene set that is robust, versatile, and useful for predicting drug resistance and cancer prognosis.

SIGNIFICANCE: The frequent activation of Nrf2 observed in cancer cells confers protection from oxidative stress, cytotoxic chemotherapeutic agents and radiation. We define a 'Nrf2 gene signature' that is comprised of 14 target genes and faithfully predicts increased Nrf2 activity, selective drug and radiation resistance in cancer cell lines, and a worse clinical prognosis in a variety of human cancers.

INTRODUCTION

Nuclear factor-erythroid 2-related factor 2 (*NFE2L2*, Nrf2) is a transcription factor that plays a key role in the regulation of cellular defense mechanisms against oxidative stress and other forms of cellular damage (1). Nrf2 is responsible for the activation of a wide range of antioxidant and detoxifying enzymes, which help to protect cells from damage caused by reactive oxygen species (ROS) and other toxic molecules (1). Kelch-like ECH-associated protein 1 (KEAP1) is an adaptor protein that acts as a negative regulator of Nrf2 by targeting it for ubiquitination and degradation (1, 2). In normal cells, KEAP1 acts to prevent Nrf2 from becoming overactive and inducing excessive oxidative stress (1, 2). However, in cancer cells and other disease states, the Nrf2-KEAP1 pathway can become dysregulated, leading to the activation of Nrf2 and the increased expression of its Nrf2 target genes (2, 3). The consequences of constitutive Nrf2 activation in cancer include increased chemoresistance and a poor overall prognosis (2-5).

The Nrf2-KEAP1 pathway is therefore a crucial component of cellular defense mechanisms, and its dysregulation can play a major role in the development of diseases such as cancer, neurodegeneration, and inflammation (5-8). In recent years, the Nrf2-KEAP1 pathway has become a target for the development of new drugs for the treatment of these and other diseases (9, 10). Thus, a deeper understanding of the Nrf2-KEAP1 pathway and its role in cellular biology is of significant interest to both basic science and clinical researchers. However, despite the interest and attention of this pathway, there are few reliable biomarkers that can be used to measure Nrf2 activity for the wide range of context in which it is being researched (11, 12).

Here we first address this issue by defining a core set of Nrf2 target genes that

faithfully predicts Nrf2 activation in a wide array of contexts through analyses of a variety

of cancer cell types and normal cells, as well as both pharmacological and genetic

activation of Nrf2. Generating these databases allowed us to find the most universally

upregulated target genes of Nrf2 while also enabling us to compare differences between

the activation of Nrf2 in these various contexts. We define a list of 14 core upregulated

target genes of Nrf2 which are present in every database that we generated, which we

further validate in other published databases. This analysis allowed us to use this gene

set as a biomarker and representation of Nrf2 activation.

Using these core target genes, we generate a Nrf2 change score which is useful

for identifying both which and to what degree specific genetic modifications and drugs

activate Nrf2. A similar but distinct Nrf2 activity score can predict which drugs are affected

by Nrf2 activation and which cell lines are most resistant to these drugs. We validate the

modeling of selective drug resistance in our own cell lines. Finally, we investigated

whether this Nrf2 activity score is associated with prognosis in a variety of cancers. We

trained the Nrf2 activity score in 22 The Cancer Genome Atlas (TCGA) cancers (13), and

found that it passed training in twelve cancers. We were then able to test validation in

eight of these twelve cancers with independent cohorts and demonstrated that four

cancers have significantly poorer prognosis associated with Nrf2 activity, as defined by

our Nrf2 activity score. Overall, we identify a Nrf2 signature that can be used in a wide

range of contexts including predicting Nrf2-associated drug resistance and as a biomarker

of a poor cancer prognosis.

RESULTS

Generation of Seven Transcriptomic Databases of Nrf2 Activation

In order to determine the core target genes of Nrf2, we investigated gene expression changes from both genetic and pharmacological targeting of the Nrf2/KEAP1 pathway (Fig. 1A). We treated myeloma (RPMI-8226), ovarian (OVCAR-8), glioma (SF8286), and primary dermal fibroblast cells with an Nrf2 activator (14), the synthetic oleanane triterpenoid CDDO-2P-Im (1-[2-Cyano-3,12-dioxooleana-1,9(11)-dien-28-oyl]-4(-pyridin-2-yl)-1H-imidazole), for six hours before extracting RNA for RNA-sequencing. We then confirmed activation of Nrf2 through upregulation of *NQO1*, a well-established target of Nrf2 (1, 2), in each of these cells (Fig. 1B-E). We also used CRISPR-Cas9 to knock out *KEAP1* in ARH-77, a B-cell lymphoblastic cell line, and verified the *KEAP1* deletion at the DNA, RNA, and protein levels (Fig. 1F-H). Furthermore, NQO1 was upregulated at baseline compared to the wild-type (WT) ARH-77 cells (Fig. 1I and J).

We also leveraged the available public databases of the Cancer Cell Line Encyclopedia (CCLE) (15) and TCGA tumor samples (13) to explore activation of Nrf2. Over 20% of non-small lung cancer cells (NSCLC) had mutations in *KEAP1* or *NFE2L2* leading us to analyze NSCLC cell lines and tumor samples (Supplemental Fig. 1A). After separating TCGA lung adenocarcinoma (LUAD) into WT and KEAP1/Nrf2 mutated samples, we investigated the top differentially expressed genes (DEGs) of which *NQO1* was among the top genes identified (Fig. 1K). We repeated the analysis for NSCLC cell lines found in the CCLE on DepMap.com (16) and verified *NQO1* was upregulated in the KEAP1/Nrf2 mutated group (Fig. 1L). Furthermore, we confirmed ROS pathway activation, a key Nrf2 related pathway, through gene set enrichment analysis (GSEA) for

our RNA-Sequencing generated databases (Supplemental Fig. 1). These databases all

demonstrate Nrf2 activation in different cellular and molecular contexts which enables us

to develop a more accurate and comprehensive perspective of Nrf2 activation.

A Core Nrf2 Transcriptional Signature is Present after Nrf2 Activation

After we created different databases of Nrf2 activation, we investigated the most

frequently upregulated targets of Nrf2 activation, a list of 15 candidate core Nrf2 genes

(Fig. 2A). Additionally, many other genes were upregulated in most scenarios of Nrf2

activation (Fig. 2B). Many of these core Nrf2 genes such as *NQO1*, *GCLC*, *GCLM*, and

SRXN1 have been well-documented as Nrf2 target genes in the literature (Supplemental

Fig. 2). Interestingly, we noticed most of these core genes, such as *ABHD4* and *SRXN1,*

showed similar upregulation across all cell types and methods of Nrf2 activation (Fig. 2C,

2D, and Supplemental Fig. 3). However, *AKR1C3* and other members of the *AKR1C*

family displayed considerably higher activation in genetic models of Nrf2 activation

compared to short term drug-induced activation (Fig. 2E, Supplemental Fig. 4, and

Supplemental Table 1). Another interesting observation is that higher concentrations of

drug does not equal higher activation of Nrf2 target genes. This appears to be somewhat

cell line dependent and gene specific as other core Nrf2 genes show elevated

upregulation at high doses (Supplemental Fig. 4A and B). We also used an *in silico* model

of motif binding, LASAGNA-Search 2.0, to identify Nrf2-ARE (Antioxidant Response

Element) sequences near the promoters of these core and nearly universal Nrf2 genes

(Fig. 2F and Supplemental Fig. 5A). This analysis suggests that Nrf2 directly binds to

promoters of these core genes and directly upregulated transcription, supporting their

positions within an initial list of core Nrf2 target genes.

Defining Conditionally Upregulated and Novels Target Genes of Nrf2 Activation

Besides finding the most frequently upregulated genes, we examined if certain genes were specifically activated in one condition or another (Fig. 3A). We found certain genes such as *HMOX1* and *AKIRIN2* which are primarily upregulated in drug induced Nrf2 activation but not present in either the TCGA LUAD or NSCLC cell lines (Fig. 3B and C), consistent with the literature (17, 18). On the other hand, genes related to enzymes in the pentose phosphate pathway (*G6PD*, *TKT*, *TALDO1*, and *PGD*) were consistently upregulated in the genetically activated Nrf2 models but infrequently in the drug-induced models (Fig. 3D-G). However, this may be due to the relatively short duration of drug exposure. The pentose phosphate pathway is of particular importance as it is thought to help protect cells from oxidative stress and apoptosis (19).

We also uncovered two putative novel Nrf2 target genes that are upregulated in most of our models, *B4GALNT1* and *POPDC3* (Fig. 3H-I). We found a putative Nrf2/ARE motif near the promoter of B4GALNT1 but not POPDC3 (Supplemental Fig. 5B). Finally, we also found novel downregulated genes (*PLAT, CDC42EP1, EVA1A)* in our models of Nrf2 activation but these genes were not always downregulated or expressed (Fig. 3J-L and Supplemental Fig. 5B). Nrf2 has been reported to interfere with transcriptional regulation of proinflammatory cytokines such as IL-6 and IL-1B (20), but we did not investigate immune cells in these experiments. Nevertheless, we observed decreased expression of immune related genes such as *CXCL6* and *CXCL1* in TCGA LUAD and KEAP1 knockout ARH-77 cells supporting the anti-inflammatory effects of Nrf2 activation (Supplemental Fig. 6). Because synthetic triterpenoids such as CDDO-2P-Im also inhibit

NF-κB (21), it is difficult to comment on the effects of drug-induced Nrf2 activation on

inflammation, although suppression of inflammatory markers is observed.

Validation of the Core Nrf2 Gene Set in Other Published Datasets

Although we believe our core Nrf2 gene set can faithfully represent Nrf2 activity,

we needed to validate its utility in other contexts of Nrf2 activation. For this, we first tested

other TCGA cancers which had considerable Nrf2/KEAP1 mutations. We found lung

squamous cell carcinoma (LUSC), head and neck squamous cell carcinoma (HNSC) and

liver hepatocellular carcinoma (LIHC) to have significant numbers of mutations in

Nrf2/KEAP1. We found that both LUSC and HNSC validated all our core and nearly

universal Nrf2 gene set (Fig. 4A-B and Supplemental Fig. 7A-B). However, LIHC only

validated 13/15 core Nrf2 genes and 10/16 of the nearly universal Nrf2 gene set (Fig. 4C

and Supplemental Fig. 7C), though LIHC also had the lower number of Nrf2/KEAP1

mutations.

We selected two published studies (22, 23) to investigate the validity our Nrf2 gene

set in other databases. In three different prostate cell lines treated with sulforaphane by

Beaver *et al.*, we found upregulation of at least 14/15 of the core Nrf2 gene sets (Fig. 4D).

CBR3 was the only Nrf2 gene in our initial core set that did not pass multiple validation

sets so we chose to omit *CBR3* from the final core Nrf2 biomarker gene set. Overall,

these two different validations strongly demonstrate the validity and versatility of our 14

core Nrf2 gene set in human tissues.

In the rat model of Nrf2 gene deletion by Taguki *et al.*, two different germline Nrf2

KO mutations were generated and used to investigate Nrf2 dependent gene expression

changes (23). Mice were divided into the following groups and treated with vehicle or Nrf2 activator, CDDO-Im: WT + vehicle, WT + CDDO-IM, Nrf2 KO + vehicle, Nrf2 KO + CDDO-Im. We found that the majority of core Nrf2 target genes were upregulated by CDDO-Im treatment but unchanged in the setting of Nrf2 gene deletion (Fig. 4E and Supplemental Fig. 7D). However, other genes (*OSGIN1, SLC7A11, ME1, FTH1, FTL*) showed a similar trend of activation following CDDO-Im treatment but did not reach significance in every case. Finally, we observed that *AKR1C3* did not display any trend suggesting Nrf2 dependency which is similar to literature reports that AKR1C genes are not homologous between human and mice (24). We also performed analysis with the nearly universal Nrf2 gene set, finding most exhibited a trend suggesting Nrf2 dependency (Supplemental Fig. 7E and F). Therefore, we believe our core Nrf2 gene can also work for rodent models, with a few adjustments for genes known to be nonhomologous.

We introduce the Nrf2 Change Score (N2CS), which is a useful measure of drug or genetic effects on Nrf2 activity. Using gene expression changes of the 14 core Nrf2 genes, we can approximate the degree to which Nrf2 is changed with scores above 1 suggesting high activation with a score of 0 as the baseline (See Methods for calculations). We demonstrate the utility of the N2CS in the various conditions in which we examined the response to CDDO-2P-Im in our study, in the KEAP1 knockout, in the response to sulforaphane from validation studies, as well as in the response to other anti-cancer agents from published datasets (Fig. 4F and G). Interestingly, we also noticed that higher concentrations of CDDO-2P-Im do not further activate Nrf2 which is also the case with other Nrf2 activators (25). We found that both bortezomib and doxorubicin induce

218 Nrf2 activation (26, 27), but paclitaxel does not significantly affect Nrf2 activity. Thus, the

219 N2CS can be used to screen RNA-Sequencing data for modulators of Nrf2.

220 Nrf2 Activity Score Correlates with Resistance to Radiation and Certain Drugs

Using the final core Nrf2 target genes, we can also generate Nrf2 activity scores

(N2AS) to represent the degree of Nrf2 activation for individual cell lines and tumor

samples. The N2AS is useful for comparing cells or samples of the same group for

differences in Nrf2 activation. In the following experiments, we demonstrate the utility of

the N2AS.

The N2AS can be used to find whether specific non-conservative mutations in

Nrf2/KEAP1 affect Nrf2 activity. We first used N2AS to rank all CCLE NSCLC cell lines in

DepMap and investigated how mutation status of Nrf2/KEAP1 was related to N2AS (Fig.

5A). While most mutations are associated with higher N2AS, we do observe several

mutations are associated with medium or low N2AS, suggesting these mutations do not

activate Nrf2. Furthermore, we correlated N2AS with a functional output, sensitivity of

oxidative stress-mediated drugs from the DepMap database (106). We found N2AS

significantly correlated positively with resistance to PX-12 and necrosulfonamide in

NSCLC cells (Fig. 5B and C). Not surprisingly, PX-12 has been previously demonstrated

to induce apoptosis in an oxidative stress dependent manner (28) while necrosulfonamide

has been linked to reacting with cystines previously (29). These two drugs were among

the top drug correlations with N2AS and we further validated this observation in our

KEAP1 knockout cell line (Fig. 5D and E, Supplemental Table 2). We also tested

correlation with individual Nrf2 core genes and our gene set had a higher correlation than

any single gene and as high correlation as other published Nrf2 gene sets (Supplemental

Fig. 8A-B). Although radiation is not in the database, we also tested if Nrf2 activation

would lead to resistance against radiation (Fig. 5F). Our findings confirmed

radioresistance associated with Nrf2 activation, and other studies report similar findings

(30). Interestingly, we also observe enhanced colony formation with KEAP1 knockout in

ARH-77 cells (Supplemental Fig. 8C-D), a cell line with poor colony formation efficiency

(31).

We also investigated drugs that were not correlated with the N2AS (Supplemental

Table 2). We found both paclitaxel and bardoxolone methyl (CDDO-Me), a synthetic

oleanane triterpenoid, to have little correlation with N2AS (Fig 5G and H). We then tested

both drugs in our cell lines and found the resistance of KEAP1 knockout cells to paclitaxel

and bardoxolone methyl was absent in comparison to resistance to both PX-12 and

necrosulfonamide (Fig 5I and J). This suggests that our modeling of Nrf2 activity and drug

resistance is appropriate and can predict relative resistance mediated by Nrf2 for drugs

in the DepMap CTRP2 database.

Nrf2 Activity is Associated with Poor Prognosis in a Variety of Cancers

Although there have been many studies linking mutations causing Nrf2 activations

to chemoresistance and poor clinical outcomes, many of these previous studies were

concentrated on advanced stage patients with small sample sizes (32, 33). Recently, a

large study found KEAP1 mutations were associated with a worse prognosis, confirming

the clinical importance of Nrf2 activation in NSCLC (34). However, despite this finding,

mutation stratification alone is not significantly associated with prognosis difference in

TCGA LUAD or LUSC datasets (Supplemental Fig. 9A and B). We believe this

observation can be explained by our analyses of cell lines and truly demonstrates that not

all mutations of Nrf2/KEAP1 are equal and should be evaluated individually.

By using the N2AS to help categorize patients, we investigated whether our Nrf2

activity score had prognostic significance in 22 different TCGA cancers (Fig. 6A). During

the training, we focused on finding an appropriate N2AS cutoff for each cancer and a

highly significant p value ($p < 0.01$). We found 12 cancers passed training with 10 having

a poor prognosis associated with higher Nrf2 activity score while Uterine Corpus

Endometrial (UCEC) and Ovarian cancers had favorable prognosis associated with

higher Nrf2 activity (Fig. 6B). Under further examination, PTEN mutation is a common

feature in UCEC and is associated with both Nrf2 activation and a favorable prognosis

(Supplemental Fig. 10C) (34,35). PTEN inactivation can directly lead to higher Nrf2

activation and thus may be simultaneously driving both Nrf2 activity and a favorable

prognosis.

To help analyze the categorization and significance of the N2AS in training and

validation, we developed LOCC (Luo's Optimization Categorization Curve) which

visualizes all hazard ratios (HR) and p-values of all possible cutoffs (35). This method

allows us to understand the selection of the cutoff while also understanding if there are

other possible windows for categorization. Using this method on LUAD, we first locate

cutoffs that are highly significant, where the yellow line, representing the –log (p value),

is above the red line (representing a p value of 0.01) (Fig. 6C). At this cutoff, the black

line reveals the HR (the red line simultaneously represents an HR of 1). The x-axes of

both graphs are lined up such that the sample distribution of N2AS can be visualized. The

highest yellow peak is the most significant p value and is in most cases, the ideal cutoff

point. After choosing a N2AS cutoff of 0.463 for LUAD, the N2AS is finalized for the

training and a Kaplan-Meier Curve can be generated with appropriate additional

calculations (HR – 1.63, p < 0.01).

Using published East Asian cohort of LUAD (OncoSG) (36), we applied the

previous N2AS cutoff to test if it can stratify patients into prognostic significant groups

(Fig. 6D). Indeed, we found higher Nrf2 activity was associated with a worse prognosis

(HR – 2.60, p – 0.01). Using LOCC (35), we observed the cutoff was highly significant

and that the cutoff was appropriate despite using a very different patient population as

well as a drastically lower percentage of patients with Nrf2/KEAP1 mutation

(Supplemental Fig. 9E). These results demonstrate that N2AS could be used as a

prognostic biomarker for LUAD in various cohorts even without the presence of Nrf2

mutations.

Following the validation for LUAD, we used other public databases to validate

N2AS prognostic values for other cancers. Through the same methods of optimizing

cutoffs, we were able to successfully validate N2AS prognostic value in renal clear cell

carcinoma (KIRC, Fig. 6E and F), liver hepatocellular carcinoma (LIHC, Fig. 6G and H),

and acute myeloid leukemia (LAML, Fig. 6I and J). For all three cancers, we found high

Nrf2 activity was associated with lower overall survival (KIRC HR – 2.12, p < 0.001, LIHC

– 2.16, p < 0.001, LAML – HR 3.87, p - 0.001). Previous studies in small cohorts have

found high Nrf2 activity is associated with poor prognosis in liver and renal cancers but

have not been validated in larger cohorts (37, 38).

We also tested validation in head and neck squamous cell carcinoma (HNSC),

pancreatic adenocarcinoma (PAAD), melanoma (SKCM), and bladder carcinoma

(BLCA). Although these cancers passed training, they showed no significant poor

prognosis correlated with N2AS in the validation cohort (Supplemental Fig. 11). For all

other TCGA cancers examined, they either did not pass training or we did not find a large

suitable cohort to validate the training (Supplemental Fig. 12). Finally, we used other Nrf2

gene sets to compare if we would obtain similar prognostic results. We found that other

Nrf2 gene sets could obtain similar results in LUAD but not LAML (Supplemental Fig. 13),

showing that our core Nrf2 gene set is better suited toward a broader range of applications

than those selected gene markers of Nrf2 activity found in the existing literature.

Discussion

The KEAP1/Nrf2 pathway plays essential but complex roles in the maintenance of

metabolic homeostasis, regulation of the cellular redox state and in the modulation of the

inflammatory response (1, 6, 7). These activities contribute to neuroprotection in the

context of neurodegenerative disease (39), and to the suppression of other chronic

diseases affecting multiple organs including the liver and lung (40, 41). However,

increasing evidence linking constitutive Nrf2 activity to therapy resistance in cancer

highlight the challenges that have influenced the clinical development of activators of this

pathway (12). In our study we define a core Nrf2 target gene set as well as several

databases of Nrf2 activations that will aid preclinical efforts designed to clarify the context-

dependent roles of Nrf2 in health and disease and to define appropriate clinical indications

for therapeutics targeting the KEAP1/Nrf2 complex. We rigorously applied our core Nrf2

gene set through Nrf2 activation validation studies, drug resistance modeling with further

validation, and finally cancer prognosis with validation in multiple patient cohorts. During

this process, we also utilized other published gene sets for comparison and found that

while some could perform in certain situations, none were as useful for clinical prognosis.

This discrepancy is likely because of the high noise from conditional or non-Nrf2 target

genes. While our core Nrf2 gene set may not be perfect for all applications, it is a solid

foundation as a biomarker that can be used in a variety of settings for the assessment of

Nrf2 activity.

While cancer research has become increasingly focused on defining the unique

genetic alterations that characterize distinct and often rare tumor histologies, we believe

it is imperative to understand the commonality between groups as the latter may be

informative regarding therapeutic resistance. Although Nrf2 has been studied in many

cancer contexts, particularly in NSCLC and HNSC, we observe the same principal target

genes are upregulated in nearly all human cancer cells and that drug resistance modeled

in NSCLC can be validated in a B cell model. Furthermore, our investigation of cancer

prognosis demonstrates that in our successful training and validation, the trend is that

high Nrf2 activity is associated with a poor prognosis. As such, our understanding of Nrf2

targets and drug resistance may not be limited to the cancers we validated but to many

others which may have a larger impact on our understanding of therapeutic efficacy and

resistance. Furthermore, the literature of chemoresistance mediated by Nrf2 not only

directly applies to cancer resistance but also to prevention of toxicities by these agents

(42). Therefore, a broad application of precision medicine and Nrf2 targeting can be

investigated in the future.

The literature has described many instances of chemoresistance by Nrf2 (43), but

we demonstrate that not all chemoresistance by Nrf2 activation is comparable.

Furthermore, strategies designed to repress, knock down, or completely delete Nrf2 gene

expression may have very different effects on drug resistance when compared to

activation of Nrf2. Moreover, if Nrf2 acts to protect both normal and transformed cells, it

will be imperative to design strategies that obviate the effect of Nrf2 specifically in tumor

cells, as global repression of Nrf2 will most certainly increase side effects and toxicities

associated with chemotherapy and. radiation therapy.   Interestingly, the pentose

phosphate pathway activated by constitutive Nrf2 activation may be a targetable option.

Multiple studies have demonstrated CB-839 (Telaglenastat), a glutaminase inhibitor, can

inhibit cancer cells with high Nrf2 activity better than those without high Nrf2 activity (43-

46). While this may represent an exploitable 'flaw in the protective armor' that Nrf2

provides cancer cells, in the absence of a clear Nrf2 biomarker, this may not have a large

clinical benefit. Thus, it is not shocking that CB-839 recently failed a clinical trial in renal

cell carcinoma, a cancer in which Nrf2 activation may play a significant role in prognosis,

given the lack of a clear biomarker to select patients for whom the drug would have

potential clinical benefit (47, 48).

There are still many important questions to address before an Nrf2 activity score

can be used as an indicator for therapy. Many anti-cancer drugs can activate Nrf2 through

ROS generation, but this acute Nrf2 activation is clearly different from the constitutive

Nrf2 activation observed with genetic mutations. We showed a short-term exposure to

Nrf2 activators will upregulate many core Nrf2 targets similarly but other genes such as

AKR1C, *G6PD*, *PGD*, *TKT*, *TALDO1* exhibit much stronger upregulation in the genetic

models. Furthermore, synthetic oleaneane triterpenoids (SOTs), which activate Nrf2 but

also have anti-cancer properties (49), synergize with several anti-cancer agents (50). The

latter likely reflects the fact that these SOTs are multi-targeted agents, modulating the

activity of NFκB, STAT3, mTOR and other key regulators of cancer cell growth and

viability (49). Given the likelihood that individual Nrf2 target genes may exhibit time-

dependent and bi-modal regulation, it is conceivable that therapy might be tailored such

that different drugs regimens that exploit low Nrf2 activity with oxidative stress-mediated

therapy (drugs that have resistance correlated with N2AS) may be followed by

therapeutics that are less affected by N2AS.

Computational biology has been a field of great promise and interest but also

presents many questions and problems as well (51). Despite knowing the mutations of

cancer cells in the CCLE and tumor samples from TCGA, we can find the general

differences, yet it is hard to estimate the effect of specific unique mutations. Here we

demonstrate that a Nrf2 gene expression signature can help differentiate Nrf2 activity in

both cell lines and patient samples. However, instead of traditional methods of selecting

cutoffs such as median or quartiles, we chose a training method to find the optimal cutoff

and employed validation methodology which has been implemented by other studies (52,

53). Through this novel approach, a visualization of the cutoff selection process was

implemented to facilitate analysis and an enhanced understanding of the entire picture of

prognosis predicted by this gene set (LOCC) (35). Previous studies have used the

receiver operating characterization (ROC) curve to help continuous variable cutoffs

validation despite its better applicability to diagnosis as opposed to prognosis (53).

However, the ROC does not help us understand why biomarkers fail or succeed in

validation and is suited for large differences in odds or hazard ratio (54). With LOCC (35),

there is an abundance of information which can be seen and then calculated for

comparison across different genes or gene sets. When validation is unsuccessful or partly

successful, the reason can be seen: e.g., no peak throughout, correctly identified peak

but not enough significance or incorrect cutoff leading to low significance. We believe

machine learning is the future of computational biology, but we can also supervise and

understand the process which will lead to less misinterpretations and more awareness of

the validations and failures.

Our N2AS is a simple unweighted average of the z-scores of the core Nrf2 target

genes, which is both foundational and imperfect. We tried to be as comprehensive and

unbiased in our searching for core Nrf2 target genes and modeling of N2AS activity to

find results. Even though we did not optimize the N2AS for any application, we were still

able to find and validate significant results from drug resistance to cancer prognosis,

suggesting that this pathway is critical for these outcomes. With a clear direction, we can

optimize the analyses for each cancer with slightly different Nrf2 target genes and used

weighted scoring knowing that our fundamental goal is still the quantitation of Nrf2

activation, which is validated. If we optimize using machine learning without knowing

whether Nrf2 activation is important, we may find significant differences, yet it may not be

directly related to Nrf2 activity, as many genes exhibit complicated transcriptional

regulation orchestrated by distinct activators and repressors acting in concert. Therefore,

our simplistic N2AS is important but we understand that future research should

investigate appropriate optimization with regards to each cancer's context including

population differences and multivariate analysis.

Finally, we ponder why a universal pathway that directly affects chemoresistance

such as Nrf2/KEAP1 is prognostic in only some cancers and not others. Even though our

results are novel in finding Nrf2 is prognostic for large cohorts of AML, LIHC, KIRC, and

possibly other cancers, there are still many questions as to why we observed this effect

in these cancers specifically. For most of the cancers which were either validated or

promising candidates, we found a skewed distribution in which Nrf2 activity was very high

in certain patient samples. This also correlated with high numbers of mutations in those

cancers related to Nrf2/KEAP1. Yet, for LUSC, we do not observe any prognostic

significance, similar to what has been shown in other studies with Nrf2 in TCGA LUSC

(55). However, most cancers do not have a skewed distribution likely owing to fewer

mutations that dramatically activate Nrf2. Furthermore, while not all cancers rely on

oxidative stress-dependent mechanisms, if therapy is dependent on oxidative stress or

impaired by Nrf2 activity, then surviving cancer cells would likely have a selection

pressure favoring Nrf2 activation. However, such an escape mechanism may not be

found using bulk RNA-seq and even be difficult to discern with single cell RNA-seq. It

may be discovered through sequential RNA-Sequencing of initial and relapsed tumor

samples in therapies that significantly favor Nrf2 activation, a phenomenon which has

been demonstrated in breast cancer (56). As such, there is still a need for research to be

done in many cancers regarding the potential role of Nrf2 in affecting outcomes.

Methods

Cell lines

SF8628 (RRID:CVCL_IT46), OVCAR-8 (RRID:CVCL_1629), RPMI-8226 (RRID:CVCL_0014), ARH-77(RRID:CVCL_1072), and primary dermal fibroblast cells were all purchased from American Type Culture Collection (ATCC). Cells were tested negative for mycoplasma with ABM Mycoplasma PCR detection kit (catalog # G238) in February 2023. Cancer cell lines were authenticated using STR profiling at the CWRU genomic facility.

Cell Treatment for RNA

Cells were treated either with a control of media plus DMSO (1%) or with CDDO-2P-Im (gift from Dr. Michael Sporn, 96.3% purity by HPLC) for 6 hours in 6 well plates before RNA extraction at concentrations indicated in Supplemental table 3. RNA was measured by Nanodrop (Thermo Fisher Scientific) and then sent to a sequencing service, LC Sciences, for RNA-Sequencing or prepared for cDNA synthesis (#4368813, Thermo Fisher Scientific).

RNA-Sequencing

A poly(A) RNA sequencing library was prepared following Illumina's TruSeq-stranded-mRNA sample preparation protocol as previously described (57). RNA integrity was checked with Agilent Technologies 2100 Bioanalyzer. Poly(A) tail-containing mRNAs were purified using oligo-(dT) magnetic beads with two rounds of purification. After purification, poly(A) RNA was fragmented using divalent cation buffer in elevated

temperature. Quality control analysis and quantification of the sequencing library were performed using Agilent Technologies 2100 Bioanalyzer High Sensitivity DNA Chip. Paired-ended sequencing was performed on Illumina's NovaSeq 6000 sequencing system.

Cutadapt was used to remove the reads that contained contamination, low quality and undetermined bases (58). Then sequence quality was verified with FastQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/). HISAT2 was used to map reads to the genome of ftp://ftp.ensembl.org/pub/release-96/fasta/homo_sapiens/dna/ (59). The mapped reads were assembled using StringTie (60). The transcriptomes were merged to reconstruct a comprehensive transcriptome using perl scripts and gffcompare. After the final transcriptome was generated, StringTie and edgeR were used to estimate the expression levels of all transcripts (60, 61). Additional RNA-Sequencing analysis was performed using GSEA (Broad Institute). RNA-Sequencing has been deposited to GEO (GSE228434, GSE221526, GSE229914, GSE229704, GSE229652).

Immunoblotting

Cells were lysed with cell lysis solution (#9803, Cell Signaling, MA, USA) with protease and phosphatase inhibitor (A32961, Thermo Fisher Scientific) according to manufacturer's instruction. After lysis, cells were measured using bicinchoninic acid assay to determine protein concentrations and equal amounts of proteins were mixed with Nupage LDS sample buffer (NP0007, Thermo Fisher Scientific) according to manufacturer's instructions. Samples were loaded into sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE) gel, and then transferred to PVDF

membranes (Catalog #IPFL00010 Millipore, MA, USA), and incubated with blocking

buffer (927-60001, Li-cor). Membranes were incubated with primary antibodies overnight

at 4°C, washed 3 times with Tris-buffered saline with 0.05% Tween-20, and then with the

corresponding secondary antibody for 1 hour at room temperature. Membranes were

washed 3 times with Tris-buffered saline with 0.05% Tween-20. Finally, membranes were

visualized with Li-Cor Odyssey DLx using Image Studio. Antibodies against KEAP1

(Thermo Fisher Scientific, #10503-2-AP), NQO1 (Thermo Fisher Scientific, #11451-1-

AP), β-actin (66009-1-Ig, Proteintech) and GAPDH (sc-32233, Santa Cruz Biotechnology)

were used.

KEAP1 Knockout Generation

ARH-77 with deletion of the genes encoding KEAP1 was generated through

CRISPR-Cas9 deletion. Cells were electroporated in R Buffer (Thermo Fisher Scientific,

BR5) containing 150 ng/µl sgRNA and 500 ng/ul recombinant TrueCut Cas9 protein

(Thermo Fisher, A36496) using Invitrogen Neon Electroporator with 2 pulses of 1200 V,

2 ms duration. Three sgRNA sequences, 5' CCGUGUAGGCGAAUUCAAUG 3', 5'

ACCAACGGGCUGCGGGAGCA 3', and 5' UGGGCCAUGAACUGGGCGGC 3', were

used for electroporation. Cells were then seeded at a density of one cell per well in 96

well plates and grown until colonies were established. Colonies were tested for DNA

changes through PCR using KEAP1 primers, 5' AGGTGCTGGCCCAGTCCCAA 3' and

5' GCCGATGGCATTGCTGGGGT 3'. RNA was extracted from cells and tested for *NQO1*

(Thermo Fisher Scientific, Hs01045993_g1) and *KEAP1* (Thermo Fisher Scientific,

Hs01003506 _m1) expression with *GAPDH* normalization (Hs02758991_g1).

RNA-Seq Dataset Analysis

RNA-Sequencing data for each sample in each cell line was analyzed in Matlab.

Data were processed such that any genes with average expression values less than 1

fragments per kilobase of exon per million mapped fragments (FPKM) for all

concentrations were removed. Mattest was used to conduct an unpaired t-test. PCA

(principal component analysis) helped generate principal component coefficients to

quantify the two most important principal components, which could then be visualized.

The visualization helped determine the exclusion of sample 1 in control RPMI-8226, and

sample 3 in the wildtype ARH-77 (Supplemental Fig. 14). Non-small lung cancer cells

data from Depmap and cancer patient TCGA data from cBioportal, (62), were analyzed

in R. Expression of cell lines from Depmap were processed in $\log_2$(Transcripts Per Million

+ 1) while TCGA expression used $\log_2$(normalized counts + 1). Normalized counts were

calculated using RSEM (batch normalized) and downloaded from cbioportal. Cell lines

and tumor samples were classified as Nrf2/KEAP1 wildtype or mutant if non-conservative

mutations were present in the DNA sequencing data. RNA expression was compared

between wildtype and knockout samples for differential expression. Gene set enrichment

analysis (Broad Institute Software) was performed on RNA-sequencing data from cell

lines.

Nrf2 Target Genes Identification

After processing, genes that were upregulated more than 25% on average from

DMSO to low CDDO-2P-Im dose were identified as candidate Nrf2 upregulated genes. A

two-way student t-test using unpaired unequal variance was used to calculate a P value

for each comparison. Genes that had upregulation and a p value of less than 0.1 (one

way t-test of 0.05) were considered candidate Nrf2 target genes for each cell line. A

similar classification was used for the KEAP1 KO ARH-77 model between WT and KO

samples. Nrf2-induced genes in LUAD TCGA samples, from the cBioportal, and the

NSCLC CCLE samples, from the DepMap, were also found by selecting the top 500

upregulated genes in each sample, ordered by lowest p value.

These lists of candidate Nrf2 target genes were compared to each other to

determine which genes that were induced across cell lines. Genes were determined to

be highly upregulated if their expression level increased two-fold after drug treatment,

following genetic modification, or if they were part of the top 100 most significant

upregulated genes by p value in the TCGA or CCLE databases. A similar analysis was

repeated for downregulated genes for each cell line, in which all genes with a significant

(p-value < 0.1) reduction in expression were compared with no minimum threshold for

downregulation. In the TCGA and CCLE datasets, the top 250 downregulated genes were

chosen as targets.

For the individual Nrf2 target genes, LASAGNA-search 2.0 tool (63) was used to

predict *NFE2L2*/ARE binding sites. For each gene, a distance of 0 to -5000 of the

transcription start site (TSS) was selected and a cutoff of 0.0005 p value was used.

Binding sites were numbered according to lowest estimated p value.

N2CS and N2AS Calculation

To calculate the N2CS and N2AS, the 14 core Nrf2 biomarker gene expressions

were used. Gene expression was standardized using FPKM for each set of calculations.

For the N2CS, the control is set as the baseline and all gene expression changes were

expressed as Log$_2$ Fold Change relative to the baseline. The fold changes of the 14 core

Nrf2 genes are used to calculate the N2CS.

557
$$N2CS = \frac{\sum FC_N}{n}$$

558   In this calculation, FC$_N$ is the individual fold changes for each core Nrf2 gene, and n is

559   the number of core Nrf2 genes which is 14 in our case. For N2AS, a baseline expression

560   must be established from the group of samples. The mean expression for each gene

561   from the control is set as the baseline and the standard deviation is used to help calculate

562   a z-score for each gene expression.

563
$$Z = \frac{x - \mu}{\sigma}$$

564   Thus, the z-score is calculated by the taking the difference of the expression value, x, and

565   the mean, μ, and divide the difference by σ, the standard deviation of this gene

566   expression. The final N2AS is the average of the z-scores of the 14 core Nrf2 genes.

567
$$N2AS = \frac{\sum Z_N}{n}$$

568   The Z$_n$ is the individual z-scores for each gene which is summed up and divided by the total

569   number of genes in the gene set, *n*. When evaluating other Nrf2 gene sets, instead of the 14 core

570   genes, other genes were substituted to calculated a new Nrf2 activity score.

Drug Resistance Modeling

Data from Depmap of gene expression and CTRP2 experimental data was

downloaded and analyzed in R (64). Information about gene expression and drug

sensitivity (Area Under the Curve) non-small lung cancer cell lines data were collected

and merged in R. Cell line expression data was paired for each cell and the N2AS score

was calculated for each cell line. A Pearson correlation was performed for every gene or

gene set and the drug AUC. The graphs of the area under the curve and N2AS score was

plotted for all cell lines using ggplot in R. For analysis of other drugs, the datasets

(GSE41929, GSE48812, GSE135842, GSE199779) were used for analysis. Code for

analysis will be published.

Cell Viability Assay

KEAP1 knockout and wildtype ARH-77 cells were plated in 96 well plates with 2.5

x 10^4 cells per well. 1% DMSO or various concentrations of drugs were added the well.

After 48 hours, CellTiter-Glo™ (Promega Corp, Madison, WI) was used to investigate cell

viability according to manufacturer's instructions. Necrosulfonamide (34702, Cayman

Chemical), PX-12 (14192, Cayman Chemical), Paclitaxel (10461, Cayman Chemical),

and bardoxolone methyl (SMB00376, Sigma Aldrich) Experiments were done in

triplicates and repeated twice. IC_{50} was calculated using CompuSyn software (65).

Colony Forming Assay

KEAP1 knockout and wildtype ARH-77 cells were plated in 6 well plates with 1000

cells per well. Cells were mixed in MethoCult™ H4434 Classic media (StemCell

Technologies). Cells were allowed to grow for 14 days before they were imaged using

Keyence BZ-X810. Colonies of more than 10 cells were counted on 10 different views at

x20 focus for each well. Experiments were done in triplicates and repeated twice.

Cancer Prognosis Modeling

TCGA Data from cBioportal was collected and analyzed in R. Additional data from

GEO, Biostudies (66), and International Cancer Genome Consortium (ICGC) was also

collected and processed as needed (Supplemental Table 4). RNA-Seq data was

processed to Transcripts per million (TPM) while microarray array data was used as is. R

packages Survival and Cutpointr were used to help analyze survival data.

Information about patients' clinical outcomes and N2AS gene expression profiles

were combined in R. Z scores for gene expression was used when present or calculated

using TPM or microarray data as needed. Patients were subsequently ranked based on

their N2AS and corresponding hazard ratios and ranked log p-values were calculated for

every cutoff using Survival. These numbers were graphed onto a LOCC graph.

For training, the optimal cutoff consisting of at least 10% of the cohort was selected

by Cutpointr (lowest p-value) and manually verified to be appropriate by LOCC

visualization. This N2AS cutoff was selected for this cancer and used in the subsequent

validation. For a cancer to pass training, the p value must be lower than $p = 0.01$ for at

least one cutoff. For validation, instead of using the optimal cutoff, the previous N2AS

activity score was used to separate groups. For validation, a p value of 0.05 was

considered significant. Kaplan-Meier plots were generated at each cutoff and calculations

for hazard ratio and p value were performed using Survival. LOCC graphs were

generated according to the previous study (35).

Data Availability

All RNA-Sequencing data is online through GEO (GSE228434, GSE221526,

GSE229914, GSE229704, GSE229652). Other published datasets used in this study are

listed in supplemental table 4.

Author's Disclosures

The authors declare no potential conflicts of interest.

Authors' Contributions

Conceptualization, G.L, J.J.L; Methodology, G.L., J.J.L., H.K., E.J., E.R.C; Formal

Analysis, G.L., H.K., E.J., K.A; Investigation, G.L., H.K., K.A., E.J., S.R, Resources, G.L.,

K.A., S.R, A.S., H.M., Writing-Original Draft, G.L., J.L., H.K., Writing – Review & Edits,

G.L., J.L., K.A., Supervision, G.L., J.J.L; Funding Acquisition, J.J.L, H.M.

Acknowledgments

G.L is supported by NIH MSTP training grant 5T32GM007250. J.J.L. is supported by the

Jane and Lee Seidman Chair in Pediatric Cancer Innovation. Graphics were made with

the aid of Biorender. We would like to acknowledge Dr. Michael Sporn for his donation of

CDDO-2P-Im and contribution to the field of Nrf2 and synthetic triterpenoids. Dr. Sporn

passed away in late 2022.

REFERENCES

1. Ma Q. Role of nrf2 in oxidative stress and toxicity. Annu Rev Pharmacol Toxicol. 2013;53:401-26.
2. Taguchi K, Yamamoto M. The KEAP1-NRF2 System in Cancer. Front Oncol. 2017;7:85.
3. Singh A, Misra V, Thimmulappa RK, Lee H, Ames S, Hoque MO, et al. Dysfunctional KEAP1-NRF2 interaction in non-small-cell lung cancer. PLoS Med. 2006;3(10):e420.
4. Shibata T, Kokubu A, Gotoh M, Ojima H, Ohta T, Yamamoto M, et al. Genetic alteration of Keap1 confers constitutive Nrf2 activation and resistance to chemotherapy in gallbladder cancer. Gastroenterology. 2008;135(4):1358-68, 68 e1-4.
5. Shibata T, Ohta T, Tong KI, Kokubu A, Odogawa R, Tsuta K, et al. Cancer related mutations in NRF2 impair its recognition by Keap1-Cul3 E3 ligase and promote malignancy. Proc Natl Acad Sci U S A. 2008;105(36):13568-73.
6. Dinkova-Kostova AT, Kostov RV, Kazantsev AG. The role of Nrf2 signaling in counteracting neurodegenerative diseases. FEBS J. 2018;285(19):3576-90.
7. Saha S, Buttari B, Panieri E, Profumo E, Saso L. An Overview of Nrf2 Signaling Pathway and Its Role in Inflammation. Molecules. 2020;25(22).
8. Pillai R, Hayashi M, Zavitsanou AM, Papagiannakopoulos T. NRF2: KEAPing Tumors Protected. Cancer Discov. 2022;12(3):625-43.
9. Cuadrado A, Rojo AI, Wells G, Hayes JD, Cousin SP, Rumsey WL, et al. Therapeutic targeting of the NRF2 and KEAP1 partnership in chronic diseases. Nat Rev Drug Discov. 2019;18(4):295-317.
10. Cuadrado A, Manda G, Hassan A, Alcaraz MJ, Barbas C, Daiber A, et al. Transcription Factor NRF2 as a Therapeutic Target for Chronic Diseases: A Systems Medicine Approach. Pharmacol Rev. 2018;70(2):348-83.
11. Yagishita Y, Gatbonton-Schwager TN, McCallum ML, Kensler TW. Current Landscape of NRF2 Biomarkers in Clinical Trials. Antioxidants (Basel). 2020;9(8).
12. Dinkova-Kostova AT, Copple IM. Advances and challenges in therapeutic targeting of NRF2. Trends Pharmacol Sci. 2023;44(3):137-49.
13. Liu J, Lichtenberg T, Hoadley KA, Poisson LM, Lazar AJ, Cherniack AD, et al. An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. Cell. 2018;173(2):400-16 e11.
14. Cao M, Onyango EO, Williams CR, Royce DB, Gribble GW, Sporn MB, et al. Novel synthetic pyridyl analogues of CDDO-Imidazolide are useful new tools in cancer prevention. Pharmacol Res. 2015;100:135-47.
15. Cancer Cell Line Encyclopedia C, Genomics of Drug Sensitivity in Cancer C. Pharmacogenomic agreement between two cancer cell line data sets. Nature. 2015;528(7580):84-7.
16. Tsherniak A, Vazquez F, Montgomery PG, Weir BA, Kryukov G, Cowley GS, et al. Defining a Cancer Dependency Map. Cell. 2017;170(3):564-76 e16.
17. Rooney JP, Chorley B, Hiemstra S, Wink S, Wang X, Bell DA, et al. Mining a human transcriptome database for chemical modulators of NRF2. PLoS One. 2020;15(9):e0239367.

18. Chorley BN, Campbell MR, Wang X, Karaca M, Sambandan D, Bangura F, et al. Identification of novel NRF2-regulated genes by ChIP-Seq: influence on retinoid X receptor alpha. Nucleic Acids Res. 2012;40(15):7416-29.

19. Patra KC, Hay N. The pentose phosphate pathway and cancer. Trends Biochem Sci. 2014;39(8):347-54.

20. Kobayashi EH, Suzuki T, Funayama R, Nagashima T, Hayashi M, Sekine H, et al. Nrf2 suppresses macrophage inflammatory response by blocking proinflammatory cytokine transcription. Nat Commun. 2016;7:11624.

21. Ahmad R, Raina D, Meyer C, Kharbanda S, Kufe D. Triterpenoid CDDO-Me blocks the NF-kappaB pathway by direct inhibition of IKKbeta on Cys-179. J Biol Chem. 2006;281(47):35764-9.

22. Beaver LM, Buchanan A, Sokolowski EI, Riscoe AN, Wong CP, Chang JH, et al. Transcriptome analysis reveals a dynamic and differential transcriptional response to sulforaphane in normal and prostate cancer cells and suggests a role for Sp1 in chemoprevention. Mol Nutr Food Res. 2014;58(10):2001-13.

23. Taguchi K, Takaku M, Egner PA, Morita M, Kaneko T, Mashimo T, et al. Generation of a New Model Rat: Nrf2 Knockout Rats Are Sensitive to Aflatoxin B1 Toxicity. Toxicol Sci. 2016;152(1):40-52.

24. Velica P, Davies NJ, Rocha PP, Schrewe H, Ride JP, Bunce CM. Lack of functional and expression homology between human and mouse aldo-keto reductase 1C enzymes: implications for modelling human cancers. Mol Cancer. 2009;8:121.

25. Copple IM, Shelton LM, Walsh J, Kratschmar DV, Lister A, Odermatt A, et al. Chemical tuning enhances both potency toward nrf2 and in vitro therapeutic index of triterpenoids. Toxicol Sci. 2014;140(2):462-9.

26. Weniger MA, Rizzatti EG, Perez-Galan P, Liu D, Wang Q, Munson PJ, et al. Treatment-induced oxidative stress and cellular antioxidant capacity determine response to bortezomib in mantle cell lymphoma. Clin Cancer Res. 2011;17(15):5101-12.

27. Songbo M, Lang H, Xinyong C, Bin X, Ping Z, Liang S. Oxidative stress injury in doxorubicin-induced cardiotoxicity. Toxicol Lett. 2019;307:41-8.

28. You BR, Shin HR, Han BR, Park WH. PX-12 induces apoptosis in Calu-6 cells in an oxidative stress-dependent manner. Tumour Biol. 2015;36(3):2087-95.

29. Sun L, Wang H, Wang Z, He S, Chen S, Liao D, et al. Mixed lineage kinase domain-like protein mediates necrosis signaling downstream of RIP3 kinase. Cell. 2012;148(1-2):213-27.

30. Guan L, Nambiar DK, Cao H, Viswanathan V, Kwok S, Hui AB, et al. NFE2L2 Mutations Enhance Radioresistance in Head and Neck Cancer by Modulating Intratumoral Myeloid Cells. Cancer Res. 2023;83(6):861-74.

31. Drewinko B, Mars W, Stragand JJ, Henderson SD, Latreille J, Barlogie B, et al. ARH-77, an established human IgG-producing myeloma cell line. II. Growth kinetics, clonogenic capacity, chalone production, xenogeneic transplantations, and response to melphalan. Cancer. 1984;54(9):1893-903.

32. Takahashi T, Sonobe M, Menju T, Nakayama E, Mino N, Iwakiri S, et al. Mutations in Keap1 are a potential prognostic factor in resected non-small cell lung cancer. J Surg Oncol. 2010;101(6):500-6.

33.   Solis LM, Behrens C, Dong W, Suraokar M, Ozburn NC, Moran CA, et al. Nrf2
and Keap1 abnormalities in non-small cell lung carcinoma and association with
clinicopathologic features. Clin Cancer Res. 2010;16(14):3743-53.
34.   Saleh MM, Scheffler M, Merkelbach-Bruse S, Scheel AH, Ulmer B, Wolf J, et al.
Comprehensive Analysis of TP53 and KEAP1 Mutations and Their Impact on Survival in
Localized- and Advanced-Stage NSCLC. J Thorac Oncol. 2022;17(1):76-88.
35.   Luo G, Letterio, J. LOCC: a novel visualization and scoring of cutoffs for
continuous variables. bioRxiv. 2023.
36.   Chen J, Yang H, Teo ASM, Amer LB, Sherbaf FG, Tan CQ, et al. Genomic
landscape of lung adenocarcinoma in East Asians. Nat Genet. 2020;52(2):177-86.
37.   Zhang M, Zhang C, Zhang L, Yang Q, Zhou S, Wen Q, et al. Nrf2 is a potential
prognostic marker and promotes proliferation and invasion in human hepatocellular
carcinoma. BMC Cancer. 2015;15:531.
38.   Yuki H, Kamai T, Murakami S, Higashi S, Narimatsu T, Kambara T, et al.
Increased Nrf2 expression by renal cell carcinoma is associated with postoperative
chronic kidney disease and an unfavorable prognosis. Oncotarget. 2018;9(47):28351-
63.
39.   Uruno A, Yamamoto M. The KEAP1-NRF2 System and Neurodegenerative
Diseases. Antioxid Redox Signal. 2023.
40.   Tang W, Jiang YF, Ponnusamy M, Diallo M. Role of Nrf2 in chronic liver disease.
World J Gastroenterol. 2014;20(36):13079-87.
41.   Rangasamy T, Guo J, Mitzner WA, Roman J, Singh A, Fryer AD, et al. Disruption
of Nrf2 enhances susceptibility to severe airway inflammation and asthma in mice. J
Exp Med. 2005;202(1):47-59.
42.   Sharma A, Parikh M, Shah H, Gandhi T. Modulation of Nrf2 by quercetin in
doxorubicin-treated rats. Heliyon. 2020;6(4):e03803.
43.   Wang XJ, Sun Z, Villeneuve NF, Zhang S, Zhao F, Li Y, et al. Nrf2 enhances
resistance of cancer cells to chemotherapeutic drugs, the dark side of Nrf2.
Carcinogenesis. 2008;29(6):1235-43.
44.   Galan-Cobo A, Sitthideatphaiboon P, Qu X, Poteete A, Pisegna MA, Tong P, et
al. LKB1 and KEAP1/NRF2 Pathways Cooperatively Promote Metabolic
Reprogramming with Enhanced Glutamine Dependence in KRAS-Mutant Lung
Adenocarcinoma. Cancer Res. 2019;79(13):3251-67.
45.   LeBoeuf SE, Wu WL, Karakousi TR, Karadal B, Jackson SR, Davidson SM, et al.
Activation of Oxidative Stress Response in Cancer Generates a Druggable Dependency
on Exogenous Non-essential Amino Acids. Cell Metab. 2020;31(2):339-50 e4.
46.   Ding H, Chen Z, Wu K, Huang SM, Wu WL, LeBoeuf SE, et al. Activation of the
NRF2 antioxidant program sensitizes tumors to G6PD inhibition. Sci Adv.
2021;7(47):eabk1023.
47.   Tannir NM, Agarwal N, Porta C, Lawrence NJ, Motzer R, McGregor B, et al.
Efficacy and Safety of Telaglenastat Plus Cabozantinib vs Placebo Plus Cabozantinib in
Patients With Advanced Renal Cell Carcinoma: The CANTATA Randomized Clinical
Trial. JAMA Oncol. 2022;8(10):1411-8.
48.   Brown JE. Glutaminase Inhibitors-Do They Have a Role in the Treatment of
Metastatic Clear-Cell Renal Cell Carcinoma? JAMA Oncol. 2022;8(10):1418-9.

49.   Liby KT, Sporn MB. Synthetic oleanane triterpenoids: multifunctional drugs with a
broad range of applications for prevention and treatment of chronic disease. Pharmacol
Rev. 2012;64(4):972-1003.
50.   Chauhan D, Li G, Podar K, Hideshima T, Shringarpure R, Catley L, et al. The
bortezomib/proteasome inhibitor PS-341 and triterpenoid CDDO-Im induce synergistic
anti-multiple myeloma (MM) activity and overcome bortezomib resistance. Blood.
2004;103(8):3158-66.
51.   Computational Pan-Genomics C. Computational pan-genomics: status, promises
and challenges. Brief Bioinform. 2018;19(1):118-35.
52.   Budczies J, Klauschen F, Sinn BV, Gyorffy B, Schmitt WD, Darb-Esfahani S, et
al. Cutoff Finder: a comprehensive and straightforward Web application enabling rapid
biomarker cutoff optimization. PLoS One. 2012;7(12):e51862.
53.   Ogluszka M, Orzechowska M, Jedroszka D, Witas P, Bednarek AK. Evaluate
Cutpoints: Adaptable continuous data distribution system for determining survival in
Kaplan-Meier estimator. Comput Methods Programs Biomed. 2019;177:133-9.
54.   Pepe MS, Janes H, Longton G, Leisenring W, Newcomb P. Limitations of the
odds ratio in gauging the performance of a diagnostic, prognostic, or screening marker.
Am J Epidemiol. 2004;159(9):882-90.
55.   Liu Z, Deng M, Wu L, Zhang S. An integrative investigation on significant
mutations and their down-stream pathways in lung squamous cell carcinoma reveals
CUL3/KEAP1/NRF2 relevant subtypes. Mol Med. 2020;26(1):48.
56.   Fox DB, Garcia NMG, McKinney BJ, Lupo R, Noteware LC, Newcomb R, et al.
NRF2 activation promotes the recurrence of dormant tumour cells through regulation of
redox and nucleotide metabolism. Nat Metab. 2020;2(4):318-34.
57.   Reed DA, Zhao Y, Bagheri Varzaneh M, Shin JS, Rozynek J, Miloro M, et al.
NG2/CSPG4 regulates cartilage degeneration during TMJ osteoarthritis. Front Dent
Med. 2022;3.
58.   Kechin A, Boyarskikh U, Kel A, Filipenko M. cutPrimers: A New Tool for Accurate
Cutting of Primers from Reads of Targeted Next Generation Sequencing. J Comput
Biol. 2017;24(11):1138-43.
59.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory
requirements. Nat Methods. 2015;12(4):357-60.
60.   Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL.
StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. Nat
Biotechnol. 2015;33(3):290-5.
61.   Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics.
2010;26(1):139-40.
62.   Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, et al. The cBio
cancer genomics portal: an open platform for exploring multidimensional cancer
genomics data. Cancer Discov. 2012;2(5):401-4.
63.   Lee C, Huang CH. LASAGNA-Search 2.0: integrated transcription factor binding
site search and visualization in a browser. Bioinformatics. 2014;30(13):1923-5.
64.   Rees MG, Seashore-Ludlow B, Cheah JH, Adams DJ, Price EV, Gill S, et al.
Correlating chemical sensitivity and basal gene expression reveals mechanism of
action. Nat Chem Biol. 2016;12(2):109-16.

65.   Chou T-C, Martin, N. A Computer Program for Quantitation of Synergism and
Antagonism in Drug Combinations, and the Determination of IC50 and ED50 Values.
CompuSyn For Drug Combinations User's Guide.
66.   Sarkans U, Gostev M, Athar A, Behrangi E, Melnichuk O, Ali A, et al. The
BioStudies database-one stop shop for all data supporting a life sciences study. Nucleic
Acids Res. 2018;46(D1):D1266-D70.

Supporting Information

Supplemental Figure S1. Gene Set Enrichment Analysis of Pharmacological or
Genetic Induction of Nrf2.

Supplemental Figure S2. Literature references of candidate core Nrf2 genes as
target genes.

Supplemental Figure S3. Induction of candidate core Nrf2 gene expression after
drug treatment or genetic modification.

Supplemental Figure S4. AKR1C family gene show significant differences in gene
induction in drug treatment and genetic modification.

Supplemental Fig S5. Predicted Nrf2 binding sites in near universal and
conditional Nrf2 genes.

Supplemental Fig S6. Activation of Nrf2 by genetic modifications may lead to
suppression of inflammatory genes.

Supplemental Fig S7. Validation of the near universal Nrf2 Gene Set in Other
Published Datasets.

Supplemental Fig S8. Activation of Nrf2 is correlated to drug resistance and
increased colony formation *in vitro*.

Supplemental Fig S9. Mutations of NFE2L2 and KEAP1 were not significantly
associated with prognosis in TCGA datasets.

Supplemental Fig S10. Analysis of TCGA cancers that did not pass training with
Nrf2 activity score.

Supplemental Fig S11. Analysis of TCGA cancers that passed training with Nrf2
activity score but did not have a validation cohort.

Supplemental Fig S12. Analysis of TCGA cancers that passed training with Nrf2
activity score but not validation.

Supplemental Fig S13. Other Nrf2 gene sets is associated with survival for some
cancers but not others.

Supplemental Figure S14. Exclusion of Irregular Control Samples Affecting Nrf2
Target Genes.

Supplemental Table S1. Core Nrf2 Target Gene Fold Changes

Supplemental Table S2. Near Universal Nrf2 Target Gene Fold Changes

Supplemental Table S3. Conditional Nrf2 Target Gene Fold Changes

Supplemental Table S4. OVCAR8 Low Concentration CDDO-2P-Im Upregulated
Gene Fold Changes

Supplemental Table S5. SF8628 Low Concentration CDDO-2P-Im Upregulated
Gene Fold Changes

Supplemental Table S6. Primary Dermal Fibroblast Low Concentration CDDO-2P-
Im Upregulated Gene Fold Changes

Supplemental Table S7. RPMI-8226 Low Concentration CDDO-2P-Im Upregulated
Gene Fold Changes

Supplemental Table S8. ARH-77 KEAP1 Knockout Upregulated Gene Fold
Changes

Supplemental Table S9. TCGA LUAD Nrf2 Gene Fold Changes

Supplemental Table S10. CCLE NSCLC Nrf2 Gene Fold Changes

Supplemental Table S11. Nrf2 Activity Score Drug Correlations

Supplemental Table S12. CDDO-2P-Im Drug Concentrations

Supplemental Table S13. Nrf2 Validation and Change Score Datasets

Supplemental Table S14. Nrf2 Cancer Prognosis Datasets

FIGURE LEGENDS

Figure 1. Generation of Transcriptomic Databases of Nrf2 Activation. **A,** Schematic

overview of the Nrf2/KEAP1 pathway and our approach towards understanding

transcriptional consequences. Cells are either treated with CDDO-2P-Im to inhibit

KEAP1-Nrf2 binding or have mutations leading to Nrf2 activation. Transcripts are taken

from cells and then analyzed by RNA-Sequencing. **B-E,** *NQO1* expression by RNA-

Sequencing is significantly increased after treatment with CDDO-2P-Im in (**B**) SF8628,

(**C**) OVCAR8, (**D**) Primary Dermal Fibroblasts, (**E**) RPMI-8226. **F,** DNA of KEAP1

knockout (KO) ARH-77 cells is sequenced following using CRISPR-Cas9 editing. **G,**

KEAP1 exon 5 RNA is evaluated by RT-qPCR following CRISPR-Cas9 editing. **H,**

Western blot of KEAP1 protein is performed after CRISPR-Cas9 editing. **I,** *NQO1* RNA

is evaluated by RT-qPCR following CRISPR-Cas9 editing. **J,** NQO1 protein is evaluated

by western blot following CRISPR-Cas9 editing. **K-L**, *NQO1* RNA expression is

evaluated between *KEAP1* or *NFE2L2* mutant and WT tumor samples using (**K**) TCGA

LUAD database or (**L**) Depmap's NSCLC CCLE Database. Cancer cell lines are from

the following cell types: SF8628 – Diffuse Intrinsic Pontine Glioma, OVCAR8 – Ovarian

Cancer, RPMI-8226 – Multiple Myeloma, ARH-77 – B cell lymphoma. LUAD – Lung

Adenocarcinoma, NSCLC – Non-small cell lung cancer. All RNA-Sequencing

experiments were performed once with 3 replicates. Other experiments were performed

independently twice with similar results. Values = means ± SD (n = 3). ns, P ≥ 0.05; *, P

< 0.05; **, P < 0.01.

Figure 2. A Core Nrf2 Transcriptional Signature is Present after Nrf2 Activation. **A,**

Heatmap of candidate core Nrf2 genes following low dose of CDDO-2P-Im (underlined

cell lines) or genetic alteration of Nrf2/KEAP1. Highly upregulated in red represents at

least a 2 fold change while upregulated represents at least a 1.25 fold change. Not

expressed means that gene expression is less than 1 fragments per kilobase of exon

per million mapped fragments (FPKM). **B,** Heatmap of near universal upregulated Nrf2

genes following Nrf2 activation. **C-E**, Gene expression fold changes by RNA-

sequencing were evaluated for the following genes: (**C**) *SRXN1*, (**D**) *ABHD4*, and (**E**)

AKR1C3. **F,** Nrf2/ARE (Antioxidant Response Element) binding motif analysis was

performed on candidate core Nrf2 genes from transcription start site (TSS) to -5000

base pairs. ARE numbers are ranked based on highest likelihood of Nrf2 binding based

on sequence similarity to consensus sequence. Only binding sequences that have a p

value of 0.0005 or lower are shown. All RNA-Sequencing experiments were performed

once. Abbreviations: C – Control, S_L – SF8268 low dose 2P, S_H – SF8268 high dose

2P, O_L – OVCAR8 low dose 2P, O_H – OVCAR8 high dose 2P, F_L – Primary Dermal

Fibroblast low dose 2P, F_H – Primary Dermal Fibroblast high dose 2P, R_L – RPMI-8226

low dose 2P, R_H – RPMI-8226 high dose 2P, A_K – ARH-77 KEAP1 knockout. Values =

means ± SD (n = 3). ns, $P ≥ 0.05$; *, $P < 0.05$; **, $P < 0.01$. ~ indicates this gene did not

pass subsequent validation with other datasets.

Figure 3. Defining Conditionally Upregulated and Novels Targets Genes of Nrf2

Activation. **A,** Heatmap of conditional, novel Nrf2 target genes following low dose of

CDDO-2P-Im (underlined cells line) or genetic alteration of Nrf2/KEAP1. **B-C,** Gene

expression changes of primarily drug-induced Nrf2 target genes, (**B**) *HMOX1* and (**C**)

AKIRIN2. **D-G,** Gene expression changes of pentose phosphate pathway enzymes, (**D**)

G6PD, (**E**) *TKT*, (**F**) *TALDO1*, (**G**) *PGD*). **H-I,** Gene expression changes of novel

upregulated Nrf2 target genes, (**H**) *B4GALNT1* and (**I**) *POPDC3*). **J-L,** Gene expression

changes of novel downregulated Nrf2 target genes, (**J**) *PLAT,* (**K**) *CDC42EP1,* (**L**)

EVA1A). All RNA-Sequencing experiments were performed once. Abbreviations: C –

Control, S_L – SF8268 low dose 2P, S_H – SF8268 high dose 2P, O_L – OVCAR8 low dose

2P, O_H – OVCAR8 high dose 2P, P_L – Primary Dermal Fibroblast low dose 2P, P_H –

Primary Dermal Fibroblast high dose 2P, R_L – RPMI-8226 low dose 2P, R_H – RPMI-

8226 high dose 2P, A_K – ARH-77 KEAP1 knockout. Values = means ± SD (n = 3). ns,

$P \geq 0.05$; *, $P < 0.05$; **, $P < 0.01$. ~ indicates this gene did not pass subsequent

validation with other datasets.

Figure 4. Validation of the Core Nrf2 Gene Set in Other Published Datasets. **A-C,**

Candidate core Nrf2 genes are evaluated in TCGA (**A**) head neck squamous cell

carcinoma (HNSC), (**B**) lung squamous cell carcinoma (LUSC), (**C**) liver hepatocellular

carcinoma (LIHC) cancer datasets with significant number of *KEAP1*/*NFE2L2*

mutations. **D,** Candidate core Nrf2 genes are evaluated using published data which

uses Nrf2 activator, sulforaphane, for 24 hours. **E,** In a previously published rat model of

WT and Nrf2 KO, candidate core Nrf2 were evaluated to see if gene expression

changes were observed after treatment with a Nrf2 activator, CDDO-Im, and if Nrf2

knockout affect upregulation. **F,** Nrf2 change score (N2CS) was evaluated across

different treatments of CDDO-2P-Im or KEAP1 knockout. **G**, N2CS was evaluated for

other drugs from published datasets. Abbreviations: C – Control, S_L – SF8268 low dose

2P, S_H – SF8268 high dose 2P, O_L – OVCAR8 low dose 2P, O_H – OVCAR8 high dose

2P, P_L – Primary Dermal Fibroblast low dose 2P, P_H – Primary Dermal Fibroblast high

dose 2P, R_L – RPMI-8226 low dose 2P, R_H – RPMI-8226 high dose 2P, A_K – ARH-77

KEAP1 knockout, BTZ – Bortezomib, DOXO – doxorubicin, PTX – Paclitaxel, SFN –

Sulforaphane. Values = means ± SD (n = 3). ns, $P \geq 0.05$; *, $P < 0.05$; **, $P < 0.01$. ~

means this gene did not pass validation with other datasets.

Figure 5. A Nrf2 Activity Score is Correlated with Resistance to Radiation and Certain

Drugs. **A,** Nrf2 Activity Score (N2AS) is calculated for each sample from the TCGA

LUAD and color coded by mutation status of Nrf2/KEAP1. **B-C**, N2AS for each cell line

was plotted against sensitivity to (**B**) necrosulfonamide and (**C**) PX-12 represented by

area under the curve (AUC). The correlation was calculated for N2AS and the AUC. **D-**

E, WT and KEAP1 KO ARH-77 were treated with (**D**) necrosulfonamide or (**E**) PX-12 for

48 hours and evaluated for cell viability by Cell-Titer Glo. **F,** Radiation was performed

on WT and KEAP1 KO ARH-77 cells at indicated radiation doses. Cells were incubated

in 96 well plates for 72 hours and cell viability was measured by Cell-Titer Glo. **G-H**,

N2AS for each cell line was plotted against sensitivity to (**G**) paclitaxel and (**H**)

bardoxolone methyl represented by area under the curve (AUC). The correlation was

calculated for N2AS and the AUC. **I-J**, WT and KEAP1 KO ARH-77 were treated with

(**I**) paclitaxel or (**J**) bardoxolone methyl for 48 hours and evaluated for cell viability by

Cell-Titer Glo. Values = means ± SD (n = 3). ns, $P \geq 0.05$; *, $P < 0.05$; **, $P < 0.01$. All

Cell-Titer Glo experiments were repeated independently twice with similar results.

Figure 6. Nrf2 Activity is Associated with Poor Prognosis in a Variety of Cancers. **A**,

Flow chart of the training and validation results using N2AS to evaluate cancer

prognosis. **B**, Results for individual TCGA cancers are categorized. Cancers colored in

red represent a high N2AS is associated with a poor prognosis while those in green

represent a high N2AS is associated with a good prognosis. **C,** TCGA LUAD was

trained using LOCC to find the most significant cutoff at N2AS of 0.463. **D,** OncoSG

LUAD was used as validation with the cutoff of N2AS of 0.463 to determine significance

and reproducibility. **E**, TCGA KIRC was trained using LOCC to find the most significant

cutoff at N2AS of 0.488. **F,** MTAB-1980 was used as validation with the cutoff of N2AS

of 0.488 to determine significance and reproducibility. **G**, TCGA LIHC was trained using

LOCC to find the most significant cutoff at N2AS of 0.604. **H,** LIKI-JP was used as

validation with the cutoff of N2AS of 0.604 to determine significance and reproducibility.

I, TCGA LAML was trained using LOCC to find the most significant cutoff at N2AS of

0.432. **J,** OSHU AML was used as validation with the cutoff of N2AS of 0.432 to

determine significance and reproducibility. Abbreviations: LUAD – Lung

adenocarcinoma, KIRC/CCRCC – Kidney renal clear cell carcinoma, LIHC/LIKI – Liver

hepatocellular carcinoma, LAML/AML – Acute myeloid leukemia, HNSC – Head and

neck squamous cell carcinoma, SKMC – skin cutaneous melanoma, PAAD - pancreatic

adenocarcinoma, BLCA - Bladder urothelial carcinoma, KIRP – Kidney renal papillary

cell carcinoma, LGG - Brain lower grade glioma, UCEC - Uterine corpus endometrial

carcinoma, OV - Ovarian serous cystadenocarcinoma.

Fig 1.

Fig 2.

A Candidate Core Genes

B Near Universal Genes

C SRXN1

TCGA LUAD   NSCLC CCLE

p = 4.82e-36   p = 2.25E-13

D ABHD4

TCGA LUAD   NSCLC CCLE

p = 6.54E-16   p = 1.32E-05

E AKR1C3

TCGA LUAD   NSCLC CCLE

p = 1.40E-38   p = 2.73E-11

F Predicted NRF2 Sites in Universal Genes

Fig 3.

Fig 4.

Fig 5.

Fig 6.