Whither unisensory olfactory cortex: processing of visual and auditory stimuli in olfactory cortex, independently of odor associations

Evelina Thunell[1,2], Moa G. Peter[1], Behzad Iravani[1,3] Danja K. Porada[1], Katharina Prenner[1], Fahimeh Darki[1], and Johan N. Lundström[1,4,5]

[1]Department of Clinical Neuroscience, Karolinska Institutet, 17177 Stockholm, Sweden
[2]Department of Psychological Sciences, Purdue University, West Lafayette, IN 47907, USA
[3]Department of Neurology and Neurological Sciences, Stanford University, CA 94305, USA
[4]Monell Chemical Senses Center, Philadelphia, PA 19104, USA
[5]Stockholm University Brain Imaging Centre, Stockholm University, 106 91 Stockholm, Sweden

Corresponding authors:

Evelina Thunell
Division of Psychology
Department of Clinical Neuroscience
Karolinska Institutet
Nobels väg 9, 171 77
Stockholm, Sweden
Email: evelina.thunell@ki.se

Johan N. Lundström
Division of Psychology
Department of Clinical Neuroscience
Karolinska Institutet
Nobels väg 9, 171 77
Stockholm, Sweden
Email: johan.lundstrom@ki.se

Abstract

Primary sensory cortices process atypical sensory objects in some conditions, which brings into question their presumed sensory specificity. Whether this atypical sensory processing originates from true atypical sensory processing or from cross-modal associations to objects is not known. Visual/auditory objects typically have strong reciprocal associations; hence, it is difficult to address this question in these modalities. Here, we instead use the olfactory system as a model system. We show that even in total absence of any odor reference, the posterior piriform cortex (PPC; odor object processing) is activated by both sounds and pictures of objects. This activation is independent of the objects' odor associations, thereby demonstrating that the activity is not due to cross-modal associations. In the anterior piriform cortex (APC; low-level odor processing) we found activation by sounds, again independent of odor associations. Analogously, we found clear cross-modal effects in visual and auditory brain regions: Pictures activated higher-level auditory cortex and caused a decreased signal in primary auditory cortex, whereas sounds activated primary visual cortex and caused a decreased signal in higher-level visual cortex. Our findings demonstrate an atypical sensory processing that is independent of cross-modal associations in olfactory cortices and contribute to a more nuanced view of modality specificity in olfactory, auditory, and visual cortices.

Introduction

The concept of a unisensory cortex has recently been challenged by demonstrations of primary sensory cortices in healthy individuals sometimes responding to atypical unisensory stimuli. For example, the primary visual cortex (V1) is commonly activated by visual stimulation but, under certain circumstances, also responds to auditory and tactile stimulation: in humans for abstract spoken words (Seydell-Greenwald et al., 2020) and, in rats, during tactile exploration (Vasconcelos et al., 2011). Two alternative explanations have emerged as to why a primary cortical area responds to atypical sensory stimuli: activations are due to *cross-modal associations* where the atypical sensory stimulus induces an association to a typical sensory stimulus, or, alternatively, activations constitute *true atypical unisensory processing*, where the atypical stimulus itself is processed in the primary area. Dissociating these theories in the audio-visual domain is a non-trivial task given the tight ecological connection between the visual and auditory systems where even simple stimuli, such as a light flash and a beep, are inherently associated when synchronous. The olfactory system does not come with the same caveat. It is difficult to identify odors without cues, odor sensations often lack spontaneous associations to other modalities, and objects that do not contain organic materials, such as a clock, are very seldom associated with odors (Cain, 1979). The unique olfactory system thus provides a model system for dissociating the two competing theories of why main sensory cortical areas respond to atypical stimuli.

The main olfactory cortex has long been considered to comprise the piriform cortex due to its role as the primary recipient of direct projections from the olfactory bulb (Zald and Pardo, 2000; Mainland et al., 2014). Much like the activation by atypical stimuli in our other sensory cortices, there are several accounts of the piriform cortex processing both unisensory non-olfactory stimuli (Gottfried et al., 2004; Maier et al., 2012; Porada et al., 2019; Zhou et al., 2019) and integrating multisensory information (Lundström et al., 2019; Porada et al., 2019). Indeed, there are direct anatomical connections to piriform cortex from the peripheral visual system (Cooper et al., 1994), auditory system (Budinger et al., 2000; Budinger and Scheich, 2009), and gustatory system (Datiche et al., 1996; Johnson et al., 2000), providing a potential route for atypical sensory information into initial processing stages of the olfactory system. However, as for other sensory cortices, it is unclear whether the piriform cortex responds to the unisensory stimuli per se (true atypical unisensory processing) or whether cross-modal

associations drive the activation (Ohla et al., 2017). In most studies that demonstrate piriform cortex activation by atypical unisensory stimuli, the stimuli were associated with odors either by presentation of odors during the experiment or by ecological or experimentally induced multisensory associations (e.g., Schulze et al., 2017; Olofsson et al., 2019). Moreover, associative processes, such as odor memory (Yang et al., 2021), attention (Zelano et al., 2005; Veldhuizen and Small, 2011), and cross-modal processing (Gottfried et al., 2004; Lundström et al., 2019; Zhou et al., 2019), are linked to piriform cortex activity.

To dissociate between the "cross-modal association" and the "true atypical unisensory processing" hypotheses, we exposed participants (*n*= 47) to unisensory visual and auditory representations of objects with varying degree of odor associations, from very low (e.g., a church bell) to strong (e.g., popcorn), while assessing neural responses with functional magnetic resonance imaging (fMRI). Critically, odors were not mentioned until after the scanning session when participants rated how much they associated each visual and auditory object to an odor. This design allowed us to dissociate between the two competing hypotheses because if potential activity in piriform cortex is dependent on cross-modal association, the response to an object should be linked to how much the individual associates it to an odor. On the other hand, if activity in piriform cortex is due to atypical unisensory processing, the neural response should be independent of odor association.

Methods

Participants

A total of 54 individuals participated in the experiment. Data from seven of these were excluded due to excessive sleepiness as defined by large amount of eye closure, judged from eye-tracker monitoring, and excessive head movement detected during pre-processing of the data. The remaining 47 individuals (28 women; mean age 33.5 years, *SD* = 10.1, range 18-46 years) reported being healthy and having normal or corrected-to-normal visual acuity, normal hearing (except for one who had lowered hearing on one ear), and normal olfactory functions. The procedures were in accordance with the Declaration of Helsinki and approved by the

Swedish Ethical Review Authority (Dnr. 2020-06533). All participants provided written informed consent prior to the experiment.

MR and physiological data acquisition

Neural responses to visual and auditory stimuli were assessed using a 3 T Magnetom Prisma Scanner (Siemens) with a 20-channel head coil. We acquired functional data using a 3D echo planar (EP) sequence (56 slices, repetition time (TR) = 1700 ms, echo time (TE) = 30 ms, flip angle (FA) = 70°, voxel size = 2.234 mm x 2.234 mm x 2.200 mm, field of view (FoV) = 94 x 94 voxels). Slices were tilted 20 degrees. A structural image was also obtained using a 3D MPRAGE T1-weighted sequence (208 slices, TR = 2300 ms, TE = 2.89 ms, FA = 9°, voxel size = 1 mm x 1 mm x 1 mm, FoV = 256 x 256 voxels). At the end, we acquired a narrow T2-weighted image that was not used in this study. Participants were instructed to breathe through their nose throughout the experiment and their respiratory trace was continuously recorded during the functional runs using birhinal nasal cannulas attached to a PowerLab 4/26 unit via a spirometer (ADInstruments Inc., Colorado Springs CO, USA). To evaluate participants' wakefulness during the functional runs, we continuously monitored their eyes using a TRACKPixx3 2kHz binocular distance eye tracker (VPixx Technologies Inc., Saint-Bruno, QC Canada).

Stimuli and task

The stimulus set comprised grayscale pictures and sounds of 48 easily identifiable objects, chosen to evoke various degrees of odor association. We selected this set based on a pilot study where 23 healthy participants identified and rated the level of their odor association to the pictures and sounds of 60 everyday objects. Sounds were obtained from the BBC Sounds Database, http://bbcsfx.acropolis.org.uk, and Freesound, https://freesound.org, and pictures were taken from the Open Image Dataset, https://opensource.google, and ImageNet, http://www.image-net.org. The sounds were seamlessly looped or trimmed to last 3 seconds and equalized with a root mean square equalization to -23 dB loudness (Regenbogen et al., 2016). Pictures were cropped symmetrically from all sides to square shape with a side length equal to the shortest of the original height and width and converted to grayscale using the

rgb2gray function in Matlab. The visual stimulation was presented on a 40'' 4K ultra-HD LCD display (NordicNeuroLab AS, Bergen, Norway) placed behind the scanner bore and viewed via a head coil mirror. The viewing distance was approximately 180 cm, and the pictures spanned 4.5 degrees of visual angle. The auditory stimulation was presented via binaural headphones (NordicNeuroLab AS, Bergen, Norway), at a loud volume that was adjusted when needed to be tolerable to the participant.

Procedure and design

Practice

To ensure that participants would recognize the objects, all stimuli were presented prior to MR scanning one at a time together with the name of the object. During this pre-scanning practice, pictures appeared with the object name written underneath, and sounds were played with the object name written in the middle of the screen. The participants were instructed to learn to identify the objects and erroneously told that they would later be tested on this task to increase motivation and attention.

MR scanning

The MR session included four functional runs. Each run lasted 10 min and 15 s and contained eight blocks lasting 1 min each. Within a block, six stimuli were presented, each with a duration of 3 s and preceded by 7 s ± 3 s of no stimulus. All blocks were preceded and followed by 15 s of no stimulus. The stimuli were grouped into pure sound and pure picture blocks. Further, the stimuli were grouped according to the average odor association level across the two modalities in the pilot study, such that the stimuli in each block belonged to either the highest or lowest rated half. The low and high odor association blocks were interleaved, with half of the participants starting with a low and the other half starting with a high odor association block. All 96 stimuli (48 sounds and 48 pictures) were presented once in the first half of the runs (runs 1 and 2), and a second time in the second half of the runs (runs 3 and 4), in a random order. The participants' task was to fixate a cross that was presented throughout each run except during picture presentation, and to identify each presented object without giving any response.

Odor association

After the MR session, participants rated how much they associated each stimulus with an odor using a visual analogue scale ranging from 0 (no odor association) to 100 (very strong odor association). Pictures and sounds were presented in alternating blocks with 12 stimuli per block. Critically, this task was the first mention of odors to the participants and no odors were presented at any point during the MR scanning. We also inquired participants whether they had thought about the smells of the objects during the MR scanning.

Imaging data processing and statistical analyses

Pre-processing was carried out separately for each participant using SPM12 (version 7487; Penny et al., 2011), including slice time correction, motion correction, coregistration of the functional runs, coregistration of the anatomical and functional data, and normalization to MNI space. The functional data were smoothed for whole-brain analyses using a 6 mm full width at half maximum kernel, while region of interest (ROI) analysis was performed on non-smoothed data. For one participant, the first functional run was excluded due to a headphone problem which was remedied before the three remaining runs.

First level statistical analyses were conducted using a General Linear Model (GLM), as implemented in SPM12. The BOLD signal responses to the pictures and sounds were modeled separately as a boxcar convolved with a canonical hemodynamic response function (HRF) as basis function. We used each participant's individual odor association ratings of the sounds and pictures, which were collected *after* the MR data acquisition, as parametric modulators (orthogonalized). To account for potential confounding motion effects, we added the six rigid-body motion parameters (i.e., 3 translational and 3 rotational variables) from the motion correction as regressors of no interest. The four regressors of interest were response to pictures, parametric modulation of the response to pictures by odor association, response to sounds, and parametric modulation of the response to sounds by odor association. We computed simple t-contrasts of each of these against implicit baseline.

Second level whole-brain analyses were conducted using one-sample t-tests on the four regressors of interest (participants' responses to pictures, modulation of these responses by the odor association ratings of the pictures, responses to the sounds, and modulation of these

responses by the odor association ratings of sounds). In a ROI analysis, we extracted mean contrast estimates in six bilateral ROIs representing basic and object-related neural processing within each sense. Namely, anterior piriform cortex (APC) and posterior piriform cortex (PPC) for olfaction, primary visual cortex (V1) and lateral occipital complex (LOC) for vision, and primary auditory cortex (A1) and higher order auditory cortex (hAC) for audition. These ROIs (Porada et al. 2019) are based on a combination of functional activations in previous studies and standardized anatomical masks to maximize specificity and independence. For each participant and ROI, we extracted mean contrast averages from the left and right hemispheres separately and subsequently averaged over the two hemispheres. We performed one-sample t-tests on the extracted voxel-averages for each ROI and contrast using an α level of 0.05. Whole-brain activations for these four effects are presented corrected for multiple comparisons using a peak-level threshold at p = .05 family-wise error (FWE). The imaging data had acceptable coverage in all ROIs (APC: 76%±12, PPC: 64%±13, V1: 98%±2, LOC: 89%±6, A1: 98%±2, hAC: 82%±8).

In an additional analysis, the ROI analysis described above was repeated with all conditions split up in low and high odor association groups based on the participants' individual ratings, separately for pictures and sounds. The 4 regressors of interest were thus replaced by 8 regressors (response to pictures with weak and strong odor association, parametric modulation by odor association of the response to weak and strong odor association pictures, response to sounds with weak and strong odor association, and parametric modulation by odor association of the response to sounds with weak and strong odor association). We also performed an additional analysis where each stimulus was assigned a separate regressor. In this GLM, there were in total 96 regressors of interest (48 for pictures and 48 for sounds); no parametric modulation was added. We ranked the stimuli according to each individual's smell association ratings, with random ordering in case of identical ratings for more than one stimulus, and correlated these rankings with the ROI activations, separately for pictures and sounds.

Stimuli, motion parameters, and ROI results are available on the open science framework (OSF) repository: https://osf.io/duwrb/?view_only=d214b458aee64204bf6e09df38249631.

Dynamic functional connectivity analysis

To assess the functional connectivity between PPC and both visual (LOC) and auditory (hAC) object processing areas, we used a dynamic functional connectivity (dFC) analysis. We opted to focus on LOC and hAC based on the assumption that object-related information constitutes more pertinent input for the olfactory system than the low-level features processed in the primary visual and auditory areas (V1 and A1). In general, the PPC receives more input from other cortical areas than does APC (Cohen et al., 2017), and we therefore consider only PPC in this analysis. The dFC was determined using a Hanning sliding window with a 5 time-points length and 50% overlap across time-points. Subsequently, we computed the Pearson correlation between different brain areas within each window. To implement this analysis, BOLD activity from 272 spherical ROIs with a radius of 18 mm were extracted across the whole brain. Only the BOLD activity of ROIs with a gray matter probability of at least 40% were assessed further. Next, a design matrix with intervals labeled either auditory or visual was created. These intervals represent the consecutive time points where each corresponding stimulus was presented to individuals. To account for the hemodynamic response, the design matrix was convolved with a HRF, and the 6 motion regressors were added to a linear regression model to remove the effects of motion. On the group level, we applied t-tests to the auditory and visual beta connectivity maps to produce group-level t-maps for visual and auditory stimuli. These maps were then averaged across hemispheres and arbitrarily thresholded to remove false positive connections so that only the 20% strongest connection were kept. The resulting functional connections were subsequently converted to distances matrices, by inverse transform, for which Floyd–Warshall algorithms (Cf. Fransson et al., 2011) were used to find the shortest path between each pair of nodes, i.e., between auditory and visual object processing areas and the PPC.

RESULTS

Olfactory cortex processes unisensory visual and auditory stimuli, independent of odor association

The anatomical separation between the anterior and posterior portions of the piriform cortex has been suggested to correspond to a functional specialization where the anterior portion

(APC) processes mainly low-level olfactory features whereas the posterior portion (PPC) is more involved in odor object processing (Gottfried, 2010; Lundström et al., 2011). This separation is analogous to the hierarchical organization of both the visual and auditory systems where activity in the primary visual cortex (V1) and primary auditory cortex (A1) reflects mostly low-level features whereas the lateral occipital complex (LOC) and higher order auditory cortex (hAC) process objects (Porada et al., 2019). Here, we used fMRI to assess potential atypical processing in both low- and high-level sensory cortices with both region of interest (ROI) and whole-brain analyses.

We first set out to test whether the PPC processes atypical unisensory objects using ROI specific analysis. We found an activation by pictures of objects in the PPC ROI (Figure 1A-B, $t(46) = 4.96$, $p = 9.99*10^{-6}$, Cohen's $d = 0.72$). Similarly, we found an activation by sounds of objects in PPC (Figure 1B, $t(46) = 8.82$, $p = 1.89*10^{-11}$, Cohen's $d = 1.29$). Next, we assessed whether the pictures and sounds also activated the APC. While we did find a significant activation in APC by sounds (Figure 1B, $t(46) = 6.04$, $p = 2.51*10^{-7}$, Cohen's $d = 0.88$), there was no apparent activation by pictures (Figure 1B, $t(46) = 0.24$, $p = .81$, Cohen's $d = 0.04$).

Figure 1. Region of interest MR imaging results for olfactory brain areas. A) Location of olfactory ROIs. **B)** In Anterior Piriform Cortex (APC), the extracted mean contrast estimates are significantly above zero for sounds (ear symbol) but not for pictures (eye symbol). In Posterior Piriform Cortex (PPC), both pictures and sounds elicited significant activations. **C)** In both APC and PPC, the mean parametric modulation by odor association is not significantly different from zero for sounds (ear symbol) nor for pictures (eye symbol). Asterisks mark statistical significance ($p < 9.99*10^{-6}$). Error bars depict ± 1 Standard Error of the Mean (SEM).

We used an ROI-based approach due to our region-specific hypothesis. A more conservative approach is a whole-brain analysis with correction for multiple comparisons across voxels. Using a whole-brain analysis, we could confirm two of our three ROI effects: We observed a response to sounds in both PPC and APC using a conservative peak-level threshold of $p < .05$ FWE (Figure 2 A-B).

Figure 2. Whole-brain MR imaging results. **A**) The location of the Anterior Piriform Cortex (APC) ROI is shown in top panel and is outlined in dark blue. The two cut-outs show activations in the ROIs in detail (pictures: eye symbol, sounds: ear symbol). The activation by sounds, but not that by pictures, is significant. **B**) The Posterior Piriform Cortex (PPC) ROI is outlined in dark blue in top pane and results in the two cut-outs. As for the APC, activation by sounds, but not pictures, is significant in the PPC. All results are corrected for multiple comparisons using a peak-level threshold at p = .05 family-wise error (FWE) across the whole field of view and visually masked in figure for inclusion using the two ROIs. ROIs are overlaid on MR template image in MRIcroGL.

These results demonstrate that olfactory cortex responds to unisensory visual and auditory objects. Critically, the activations occurred in the absence of any olfactory stimulation or explicit odor associations. No odors were presented at any point during the experiment and odors were first mentioned to the participants during the post-scanning association task. In the post-scanning interview, 30 out of the 47 participants (64%) said that they had not thought about the smells of the objects during the scanning. The remaining 17 participants (36%) reported that they had thought of, or might have thought of, the smell of one or a few objects. To test whether the activations were driven by odor associations that might have been automatically evoked when an object was viewed or heard, we assessed the neural modulation by participants' degree of odor association to each object. Importantly, these ratings were obtained post-scanning and therefore could not influence their neural activation (Figure 3).

Figure 3. Average odor association ratings for the 48 objects. Overall, the picture and sound representation of the same object have similar odor association. Critically, values span a large portion of the scale. Eight example objects are named below the graph. Error bars denote ±1 Standard Deviation (SD).

There was no significant parametric modulation by odor association in the PPC for either pictures (Figure 1C, $t(46) = 0.96$, $p = .34$, Cohen's $d = 0.14$) or sounds (Figure 1C, $t(46) = 0.75$, $p = .46$, Cohen's $d = 0.11$). Likewise, the modulation by odor association of the response to sounds in the APC was non-significant (Figure 1C, $t(46) = 0.65$, $p = .52$, Cohen's $d = .10$), as was the modulation of the (non-significant) response to pictures in the APC (Figure 1C, $t(46) = 1.94$, $p = .06$, Cohen's $d = 0.28$). Null results cannot be interpreted as absence of effects, so we performed additional Bayesian one-sample t-tests to evaluate the support for the null model (absence of effect). These indicated moderate evidence for the null effect for odor association modulation in the PPC, both in the case of visual stimulation ($B_{01} = 4.1$) and auditory stimulation ($B_{01} = 4.9$). Similarly, there was weak evidence for the null model of odor association modulation in the APC for pictures ($B_{01} = 1.1$) and moderate support for the null model for sounds ($B_{01} = 5.2$).

Although we did not detect any effect of odor association on the activations in piriform cortex when considering the parametric modulations, a subset of objects with strong odor associations could potentially have produced an odor association-dependent activity in a non-linear fashion. To further investigate this possibility, we performed two additional analyses based on new GLMs. In the first analysis, we split the pictures and sounds into two groups, based on each individual's odor association ratings, where one group comprised the lowest and one group the highest rated half of the stimuli. In the PPC ROI, we found that the visual and auditory activations persisted both for objects with weak and strong odor associations

(weak: pictures with odor association 11.8 ± 10.9: $t(40) = 4.19$, $p = .00015$, Cohen's $d = 0.65$; sounds with odor association 12.3 ± 11.7: $t(40) = 7.52$, $p = 3.6*10^{-9}$, Cohen's $d = 1.17$; strong: pictures with odor association 71.3 ± 16.1: $t(40) = 3.79$, $p = .00050$, Cohen's $d = 0.59$; sounds with odor association 68.8 ± 16.1: $t(40) = 7.68$, $p = 2.2*10^{-9}$, Cohen's $d = 1.20$). Further, a Bayesian paired t-test showed moderate evidence for the null effect of no difference between these activations for both pictures ($B_{01} = 5.26$) and sounds ($B_{01} = 5.89$).

In the APC ROI, the auditory activation also remained significant for both objects with weak and strong odor associations (weak: $t(40) = 6.48$, $p = 1.0*10^{-7}$, Cohen's $d = 1.01$; strong: $t(40) = 6.51$, $p = 9.2*10^{-8}$, Cohen's $d = 1.02$). Again, a Bayesian paired t-test showed moderate evidence for the null effect of no difference between these activations ($B_{01} = 5.61$). In relation to the non-significant response to sounds in the APC seen in the main analysis, this remained non-significant both for objects with weak and strong odor associations (weak: $t(40) = .42$, $p = 0.67$, Cohen's $d = 0.07$; strong: $t(40) = 1.43$, $p = 0.16$, Cohen's $d = 0.22$), and the Bayesian paired t-test showed moderate evidence for the null effect of no difference between these responses ($B_{01} = 4.53$).

In the second analysis, we treated each stimulus as a separate condition and assessed the respective PPC ROI activations. We then ranked the stimuli according to each individual's odor association ratings. These rankings did not significantly correlate with the activation for either pictures or sounds in the PPC (pictures: Spearman's $\rho = .19$, $p = .20$, sounds: Spearman's $\rho = -.039$, $p = .79$). In the APC, there was a significant correlation for pictures (Spearman's $\rho = .42$, $p = .0033$), but not for sounds (Spearman's $\rho = -.073$, $p = .62$).

Piriform cortex is not activated by nasal breathing but can be activated by sniffing alone, even in the absence of other sensory stimuli (Sobel et al., 1998). Although participants were instructed to breathe normally through their nose and were not asked to sniff when presented with a stimulus, a potential confound is that participants might have non-consciously sniffed when presented with a visual or auditory stimuli (Perl et al., 2019), independent of its odor association. As final control analysis, we therefore assessed whether piriform activation was associated with a sniff response synchronized to stimulus onset; in other words, whether the act of seeing or hearing a stimulus made participants initiate a sniff. To this end, we used the nasal respiratory data collected throughout the experiment and operationalized "a sniff" as a peak occurring at some point 0-1s after stimulus onset, a definition that bias our analyses

towards finding sniff synchronization. This operationalization was used because the presented stimuli were not synchronized with breathing or presented with an inhalation cue. We then assessed the probability, considering the total number of nasal breaths throughout the recording, of a sniff occurring 0-1s after presenting a stimulus. There was no significant increase in nasal inspiration time-locked to stimulus onset to either visual or auditory stimuli. Across all participants, the maximum probability of a peak occurring after a visual stimulus was .047 and occurring after a sound stimulus .050.

Visual cortex processes auditory unisensory stimuli

We found cross effects in the two ROIs covering visual brain regions: Sounds activated V1 (t(46) = 8.01, p = 2.8*10^{-10}, Cohen's d = 1.17) and deactivated LOC (t(46) = -6.44, p = 6.4*10^{-8}, Cohen's d =-0.94; Figure 4A-B). Naturally, the sounds also activated multiple known nodes of the auditory processing network in the temporal lobe and frontotemporal junction (Figure 5). These expected activations not only indicate that our stimulation and analysis were effective but also serve as a data sanity check. Accordingly, the activations were both significant in A1 (t(46) = 21.76, p = 7.4*10^{-26}, Cohen's d = 3.17) and hAC (t(46) = 18.97, p = 2.2*10^{-23}, Cohen's d = 2.77; Figure 4D-E). In A1, the response to sounds was positively modulated by odor association (t(46) = 2.01, p = .050, Cohen's d = 0.29), whereas the modulation of the response to pictures was non-significant (t(46) = -0.13, p = .90, Cohen's d = -0.02; Figure 4F). In hAC, none of the parametric modulations were significant (pictures: t(46) = -0.26, p = .79, Cohen's d = -0.04; sounds: t(46) = 0.36, p = 0.72, Cohen's d = 0.05).

Figure 4. Region of interest MR imaging results for visual and auditory brain regions. **A**) Location of visual ROIs. **B**) Both sounds and pictures activated primary visual cortex (V1). In lateral occipital complex (LOC), activity increased in response to pictures and decreased in response to sounds. **C**) In V1, the modulation by odor association was significantly above zero for pictures, but not for sounds. In LOC, there was no significant modulation for either pictures or sounds. **D**) Location of auditory ROIs. **E**) In primary auditory cortex (A1), the BOLD signal increased in response to sounds and decreased in response to pictures. Both pictures and sounds elicited significant activations in higher order auditory cortex (hAC). **F**) In A1, the modulation by rated odor association was significantly above zero for sounds, but not for pictures. Asterisks mark statistical significance and error bars depict ± 1 Standard Error of the Mean (SEM). Note the varying scale on the y axes.

Auditory cortex processes visual unisensory stimuli

Cross-modal effects were also noted in the auditory ROIs: Pictures elicited a decreased neural activity in primary auditory cortex (A1; $t(46)$ = -2.45, p = .018, Cohen's d = -0.36) and an increased activity in higher auditory cortices (hAC; $t(46)$ = 2.42, p = .019, Cohen's d =0.35; Figure 4E). Naturally, the visual stimulation also activated multiple regions in the occipital and parietal cortices that are all known nodes in the visual processing network (Figure 5), again indicating that our stimulation and analysis worked as intended. These activations were significant both in primary visual cortex (V1; $t(46)$ =9.79, p = 7.97*10^{-13}, Cohen's d = 1.43) and in the visual object-processing area LOC ($t(46)$ = 8.01, p = 2.83*10^{-10}, Cohen's d = 1.17; Figure 4B). In V1, the activation by pictures was positively modulated by odor association ($t(46)$ = 2.16, p = .036, Cohen's d = 0.32), whereas the activation by sounds was not ($t(46)$ = 1.53, p = .13, Cohen's d = 0.22; Figure 4C). In LOC, none of the parametric modulations were significant (pictures: $t(46)$ = -0.89, p = .38, Cohen's d = -0.13; sounds: ($t(46)$ = 0.68, p = .50, Cohen's d = 0.10).

Figure 5: Whole-brain fMRI results. *Upper panel*: Visual activation. The outlines of the visual ROIs are overlaid in cyan (V1) and blue (LOC). *Lower panel*: Auditory activation. The outlines of the auditory ROIs are overlaid in cyan (A1) and blue (hAC). Both the visual and the auditory activations are corrected for multiple comparisons using a peak-level threshold at p = .05 family-wise error (FWE) across the whole field of view.

Auditory and visual object information are transferred through mediator nodes to PPC

We demonstrated above that PPC processes unisensory visual and auditory objects. We next sought to understand which cerebral node might mediate the information to the PPC. Using a Floyd–Warshall algorithm of the dynamic functional connectivity between our ROIs, separately for each input stimulus. We found that both auditory and visual object areas are connected to PPC via known mediator nodes. For auditory stimuli, information about the auditory object is conveyed from the hAC via the planum polare (x ±45, y 9, z -9; p < .05, FDR corrected) and for visual objects, information is conveyed from the LOC to the PPC via the retrosplenial cortex (x ±9, y -45, z -9; p < .05, FDR corrected).

DISCUSSION

We here aimed to distinguish between two main candidate explanations of atypical unisensory activation of sensory cortices: activation by the atypical stimulus per se (true atypical unisensory processing), and activation due to association of the atypical stimulus to a typical one (cross-modal association). We first demonstrated that piriform (olfactory) cortex responded to both unimodal visual and unimodal auditory stimuli in form of pictures and sounds of object which were neither explicitly nor implicitly linked to odors during the experiment. Specifically, we found that the posterior piriform cortex (PPC), an area known to

process odor objects (Howard et al., 2009; Gottfried, 2010), responded to both pictures and sounds of objects. The anterior piriform cortex (APC), which is thought to code mainly for low-level odor properties (Gottfried, 2010), responded to sounds of objects. Thereafter, we demonstrated that the atypical unisensory activation in piriform cortex did not depend on either the odor association of the presented objects or a sniff response to their presentation. Together, these results provide strong support for true atypical unisensory processing in olfactory cortex.

Our results demonstrate that unisensory auditory and visual objects can activate piriform cortex, even in the absence of any odor stimulation or mention of odors. Critically, the observed cross-modal processing in piriform cortex was independent of the individual's level of odor association to the different objects (rated post fMRI acquisition), demonstrated by three independent analyses. First, there was a moderate support for the null model of an absence of modulation by odor association. Second, there was clear activation of piriform cortex even for objects that the participant associated only weakly with odors, which were also comparable to the effects for objects with strong odor associations. Third, we found no correlation between odor association and piriform activation. Our results extend prior findings of cross-modal processing in human piriform cortex (Gottfried et al., 2004; Karunanayaka et al., 2015; Schulze et al., 2017; Porada et al., 2019) in two important ways. We show both auditory and visual cross-modal activity in the piriform cortex whereas previous studies have only reported visual cross-modal activity. A recent study by Porada and colleagues (2019) included unisensory auditory stimulation but did not find auditory evoked activation of piriform cortex, possibly due to a lack of sufficient statistical power to detect small effects. The present study has considerably higher statistical power with three times as many participants and roughly five times as many trials per participant. Second, and more importantly, all previous studies associated the presented non-olfactory stimuli with olfactory stimuli before and/or during neuroimaging data collection. For example, in both Porada and colleagues (2019) and Gottfried and colleagues (2004), experiments were preceded by a session where the associations between the non-olfactory and olfactory stimuli were learned. The non-olfactory stimuli giving rise to piriform activity were thus explicitly associated with odors and it is likely that participants associated the stimulation to odors, even when no odor was simultaneously presented. We did not present or mention odors prior to or during fMRI

data acquisition, making it possible to distinguish between true atypical sensory processing and cross-modal association.

We have clearly demonstrated that the cross-modal processing of auditory and visual stimuli in piriform cortex is not dependent on how much the participant associates the presented object with an odor. That said, the cross-modal activity may still depend on the fact that our unimodal auditory and visual stimulation depicted objects. The activation was significant for both auditory and visual object stimuli in the PPC. Given that this region is commonly linked to olfactory object and category processing (Howard et al., 2009; Gottfried, 2010), it is plausible that the processing of non-olfactory input should be equally dependent on object or category specific content. This hypothesis is supported by our recent study demonstrating multisensory integration of object information in PPC (Porada et al., 2019). Specifically, olfactory, auditory, and visual object stimuli were presented uni-, bi-, and trimodally and the activity in PPC increased with the number of sensory modalities providing object information, but only when the object information was congruent (i.e., the same object presented) in the different sensory modalities. This indicates that specific object-related information, rather than low-level stimulus properties alone, affects the activity in PPC also in the case of non-olfactory stimuli. We propose that this non-olfactory object processing is what we observe in PPC in the current study. We furthermore speculate that the auditory activation of APC might be caused by the relatively high auditory activation of PPC as a reflection of the strong internal connectivity between these two regions (Carmichael et al., 1994). Based on the current data, we cannot conclusively determine whether the piriform activity evoked by pictures and sounds of objects indeed reflects processing of object information. Future studies could address this by either contrast the evoked activity from auditory and visual object and non-object stimuli with similar low-level perceptual properties or determine whether visual/auditory objects can be decoded based on piriform cortex activation patters.

An alternative interpretation of the current findings is that the cross-modal processing in piriform cortex is not related to the specific object information or low-level properties of the stimuli, but instead is an effect caused by increased attention or arousal evoked by the onset of the auditory and visual stimuli. Indeed, attention-dependent activity in piriform cortex has been demonstrated, but only in the APC (Zelano et al., 2005), a region where we in the current study find cross-modal activation only by auditory stimuli. It should also be noted that the

attention effect demonstrated by Zelano and colleagues (2005) was evident when attending to odors, whereas the participants in the current study were unaware that the study in fact assessed phenomena related to odors and were presumably not engaged in any conscious odor-searching behavior (the majority of our participants reported not having thought about odors during the experiment). Speaking against the notion that our obtained results are mediated by attentional processes is the fact that we did not find a stimuli-synchronized sniff response, a phenomenon argued to be associated with attentional allocation (Perl et al., 2019). Nonetheless, it is possible that our findings are generated by a general preparatory or prediction coding effect where non-olfactory sensory input prepares olfactory cortex for potential incoming olfactory input in response to new non-olfactory information (Ohla et al., 2017).

The functional connectivity analysis provides us with potential paths that the auditory and visual object information travels from object processing areas in the auditory (hAC) and visual cortex (LOC) to the olfactory object processing region (PPC). The auditory information seemingly connects in the planum polare, and the visual information in the retrosplenial cortex, before it reaches the PPC. The planum polare is a known hub in the auditory "what" pathway (Warren and Griffiths, 2003)(Ahveninen et al., 2006) whereas the retrosplenial cortex is associated with various functions, such as episodic memory, navigation, and planning(Vann et al., 2009). The latter functions are linked to olfaction. Morover, the retrosplenial cortex has been identified as the central node through which information flows between medial temporal lobe, including olfactory related entorhinal and perirhinal cortices, and other the brain networks (Kaboodvand et al., 2018). Additional to functional connections, retrosplenial is structurally connected to visual regions as well as multiple regions important for olfaction, such as the hippocampus, entorhinal cortex, and the orbitofrontal cortex (Vann et al., 2009). Given the finding of these two connection nodes, one can speculate whether the function of the visual/auditory activation of the piriform cortex is to subserve the olfactory system's task of assisting us to make rapid approach/avoidance decisions (Arshamian et al., 2017; Iravani et al., 2021).

Analogously to the crossmodal activity of the olfactory cortex, we found cross-modal effects also in the visual and auditory ROIs. Primary visual cortex (V1) was activated by the sounds of objects, and hAC was activated by the pictures. It is here interesting to note that the latter of

these cross-activations was apparent also in the aforementioned study by Porada and colleagues (2019). In primary auditory cortex (A1) and LOC, we instead found decreased BOLD signal in response to the atypical stimulation (pictures and sounds, respectively). Similar effects have previously been described in auditory and extrastriate visual areas (Laurienti et al., 2002). Moreover, there are several other accounts of cross-modal effects in the auditory and visual cortices on a finer spatial scale. For example, over 40% of cells in the cat visual parastriate cortex responds to unimodal localized auditory stimuli (Morrell, 1972), auditory stimulation influences the preferences of cells in the cat V1 area (Chanauria et al., 2019), and visual and auditory objects can be decoded from the their respective atypical sensory cortex in humans (Meyer et al., 2010; Hsieh et al., 2012; Vetter et al., 2014). Thus, previous work has demonstrated stimulus-specific activations in auditory cortex by unimodal visual stimulation, and stimulus-specific activation in visual cortex by unimodal auditory stimulation. We can here conceptually replicate the general finding of visual-auditory cross-activation of respective atypical sensory cortices.

It has previously been argued that the neocortex, including both higher-order processing regions and regions traditionally thought of as unisensory, is in essence multisensory (Ghazanfar and Schroeder, 2006). We can here provide support to this notion and extend this concept of a multisensory neocortex by demonstrating that the olfactory cortex, which is evolutionarily the oldest sensory cortex and not part of the neocortex which evolved from the olfactory-dominated cerebral hemispheres (Northcutt and Kaas, 1995), also appear to respond in a multisensory fashion: It appears to process non-olfactory input independently of odor association. Moreover, it has been proposed that the brain is organized according to tasks rather than sensory input, with the possible exception of primary sensory cortices (Amedi et al., 2017). Studies on congenitally blind individuals have shown that they demonstrate the same cortical specialization in 'visual' cortex for non-visual stimuli as is normally seen in visual processing, such as intact ventral/dorsal pathways (Cecchetti et al., 2016). Assuming that the cross-modal activity in the PPC that we have demonstrated reflects processing of object information in the atypical (non-olfactory) stimuli, our findings are consistent with a task-oriented brain organization in which the role of PPC is object processing, with a preference for olfactory input. Another interpretation of our results is that the PPC is an olfactory region that can be activated by non-olfactory input in a preparatory way, possibly

driven by a lifetime of multisensory input. Regardless of which theory that will find future support, our findings support the notion of a multimodal cortex and highlights that this extends to the olfactory cortex.

In summary, we here conclusively demonstrate true unisensory processing of sounds and pictures of objects in olfactory cortex that is independent of odor associations. These results provide additional support against the notion of piriform cortex as a true unisensory olfactory region and for the argument that sensory processing is best characterized as relying on multisensory networks.

Acknowledgments

Funding provided by the Knut and Alice Wallenberg Foundation (KAW 2018.0152) and the Swedish Research Council (2021-06527), awarded to JNL. Data acquisition was supported by a grant to the Stockholm University Brain Imaging Centre (SU FV-5.1.2-1035-15).

References

Ahveninen J, Jääskeläinen IP, Raij T, Bonmassar G, Devore S, Hämäläinen M, Levänen S, Lin F-H, Sams M, Shinn-Cunningham BG, Witzel T, Belliveau JW (2006) Task-modulated "what" and "where" pathways in human auditory cortex. Proc Natl Acad Sci USA 103:14608–14613.

Amedi A, Hofstetter S, Maidenbaum S, Heimler B (2017) Task selectivity as a comprehensive principle for brain organization. Trends Cogn Sci (Regul Ed) 21:307–310.

Arshamian A, Laska M, Gordon AR, Norberg M, Lahger C, Porada DK, Jelvez Serra N, Johansson E, Schaefer M, Amundin M, Melin H, Olsson A, Olsson MJ, Stensmyr M, Lundström JN (2017) A mammalian blood odor component serves as an approach-avoidance cue across phylum border - from flies to humans. Sci Rep 7:13635.

Budinger E, Heil P, Scheich H (2000) Functional organization of auditory cortex in the Mongolian gerbil (Meriones unguiculatus). III. Anatomical subdivisions and corticocortical connections. Eur J Neurosci 12:2425–2451.

Budinger E, Scheich H (2009) Anatomical connections suitable for the direct processing of neuronal information of different modalities via the rodent primary auditory cortex. Hear Res 258:16–27.

Cain WS (1979) To know with the nose: keys to odor identification. Science 203:467–470.

Carmichael ST, Clugnet MC, Price JL (1994) Central olfactory connections in the macaque monkey. J Comp Neurol 346:403–434.

Cecchetti L, Kupers R, Ptito M, Pietrini P, Ricciardi E (2016) Are Supramodality and Cross-Modal Plasticity the Yin and Yang of Brain Development? From Blindness to Rehabilitation. Front Syst Neurosci 10:89.

Chanauria N, Bharmauria V, Bachatene L, Cattan S, Rouat J, Molotchnikoff S (2019) Sound Induces Change in Orientation Preference of V1 Neurons: Audio-Visual Cross-Influence. Neuroscience 404:48–61.

Cohen Y, Courtiol E, Sullivan RM, Wilson DA (2017) Cortical processing of odorants. In: Oxford research encyclopedia of neuroscience. Oxford University Press.

Cooper HM, Parvopassu F, Herbin M, Magnin M (1994) Neuroanatomical pathways linking vision and olfaction in mammals. Psychoneuroendocrinology 19:623–639.

Datiche F, Litaudon P, Cattarelli M (1996) Intrinsic association fiber system of the piriform cortex: a quantitative study based on a cholera toxin B subunit tracing in the rat. J Comp Neurol 376:265–277.

Fransson P, Aden U, Blennow M, Lagercrantz H (2011) The functional architecture of the infant brain as revealed by resting-state fMRI. Cereb Cortex 21:145–154.

Ghazanfar AA, Schroeder CE (2006) Is neocortex essentially multisensory? Trends Cogn Sci (Regul Ed) 10:278–285.

Gottfried JA, Smith APR, Rugg MD, Dolan RJ (2004) Remembrance of odors past: human olfactory cortex in cross-modal recognition memory. Neuron 42:687–695.

Gottfried JA (2010) Central mechanisms of odour object perception. Nat Rev Neurosci 11:628–641.

Howard JD, Plailly J, Grueschow M, Haynes J-D, Gottfried JA (2009) Odor quality coding and categorization in human posterior piriform cortex. Nat Neurosci 12:932–938.

Hsieh PJ, Colas JT, Kanwisher N (2012) Spatial pattern of BOLD fMRI activation reveals cross-modal information in auditory cortex. J Neurophysiol 107:3428–3432.

Iravani B, Schaefer M, Wilson DA, Arshamian A, Lundström JN (2021) The human olfactory bulb processes odor valence representation and cues motor avoidance behavior. Proc Natl Acad Sci USA 118.

Johnson DM, Illig KR, Behan M, Haberly LB (2000) New features of connectivity in piriform cortex visualized by intracellular injection of pyramidal cells suggest that "primary" olfactory cortex functions like "association" cortex in other sensory systems. J Neurosci 20:6974–6982.

Kaboodvand N, Bäckman L, Nyberg L, Salami A (2018) The retrosplenial cortex: A memory gateway between the cortical default mode network and the medial temporal lobe. Hum Brain Mapp 39:2020–2034.

Karunanayaka PR, Wilson DA, Vasavada M, Wang J, Martinez B, Tobia MJ, Kong L, Eslinger P, Yang QX (2015) Rapidly acquired multisensory association in the olfactory cortex. Brain Behav 5:e00390.

Laurienti PJ, Burdette JH, Wallace MT, Yen Y-F, Field AS, Stein BE (2002) Deactivation of sensory-specific cortex by cross-modal stimuli. J Cogn Neurosci 14:420–429.

Lundström JN, Boesveldt S, Albrecht J (2011) Central processing of the chemical senses: an overview. ACS Chem Neurosci 2:5–16.

Lundström JN, Regenbogen C, Ohla K, Seubert J (2019) Prefrontal control over occipital responses to crossmodal overlap varies across the congruency spectrum. Cereb Cortex 29:3023–3033.

Maier JX, Wachowiak M, Katz DB (2012) Chemosensory convergence on primary olfactory cortex. J Neurosci 32:17037–17047.

Mainland JD, Lundström JN, Reisert J, Lowe G (2014) From molecule to mind: an integrative perspective on odor intensity. Trends Neurosci 37:443–454.

Meyer K, Kaplan JT, Essex R, Webber C, Damasio H, Damasio A (2010) Predicting visual stimuli on the basis of activity in auditory cortices. Nat Neurosci 13:667–668.

Morrell F (1972) Visual system's view of acoustic space. Nature 238:44–46.

Northcutt RG, Kaas JH (1995) The emergence and evolution of mammalian neocortex. Trends Neurosci 18:373–379.

Ohla K, Höchenberger R, Freiherr J, Lundström JN (2017) Superadditive and Subadditive Neural Processing of Dynamic Auditory-Visual Objects in the Presence of Congruent Odors. Chem Senses 43:35–44.

Olofsson JK, Zhou G, East BS, Zelano C, Wilson DA (2019) Odor Identification in Rats: Behavioral and Electrophysiological Evidence of Learned Olfactory-Auditory Associations. eNeuro 6.

Perl O, Ravia A, Rubinson M, Eisen A, Soroka T, Mor N, Secundo L, Sobel N (2019) Human non-olfactory cognition phase-locked with inhalation. Nat Hum Behav 3:501–512.

Porada DK, Regenbogen C, Seubert J, Freiherr J, Lundström JN (2019) Multisensory enhancement of odor object processing in primary olfactory cortex. Neuroscience 418:254–265.

Regenbogen C, Johansson E, Andersson P, Olsson MJ, Lundström JN (2016) Bayesian-based integration of multisensory naturalistic perithreshold stimuli. Neuropsychologia 88:123–130.

Schulze P, Bestgen A-K, Lech RK, Kuchinke L, Suchan B (2017) Preprocessing of emotional visual information in the human piriform cortex. Sci Rep 7:9191.

Seydell-Greenwald A, Wang X, Newport E, Bi Y, Striem-Amit E (2020) Spoken language comprehension activates the primary visual cortex. BioRxiv.

Sobel N, Prabhakaran V, Desmond JE, Glover GH, Goode RL, Sullivan EV, Gabrieli JD (1998) Sniffing and smelling: separate subsystems in the human olfactory cortex. Nature 392:282–286.

Vann SD, Aggleton JP, Maguire EA (2009) What does the retrosplenial cortex do? Nat Rev Neurosci 10:792–802.

Vasconcelos N, Pantoja J, Belchior H, Caixeta FV, Faber J, Freire MAM, Cota VR, Anibal de Macedo E, Laplagne DA, Gomes HM, Ribeiro S (2011) Cross-modal responses in the primary visual cortex encode complex objects and correlate with tactile discrimination. Proc Natl Acad Sci USA 108:15408–15413.

Veldhuizen MG, Small DM (2011) Modality-specific neural effects of selective attention to taste and odor. Chem Senses 36:747–760.

Vetter P, Smith FW, Muckli L (2014) Decoding sound and imagery content in early visual cortex. Curr Biol 24:1256–1262.

Warren JD, Griffiths TD (2003) Distinct mechanisms for processing spatial sequences and pitch sequences in the human auditory brain. J Neurosci 23:5799–5804.

Yang AI, Dikecligil GN, Jiang H, Das SR, Stein JM, Schuele SU, Rosenow JM, Davis KA, Lucas TH, Gottfried JA (2021) The what and when of olfactory working memory in humans. Curr Biol 31:4499-4511.e8.

Zald DH, Pardo JV (2000) Functional neuroimaging of the olfactory system in humans. Int J Psychophysiol 36:165–181.

Zelano C, Bensafi M, Porter J, Mainland J, Johnson B, Bremner E, Telles C, Khan R, Sobel N (2005) Attentional modulation in human primary olfactory cortex. Nat Neurosci 8:114–120.

Zhou G, Lane G, Cooper SL, Kahnt T, Zelano C (2019) Characterizing functional pathways of the human olfactory system. eLife 8.