1

Alterations in cortical excitability during pain: A combined TMS-EEG Study

Nahian S Chowdhury[1,2], Alan KI Chiang[1,2], Samantha K Millard[1,2], Patrick Skippen[1], Wei-Ju

Chang[1,3], David A Seminowicz[1, 6], Siobhan M Schabrun[1,7]

Affiliations

[1]Center for Pain IMPACT, Neuroscience Research Australia, Sydney, New South Wales,

Australia, [2]University of New South Wales, Sydney, New South Wales, Australia, [3]School of

Health Sciences, College of Health, Medicine and Wellbeing, The University of Newcastle,

Callaghan, New South Wales, Australia,[6]Department of Medical Biophysics, Schulich School

of Medicine & Dentistry, University of Western Ontario, London, Canada, [7]The Gray Centre

for Mobility and Activity, University of Western Ontario, London, Canada.

Corresponding author

Nahian S Chowdhury

Email: n.chowdhury@neura.edu.au

Neuroscience Research Australia, 139 Barker Street Randwick 2031 NSW

Disclosures

The authors have no conflicts of interest to declare.

Abstract

Transcranial magnetic stimulation (TMS) has been used to examine inhibitory and facilitatory circuits during experimental pain and in chronic pain populations. However, current applications of TMS to pain have been restricted to measurements of motor evoked potentials (MEPs) from peripheral muscles. Here, TMS was combined with electroencephalography (EEG) to determine whether experimental pain could induce alterations in cortical inhibitory/facilitatory activity observed in TMS-evoked potentials (TEPs). In Experiment 1 (n = 29), multiple sustained thermal stimuli were administered to the forearm, with the first, second and third block of thermal stimuli consisting of warm but non-painful (pre-pain block), painful (pain block) and warm but non-painful (post-pain block) temperatures respectively. During each stimulus, TMS pulses were delivered while EEG (64 channels) was simultaneously recorded. Verbal pain ratings were collected between TMS pulses. Relative to pre-pain warm stimuli, painful stimuli led to an increase in the amplitude of the frontocentral negative peak ~45ms post-TMS (N45), with a larger increase associated with higher pain ratings. Experiments 2 and 3 (n = 10 in each) showed that the increase in the N45 in response to pain was not due to changes in sensory potentials associated with TMS, or a result of stronger reafferent muscle feedback during pain. This is the first study to use combined TMS-EEG to examine alterations in cortical excitability in response to pain. These results suggest that the N45 TEP peak, which indexes GABAergic neurotransmission, is implicated in pain perception and is a potential marker of individual differences in pain sensitivity.

Introduction

Pain is a complex subjective experience, and understanding how pain is processed

remains a challenge (Apkarian, 2021). Several neuroimaging techniques have been applied

to disentangle these complexities: functional magnetic resonance imaging has assisted in

identifying brain structures implicated in pain processing (Reddan & Wager, 2018), while

electroencephalography (EEG) has contributed to our understanding of the temporal sequence

of pain processing (Ploner & May, 2018). Another useful technique that has been used to

examine the role of inhibitory and facilitatory neural circuits in pain has been transcranial

magnetic stimulation (TMS) delivered to the brain (Chang et al., 2018; Schabrun & Hodges,

2012). However, current applications of TMS to pain have involved recording the output of

TMS from a muscle, a signal that could be influenced by many intermediate (subcortical,

spinal, peripheral) factors, and which restricts investigations to the motor system only. Here,

we use a combined TMS-EEG measure to record output of TMS directly from the cortex and

from multiple brain regions, in pain-free and tonic pain conditions.

When TMS is delivered over the primary motor cortex (M1), a magnetic pulse

induces an electrical current in underlying cortical tissue that, if the intensity is sufficient,

activates corticomotor pathways, inducing a motor evoked potential (MEP) in a target

muscle. The magnitude of the MEP serves as an index of corticomotor excitability. Past

systematic reviews on studies measuring MEPs during acute experimental pain (Bank, Peper,

Marinus, Beek, & Van Hilten, 2013; Burns, Chipchase, & Schabrun, 2016; Chowdhury et al.,

2022; Rohel et al., 2021) have shown a reduction in MEP amplitude during pain and after

pain resolution, with stronger reductions in MEP amplitude associated with lower acute pain

severity (Chowdhury, et al., 2022). It has been hypothesised that this reduction in MEP

amplitude is an adaptive mechanism that restricts movement in the pain-afflicted area, to

protect the area from further pain and injury (Hodges & Tucker, 2011).

While previous findings show promise for the use of TMS to discover and validate

potential biomarkers for pain, limitations exist when using TMS to measure MEPs. First,

MEP responses to TMS reflect the net sum of cortical, spinal, and peripheral activity within

the corticomotor pathway. This makes it unclear as to whether pain processes occur at the

cortical, spinal or peripheral level. Further, measurement of MEPs restricts investigations to

M1. One way of overcoming these limitations is by combining TMS and EEG to measure

TMS-evoked potentials (TEPs). TEPs index cortical excitability *directly* from the cortex (i.e.

without influence of subcortical, spinal and peripheral processes), as well as from regions

outside M1 (Farzan et al., 2016). TEPs also provide an index of the activity of specific

neurotransmitter circuits within the cortex. For example, TEP peaks that occur at ~45ms and

100ms post-stimulation are linked to $GABA_A$ and $GABA_B$ neurotransmission respectively

(Premoli et al., 2014), while the TEP peak ~60ms post-stimulation is linked to glutamatergic

neurotransmission (Belardinelli et al., 2021). Overall, TEPs provides additional spatial and

temporal information about cortical activity over MEPs, making it ideal for understanding the

brain mechanisms involved in pain perception.

TEPs have already shown potential to serve as a biomarker for the development and

prognosis of various neurological and psychiatric conditions (for reviews see (Kallioniemi &

Daskalakis, 2022; Tremblay et al., 2019)). However, the applicability of TEPs to pain

research is yet to be established. While GABAergic processes indexed by TEPs have been

hypothesised to be involved in pain (Barr, Farzan, Davis, Fitzgerald, & Daskalakis, 2013),

direct evidence is scarce. Two studies (Che et al., 2019; Ye, Wang, & Che, 2022) examined

whether the potential analgesic effects of repetitive TMS (rTMS) over the dorsal prefrontal

cortex are associated with plasticity in TEPs. These studies separately measured TEPs and

ratings to painful stimuli, before and after rTMS, with one finding that increases in pain

thresholds following rTMS were associated with changes in TEPs that index GABAergic

processes (Ye et al., 2022). While these studies assist us in understanding whether TEPs

might mediate rTMS-induced pain reductions, no study has investigated whether TEPs are

altered in direct response to pain.

The aim of the present study was to use TMS-EEG to determine whether acute

experimental pain induces alterations in cortical inhibitory and facilitatory peaks observed

using TEPs. We used a tonic heat pain paradigm (Furman et al., 2020; Granot, Granovsky,

Sprecher, Nir, & Yarnitsky, 2006), in which multiple thermal stimuli were applied over the

right extensor carpi radialis brevis (ECRB) muscle via a thermode. For each thermal

stimulus, the temperature increased from a neutral baseline of 32°C to either a warm non-

painful or a painful (46°C) temperature, with this temperature maintained for 40 seconds.

During this time, TMS was administered to the left M1 with concurrent EEG to obtain TEPs

from 63 scalp channels, and MEPs from the ECRB muscle (see Figure 1). Verbal pain ratings

were obtained between pulses. It was hypothesized that TEP peaks that index GABAergic

processes, including the peaks at ~45 and 100ms after TMS, would increase in response to

painful stimuli relative to warm non-painful stimuli.

<div align="center">**Results**</div>

Experiment 1 – Does acute pain alter cortical excitability?

In Experiment 1 (n = 29), we determined whether painful thermal stimuli induced

alterations in TEP peaks relative to a non-painful baseline. The protocol (Figure 2) consisted

of three blocks of stimuli, in chronological order: pre-pain, pain, and post-pain blocks. The

pre-pain and post-pain blocks each consisted of six 40s thermal stimuli (20s interstimulus

interval) delivered at a non-painful temperature (calibrated to each participant's warmth

detection threshold), while the pain block consisted of six 40s thermal stimuli delivered at

46°C. The pre-pain/pain/post-pain design has been commonly used in the TMS-MEP pain

literature, as many studies have demonstrated strong changes in corticomotor excitability that

persist beyond the painful period. Indeed, in a systematic review, we showed effect sizes of

0.55-0.9 for MEP reductions 0-30 minutes after pain had resolved (Chowdhury et al., 2022).

As such, if we had used an alternative design with blocks of warm stimuli intermixed with

blocks of painful stimuli, the warm stimuli blocks would not serve as a valid non-painful

baseline. Based on a previous study (Dubé & Mercier, 2011) which also used sequences of

painful (50°C) and warm (36°C) thermal stimuli, we did not anticipate that the stimulus in the

pain block would entrain pain in the post-pain block.

Prior to the test blocks, we measured warmth, cool, and pain detection thresholds to

ascertain whether: a) participants could perceive increases or decreases in the thermode

temperature relative to a neutral baseline of 32°C, and b) the pain detection threshold was

below 46°C. All participants were able to detect increases or decreases in temperature from

baseline. The mean ($\pm$ SD) cool and warmth detection threshold was 28.6 $\pm$ 1.9°C and 35.1 $\pm$

1.5°C respectively. All participants reported a heat pain threshold that was above their

warmth detection threshold and below the test temperature of 46°C. The mean heat pain

threshold was 41.2 $\pm$ 2.8°C.

All participants reported 0/10 pain during the pre-pain and post-pain blocks, and pain

ratings varying between 1-10 during the pain block. Figure 3 shows the mean pain ratings for

the ten pain measurements of each of the six painful stimuli delivered during the pain block

(~4 seconds in between pain measurements). A 6 (stimulus number: 1-6) x 10 (timepoint:1-

10) Bayesian repeated measures ANOVA revealed anecdotal evidence (i.e., no conclusive

evidence) of a difference in pain between six thermal stimuli ($BF_{10} = 2.86$). However, there

was very strong evidence for a difference in pain ratings between the ten timepoints ($BF_{10} =$

6.1^{30}). There was also strong evidence of an interaction between stimulus number and

timepoint, suggesting the time course of pain across the 40 seconds thermal stimulus differed

across the six thermal stimuli of the pain block ($BF_{10} = 19.6$). Overall, although there was no

conclusive evidence for pain differing *between* successive stimuli, there was evidence that

pain fluctuated *during* each 40 second stimulus.

156   The mean resting motor threshold (RMT) and test intensity of TMS was (mean $\pm$ SD)

$70.7 \pm 8.5\%$ and $77.7 \pm 9.2\%$ of maximum stimulus output respectively. We note that the

relatively high RMTs are likely due to aspects of the experimental setup that increased the

distance between the TMS coil and the scalp, including the layer of foam placed over the coil,

the EEG cap and relatively thick electrodes (6mm). Three participants were excluded from

the MEP analysis due to EMG software failure – these participants were still included in the

TEP analysis. A Bayesian repeated-measures ANOVA was run to compare MEP amplitudes

between pre-pain, pain and post-pain conditions. There was anecdotal evidence of a

difference in MEP amplitude between blocks ($BF_{10} = 1.02$) (Figure 4A). A Bayesian

correlation test was also run to determine whether the mean pain rating (across blocks and

timepoints) was associated with the change in MEP amplitude during pain as a proportion of

pre-pain. These MEP change values were log-transformed as they were not normally

distributed according to a Shapiro-Wilk Test ($W = .58$, $p < .001$). There was strong evidence

for a positive relationship ($r_{26} = 0.54$, $BF_{10} = 11.17$) (Figure 4B). such that participants who

showed a larger reduction in MEP amplitude during pain reported lower pain ratings.

171   One participant was excluded from the TEP analysis due to failure to save the

recording during the experiment, though this participant was still included in the MEP

analysis. One participant had missing post-pain data as the TMS coil had overheated during

this portion of the experiment – their data were still included for the pre-pain vs. pain

comparison. Figure 5 shows the grand average TEPs for all 63 channels, across pre-pain, pain

and post-pain conditions, as well as the scalp topographies at timepoints where TEP peaks are

typically observed – N15, P30, N45, P60, N100 and P180ms (Farzan et al., 2016). Source

reconstruction using a co-registered template brain model was also conducted to characterise

source activity at each timepoint (Figure 5). For the N15 and P30 peaks, there was higher

current density in the left motor areas, consistent with previous studies suggesting that TMS

evoked activity at 15 and 30ms after the TMS pulse reflect early excitation of motor areas

ipsilateral to the stimulated region (Farzan & Bortoletto, 2022). For the N45 and P60 peaks,

there was higher current density in the left motor and somatosensory areas at 45 and 60ms

after the TMS pulse, consistent with previous studies showing a sensorimotor origin a these

timepoints (Ahn & Fröhlich, 2021), however higher current density was also present in the

left parietal and right sensorimotor areas. For the N100 and P180 peaks, there was higher

current density in the central regions, mostly contralateral to the stimulated cortex. Overall,

we found consistencies in the source localization with previous studies, including a

sensorimotor origin of early peaks from 15-60ms. However, we did not directly compare

source activity between conditions due to the inaccuracies involved in source estimation in

the absence of co-registered magnetic resonance imaging (MRI) scans (Brodbeck et al., 2011;

Michel & Brunet, 2019) and EEG electrode location digitization (Shirazi & Huang, 2019).

Comparisons of TEP amplitude between conditions were based on the electrode-space

data. However, as this is the first study investigating the effects of experimental pain on TEPs

amplitude, there were no *a priori* regions or timepoints of interest to compare between

conditions. A statistically robust starting point in these situations is to use a cluster-based

permutation analysis (Frömer, Maier, & Abdel Rahman, 2018). This analysis was used to

compare amplitudes between pre-pain and pain, and pre-pain and post-pain, at each timepoint

and for each electrode. We found that during pain relative to pre-pain, there was a

significantly larger negative amplitude ($p = .021$) at frontocentral electrodes and a

significantly larger positive amplitude at parietal-occipital electrodes ($p = .028$), specifically

between 43-90ms after the TMS pulse. No significant differences in TEP amplitude were

found when comparing the pre-pain and post-pain conditions, and pain and post-pain

conditions. As such, the subsequent TEP peak analyses were focused on the pre-pain vs. pain

comparison, while the pre-pain vs. post-pain and pain vs. postpain comparisons are presented

in the supplementary material.

Figure 6A shows the grand average TEP waveform at the frontocentral electrodes

('AF3','AFz','AF4','F1','Fz','F2','F4','FC2','FC4') identified from the cluster analysis for the

pre-pain vs. pain conditions (Supplementary Figure 1 shows the pre-pain vs. post-pain

comparison). Two peaks at ~45 and 85ms after the TMS pulse are visible in the time window

where the significant cluster was detected. Given the approximate timing, these peaks are

likely to be the N45 peak and an early N100 peak. The amplitude of these peaks was

identified for each participant using the TESA peak function (Rogasch et al., 2017) with

defined time windows of 40-70 and 75-95ms for the first and second peak respectively.

These time windows were chosen to account for variation between participants in the latency

of the first and second peak. Bayesian paired-sample t-tests showed very strong evidence that

the first peak at ~45ms (BF_{10} = 57.21) and moderate evidence that the second peak at ~85ms

(BF_{10} = 6.77) had larger amplitude during the pain block compared to the pre-pain block.

Figure 6B shows the individual level relationship between the mean pain rating, and the

difference in N45 and N100 amplitude between pain and pre-pain. There was strong evidence

that participants who reported higher pain ratings also showed a larger increase in N45 peak

amplitude during the pain block (r_{26} = 0.52, BF_{10} = 10.64). There was anecdotal evidence for

no association between pain ratings and changes in the N100 peak amplitude during pain (r_{26}

= 0.24, BF_{10} = 0.48).

Figure 6C shows the mean TEP waveform of the parietal-occipital electrodes

('P1','PO3','O1','CPz','Pz','Pz','Oz','CP2','P2', 'PO4','O2','CP4','P4') identified from the cluster

analysis for the pre-pain vs. pain conditions (Supplementary Figure 1 shows the pre-pain vs.

post-pain comparison). One peak at ~50ms is visible in the time window where the

significant cluster was detected. The approximate timing of this peak is consistent with the

commonly identified P60. The amplitude of this peak was identified for each participant with

a defined time window of 35 to 65ms. This time window was chosen to account for variation

between participants in the latency of the peak. There was moderate evidence that the peak at

~50ms was stronger during the pain block compared to pre-pain block (BF_{10} = 5.56). Figure

6D shows the individual level relationship between the mean pain rating and the difference in

the P60 amplitude between pain and pre-pain. There was anecdotal evidence in favour of no

relationship between pain ratings and changes in P60 amplitude during the pain block (r_{26} =

0.21, BF_{10} = 0.407). There was no conclusive evidence of any relationship between

alterations in MEP amplitude during pain, and alterations in N100, N45 and P60 amplitude

during pain (see supplementary material).

Experiment 2 – Does acute pain alter cortical excitability or sensory potentials?

Several studies have shown that a significant portion of TEPs do not reflect the direct

cortical response to TMS, but rather auditory potentials elicited by the "clicking" sound from

the TMS coil, and somatosensory potentials elicited by the "flicking" sensation on the skin of

the scalp (Biabani, Fornito, Mutanen, Morrow, & Rogasch, 2019; Chowdhury, et al., 2022;

Conde et al., 2019; Rocchi et al., 2021). Indeed, the signal at ~100ms post-TMS from

Experiment 1 may reflect an auditory N100 response. As it is extremely challenging to isolate

and filter these auditory and somatosensory evoked potentials using pre-processing pipelines,

masking methods have been used to suppress these sensory inputs, (Ilmoniemi & Kičić,

2010; Massimini et al., 2005). However recent studies have shown that even when these

methods are used, sensory contamination of TEPs is still present, as shown by commonalities

in the signal between active and sensory sham conditions that mimic the

auditory/somatosensory aspects of real TMS (Biabani et al., 2019; Conde et al., 2019; Rocchi

et al., 2021). This has led many leading authors (Biabani et al., 2019; Conde et al., 2019) to

recommend the use of sham conditions to control for sensory contamination. To separate the

direct cortical response to TMS from sensory evoked activity, Experiment 2 (n = 10) included

a sham TMS condition that mimicked the auditory/somatosensory aspects of active TMS to

determine whether any alterations in the TEP peaks in response to pain were due to changes

in sensory evoked activity associated with TMS, as opposed to changes in cortical

excitability. A similar design was used to Experiment 1, with the inclusion of a sham TMS

condition within the pre-pain and pain blocks, and exclusion of the post-pain block, since the

aim was to identify the source of the pain effect from Experiment 1. The sham TMS

condition was similar to a recent study (Gordon et al., 2021), involving the delivery of the

TMS coil rotated 90 degrees to the scalp to simulate the auditory component associated with

real TMS, and concurrent electrical stimulation beneath a sham coil to simulate the

somatosensory component associated with real TMS (see Figure 7A).

Figure 7B shows the grand average TEPs, scalp topographies and estimated source

activity for active and sham TMS, across pre-pain and pain conditions. Figure 8 shows the

mean TEP waveform for the frontocentral and parietal-occipital clusters identified from

Experiment 1, across active and sham conditions. There was moderate evidence that the

amplitude of the N45 peak was increased during pain vs. pre-pain blocks for active TMS

($BF_{10} = 3.26$), and moderate evidence for no difference between pain and pre-pain blocks for

sham TMS ($BF_{10} = 0.309$). When comparing pain and pre-pain blocks, there was,

respectively, moderate and anecdotal evidence for no alterations in the frontocentral N100 for

active ($BF_{10} = 0.31$) and sham TMS ($BF_{10} = 0.42$). There was anecdotal evidence for no

alteration in the parietal occipital P60 for both active ($BF_{10} = 0.786$) and sham TMS ($BF_{10} =$

0.42). Overall, the results showed that the N45 peak was altered in response to pain for active

but not sham TMS, suggesting the experience of pain led to an alteration in the excitability of

the cortex, and not the auditory/somatosensory aspects of TMS.

Experiment 3 – Does acute pain alter cortical excitability or reafferent muscle activity?

Previous studies have shown that a significant portion of the TEP peaks at 45 and 60ms post-TMS reflect reafferent feedback from the muscle twitch in response to suprathreshold TMS applied over M1. This comes from MRI-informed EEG studies showing source localization of the N45 and P60 peaks to the somatosensory areas, as well as correlations between MEP amplitude and N45/P60 amplitude (Ahn & Fröhlich, 2021; Petrichella, Johnson, & He, 2017). Indeed, Experiments 1 and 2 also showed localization of the N45 and P60 to sensorimotor areas. As such, Experiment 3 recruited a further ten participants to determine whether the pain-induced increase in the N45 peak was due to stronger reafferent feedback from muscle twitches. A design similar to Experiment 1 was used, with the inclusion of a subthreshold TMS condition (90% RMT) within the pre-pain and pain blocks.

Figure 9 shows the grand average TEPs, scalp topographies and estimated source activity for supra- and subthreshold TMS, across pre-pain and pain blocks. Figure 10 shows the mean TEP waveform for the frontocentral and parietal-occipital clusters identified from Experiment 1, across suprathreshold and subthreshold TMS conditions. When comparing the pain with pre-pain blocks, there was moderate evidence that the frontocentral N45 was increased during subthreshold TMS ($BF_{10} = 3.05$) and suprathreshold TMS ($BF_{10} = 3.01$). When comparing pain with pre-pain blocks, there was anecdotal evidence for no alterations in the frontocentral N100 peak during suprathreshold TMS ($BF_{10} = 0.42$) and subthreshold TMS ($BF_{10} = 0.36$). When comparing pain with pre-pain blocks, there was anecdotal evidence for an increase in the parietal occipital P60 peak during suprathreshold TMS ($BF_{10} = 2.71$) and anecdotal evidence for no alteration in the P60 peak during subthreshold TMS ($BF_{10} = 0.72$). Overall, there was evidence that that the N45 peak was altered in response to

both supra- and subthreshold TMS, suggesting the pain-induced increase in the N45 peak was

not a result of stronger reafferent feedback from the muscle twitches.

Discussion

The present study determined whether acute experimental pain induces alterations in

cortical inhibitory and/or facilitatory activity observed in TMS-evoked potentials. Across

three experiments, there was Bayesian evidence (varying between moderate to very strong)

for an increase in the amplitude of the N45 peak during painful stimuli compared to a non-

painful baseline. Experiment 1 showed very strong evidence that a larger increase in the N45

peak in response to pain was correlated with higher pain ratings. Experiment 2 showed that

the increase in the N45 peak during pain was not a result of alterations in sensory potentials

associated with the TMS pulses, but rather, changes in cortical excitability. Experiment 3

showed that the increase in the N45 peak was not a result of stronger reafferent feedback

from muscle twitches evoked by TMS during painful stimuli. While Experiment 1 showed

moderate evidence for an increase in N100 and P60 peaks during pain relative to pre-pain

baseline, this was not replicated in the follow-up experiments. Experiment 1 showed

anecdotal evidence for group-level alterations in MEP amplitude during pain, however there

was very strong evidence that a larger reduction in MEP amplitude during pain was

correlated with lower pain ratings.

Increased GABAergic activity during tonic pain

This study is the first to use TMS-EEG methodology to examine the direct cortical

response to acute pain, extending previous studies that have used TMS to measure MEPs in

response to pain (Chowdhury, et al., 2022). The key finding was an increase in the amplitude

of the N45 peak in response to pain. This result was replicated across three experiments,

providing robust evidence for the effect. Furthermore, we accounted for major confounds that

have caused significant data interpretation issues in the TMS-EEG literature in recent years,

namely the contamination of TEPs by sensory potentials associated with TMS pulses

(Biabani et al., 2019; Chowdhury, Rogasch, et al., 2022) and the presence of reafferent

feedback from muscle twitches (Ahn & Fröhlich, 2021).

The finding of a reliable increase in the amplitude of the N45 peak during pain

suggests a role for GABA$_A$ neurotransmission in pain processing, as previous work has

shown that the amplitude of the N45 peak is increased in response to GABA$_A$ agonists

(Premoli et al., 2014). Our source reconstruction results suggest that around this timepoint,

the current density was stronger in the sensorimotor area, consistent with the idea that the

N45 peak reflects GABAergic activity within the sensorimotor cortex (Farzan & Bortoletto,

2022). While it has been shown that reafferent muscle activity also contributes to the N45

peak (Ahn & Fröhlich, 2021), Experiment 3 showed that pain increased the amplitude of the

N45 peak even during subthreshold TMS. Taken together, these findings suggest that the

increased amplitude of the N45 peak in response to pain reflects an increase in GABAergic

activity within the sensorimotor cortex.

GABAergic neurons play a critical role in pain-related brain networks (Barr et al.,

2013; Ong, Stohler, & Herr, 2019). They are involved in the generation of gamma

oscillations (Buzsáki & Wang, 2012), which have been strongly implicated in pain perception

(Barr et al., 2013; Li, Zhang, Zeng, Zhao, & Hu, 2023). Indeed, previous work has shown an

increase in gamma oscillations in response to painful thermal stimuli comparable to the

present study, across a wide range of brain regions such as the prefrontal (Schulz et al., 2015)

and sensorimotor cortices (Gross, Schnitzler, Timmermann, & Ploner, 2007). It is therefore

possible that increases in the N45 peak during pain reflect increased sensorimotor gamma

oscillations. Further multimodal work is required to confirm this finding.

While our findings are consistent with some studies that show increases in

GABAergic activity in response to pain (Gross et al., 2007; Kupers, Danielsen, Kehlet,

Christensen, & Thomsen, 2009; Schulz et al., 2015), other studies have also reported reduced

GABAergic activity in response to experimental pain (Cleve, Gussew, & Reichenbach, 2015;

De Matos, Hock, Wyss, Ettlin, & Brügger, 2017). Differences between studies can be

attributed to the duration of the noxious stimulus (tonic pain lasting several seconds/minutes

vs. transient pain stimuli lasting <1second). Indeed, pooled data have shown that the cerebral

response to pain is highly dependent on the duration of the painful stimuli, as the adaptive

response (to suppress or increase cortical activity) changes depending on the duration of pain

(Chowdhury, et al., 2022). This highlights the need for further work to replicate our findings

using different durations of experimental and clinical pain.

363    Another finding of Experiment 1 was the increase in the amplitude of N100 peak, a

marker of $GABA_B$ neurotransmission (Premoli et al., 2014), and the parietal-occipital P60

peak, a marker of glutamatergic neurotransmission (Belardinelli et al., 2021). However, this

was not replicated in Experiments 2 and 3, potentially due to the smaller sample size.

Nonetheless, we encourage further investigations of alterations in these peaks during pain,

particularly P60 peak, as several magnetic resonance spectroscopy studies have reported

increases in glutamate concentration during experimental pain (Archibald et al., 2020).

Predicting individual differences in pain using TEPs

371    Experimental pain models are useful tools to explore brain measures that may predict

individual differences in pain sensitivity, with an ultimate goal of determining whether such

measures explain why some people develop chronic pain. Experiment 1 showed that higher

pain ratings were associated with a larger increase in the N45 peak during pain. This analysis

was not conducted in Experiments 2 and 3 due to smaller sample sizes and given the primary

aims of Experiments 2 and 3 were to isolate the source of the group-level effect. Nonetheless,

our results suggest that the N45 peak is a potential marker of sensorimotor GABAergic

activity and may be associated with individual differences in pain sensitivity. This is

consistent with other studies measuring GABAergic responses to pain, showing associations

between higher pain sensitivity and larger sensorimotor gamma oscillations (Barr et al., 2013)

and higher left somatosensory cortical GABA laterality (Niddam, Wang, & Tsai, 2021).

However, the direction of this relationship likely depends on the duration of pain

(Chowdhury, et al., 2022). Our results have implications for understanding the development

and maintenance of chronic pain. Further TMS-EEG studies are required to determine

whether the N45 peak is altered in chronic pain populations and whether the N45 peak can

explain why some individuals in the acute stages of pain transition to chronic pain.

The TEP vs. MEP response to pain

The present study showed that a larger reduction in MEP amplitude during pain was

correlated with lower pain ratings, consistent with a recent systematic review (Chowdhury et

al., 2022) and the idea that reduced MEP amplitude is an adaptive mechanism that restricts

movement in the pain-afflicted area, to protect the area from further pain and injury (Hodges

& Tucker, 2011). The novelty of this study was the use of an experimental heat pain

paradigm that has not yet been used in combination with TMS research, and a paradigm that

controls for non-painful somatosensory stimulation.

The finding of a pain-induced increase in the amplitude of the N45 peak, which

indexes $GABA_A$ receptor activity, is consistent with TMS research showing pain-induced

increases in short-interval intracortical inhibition (SICI) (Salo, Vaalto, Koponen, Nieminen,

& Ilmoniemi, 2019; Schabrun & Hodges, 2012). SICI refers to the reduction in MEP

amplitude to a TMS pulse that is preceded 1-5ms by a subthreshold pulse, with this reduction

believed to be mediated by $GABA_A$ neurotransmission (Kujirai et al., 1993). Some studies

have reported associations between SICI and the TEP N45 peak (Leodori et al., 2019; Rawji,

Kaczmarczyk, Rocchi, Rothwell, & Sharma, 2019), suggesting the two may share common

neurophysiological mechanisms. However, we also found that a larger reduction in MEP

amplitude during pain was associated with less pain, while a larger increase in the TEP N45

peak during pain was associated with stronger pain ratings, suggesting that inhibitory

processes mediating MEPs and the TEP N45 peak during pain are distinct. Further work is

required to disentangle the relationship between corticomotor excitability measured by MEPs

and cortical activity measured by TEPs.

Study limitations

Some methodological limitations should be noted. Firstly, while there was no

conclusive evidence for a difference in pain ratings between the six thermal stimuli of the

pain block, there was evidence for fluctuations in pain ratings during each painful stimulus.

This suggests that the perceived pain intensity was not stable across 40 seconds, which may

have introduced noise in the TEP data. Future studies could use pain paradigms that can more

effectively maintain a constant level of pain e.g., hypertonic saline infusion paradigms

(Svensson, Cairns, Wang, & Arendt-Nielsen, 2003). Secondly, the use of verbal pain ratings

prevented the characterisation of pain on a finer time scale. However, verbal ratings were

used to eliminate potential contamination of MEPs introduced by using the hand for

providing pain rating. Thirdly, the increased N45 peak amplitude in response to pain may

reflect increased alertness/arousal during pain. However, a recent study showed that higher

alertness is associated with reduced TEP amplitude (Noreika et al., 2020), suggesting the

increase in the N45 peak amplitude is not related to pain-induced arousal. Lastly, future

research should consider replicating our experiment using intermixed pain and no pain

blocks, as opposed to fixed pre-pain and pain blocks, to control for order effects i.e. the

explanation that successive thermal stimuli applied to the skin results an increase in the N45

peak, regardless of whether the stimuli are painful or not. However, we note that there was no

conclusive evidence for a difference in N45 peak amplitude between pre-pain and post-pain

conditions of Experiment 1 (Supplementary Figure 1), suggesting it is unlikely that the

observed effects were an artefact of time.

Conclusion

This study is the first to use TMS-EEG methodology to examine alterations in cortical

activity in direct response to acute pain. Findings across three experiments suggest that tonic

heat pain leads to an increase in the amplitude of the frontocentral TEP N45 peak (associated

with GABAergic neurotransmission), and that larger increases in this peak are associated

with higher pain ratings. The findings suggest that TEP indices of GABAergic

neurotransmission have the potential as predictive markers of pain severity.

Materials and Methods

Participants

Experiment 1 consisted of 29 healthy participants (18 males, 11 females, mean age; 26.24 ± 5.5). Participants were excluded if they had a history of chronic pain condition or any current acute pain, any contraindication to TMS such as pregnancy, mental implants in the skull, seizure, or if they reported a history of neurological or psychiatric conditions, or were taking psychoactive medication. Participants completed a TMS safety screen (Rossi, Hallett, Rossini, Pascual-Leone, & Group, 2009). Procedures adhered to the Declaration of Helsinki and were approved by the human research ethics committee of UNSW (HC200328). All participants provided informed written consent.

The sample size calculation was done in G*power 3.1.9.7 with 80% power. As there were no prior TMS-EEG studies, we used pooled data from our systematic review on TMS studies (Chowdhury et al., 2022) showing that the weighted effect size of changes in MEP amplitude in response to tonic experimental pain was 0.56. Using this value, 28 participants were required to detect a significant difference between pain and pre-pain blocks.

To determine the sample size of Experiments 2 and 3, we computed the effect sizes of the N45, P60 and N100 changes (pain vs. prepain) from Experiment 1 (Cohen's $d_{RM} = 1.76$, 0.99, 0.83 for N45, P60 and N100 respectively). With a power of 80%, the required sample size was 4-11 participants to detect a significant difference. Experiment 2 recruited a further ten healthy participants (four males, six females, mean age: 26.8 ± 5.9) and Experiment 3 consisted of ten healthy participants (four males, six females, mean age: 28 ± 5.9).

Experimental protocol

Participants were seated comfortably in a shielded room. They viewed a fixation cross to minimise eye movements. TMS was applied to left M1 while participants wore an EEG cap containing 63 scalp electrodes to record TEPs. Surface electromyographic (EMG)

electrodes were placed over the distal region of the right ECRB to record MEPs. EMG

signals were amplified (x 1000) and filtered (16 to 1000Hz), and digitally sampled at 2000

Hz (Spike2, CED). A thermode was attached over the proximal region of the right ECRB in

close proximity to the EMG electrodes (Figure 1).The TMS coil was covered in a layer of

foam (5mm thickness) to minimize decay artefacts (Rogasch et al., 2017). Participants also

wore both foam earplugs and headphones to reduce any potential discomfort from the TMS

click. Auditory masking was not used. Instead, auditory evoked potentials resulting from the

TMS click sound were controlled for in Experiment 2.

The protocol for each experiment is illustrated in Figure 2 (Furman et al., 2020;

Granot et al., 2006). In Experiment 1, participants experienced three blocks of thermal

stimuli, in chronological order: pre-pain, pain, and post-pain block. Each block consisted of

multiple thermal stimuli delivered 40s at a time during which suprathreshold (110% RMT)

TMS measurements and verbal pain ratings were obtained. The thermode commenced at a

baseline temperature of 32°C. The pre-pain and post-pain blocks consisted of six thermal

stimuli delivered at the warm threshold (the temperature that led to any detectable change in

skin temperature from baseline). In the pain block, six thermal stimuli were delivered at

46°C, which has been shown to produce lasting pain with a mean rating of ~5/10 (Furman et

al., 2020). Given we were interested in the individual relationship between pain and

excitability changes, the fixed temperature of 46ºC ensured larger variability in pain ratings

as opposed to calibrating the temperature of the thermode for each participant (Adamczyk et

al., 2022). The inclusion of blocks with warm stimuli allowed for control for changes in

cortical excitability due to non-painful somatosensory stimulation.

The protocol for Experiment 2 and 3 were identical to Experiment 1 with two

differences: the exclusion of the post-pain block (as the aim was to disentangle the source of

the pain vs. pre-pain effect from Experiment 1) and the inclusion of a sham TMS condition

(Experiment 2) or subthreshold (90% RMT) TMS (Experiment 3) intermixed within both the

pre-pain and pain blocks.

Electrical stimulation setup (Experiment 2 Only)

Electrical stimulation was based on previous studies attempting to simulate the

somatosensory component of active TMS (Chowdhury et al., 2022; Gordon et al., 2022;

Rocchi et al., 2021). Prior to EEG setup, 8mm Ag/AgCl electrodes were placed directly over

the scalp. "Snap on" lead wires were then clipped in place and connected to the electrical

stimulator (Digitimer DS7AH, Digitimer Ltd., UK). To keep the electrodes and lead wires

firmly in position, participants were fitted with a tight netted wig cap, which sat on top of the

electrodes but underneath the EEG cap. Consistent with previous research (Chowdhury,

Rogasch, et al., 2022; Rocchi et al., 2021), and to minimize EEG artefacts caused by

electrical stimulation, the stimulating electrodes were not placed directly underneath the EEG

electrodes. Rather, stimulating electrodes were positioned in the middle of the EEG electrode

cluster located in closest proximity to the motor hotspot. This roughly corresponded to an

anode position between FC1 and FC3 and a cathode position between C1 and C3. Scalp

electrical stimulation was delivered using a 200 µs square wave via with a compliance of

200V.

Electroencephalography

EEG was recorded using a DC-coupled, TMS-compatible amplifier (ActiChamp Plus,

Brain Products, Germany) at a sampling rate of 25000 Hz. Signals were recorded from 63

TMS-compatible active electrodes (6mm height, 13mm width), embedded in an elastic cap

(ActiCap, Brain Products, Germany), in line with the international 10-10 system. Active

electrodes result in similar TEPs (both magnitude and peaks) to more commonly used passive

electrodes (Mancuso et al., 2021). There is also evidence that active electrodes have higher

signal quality than passive electrodes at higher impedance levels (Laszlo, Ruiz-Blondet,

Khalifian, Chu, & Jin, 2014). Recordings were referenced online to 'FCz' and the ground

electrode placed on 'FPz'. Electrolyte gel was used to reduce electrode impedances below

~5kOhms. Online TEP monitoring was not available with the EEG software.

Transcranial magnetic stimulation.

Single, monophasic stimuli were delivered using a Magstim unit (Magstim Ltd., UK)

and 70mm figure-of-eight flat coil. The coil was oriented at 45° to the midline, inducing a

current in the posterior-anterior direction. The scalp site that evoked the largest MEP

measured at the ECRB ('hotspot') was determined and marked. The RMT was determined

using the TMS motor thresholding assessment tool, which estimates the TMS intensity

required to induce an MEP of 50 microvolts with a 50% probability using maximum

likelihood parametric estimation by sequential testing (Awiszus, 2003; Awiszus & Borckardt,

2011). This method has been shown to achieve the accuracy of methods such as the Rossini-

Rothwell method (Rossini et al., 1994; Rothwell et al., 1999) but with fewer pulses (Qi, Wu,

& Schweighofer, 2011; Silbert, Patterson, Pevcic, Windnagel, & Thickbroom, 2013). The test

stimulus intensity was set at 110% RMT to concurrently measure MEPs and TEPs during

pre-pain, pain and post-pain blocks.

Thermal pain

Thermal stimuli were delivered over the proximal region of the right ECRB using a

contact heat stimulator (27-mm diameter Medoc Pathway CHEPS Peltier device; Medoc

Advanced Medical Systems Ltd). Pain ratings were obtained after each TMS pulse using a

verbal rating scale (0 = no pain, and 10 = most pain imaginable). Verbal ratings were

collected rather than pain ratings provided on the computer by hand to avoid contamination

of MEP measures from motor processes of hand movements. Verbal pain ratings have been shown to yield excellent test-retest reliability (Alghadir, Anwer, Iqbal, & Iqbal, 2018).

Quantitative sensory testing

Warmth, cold and pain thresholds were assessed in line with a previous study (Furman et al., 2020). With the baseline temperature set at a neutral skin temperature of 32°C, participants completed three threshold tests: to report when they felt a temperature increase (warmth detection threshold) (Furman et al., 2020), to report when they felt a temperature decrease (cool detection threshold) (Furman et al., 2020); 3) to report when an increasing temperature first became painful (heat pain threshold) (Furman et al., 2020). A total of three trials was conducted for each test to obtain an average, with an interstimulus interval of six seconds (Furman et al., 2020). The sequence of cold, warmth and pain threshold was the same for all participants. Participants provided feedback for each trial by pressing a button (with their left hand) on a hand-held device connected to the Medoc Pathway. Temperatures were applied with a rise/decrease rate of 1°C/s and return rate of 2°C/s (initiated by the button click).

Matching task (Experiment 2 only)

As the aim of Experiment 2 was to perceptually match the somatosensory aspects of active and sham TMS, a 2-Alternative Forced Choice task was used to determine the electrical stimulation intensity that led to a similar flicking sensation to active TMS (Chowdhury, et al., 2022). Participants received either electrical stimulation or active TMS in a randomized order and were asked whether the first or second stimulus led to a stronger flick sensation. The electrical stimulation intensity was then increased or decreased until participants could no longer judge the first or second stimulus as stronger. This intensity was then applied during the test blocks.

Test blocks

Experiment 1. The temperature of the thermode commenced at a neutral skin temperature (32°C). Participants were exposed to 18 sustained thermal stimuli with a 20 second interstimulus interval. For each thermal stimulus, a single temperature (rise rate of 1 °C/s, return rate of 2 °C/s) was applied over the proximal region of the ECRB for 40 seconds. Thermal stimuli 1-6 (pre-pain block) were delivered at the participant's individually determined warmth detection threshold, Thermal stimuli 7-12 at 46 degrees, and Thermal stimuli 13-18 again at the participant's warmth detection threshold. Participants were not informed of the order of the warm and painful stimuli to minimize the influence of expectation of pain on TEPs and MEPs. During each 40 second thermal stimulus, TMS pulses were manually delivered, with a verbal pain rating score (0 = no pain, and 10 = worst pain imaginable) obtained between pulses. To avoid contamination of TEPs by verbal ratings, the subsequent TMS pulse was not delivered until the verbal rating was complete, and the participant was cued by the experimenter to provide the pain rating after each pulse. As TMS was delivered manually, there was no set interpulse interval. However, the 40 second thermal stimulus duration allowed for 11 pulses for each thermal stimulus (hence 66 TMS pulses for each of the pre-pain, pain and post-pain blocks), and 10 verbal pain ratings between each TMS pulse (~ 4 seconds in between pain ratings). Current recommendations (Hernandez-Pavon et al., 2023) suggest basing the number of TMS trials per condition on the key outcome measure (e.g., TEP peaks vs. frequency measures) and based on previous test-retest reliability studies. In our study the number of trials was based on a test-retest reliability study by (Kerwin, Keller, Wu, Narayan, & Etkin, 2018) which showed that 60 TMS pulses (delivered in the same run) was sufficient to obtain reliable TEP peaks (i.e., sufficient within-individual concordance between the resultant TEP peaks of each trial).

Experiment 2. Participants were exposed to 24 sustained thermal stimuli (40 seconds

each). Thermal stimuli 1-6 and 7-12 consisted of warm stimuli (pre-pain block), while

Thermal stimuli 13-18 and 19-24 consisted of stimuli delivered at 46°C (pain block). Active

or sham TMS was delivered during either thermal stimuli 1-6 or 7-12, with the order of active

and sham randomly determined for each participant. The same applied for thermal stimuli 13-

18 and 19-24. The active and sham TMS conditions were similar to that used in a recent

TMS-EEG study (Gordon et al., 2021). Sham TMS involved the active TMS coil rotated 90

degrees to the scalp, and a sham coil (identical in shape/weight) placed underneath the active

coil and tangentially over the scalp. The active TMS coil was then triggered with the

electrical stimulation unit to simultaneously simulate the auditory and somatosensory

components of active TMS respectively. Active TMS involved the delivery of the TMS coil

placed tangentially over the scalp, and the sham TMS coil above the active coil rotated 90

degrees to the scalp (see Figure 9). The design allowed for 11 pulses for each thermal

stimulus and ten pain ratings (hence 66 TMS pulses for active pre-pain, sham pre-pain, active

pain and sham pain blocks).

Experiment 3. Participants were exposed to 24 sustained thermal stimuli (40 seconds

each). Thermal stimuli 1-6 and 7-12 consisted of warm stimuli (pre-pain block), while

thermal stimuli 13-18 and 19-24 consisted of thermal stimuli delivered at 46°C (pain block).

Suprathreshold or subthreshold TMS (90% RMT) was delivered during either thermal stimuli

1-6 or 7-12, with the order of supra- and subthreshold TMS randomly determined for each

participant. The same applied for thermal stimuli 13-18 and 19-24. In addition to the pain

rating in between TMS pulses, we collected a second rating for warmth of the thermal

stimulus (0 = neutral, 10 = very warm) to confirm that the participants felt some difference in

sensation relative to baseline during the pre-pain block. This data is presented in the

supplementary material. Overall, the design allowed for 11 pulses for each thermal stimulus

and ten pain/warmth ratings (hence 66 TMS pulses for suprathreshold pre-pain, subthreshold

pre-pain, suprathreshold pain and subthreshold pain blocks).

Data processing

Motor evoked potentials. The amplitude of each MEP was determined using a custom

MATLAB script. The onsets and offsets of the MEPs were manually determined for each

trial. In some participants, background EMG activity was observed due to placement of the

thermode close to the EMG electrodes, which can influence MEP amplitude (Ruddy et al.,

2018). To account for this, MEP amplitude was calculated by subtracting the root mean

square (RMS) of background EMG noise from the RMS of the MEP window using a fixed

window between 55 and 5ms before the TMS pulse (Chowdhury et al., 2023; Schabrun et al.,

2017; Tsao et al., 2011).

TMS-evoked potentials. Pre-processing of the TEPs was completed using EEGLAB

(Delorme & Makeig, 2004) and TESA (Rogasch et al., 2017) in MATLAB (R2021b, The

Math works, USA), and based on previously described methods (Chowdhury et al., 2022;

Mutanen et al., 2018; Rogasch et al., 2017). First, bad channels were removed. The period

between -5 and ~14ms after the TMS pulse was removed and interpolated using the ARFIT

function for continuous data (Neumaier & Schneider, 2001; Schneider & Neumaier, 2001).

The exact interval was based on the duration of decay artefacts. Data was epoched 1000ms

before and after the TMS pulse, and baseline corrected between -1000 and -5ms before the

TMS pulse. Noisy epochs were identified via the EEGLAB auto-trial rejection function

(Delorme, Sejnowski, & Makeig, 2007) and then visually confirmed. The fastICA algorithm

with auto-component rejection used to remove eyeblink and muscle artefacts (Rogasch et al.,

2017). The source-estimation noise-discarding (SOUND) algorithm was applied (Mutanen,

Biabani, Sarvas, Ilmoniemi, & Rogasch, 2020; Mutanen, Metsomaa, Liljander, & Ilmoniemi,

2018), which estimates and supresses noise at each channel based on the most likely cortical

current distribution given the recording of other channels. This signal was then re-referenced

(to average). A band-pass (1-100Hz) and band-stop (48-52Hz) Butterworth filter was then

applied. Any lost channels were interpolated.

Source localization. Source localization of TEPs was conducted using Brainstorm

(Tadel, Baillet, Mosher, Pantazis, & Leahy, 2011). A template brain model (ICBM 152) was

co-registered with the TMS-EEG data. Noise estimation was used to determine sensor

weighting and regularization parameter of the current density construction. The forward

model involved use of the Symmetric Boundary Element Method with the head having 3

compartments of fixed conductivities, implemented in OpenMEEG software (Gramfort,

Papadopoulo, Olivi, & Clerc, 2010), and inverse model involved use of Minimum Norm

Estimations.

Statistical Analysis

Given we were interested in determining the evidence for pain altering TEP peaks in

certain conditions (e.g., active TMS) and pain not altering TEP peaks in other conditions

(sham TMS), we used a Bayesian approach as opposed to a frequentist approach, which

considers the strength of the evidence for the alternative vs. null hypothesis. Bayesian

inference was used to analyse the data using JASP software (Version 0.12.2.0, JASP Team,

2020). Bayes factors were expressed as BF_{10} values, where BF_{10}'s of 1-3, 3-10, 10-30 and 30-

100 indicated "weak", "moderate", "strong" and "very strong" evidence for the alternative

hypothesis, while BF_{10}'s of 1/3-1, 1/10-1/3, 1/30-1/10 and 1/100-1/30 indicated "anecdotal",

"moderate", "strong" and "very strong" evidence in favour of the null hypothesis (van Doorn

et al., 2021).

Pain ratings. A 6 (thermal stimulus number: 1-6) x 10 (timepoint:1-10) Bayesian

repeated measures ANOVA with default priors in JASP (r scale fixed effects = .5, r scale

random effects = 1, r scale covariates = .354) was conducted on the pain ratings during the

pain block. This was to assess differences in pain ratings between the six painful stimuli and

whether pain ratings differed between the timepoints of each painful stimulus.

MEPs. A Bayesian one-way repeated measures ANOVA with default priors in JASP

was performed to assess differences in MEP amplitudes between pre-pain, pain and post-pain

blocks of Experiment 1. As there is now increasing emphasis placed on investigating the

individual level relationship between changes in cortical excitability and pain and not only

the group level effect, (Chowdhury et al., 2022; Seminowicz et al., 2018; Seminowicz,

Thapa, & Schabrun, 2019; Summers et al., 2019) we also investigated the correlations

between pain ratings and changes in MEP (and TEP) amplitude, A Bayesian correlation

analysis with default priors in JASP (stretched Beta prior width = 1) was run to determine

whether the change in mean MEP amplitude during pain (as a proportion of pre-pain) was

associated with the mean verbal pain rating score. Data were checked for assumptions of

normally distributed data using a Shapiro-Wilk test. Where assumptions were violated, data

were log-transformed.

TEPs. The grand-averaged signals for the pre-pain, pain and post-pain condition were

obtained. For Experiment 1, a cluster-based permutation analysis was used to compare

amplitude levels between pre-pain and pain, and pre-pain and post-pain, at each time-point

and for each electrode. For all experiments, the mean TEP waveform of any identified

clusters from Experiment 1 were plotted, and peaks (e.g., N15, P30, N45, P60, N100) were

identified using the TESA peak function (Rogasch et al., 2017). Any identified peaks were

then compared between conditions using Bayes paired sample t-tests with default priors in

JASP (Cauchy scale = .707). A Bayesian correlation analysis with default priors in JASP was

performed to determine whether the difference in identified peaks between pre-pain and pain

blocks was associated with the mean pain rating score. Data were checked for assumptions of

normally distributed data using a Shapiro-Wilk test. Where assumptions were violated, data

were log-transformed.

Acknowledgements

This work was supported by 1R61NS113269-01 from The National Institutes of Health to

DAS, SMS, and The Four Borders Foundation to DAS.

References

Adamczyk, W. M., Szikszay, T. M., Nahman-Averbuch, H., Skalski, J., Nastaj, J., Gouverneur, P., &
Luedtke, K. (2022). To calibrate or not to calibrate? A methodological dilemma in
experimental pain research. *The journal of Pain, 23*(11), 1823-1832.
Ahn, S., & Fröhlich, F. (2021). Pinging the brain with transcranial magnetic stimulation reveals
cortical reactivity in time and space. *Brain stimulation, 14*(2), 304-315.
Alghadir, A. H., Anwer, S., Iqbal, A., & Iqbal, Z. A. (2018). Test–retest reliability, validity, and
minimum detectable change of visual analog, numerical rating, and verbal rating scales for
measurement of osteoarthritic knee pain. *Journal of pain research, 11*, 851.
Apkarian, A. V. (2021). The Necessity of Methodological Advances in Pain Research: Challenges and
Opportunities. *Frontiers in Pain Research, 2*, 10.
Archibald, J., MacMillan, E. L., Graf, C., Kozlowski, P., Laule, C., & Kramer, J. L. K. (2020). Metabolite
activity in the anterior cingulate cortex during a painful stimulus using functional MRS.
Scientific Reports, 10(1), 19218. doi:10.1038/s41598-020-76263-3
Awiszus, F. (2003). TMS and threshold hunting. In *Supplements to Clinical neurophysiology* (Vol. 56,
pp. 13-23): Elsevier.
Awiszus, F., & Borckardt, J. (2011). TMS motor threshold assessment tool (MTAT 2.0). *Brain
Stimulation Laboratory, Medical University of South Carolina, USA*.
Bank, P. J., Peper, C., Marinus, J., Beek, P. J., & Van Hilten, J. (2013). Motor consequences of
experimentally induced limb pain: a systematic review. *European journal of pain, 17*(2), 145-
157.
Barr, M. S., Farzan, F., Davis, K. D., Fitzgerald, P. B., & Daskalakis, Z. J. (2013). Measuring GABAergic
inhibitory activity with TMS-EEG and its potential clinical application for chronic pain. *Journal
of Neuroimmune Pharmacology, 8*(3), 535-546.
Belardinelli, P., König, F., Liang, C., Premoli, I., Desideri, D., Müller-Dahlhaus, F., . . . Ziemann, U.
(2021). TMS-EEG signatures of glutamatergic neurotransmission in human cortex. *Scientific
Reports, 11*(1), 1-14.
Biabani, M., Fornito, A., Mutanen, T. P., Morrow, J., & Rogasch, N. C. (2019). Characterizing and
minimizing the contribution of sensory inputs to TMS-evoked potentials. *Brain stimulation,
12*(6), 1537-1552.
Brodbeck, V., Spinelli, L., Lascano, A. M., Wissmeier, M., Vargas, M.-I., Vulliemoz, S., . . . Seeck, M.
(2011). Electroencephalographic source imaging: a prospective study of 152 operated
epileptic patients. *Brain, 134*(10), 2887-2897. doi:10.1093/brain/awr243
Burns, E., Chipchase, L. S., & Schabrun, S. M. (2016). Primary sensory and motor cortex function in
response to acute muscle pain: A systematic review and meta-analysis. *European Journal of
pain, 20*(8), 1203-1213. doi:10.1002/ejp.859
Buzsáki, G., & Wang, X.-J. (2012). Mechanisms of gamma oscillations. *Annual review of neuroscience,
35*, 203.
Chang, W.-J., O'Connell, N. E., Beckenkamp, P. R., Alhassani, G., Liston, M. B., & Schabrun, S. M.
(2018). Altered primary motor cortex structure, organization, and function in chronic pain: a
systematic review and meta-analysis. *The journal of Pain, 19*(4), 341-359.
Che, X., Cash, R., Chung, S. W., Bailey, N., Fitzgerald, P. B., & Fitzgibbon, B. M. (2019). The
dorsomedial prefrontal cortex as a flexible hub mediating behavioral as well as local and
distributed neural effects of social support context on pain: A Theta Burst Stimulation and
TMS-EEG study. *Neuroimage, 201*, 116053.
Chowdhury, N. S., Chang, W.-J., Millard, S. K., Skippen, P., Bilska, K., Seminowicz, D. S., & Schabrn, S.
M. (2022). The Effect of Acute and Sustained Pain on Corticomotor Excitability: A Systematic
Review and Meta-analysis of group-and individual-level data. *The journal of Pain*.

Chowdhury, N. S., Rogasch, N. C., Chiang, A. K., Millard, S. K., Skippen, P., Chang, W.-J., . . . Schabrun,
S. M. (2022). The influence of sensory potentials on transcranial magnetic stimulation–
Electroencephalography recordings. *Clinical Neurophysiology, 140*, 98-109.

Cleve, M., Gussew, A., & Reichenbach, J. R. (2015). In vivo detection of acute pain-induced changes
of GABA+ and Glx in the human brain by using functional 1H MEGA-PRESS MR spectroscopy.
Neuroimage, 105, 67-75.

Conde, V., Tomasevic, L., Akopian, I., Stanek, K., Saturnino, G. B., Thielscher, A., . . . Siebner, H. R.
(2019). The non-transcranial TMS-evoked potential is an inherent source of ambiguity in
TMS-EEG studies. *Neuroimage, 185*, 300-312.

De Matos, N. M., Hock, A., Wyss, M., Ettlin, D. A., & Brügger, M. (2017). Neurochemical dynamics of
acute orofacial pain in the human trigeminal brainstem nuclear complex. *Neuroimage, 162*,
162-172.

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG
dynamics including independent component analysis. *Journal of neuroscience methods,*
134(1), 9-21.

Delorme, A., Sejnowski, T., & Makeig, S. (2007). Enhanced detection of artifacts in EEG data using
higher-order statistics and independent component analysis. *Neuroimage, 34*(4), 1443-1449.

Dubé, J. A., & Mercier, C. (2011). Effect of pain and pain expectation on primary motor cortex
excitability. *Clinical Neurophysiology, 122*(11), 2318-2323.

Farzan, F., & Bortoletto, M. (2022). Identification and verification of a'true'TMS evoked potential in
TMS-EEG. *Journal of neuroscience methods*, 109651.

Farzan, F., Vernet, M., Shafi, M. M., Rotenberg, A., Daskalakis, Z. J., & Pascual-Leone, A. (2016).
Characterizing and modulating brain circuitry through transcranial magnetic stimulation
combined with electroencephalography. *Frontiers in neural circuits, 10*, 73.

Frömer, R., Maier, M., & Abdel Rahman, R. (2018). Group-level EEG-processing pipeline for flexible
single trial-based analyses including linear mixed models. *Frontiers in neuroscience, 12*, 48.

Furman, A. J., Prokhorenko, M., Keaser, M. L., Zhang, J., Chen, S., Mazaheri, A., & Seminowicz, D. A.
(2020). Sensorimotor peak alpha frequency is a reliable biomarker of prolonged pain
sensitivity. *Cerebral Cortex, 30*(12), 6069-6082.

Gordon, P. C., Jovellar, D. B., Song, Y., Zrenner, C., Belardinelli, P., Siebner, H. R., & Ziemann, U.
(2021). Recording brain responses to TMS of primary motor cortex by EEG–utility of an
optimized sham procedure. *Neuroimage*, 118708.

Gramfort, A., Papadopoulo, T., Olivi, E., & Clerc, M. (2010). OpenMEEG: opensource software for
quasistatic bioelectromagnetics. *Biomedical engineering online, 9*(1), 1-20.

Granot, M., Granovsky, Y., Sprecher, E., Nir, R.-R., & Yarnitsky, D. (2006). Contact heat-evoked
temporal summation: tonic versus repetitive-phasic stimulation. *Pain, 122*(3), 295-305.

Gross, J., Schnitzler, A., Timmermann, L., & Ploner, M. (2007). Gamma oscillations in human primary
somatosensory cortex reflect pain perception. *PLoS biology, 5*(5), e133.

Hernandez-Pavon, J. C., Veniero, D., Bergmann, T. O., Belardinelli, P., Bortoletto, M., Casarotto, S., . .
. Julkunen, P. (2023). TMS combined with EEG: Recommendations and open issues for data
collection and analysis. *Brain Stimulation*.

Hodges, P. W., & Tucker, K. (2011). Moving differently in pain: A new theory to explain the
adaptation to pain. *Pain, 152*(SUPPL.3), S90-S98. doi:10.1016/j.pain.2010.10.020

Ilmoniemi, R. J., & Kičić, D. (2010). Methodology for combined TMS and EEG. *Brain Topography, 22*,
233-248.

Kallioniemi, E., & Daskalakis, Z. J. (2022). Identifying novel biomarkers with TMS-EEG -
Methodological possibilities and challenges. *J Neurosci Methods, 377*, 109631.
doi:10.1016/j.jneumeth.2022.109631

Kerwin, L. J., Keller, C. J., Wu, W., Narayan, M., & Etkin, A. (2018). Test-retest reliability of
transcranial magnetic stimulation EEG evoked potentials. *Brain stimulation, 11*(3), 536-544.
doi:https://doi.org/10.1016/j.brs.2017.12.010

Kujirai, T., Caramia, M., Rothwell, J. C., Day, B., Thompson, P., Ferbert, A., . . . Marsden, C. D. (1993).
Corticocortical inhibition in human motor cortex. *The Journal of Physiology, 471*(1), 501-519.
Kupers, R., Danielsen, E. R., Kehlet, H., Christensen, R., & Thomsen, C. (2009). Painful tonic heat
stimulation induces GABA accumulation in the prefrontal cortex in man. *Pain, 142*(1-2), 89-
93.
Laszlo, S., Ruiz-Blondet, M., Khalifian, N., Chu, F., & Jin, Z. (2014). A direct comparison of active and
passive amplification electrodes in the same amplifier system. *Journal of neuroscience*
methods, 235, 298-307.
Leodori, G., Thirugnanasambandam, N., Conn, H., Popa, T., Berardelli, A., & Hallett, M. (2019).
Intracortical inhibition and surround inhibition in the motor cortex: a TMS-EEG study.
Frontiers in neuroscience, 13, 612.
Li, Z., Zhang, L., Zeng, Y., Zhao, Q., & Hu, L. (2023). Gamma-band oscillations of pain and nociception:
A systematic review and meta-analysis of human and rodent studies. *Neuroscience &*
Biobehavioral Reviews, 146, 105062. doi:https://doi.org/10.1016/j.neubiorev.2023.105062
Mancuso, M., Sveva, V., Cruciani, A., Brown, K., Ibáñez, J., Rawji, V., . . . Rothwell, J. (2021).
Transcranial evoked potentials can be reliably recorded with active electrodes. *Brain*
Sciences, 11(2), 145.
Massimini, M., Ferrarelli, F., Huber, R., Esser, S. K., Singh, H., & Tononi, G. (2005). Breakdown of
cortical effective connectivity during sleep. *Science, 309*(5744), 2228-2232.
Michel, C. M., & Brunet, D. (2019). EEG source imaging: a practical review of the analysis steps.
Frontiers in Neurology, 10, 325.
Mutanen, T. P., Biabani, M., Sarvas, J., Ilmoniemi, R. J., & Rogasch, N. C. (2020). Source-based
artifact-rejection techniques available in TESA, an open-source TMS–EEG toolbox. *Brain*
Stimulation: Basic, Translational, and Clinical Research in Neuromodulation, 13(5), 1349-
1351.
Mutanen, T. P., Metsomaa, J., Liljander, S., & Ilmoniemi, R. J. (2018). Automatic and robust noise
suppression in EEG and MEG: The SOUND algorithm. *Neuroimage, 166*, 135-151.
Neumaier, A., & Schneider, T. (2001). Estimation of parameters and eigenmodes of multivariate
autoregressive models. *ACM Transactions on Mathematical Software (TOMS), 27*(1), 27-57.
Niddam, D. M., Wang, S.-J., & Tsai, S.-Y. (2021). Pain sensitivity and the primary sensorimotor
cortices: a multimodal neuroimaging study. *Pain, 162*(3), 846-855.
Ong, W.-Y., Stohler, C. S., & Herr, D. R. (2019). Role of the Prefrontal Cortex in Pain Processing.
Molecular Neurobiology, 56(2), 1137-1166. doi:10.1007/s12035-018-1130-9
Petrichella, S., Johnson, N., & He, B. (2017). The influence of corticospinal activity on TMS-evoked
activity and connectivity in healthy subjects: A TMS-EEG study. *PloS one, 12*(4), e0174879.
Ploner, M., & May, E. S. (2018). Electroencephalography and magnetoencephalography in pain
research—current state and future perspectives. *Pain, 159*(2), 206-211.
Premoli, I., Castellanos, N., Rivolta, D., Belardinelli, P., Bajo, R., Zipser, C., . . . Ziemann, U. (2014).
TMS-EEG signatures of GABAergic neurotransmission in the human cortex. *Journal of*
Neuroscience, 34(16), 5603-5612.
Qi, F., Wu, A. D., & Schweighofer, N. (2011). Fast estimation of transcranial magnetic stimulation
motor threshold. *Brain stimulation, 4*(1), 50-57.
Rawji, V., Kaczmarczyk, I., Rocchi, L., Rothwell, J. C., & Sharma, N. (2019). Short interval intracortical
inhibition as measured by TMS-EEG. *BioRxiv*, 802504.
Reddan, M. C., & Wager, T. D. (2018). Modeling pain using fMRI: from regions to biomarkers.
Neuroscience bulletin, 34(1), 208-215.
Rocchi, L., Di Santo, A., Brown, K., Ibáñez, J., Casula, E., Rawji, V., . . . Rothwell, J. (2021).
Disentangling EEG responses to TMS due to cortical and peripheral activations. *Brain*
stimulation, 14(1), 4-18.
Rogasch, N. C., Sullivan, C., Thomson, R. H., Rose, N. S., Bailey, N. W., Fitzgerald, P. B., . . . Hernandez-
Pavon, J. C. (2017). Analysing concurrent transcranial magnetic stimulation and

electroencephalographic data: A review and introduction to the open-source TESA software. *Neuroimage, 147*, 934-951.

Rohel, A., Bouffard, J., Patricio, P., Mavromatis, N., Billot, M., Roy, J.-S., . . . Masse-Alarie, H. (2021). The effect of experimental pain on the excitability of the corticospinal tract in humans: A systematic review and meta-analysis: *European Journal of Pain, 25*(6), 1209-1226..

Rossi, S., Hallett, M., Rossini, P. M., Pascual-Leone, A., & Group, S. o. T. C. (2009). Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clinical Neurophysiology, 120*(12), 2008-2039.

Rossini, P. M., Barker, A., Berardelli, A., Caramia, M., Caruso, G., Cracco, R., . . . Lücking, C. (1994). Non-invasive electrical and magnetic stimulation of the brain, spinal cord and roots: basic principles and procedures for routine clinical application. Report of an IFCN committee. *Electroencephalography and clinical Neurophysiology, 91*(2), 79-92.

Rothwell, J. C., Hallett, M., Berardelli, A., Eisen, A., Rossini, P., & Paulus, W. (1999). Magnetic stimulation: motor evoked potentials. The International Federation of Clinical Neurophysiology. *Electroencephalography and clinical neurophysiology. Supplement, 52*, 97-103.

Ruddy, K., Balsters, J., Mantini, D., Liu, Q., Kassraian-Fard, P., Enz, N., . . . Wenderoth, N. (2018). Neural activity related to volitional regulation of cortical excitability. *Elife, 7*.

Salo, K. S.-T., Vaalto, S. M., Koponen, L. M., Nieminen, J. O., & Ilmoniemi, R. J. (2019). The effect of experimental pain on short-interval intracortical inhibition with multi-locus transcranial magnetic stimulation. *Experimental Brain Research, 237*(6), 1503-1510.

Schabrun, S. M., & Hodges, P. W. (2012). Muscle pain differentially modulates short interval intracortical inhibition and intracortical facilitation in primary motor cortex. *Journal of Pain, 13*(2), 187-194. doi:10.1016/j.jpain.2011.10.013

Schneider, T., & Neumaier, A. (2001). Algorithm 808: ARfit—A Matlab package for the estimation of parameters and eigenmodes of multivariate autoregressive models. *ACM Transactions on Mathematical Software (TOMS), 27*(1), 58-65.

Schulz, E., May, E. S., Postorino, M., Tiemann, L., Nickel, M. M., Witkovsky, V., . . . Ploner, M. (2015). Prefrontal gamma oscillations encode tonic pain in humans. *Cerebral cortex, 25*(11), 4407-4414.

Seminowicz, D. A., Thapa, T., Furman, A. J., Summers, S. J., Cavaleri, R., Fogarty, J. S., . . . Schabrun, S. M. (2018). Slow peak alpha frequency and corticomotor depression linked to high pain susceptibility in transition to sustained pain. *BioRxiv*, 278598.

Seminowicz, D. A., Thapa, T., & Schabrun, S. M. (2019). Corticomotor depression is associated with higher pain severity in the transition to sustained pain: A longitudinal exploratory study of individual differences. *The journal of Pain, 20*(12), 1498-1506.

Shirazi, S. Y., & Huang, H. J. (2019). More reliable EEG electrode digitizing methods can reduce source estimation uncertainty, but current methods already accurately identify brodmann areas. *Frontiers in neuroscience, 13*, 1159.

Silbert, B., Patterson, H., Pevcic, D., Windnagel, K., & Thickbroom, G. (2013). A comparison of relative-frequency and threshold-hunting methods to determine stimulus intensity in transcranial magnetic stimulation. *Clinical Neurophysiology, 124*(4), 708-712.

Summers, S. J., Chipchase, L. S., Hirata, R., Graven-Nielsen, T., Cavaleri, R., & Schabrun, S. M. (2019). Motor adaptation varies between individuals in the transition to sustained pain. *Pain, 160*(9), 2115-2125.

Svensson, P., Cairns, B. E., Wang, K., & Arendt-Nielsen, L. (2003). Injection of nerve growth factor into human masseter muscle evokes long-lasting mechanical allodynia and hyperalgesia. *Pain, 104*(1-2), 241-247.

Tadel, F., Baillet, S., Mosher, J. C., Pantazis, D., & Leahy, R. M. (2011). Brainstorm: a user-friendly application for MEG/EEG analysis. *Computational Intelligence and Neuroscience, 2011*.

Tremblay, S., Rogasch, N. C., Premoli, I., Blumberger, D. M., Casarotto, S., Chen, R., . . . Fitzgerald, P.
B. (2019). Clinical utility and prospective of TMS–EEG. *Clinical Neurophysiology, 130*(5), 802-
844.
van Doorn, J., van den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., . . . Haaf, J. M. (2021).
The JASP guidelines for conducting and reporting a Bayesian analysis. *Psychonomic Bulletin*
& Review, 28, 813-826.
Ye, Y., Wang, J., & Che, X. (2022). Concurrent TMS-EEG to reveal the neuroplastic changes in the
prefrontal and insular cortices in the analgesic effects of DLPFC-rTMS. *Cerebral Cortex,*
32(20), 4436-4446.

956 **Figures**

Figure 1. **Schematic of experimental apparatus.** This consisted of transcranial magnetic
stimulation (TMS) during concurrent electroencephalography (EEG) to simultaneously
record motor evoked potentials (MEPs) and TMS-evoked potentials (TEPs). MEPs were
recorded using electromyographic (EMG) electrodes placed over the distal region of the
extensor carpi radialis brevis (ECRB) while thermal pain was delivered over the proximal
region of the ECRB.

Figure 2. **Schematic of the experimental protocols.** In Experiment 1, participants
experienced three blocks of thermal stimuli: a pre-pain, pain, and post-pain block, with each
block consisting of multiple thermal stimuli delivered 40s at a time, and during which TMS
measurements (indicated by blue arrows) and verbal pain ratings were obtained. The pre-
pain and post-pain blocks involved thermal stimuli delivered at the warm threshold (i.e., the
temperature that leads to any perceived change in skin temperature from baseline). In the pain
block, thermal stimuli were delivered at 46°C. For Experiment 2, the post-pain block was
excluded, and an additional sham TMS condition was intermixed within both the pre-pain
and pain blocks. For Experiment 3, the post-pain block was also excluded, and an additional
subthreshold TMS condition intermixed within both the pre-pain and pain blocks.
982
983

Figure 3. **No conclusive evidence of a difference in pain ratings between successive 46°C 40s thermal stimuli.** Mean (± SD) pain ratings during the 6 thermal stimuli delivered during the pain block (thermal stimuli delivered at 46°C) of Experiment 1. 10 pain ratings were collected over each 40 second thermal stimulus ~ every 4 second.

1006

1007   *Figure 4.* **No conclusive evidence of MEP amplitude differences between conditions;**
1008   **however individual pain sensitivity was predicted by changes in MEP amplitude.** A:
1009   Mean (± SD) MEP amplitude during the pre-pain, pain and post-pain blocks of Experiment 1.
1010   B: Individual-level Relationship between change in MEP amplitude during pain (proportion
1011   of pre-pain) and mean verbal pain rating provided by each participant.

1012

1013

1014

1015

1016

1017

1018

1019

1020

1021

Figure 5. **Pain led to increased negative and positive amplitude in frontocentral and parietal-occipital sites respectively, 43-90ms after the TMS pulse.** A: Grand-average TEPs during the pre-pain, pain and post-pain blocks of Experiment 1. The grey shaded area represents the window of interpolation around the TMS pulse. B: Scalp topographies and estimated source activity at timepoints where TEP peaks are commonly observed, including the N15, P30, N45, P60, N100 and P180. A cluster plot is also shown on the right comparing signal amplitude between the pain and pre-pain conditions at a representative timepoint (48ms) between 43-90ms, which is where significant amplitude differences were observed. The black stars demonstrate the presence of significant positive (yellow) or negative (blue) clusters.

10:

Figure 6. **Pain led to increases in N45, P60 and N100 peak amplitude, and individual pain sensitivity was predicted by changes in the N45 peak.** TEPs across pain and pre-pain condition for the frontocentral electrodes (A) and parietal-occipital electrodes (C) identified from the cluster analysis of Experiment 1. The grey shaded area represents the window of interpolation around the transcranial magnetic stimulation (TMS) pulse. For the frontocentral electrodes, two significantly stronger negative peaks were identified at ~45 and 85ms post-TMS. For the parietal-occipital electrodes, a significantly stronger positive peak was identified at ~50ms post-TMS. * indicates at least moderate evidence the alternative hypothesis that the amplitude is larger in pain vs. pre-pain ($BF_{10} > 3$). Individual-level relationship between mean verbal pain ratings provided by each participant and change in peak amplitudes at ~45ms (N45), ~85ms (N100) post-TMS (B), and ~50ms (P60) post-TMS (D).

Figure 7. **TMS-evoked potentials for Active and Sham TMS.** A: Schematics showing the delivery of active and sham TMS of Experiment 2. Sham TMS involved scalp electrical stimulation (in red) beneath a sham coil (in dotted blue) to mimic somatosensory stimulation associated with active TMS, and concurrent delivery of active TMS 90 degrees to the scalp (in shaded blue) to mimic auditory stimulation associated with TMS. B: Left: TEPs during the pre-pain and pain blocks, for both active and sham stimulation. The grey shaded area represents the window of interpolation around the TMS pulse. Right: Scalp topographies and estimated source activity at timepoints where TEP peaks are commonly observed, including the N15, P30, N45, P60, N100 and P180.

Figure 8. **Pain led to an increase in the N45 peak amplitude during active TMS but not sham TMS.** TEPs during pain and pre-pain blocks, across active and sham TMS conditions of Experiment 2, for the frontocentral electrodes (left) and parietal occipital electrodes (right) identified from the cluster analysis in the main experiment. A significantly stronger frontocentral negative peak was identified ~45ms post-TMS during pain compared to pre-pain, for the active TMS condition. * indicates at least moderate evidence the alternative hypothesis that the amplitude is larger in pain vs. pre-pain ($BF_{10} > 3$). The dotted line shows the timing of the peak.

1070

1071

Figure 9. **TMS-evoked potentials for supra- and subthreshold TMS.** Left: TEPs during the pre-pain and pain blocks, for both supra- and subthreshold TMS of Experiment 3. The grey shaded area represents the window of interpolation around the transcranial magnetic stimulation TMS pulse. Right: Scalp topographies and estimated source activity at timepoints where TEP peaks are commonly observed, including the N15, P30, N45, P60, N100 and P180.

1078

1079

1080

Figure 10. **Pain led to an increase in the N45 peak amplitude for both suprathreshold**
and subthreshold TMS. TEPs during pain and pre-pain blocks of Experiment 3, across
supra- and subthreshold TMS conditions, for the frontocentral electrodes (left) and parietal
occipital electrodes (right) identified from the cluster analysis in Experiment 1. A
significantly stronger frontocentral negative peak was identified ~45ms post-TMS during
pain compared to pre-pain for both supra- and subthreshold stimulation. * indicates at least
moderate evidence the alternative hypothesis that the amplitude is larger in pain vs. pre-pain
($BF_{10} > 3$). The dotted line shows the timing of the peak.
1089

1090

1091

1092

1093

1094

1095

1096

1097