Alterations in cortical excitability during pain: A combined TMS-EEG Study

Nahian S Chowdhury[1,2], Alan KI Chiang[1,2], Samantha K Millard[1,2], Patrick Skippen[1], Wei-Ju Chang[1,3], David A Seminowicz[1,4,5,6], Siobhan M Schabrun[1,7]

Affiliations

[1]Center for Pain IMPACT, Neuroscience Research Australia, Sydney, New South Wales, Australia, [2]University of New South Wales, Sydney, New South Wales, Australia, [3]School of Health Sciences, College of Health, Medicine and Wellbeing, The University of Newcastle, Callaghan, New South Wales, Australia, [4]Department of Neural and Pain Sciences, School of Dentistry, University of Maryland Baltimore, [5]Center to Advance Chronic Pain Research, University of Maryland Baltimore, [6]Department of Medical Biophysics, University of Western Ontario, Canada, [7]The Gray Centre for Mobility and Activity, University of Western Ontario, London, Canada.

Corresponding author

Nahian S Chowdhury

Email: n.chowdhury@neura.edu.au

Neuroscience Research Australia, 139 Barker Street Randwick 2031 NSW

Disclosures

DAS is an advisor to Empower Therapeutics. The other authors have no conflicts of interest to declare.

Abstract

Transcranial magnetic stimulation (TMS) has been used to examine the inhibitory and facilitatory circuits during experimental pain and in chronic pain populations. However, current applications of TMS to pain have been restricted to measurements of motor evoked potentials (MEPs) from peripheral muscles. Here, TMS was combined with electroencephalography (EEG) to determine whether experimental pain could induce alterations in cortical inhibitory/facilitatory activity observed in TMS-evoked potentials (TEPs). In Experiment 1 (n = 29), multiple sustained thermal stimuli were administered over the forearm, with the first, second and third block of stimuli consisting of warm but non-painful (pre-pain block), painful heat (pain block) and warm but non-painful (post-pain block) temperatures respectively. During each stimulus, TMS pulses were delivered while EEG (64 channels) was simultaneously recorded. Verbal pain ratings were collected between TMS pulses. Relative to pre-pain warm stimuli, painful stimuli led to an increase in the amplitude of the frontocentral negative peak ~45ms post-TMS (N45), with a larger increase associated with higher pain ratings. Experiments 2 and 3 (n = 10 in each) showed that the increase in the N45 in response to pain was not due to changes in sensory potentials associated with TMS, or a result of stronger reafferent muscle feedback during pain. This is the first study to use combined TMS-EEG to examine alterations in cortical excitability in response to pain. These results suggest that the N45 TEP peak, which indexes GABAergic neurotransmission, is implicated in pain perception and is a potential marker of individual differences in pain sensitivity.

Introduction

Pain is a complex subjective experience, and understanding how pain is processed remains a challenge (Apkarian, 2021). Several neuroimaging techniques have been applied to disentangle these complexities: functional magnetic resonance imaging has helped us understand where in the brain pain is processed (Reddan & Wager, 2018), while electroencephalography (EEG) has contributed to our understanding of the temporal sequence of pain processing (Ploner & May, 2018). Another useful technique that has been used to examine the role of inhibitory and facilitatory neural circuits in pain has been transcranial magnetic stimulation (TMS) (Chang et al., 2018; Schabrun & Hodges, 2012). However, a limitation of TMS studies has been the recording of output from the muscle, a signal that could be influenced by many intermediate (subcortical, spinal, peripheral) factors, and which restricts investigations to the motor system only. Here, we use a combined TMS-EEG measure to record output of TMS directly from the cortex and from multiple brain regions, in pain-free and tonic pain conditions.

When TMS is delivered over the primary motor cortex (M1), a magnetic pulse induces an electrical current in underlying cortical tissue that, if the intensity is sufficient, activates corticomotor pathways, inducing a motor evoked potential (MEP) in a target muscle. The magnitude of the MEP serves as an index of the excitability of the pathway between M1 and the muscle. Past systematic reviews on studies using TMS during acute experimental pain (Bank, Peper, Marinus, Beek, & Van Hilten, 2013; Burns, Chipchase, & Schabrun, 2016; Chowdhury et al., 2022; Rohel et al., 2021) have shown a reduction in corticomotor excitability during pain and after pain resolution, with stronger reductions in corticomotor excitability associated with lower acute pain severity (Chowdhury et al., 2022). It has been hypothesised that this reduction in MEP amplitude is an adaptive mechanism that

restricts movement in the pain-afflicted area, to protect the area from further pain and injury (Hodges & Tucker, 2011).

While previous findings show promise for the use of the TMS to discover and validate potential biomarkers for pain, limitations exist when using TMS alone. First, MEP responses to TMS reflect the net sum of cortical, spinal, and peripheral activity within the corticomotor pathway. As such, the precise origin of the pain mechanism cannot be localized. Further, measurement of MEPs restricts investigations to M1. One way of overcoming the limitations of using TMS alone is by combining TMS and EEG. TMS-EEG is an increasingly popular tool that allows for the non-invasive investigation of cortical excitability, connectivity and plasticity *directly* from the cortex (i.e. without influence of subcortical, spinal and peripheral processes), as well as from regions *outside M1* (Farzan et al., 2016). TMS-EEG also provides an index of the activity of specific neurotransmitter circuits within the cortex. For example, TMS-evoked potential (TEP) peaks that occur at ~45ms and 100ms post-stimulation are linked to $GABA_A$ and $GABA_B$ neurotransmission respectively (Premoli et al., 2014), while the TEP peak ~60ms post-stimulation is linked to glutamatergic neurotransmission (Belardinelli et al., 2021). Overall, TMS-EEG provides additional spatial and temporal information about cortical activity over TMS alone, making it ideal for understanding the brain mechanisms involved in pain perception.

TMS-EEG has already shown potential to serve as a biomarker for the development and prognosis of various neurological and psychiatric conditions (for reviews see (Kallioniemi & Daskalakis, 2022; Tremblay et al., 2019)). However, the applicability of TMS-EEG to pain research is yet to be established. While GABAergic processes indexed by TMS-EEG have been hypothesised to be involved in pain (Barr, Farzan, Davis, Fitzgerald, & Daskalakis, 2013), direct evidence is scarce. Two studies (Ye et al., 2022; Che et al, 2019) examined whether the potential analgesic effects of repetitive TMS (rTMS) over the dorsal

prefrontal cortex are associated with plasticity in TEPs. These studies separately measured TEPs and ratings to painful stimuli, before and after rTMS, with one finding that increases in pain thresholds following rTMS were associated with changes in TEPs that index GABAergic processes (Ye et al., 2022). However, no study has investigated whether cortical activity is altered in direct response to pain.

The aim of the present study was to use TMS-EEG to determine whether acute experimental pain induces alterations in cortical inhibitory and facilitatory peaks observed using TEPs. We used a tonic heat pain paradigm (Furman et al., 2020; Granot, Granovsky, Sprecher, Nir, & Yarnitsky, 2006), in which multiple thermal stimuli were applied over the right extensor carpi radialis brevis (ECRB) muscle via a thermode. For each thermal stimulus, the temperature increased from a neural baseline of 32°C to either a warm non-painful or a painful (46°C) temperature, with this temperature maintained for 40 seconds. During this time, TMS was administered to the left M1 with concurrent EEG to obtain TEPs from 63 scalp channels, and MEPs from the ECRB muscle (see Figure 1). Verbal pain ratings were obtained between pulses. It was hypothesized that TEP peaks that index GABAergic processes, including the peaks at ~45 and 100ms after TMS, would increase in response to painful stimuli relative to warm non-painful stimuli.

Results

Experiment 1 – Does acute pain alter cortical excitability?

In Experiment 1 (n = 29), we determined whether painful thermal stimuli induced alterations in TEP peaks relative to a non-painful baseline. The protocol (Figure 2) consisted of three blocks of stimuli, in chronological order: pre-pain, pain, and post-pain blocks. The pre-pain and post-pain blocks each consisted of six 40s thermal stimuli (20s interstimulus interval) delivered at a non-painful temperature (calibrated to each participant's warmth detection threshold), while the pain block consisted of six 40s thermal stimuli delivered at 46°C. The pre-pain/pain/post-pain design has been commonly used in the TMS-MEP pain literature, as many studies have demonstrated strong changes in corticomotor excitability that persist beyond the painful period (for a review see (Chowdhury et al., 2022)). As such, if we had used an alternative design with blocks of warm stimuli intermixed with blocks of painful stimuli, the warm stimuli blocks would not serve as a valid non-painful baseline.

Prior to the test blocks, we measured warmth, cool, and pain detection thresholds to ascertain whether: a) participants could perceive increases or decreases in the thermode temperature relative to a neutral baseline of 32°C, and b) the pain detection threshold was below 46°C. All participants were able to detect increases or decreases in temperature from baseline. The mean (± SD) cool and warmth detection threshold was 28.6 ± 1.9°C and 35.1 ± 1.5°C respectively. All participants reported a heat pain threshold that was above their warmth detection threshold and below the test temperature of 46°C. The mean heat pain threshold was 41.2 ± 2.8°C.

All participants reported 0/10 pain during the pre-pain and post-pain blocks, and pain ratings varying between 1-10 during the pain block. Figure 3 shows the mean pain ratings for the ten pain measurements of each of the six painful stimuli delivered during the pain block (~4 seconds in between pain measurements). A 6 (stimulus number: 1-6) x 10 (timepoint:1-

10) Bayesian repeated measures ANOVA revealed anecdotal evidence (i.e., no conclusive evidence) of a difference in pain between six thermal stimuli ($BF_{10} = 2.86$). However, there was very strong evidence of a difference in pain ratings between ten timepoints ($BF_{10} = 6.1^{30}$). There was also strong evidence of an interaction between stimulus number and timepoint, suggesting the time course of pain across the 40 seconds heat stimulus differed across six thermal stimuli ($BF_{10} = 19.6$). Overall, there was no conclusive evidence for pain differing *between* successive stimuli, however there was evidence that pain fluctuated *during* each 40 second stimulus.

The mean resting motor threshold (RMT) and test intensity of TMS was (mean $\pm$ SD) $70.7 \pm 8.5\%$ and $77.7 \pm 9.2\%$ of maximum stimulus output respectively. Three participants were excluded from the MEP analysis due to EMG software failure – these participants were still included in the TEP analysis. A Bayesian repeated-measures ANOVA was run to compare MEP amplitudes between pre-pain, pain and post-pain conditions. There was anecdotal evidence of a difference in MEP amplitude between blocks ($BF_{10} = 1.015$) (Figure 4A). A Bayesian correlation test was also run to determine whether the mean pain rating (across blocks and timepoints) was associated with the change in MEP amplitude during pain as a proportion of pre-pain. These MEP change values were log-transformed as they were not normally distributed according to a Shapiro-Wilk Test ($W = .572$, $p < .001$). There was strong evidence for a positive relationship ($r_{26} = 0.54$, $BF_{10} = 11.17$) (Figure 4B). such that participants who showed a larger reduction in MEP amplitude during pain reported lower pain ratings.

One participant was excluded from the TEP analysis due to failure to save the recording during the experiment, though this participant was still included in the MEP analysis. One participant had missing post-pain data as the TMS coil had overheated during this portion of the experiment – their data were still included for the pre-pain vs. pain

comparison. Figure 5 shows the grand average TEPs for all 63 channels, across pre-pain, pain and post-pain conditions, as well as the scalp topographies at timepoints where TEP peaks are typically observed – N15, P30, N45, P60, N100 and P180ms (Farzan et al., 2016). Source reconstruction using a co-registered template brain model was also conducted to characterise source activity at each timepoint (Figure 5). For the N15 and P30 peaks, there was higher current density in the left motor areas, consistent with previous studies suggesting that TMS evoked activity at 15 and 30ms after the TMS pulse reflect early excitation of motor areas ipsilateral to the stimulated region (Farzan & Bortoletto, 2022). For the N45 and P60 peaks, there was higher current density in the left motor and somatosensory areas at 45 and 60ms after the TMS pulse, consistent with previous studies showing a sensorimotor origin a these timepoints (Ahn & Fröhlich, 2021), however higher current density was also present in the left parietal and right sensorimotor areas. For the N100 and P180 peaks, there was higher current density in the central regions, mostly contralateral to the stimulated cortex. Overall, we found consistencies in the source localization with previous studies, including a sensorimotor origin of early peaks from 15-60ms. However, we did not directly compare source activity between conditions due to the inaccuracies involved in source estimation in the absence of co-registered magnetic resonance imaging (MRI) scans (Brodbeck et al., 2011; Michel & Brunet, 2019) and EEG electrode location digitization (Shirazi & Huang, 2019).

Comparisons of TEP amplitude between conditions were based on the electrode-space data. However, as this is the first study investigating the effects of experimental pain on TEPs amplitude, there were no *a priori* regions or timepoints of interest to compare between conditions. A statistically robust starting point in these situations is to use a cluster-based permutation analysis (Frömer, Maier, & Abdel Rahman, 2018). This analysis was used to compare amplitudes between pre-pain and pain, and pre-pain and post-pain, at each timepoint and for each electrode. We found that during pain relative to pre-pain, there was a

significantly larger negative amplitude ($p = .021$) at frontocentral electrodes and a significantly larger positive amplitude at parietal-occipital electrodes ($p = .028$), specifically between 43-90ms after the TMS pulse. No significant differences in TEP amplitude were found when comparing the pre-pain and post-pain conditions. As such, the subsequent TEP peak analyses were focused on the pre-pain vs. pain comparison, while the pre-pain vs. post-pain comparisons are presented in the supplementary material.

Figure 6A shows the grand average TEP waveform at the frontocentral electrodes ('AF3','AFz','AF4','F1','Fz','F2','F4','FC2','FC4') identified from the cluster analysis for the pre-pain vs. pain conditions (Supplementary Figure 1 shows the pre-pain vs. post-pain comparison). Two peaks at ~45 and 85ms after the TMS pulse are visible in the time window where the significant cluster was detected. Given the approximate timing, these peaks are likely to be the N45 peak and an early N100 peak. The amplitude of these peaks was identified for each participant using the TESA peak function (Rogasch et al., 2017) with defined time windows of 40-70 and 75-95ms for the first and second peak respectively. These time windows were chosen to account for variation between participants in the latency of the first and second peak. Bayesian paired-sample t-tests showed very strong evidence that the first peak at ~45ms ($BF_{10} = 57.21$) and moderate evidence that the second peak at ~85ms ($BF_{10} = 6.77$) had larger amplitude during pain block compared to pre-pain block. Figure 6B shows the individual level relationship between the mean pain rating, and the difference in N45 and N100 amplitude between pain and pre-pain. There was strong evidence that participants who reported higher pain ratings also showed a larger increase in N45 peak amplitude during pain block ($r_{26} = 0.52$, $BF_{10} = 10.64$). There was anecdotal evidence for no association between pain ratings and changes in the N100 peak amplitude during pain ($r_{26} = 0.24$, $BF_{10} = 0.48$).

Figure 6C shows the mean TEP waveform of the parietal-occipital electrodes ('P1','PO3','O1','CPz','Pz','Pz','Oz','CP2','P2', 'PO4','O2','CP4','P4') identified from the cluster analysis for the pre-pain vs. pain conditions (Supplementary Figure 1 shows the pre-pain vs. post-pain comparison). One peak at ~50ms is visible in the time window where the significant cluster was detected. The approximate timing of this peak is consistent with the commonly identified P60. The amplitude of this peak was identified for each participant with a defined time window of 35 to 65ms. This time window was chosen to account for variation between participants in the latency of the peak. There was moderate evidence that the peak at ~50ms was stronger during pain block compared to pre-pain block ($BF_{10} = 5.56$). Figure 6D shows the individual level relationship between the mean pain rating and the difference in the P60 amplitude between pain and pre-pain. There was anecdotal evidence in favour of no relationship between pain ratings and changes in P60 amplitude during pain block ($r_{26} = 0.21$, $BF_{10} = 0.407$).

Experiment 2 – Does acute pain alter cortical excitability or sensory potentials?

Several studies have shown that a significant portion of TEPs do not reflect the direct cortical response to TMS, but rather auditory potentials elicited by the "clicking" sound from the TMS coil, and somatosensory potentials elicited by the "flicking" sensation on the skin of the scalp (Biabani, Fornito, Mutanen, Morrow, & Rogasch, 2019; Chowdhury et al., 2022; Conde et al., 2019; Rocchi et al., 2021). To separate the direct cortical response to TMS from sensory evoked activity, Experiment 2 (n = 10) included a sham TMS condition that mimicked the auditory/somatosensory aspects of active TMS to determine whether any alterations in the TEP peaks in response to pain were due to changes in sensory evoked activity associated with TMS, as opposed to changes in cortical excitability. A similar design was used to Experiment 1, with the inclusion of a sham TMS condition within the pre-pain and pain blocks, and exclusion of the post-pain block, since the aim was to identify the source

of the pain effect from Experiment 1. The sham TMS condition was similar to a recent study (Gordon et al., 2021), involving the delivery of the TMS coil rotated 90 degrees to the scalp to simulate the auditory component associated with real TMS, and concurrent electrical stimulation beneath a sham coil to simulate the somatosensory component associated with real TMS (see Figure 7A).

Figure 7B shows the grand average TEPs, scalp topographies and estimated source activity for active and sham TMS, across pre-pain and pain conditions. Figure 8 shows the mean TEP waveform for the frontocentral and parietal-occipital clusters identified from Experiment 1, across active and sham conditions. There was moderate evidence that the amplitude of the frontocentral N45 was increased during pain vs. pre-pain blocks for active TMS condition ($BF_{10} = 3.26$), and moderate evidence for no difference between pain and pre-pain blocks for sham TMS condition ($BF_{10} = 0.309$). When comparing pain and pre-pain blocks, there was, respectively, moderate and anecdotal evidence for no alterations in the frontocentral N100 for active ($BF_{10} = 0.31$) and sham TMS ($BF_{10} = 0.42$). There was anecdotal evidence for no alteration in the parietal occipital P60 for both active ($BF_{10} = 0.786$) and sham TMS ($BF_{10} = 0.42$). Overall, the results showed that the N45 peak was altered in response to pain for active but not sham TMS, suggesting the experience of pain led to an alteration in the excitability of the cortex, and not the auditory/somatosensory aspects of TMS.

Experiment 3 – Does acute pain alter cortical excitability or reafferent muscle activity?

Previous studies have shown that a significant portion of the TEP peaks at 45 and 60ms post-TMS reflect reafferent feedback from the muscle twitch in response to suprathreshold TMS applied over M1. This comes from MRI-informed EEG studies showing source localization of the N45 and P60 TEP peaks to the somatosensory areas, as well as correlations between MEP amplitude and N45/P60 amplitude (Ahn & Fröhlich, 2021;

Petrichella, Johnson, & He, 2017). Indeed, Experiments 1 and 2 also showed localization of the N45 and P60 to sensorimotor areas. As such, Experiment 3 recruited a further ten participants to determine whether the pain-induced increase in the N45 peak was due to stronger reafferent feedback from muscle twitches. A design similar to Experiment 1 was used, with the inclusion of a subthreshold TMS condition (90% RMT) within the pre-pain and pain blocks.

Figure 9 shows the grand average TEPs, scalp topographies and estimated source activity for supra- and subthreshold TMS, across pre-pain and pain blocks. Figure 10 shows the mean TEP waveform for the frontocentral and parietal-occipital clusters identified from Experiment 1, across suprathreshold and subthreshold TMS conditions. When comparing the pain with pre-pain blocks, there was moderate evidence that the frontocentral N45 was increased during subthreshold TMS ($BF_{10} = 3.05$) and suprathreshold TMS ($BF_{10} = 3.01$). When comparing pain and pre-pain blocks, there was anecdotal evidence for no alterations in the frontocentral N100 peak during suprathreshold TMS ($BF_{10} = 0.42$) and subthreshold TMS ($BF_{10} = 0.36$). When comparing pain with pre-pain blocks, there was anecdotal evidence for an increase in the parietal occipital P60 peak during suprathreshold TMS ($BF_{10} = 2.71$) and anecdotal evidence for no alteration in P60 peak during subthreshold TMS ($BF_{10} = 0.72$). Overall, there was evidence that that the N45 peak was altered in response to both supra- and subthreshold TMS, suggesting the pain-induced increase in the N45 peak was not a result of stronger reafferent feedback from the muscle twitches.

Discussion

The present study determined whether acute experimental pain induces alterations in cortical inhibitory and/or facilitatory activity observed in TMS-evoked potentials. Across three experiments, there was Bayesian evidence (varying between moderate to very strong) for an increase in the amplitude of N45 peak during painful stimuli compared to a non-painful baseline. Experiment 1 showed very strong evidence that a larger increase in N45 peak in response to pain was correlated with higher pain ratings. Experiment 2 showed that the increase in N45 peak during pain was not a result of alterations in sensory potentials associated with TMS pulses, but rather, changes in cortical excitability. Experiment 3 showed that the increase in N45 peak was not a result of stronger reafferent feedback from muscle twitches evoked by TMS during painful stimuli. While Experiment 1 showed moderate evidence for an increase in N100 and P60 peaks during pain relative to pre-pain baseline, this was not replicated in the follow-up experiments. Experiment 1 showed anecdotal evidence for group-level alterations in MEP amplitude during pain, however there was very strong evidence that a stronger reduction in MEP amplitude during pain was correlated with lower pain ratings.

Increased GABAergic activity during tonic pain

This study is the first to use TMS-EEG methodology to examine the direct cortical response to acute pain, extending previous studies that have used TMS to measure MEPs in response to pain (Chowdhury et al., 2022). The key finding was an increase in the amplitude of N45 peak in response to pain. This result was replicated across three experiments, providing robust evidence for the effect. Furthermore, we also accounted for major confounds that have caused significant data interpretation issues in the TMS-EEG literature in recent years, namely the contamination of TEPs by sensory potentials associated with TMS pulses (Biabani et al., 2019; Chowdhury et al., 2022) and the presence of reafferent feedback from muscle twitches (Ahn & Fröhlich, 2021).

The finding of a reliable increase in the amplitude of N45 peak during pain suggests a role for GABA$_A$ neurotransmission in pain processing, as previous work has shown that the amplitude of N45 peak is increased in response to GABA$_A$ agonists (Premoli et al., 2014). Our source reconstruction results suggest that around this timepoint, the current density was stronger in the sensorimotor area, consistent with the idea that N45 peak reflects GABAergic activity within the sensorimotor cortex (Farzan & Bortoletto, 2022). While it has been shown that reafferent muscle activity also contributes to N45 peak (Ahn & Fröhlich, 2021), Experiment 3 showed that pain increased the amplitude of N45 peak even during subthreshold TMS. Taken together, these findings suggest that the increased amplitude of N45 peak in response to pain reflect a stronger GABAergic activity within the sensorimotor cortex.

GABAergic neurons play a critical role in pain-related brain networks (Barr et al., 2013; Ong, Stohler, & Herr, 2019). They are involved in the generation of gamma oscillations (Buzsáki & Wang, 2012), which have been strongly implicated in pain perception (Barr et al., 2013; Li, Zhang, Zeng, Zhao, & Hu, 2023). Indeed, previous work has shown an increase in gamma oscillations in response to painful thermal stimuli comparable to the present study, across a wide range of brain regions such as the prefrontal (Schulz et al., 2015) and sensorimotor cortices (Gross, Schnitzler, Timmermann, & Ploner, 2007). It is therefore possible that increases in N45 peak during pain reflect increased sensorimotor gamma oscillations. Further multimodal work is required to confirm this finding.

While our findings are consistent with some studies that show increases in GABAergic activity in response to pain (Gross et al., 2007; Kupers, Danielsen, Kehlet, Christensen, & Thomsen, 2009; Schulz et al., 2015), other studies have also reported reduced GABAergic activity in response to experimental pain (Cleve, Gussew, & Reichenbach, 2015; De Matos, Hock, Wyss, Ettlin, & Brügger, 2017). Differences between studies can be

attributed to the duration of the noxious stimulus (tonic pain lasting several seconds/minutes vs. transient pain stimuli lasting <1second). Indeed, pooled data have shown that the cerebral response to pain is highly dependent on the duration of the painful stimuli, as the adaptive response (to suppress or increase cortical activity) changes depending on the duration of pain (Chowdhury et al., 2022). This highlights the need for further work to replicate our findings using different durations of experimental and clinical pain.

Another finding of Experiment 1 was the increase in the amplitude of N100 peak, a marker of $GABA_B$ neurotransmission (Premoli et al., 2014), and the parietal-occipital P60 peak, a marker of glutamatergic neurotransmission (Belardinelli et al., 2021). However, this was not replicated in Experiments 2 and 3, potentially due to the smaller effect size. Nonetheless, we encourage further investigations of alterations in these peaks during pain, particularly P60 peak, as several magnetic resonance spectroscopy studies have reported increases in glutamate concentration during experimental pain (Archibald et al., 2020).

Predicting individual differences in pain using TEPs

Experimental pain models are useful tools to explore brain measures that may predict individual differences in pain sensitivity, with an ultimate goal of determining whether such measures explain why some people develop persistent pain. Experiment 1 showed that higher pain ratings were associated with a larger increase in N45 peak during pain. This analysis was not conducted in Experiments 2 and 3 due to smaller sample sizes and given the primary aims of Experiments 2 and 3 were to isolate the source of the group-level effect. Nonetheless, our results suggest that N45 peak is a potential marker of sensorimotor GABAergic activity and may be associated with individual differences in pain sensitivity. This is consistent with other studies measuring GABAergic responses to pain, showing associations between higher pain sensitivity and stronger sensorimotor gamma oscillations (Barr et al., 2013) and higher left somatosensory cortical GABA laterality (Niddam, Wang, & Tsai, 2021). However, the

direction of this relationship likely depends on the duration of pain (Chowdhury et al., 2022). Our results have implications for understanding the development and maintenance of chronic pain. Further TMS-EEG studies are required to determine whether N45 peak is altered in chronic pain populations and whether N45 peak can explain why some individuals in the acute stages of pain transition to chronic pain.

The TEP vs. MEP response to pain

The present study showed that a larger reduction in MEP amplitude during pain was correlated with lower pain ratings, consistent with a recent systematic review (Chowdhury et al., 2022) and the idea that reduced corticomotor excitability is an adaptive mechanism that restricts movement in the pain-afflicted area, to protect the area from further pain and injury (Hodges & Tucker, 2011). The novelty of this study was the use of an experimental heat pain paradigm that has not yet been used in combination with TMS research, and a paradigm that controls for non-painful somatosensory stimulation.

The finding of a pain-induced increase in the amplitude of N45 peak, which indexes $GABA_A$ receptor activity, is consistent with TMS research showing pain-induced increases in short-interval intracortical inhibition (SICI) (Salo, Vaalto, Koponen, Nieminen, & Ilmoniemi, 2019; Schabrun & Hodges, 2012). SICI refers to the reduction in MEP amplitude to a TMS pulse that is preceded 1-5ms by a subthreshold pulse, due to activation of $GABA_A$ receptors (Kujirai et al., 1993). Some studies have reported associations between SICI and the TEP N45 (Leodori et al., 2019; Rawji, Kaczmarczyk, Rocchi, Rothwell, & Sharma, 2019), suggesting the two may share common neurophysiological mechanisms. However, we also found that a larger reduction in MEP amplitude during pain was associated with less pain, while a larger increase in N45 peak during pain was associated with stronger pain ratings, suggesting that inhibitory processes mediating corticomotor excitability and the N45 peak

during pain are distinct. Further work is required to disentangle the relationship between corticomotor excitability measured by MEPs and cortical activity measured by TEPs.

Study limitations

This study has some methodological limitations. Firstly, while there was no conclusive evidence for a difference in pain ratings between six thermal stimuli, there was evidence for fluctuations in pain ratings during each painful stimulus. This suggests the perceived pain intensity was not stable, which may have introduced noise in the TEP data. Future studies could use pain paradigms that can more effectively maintain a constant level of pain e.g., hypertonic saline infusion paradigms (Svensson, Cairns, Wang, & Arendt-Nielsen, 2003). Secondly, the use of verbal pain ratings prevented the characterisation of pain on a finer time scale. However, verbal ratings were used to eliminate potential contamination of MEPs introduced by using the hand for providing pain rating. Thirdly, while this study attempted to mimic the auditory and somatosensory potentials associated with TMS in the sham condition of Experiment 2, the facial/cranial nerve stimulation associated with active TMS was not accounted for (Mutanen, Mäki, & Ilmoniemi, 2013; Rogasch & Fitzgerald, 2013). However, we note that facial/cranial nerve twitches are more strongly associated with stimulation of lateral brain areas rather than central brain areas (Mutanen et al., 2013). Lastly, the increased N45 peak amplitude in response to pain may reflect increased alertness/arousal during pain. However, a recent study showed that higher alertness is associated with reduced TEP amplitude (Noreika et al., 2020), suggesting the increase in N45 peak amplitude is not related to pain-induced arousal.

Conclusion

This study is the first to use TMS-EEG methodology to examine alterations in cortical activity in direct response to acute pain. Findings across three experiments suggest that tonic heat pain leads to an increase in the amplitude of the frontocentral TEP N45 peak (associated

with GABAergic neurotransmission), and that larger increases in this peak are associated

with higher pain ratings. The findings suggest that TEP indices of GABAergic

neurotransmission have the potential as predictive markers of pain severity.

Materials and Methods

Participants

Experiment 1 consisted of 29 healthy participants (18 males, 11 females, mean age; 26.24 ± 5.5). Participants were excluded if they had any contraindication to TMS such as pregnancy, mental implants in the skull, seizure, or if they reported a history of neurological or psychiatric conditions, or were taking psychoactive medication. Participants completed a TMS safety screen (Rossi, Hallett, Rossini, Pascual-Leone, & Group, 2009). Procedures adhered to the Declaration of Helsinki and were approved by the human research ethics committee of UNSW (HC200328). All participants provided informed written consent.

The sample size calculation was done in G*power 3.1.9.7 with 80% power. As there were no prior TMS-EEG studies, we used pooled data from our systematic review on TMS studies (Chowdhury et al., 2022) showing that the weighted effect size of changes in MEP amplitude in response to tonic experimental pain was 0.56. Using this value, 28 participants were required to detect a significant difference between pain and pre-pain blocks.

To determine the sample size of Experiments 2 and 3, we computed the effect sizes of the N45, P60 and N100 changes (pain vs. prepain) from Experiment 1 (Cohen's d_{RM} = 1.76, 0.99, 0.83 for N45, P60 and N100 respectively). With a power of 80%, the required sample size was 4-11 participants to detect a significant difference. Experiment 2 recruited a further ten healthy participants (four males, six females, mean age: 26.8 ± 5.9) and Experiment 3 consisted of ten healthy participants (four males, six females, mean age: 28 ± 5.9).

Experimental protocol

Participants were seated comfortably in a shielded room. They viewed a fixation cross to minimise eye movements. TMS was applied to left M1 while participants wore an EEG cap containing 63 scalp electrodes to record TEPs. Surface electromyographic (EMG) electrodes were placed over the distal region of the right ECRB to record MEPs. EMG

signals were amplified (x 1000) and filtered (16 to 1000Hz), and digitally sampled at 2000 Hz (Spike2, CED). A thermode was attached over the proximal region of the right ECRB in close proximity to the EMG electrodes (Figure 1).The TMS coil was covered in a layer of foam (5mm thickness) to minimize decay artefacts (Rogasch et al., 2017). Participants also wore both foam earplugs and headphones to reduce any potential discomfort from the TMS click.

The protocol for each experiment is illustrated in Figure 2 (Furman et al., 2020; Granot et al., 2006). In Experiment 1, participants experienced three blocks of heat stimuli, in chronological order: pre-pain, pain, and post-pain block. Each block consisted of multiple heat stimuli delivered 40s at a time during which TMS measurements and verbal pain ratings were obtained. The thermode commenced at a baseline temperature of 32°C. The pre-pain and post-pain blocks consisted of six heat stimuli delivered at the warm threshold (the temperature that led to any detectable change in skin temperature from baseline). In the pain block, six heat stimuli were delivered at 46°C, which has been shown to produce lasting pain with a mean rating of ~5/10 (Furman et al., 2020). The inclusion of blocks with warm stimuli allowed for control for changes in cortical excitability due to non-painful somatosensory stimulation. The protocol for Experiment 2 were identical to Experiment 1 with two differences: the exclusion of the post-pain block (as the aim was to disentangle the source of the pain vs. pre-pain effect from Experiment 1) and the inclusion of a sham TMS condition intermixed within both the pre-pain and pain blocks. The protocol for Experiment 3 was identical to Experiment 1 with two differences: the exclusion of the post-pain block (as the aim was to disentangle the source of the pain vs. pre-pain effect from Experiment 1) and the inclusion of a subthreshold TMS condition intermixed within both the pre-pain and pain blocks.

Electrical stimulation setup (Experiment 2 Only)

Electrical stimulation was used to simulate the somatosensory component of active TMS. Prior to EEG setup, 8mm Ag/AgCl electrodes were placed directly over the scalp. "Snap on" lead wires were then clipped in place and connected to the electrical stimulator (Digitimer DS7AH, Digitimer Ltd., UK). To keep the electrodes and lead wires firmly in position, participants were fitted with a tight netted wig cap, which sat on top of the electrodes but underneath the EEG cap. Consistent with previous research (Chowdhury et al., 2022; Rocchi et al., 2021), and to minimize EEG artefacts caused by electrical stimulation, the stimulating electrodes were not placed directly underneath the EEG electrodes. Rather, stimulating electrodes were positioned in the middle of the EEG electrode cluster located in closest proximity to the motor hotspot. This roughly corresponded to an anode position between FC1 and FC3 and a cathode position between C1 and C3. Scalp electrical stimulation was delivered using a 200 µs square wave via with a compliance of 200V.

Electroencephalography

EEG was recorded using a DC-coupled, TMS-compatible amplifier (ActiChamp Plus, Brain Products, Germany) at a sampling rate of 25000 Hz. Signals were recorded from 63 active electrodes, embedded in an elastic cap (ActiCap, Brain Products, Germany), in line with the 10-10 system. Active electrodes result in similar TEPs (both magnitude and peaks) to more commonly used passive electrodes (Mancuso et al., 2021). Recordings were referenced online to 'FCz' and the ground electrode placed on 'FPz'. Electrolyte gel was used to reduce electrode impedances below ~5kOhms.

Transcranial magnetic stimulation.

Single, monophasic stimuli were delivered using a Magstim unit (Magstim Ltd., UK) and 70mm figure-of-eight flat coil. The coil was oriented at 45° to the midline, inducing a current in the posterior-anterior direction. The scalp site that evoked the largest MEP

measured at the ECRB ('hotspot') was determined and marked. The RMT was determined using the TMS motor thresholding assessment tool, which estimates the lowest TMS intensity required to reliably induce an MEP (Awiszus and Borckardt, 2011), with the MEP amplitude threshold set at 50 microvolts (Rossini et al., 1994; Rothwell et al., 1999). The test stimulus intensity was set at 110% RMT.

Thermal pain

Thermal stimuli were delivered over the proximal region of the right ECRB using a contact heat stimulator (27-mm diameter Medoc Pathway CHEPS Peltier device; Medoc Advanced Medical Systems Ltd). Pain ratings were obtained after each TMS pulse using a verbal rating scale (0 = no pain, and 10 = most pain imaginable). Verbal ratings were collected rather than pain ratings provided on the computer by hand to avoid contamination of MEP measures from motor processes of hand movements. Verbal pain ratings have been shown to yield excellent test-retest reliability (Alghadir, Anwer, Iqbal, & Iqbal, 2018).

Quantitative sensory testing

Quantitative sensory testing was performed in line with a previous study (Furman et al., 2020). With the baseline temperature set at a neutral skin temperature of 32°C, participants completed three threshold tests: to report when they felt a temperature increase (warmth detection threshold), to report when they felt a temperature decrease (cool detection threshold); 3) to report when an increasing temperature first became painful (heat pain threshold). A total of three trials was conducted for each test to obtain an average, with an interstimulus interval of six seconds. Participants provided feedback for each trial by pressing a button (with their left hand) on a hand-held device connected to the Medoc Pathway. Temperatures were applied with a rise/decrease rate of 1°C/s and return rate of 2°C/s (initiated by the button click).

Matching task (Experiment 2 only)

As the aim of Experiment 2 was to perceptually match the somatosensory aspects of active and sham TMS, a 2-Alternative Forced Choice task was used to determine the electrical stimulation intensity that led to a similar flicking sensation to active TMS (Chowdhury et al., 2022). Participants received either electrical stimulation or active TMS in a randomized order and were asked whether the first or second stimulus led to a stronger flick sensation. The electrical stimulation intensity was then increased or decreased until participants could no longer judge the first or second stimulus as stronger. This intensity was then applied during the test blocks.

Test blocks

Experiment 1. The temperature of the thermode commenced at a neutral skin temperature (32°C). Participants were exposed to 18 sustained heat stimuli with a 20 second interstimulus interval. For each stimulus, a single temperature (rise rate of 1 °C/s, return rate of 2 °C/s) was applied over the proximal region of the ECRB for 40 seconds. Stimuli 1-6 (pre-pain block) were delivered at the participant's individually determined warmth detection threshold, Stimuli 7-12 at 46 degrees, and Stimuli 13-18 again at the participant's warmth detection threshold. Participants were not informed of the order of the warm and painful stimuli to minimize the influence of expectation of pain on TEPs and MEPs. During each 40 second stimulus, TMS pulses were manually delivered, with a verbal pain rating score obtained between pulses. As TMS was delivered manually, there was no set interpulse interval. However, the 40 second stimulus duration allowed for 11 pulses for each heat stimulus (hence 66 TMS pulses for each of the pre-pain, pain and post-pain blocks), and 10 verbal pain ratings between each TMS pulse (~ 4 seconds in between pain ratings). A test-retest reliability study by (Kerwin, Keller, Wu, Narayan, & Etkin, 2018) showed that 60 TMS

pulses (delivered in the same run) was sufficient to obtain reliable TEPs (i.e., sufficient within-individual concordance between the resultant TEP peaks of each trial).

Experiment 2. Participants were exposed to 24 sustained heat stimuli (40 seconds each). Stimuli 1-6 and 7-12 consisted of warm stimuli (pre-pain block), while Stimuli 13-18 and 19-24 consisted of stimuli delivered at 46°C (pain block). Active or sham TMS was delivered during either Stimuli 1-6 or 7-12, with the order of active and sham randomly determined for each participant. The same applied for Stimuli 13-18 and 19-24. The active and sham TMS conditions were similar to that used in a recent TMS-EEG study (Gordon et al., 2021). Sham TMS involved the active TMS coil rotated 90 degrees to the scalp, and a sham coil (identical in shape/weight) placed underneath the active coil and tangentially over the scalp. The active TMS coil was then triggered with the electrical stimulation unit to simultaneously simulate the auditory and somatosensory components of active TMS respectively. Active TMS involved the delivery of the TMS coil placed tangentially over the scalp, and the sham TMS coil above the active coil rotated 90 degrees to the scalp (see Figure 9). The design allowed for 11 pulses for each heat stimulus and ten pain ratings (hence 66 TMS pulses for active pre-pain, sham pre-pain, active pain and sham pain blocks).

Experiment 3. Participants were exposed to 24 sustained heat stimuli (40 seconds each). Stimuli 1-6 and 7-12 consisted of warm stimuli (pre-pain block), while Stimuli 13-18 and 19-24 consisted of stimuli delivered at 46°C (pain block). Suprathreshold or subthreshold TMS (90% RMT) was delivered during either Stimuli 1-6 or 7-12, with the order of supra- and subthreshold TMS randomly determined for each participant. The same applied for Stimuli 13-18 and 19-24. The design allowed for 11 pulses for each heat stimulus and ten pain ratings (hence 66 TMS pulses for suprathreshold pre-pain, subthreshold pre-pain, suprathreshold pain and subthreshold pain blocks).

Data processing

Motor evoked potentials. The amplitude of each MEP was determined using a custom MATLAB script. The onsets and offsets of the MEPs were manually determined for each trial. In some participants, background EMG activity was observed due to placement of the thermode close to the EMG electrodes, which can influence MEP amplitude (Ruddy et al., 2018). To account for this, MEP amplitude was calculated by subtracting the root mean square (RMS) of background EMG noise from the RMS of the MEP window using a fixed window between 55 and 5ms before the TMS pulse (Chowdhury et al., 2023; Schabrun et al., 2017; Tsao et al., 2011).

TMS-evoked potentials. Pre-processing of the TEPs was completed using EEGLAB (Delorme & Makeig, 2004) and TESA (Rogasch et al., 2017) in MATLAB (R2021b, The Math works, USA), and based on previously described methods (Chowdhury et al., 2022; Mutanen et al., 2018; Rogasch et al., 2017). First, bad channels were removed. The period between -5 and ~14ms after the TMS pulse was removed and interpolated using the ARFIT function for continuous data (Neumaier & Schneider, 2001; Schneider & Neumaier, 2001). The exact interval was based on the duration of decay artefacts. Data was epoched 1000ms before and after the TMS pulse, and baseline corrected between -1000 and -5ms before the TMS pulse. Noisy epochs were identified via the EEGLAB auto-trial rejection function (Delorme, Sejnowski, & Makeig, 2007) and then visually confirmed. The fastICA algorithm with auto-component rejection used to remove eyeblink and muscle artefacts (Rogasch et al., 2017). The source-estimation noise-discarding (SOUND) algorithm was applied (Mutanen, Biabani, Sarvas, Ilmoniemi, & Rogasch, 2020; Mutanen et al., 2018), which estimates and supresses noise at each channel based on the most likely cortical current distribution given the recording of other channels. This signal was then re-referenced (to average). A band-pass

(1-100Hz) and band-stop (48-52Hz) Butterworth filter was then applied. Any lost channels were interpolated.

Source localization. Source localization of TEPs was conducted using Brainstorm (Tadel, Baillet, Mosher, Pantazis, & Leahy, 2011). A template brain model (ICBM 152) was co-registered with the TMS-EEG data. Noise estimation was used to determine sensor weighting and regularization parameter of the current density construction. The forward model involved use of the Symmetric Boundary Element Method with the head having 3 compartments of fixed conductivities, implemented in OpenMEEG software (Gramfort, Papadopoulo, Olivi, & Clerc, 2010), and inverse model involved use of Minimum Norm Estimations.

Statistical Analysis

Bayesian inference was used to analyse the data using JASP software (Version 0.12.2.0, JASP Team, 2020). Bayes factors were expressed as BF_{10} values, where BF_{10}'s of 1-3, 3-10, 10-30 and 30-100 indicated "weak", "moderate", "strong" and "very strong" evidence for the alternative hypothesis, while BF_{10}'s of 1/3-1, 1/10-1/3, 1/30-1/10 and 1/100-1/30 indicated "anecdotal", "moderate", "strong" and "very strong" evidence in favour of the null hypothesis (van Doorn et al., 2021).

Pain ratings. A 6 (stimulus number: 1-6) x 10 (timepoint:1-10) Bayesian repeated measures ANOVA with default priors in JASP (r scale fixed effects = .5, r scale random effects = 1, r scale covariates = .354) was conducted on the pain ratings during the pain block. This was to assess differences in pain ratings between the six painful stimuli and whether pain ratings differed between the timepoints of each painful stimulus.

MEPs. A Bayesian one-way repeated measures ANOVA with default priors in JASP was performed to assess differences in MEP amplitudes between pre-pain, pain and post-pain blocks of Experiment 1. A Bayesian correlation analysis with default priors in JASP

(stretched Beta prior width = 1) was run to determine whether the change in mean MEP amplitude during pain (as a proportion of pre-pain) was associated with the mean verbal pain rating score. Data were checked for assumptions of normally distributed data using a Shapiro-Wilk test. Where assumptions were violated, data were log-transformed.

    TEPs. The grand-averaged signals for the pre-pain, pain and post-pain condition were obtained. For Experiment 1, a cluster-based permutation analysis was used to compare amplitude levels between pre-pain and pain, and pre-pain and post-pain, at each time-point and for each electrode. For all experiments, the mean TEP waveform of any identified clusters from Experiment 1 were plotted, and peaks were identified using the TESA peak function (Rogasch et al., 2017). Any identified peaks were then compared between conditions using Bayes paired sample t-tests with default priors in JASP (Cauchy scale = .707). A Bayesian correlation analysis with default priors in JASP was performed to determine whether the difference in identified peaks between pre-pain and pain blocks was associated with the mean pain rating score. Data were checked for assumptions of normally distributed data using a Shapiro-Wilk test. Where assumptions were violated, data were log-transformed.

Acknowledgements

This work was supported by 1R61NS113269-01 from The National Institutes of Health to DAS, SMS, and The Four Borders Foundation to DAS.

References

Ahn, S., & Fröhlich, F. (2021). Pinging the brain with transcranial magnetic stimulation reveals cortical reactivity in time and space. *Brain stimulation, 14*(2), 304-315.

Alghadir, A. H., Anwer, S., Iqbal, A., & Iqbal, Z. A. (2018). Test–retest reliability, validity, and minimum detectable change of visual analog, numerical rating, and verbal rating scales for measurement of osteoarthritic knee pain. *Journal of pain research, 11*, 851.

Apkarian, A. V. (2021). The Necessity of Methodological Advances in Pain Research: Challenges and Opportunities. *Frontiers in Pain Research, 2*, 10.

Archibald, J., MacMillan, E. L., Graf, C., Kozlowski, P., Laule, C., & Kramer, J. L. K. (2020). Metabolite activity in the anterior cingulate cortex during a painful stimulus using functional MRS. *Scientific Reports, 10*(1), 19218. doi:10.1038/s41598-020-76263-3

Bank, P. J., Peper, C., Marinus, J., Beek, P. J., & Van Hilten, J. (2013). Motor consequences of experimentally induced limb pain: a systematic review. *European journal of pain, 17*(2), 145-157.

Barr, M. S., Farzan, F., Davis, K. D., Fitzgerald, P. B., & Daskalakis, Z. J. (2013). Measuring GABAergic inhibitory activity with TMS-EEG and its potential clinical application for chronic pain. *Journal of Neuroimmune Pharmacology, 8*(3), 535-546.

Belardinelli, P., König, F., Liang, C., Premoli, I., Desideri, D., Müller-Dahlhaus, F., . . . Ziemann, U. (2021). TMS-EEG signatures of glutamatergic neurotransmission in human cortex. *Scientific Reports, 11*(1), 1-14.

Biabani, M., Fornito, A., Mutanen, T. P., Morrow, J., & Rogasch, N. C. (2019). Characterizing and minimizing the contribution of sensory inputs to TMS-evoked potentials. *Brain stimulation, 12*(6), 1537-1552.

Brodbeck, V., Spinelli, L., Lascano, A. M., Wissmeier, M., Vargas, M.-I., Vulliemoz, S., . . . Seeck, M. (2011). Electroencephalographic source imaging: a prospective study of 152 operated epileptic patients. *Brain, 134*(10), 2887-2897. doi:10.1093/brain/awr243

Burns, E., Chipchase, L. S., & Schabrun, S. M. (2016). Primary sensory and motor cortex function in response to acute muscle pain: A systematic review and meta-analysis. *Eur J Pain, 20*(8), 1203-1213. doi:10.1002/ejp.859

Buzsáki, G., & Wang, X.-J. (2012). Mechanisms of gamma oscillations. *Annual review of neuroscience, 35*, 203.

Cavaleri, R., Chipchase, L. S., Summers, S. J., Chalmers, J., & Schabrun, S. M. (2021). The Relationship Between Corticomotor Reorganization and Acute Pain Severity: A Randomized, Controlled Study Using Rapid Transcranial Magnetic Stimulation Mapping. *Pain Medicine, 22*(6), 1312-1323.

Chang, W.-J., O'Connell, N. E., Beckenkamp, P. R., Alhassani, G., Liston, M. B., & Schabrun, S. M. (2018). Altered primary motor cortex structure, organization, and function in chronic pain: a systematic review and meta-analysis. *The journal of Pain, 19*(4), 341-359.

Chowdhury, N. S., Chang, W.-J., Millard, S. K., Skippen, P., Bilska, K., Seminowicz, D. S., & Schabrn, S. M. (2022). The Effect of Acute and Sustained Pain on Corticomotor Excitability: A Systematic Review and Meta-analysis of group-and individual-level data. *The Journal of Pain*.

Chowdhury, N. S., Rogasch, N. C., Chiang, A. K., Millard, S. K., Skippen, P., Chang, W.-J., . . . Schabrun, S. M. (2022). The influence of sensory potentials on transcranial magnetic stimulation–Electroencephalography recordings. *Clinical Neurophysiology, 140*, 98-109.

Chowdhury, N. S., Skippen, P., Si, E., Chiang, A. K. I., Millard, S. K., Furman, A. J., . . . Seminowicz, D. A. (2023). The reliability of two prospective cortical biomarkers for pain: EEG peak alpha frequency and TMS corticomotor excitability. *J Neurosci Methods, 385*, 109766. doi:10.1016/j.jneumeth.2022.109766

Cleve, M., Gussew, A., & Reichenbach, J. R. (2015). In vivo detection of acute pain-induced changes of GABA+ and Glx in the human brain by using functional 1H MEGA-PRESS MR spectroscopy. *Neuroimage, 105*, 67-75.

Conde, V., Tomasevic, L., Akopian, I., Stanek, K., Saturnino, G. B., Thielscher, A., . . . Siebner, H. R. (2019). The non-transcranial TMS-evoked potential is an inherent source of ambiguity in TMS-EEG studies. *Neuroimage, 185*, 300-312.

De Matos, N. M., Hock, A., Wyss, M., Ettlin, D. A., & Brügger, M. (2017). Neurochemical dynamics of acute orofacial pain in the human trigeminal brainstem nuclear complex. *Neuroimage, 162*, 162-172.

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of neuroscience methods, 134*(1), 9-21.

Delorme, A., Sejnowski, T., & Makeig, S. (2007). Enhanced detection of artifacts in EEG data using higher-order statistics and independent component analysis. *Neuroimage, 34*(4), 1443-1449.

Farzan, F., & Bortoletto, M. (2022). Identification and verification of a'true'TMS evoked potential in TMS-EEG. *Journal of neuroscience methods*, 109651.

Farzan, F., Vernet, M., Shafi, M. M., Rotenberg, A., Daskalakis, Z. J., & Pascual-Leone, A. (2016). Characterizing and modulating brain circuitry through transcranial magnetic stimulation combined with electroencephalography. *Frontiers in neural circuits, 10*, 73.

Frömer, R., Maier, M., & Abdel Rahman, R. (2018). Group-level EEG-processing pipeline for flexible single trial-based analyses including linear mixed models. *Frontiers in neuroscience, 12*, 48.

Furman, A. J., Prokhorenko, M., Keaser, M. L., Zhang, J., Chen, S., Mazaheri, A., & Seminowicz, D. A. (2020). Sensorimotor peak alpha frequency is a reliable biomarker of prolonged pain sensitivity. *Cerebral Cortex, 30*(12), 6069-6082.

Gordon, P. C., Jovellar, D. B., Song, Y., Zrenner, C., Belardinelli, P., Siebner, H. R., & Ziemann, U. (2021). Recording brain responses to TMS of primary motor cortex by EEG–utility of an optimized sham procedure. *Neuroimage*, 118708.

Gramfort, A., Papadopoulo, T., Olivi, E., & Clerc, M. (2010). OpenMEEG: opensource software for quasistatic bioelectromagnetics. *Biomedical engineering online, 9*(1), 1-20.

Granot, M., Granovsky, Y., Sprecher, E., Nir, R.-R., & Yarnitsky, D. (2006). Contact heat-evoked temporal summation: tonic versus repetitive-phasic stimulation. *Pain, 122*(3), 295-305.

Gross, J., Schnitzler, A., Timmermann, L., & Ploner, M. (2007). Gamma oscillations in human primary somatosensory cortex reflect pain perception. *PLoS biology, 5*(5), e133.

Hodges, P. W., & Tucker, K. (2011). Moving differently in pain: A new theory to explain the adaptation to pain. *Pain, 152*(SUPPL.3), S90-S98. doi:10.1016/j.pain.2010.10.020

Kallioniemi, E., & Daskalakis, Z. J. (2022). Identifying novel biomarkers with TMS-EEG - Methodological possibilities and challenges. *J Neurosci Methods, 377*, 109631. doi:10.1016/j.jneumeth.2022.109631

Kerwin, L. J., Keller, C. J., Wu, W., Narayan, M., & Etkin, A. (2018). Test-retest reliability of transcranial magnetic stimulation EEG evoked potentials. *Brain stimulation, 11*(3), 536-544. doi:https://doi.org/10.1016/j.brs.2017.12.010

Kujirai, T., Caramia, M., Rothwell, J. C., Day, B., Thompson, P., Ferbert, A., . . . Marsden, C. D. (1993). Corticocortical inhibition in human motor cortex. *The Journal of Physiology, 471*(1), 501-519.

Kupers, R., Danielsen, E. R., Kehlet, H., Christensen, R., & Thomsen, C. (2009). Painful tonic heat stimulation induces GABA accumulation in the prefrontal cortex in man. *Pain, 142*(1-2), 89-93.

Leodori, G., Thirugnanasambandam, N., Conn, H., Popa, T., Berardelli, A., & Hallett, M. (2019). Intracortical inhibition and surround inhibition in the motor cortex: a TMS-EEG study. *Frontiers in neuroscience, 13*, 612.

Li, Z., Zhang, L., Zeng, Y., Zhao, Q., & Hu, L. (2023). Gamma-band oscillations of pain and nociception: A systematic review and meta-analysis of human and rodent studies. *Neuroscience & Biobehavioral Reviews, 146*, 105062. doi:https://doi.org/10.1016/j.neubiorev.2023.105062

Mancuso, M., Sveva, V., Cruciani, A., Brown, K., Ibáñez, J., Rawji, V., . . . Rothwell, J. (2021). Transcranial evoked potentials can be reliably recorded with active electrodes. *Brain Sciences, 11*(2), 145.

Michel, C. M., & Brunet, D. (2019). EEG source imaging: a practical review of the analysis steps. *Frontiers in Neurology, 10*, 325.

Mutanen, T., Mäki, H., & Ilmoniemi, R. J. (2013). The effect of stimulus parameters on TMS–EEG muscle artifacts. *Brain stimulation, 6*(3), 371-376.

Mutanen, T. P., Biabani, M., Sarvas, J., Ilmoniemi, R. J., & Rogasch, N. C. (2020). Source-based artifact-rejection techniques available in TESA, an open-source TMS–EEG toolbox. *Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation, 13*(5), 1349-1351.

Mutanen, T. P., Metsomaa, J., Liljander, S., & Ilmoniemi, R. J. (2018). Automatic and robust noise suppression in EEG and MEG: The SOUND algorithm. *Neuroimage, 166*, 135-151.

Neumaier, A., & Schneider, T. (2001). Estimation of parameters and eigenmodes of multivariate autoregressive models. *ACM Transactions on Mathematical Software (TOMS), 27*(1), 27-57.

Niddam, D. M., Wang, S.-J., & Tsai, S.-Y. (2021). Pain sensitivity and the primary sensorimotor cortices: a multimodal neuroimaging study. *Pain, 162*(3), 846-855.

Ong, W.-Y., Stohler, C. S., & Herr, D. R. (2019). Role of the Prefrontal Cortex in Pain Processing. *Molecular Neurobiology, 56*(2), 1137-1166. doi:10.1007/s12035-018-1130-9

Petrichella, S., Johnson, N., & He, B. (2017). The influence of corticospinal activity on TMS-evoked activity and connectivity in healthy subjects: A TMS-EEG study. *PloS one, 12*(4), e0174879.

Ploner, M., & May, E. S. (2018). Electroencephalography and magnetoencephalography in pain research—current state and future perspectives. *Pain, 159*(2), 206-211.

Premoli, I., Castellanos, N., Rivolta, D., Belardinelli, P., Bajo, R., Zipser, C., . . . Ziemann, U. (2014). TMS-EEG signatures of GABAergic neurotransmission in the human cortex. *Journal of Neuroscience, 34*(16), 5603-5612.

Rawji, V., Kaczmarczyk, I., Rocchi, L., Rothwell, J. C., & Sharma, N. (2019). Short interval intracortical inhibition as measured by TMS-EEG. *BioRxiv*, 802504.

Reddan, M. C., & Wager, T. D. (2018). Modeling pain using fMRI: from regions to biomarkers. *Neuroscience bulletin, 34*(1), 208-215.

Rocchi, L., Di Santo, A., Brown, K., Ibáñez, J., Casula, E., Rawji, V., . . . Rothwell, J. (2021). Disentangling EEG responses to TMS due to cortical and peripheral activations. *Brain stimulation, 14*(1), 4-18.

Rogasch, N. C., & Fitzgerald, P. B. (2013). Assessing cortical network properties using TMS–EEG. *Human brain mapping, 34*(7), 1652-1669.

Rogasch, N. C., Sullivan, C., Thomson, R. H., Rose, N. S., Bailey, N. W., Fitzgerald, P. B., . . . Hernandez-Pavon, J. C. (2017). Analysing concurrent transcranial magnetic stimulation and electroencephalographic data: A review and introduction to the open-source TESA software. *Neuroimage, 147*, 934-951.

Rohel, A., Bouffard, J., Patricio, P., Mavromatis, N., Billot, M., Roy, J.-S., . . . Masse-Alarie, H. (2021). *The effect of experimental pain on the excitability of the corticospinal tract in humans: A systematic review and meta-analysis*: European Journal of Pain. 2021, pp. No Pagination Specified.

Rossi, S., Hallett, M., Rossini, P. M., Pascual-Leone, A., & Group, S. o. T. C. (2009). Safety, ethical considerations, and application guidelines for the use of transcranial magnetic stimulation in clinical practice and research. *Clinical Neurophysiology, 120*(12), 2008-2039.

Rossini, P. M., Barker, A., Berardelli, A., Caramia, M., Caruso, G., Cracco, R., . . . Lücking, C. (1994). Non-invasive electrical and magnetic stimulation of the brain, spinal cord and roots: basic

principles and procedures for routine clinical application. Report of an IFCN committee. *Electroencephalography and clinical Neurophysiology, 91*(2), 79-92.

Rothwell, J. C., Hallett, M., Berardelli, A., Eisen, A., Rossini, P., & Paulus, W. (1999). Magnetic stimulation: motor evoked potentials. The International Federation of Clinical Neurophysiology. *Electroencephalography and clinical neurophysiology. Supplement, 52*, 97-103.

Ruddy, K., Balsters, J., Mantini, D., Liu, Q., Kassraian-Fard, P., Enz, N., . . . Wenderoth, N. (2018). Neural activity related to volitional regulation of cortical excitability. *Elife, 7*.

Salo, K. S.-T., Vaalto, S. M., Koponen, L. M., Nieminen, J. O., & Ilmoniemi, R. J. (2019). The effect of experimental pain on short-interval intracortical inhibition with multi-locus transcranial magnetic stimulation. *Experimental Brain Research, 237*(6), 1503-1510.

Schabrun, S. M., & Hodges, P. W. (2012). Muscle pain differentially modulates short interval intracortical inhibition and intracortical facilitation in primary motor cortex. *J Pain, 13*(2), 187-194. doi:10.1016/j.jpain.2011.10.013

Schneider, T., & Neumaier, A. (2001). Algorithm 808: ARfit—A Matlab package for the estimation of parameters and eigenmodes of multivariate autoregressive models. *ACM Transactions on Mathematical Software (TOMS), 27*(1), 58-65.

Schulz, E., May, E. S., Postorino, M., Tiemann, L., Nickel, M. M., Witkovsky, V., . . . Ploner, M. (2015). Prefrontal gamma oscillations encode tonic pain in humans. *Cerebral cortex, 25*(11), 4407-4414.

Shirazi, S. Y., & Huang, H. J. (2019). More reliable EEG electrode digitizing methods can reduce source estimation uncertainty, but current methods already accurately identify brodmann areas. *Frontiers in neuroscience, 13*, 1159.

Svensson, P., Cairns, B. E., Wang, K., & Arendt-Nielsen, L. (2003). Injection of nerve growth factor into human masseter muscle evokes long-lasting mechanical allodynia and hyperalgesia. *Pain, 104*(1-2), 241-247.

Tadel, F., Baillet, S., Mosher, J. C., Pantazis, D., & Leahy, R. M. (2011). Brainstorm: a user-friendly application for MEG/EEG analysis. *Computational Intelligence and Neuroscience, 2011*.

Tremblay, S., Rogasch, N. C., Premoli, I., Blumberger, D. M., Casarotto, S., Chen, R., . . . Fitzgerald, P. B. (2019). Clinical utility and prospective of TMS–EEG. *Clinical Neurophysiology, 130*(5), 802-844.

van Doorn, J., van den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., . . . Haaf, J. M. (2021). The JASP guidelines for conducting and reporting a Bayesian analysis. *Psychonomic Bulletin & Review, 28*, 813-826.

Figures

Figure 1. **Schematic of experimental apparatus.** This consisted of transcranial magnetic stimulation (TMS) during concurrent electroencephalography (EEG) to simultaneously record motor evoked potentials (MEPs) and TMS-evoked potentials (TEPs). MEPs were recorded using electromyographic (EMG) electrodes placed over the distal region of the extensor carpi radialis brevis (ECRB) while thermal pain was delivered over the proximal region of the ECRB.

Figure 2. **Schematic of the experimental protocols.** In Experiment 1, participants experienced three blocks of stimuli: a pre-pain, pain, and post-pain block, with each block consisting of multiple heat stimuli delivered 40s at a time, and during which TMS measurements (indicated by blue arrows) and verbal pain ratings were obtained. The pre-pain and post-pain blocks involved thermal stimuli delivered at the warm threshold (i.e., the temperature that leads to any perceived change in skin temperature from baseline). In the pain block, thermal stimuli were delivered at 46°C. For Experiment 2, the post-pain block was excluded, and an additional sham TMS condition was intermixed within both the pre-pain and pain blocks. For Experiment 3, the post-pain block was also excluded, and an additional subthreshold TMS condition intermixed within both the pre-pain and pain blocks.

Figure 3. **No conclusive evidence of a difference in pain ratings between successive 46°C 40s thermal stimuli.** Mean (± SD) pain ratings during the 6 thermal stimuli delivered during the pain block (heat stimuli delivered at 46°C) of Experiment 1. 10 pain ratings were collected over each 40 second stimulus ~ every 4 second.

Figure 4. **No conclusive evidence of MEP amplitude differences between conditions; however individual pain sensitivity was predicted by changes in MEP amplitude.** A: Mean (± SD) MEP amplitude during the pre-pain, pain and post-pain blocks of Experiment 1. B: Individual-level Relationship between change in MEP amplitude during pain (proportion of pre-pain) and mean verbal pain rating provided by each participant.

Figure 5. **Pain led to increased negative and positive amplitude in frontocentral and parietal-occipital sites respectively, 43-90ms after the TMS pulse.** A: Grand-average TEPs during the pre-pain, pain and post-pain blocks of Experiment 1. The grey shaded area represents the window of interpolation around the TMS pulse. B: Scalp topographies and estimated source activity at timepoints where TEP peaks are commonly observed, including the N15, P30, N45, P60, N100 and P180. A cluster plot is also shown on the right comparing signal amplitude between the pain and pre-pain conditions at a representative timepoint (48ms) between 43-90ms, which is where significant amplitude differences were observed. The black stars demonstrate the presence of significant positive (yellow) or negative (blue) clusters.

Figure 6. **Pain led to increases in N45, P60 and N100 peak amplitude, and individual pain sensitivity was predicted by changes in the N45 peak.** A: TEPs across pain and pre-pain condition for the frontocentral electrodes and parietal-occipital electrodes identified from the cluster analysis of Experiment 1. The grey shaded area represents the window of interpolation around the transcranial magnetic stimulation (TMS) pulse. For the frontocentral electrodes, two significantly stronger negative peaks were identified at ~45 and 85ms post-TMS. For the parietal-occipital electrodes, a significantly stronger positive peak was identified at ~50ms post-TMS. * indicates at least moderate evidence the alternative hypothesis that the amplitude is larger in pain vs. pre-pain ($BF_{10} > 3$). B: Individual-level relationship between change in peak amplitudes at ~45ms (N45), ~50ms (P60) and ~85ms (N100) post-TMS, and mean verbal pain ratings provided by each participant.

Figure 7. **TMS-evoked potentials for Active and Sham TMS.** A: Schematics showing the delivery of active and sham TMS of Experiment 2. Sham TMS involved scalp electrical stimulation (in red) beneath a sham coil (in dotted blue) to mimic somatosensory stimulation associated with active TMS, and concurrent delivery of active TMS 90 degrees to the scalp (in shaded blue) to mimic auditory stimulation associated with TMS. B: Left: TEPs during the pre-pain and pain blocks, for both active and sham stimulation. The grey shaded area represents the window of interpolation around the TMS pulse. Right: Scalp topographies and estimated source activity at timepoints where TEP peaks are commonly observed, including the N15, P30, N45, P60, N100 and P180.

Figure 8. **Pain led to an increase in the N45 peak amplitude during active TMS but not sham TMS.** TEPs during pain and pre-pain blocks, across active and sham TMS conditions of Experiment 2, for the frontocentral electrodes (left) and parietal occipital electrodes (right) identified from the cluster analysis in the main experiment. A significantly stronger frontocentral negative peak was identified ~45ms post-TMS during pain compared to pre-pain, for the active TMS condition. * indicates at least moderate evidence the alternative hypothesis that the amplitude is larger in pain vs. pre-pain ($BF_{10} > 3$). The dotted line shows the timing of the peak.

Figure 9. **TMS-evoked potentials for supra- and subthreshold TMS.** Left: TEPs during the pre-pain and pain blocks, for both supra- and subthreshold TMS of Experiment 3. The grey shaded area represents the window of interpolation around the transcranial magnetic stimulation TMS pulse. Right: Scalp topographies and estimated source activity at timepoints where TEP peaks are commonly observed, including the N15, P30, N45, P60, N100 and P180.

Figure 10. **Pain led to an increase in the N45 peak amplitude for both suprathreshold and subthreshold TMS.** TEPs during pain and pre-pain blocks of Experiment 3, across supra- and subthreshold TMS conditions, for the frontocentral electrodes (left) and parietal occipital electrodes (right) identified from the cluster analysis in Experiment 1. A significantly stronger frontocentral negative peak was identified ~45ms post-TMS during pain compared to pre-pain for both supra- and subthreshold stimulation. * indicates at least moderate evidence the alternative hypothesis that the amplitude is larger in pain vs. pre-pain ($BF_{10} > 3$). The dotted line shows the timing of the peak.