Lean female mice producing *trans* 10, *cis* 12-conjugated linoleic acid present excessive fibroblast growth factor 21, adrenaline, prostaglandin E2, corticosterone, and glucagon

Yu Rao[1,2, #], Lu-Wen Liang[1, #], Mei-Juan Li[3, #], Yang-Yang Wang[1], Bao-Zhu Wang[1], Ke-Mian Gou[1,2, *]

[1]*Institute of Comparative Medicine,* [2]*Jiangsu Co-innovation Center for Prevention and Control of Important Animal Infectious Diseases and Zoonoses, Department of Experimental Zoology, College of Veterinary Medicine, Yangzhou University, Yangzhou 225009, China;*

[3]*Institute of Animal Husbandry and Veterinary Science, Guizhou Academy of Agricultural Sciences, Guiyang 550005, China.*

# These authors contributed equally to this work.

*Corresponding author: Kemian Gou, College of Veterinary Medicine, Yangzhou University, Wenhui South Road, Yangzhou 225009, China.

Email: gou@yzu.edu.cn

ABSTRACT

Dietary *trans* 10, *cis* 12-conjugated linoleic acid (t10c12-CLA) has been shown to
induce white fat reduction. In this study, the transgenic mice that produce endogenous
t10c12-CLA were used to determine the long-term impact of this substance on lipid
metabolism in lean female mammals. Phenotype profiling showed that the t10c12-CLA did
not induce fat loss but resulted in less heat release and oxygen consumption, as well as
excessive adrenaline, tumour necrosis factor-alpha, and interleukin-6 in either biallelic
Pai/Pai or monoallelic Pai/wt mice compared to their wild-type littermates. Furthermore,
the following effects of t10c12-CLA were genotype-specific. Pai/wt mice specifically
exhibited increased chronic fibroblast growth factor 21 (FGF21) and reduced physical
activities. Contrarily, Pai/Pai mice showed the overproduction of prostaglandin E2 (PGE2),
corticosterone, and glucagon, accompanied by hypertriglyceridemia and fatty liver. Further
analysis of Pai/Pai livers revealed that the steatosis resulted from attenuated lipid
metabolism via down-regulating phosphorylated AMP-activated protein kinase (AMPK)
and fatty acid synthase. Analysis of Pai/Pai brown adipose tissue indicated stimulated
thermogenesis and lipolysis by up-regulating uncoupling proteins 1 and 2, phosphorylated
hormone-sensitive lipase, carnitine palmitoyltransferase 1B, and AMPK, as well as down-
regulating phosphorylated AMPK. These results suggest that chronic t10c12-CLA may affect
lipid metabolism by stimulating the overproduction of multiple hormones (FGF21, PEG2,
adrenaline, etc.) in a dose-dependent manner in lean female mammals. It also suggests that
t10c12-CLA-induced fatty liver, inflammation, hypothermia and excess hormones are major
health concerns.

Keywords: conjugated linoleic acid; prostaglandin E2; steatosis; adrenaline;
inflammation; brown adipose tissue

1. INTRODUCTION

Trans 10, *cis* 12-conjugated linoleic acid (t10c12-CLA) occurs naturally in small amounts in foods such as beef and dairy products. Its popular health benefit is reducing body fat by increasing energy expenditure via browning of white adipose tissue (WAT) [1,2] or uncoupling protein (UCP) 1 & 2-mediated thermogenesis in brown adipose tissue (BAT) [3], promoting biosynthesis of N-lactoyl-phenylalanine via the up-regulation of cytosolic nonspecific dipeptidase 2 in WAT [4], or increasing intramuscular fat [5]. In addition, it has other potential health influences, including protecting against atherosclerosis by modulating the macrophage phenotype [6], promoting beneficial effects on cholesterol efflux [7], reducing oxidative stress [8], changing intestinal microbiota [9] and metabolite profiles [5], even down-regulating Alzheimer's hallmarks in an aluminium mouse through an Nrf2-mediated adaptive response and increasing brain glucose transporter levels [10].

Although the potential health benefits, especially the fat loss properties of dietary t10c12-CLA, have been extensively studied in obese animals, the potential health benefits in lean animals have not been clarified fully. A study in lean male mice treated with dietary t10c12-CLA observed a dramatic reduction in fat mass in the first two weeks which interestingly rebounded during the following 3-5 weeks due to a massive accumulation of lipids in the liver [11]. At the same time, other studies have confirmed its fat loss properties but indicated its side effects result in fatty liver in male [9,11] or female lean mice [12-16].

In our previous study, we produced a transgenic mouse that can produce t10c12-CLA after inserting the foreign gene encoding the *Propionibacterium acnes* isomerase (PAI) into the Rosa26 locus [3]. This Rosa26-Pai knock-in mouse exhibited continuous production of t10c12-CLA and can be used to elucidate its enduring influence on overall health. In its lean male offspring, fat reduction in biallelic Pai/Pai, not in monoallelic Pai/wt genotypes suggests t10c12-CLA's impact is dose-dependent. Additionally, low doses of t10c12-CLA possibly play an active role in lipid metabolism by stimulating the fibroblast growth factor 21 (FGF21) secretion. Whereas the high doses of t10c12-CLA possibly induce fat reduction by promoting energy expenditure via UCP1/2-mediated BAT thermogenesis [3].

The current study continued to investigate the adiposity characteristics and metabolic profiles of these lean female mice. In contrast to their male counterparts, the findings of this study demonstrate that Pai female mice did not experience changes in body fat but did experience a decrease in heat release. Moreover, specific metabolic features in

Pai/wt female mice have confirmed that the low doses of t10c12-CLA can induce the
overproduction of FGF21, while the high doses of t10c12-CLA stimulate excessive
prostaglandin E2 (PGE2), corticosterone, and glucagon which probably affect the lipid
metabolism via varies of pathways. The results suggest that Pai mice may be a useful model
to investigate the chronic t10c12-CLA's effects and its mechanisms of action in the body. It
also indicates that t10c12-CLA-induced fatty liver, inflammation, hypothermia and excess
hormones are major health concerns.

2. MATERIALS AND METHODS

2. 1. Mice and diet

The transgenic founder from a hybrid embryo of DBA/2 and C57BL/6J was backcrossed to C57BL/6J mice for more than eight generations, and then the monoallelic Pai/wt matings were used to produce the Pai/Pai, Pai/wt, and wild-type (wt) offspring for this study. Genotyping was performed using genomic DNA from tail tissues, with the presence of the Pai gene being assayed by amplifying the 514-bp Pai fragments (forward: 5-taaccatgttcatgccttcttc-3; reverse: 5-caccttgttgtagtgtccgttt-3) and the wt Rosa26 sequence being assayed by amplifying the 605-bp fragments that spanned the insertional site of the Rosa26 locus (forward: 5-ccaaagtcgctctgagttgttatcagt-3; reverse: 5-ggagcgggagaaatggatatgaag-3). PCR amplification was performed for 30 cycles at 94°C for 30 sec, 57°C for 30 sec, and 72°C for 40 sec.

Table 1. Sources of dietary ingredients

Ingredients	Grams	Sources
Lactic casein	200.0	Gansu Hualing Dairy Co. Ltd, Hezuo city, China
L-cystine	3.0	Zhongtuan shihua Food Reagents Co. Ltd, Hangzhou, China
Sucrose	176.8	Jinhuitaiya, Chemcal Reagents Co. Ltd, Tianjin, China
Corn Starch	452.2	Zhucheng Xingmao Corn Co. Ltd, Weifang city, China
Maltodextrin 10	75.0	Dongzhu Food Co. Ltd, Guilin city, China
Cellulose	50.0	Yisheng Kangyuan Biotech Co. Ltd, Zibo city, China
Soybean oil	25.0	Shandong Luhua Co. Ltd, Leiyang, China
Lard	20.0	Yaodong Baofengyuan Co. Ltd, Qingdao, China
Mineral Mix (20x)	50.0	Mingxin Chemical Reagents Co. Ltd, Zhenzhou, China
Choline bitartrate	2.0	Zhejiang Linuo Biotech Co. Ltd, Yiwu, China
Vitamin Mix (1000x)	1.0	Mingxin Chemical Reagents Co. Ltd, Zhenzhou, China
Total Gram	1055	

Mice of the above three genotypes aged 11-15 weeks were used for analysis in this study unless otherwise specified. All animal experiments were conducted in accordance with the guidance provided by the Committee for Experimental Animals of Yangzhou University. The animal study protocol was approved by the Ethics Committee of Yangzhou University (protocol code NSFC2020-SYXY-20 and dated 25 March 2020). The mice were housed in a

light-controlled room (12L:12D, lights on at 0700 h) at a temperature of 22-23°C and fed ad
libitum with a standard diet containing 10% kcal% fat. Fresh diets were prepared every
month in accordance with formula No. D12450H of OpenSource DIETS™ (Research Diets
Inc., NJ, USA) under sterile conditions. The composition of the D12450H and the sources of
the food-grade ingredients used in the diets are provided in Table 1.

2. 2. Major methods

The procedures utilised in this study, including gas chromatography, real-time RNA
and proteins analyses, histological analysis, intraperitoneal glucose (2g/kg body weight) and
insulin (0.5 IU/kg body weight) tolerance tests, hepatic and blood parameter measurements,
and experiments conducted in metabolic cages, were performed following the standard
methods as described in our previous study of transgenic male mice [3].

2.3. Measurement of serum estradiol

Three-week-old mice were injected intraperitoneally with 5 IU pregnant mare serum
gonadotrophin (Jinmu Biotech Co. Ltd., Wuxi, China), and 48 h later, blood was collected
from the orbital vein of anaesthetised mice and used to measure the serum estradiol levels
using the E2 ELISA kit (Meimian Industrial Co. Ltd., Jiangsu, China).

2. 4. Statistical analysis.

Analysis was conducted using GraphPad Prism 8.0 (GraphPad Software Inc.; La
Jolla, CA, USA). Comparisons between wt and Pai/Pai samples were performed using an
unpaired two-tailed t-test with Welch's correction where appropriate. Multiple comparisons
among wt, Pai/wt, and Pai/Pai groups were performed using one-way ANOVA and Brown-
Forsythe and Welch tests. Data are presented as mean ± standard deviation (SD), and $p <$
0.05 was considered statistically significant.

3. RESULTS

3. 1. Fatty acids analysis

Gas chromatography was used to analyse the compositions of fatty acids (FAs) in the livers, kidneys, hearts, tibialis anterior muscle, and interscapular BAT. The content of t10c12-CLA had increased by 69% in the Pai/wt kidneys compared to wt littermates ($p < 0.05$; Table 2). However, the quantities of substrate linoleic acid had decreased in the Pai/Pai livers by 23% and skeletal muscle by 34% and had increased by 36% in the Pai/wt kidneys ($p < 0.05$; Supplementary Table 1).

The contents of other FAs such as myristic (14:0), palmitic (16:0), stearic (18:0), palmitoleic (16:1n-7), cis-vaccenic (18:1n-7), oleic (18:1n-9), arachidonic (20:4n-6), and docosahexaenoic (22:6n-3) acids were changed to varying degrees in one or more tissues (Supplementary Table 1), This suggests that the changes in the content of each FA were genotype- or tissue-specific. Additionally, the amounts of total FAs (mg/g) increased in the Pai/wt kidneys (10.2 ± 1.39 versus 13.3 ± 1.69; $p < 0.05$), similar to our previous study in Pai/wt male mice [3]. These results suggest lipopathy in the Pai/wt kidney.

Table 2. Content comparison of t10c12-CLA (μg/g) in wild-type and Pai mice

	wild-type (n)	Pai/wt (n)	Pai/Pai (n)
Liver	2.56 ± 3.24 (7)	4.30 ± 4.10 (5)	ND (5)
Kidney	6.50 ± 1.25[a] (5)	11.0 ± 3.65[b] (6)	6.09 ± 0.42[a] (4)
Heart	ND (5)	ND (4)	ND (4)
Muscle	ND (4)	ND (4)	ND (4)
Brown adipose tissue	22.4 ± 15.9 (4)	29.5 ± 7.5 (4)	29.8 ± 0.5 (4)

Note: Tissues from 11-week-old mice was analysed by gas chromatography and 19:0 is an internal standard. ND, not detected. Each value represents the mean ± SD and the numbers in brackets indicate the size of the tested samples. Different letters in superscript indicate a level of statistical significance of $p < 0.05$.

3. 2. Growth analysis

There was a significant difference in weaning weight among the three genotypes (wt > Pai/wt > Pai/Pai, respectively; $p < 0.05$; Fig. 1), similar to our previous study in male mice [3]. However, this bodyweight difference disappeared from five weeks of age onwards.

Figure 1. Comparison of growth of wild-type and Pai mice. Each group contained 11-29 mice. Bars represent the mean ± SD. # indicates $p < 0.05$ among three genotypes.

3. 3. Energy metabolism and activity studies

Energy metabolism and activity parameters were assessed in consideration of body weight. Compared to their wt littermates, both Pai/wt and Pai/Pai mice exhibited no significant differences in body weight gain, food and water intake (Fig. 2 A-C), total distance travelled during the whole 72-h observation (Fig. 2 D), and respiratory exchange ratio (Fig. 2 E-F). However, the other parameters showed varying differences.

The Pai/Pai mice covered a longer distance per sampling duration (~50% longer) and exhibited 52.5% higher speed during the light phase ($p < 0.05$; Fig. 2 G-H) but did not decrease their total locomotor activities in either the centre (Fig. 2 I-L) or margin/corners (Fig. 2 M-P). Nevertheless, during both the light ($p < 0.05$) and dark ($p > 0.05$) periods, they consumed 8.9% and 9.2% less O_2 (Fig. 2 Q) and produced 10.8% and 10.1% less CO_2 (Fig. 2 R) as well as 9.3% and 9.4% less heat (Fig. 2 S-T), respectively. The results suggest that Pai/Pai mice simultaneously reduce oxygen consumption and heat release while maintaining normal locomotor activities.

On the other hand, Pai/wt females did not reduce their central activities ($p > 0.05$; Fig. 2 I-L) but displayed 30.1% fewer marginal activities, such as marginal ambulating and stereotypic activities during the dark period ($p < 0.05$; Fig. 2 M-P). Moreover, during both the light and dark periods, they consumed 10.7% and 14.6% less O_2, produced 9.7% and 15.2% less CO_2, as well as 10.5% and 14.7% less heat, respectively ($p < 0.05$, Fig. 2 Q-T), suggesting that Pai/wt mice simultaneously reduce oxygen consumption and heat release while exhibiting fewer marginal activities.

Figure 2. Abnormal energy metabolism and activities in Pai mice. Comparison of body weight gain during a 72-h period (A), food intake (B), water intake (C), total distance travelled during a 72-h period (D), respiratory exchange ratio (RER) and RER time course (E-F), distance travelled during a 39-min sampling duration (G), speed (H), locomotor activities in the centre of the cage and its time course (I-J), central ambulating and stereotypic activities

(K-L), locomotor activities in the margin or corners of the cage and its time course (M-N),

marginal ambulating and stereotypic activities (O-P), O_2 consumption (Q), CO_2 production

(R), heat release and its time course (S-T) among wild-type (n = 4), Pai/wt (n = 4) and Pai/Pai

(n = 4) mice at 11 weeks. The lights turned on and off at 7:00 am and 7:00 pm, respectively.

The data are normalised to lean body weight. The bars represent the mean ± SD. Different

letters indicate $p < 0.05$ within the same phase.

3. 4. Metabolic complications

Concentrations of circulating glucose, insulin, leptin, and ghrelin (Table 3), as well as

small intestine length (Supplementary Table 2), showed no significant differences ($p > 0.05$)

between Pai mice and their wt littermates. This suggests that t10c12-CLA did not have any

significant impact on energy intake, which is consistent with the observation in the metabolic

cage.

Table 3. Circulating factors in wild-type and Pai mice

	wild-type (n)	Pai/wt (n)	Pai/Pai (n)
Blood glucose (mM)	7.35 ± 0.70 (12)	7.37 ± 1.18 (15)	7.55 ± 1.35 (13)
Serum insulin (μg/L)	0.21 ± 0.08 (9)	0.27 ± 0.11 (17)	0.32 ± 0.18 (8)
Serum leptin (μg/L)	1.04 ± 0.41 (8)	0.92 ± 0.21 (14)	0.91 ± 0.31 (7)
Serum ghrelin (ng/L)	181 ± 44 (7)	225 ± 108 (20)	210 ± 152 (6)
Serum fibroblast growth factor 21 (ng/L)	403 ± 29[a] (16)	441 ± 49[b] (10)	421 ± 34[ab] (18)
Serum total cholesterol (μM)	311 ± 40 (17)	324 ± 40 (8)	321 ± 30 (16)
Plasma triglycerides (mg/dL)	107 ± 35[a] (34)	105 ± 27[a] (29)	129 ± 42[b] (29)
Plasma high-density lipoprotein (mg/dL)	48.6 ± 13.9 (34)	47.7 ± 16.2 (29)	52.5 ± 19.6 (29)
Serum free fatty acids (μM)	471 ± 95 (8)	527 ± 79 (16)	517 ± 136 (8)
Serum Estradiol (pM)	71.0 ± 8.8 (15)	64.9 ± 6.8 (7)	68.4 ± 8.1 (15)
Serum prostaglandin E2 (ng/L)	349 ± 33[a] (16)	356 ± 38[ab] (8)	380 ± 38[b] (16)
Serum glucagon (mU/L)	54.3 ± 62.3[a] (20)	55.0 ± 38.4[a] (17)	104 ± 67[b] (16)
Serum cortosterone (μg/L)	7.51 ± 1.27[a] (16)	7.88 ± 1.64[ab] (8)	8.68 ± 1.61[b] (16)
Serum adrenaline (μg/L)	3.01 ± 1.64[a] (7)	10.5 ± 9.4[b] (14)	13.7 ± 7.5[b] (7)
Serum interleukin-6 (ng/L)	24.7 ± 17.7[a] (9)	47.8 ± 27.8[b] (14)	60.6 ± 32.8[b] (6)
Serum tumour necrosis factor α (ng/L)	69.1 ± 54.9[a] (12)	174 ± 161[b] (16)	205 ± 175[b] (13)
Serum C reactive protein (μg/L)	34.3 ± 19.2[a] (16)	49.1 ± 31.4[a] (17)	107 ± 72[b] (10)
Serum lactate dehydrogenase (U/L)	2592 ± 646 (7)	2234 ± 627 (7)	2518 ± 454 (6)

Note: Blood samples were collected from non-fasted mice at the ages of 3 (specific for estradiol) or 11~15 weeks. Each value represents the mean ± SD and the numbers in brackets indicate the size of the tested samples. Different letters in superscript indicate a level of statistical significance of $p < 0.05$.

Although the circulating levels of total cholesterol, free FAs, high-density lipoprotein
(HDL), and estradiol showed no significant differences among the three genotypes, the levels
of plasma triglycerides (TGs) were significantly elevated in Pai/Pai mice when compared to
wt or Pai/wt genotypes ($p < 0.05$, Table 3), indicating hypertriglyceridemia in Pai/Pai mice.

In Pai/Pai mice, serum concentrations of prostaglandin E2 (PGE2, 109%), glucagon
(2-fold), corticosterone (116%), adrenaline (4.5-fold), tumour necrosis factor-alpha (TNFα, 3-
fold), interleukin-6 (IL-6, 2.5-fold), and C-reactive protein (CRP, 3-fold) were significantly
increased ($p < 0.05$) when compared to their wt littermates. However, only FGF21 (109%),
adrenaline (3.5-fold), TNFα (2.5-fold), and IL-6 (2-fold) were significantly ($p < 0.05$) elevated
in Pai/wt mice (Table 3). Additionally, there was no increase in blood levels of lactic acid
dehydrogenase in either Pai mice, indicating no cellular injury caused by the PAI protein.
These results suggest that the t10c12-CLA can lead to an inflammation reaction and response
to illness-related stimuli in female mammals.

Figure 3. Comparisons of glucose (A) and insulin (B) tolerance tests and dynamic glucose
levels (C) in wild-type (wt) and Pai mice. (A) Absolute blood glucose levels and the area
under the curve are compared among wt (n = 5), Pai/wt (n = 10), and Pai/Pai (n = 5) mice.
(B) Blood glucose levels relative to initial values and the area under the curve are compared
among wt (n = 10), Pai/wt (n = 12), and Pai/Pai (n = 7) mice. (C) Dynamic blood glucose
levels and the area under the curve are measured in wt (n = 12), Pai/wt (n = 13), and
Pai/Pai (n = 10) fasting mice starved from Zeitgeber time 0 to 24. Zeitgeber times 0 and 12

are lights-on and -off times, respectively. The bars represent the mean ± SD. † indicates p <
0.05 between Pai/wt and wt or Pai/Pai mice; # indicates p < 0.05 between wt and Pai/Pai
mice; * indicates p < 0.05 between wt and Pai/wt mice.

Both glucose and insulin tolerance tests showed no difference in the circulating
glucose concentrations or the area under the curve between wt and two Pai genotypes.
However, the glucose levels before glucose injection were significantly (p < 0.05) higher in
Pai/wt mice compared to wt or Pai/Pai littermates (Fig. 3 A-B). To investigate further, we
measured blood glucose levels during a 24-h fasting duration and found that Pai/wt mice
maintained higher glucose levels continuously compared to their wt littermates (p < 0.05;
Fig. 3 C). This suggests that t10c12-CLA can ameliorate the circulating glucose level in Pai/wt
mice during starvation, which may be associated with the elevated FGF21 exclusively in
Pai/wt female mice (Table 3), similar to high FGF21 levels in Pai/wt male mice [3], for its
beneficial insulin-sensitising and glucose-lowering effects [17].

3. 5. WAT characteristics

WAT features were determined in the mice at 11 weeks of age. Compared to their wt
littermates, Pai/Pai mice showed generalised organomegaly, including significantly
enlarged livers, spleens, hearts, and ovaries, when body weight was considered (p < 0.05;
Supplementary Table 2). However, no fat mass loss was observed (Fig. 4 A-B). The
histological analysis did not reveal any changes in the cross-sectional area of white
adipocytes (Fig. 4 C-D). Pai/wt mice showed enlarged ovaries only and no fat loss whereas
its cross-sectional area per adipocyte was increased (p < 0.05), which was also confirmed by
the increased mRNA level of leptin (Fig. 4 E).

Total RNA levels of 14 (42%) genes, including seven up-regulated and seven down-
regulated, were changed in Pai/Pai white adipocytes among 33 examined genes (Table 4).
The seven up-regulated genes were corticosterone receptor Nr3c1, F4/80, Adiponectin, G6p,
Fasn, Ppar-γ, and Hsl. The seven down-regulated genes were Lpl, Cpt1a, Plin1, Prdm16, Insr,
Glut4, and Chrebp. These aberrant gene expressions in white adipocytes suggest that t10c12-
CLA can affect a series of metabolic processes, such as lipid metabolism, lipid and glucose
uptake, energy expenditure, stimuli response and glucose homeostasis.

Figure 4. Magnetic resonance imaging and histological analysis of white adipose tissues.
(A) Coronal sections were obtained on the whole-body midsection for four wild-type or
Pai/Pai mice at 11 weeks in each group during magnetic resonance imaging (bar = 1 cm).
The wild-type mice show adipose tissue in the subcutaneous and intra-abdominal regions
as areas of increased signal intensity on images. The Pai/Pai mice show normal body fat.
(B) There is no difference in WAT mass among wild-type (n = 21), Pai/wt (n = 17), and
Pai/Pai (n = 13) mice. (C) Imaging of hematoxylin-eosin staining (bar = 10 μm), (D) analysis
of cross-sectional area per cell, and (E) RNA level of leptin gene show that the volume of
white adipocytes is specifically increased in Pai/wt mice (n = 5). The bars represent the
mean ± SD. Different letters indicate p < 0.05.

Similar to Pai/Pai cells, Ppar-γ (1.52 ± 0.34, n = 10), Nr3c1 (1.53 ± 0.21, n = 12), and
F4/80 (1.98 ± 0.64, n = 10) were also up-regulated (p < 0.05), and Cpt1a (0.58 ± 0.27, n = 5) and
Chrebp (0.43 ± 0.45, n = 5) were down-regulated in Pai/wt adipocytes compared to wt
samples (p < 0.05), respectively. The RNA levels of Fasn and Adiponectin were not different
from those of wt samples (p > 0.05).

Table 4. Relative Expression of mRNAs in the white adipose tissue

Genes	Wild-type	Pai/Pai
Fatty acid synthase (*Fasn*)	1.04 ± 0.30 (6)	1.80 ± 0.47*** (8)
Peroxisome proliferator-activated receptor-γ (*Ppar-γ*)	1.03 ± 0.24 (12)	1.54 ± 0.61* (10)
Adipose triglyceride lipase (*Atgl*)	1.01 ± 0.16 (7)	1.06 ± 0.19 (7)
Hormone-sensitive lipase (*Hsl*)	1.02 ± 0.23 (9)	1.44 ± 0.29* (8)
Monoacylglycerol lipase (*Mgl*)	0.93 ± 0.25 (9)	0.99 ± 0.32 (8)
Perilipin 1a (*Plin1*)	1.07 ± 0.44 (8)	0.43 ± 0.16** (8)
Comparative gene identification 58 (*Cgi58*)	1.04 ± 0.30 (8)	0.81 ± 0.35 (8)
Lipoprotein lipase (*Lpl*)	1.03 ± 0.27 (14)	0.69 ± 0.24** (15)
Carnitine palmitoyltransferase Ia (*Cpt1a*)	1.06 ± 0.36 (8)	0.56 ± 0.15** (8)
Long-chain acyl-CoA dehydrogenase (*Lcad*)	1.12 ± 0.58 (7)	0.95 ± 0.44 (8)
Medium-chain acyl-CoA dehydrogenase (*Mcad*)	1.01 ± 0.17 (6)	1.07 ± 0.18 (9)
Uncoupling protein 1 (*Ucp1*)	1.09 ± 0.50 (6)	5.39 ± 5.88 (7)
Ppar-gamma coactivator 1 alpha (Pgc1α)	1.05 ± 0.39 (8)	1.18 ± 0.37 (8)
PR domain containing 16 (*Prdm16*)	1.01 ± 0.17 (7)	0.70 ± 0.20** (7)
Insulin receptor (*Insr*)	1.03 ± 0.25 (11)	0.75 ± 0.22** (13)
Insulin receptor substrate-1 (*Irs1*)	0.98 ± 0.32 (15)	1.03 ± 0.34 (20)
Insulin receptor substrate-2 (*Irs2*)	1.17 ± 0.69 (5)	2.28 ± 1.29 (6)
Phosphoinositide 3-kinase (*Pi3k*)	1.05 ± 0.39 (5)	0.78 ± 0.37 (8)
IGF binding protein 1 (*Igfbp1*)	1.80 ± 2.00 (6)	0.55 ± 0.14 (5)
Glucose transporter type 4 (*Glut4*)	1.02 ± 0.24 (6)	0.72 ± 0.17* (6)
Carbohydrate response element binding protein (Chrebp)	1.11 ± 0.51 (8)	0.35 ± 0.17** (8)
Fibroblast growth factor 21 (*Fgf21*)	1.08 ± 0.46 (6)	0.71 ± 0.39 (7)
Glucose-6-phosphatase (*G6p*)	1.24 ± 0.87 (6)	2.68 ± 0.83* (6)
Phosphoenolpyruvate carboxykinase (*Pepck*)	1.07 ± 0.42 (8)	1.11 ± 0.42 (8)
AMP-activated protein kinase (*Ampk*)	1.10 ± 0.49 (5)	1.45 ± 0.58 (8)
Adiponectin	0.99 ± 0.24 (9)	1.42 ± 0.38** (10)
Nuclear receptor subfamily 3 group C member 1 (*Nr3c1*)	1.04 ± 0.29 (22)	1.64 ± 0.81** (22)
F4/80	1.03 ± 0.29 (12)	1.54 ± 0.58* (11)
CD68	1.06 ± 0.38 (14)	1.03 ± 0.61 (13)
CD11c	1.05 ± 0.29 (14)	0.95 ± 0.44 (16)
CCAAT/enhancer-binding-protein beta (*Cebp*)	1.04 ± 0.31 (6)	1.13 ± 0.60 (8)
Forkhead box protein c2 (*Foxc2*)	1.02 ± 0.21 (5)	1.15 ± 0.45 (7)
Pai	1.24 ± 1.23 (23)	393 ± 270*** (25)

Numbers in brackets indicate the size of tested samples. Each value represents the mean ± SD. * indicates $p < 0.05$; ** indicates $p < 0.01$; *** indicates $p < 0.001$, respectively.

3. 6. BAT thermoregulation

To investigate the possible reason for hypothermia in Pai/Pai mice, BAT features
were measured. Magnetic resonance imaging and dissection analysis did not reveal any mass
changes of BAT in Pai mice (Fig. 5 A-B). HE staining showed Pai/Pai adipocytes with
uniform and small-sized lipid drops, while Pai/wt adipocytes had size-irregular lipid drops
compared to wt adipocytes containing uniform and large-sized lipid drops (Fig. 5 C). The

 reduced cross-sectional areas per adipocyte indicated that the volumes of brown adipocytes

269   decreased significantly in both Pai genotypes (p < 0.05; Fig. 5 D).

Figure 5. **Aspects of brown adipose tissues**. (A) Axial sections of the interscapular BAT show similar areas of grey signal intensity on magnetic resonance images (brown arrows) between four Pai/Pai and four wild-type mice at 11 weeks. (B) There is no BAT mass difference among wild-type (n = 21), Pai/wt (n = 17), and Pai/Pai (n = 13) mice. (C-D) Analysis of hematoxylin-eosin staining shows that the sizes of lipid drops become small in the Pai/wt or Pai/Pai adipocytes (bar = 10 μm) and the cross-sectional area per cell is decreased in Pai/wt and Pai/Pai mice (n = 5). (E) Analyses of the relative expression of mRNAs of critical genes in wild-type (n = 5~6) and in Pai/Pai (n = 6~8) mice. (F-G) Western

blot analysis of proteins and their relative intensities in wild-type (n = 3) and Pai/Pai mice (n = 3). See Table 4 for full names of genes/proteins. pAMPK, phosphorylated AMPK; pHSL, phosphorylated hormone-sensitive lipase. The bars represent the mean ± SD. * indicates p < 0.05; ** indicates p < 0.01; and *** indicates p < 0.001 vs. wild-type, respectively.

In Pai/Pai brown adipocytes, among the 18 tested genes, the transcriptional levels were changed in eight genes, which were up-regulated Fasn, Atgl, Plin1, Fgf21, and Ampk, as well as down-regulated Hsl, Mgl, and Pgc1α (p < 0.05; Fig. 5 E). In contrast, the other critical genes involved in lipolysis and thermogenesis, such as Ppar-γ, Ucp1, Ucp2, Prdm16, and Ppar-δ, showed no changes in RNA levels. Western blot analysis revealed that UCP1 and UCP2, phosphorylated HSL and CPT1B were over-expressed (p < 0.05; Fig. 5 F-G). Simultaneously, an increase in total AMPK and a decrease in phosphorylated AMPK were observed in Pai/Pai BAT. These results indicate that the t10c12-CLA effect is associated with BAT's thermogenesis, lipolysis, and beta-oxidation processes. The discrepancy of levels of some mRNA and their encoded proteins also suggests the complex regulation by feedback or adaptive response.

3. 7. Hepatic features

To investigate if t10c12-CLA affects hepatic function, we compared the Pai liver features and the molecular basis of lipid and glucose metabolism with wt samples.

In Pai/Pai mice, hepatic hypertrophy was observed (Fig. 6 A). The levels of hepatic TC remained unchanged (Fig. 6 B), but the transcriptional levels of critical genes involved in the sterol pathway were changed, which were down-regulated Srebp1a, Insig2, Hmgcr, and Ldlr; as well as up-regulated Srebp1c and Insig1 (p < 0.05; Table 5). On the other hand, the levels of hepatic TGs increased by 15.4%, and histological staining showed profound fat accumulation (~2.3 fold) in swollen hepatocytes (p < 0.05; Fig. 6 C-G), indicating hepatic steatosis in Pai/Pai mice. RNA levels of 48% (11/23) genes involved in lipid metabolism were significantly (p < 0.05) altered in Pai/Pai livers, including three up-regulated (Fasn, Mgl, Cgi58) and eight down-regulated genes, which were down-regulate Ppar-γ, Dgat1, Dgat2, Atgl, Lpl, Cpt1a, Mcad, and Acox1; as well as up-regulated Mgl and Cgi58 (Table 5). Western blot analysis revealed that the protein levels of pAMPK and FASN were reduced, and CPT1A was increased while the levels of AMPK, ATGL, and CPA1B did not change (Fig. 6 H-I). The result indicates that the chronic effect of t10c12-CLA attenuates the pathway modified by

pAMPK. It also suggests that hepatic steatosis causes the down-regulation of fatty acid synthesis and up-regulation of beta-oxidation through negative feedback regulation in the liver.

Figure 6. Lipid and histological analysis of livers. (A) Dissection analysis shows an increased liver mass in Pai/Pai (n = 13) but not in Pai/wt (n = 17) mice compared to wild-type mice (n = 21). (B-C) ELISA analyses indicate similar levels of hepatic total cholesterol (TC) and increased levels of triglycerides (TGs) in Pai/Pai livers (n = 12) compared to wild-type (n = 14) and Pai/wt (n = 14). (D-E) Analyses of hematoxylin-eosin staining show the abnormal morphology and hepatocyte oedema in Pai/Pai livers and the enlarged cross-sectional area per cell in Pai/Pai and Pai/wt hepatocytes compared to wild-type samples (n

= 5). (F-G) Analyses of Oil red staining show hepatic lipid accumulation in Pai/Pai but not
in Pai/wt mice compared to wild-type mice (n = 5), Bar = 10 μm. (H-I) Western blot analysis
of proteins and their relative intensities in wild-type (n=3) and Pai/Pai mice (n = 3). The bars
represent the mean ± SD, and different letters indicate p < 0.05.

Moreover, this study analysed a range of crucial enzymes related to insulin/IGF
signalling and glucose metabolism. The mRNA levels of Igfbp1, Glut4, Fgf21, G6p, and Pepck
were decreased, but western blot analysis showed that the FGF21 level did not alter. These
findings suggest that t10c12-CLA may have a significant impact on glucose metabolism and
insulin sensitivity at least on transcriptional levels in female livers. Additionally, two
NADPH-producing enzymes, Malic and Pgd, were significantly up-regulated, and the
inflammatory factor CD11c was down-regulated in the Pai/Pai livers (p < 0.05; Table 5).

Table 5. Relative Expression of mRNAs in the liver

Genes	wild-type	Pai/wt	Pai/Pai
Sterol pathway			
Liver X receptor α (*Lxrα*)	1.00 ± 0.09 (6)	0.91 ± 0.26 (6)	1.08 ± 0.45 (8)
Liver X receptor β (*Lxrβ*)	1.04 ± 0.31 (9)	0.91 ± 0.42 (6)	1.11 ± 0.26 (7)
SREBP cleavage-activating protein (*Scap*)	1.01 ± 0.15 (8)	0.92 ± 0.15 (6)	0.90 ± 0.25 (6)
Sterol Regulatory Element Binding Protein-1a (*Srebp1a*)	1.02 ± 0.22[a] (7)	0.46 ± 0.25[b] (6)	0.61 ± 0.20[b] (7)
Srebp1c	0.96 ± 0.33[a] (6)	1.61 ± 0.55[b] (7)	1.60 ± 0.42[b] (5)
Srebp2	1.07 ± 0.39 (8)	1.06 ± 0.22 (7)	1.25 ± 0.52 (8)
Insulin induced gene 1 (*Insig1*)	1.03 ± 0.25[a] (7)	1.97 ± 0.52[b] (7)	1.47 ± 0.26[b] (5)
Insulin induced gene 2a (*Insig2*)	1.07 ± 0.38[a] (6)	0.20 ± 0.34[b] (6)	0.59 ± 0.31[b] (5)
HMG-CoA reductase (*Hmgcr*)	1.01 ± 0.16[a] (6)	1.23 ± 0.32[a] (5)	0.62 ± 0.34[b] (6)
LDL receptor (*Ldlr*)	1.04 ± 0.33[a] (8)	1.23 ± 0.36[a] (6)	0.71 ± 0.10[b] (5)
Fatty acid and TGs synthesis			
Fatty acid synthase (*Fasn*)	1.01 ± 0.18[a] (6)		2.65 ± 1.32[b] (6)
Stearoyl-CoA desaturase 1 (*Scd1*)	1.07 ± 0.43 (7)	0.97 ± 0.49 (6)	1.11 ± 0.57 (6)
Peroxisome proliferator-activated receptor-γ (*Ppar-γ*)	1.01 ± 0.46[a] (14)	1.06 ± 0.27[a] (7)	0.66 ± 0.44[b] (12)
Glycerol-3-phosphate acyltransferase 1 (*Gpat1*)	1.04 ± 0.33 (7)	1.44 ± 0.46 (6)	1.22 ± 0.31 (6)
1-acylglycerol-3-phosphate O-acyltransferase 2 (*Agpat2*)	0.97 ± 0.25 (8)	0.80 ± 0.22 (6)	1.11 ± 0.31 (7)
Diacylglycerol acyltransferase-1 (*Dgat1*)	1.00 ± 0.16[a] (7)	0.81 ± 0.18[ab] (6)	0.70 ± 0.13[b] (7)
Diacylglycerol acyltransferase-2 (*Dgat2*)	1.00 ± 0.08[a] (6)	1.04 ± 0.15[a] (6)	0.85 ± 0.11[b] (5)
CD36 (Cd36)	1.06 ± 0.41 (6)		0.83 ± 0.31 (6)

Table 5. Relative Expression of mRNAs in the liver

Genes	wild-type	Pai/wt	Pai/Pai
Lipolysis and fatty acid oxidation			
Hepatic triglyceride lipase (*Htgl*)	0.98 ± 0.27 (7)	0.90 ± 0.28 (6)	1.13 ± 0.34 (7)
Adipose triglyceride lipase (*Atgl*)	1.02 ± 0.21[a] (6)	0.79 ± 0.36[ab] (6)	0.56 ± 0.14[b] (6)
Hormone-sensitive lipase (*Hsl*)	1.05 ± 0.33 (8)	1.29 ± 0.41 (6)	1.16 ± 0.49 (8)
Monoacylglycerol lipase (*Mgl*)	1.02 ± 0.19[a] (8)	1.65 ± 0.55[b] (6)	1.56 ± 0.58[b] (7)
Comparative gene identification 58 (*Cgi58*)	1.05 ± 0.34[a] (7)	1.38 ± 0.66[ab] (6)	1.75 ± 0.59[b] (7)
Adenylate cyclase 3 (*Adcy3*)	1.07 ± 0.42 (8)		1.08 ± 0.23 (8)
Lipoprotein lipase (*Lpl*)	1.02 ± 0.23[a] (7)	0.60 ± 0.17[b] (6)	0.50 ± 0.23[b] (6)
Carnitine palmitoyltransferase Ia (*Cpt1a*)	1.01 ± 0.19[a] (14)	0.83 ± 0.17[b] (7)	0.68 ± 0.18[b] (11)
Long-chain 3-hydroxyacyl-CoA dehydrogenase (*Lchad*)	1.04 ± 0.28 (9)		1.01 ± 0.25 (9)
Acetyl-CoA acyltransferase 2 (*Acaa2*)	1.08 ± 0.44 (7)		1.23 ± 0.23 (6)
Long-chain acyl-CoA dehydrogenase (*Lcad*)	1.00 ± 0.09 (8)	0.93 ± 0.22 (7)	0.78 ± 0.30 (8)
Medium-chain acyl-CoA dehydrogenase (*Mcad*)	1.03 ± 0.27[a] (9)	0.57 ± 0.12[b] (6)	0.65 ± 0.20[b] (6)
Uncoupling protein 1 (*Ucp1*)	1.04 ± 0.32 (7)	1.12 ± 0.40 (7)	0.98 ± 0.34 (7)
Acyl-CoA oxidase-1 (*Acox1*)	1.04 ± 0.32[a] (8)	0.89 ± 0.32[a] (7)	0.69 ± 0.32[b] (8)
Acyl-CoA oxidase-2 (*Acox2*)	1.07 ± 0.42 (9)		1.17 ± 0.43 (9)
Insulin and IGF-1 signaling			
Insulin receptor (*Insr*)	1.05 ± 0.37 (9)	1.35 ± 0.42 (6)	1.36 ± 0.70 (8)
Insulin receptor substrate-1 (*Irs1*)	0.97 ± 0.43 (6)	0.97 ± 0.43 (4)	0.94 ± 0.59 (6)
Insulin receptor substrate-2 (*Irs2*)	1.27 ± 1.01 (9)	1.27 ± 0.87 (7)	1.41 ± 0.99 (8)
Phosphoinositide 3-kinase (*Pi3k*)	1.04 ± 0.28 (7)		1.17 ± 0.24 (7)
Insulin-like growth factor-1 (*Igf1*)	1.04 ± 0.31[a] (8)	1.16 ± 0.29[a] (7)	2.09 ± 0.80[b] (6)
IGF binding protein 1 (*Igfbp1*)	1.04 ± 0.30[a] (6)	0.45 ± 0.33[b] (6)	0.09 ± 0.02[b] (5)
Forkhead box protein a2 (*Foxa2*)	1.06 ± 0.37[a] (9)	1.98 ± 0.86[b] (6)	1.76 ± 0.48[b] (7)
Glucose metabolism and gluconeogenesis			
Glucose transporter type 4 (*Glut4*)	1.06 ± 0.38[a] (6)	0.88 ± 0.28[a] (5)	0.61 ± 0.25[b] (6)
Carbohydrate response element binding protein (*Chrebp*)	1.03 ± 0.26 (5)		1.33 ± 0.43 (6)
Fibroblast growth factor 21 (*Fgf21*)	1.07 ± 0.46[a] (5)		0.19 ± 0.12[b] (7)
Glucose-6-phosphatase (*G6p*)	1.07 ± 0.41[a] (9)	0.47 ± 0.17[b] (7)	0.66 ± 0.19[b] (6)
Phosphoenolpyruvate carboxykinase (*Pepck*)	1.04 ± 0.29[a] (8)	0.53 ± 0.22[b] (6)	0.51 ± 0.19[b] (7)
NADPH-producing			
Malic enzyme (*Malic*)	0.99 ± 0.32[a] (8)	0.98 ± 0.39[a] (7)	1.72 ± 0.70[b] (7)
Glucose-6-phosphate dehydrogenase (*G6pd*)	1.06 ± 0.38 (6)	1.14 ± 0.34 (4)	1.05 ± 0.46 (4)
6-Phosphogluconate dehydrogenase (*Pgd*)	1.05 ± 0.35[a] (7)		1.73 ± 0.56[b] (6)
Others			
AMP-activated protein kinase (*Ampk*)	1.07 ± 0.41 (7)		1.48 ± 0.32 (7)
Liver kinase B1 (*Lkb1*)	1.02 ± 0.23 (5)		1.10 ± 0.12 (5)
Nuclear receptor subfamily 3 group C member 1 (*Nr3c1*)	1.07 ± 0.39 (6)		2.01 ± 1.05 (6)

Table 5. Relative Expression of mRNAs in the liver

Genes	wild-type	Pai/wt	Pai/Pai
CD11c	1.04 ± 0.31^a (7)	0.57 ± 0.26^b (6)	0.53 ± 0.27^b (7)
Cyclophilin	1.01 ± 0.17 (9)	1.04 ± 0.28 (6)	1.01 ± 0.16 (6)
Pai	1.01 ± 0.43^a (5)	1674 ± 569^b (6)	6059 ± 2317^c (6)

Numbers in brackets indicate the size of tested samples. Each value represents the mean ± SD. Different letters in superscript indicate $p < 0.05$.

Compared to wt samples, Pai/wt livers exhibited normal except for swollen cells
(Fig. 6 E). However, among the forty tested genes (Table 5), the mRNA levels of eight genes
(Hmgcr, Ldlr, Ppar-γ, Dgat2, Acox1, Igf1, Glut4, and Malic) were significantly ($p < 0.05$)
different from those in Pai/Pai livers, but not different from those in wt samples. The
remaining 32 genes had similar transcription levels to Pai/Pai samples, and 13 (33%) of them
were markedly different from those in wt samples ($p < 0.05$). Specifically, these 13 genes that
exhibited different expressions from wt livers included four up-regulated (Mgl, Srebp1c,
Insig1, and Foxa2) and nine down-regulated (Lpl, Cpt1a, Mcad, Srebp1a, Insig2, Igfbp1, G6p,
Pepck, and Cd11c). These results suggest that gene expression is genotype-specific and that
the process of lipolysis and gluconeogenesis is impaired in Pai/wt mice.

3. 8. Hypothalamic analysis

To determine whether t10c12-CLA affected the hypothalamus, critical hypothalamic
genes and proteins were also measured. The mRNA levels of the leptin receptor and agouti-
related peptide were increased, while the transcription levels of the ghrelin receptor and
Orexins decreased in the Pai/wt or Pai/Pai hypothalamus when compared to wt mice (Fig.
7 A). Western blot analysis revealed no changes in AMPK, phosphorylated AMPK, and
GRP78 proteins in the Pai/Pai hypothalamus; additionally, the ratio of pAMPK/AMPK did
not show a significant difference between wt and Pai/Pai samples ($p > 0.05$; Fig. 7 B-C). These
findings suggest that the central regulation of energy intake may only be affected at the
transcriptional level.

Figure 7. Analyses of the relative mRNAs (A), protein levels and their relative intensities in the hypothalamus (B-C). Agrp, Agouti-related peptide; Ampk, AMP-activated protein kinase; Ghsr, Ghrelin receptor; Grp78, glucose-regulated protein 78; Insr, Insulin receptor; Lepr, Leptin receptor; Mc4r, Melanocortic 4 receptor; Nr3c1, Nuclear receptor subfamily 3 group C member 1; (p)AMPK, (phosphorylated) AMP-activated protein kinase; Pomc, Proopiomelanocortin. The sample sizes for RNA analysis are 6~9 in each group. The bars represent the mean ± SD, and different letters indicate p < 0.05.

4. DISCUSSION

Our previous observation in Pai/Pai male mice showed that t10c12-CLA reduced body fat by releasing heat via BAT activation with lower blood TGs [3]. Conversely, in the Pai female mice, t10c12-CLA did not cause body fat loss but resulted in complicated symptoms, including hypothermia, fatty liver and hypertriglyceridemia, and was highly responsive to stress and inflammation in a dose-dependent manner. The different physiological and pathological characteristics of both sexes can help us understand the mechanism of action of t10c12-CLA in gender-specific and dose-dependent manners. As for Pai mice, there are two problems we have not fixed until now. Firstly, we have reported as many results as possible for a comprehensive understanding of the Pai mouse, but a few parameters are difficult to explain well and even show contradictions in some respects. Secondly, explaining why there are only minimal differences in t10c12-CLA and linoleic acid is challenging although we once observed drastic changes in an increase of product t10c12-CLA and a decrease of substrate linoleic acid in Pai-transfected 3T3 cells [18].

The metabolic pathway of t10c12-CLA, similar to that of linoleic or alpha-linolenic acids [18], suggests it can competitively interrupt the metabolic pathway of n-6 or n-3 FAs and their derived leukotrienes signals/hormonal molecules. It also indicates that t10c12-CLA can be directly converted into the analogues of n-6 FAs and their derivatives. PGE2, an arachidonic acid-derived hormone, will be easily disturbed by t10c12-CLA. PGE2 is involved in stress response and has essential effects on inflammation, fever, and renal filtration [19]. It can induce corticosterone secretion by regulating the hypothalamic-pituitary-adrenal axis activity and enhancing adrenaline synthesis in the Chromaffin cells of the adrenal medulla. Subsequently, the increased circulating corticosterone and adrenaline could stimulate excess glucagon secretion [20]. In Pai/Pai female mice, we have observed the excess PGE2 induced by t10c12-CLA, and then the PGE2 (or analogues) caused the increases in corticosterone, glucagon, adrenaline, and inflammatory factors (IL-6, TNFα, and CRP). Obviously, hypothermia in Pai/Pai female mice may be related to hyperprostaglandinemia, hyperadrenerginemia and inflammatory response; while hyperglyceridemia and fatty liver are associated with hypercorticosteroidemia and glucagon resistance.

In the hypothalamic preoptic area, PGE2 receptors EP3, coupled to cAMP decreases, can stimulate a passive defence mechanism such as less O_2 consumption and hypothermia to avoid further energy expenditure; while receptors EP4, connected to cAMP increases, can

stimulate an active defence mechanism, including more O_2 consumption and febrile
responses by increasing BAT thermogenesis, typically occurs to promote well-being and
survival [19,21]. It suggests that gender-based differences in PGE2 levels might partially result
in the opposite heat release in male and female Pai/Pai mice. In female mice, the co-elevation
of PGE2 and corticosterone suggests that t10c12-CLA induced a passive defence mechanism
including hypothermia which conversely promotes BAT thermogenesis and lipid
metabolism by negative feedback to down-regulate pAMPK and up-regulate UCP1, UCP2,
CPT1B, and pHSL. In contrast, in male mice, the lack of PGE2 elevation suggests that t10c12-
CLA induced an active defence mechanism including hyperthermia to promote survival [3].

In addition to PGE2- and adrenaline-induced heating [22], BAT thermogenesis can be
mediated through the central GRP78 [23,24] or AMPK-OREXINS-BMP8B pathway [25].
Orexin-null mice exhibited periods of decreased activity during the dark phase and
demonstrated hypothermia, reduced oxygen consumption, and blunted thermogenic
responses [26]; furthermore, BMP8B's thermogenic effect is sexually dimorphic and observed
in females only [25]. The Pai/Pai female mice exhibited similar phenotypes to orexin-null
mice and lower transcription levels of Orexins in the Pai hypothalamus. These characteristics
suggest that the t10c12-CLA-induced BAT thermogenesis may be regulated through central
pathways in female mice. Unfortunately, we did not observe evident changes in GRP78,
AMPK, pAMPK, and pAMPK/AMPK ratio in the hypothalamus.

Previous studies on mice have yielded conflicting results on whether dietary t10c12-
CLA leads to hepatic steatosis [11,16,27,28]. The current observation of fatty liver and
hypertriglyceridemia in Pai/Pai mice, which may be associated with t10c12-CLA-induced
excess corticosterone for the overproduction of corticosterone can lead to Cushing syndrome,
corticosteroid-induced lipodystrophy, hypertriglyceridemia, or hepatic steatosis [29,30]. On
the other hand, the existence of glucagon resistance can increase liver fat [31-33], although
chronic increases in plasma glucagon promote lipolysis through INSP3R1/CAMKII/ATGL
[34] or PKA/HSL pathways in hepatocytes [35] and stimulate beta-oxidation by activating
AMPK/PPAR-α/CPT1A/MCAD pathway [20]. In the current study, the serum glucagon
concentration increased nearly twofold in Pai/Pai female mice, which is similar to its
physiological levels (2~3 fold) in response to long-term fasting or hypoglycemia [36], whereas
the protein levels of unchanged ATGL, decreased pAMPK, and increased CPT1A indicate
the chronic glucagon excess did not stimulate the lipolysis and beta-oxidation in the liver.
These results suggest that the chronic impact of t10c12-CLA might give rise to glucagon
resistance in Pai/Pai female mice. In this case, the down-regulated hepatic enzymes may not

fully and immediately catalyse the FAs molecules mobilised from other tissues based on
glucagon stimulation. Consequently, it disrupts hepatic and systemic lipid homeostasis and
leads to dyslipidemia and fatty liver in female mice. Contrarily, the t10c12-CLA did not cause
the overproduction of corticosterone or glucagon in Pai/Pai male mice. Thus, the mice had
no fatty liver disease and even exhibited lower TGs levels in the blood.

FGF21 is a key regulator of the specific hormonal signalling pathway in increasing
energy expenditure, reducing hepatic and plasma TGs levels, accelerating lipoprotein
catabolism in WAT and BAT [37], and suppressing physical activity via central action [38].
FGF21 signalling enhances insulin sensitivity and lowers serum glucose by acting directly on
adipose tissues [17,39]. The Pai/wt mouse is the first animal in which t10c12-CLA can induce
FGF21 secretion. Excess FGF21 is a unique feature in Pai/wt, not in Pai/Pai genotypes,
indicating that t10c12-CLA-induced FGF21 secretion is specifically associated with low doses
in both genders and it can be suppressed to the normal levels through negative feedback
when high doses occur in Pai/Pai mice. This suggests that the health benefits of low-dose
t10c12-CLA may be due, in part, to its ability to stimulate FGF21 secretion. Therefore, the
different effects of t10c12-CLA on glucose metabolism in diabetic Zucker fatty rats [40] and
insulin resistance in mice [12,41] may be related to the different levels of FGF21 induced by
t10c12-CLA. In the current study, excess FGF21 may be associated with specific phenotypes
in Pai/wt mice, such as blood glucose homeostasis during starvation, enlarged white
adipocytes, no steatosis in hepatocytes and normal circulating TGs. The exact mechanism by
which t10c12-CLA activates FGF21 secretion remains unclear and requires further
investigation.

Based on the excesses of hormones, abnormality of the energy metabolism, and
physical activities in Pai/Pai mice in this study, we suggest that high doses of t10c12 result
in an increased PEG2 which subsequently induces increases in corticosterone, glucagon, and
adrenaline, and inflammatory factors (IL6, TNFα, and C reaction protein). The combination
of these factors causes metabolic syndrome, including fatty liver, hyperglyceridemia,
hypothermia, and inflammatory changes in lean female mice (Fig. 8). Conversely, low doses
of t10c12-CLA may cause increases in circulating adrenaline which can induce WAT lipolysis
and BAT heating through a mechanism of negative feedback, and FGF21 which can promote
energy expenditure, reduce hepatic and plasma TGs levels, accelerate lipoprotein catabolism
in adipocytes, suppress physical activity, and enhance insulin sensitivity and lower serum
glucose (Fig. 8).

Figure 8. Diagram showing the dose-dependent effect of chronic t10c12-CLA on lean female mice. Besides the common effects, such as increased adrenaline, IL-6, and TNFα, as well as decreased heat release and oxygen consumption (Middle panel), the low doses specifically increase circulating levels of FGF21 which plays active roles in maintaining energy homeostasis, suppressing physical activity, and improving serum glucose levels in Pai/wt mice (Left panel). Conversely, the high doses specifically result in excesses of prostaglandin E2, corticosterone, glucagon, and CRP; subsequently, the combination of these factors causes metabolic syndrome, including hypothermia, fatty liver, and hypertriglyceridemia in Pai/Pai mice (Right panel). Words in black and red describe physiological features and endocrinal factors, respectively.

Our studies on male and female Pai mice suggest that the chronic effects of t10c12-CLA on lean animals are dose-dependent and differ between sexes. Furthermore, the metabolic impact of t10c12-CLA involves various hormones and has system-wide effects on multiple metabolic pathways. Thus, interpreting the data from Pai mice requires careful consideration, and more work is needed to understand the mechanism of action of t10c12-CLA and its metabolic impact.

Abbreviations

BAT, brown adipose tissue; CPT1, carnitine palmitoyltransferase 1; CRP, C-reactive protein; FA, fatty acids; FASN, fatty acid synthase; FGF21, fibroblast growth factor 21; IL-6, interleukin-6; LA, linoleic acid; pai, *Propionibacterium acnes* isomerase; (p)AMPK, phosphorylated AMP-activated protein kinase; PEG2, prostaglandin E2; (p)HSL, phosphorylated hormone-sensitive lipase; t10c12-CLA, *trans* 10, *cis* 12-conjugated linoleic acid; TC, total cholesterol; TGs, triglycerides; TNFα, tumour necrosis factor-alpha; UCP, uncoupling protein; WAT, white adipose tissue; wt, wild-type;

Ethics approval and consent to participate:

All animal experiments followed the Committee for Experimental Animals of Yangzhou University. All applicable institutional and/or national guidelines for the care and use of animals were followed.

Institutional Review Board Statement:

The animal study protocol was approved by the Ethics Committee of Yangzhou University (protocol code NSFC2020-SYXY-20 and dated 25 March 2020).

Consent for publication:

Not applicable

Availability of data and materials:

The datasets used and/or analysed during the current study are available from the corresponding author upon reasonable request.

Competing interests:

The authors declare that they have no competing interests.

Funding sources

This work was supported by the National Key Research and Development Program of China (2018YFC1003800), the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD), and the Jiangsu Funding Program for Excellent Postdoctoral Talent.

Acknowledgement

We thank graduate students Qi Wang, Mei Liu, and Jiaqi Lu in our lab for their
generous assistance in partial tests and Ms Yuyang Wang in our university for technique
help in gas chromatography. We also thank Professor Hongjun Wang at the Medical
University of South Carolina, USA, for the helpful discussion during the writing process and
Ms Wenjing Gou's language assistance in preparing this manuscript.

Authors 'contributions:

Conceptualization, Y Rao and K Gou; Data curation, Y Rao, M Li, B Wang, Y Wang,
and L Liang; Formal analysis, Y Rao and K Gou; Funding acquisition, Y Rao and K Gou;
Investigation, Y Rao, M Li, B Wang, Y Wang and L Liang; Methodology, Y Rao, M Li, B Wang,
Y Wang and L Liang; Project administration, K Gou; Software, Y Rao; Supervision, K Gou;
Validation, K Gou; Visualization, Y Rao and L Liang; Writing – original draft, Y Rao; Writing
– review & editing, K Gou. All authors proofread, made comments, and approved the report.

Supplementary data

This manuscript contains supplementary data.

References

1. Shen, W.; Baldwin, J.; Collins, B.; Hixson, L.; Lee, K.T.; Herberg, T.; Stames, J.; Cooney,
P.; Chuang, C.C.; Hopkins, R., et al. Low level of trans-10, cis-12 conjugated linoleic
acid decreases adiposity and increases browning independent of inflammatory
signaling in overweight Sv129 mice. *Journal of Nutritional Biochemistry* **2015**, *26*, 616-
625, doi:10.1016/j.jnutbio.2014.12.016.

2. den Hartigh, L.J.; Wang, S.R.; Goodspeed, L.; Wietecha, T.; Houston, B.; Omer, M.;
Ogimoto, K.; Subramanian, S.; Gowda, G.A.N.; O'Brien, K.D., et al. Metabolically
distinct weight loss by 10,12 CLA and caloric restriction highlight the importance of
subcutaneous white adipose tissue for glucose homeostasis in mice. *Plos One* **2017**, *12*,
doi:10.1371/journal.pone.0172912.

3. Rao, Y.; Li, S.L.; Li, M.J.; Wang, B.Z.; Wang, Y.Y.; Liang, L.W.; Yu, S.; Liu, Z.P.; Cui, S.;
Gou, K.M. Transgenic mice producing the trans 10, cis 12-conjugated linoleic acid
present reduced adiposity and increased thermogenesis and fibroblast growth factor
21 (FGF21). *J Nutr Biochem* **2023**, 10.1016/j.jnutbio.2023.109419, 109419,
doi:10.1016/j.jnutbio.2023.109419.

4. Berkel, C.; Cacan, E. Trans-10, cis-12 conjugated linoleic acid- and caloric restriction-
mediated upregulation of CNDP2 expression in white adipose tissue in rodents, with
implications in feeding and obesity. *J Nutr Biochem* **2023**, *114*, 109269,
doi:10.1016/j.jnutbio.2023.109269.

5. Wang, L.; Zhang, S.; Huang, Y.; You, W.; Zhou, Y.; Chen, W.; Sun, Y.; Yi, W.; Sun, H.;
Xie, J., et al. CLA improves the lipo-nutritional quality of pork and regulates the gut
microbiota in Heigai pigs. *Food Funct* **2022**, *13*, 12093-12104, doi:10.1039/d2fo02549c.

6. Kanter, J.E.; Goodspeed, L.; Wang, S.; Kramer, F.; Wietecha, T.; Gomes-Kjerulf, D.; Subramanian, S.; O'Brien, K.D.; den Hartigh, L.J. 10,12 Conjugated Linoleic Acid-Driven Weight Loss Is Protective against Atherosclerosis in Mice and Is Associated with Alternative Macrophage Enrichment in Perivascular Adipose Tissue. *Nutrients* **2018**, *10*, doi:10.3390/nu10101416.

7. Vaisar, T.; Wang, S.; Omer, M.; Irwin, A.D.; Storey, C.; Tang, C.; den Hartigh, L.J. 10,12-Conjugated linoleic acid supplementation improves HDL composition and function in mice. *J Lipid Res* **2022**, *63*, 100241, doi:10.1016/j.jlr.2022.100241.

8. Kim, J.H.; Kim, Y.; Kim, Y.J.; Park, Y. Conjugated linoleic acid: Potential health benefits as a functional food ingredient. *Annu Rev Food Sci Technol* **2016**, *7*, 221-244, doi:10.1146/annurev-food-041715-033028.

9. Marques, T.M.; Wall, R.; O'Sullivan, O.; Fitzgerald, G.F.; Shanahan, F.; Quigley, E.M.; Cotter, P.D.; Cryan, J.F.; Dinan, T.G.; Ross, R.P., et al. Dietary trans-10, cis-12-conjugated linoleic acid alters fatty acid metabolism and microbiota composition in mice. *Br J Nutr* **2015**, *113*, 728-738, doi:10.1017/S0007114514004206.

10. Cuciniello, R.; Luongo, D.; Ferramosca, A.; Lunetti, P.; Rotondi-Aufiero, V.; Crispi, S.; Zara, V.; Maurano, F.; Filosa, S.; Bergamo, P. Conjugated linoleic acid downregulates Alzheimer's hallmarks in aluminum mouse model through an Nrf2-mediated adaptive response and increases brain glucose transporter levels. *Free Radic Biol Med* **2022**, *191*, 48-58, doi:10.1016/j.freeradbiomed.2022.08.027.

11. Cordoba-Chacon, J.; Sugasini, D.; Yalagala, P.C.R.; Tummala, A.; White, Z.C.; Nagao, T.; Kineman, R.D.; Subbaiah, P.V. Tissue-dependent effects of cis-9,trans-11- and trans-10,cis-12-CLA isomers on glucose and lipid metabolism in adult male mice. *J Nutr Biochem* **2019**, *67*, 90-100, doi:10.1016/j.jnutbio.2019.01.020.

12. Clement, L.; Poirier, H.; Niot, I.; Bocher, V.; Guerre-Millo, M.; Krief, S.; Staels, B.; Besnard, P. Dietary trans-10,cis-12 conjugated linoleic acid induces hyperinsulinemia and fatty liver in the mouse. *J Lipid Res* **2002**, *43*, 1400-1409, doi:10.1194/jlr.m20008-jlr200.

13. Kelley, D.S.; Bartolini, G.L.; Warren, J.M.; Simon, V.A.; Mackey, B.E.; Erickson, K.L. Contrasting effects of t10,c12- and c9,t11-conjugated linoleic acid isomers on the fatty acid profiles of mouse liver lipids. *Lipids* **2004**, *39*, 135-141, doi:10.1007/s11745-004-1211-9.

14. Jaudszus, A.; Moeckel, P.; Hamelmann, E.; Jahreis, G. Trans-10,cis-12-CLA-caused lipodystrophy is associated with profound changes of fatty acid profiles of liver, white adipose tissue and erythrocytes in mice: possible link to tissue-specific alterations of fatty acid desaturation. *Ann Nutr Metab* **2010**, *57*, 103-111, doi:10.1159/000319877.

15. Letona, A.Z.; Niot, I.; Laugerette, F.; Athias, A.; Monnot, M.C.; Portillo, M.P.; Besnard, P.; Poirier, H. CLA-enriched diet containing t10,c12-CLA alters bile acid homeostasis and increases the risk of cholelithiasis in mice. *J Nutr* **2011**, *141*, 1437-1444, doi:10.3945/jn.110.136168.

16. Li, S.L.; Ma, S.Y.; Xu, B.R.; Fan, Z.Y.; Li, M.J.; Cao, W.G.; Gou, K.M. Effects of trans-10, cis-12-conjugated linoleic acid on mice are influenced by the dietary fat content and the degree of murine obesity. *Eur J Lipid Sci Tech* **2015**, *117*, 1908-1918, doi:10.1002/ejlt.201400568.

17. Flippo, K.H.; Potthoff, M.J. Metabolic Messengers: FGF21. *Nat Metab* **2021**, *3*, 309-317, doi:10.1038/s42255-021-00354-2.

18. Subbaiah, P.V.; Gould, I.G.; Lal, S.; Aizezi, B. Incorporation profiles of conjugated linoleic acid isomers in cell membranes and their positional distribution in phospholipids. *Biochim Biophys Acta* **2011**, *1811*, 17-24, doi:10.1016/j.bbalip.2010.09.004.

19. Furuyashiki, T.; Narumiya, S. Stress responses: the contribution of prostaglandin E(2) and its receptors. *Nat Rev Endocrinol* **2011**, *7*, 163-175, doi:10.1038/nrendo.2010.194.

20. Zeigerer, A.; Sekar, R.; Kleinert, M.; Nason, S.; Habegger, K.M.; Muller, T.D. Glucagon's metabolic action in health and disease. *Compr Physiol* **2021**, *11*, 1759-1783, doi:10.1002/cphy.c200013.

21. Saper, C.B.; Lowell, B.B. The hypothalamus. *Curr Biol* **2014**, *24*, R1111-1116, doi:10.1016/j.cub.2014.10.023.

22. Cypess, A.M.; Weiner, L.S.; Roberts-Toler, C.; Franquet Elia, E.; Kessler, S.H.; Kahn, P.A.; English, J.; Chatman, K.; Trauger, S.A.; Doria, A., et al. Activation of human brown adipose tissue by a beta3-adrenergic receptor agonist. *Cell Metab* **2015**, *21*, 33-38, doi:10.1016/j.cmet.2014.12.009.

23. Contreras, C.; Gonzalez-Garcia, I.; Seoane-Collazo, P.; Martinez-Sanchez, N.; Linares-Pose, L.; Rial-Pensado, E.; Ferno, J.; Tena-Sempere, M.; Casals, N.; Dieguez, C., et al. Reduction of hypothalamic endoplasmic reticulum stress activates browning of white fat and ameliorates obesity. *Diabetes* **2017**, *66*, 87-99, doi:10.2337/db15-1547.

24. Contreras, C.; Fondevila, M.F.; Lopez, M. Hypothalamic GRP78, a new target against obesity? *Adipocyte* **2018**, *7*, 63-66, doi:10.1080/21623945.2017.1405878.

25. Martins, L.; Seoane-Collazo, P.; Contreras, C.; Gonzalez-Garcia, I.; Martinez-Sanchez, N.; Gonzalez, F.; Zalvide, J.; Gallego, R.; Dieguez, C.; Nogueiras, R., et al. A functional link between ampk and orexin mediates the effect of bmp8b on energy balance. *Cell Rep* **2016**, *16*, 2231-2242, doi:10.1016/j.celrep.2016.07.045.

26. Chemelli, R.M.; Willie, J.T.; Sinton, C.M.; Elmquist, J.K.; Scammell, T.; Lee, C.; Richardson, J.A.; Williams, S.C.; Xiong, Y.M.; Kisanuki, Y., et al. Narcolepsy in orexin knockout mice: Molecular genetics of sleep regulation. *Cell* **1999**, *98*, 437-451, doi:Doi 10.1016/S0092-8674(00)81973-X.

27. Ashwell, M.S.; Ceddia, R.P.; House, R.L.; Cassady, J.P.; Eisen, E.J.; Eling, T.E.; Collins, J.B.; Grissom, S.F.; Odle, J. Trans-10, cis-12-conjugated linoleic acid alters hepatic gene expression in a polygenic obese line of mice displaying hepatic lipidosis. *J Nutr Biochem* **2010**, *21*, 848-855, doi:10.1016/j.jnutbio.2009.06.013.

28. Vyas, D.; Kadegowda, A.K.; Erdman, R.A. Dietary conjugated linoleic acid and hepatic steatosis: species-specific effects on liver and adipose lipid metabolism and gene expression. *J Nutr Metab* **2012**, *2012*, 932928, doi:10.1155/2012/932928.

29. Laryea, G.; Schutz, G.; Muglia, L.J. Disrupting hypothalamic glucocorticoid receptors causes HPA axis hyperactivity and excess adiposity. *Mol Endocrinol* **2013**, *27*, 1655-1665, doi:10.1210/me.2013-1187.

30. Nicolaides, N.C.; Charmandari, E.; Kino, T.; Chrousos, G.P. Stress-related and circadian secretion and target tissue actions of glucocorticoids: Impact on health. *Front Endocrinol (Lausanne)* **2017**, *8*, 70, doi:10.3389/fendo.2017.00070.

31. Krilov, L.; Nguyen, A.; Miyazaki, T.; Unson, C.G.; Bouscarel, B. Glucagon receptor recycling: role of carboxyl terminus, beta-arrestins, and cytoskeleton. *Am J Physiol-Cell Ph* **2008**, *295*, C1230-C1237, doi:10.1152/ajpcell.00240.2008.

32. Charbonneau, A.; Unson, C.G.; Lavoie, J.M. High-fat diet-induced hepatic steatosis reduces glucagon receptor content in rat hepatocytes: potential interaction with acute exercise. *J Physiol-London* **2007**, *579*, 255-267, doi:10.1113/jphysiol.2006.121954.

33. Sammons, M.F.; Lee, E.C.Y. Recent progress in the development of small-molecule glucagon receptor antagonists. *Bioorg Med Chem Lett* **2015**, *25*, 4057-4064, doi:10.1016/j.bmcl.2015.07.092.

34. Perry, R.J.; Zhang, D.; Guerra, M.T.; Brill, A.L.; Goedeke, L.; Nasiri, A.R.; Rabin-Court, A.; Wang, Y.; Peng, L.; Dufour, S., et al. Glucagon stimulates gluconeogenesis by INSP3R1-mediated hepatic lipolysis. *Nature* **2020**, *579*, 279-283, doi:10.1038/s41586-020-2074-6.

35. Hayashi, Y. Glucagon regulates lipolysis and fatty acid oxidation through inositol triphosphate receptor 1 in the liver. *J Diabetes Investig* **2021**, *12*, 32-34, doi:10.1111/jdi.13315.

36. Kleinert, M.; Sachs, S.; Habegger, K.M.; Hofmann, S.M.; Muller, T.D. Glucagon regulation of energy expenditure. *Int J Mol Sci* **2019**, *20*, doi:10.3390/ijms20215407.

37. Schlein, C.; Talukdar, S.; Heine, M.; Fischer, A.W.; Krott, L.M.; Nilsson, S.K.; Brenner, M.B.; Heeren, J.; Scheja, L. Fgf21 lowers plasma triglycerides by accelerating lipoprotein catabolism in white and brown adipose tissues. *Cell Metab* **2016**, *23*, 441-453, doi:10.1016/j.cmet.2016.01.006.

38. Bookout, A.L.; de Groot, M.H.; Owen, B.M.; Lee, S.; Gautron, L.; Lawrence, H.L.; Ding, X.; Elmquist, J.K.; Takahashi, J.S.; Mangelsdorf, D.J., et al. FGF21 regulates metabolism and circadian behavior by acting on the nervous system. *Nat Med* **2013**, *19*, 1147-1152, doi:10.1038/nm.3249.

39. Holland, W.L.; Adams, A.C.; Brozinick, J.T.; Bui, H.H.; Miyauchi, Y.; Kusminski, C.M.; Bauer, S.M.; Wade, M.; Singhal, E.; Cheng, C.C., et al. An FGF21-adiponectin-ceramide axis controls energy expenditure and insulin action in mice. *Cell Metab* **2013**, *17*, 790-797, doi:10.1016/j.cmet.2013.03.019.

40. Henriksen, E.J.; Teachey, M.K.; Taylor, Z.C.; Jacob, S.; Ptock, A.; Kramer, K.; Hasselwander, O. Isomer-specific actions of conjugated linoleic acid on muscle glucose transport in the obese Zucker rat. *Am J Physiol Endocrinol Metab* **2003**, *285*, E98-E105, doi:10.1152/ajpendo.00013.2003.

41. Tsuboyama-Kasaoka, N.; Takahashi, M.; Tanemura, K.; Kim, H.J.; Tange, T.; Okuyama, H.; Kasai, M.; Ikemoto, S.; Ezaki, O. Conjugated linoleic acid supplementation reduces adipose tissue by apoptosis and develops lipodystrophy in mice. *Diabetes* **2000**, *49*, 1534-1542, doi:10.2337/diabetes.49.9.1534.