Gender-specific impact of trans 10, cis 12-conjugated linoleic acid on pai transgenic

female mice

Yu Rao[1,2, #], Luwen Liang[1, #], Meijuan Li[3, #], Yangyang Wang[1], Baozhu Wang[1], Kemian

Gou[1,2, *]

[1]*Institute of Comparative Medicine,* [2]*Jiangsu Co-innovation Center for Prevention and*

Control of Important Animal Infectious Diseases and Zoonoses, College of Veterinary

Medicine, Yangzhou University, Yangzhou 225009, China;

[3]*Institute of Animal Husbandry and Veterinary Science, Guizhou Academy of Agricultural*

Sciences, Guiyang 550005, China.

# YR, LL, and ML contributed equally to this work.

*Corresponding author: Kemian Gou, College of Veterinary Medicine, Yangzhou

University, Yangzhou 225009, China.

Email: gou@yzu.edu.cn

ABSTRACT

Trans 10, *cis* 12-conjugated linoleic acid (t10c12-CLA) has been shown to induce body fat loss in mammals. In this study, we analysed female pai transgenic mice that produced t10c12-CLA to determine the long-term impact of this substance on the lipid metabolism of lean mammals. Phenotype profiling showed that the effects of t10c12-CLA were genotype-specific in biallelic pai/pai or monoallelic pai/wt mice compared to their wild-type littermates. In both pai genotypes, white fat was not reduced, but heat release was decreased. Specifically, pai/wt mice exhibited increased chronic FGF21 and adrenaline possibly to regulate energy homeostasis and reduce physical activities. Contrarily, pai/pai mice showed hypothermia which may be related to the overproduction of prostaglandins and adrenaline, as well as an inflammatory response. Analysis of brown adipose tissue indicated that hypothermia might conversely promote thermogenesis and lipolysis by up-regulating UCP1/2, pHSL, CPT1B, and AMPK, and down-regulating pAMPK in pai/pai mice. Moreover, pai/pai mice also showed hypertriglyceridemia and fatty liver which is associated with long-term excesses of corticosterone and glucagon. Further analysis of pai/pai livers revealed that hepatic steatosis resulted from attenuated lipid metabolism via down-regulating pAMPK and fatty acid synthase. These results suggest that the systemic impact of chronic t10c12-CLA is dose-dependent in lean female mammals, low doses of t10c12-CLA may play an active role via the FGF21 and/or adrenaline pathways while its high-dose exposure may cause hormones overproduction and pose a safety risk.

Keywords: thermogenesis; FGF21; adrenaline; prostaglandins; corticosterone; glucagon

1. INTRODUCTION

Trans 10, cis 12-conjugated linoleic acid (t10c12-CLA), one of the most studied isomers of linoleic acid, occurs naturally in small amounts in some foods such as beef and dairy products. Its popular health benefit is reducing body fat by increasing energy expenditure and white adipose tissue (WAT) browning [1,2], promoting biosynthesis of N-lactoyl-phenylalanine via the up-regulation of cytosolic nonspecific dipeptidase 2 in WAT [3], or increasing intramuscular fat [4]. In addition, it has other potential health influences, including depressing milk fat production by inhibiting lipogenesis [5-7], reducing the globule size of milk fat droplets [8], protecting against atherosclerosis by modulating the macrophage phenotype [9], promoting beneficial effects on cholesterol efflux [10], reducing oxidative stress [11], changing intestinal microbiota [12] and metabolite profiles [4], down-regulating Alzheimer's hallmarks in an aluminium mouse through an Nrf2-mediated adaptive response and increasing brain glucose transporter levels [13].

Although the potential health benefits, especially the fat loss properties of t10c12-CLA, have been extensively studied, little is known about the mechanism of action of t10c12-CLA and it is also unclear what the optimal dosage for this fatty acid is, particularly under different host obesity or food fat degrees [14,15]. Moreover, some studies have not found its effects on fat reduction, and even a few studies have indicated its side effects result in fatty liver, an inflammatory response, and insulin resistance [11]. As we know, commercial CLA supplements used in most studies contain 10-20% other CLA isomers and the t10c12-CLA-mediated effect might be covered or reversed by other isomers. Additionally, commercial CLA differs significantly in the ratio of cis 9, trans 11-CLA to t10c12-CLA (approximately 1:1) when compared to levels found in food (approximately 8~9:1), therefore, CLA supplement users ingest significantly more t10c12-CLA than would be obtained from the diet [11,16]. This excessive t10c12-CLA dosage or CLA mixture supplements can bring the health impact of t10c12-CLA into conflict.

To identify the individual t10c12-CLA's impacts, we have produced a transgenic mouse that can produce t10c12-CLA by inserting the foreign gene encoding the *Propionibacterium acnes* isomerase (PAI) into the Rosa26 locus (Rao et al, 2023 bioRxiv preprint: https://doi.org/10.1101/2023.01.31.526123). In its lean male offspring, decreased white fat was observed in biallelic pai/pai, not in monoallelic

pai/wt genotypes, suggesting t10c12-CLA's impact on fat reduction in a dose-
dependent manner. The pai/wt male mice with excess FGF21 in the blood suggests
that t10c12-CLA in low doses possibly stimulate the FGF21 secretion and plays an
active role via the pathway of FGF21 on health. Whereas the lean pai/pai male mice
released more heat by promoting energy expenditure via UCP1/2-mediated
thermogenesis and increasing fatty acids oxidation through the CPT1B pathway in
brown adipose tissue (BAT) (Rao et al, 2023 bioRxiv preprint:
https://doi.org/10.1101/2023.01.31.526123). The study on male mice suggests that fat
mass loss is the result of BAT heating, and the pathways of t10c12-CLA action are
associated with its dosage.

The current study investigated the adiposity characteristics and metabolic
profiles of lean pai female mice. In contrast to their male counterparts, the findings of
this study demonstrate that pai female mice did not experience changes in body fat,
but did experience a decrease in heat release. Moreover, specific metabolic features in
female mice indicate that the systemic effects of t10c12-CLA on health are dose-
dependent and gender-specific. The results suggest that pai mice may be a useful
model to investigate the chronic t10c12-CLA's effects and its mechanisms of action in
the body.

2. MATERIALS AND METHODS

2. 1. Chemicals

Unless otherwise stated, all chemicals were purchased from Sigma (USA) or Sinopharm Chemical Reagent Co., Ltd. (Shanghai, China). Restriction endonucleases were purchased from New England Biolabs Inc (USA).

2. 2. Diet

Fresh diets were prepared every month in accordance with formula No. D12450H of OpenSource DIETS™ (10% kcal% fat, Research Diets Inc., NJ, USA) under sterile conditions. The sources of the food-grade ingredients used in the diets are listed in Supplementary Table 1. The fatty acid composition of the diets was spot-checked using gas chromatography to examine the contents of linoleic acid and t10c12-CLA. Each kilogram of the diet contains approximately 20 g of linoleic acid, and no t10c12-CLA was detected in the diet samples.

2. 3. Mice

All animal experiments were conducted in accordance with the guidance provided by the Committee for Experimental Animals of Yangzhou University. The animal study protocol was approved by the Ethics Committee of Yangzhou University (protocol code NSFC2020-SYXY-20 and dated 25 March 2020). The mice were housed in a light-controlled room (12L:12D, lights on at 0700 h) at a temperature of 22-23°C and fed ad libitum with a standard diet containing 10% kcal% fat. For the transgenic founder produced from hybrids embryos of DBA/2 and C57BL/6J, the pai/wt mice were backcrossed to C57BL/6J mice for more than eight generations before the present study. Then the pai/wt matings were used to produce the biallelic pai/pai, monoallelic pai/wt, and wild-type (wt) offspring for this study. Unless otherwise specified, mice of the above three genotypes aged 11-15 weeks were used for analysis in this study.

2. 4. Methods

The methods utilised in this study, including gas chromatography, RNA and
proteins analyses, histology, hepatic and blood parameter measurements,
intraperitoneal glucose (2g/kg body weight) and insulin (0.5 IU/kg body weight)
tolerance tests, and experiments conducted in metabolic cages (Supplementary
method), have been previously described in our study of pai male mice (Rao et al, 2023
bioRxiv preprint: https://doi.org/10.1101/2023.01.31.526123). For reviewing
convenience, the methods are copied in the Supplementary methods.

2.5. Genotyping

Genotyping was performed using genomic DNA from tail tissues, with the
presence of the pai gene being assayed by amplifying the 514-bp pai fragments
(forward: 5-taaccatgttcatgccttcttc-3; reverse: 5-caccttgttgtagtgtccgttt-3) and the wt
Rosa26 sequence being assayed by amplifying the 605-bp fragments that spanned the
insertional site of the Rosa26 locus (forward: 5-ccaaagtcgctctgagttgttatcagt-3; reverse:
5-ggagcgggagaaatggatatgaag-3). PCR amplification was performed for 30 cycles at
94°C for 30 sec, 57°C for 30 sec, and 72°C for 40 sec.

2.6. Measurement of serum estradiol

Three-week-old mice were injected intraperitoneally with 5 IU pregnant mare
serum gonadotrophin (Jinmu Biotech Co. Ltd., Wuxi, China), and 48 h later, blood was
collected from the orbital vein of anaesthetised mice and used to measure the serum
estradiol levels using the E2 ELISA kit (Meimian Industrial Co. Ltd., Jiangsu, China).

2. 5. Statistical analysis.

Statistical analysis was performed with unpaired Student's t-test or one-way
ANOVA (Brown-Forsythe and Welch) among wt, pai/wt, and pai/pai mice. All values
are presented as mean $\pm$ SD. A p-value of < 0.05 was considered statistically
significant.

3. RESULTS

3. 1. Fatty acids analysis

Gas chromatography was used to analyse the compositions of fatty acids
(FAs) in the livers, kidneys, hearts, tibialis anterior muscle, and interscapular BAT.
The content of t10c12-CLA had increased by 69% in the pai/wt kidneys compared to
wt littermates ($p < 0.05$; Table 1). However, the quantities of substrate linoleic acid had
decreased in the pai/pai livers by 23% and skeletal muscle by 34%, and had increased
by 36% in the pai/wt kidneys ($p < 0.05$; Supplementary Table 2).

Table 1. Comparison of t10c12-CLA and circulating factors in wt and pai mice

	wt	pai/wt	pai/pai
t10c12-CLA (µg/g)			
Liver	2.56 ± 3.24 (7)	4.30 ± 4.10 (5)	ND (5)
Kidney	6.50 ± 1.25[a] (5)	11.0 ± 3.65[b] (6)	6.09 ± 0.42[a] (4)
Heart	ND (5)	ND (4)	ND (4)
Skeletal muscle	ND (4)	ND (4)	ND (4)
Brown adipose tissue	22.4 ± 15.9 (4)	29.5 ± 7.5 (4)	29.8 ± 0.5 (4)
Blood/hormonal parameters			
Blood glucose (mM)	7.35 ± 0.70 (12)	7.37 ± 1.18 (15)	7.55 ± 1.35 (13)
Serum insulin (µg/L)	0.21 ± 0.08 (9)	0.27 ± 0.11 (17)	0.32 ± 0.18 (8)
Serum leptin (µg/L)	1.04 ± 0.41 (8)	0.92 ± 0.21 (14)	0.91 ± 0.31 (7)
Serum ghrelin (ng/L)	181 ± 44(7)	225 ± 108 (20)	210 ± 152 (6)
Serum FGF21 (ng/L)	403 ± 29[a] (16)	441 ± 49[b] (10)	421 ± 34[ab] (18)
Serum total cholesterol (µM)	311 ± 40 (17)	324 ± 40 (8)	321 ± 30 (16)
Plasma triglycerides (mg/dL)	107 ± 35[a] (34)	105 ± 27[a] (29)	129 ± 42[b] (29)
Plasma HDL (mg/dL)	48.6 ± 13.9 (34)	47.7 ± 16.2 (29)	52.5 ± 19.6 (29)
Serum free fatty acids (µM)	471 ± 95 (8)	527 ± 79 (16)	517 ± 136 (8)
Serum Estradiol (pM)	71.0 ± 8.8 (15)	64.9 ± 6.8 (7)	68.4 ± 8.1 (15)
Serum prostaglandin E2 (ng/L)	349 ± 33[a] (16)	356 ± 38[ab] (8)	380 ± 38[b] (16)
Serum glucagon (mU/L)	54.3 ± 62.3[a] (20)	55.0 ± 38.4[a] (17)	104 ± 67[b] (16)
Serum cortosterone (µg/L)	7.51 ± 1.27[a] (16)	7.88 ± 1.64[ab] (8)	8.68 ± 1.61[b] (16)
Serum adrenaline (µg/L)	3.01 ± 1.64[a] (7)	10.5 ± 9.4[b] (14)	13.7 ± 7.5[b] (7)
Serum interleukin-6 (ng/L)	24.7 ± 17.7[a] (9)	47.8 ± 27.8[b] (14)	60.6 ± 32.8[b] (6)
Serum TNFα (ng/L)	69.1 ± 54.9[a] (12)	174 ± 161[b] (16)	205 ± 175[b] (13)
Serum C reactive protein (µg/L)	34.3 ± 19.2[a] (16)	49.1 ± 31.4[a] (17)	107 ± 72[b] (10)
Serum LDH (U/L)	2592 ± 646 (7)	2234 ± 627 (7)	2518 ± 454 (6)

Table 1. Comparison of t10c12-CLA and circulating factors in wt and pai mice

	wt	pai/wt	pai/pai

Note: Tissues from 11-week-old mice was analysed by gas chromatography. Blood samples were collected from non-fasted mice at the ages of 3 (only estradiol analysis) or 11~15 weeks. FGF21, fibroblast growth factor 21; HDL, high-density lipoprotein; LDH, Lactate dehydrogenase; ND, not detected; TNFα, tumour necrosis factor alpha. Each value represents the mean ± SD, and the numbers in brackets indicate the size of the tested samples. Different letters in superscript indicate a level of statistical significance of $p < 0.05$.

The contents of other FAs such as myristic (14:0), palmitic (16:0), stearic (18:0),
palmitoleic (16:1n-7), cis-vaccenic (18:1n-7), oleic (18:1n-9), arachidonic (20:4n-6), and
docosahexaenoic (22:6n-3) acids were changed to varying degrees in one or more
tissues (Supplementary Table 2), This suggests that the changes in the content of each
FA were genotype- or tissue-specific. Additionally, the amounts of total FAs (mg/g)
increased in the pai/wt kidneys (10.2 ± 1.39 versus 13.3 ± 1.69; $p < 0.05$), similar to our
previous study in pai/wt male mice (https://doi.org/10.1101/2023.01.31.526123).
These results suggest lipopathy in the pai/wt kidney.

Figure 1. Comparison of growth of pups from wt (A) and pai/wt (B) mothers, as well as
their weaning (3-week-old) weight (C). Each group contains 11-29 (A, B) or 24-81 (C)
pups. The bars represent the mean ± SD. # or different letters indicate $p < 0.05$ among three
genotypes.

3. 2. Body weight analysis

There were no differences in body weight between pai/wt and wt littermates
from 3 weeks onwards in the group of wt mothers (Fig. 1A). However, in the group
of pai/wt mothers, there was a significant difference in weaning weight among the

three genotypes (wt > pai/wt > pai/pai, respectively; $p < 0.05$), and this difference
disappeared from five weeks of age onwards (Fig. 1 B-C).

Additionally, the weaning weight of wt and pai/wt offspring from pai/wt
mothers was higher than that of genotype-matched pups from wt mothers ($p < 0.05$;
Fig. 1C), similar to our previous study in male mice (Rao et al, 2023 bioRxiv preprint:
https://doi.org/10.1101/2023.01.31.526123). This further confirms that maternal
t10c12-CLA may play an active role in developing embryos and/or weaning pups.

3. 3. Energy metabolism and activity studies

Energy metabolism and activity parameters were assessed in consideration of
body weight. Compared to their wt littermates, both pai/wt and pai/pai mice
exhibited no significant differences in body weight gain, food and water intake (Fig. 2
A-C), respiratory exchange ratio (Fig. 2 F-G), and total distance travelled during the
whole 72-h observation (Fig. 2 R). However, the other parameters showed varying
differences.

The pai/pai mice did not decrease their total locomotor activities in either the
centre (Fig. 2 J-K, N-O) or margin/corners (Fig. 2 L-M, P-Q). Nevertheless, during both
the light ($p < 0.05$) and dark ($p > 0.05$) periods, they consumed 8.9% and 9.2% less O_2
(Fig. 2 D) and produced 10.8% and 10.1% less CO_2 (Fig. 2 E) as well as 9.3% and 9.4%
less heat (Fig. 2 H-I), respectively. Additionally, pai/pai mice covered a longer
distance per sampling duration (~50% longer) and exhibited 52.5% higher speed
during the light phase ($p < 0.05$; Fig. 2 S-T). The results suggest that pai/pai mice
simultaneously reduce oxygen consumption and heat release while maintaining
normal locomotor activities.

On the other hand, pai/wt females did not reduce their central activities ($p >$
0.05; Fig. 2 J-K) but displayed 30.1% fewer marginal activities, such as marginal
ambulating and stereotypic activities during the dark period ($p < 0.05$; Fig. 2 L-M, P-
Q). Moreover, during both the light and dark periods, they consumed 10.7% and
14.6% less O_2, produced 9.7% and 15.2% less CO_2, as well as 10.5% and 14.7% less heat,
respectively ($p < 0.05$, Fig. 2 D-E, H-I), suggesting that pai/wt mice simultaneously
reduce oxygen consumption and heat release while exhibiting fewer marginal
activities.

Figure 2. Abnormal energy metabolism and activities in pai mice. Comparison of body

weight gain during a 72-h period (A), food intake (B), water intake (C), O_2 consumption (D),
CO_2 production (E), respiratory exchange ratio (RER) and RER time course (F-G), heat release
and its time course (H-I), locomotor activities in the centre of the cage and its time course (J-
K), locomotor activities in the margin or corners of the cage and its time course (L-M), central
ambulating and stereotypic activities (N-O), marginal ambulating and stereotypic activities (P-
Q), total distance travelled during a 72-h period (R), distance travelled during a 39-min
sampling duration (S), and speed (T) among wt (n = 4), pai/wt (n = 4) and pai/pai (n = 4) mice
at 11 weeks. The lights turned on and off at 7:00 am and 7:00 pm, respectively. The data are
normalised to lean body weight. The bars represent the mean ± SD. Different letters indicate p
< 0.05 within the same phase.

3. 4. Metabolic complications

Concentrations of circulating glucose, insulin, leptin, and ghrelin (Table 1), as
well as small intestine length (Supplementary Table 3), showed no significant
differences (p > 0.05) between pai mice and their wt littermates. This suggests that
t10c12-CLA did not have any significant impact on energy intake, which is consistent
with the observation in the metabolic cage.

Both glucose and insulin tolerance tests showed no difference in the
circulating glucose concentrations or the area under the curve between wt and two
pai genotypes, however, the glucose levels before glucose injection were significantly
(p < 0.05) higher in pai/wt mice compared to wt or pai/pai littermates (Fig. 3 A-B).
To investigate further, we measured blood glucose levels during a 24-h fasting
duration and found that pai/wt mice maintained higher glucose levels continuously
compared to their wt littermates (p < 0.05; Fig. 3 C). This suggests that t10c12-CLA can
ameliorate the circulating glucose level in pai/wt mice during starvation, which may
be associated with the elevated FGF21 exclusively in pai/wt mice (Table 1), for its
beneficial insulin-sensitising and glucose-lowering effects [17].

Although the circulating levels of total cholesterol, free FAs, high-density
lipoprotein (HDL), and estradiol showed no significant differences among the three
genotypes, the levels of plasma triglycerides (TGs) were significantly elevated in
pai/pai mice when compared to wt or pai/wt genotypes (p < 0.05, Table 1), indicating
hypertriglyceridemia in pai/pai mice.

Figure 3. Comparisons of glucose (A) and insulin (B) tolerance tests, and dynamic glucose levels (C) in wt and pai mice. (A) Absolute blood glucose levels and the area under the curve are compared among wt (n = 5), pai/wt (n = 10), and pai/pai (n = 5) mice. (B) Blood glucose levels relative to initial values and the area under the curve are compared among wt (n = 10), pai/wt (n = 12), and pai/pai (n = 7) mice. (C) Dynamic blood glucose levels and the area under the curve are measured in wt (n = 12), pai/wt (n = 13), and pai/pai (n = 10) fasting mice starved from Zeitgeber time 0 to 24. Zeitgeber times 0 and 12 are lights-on and -off times, respectively. The bars represent the mean ± SD. # indicates $p < 0.05$

between wt and pai/pai mice; * indicates p < 0.05 between wt and pai/wt mice; #* indicates p
< 0.05 between pai/wt and wt or pai/pai mice.

In pai/pai mice, serum concentrations of prostaglandin E2 (PGE2, 109%),
glucagon (2-fold), corticosterone (116%), adrenaline (4.5-fold), tumour necrosis factor-
alpha (TNFα, 3-fold), interleukin-6 (IL-6, 2.5-fold), and C-reactive protein (CRP, 3-
fold) were significantly increased (p < 0.05; Table 1) when compared to their wt
littermates. However, only FGF21 (109%), adrenaline (3.5-fold), TNFα (2.5-fold), and
IL-6 (2-fold) were significantly (p < 0.05) elevated in pai/wt mice. Additionally, there
was no increase in blood levels of lactic acid dehydrogenase in either pai mice,
indicating no cellular injury caused by the PAI protein. These results suggest that the
t10c12-CLA can lead to an inflammation reaction and response to illness-related
stimuli in female mammals.

3. 5. WAT characteristics

WAT features were determined in the mice at 11 weeks of age. Compared to
their wt littermates, pai/pai mice showed generalised organomegaly, including
significantly enlarged livers, spleens, hearts, and ovaries, when body weight was
considered (p < 0.05; Supplementary Table 3). However, no fat mass loss was observed
(Fig. 4 A-B). The histological analysis did not reveal any changes in the cross-sectional
area of white adipocytes (Fig. 4 C-D). Pai/wt mice showed enlarged ovaries only and
no fat loss whereas its cross-sectional area per adipocyte was increased (p < 0.05),
which was also confirmed by the increased mRNA level of leptin (Fig. 4 E).

As shown in Table 2, among 33 examined genes, RNA levels of 14 (42%) genes,
including seven up-regulated and seven down-regulated, were changed in pai/pai
adipocytes. The names of these genes are listed in Table 2. The seven up-regulated
genes were corticosterone receptor Nr3c1, F4/80, Adiponectin, G6p, Fasn, Ppar-γ, and
Hsl. The seven down-regulated genes were Lpl, Cpt1a, Plin1, Prdm16, Insr, Glut4, and
Chrebp. These aberrant gene expressions in white adipocytes suggest that t10c12-CLA
can affect a series of metabolic processes, such as lipid metabolism, lipid and glucose
uptake, energy expenditure, stimuli response and glucose homeostasis.

Figure 4. Magnetic resonance imaging and histological analysis of white adipose tissues.
(A) Coronal sections were obtained on the whole-body midsection for four wt or pai/pai mice
at 11 weeks in each group during magnetic resonance imaging (bar = 1 cm). The wt mice
show adipose tissue in the subcutaneous and intra-abdominal regions as areas of increased
signal intensity on images. The pai/pai mice show normal body fat. (B) There is no difference
in WAT mass among wt (n = 21), pai/wt (n = 17), and pai/pai (n = 13) mice. (C) Imaging of

hematoxylin-eosin staining (bar = 10 μm), (D) analysis of cross-sectional area per cell, and
(E) RNA level of leptin gene show that the volume of white adipocytes is increased in pai/wt
mice (n = 5). The bars represent the mean ± SD. Different letters indicate p < 0.05.

Table 2. Relative Expression of mRNAs in the white adipose tissue

Genes	wt	pai/pai
Fatty acid synthase (*Fasn*)	1.04 ± 0.30 (6)	1.80 ± 0.47*** (8)
Peroxisome proliferator-activated receptor-γ (*Ppar-γ*)	1.03 ± 0.24 (12)	1.54 ± 0.61* (10)
Adipose triglyceride lipase (*Atgl*)	1.01 ± 0.16 (7)	1.06 ± 0.19 (7)
Hormone-sensitive lipase (*Hsl*)	1.02 ± 0.23 (9)	1.44 ± 0.29* (8)
Monoacylglycerol lipase (*Mgl*)	0.93 ± 0.25 (9)	0.99 ± 0.32 (8)
Perilipin 1a (*Plin1*)	1.07 ± 0.44 (8)	0.43 ± 0.16** (8)
Comparative gene identification 58 (*Cgi58*)	1.04 ± 0.30 (8)	0.81 ± 0.35 (8)
Lipoprotein lipase (*Lpl*)	1.03 ± 0.27 (14)	0.69 ± 0.24** (15)
Carnitine palmitoyltransferase Ia (*Cpt1a*)	1.06 ± 0.36 (8)	0.56 ± 0.15** (8)
Long-chain acyl-CoA dehydrogenase (*Lcad*)	1.12 ± 0.58 (7)	0.95 ± 0.44 (8)
Medium-chain acyl-CoA dehydrogenase (*Mcad*)	1.01 ± 0.17 (6)	1.07 ± 0.18 (9)
Uncoupling protein 1 (*Ucp1*)	1.09 ± 0.50 (6)	5.39 ± 5.88 (7)
Ppar-gamma coactivator 1 alpha (Pgc1α)	1.05 ± 0.39 (8)	1.18 ± 0.37 (8)
PR domain containing 16 (*Prdm16*)	1.01 ± 0.17 (7)	0.70 ± 0.20** (7)
Insulin receptor (*Insr*)	1.03 ± 0.25 (11)	0.75 ± 0.22** (13)
Insulin receptor substrate-1 (*Irs1*)	0.98 ± 0.32 (15)	1.03 ± 0.34 (20)
Insulin receptor substrate-2 (*Irs2*)	1.17 ± 0.69 (5)	2.28 ± 1.29 (6)
Phosphoinositide 3-kinase (*Pi3k*)	1.05 ± 0.39 (5)	0.78 ± 0.37 (8)
IGF binding protein 1 (*Igfbp1*)	1.80 ± 2.00 (6)	0.55 ± 0.14 (5)
Glucose transporter type 4 (*Glut4*)	1.02 ± 0.24 (6)	0.72 ± 0.17* (6)
Carbohydrate response element binding protein (Chrebp)	1.11 ± 0.51 (8)	0.35 ± 0.17** (8)
Fibroblast growth factor 21 (*Fgf21*)	1.08 ± 0.46 (6)	0.71 ± 0.39 (7)
Glucose-6-phosphatase (*G6p*)	1.24 ± 0.87 (6)	2.68 ± 0.83* (6)
Phosphoenolpyruvate carboxykinase (*Pepck*)	1.07 ± 0.42 (8)	1.11 ± 0.42 (8)
AMP-activated protein kinase (*Ampk*)	1.10 ± 0.49 (5)	1.45 ± 0.58 (8)
Leptin	1.03 ± 0.26 (5)	0.75 ± 0.34 (6)
Adiponecin	0.99 ± 0.24 (9)	1.42 ± 0.38** (10)
Nuclear receptor subfamily 3 group C member 1 (*Nr3c1*)	1.04 ± 0.29 (22)	1.64 ± 0.81** (22)
F4/80	1.03 ± 0.29 (12)	1.54 ± 0.58* (11)
CD68	1.06 ± 0.38 (14)	1.03 ± 0.61 (13)
CD11c	1.05 ± 0.29 (14)	0.95 ± 0.44 (16)
CCAAT/enhancer-binding-protein beta (*Cebp*)	1.04 ± 0.31 (6)	1.13 ± 0.60 (8)
Forkhead box protein c2 (*Foxc2*)	1.02 ± 0.21 (5)	1.15 ± 0.45 (7)
Pai	1.24 ± 1.23 (23)	393 ± 270*** (25)

Numbers in brackets indicate the size of tested samples. Each value represents the mean ± SD. *
indicates p < 0.05; ** indicates p < 0.01; *** indicates p < 0.001, respectively.

Similar to pai/pai cells, Ppar-γ (1.52 ± 0.34, n = 10), Nr3c1 (1.53 ± 0.21, n = 12),

and F4/80 (1.98 ± 0.64, n = 10) were also up-regulated (p < 0.05), and Cpt1a (0.58 ±

0.27, n = 5) and Chrebp (0.43 ± 0.45, n = 5) were down-regulated in pai/wt adipocytes

compared to wt samples (p < 0.05), respectively. The RNA levels of Fasn and

Adiponectin were not different from those of wt samples (p > 0.05).

3. 6. BAT thermoregulation

To investigate the possible reason for hypothermia in pai/pai mice, BAT

features were measured. Magnetic resonance imaging and dissection analysis did not

reveal any mass changes of BAT in pai mice (Fig. 5 A-B). HE staining showed pai/pai

adipocytes with uniform and small-sized lipid drops, while pai/wt adipocytes had

size-irregularly lipid drops compared to wt adipocytes containing uniform and large-

sized lipid drops (Fig. 5 C). The reduced cross-sectional areas per adipocyte indicated

that the volumes of brown adipocytes decreased significantly in both pai genotypes

(p < 0.05; Fig. 5 D).

In pai/pai brown adipocytes, among the 18 tested genes, the transcriptional

levels were changed in eight genes, which were up-regulated Fasn, Atgl, Plin1, Fgf21,

and Ampk; as well as down-regulated Hsl, Mgl, and Pgc1α (p < 0.05; Fig. 5 E). In

contrast, the other critical genes involved in lipolysis and thermogenesis, such as Ppar-

γ, Ucp1, Ucp2, Prdm16, and Ppar-δ, showed no changes in RNA levels. Western blot

analysis revealed that UCP1 and UCP2, phosphorylated HSL and CPT1B were over-

expressed (p < 0.05; Fig. 5 F-G). Simultaneously, an increase in total AMPK and a

decrease in phosphorylated AMPK were observed in pai/pai BAT. These results

indicate that the t10c12-CLA effect is associated with BAT's thermogenesis, lipolysis,

and beta-oxidation processes. The discrepancy of levels of some mRNA and their

encoded proteins also suggests the complex regulation by feedback or adaptive

response.

Figure 5. **Aspects of brown adipose tissues**. (A) Axial sections of the interscapular BAT

show similar areas of grey signal intensity on magnetic resonance images (brown arrows) between four pai/pai and four wt mice at 11 weeks. (B) There is no BAT mass difference among wt (n = 21), pai/wt (n = 17), and pai/pai (n = 13) mice. (C-D) Analysis of hematoxylin-eosin staining shows that the sizes of lipid drop become small in the pai/wt or pai/pai adipocytes (bar = 10 μm) and the cross-sectional area per cell is decreased in pai/wt and pai/pai mice (n = 5). (E) Analyses of the relative expression of mRNAs of critical genes in wt (n = 5~6) and in pai/pai (n = 6~8) mice. (F-G) Western blot analysis of proteins and their relative intensities in wt and pai/pai mice (n = 3). The full name of the gene/proteins is listed in Table 2. pAMPK, phosphorylated AMPK; pHSL, phosphorylated hormone-sensitive lipase. The bars represent the mean ± SD. Different letters or * indicate $p < 0.05$; ** indicates $p < 0.01$; *** indicates $p < 0.001$, respectively.

3. 7. Hepatic features

To investigate if t10c12-CLA affects hepatic function, we compared the pai liver features and the molecular basis of lipid and glucose metabolism with wt samples.

In pai/pai mice, hepatic hypertrophy was observed (Fig. 6 A). The levels of hepatic TC remained unchanged (Fig. 6 B), but the transcriptional levels of critical genes involved in the sterol pathway were changed, which were down-regulated Srebp1a, Insig2, Hmgcr, and Ldlr; as well as up-regulated Srebp1c and Insig1 ($p < 0.05$; Table 3). On the other hand, the levels of hepatic TGs increased by 15.4%, and histological staining showed profound fat accumulation (~2.3 fold) in swollen hepatocytes ($p < 0.05$; Fig. 6 C-F), indicating hepatic steatosis in pai/pai mice. RNA levels of 40% (10/22) genes involved in lipid metabolism were significantly ($p < 0.05$) altered in pai/pai livers, including two up-regulated (Mgl and Cgi58) and eight down-regulated genes, which were down-regulate Ppar-γ, Dgat1, Dgat2, Atgl, Lpl, Cpt1a, Mcad, and Acox1; as well as up-regulated Mgl and Cgi58 (Table 3). Western blot analysis revealed that the protein levels of pAMPK and FASN were reduced and CPT1A was increased while the levels of AMPK, ATGL, and CPA1B did not change (Fig. 6 H-I). The result indicates that the chronic effect of t10c12-CLA attenuates the pathway modified by pAMPK. It also suggests that hepatic steatosis causes the down-regulation of fatty acid synthesis and up-regulation of beta-oxidation through negative feedback in the liver.

Moreover, this study analysed a range of crucial enzymes related to
insulin/IGF signalling and glucose metabolism. The mRNA levels of Igfbp1, Glut4,
Fgf21, G6p, and Pepck were decreased, but western blot analysis showed that the
FGF21 level did not alter. These findings suggest that t10c12-CLA may have a
significant impact on glucose metabolism and insulin sensitivity at least on
transcriptional levels in female livers. Additionally, two NADPH-producing
enzymes, Malic and Pgd, were significantly up-regulated, and the inflammatory
factor CD11c was down-regulated in the pai/pai livers ($p < 0.05$; Table 3).

Figure 6. Lipid and histological analysis of livers. (A) Dissection analysis shows increased
liver mass in pai/pai (n = 13) but not in pai/wt (n = 17) mice compared to wt mice (n = 21). (B-
C) ELISA analyses indicate similar levels of hepatic total cholesterol (TC) and the increased
levels of triglycerides (TGs) in pai/pai livers (n = 12) compared to wt (n = 14) and pai/wt (n =
14). (D-E) Analyses of hematoxylin-eosin staining show the abnormal morphology and
hepatocyte oedema in pai/pai livers, and the enlarged cross-sectional area per cell in pai/pai
and pai/wt hepatocytes compared to wt samples (n = 5). (F-G) Analyses of Oil red staining
show hepatic lipid accumulation in pai/pai but not in pai/wt mice compared to wt mice (n = 5),
Bar = 10 μm. (H-I) Western blot analysis of proteins and their relative intensities in wt and
pai/pai mice (n = 3). The bars represent the mean ± SD. Different letters or * indicate p < 0.05,
** indicates p < 0.01, respectively.

Table 3. Relative Expression of mRNAs in the liver

Genes	wt	pai/wt	pai/pai
Sterol pathway			
Liver X receptor α (*Lxrα*)	1.00 ± 0.09 (6)	0.91 ± 0.26 (6)	1.08 ± 0.45 (8)
Liver X receptor β (*Lxrß*)	1.04 ± 0.31 (9)	0.91 ± 0.42 (6)	1.11 ± 0.26 (7)
SREBP cleavage-activating protein (*Scap*)	1.01 ± 0.15 (8)	0.92 ± 0.15 (6)	0.90 ± 0.25 (6)
Sterol Regulatory Element Binding Protein-1a (*Srebp1a*)	$1.02 ± 0.22^a$ (7)	$0.46 ± 0.25^b$ (6)	$0.61 ± 0.20^b$ (7)
Srebp1c	$0.96 ± 0.33^a$ (6)	$1.61 ± 0.55^b$ (7)	$1.60 ± 0.42^b$ (5)
Srebp2	1.07 ± 0.39 (8)	1.06 ± 0.22 (7)	1.25 ± 0.52 (8)
Insulin induced gene 1 (*Insig1*)	$1.03 ± 0.25^a$ (7)	$1.97 ± 0.52^b$ (7)	$1.47 ± 0.26^b$ (5)
Insulin induced gene 2a (*Insig2*)	$1.07 ± 0.38^a$ (6)	$0.20 ± 0.34^b$ (6)	$0.59 ± 0.31^b$ (5)
HMG-CoA reductase (*Hmgcr*)	$1.01 ± 0.16^a$ (6)	$1.23 ± 0.32^a$ (5)	$0.62 ± 0.34^b$ (6)
LDL receptor (*Ldlr*)	$1.04 ± 0.33^a$ (8)	$1.23 ± 0.36^a$ (6)	$0.71 ± 0.10^b$ (5)
Fatty acid and TGs synthesis			
Stearoyl-CoA desaturase 1 (*Scd1*)	1.07 ± 0.43 (7)	0.97 ± 0.49 (6)	1.11 ± 0.57 (6)
Peroxisome proliferator-activated receptor-γ (*Ppar-γ*)	$1.01 ± 0.46^a$ (14)	$1.06 ± 0.27^a$ (7)	$0.66 ± 0.44^b$ (12)
Glycerol-3-phosphate acyltransferase 1 (*Gpat1*)	1.04 ± 0.33 (7)	1.44 ± 0.46 (6)	1.22 ± 0.31 (6)
1-acylglycerol-3-phosphate O-acyltransferase 2 (*Agpat2*)	0.97 ± 0.25 (8)	0.80 ± 0.22 (6)	1.11 ± 0.31 (7)
Diacylglycerol acyltransferase-1 (*Dgat1*)	$1.00 ± 0.16^a$ (7)	$0.81 ± 0.18^{ab}$ (6)	$0.70 ± 0.13^b$ (7)
Diacylglycerol acyltransferase-2 (*Dgat2*)	$1.00 ± 0.08^a$ (6)	$1.04 ± 0.15^a$ (6)	$0.85 ± 0.11^b$ (5)
CD36 (*Cd36*)	1.06 ± 0.41 (6)		0.83 ± 0.31 (6)
Lipolysis and fatty acid oxidation			

Table 3. Relative Expression of mRNAs in the liver

Genes	wt	pai/wt	pai/pai
Hepatic triglyceride lipase (*Htgl*)	0.98 ± 0.27 (7)	0.90 ± 0.28 (6)	1.13 ± 0.34 (7)
Adipose triglyceride lipase (*Atgl*)	1.02 ± 0.21[a] (6)	0.79 ± 0.36[ab] (6)	0.56 ± 0.14[b] (6)
Hormone-sensitive lipase (*Hsl*)	1.05 ± 0.33 (8)	1.29 ± 0.41 (6)	1.16 ± 0.49 (8)
Monoacylglycerol lipase (*Mgl*)	1.02 ± 0.19[a] (8)	1.65 ± 0.55[b] (6)	1.56 ± 0.58[b] (7)
Comparative gene identification 58 (*Cgi58*)	1.05 ± 0.34[a] (7)	1.38 ± 0.66[ab] (6)	1.75 ± 0.59[b] (7)
Adenylate cyclase 3 (*Adcy3*)	1.07 ± 0.42 (8)		1.08 ± 0.23 (8)
Lipoprotein lipase (*Lpl*)	1.02 ± 0.23[a] (7)	0.60 ± 0.17[b] (6)	0.50 ± 0.23[b] (6)
Carnitine palmitoyltransferase Ia (*Cpt1a*)	1.01 ± 0.19[a] (14)	0.83 ± 0.17[b] (7)	0.68 ± 0.18[b] (11)
Long-chain 3-hydroxyacyl-CoA dehydrogenase (*Lchad*)	1.04 ± 0.28 (9)		1.01 ± 0.25 (9)
Acetyl-CoA acyltransferase 2 (*Acaa2*)	1.08 ± 0.44 (7)		1.23 ± 0.23 (6)
Long-chain acyl-CoA dehydrogenase (*Lcad*)	1.00 ± 0.09 (8)	0.93 ± 0.22 (7)	0.78 ± 0.30 (8)
Medium-chain acyl-CoA dehydrogenase (*Mcad*)	1.03 ± 0.27[a] (9)	0.57 ± 0.12[b] (6)	0.65 ± 0.20[b] (6)
Uncoupling protein 1 (*Ucp1*)	1.04 ± 0.32 (7)	1.12 ± 0.40 (7)	0.98 ± 0.34 (7)
Acyl-CoA oxidase-1 (*Acox1*)	1.04 ± 0.32[a] (8)	0.89 ± 0.32[a] (7)	0.69 ± 0.32[b] (8)
Acyl-CoA oxidase-2 (*Acox2*)	1.07 ± 0.42 (9)		1.17 ± 0.43 (9)
Insulin and IGF-1 signaling			
Insulin receptor (*Insr*)	1.05 ± 0.37 (9)	1.35 ± 0.42 (6)	1.36 ± 0.70 (8)
Insulin receptor substrate-1 (*Irs1*)	0.97 ± 0.43 (6)	0.97 ± 0.43 (4)	0.94 ± 0.59 (6)
Insulin receptor substrate-2 (*Irs2*)	1.27 ± 1.01 (9)	1.27 ± 0.87 (7)	1.41 ± 0.99 (8)
Phosphoinositide 3-kinase (*Pi3k*)	1.04 ± 0.28 (7)		1.17 ± 0.24 (7)
Insulin-like growth factor-1 (*Igf1*)	1.04 ± 0.31[a] (8)	1.16 ± 0.29[a] (7)	2.09 ± 0.80[b] (6)
IGF binding protein 1 (*Igfbp1*)	1.04 ± 0.30[a] (6)	0.45 ± 0.33[b] (6)	0.09 ± 0.02[b] (5)
Forkhead box protein a2 (*Foxa2*)	1.06 ± 0.37[a] (9)	1.98 ± 0.86[b] (6)	1.76 ± 0.48[b] (7)
Glucose metabolism and gluconeogenesis			
Glucose transporter type 4 (*Glut4*)	1.06 ± 0.38[a] (6)	0.88 ± 0.28[a] (5)	0.61 ± 0.25[b] (6)
Carbohydrate response element binding protein (*Chrebp*)	1.03 ± 0.26 (5)		1.33 ± 0.43 (6)
Fibroblast growth factor 21 (*Fgf21*)	1.07 ± 0.46[a] (5)		0.19 ± 0.12[b] (7)
Glucose-6-phosphatase (*G6p*)	1.07 ± 0.41[a] (9)	0.47 ± 0.17[b] (7)	0.66 ± 0.19[b] (6)
Phosphoenolpyruvate carboxykinase (*Pepck*)	1.04 ± 0.29[a] (8)	0.53 ± 0.22[b] (6)	0.51 ± 0.19[b] (7)
NADPH-producing			
Malic enzyme (*Malic*)	0.99 ± 0.32[a] (8)	0.98 ± 0.39[a] (7)	1.72 ± 0.70[b] (7)
Glucose-6-phosphate dehydrogenase (*G6pd*)	1.06 ± 0.38 (6)	1.14 ± 0.34 (4)	1.05 ± 0.46 (4)
6-Phosphogluconate dehydrogenase (*Pgd*)	1.05 ± 0.35[a] (7)		1.73 ± 0.56[b] (6)
Others			
AMP-activated protein kinase (*Ampk*)	1.07 ± 0.41 (7)		1.48 ± 0.32 (7)
Liver kinase B1 (*Lkb1*)	1.02 ± 0.23 (5)		1.10 ± 0.12 (5)

Table 3. Relative Expression of mRNAs in the liver

Genes	wt	pai/wt	pai/pai
Nuclear receptor subfamily 3 group C member 1 (*Nr3c1*)	1.07 ± 0.39 (6)		2.01 ± 1.05 (6)
CD11c	1.04 ± 0.31[a] (7)	0.57 ± 0.26[b] (6)	0.53 ± 0.27[b] (7)
Cyclophilin	1.01 ± 0.17 (9)	1.04 ± 0.28 (6)	1.01 ± 0.16 (6)
Pai	1.01 ± 0.43[a] (5)	1674 ± 569[b] (6)	6059 ± 2317[c] (6)

Numbers in brackets indicate the size of tested samples. Each value represents the mean ± SD. Different letters in superscript indicate $p < 0.05$.

Compared to wt samples, pai/wt livers exhibited normal except for swollen
cells (Fig. 6 E). However, among the forty tested genes (Table 3), the mRNA levels of
eight genes (Hmgcr, Ldlr, Ppar-γ, Dgat2, Acox1, Igf1, Glut4, and Malic) were
significantly ($p < 0.05$) different from those in pai/pai livers, but not different from
those in wt samples. The remaining 32 genes had similar transcription levels to
pai/pai samples, and 13 (33%) of them were markedly different from those in wt
samples ($p < 0.05$). Specifically, these 13 genes that exhibited different expressions
from wt livers included four up-regulated (Mgl, Srebp1c, Insig1, and Foxa2) and nine
down-regulated (Lpl, Cpt1a, Mcad, Srebp1a, Insig2, Igfbp1, G6p, Pepck, and Cd11c).
These results suggest that gene expression is genotype-specific and that the process of
lipolysis and gluconeogenesis is impaired in pai/wt mice.

3. 8. Hypothalamic analysis

To determine whether t10c12-CLA affected the hypothalamus, critical
hypothalamic genes and proteins were also measured. The mRNA levels of the leptin
receptor and agouti-related peptide were increased, while the transcription levels of
the ghrelin receptor and Orexins decreased in the pai/wt or pai/pai hypothalamus
when compared to wt mice (Fig. 7 A). Western blot analysis revealed no changes in
AMPK, phosphorylated AMPK, and GRP78 proteins in the pai/pai hypothalamus,
additionally, the ratio of pAMPK/AMPK did not show a significant difference
between wt and pai/pai samples ($p > 0.05$; Fig. 7 B-C). These findings suggest that the
central regulation of energy intake may only be affected at the transcriptional level.

Figure 7. Analyses of the relative mRNAs (A), proteins levels and their relative intensities in the hypothalamus (B-C). Agrp, Agouti-related peptide; Ampk, AMP-activated protein kinase; Ghsr, Ghrelin receptor; Grp78, glucose-regulated protein 78; Insr, Insulin receptor; Lepr, Leptin receptor; Mc4r, Melanocortic 4 receptor; Nr3c1, Nuclear receptor subfamily 3 group C member 1; (p)AMPK, (phosphorylated) AMP-activated protein kinase; Pomc, Proopiomelanocortin. The sample sizes for RNA analysis are 6~9 in each group. The bars represent the mean ± SD. Different letters indicate $p < 0.05$.

4. DISCUSSION

It has been reported that t10c12-CLA can depress milk fat production by inhibiting lipogenesis [5-7], reducing the globule size of milk fat droplets [8]. A study on grazing herds indicated that other fatty acids, apart from t10c12-CLA, may also inhibit milk fat synthesis during milk fat depression [18]. These results suggest that this milk fat depression may result from t10c12-CLA directly or other compositions induced by t10c12-CLA indirectly. Research on early-weaning lambs has revealed that t10c12-CLA can facilitate the weaning transition through the microbiota-host transcriptome interaction [19]. In the current study, maternal t10c12-CLA plays an active role in development in female offspring, similar to male offspring fostered by pai/wt mother (Rao et al, 2023 bioRxiv preprint: https://doi.org/10.1101/2023.01.31.526123). It suggests that maternal t10c12-CLA benefits embryo and/or offspring development.

Our previous observation showed that t10c12-CLA reduced body fat by releasing heat via BAT activation in pai/pai male mice with lower blood TGs (Rao et al, 2023 bioRxiv preprint: https://doi.org/10.1101/2023.01.31.526123). Conversely, in the pai female mice, t10c12-CLA did not cause body fat loss but resulted in complicated symptoms, including hypothermia, fatty liver and hypertriglyceridemia, and highly responsive to stress and inflammation in a dose-dependent manner. The different physiological and pathological characteristics of both sexes can help us understand the mechanism of action of t10c12-CLA in gender-specific and dose-dependent manners. As for pai mice, there are two problems we have not fixed until now. Firstly, we have reported as many results as possible for a comprehensive understanding of the pai mouse, but a few parameters are difficult to be explained well, and even show contradictions in some respects. Secondly, explaining why there are only minimal differences in t10c12-CLA and linoleic acid is challenging although we once observed drastic changes in an increase of product t10c12-CLA and a decrease of substrate linoleic acid in pai-transfected 3T3 cells [18].

The metabolic pathway of t10c12-CLA, similar to that of linoleic or alpha-linolenic acids [20], suggests it can competitively interrupt the metabolic pathway of n-6 or n-3 FAs and their derived leukotrienes signals/hormonal molecules. It also indicates that t10c12-CLA can be directly converted into the analogues of n-6 FAs and their derivatives. PGE2, an arachidonic acid-derived hormone, will be easily disturbed

by t10c12-CLA. PGE2 is involved in stress response and has essential effects on inflammation, fever, and renal filtration [21]. It can induce corticosterone secretion by regulating the hypothalamic-pituitary-adrenal axis activity and enhancing adrenaline synthesis in the Chromaffin cells of the adrenal medulla. Subsequently, the increased circulating corticosterone and adrenaline could stimulate excess glucagon secretion [22]. In pai/pai female mice, we have observed the excess PGE2 induced by t10c12-CLA, and then the PGE2 (or analogues) caused the increases in corticosterone, glucagon, adrenaline, and inflammatory factors (IL-6, TNFα, and CRP). Obviously, hypothermia in pai/pai female mice may be related to hyperprostaglandinemia, hyperadrenerginemia and inflammatory response; while hyperglyceridemia and fatty liver are associated with hypercorticosteroidemia and glucagon resistance.

In the hypothalamic preoptic area, PGE2 receptors EP3, coupled to cAMP decreases, can stimulate a passive defence mechanism such as less O_2 consumption and hypothermia to avoid further energy expenditure; while receptors EP4, connected to cAMP increases, can stimulate an active defence mechanism, including more O_2 consumption and febrile responses by increasing BAT thermogenesis, typically occurs to promote well-being and survival [21,23]. It suggests that gender-based differences in PGE2 levels might partially result in the opposite heat release in male and female pai/pai mice. In female mice, the co-elevation of PGE2 and corticosterone suggests that t10c12-CLA induced a passive defence mechanism including hypothermia which conversely promote BAT thermogenesis and lipid metabolism by negative feedback to down-regulate pAMPK and up-regulate UCP1, UCP2, CPT1B, and pHSL. In contrast, in male mice, the lack of PGE2 elevation suggests that t10c12-CLA induced an active defence mechanism including hyperthermia to promote survival (Rao et al, 2023 bioRxiv preprint: https://doi.org/10.1101/2023.01.31.526123).

In addition to PGE2- and adrenaline-induced heating [24], BAT thermogenesis can be mediated through the central GRP78 [25,26] or AMPK-OREXINS-BMP8B pathway [27]. Orexin-null mice exhibited periods of decreased activity during the dark phase and demonstrated hypothermia, reduced oxygen consumption, and blunted thermogenic responses [28]; furthermore, BMP8B's thermogenic effect is sexually dimorphic and observed in females only [27]. The pai/pai female mice exhibited similar phenotypes to orexin-null mice and lower transcription levels of Orexins in the pai hypothalamus. These characteristics suggest that the t10c12-CLA-induced BAT thermogenesis may be regulated through central pathways in female mice.

Unfortunately, we did not observe evident changes in GRP78, AMPK, pAMPK, and
pAMPK/AMPK ratio in the hypothalamus.

Previous studies on mice have yielded conflicting results on whether dietary
t10c12-CLA leads to hepatic steatosis [15,29-31]. The current observation of fatty liver
and hypertriglyceridemia in pai/pai mice, which may be associated with t10c12-CLA-
induced excess corticosterone for the overproduction of corticosterone can lead to
Cushing syndrome, corticosteroid-induced lipodystrophy, hypertriglyceridemia, or
hepatic steatosis [32,33]. On the other hand, the existence of glucagon resistance can
increase liver fat [34-36], although chronic increases in plasma glucagon promote
lipolysis through INSP3R1/CAMKII/ATGL [37] or PKA/HSL pathways in
hepatocytes [38] and stimulate beta-oxidation by activating AMPK/PPAR-
α/CPT1A/MCAD pathway [22]. In the current study, the serum glucagon
concentration increased nearly twofold in pai/pai female mice, which is similar to its
physiological levels (2~3 fold) in response to long-term fasting or hypoglycemia [39],
whereas the protein levels of unchanged ATGL, decreased pAMPK, and increased
CPT1A indicate the chronic glucagon excess did not stimulate the lipolysis and beta-
oxidation in the liver. These results suggest that the chronic impact of t10c12-CLA
might give rise to glucagon resistance in pai/pai female mice. In this case, the down-
regulated hepatic enzymes may not fully and immediately catalyse the FAs molecules
mobilised from other tissues based on glucagon stimulation. Consequently, it disrupts
hepatic and systemic lipid homeostasis and leads to dyslipidemia and fatty liver in
female mice. Contrarily, the t10c12-CLA did not cause the overproduction of
corticosterone or glucagon in pai/pai male mice, thus, the mice had no fatty liver
disease and even exhibited lower TGs levels in the blood.

FGF21 is a key regulator of the specific hormonal signalling pathway in
increasing energy expenditure, reducing hepatic and plasma TGs levels, accelerating
lipoprotein catabolism in WAT and BAT [40], and suppressing physical activity via
central action [41]. FGF21 signalling enhances insulin sensitivity and lowers serum
glucose by acting directly on adipose tissues [17,42]. As we know, the pai/wt mouse is
the first animal that t10c12-CLA can induce FGF21 secretion. Excess FGF21 is a unique
feature in pai/wt, not in pai/pai genotypes, indicating that t10c12-CLA-induced
FGF21 secretion is specifically associated with low doses in both genders and it can be
suppressed to the normal levels through negative feedback when high doses occurred
in pai/pai mice. This suggests that the health benefits of low-dose t10c12-CLA may be

due, in part, to its ability to stimulate FGF21 secretion. Therefore, the different effects

of t10c12-CLA on glucose metabolism in diabetic Zucker fatty rats [43] and insulin

resistance in mice [44,45] may be related to the different levels of FGF21 induced by

t10c12-CLA. In the current study, excess FGF21 may be associated with specific

phenotypes in pai/wt mice, such as blood glucose homeostasis during starvation,

enlarged white adipocytes, no steatosis in hepatocytes and normal circulating TGs.

The exact mechanism by which t10c12-CLA activates FGF21 secretion remains unclear

and requires further investigation.

Figure 8. Diagram showing the dose-dependent effect of chronic t10c12-CLA on lean

female mice. Besides the common effects, such as increased adrenaline, IL-6, and TNFα, as

well as decreased heat release and oxygen consumption (Middle panel), the low doses

specifically increase circulating levels of FGF21 which plays active roles in maintaining

energy homeostasis, suppressing physical activity, and improving serum glucose levels in

pai/wt mice (Left panel). Conversely, the high doses specifically result in excesses of

prostaglandin E2, corticosterone, glucagon, and C reaction protein; subsequently, the

combination of these factors causes metabolic syndrome, including hypothermia, fatty liver,

and hypertriglyceridemia in pai/pai mice (Right panel). Words in black and red describe

physiological features and endocrinal factors, respectively.

Based on the observation in female animals in this study, we suggest that low
doses of t10c12-CLA may cause increases in circulating adrenaline which can induce
WAT lipolysis and BAT heating through a mechanism of negative feedback, and
FGF21 which can promote energy expenditure, reduce hepatic and plasma TGs levels,
accelerate lipoprotein catabolism in adipocytes, suppress physical activity, and
enhance insulin sensitivity and lower serum glucose (as shown in Fig. 8). Conversely,
high doses of t10c12 results in an increased PEG2 which induces increases in
corticosterone, glucagon, adrenaline, and inflammatory factors. The combination of
these factors causes metabolic syndrome, including fatty liver and hyperglyceridemia,
hyperprostaglandinemia and hypothermia, and inflammatory changes in female mice
(as shown in Fig. 8). The metabolic impact of t10c12-CLA involves various hormones
and inflammatory factors and is system-wide effects on multiple metabolic pathways.
Thus, interpreting the data from pai mice requires careful consideration, and more
work is needed to understand the mechanism of action of t10c12-CLA and its
metabolic impact.

Totally, our results suggest that the chronic effects of t10c12-CLA on lean
animals are dose-dependent and differ between sexes and low doses of t10c12-CLA
are beneficial while high doses may become a health risk in lean female mammals.
Further research is necessary to understand the health benefits and safety risks of
t10c12-CLA on pai obese mice.

Abbreviations

t10c12-CLA, trans 10, cis 12-conjugated linoleic acid; LA, linoleic aci; pai,
Propionibacterium acnes isomerase; wt, wild-type; FGF21, fibroblast growth factor 21;
UCP, uncoupling protein; AMPK, AMP-activated protein kinase; pAMPK,
phosphorylated AMPK; HSL, hormone-sensitive lipase; pHSL, phosphorylated HSL;
CPT1, carnitine palmitoyltransferase I; BAT, brown adipose tissue; WAT, white
adipose tissue; FAs, fatty acids

Ethics approval and consent to participate:

All animal experiments followed the Committee for Experimental Animals of
Yangzhou University.

Institutional Review Board Statement:

The animal study protocol was approved by the Ethics Committee of Yangzhou University (protocol code NSFC2020-SYXY-20 and dated 25 March 2020).

Consent for publication:

Not applicable

Availability of data and materials:

The datasets used and/or analysed during the current study are available from the corresponding author upon reasonable request.

Competing interests:

The authors declare that they have no competing interests.

Funding sources

This work was supported by the National Key Research and Development Program of China (2018YFC1003800), the Project Funded by the Priority Academic Program Development of Jiangsu Higher Education Institutions (PAPD), and Jiangsu Funding Program for Excellent Postdoctoral Talent (Rao Y).

Acknowledgement

We thank graduate students Qi Wang, Mei Liu, and Jiaqi Lu in our lab for their generous assistance in partial tests and Ms Yuyang Wang in our university for technique help in gas chromatography. We also thank Professor Hongjun Wang at the Medical University of South Carolina, USA, for the helpful discussion during the writing process and Ms Wenjing Gou's language assistance in preparing this manuscript.

Authors 'contributions:

Conceptualization, Y Rao and K Gou; Data curation, Y Rao, M Li, B Wang, Y Wang, and L Liang; Formal analysis, Y Rao and K Gou; Funding acquisition, Y Rao and K Gou; Investigation, Y Rao, M Li, B Wang, Y Wang and L Liang; Methodology, Y Rao, M Li, B Wang, Y Wang and L Liang; Project administration, K Gou; Software,

Y Rao; Supervision, K Gou; Validation, K Gou; Visualization, Y Rao and L Liang; Writing – original draft, Y Rao; Writing – review & editing, K Gou. All authors proofread, made comments, and approved the report.

Supplementary data

This manuscript contains supplementary data.

References

1. Shen, W.; Baldwin, J.; Collins, B.; Hixson, L.; Lee, K.T.; Herberg, T.; Stames, J.; Cooney, P.; Chuang, C.C.; Hopkins, R., et al. Low level of trans-10, cis-12 conjugated linoleic acid decreases adiposity and increases browning independent of inflammatory signaling in overweight Sv129 mice. *Journal of Nutritional Biochemistry* **2015**, *26*, 616-625, doi:10.1016/j.jnutbio.2014.12.016.

2. den Hartigh, L.J.; Wang, S.R.; Goodspeed, L.; Wietecha, T.; Houston, B.; Omer, M.; Ogimoto, K.; Subramanian, S.; Gowda, G.A.N.; O'Brien, K.D., et al. Metabolically distinct weight loss by 10,12 CLA and caloric restriction highlight the importance of subcutaneous white adipose tissue for glucose homeostasis in mice. *Plos One* **2017**, *12*, doi:10.1371/journal.pone.0172912.

3. Berkel, C.; Cacan, E. Trans-10, cis-12 conjugated linoleic acid- and caloric restriction-mediated upregulation of CNDP2 expression in white adipose tissue in rodents, with implications in feeding and obesity. *J Nutr Biochem* **2023**, *114*, 109269, doi:10.1016/j.jnutbio.2023.109269.

4. Wang, L.; Zhang, S.; Huang, Y.; You, W.; Zhou, Y.; Chen, W.; Sun, Y.; Yi, W.; Sun, H.; Xie, J., et al. CLA improves the lipo-nutritional quality of

pork and regulates the gut microbiota in Heigai pigs. *Food Funct* **2022**,

13, 12093-12104, doi:10.1039/d2fo02549c.

5.   Baumgard, L.H.; Sangster, J.K.; Bauman, D.E. Milk fat synthesis in dairy

cows is progressively reduced by increasing supplemental amounts of

trans-10, cis-12 conjugated linoleic acid (CLA). *J Nutr* **2001**, *131*, 1764-

1769, doi:10.1093/jn/131.6.1764.

6.   Harvatine, K.J.; Robblee, M.M.; Thorn, S.R.; Boisclair, Y.R.; Bauman, D.E.

Trans-10, cis-12 CLA dose-dependently inhibits milk fat synthesis without

disruption of lactation in C57BL/6J mice. *J Nutr* **2014**, *144*, 1928-1934,

doi:10.3945/jn.114.198911.

7.   Harvatine, K.J.; Boisclair, Y.R.; Bauman, D.E. Time-dependent effect of

trans-10,cis-12 conjugated linoleic acid on gene expression of lipogenic

enzymes and regulators in mammary tissue of dairy cows. *J Dairy Sci*

2018, *101*, 7585-7592, doi:10.3168/jds.2017-13935.

8.   Han, L.; Huang, Q.; Yang, J.; Lu, W.; Hu, M.; Yang, Y.; Zhu, H.; Pang, K.;

Yang, G. Proteomic analysis of milk fat globule membranes from small-

sized milk fat globules and their function in promoting lipid droplet

fusion in bovine mammary epithelial cells. *Food Funct* **2023**, *14*, 2304-

2312, doi:10.1039/d2fo03476j.

9.   Kanter, J.E.; Goodspeed, L.; Wang, S.; Kramer, F.; Wietecha, T.; Gomes-

Kjerulf, D.; Subramanian, S.; O'Brien, K.D.; den Hartigh, L.J. 10,12

Conjugated Linoleic Acid-Driven Weight Loss Is Protective against

Atherosclerosis in Mice and Is Associated with Alternative Macrophage

Enrichment in Perivascular Adipose Tissue. *Nutrients* **2018**, *10*,

doi:10.3390/nu10101416.

10.   Vaisar, T.; Wang, S.; Omer, M.; Irwin, A.D.; Storey, C.; Tang, C.; den
Hartigh, L.J. 10,12-Conjugated linoleic acid supplementation improves
HDL composition and function in mice. *J Lipid Res* **2022**, *63*, 100241,
doi:10.1016/j.jlr.2022.100241.

11.   Kim, J.H.; Kim, Y.; Kim, Y.J.; Park, Y. Conjugated linoleic acid: Potential
health benefits as a functional food ingredient. *Annu Rev Food Sci*
Technol **2016**, *7*, 221-244, doi:10.1146/annurev-food-041715-033028.

12.   Marques, T.M.; Wall, R.; O'Sullivan, O.; Fitzgerald, G.F.; Shanahan, F.;
Quigley, E.M.; Cotter, P.D.; Cryan, J.F.; Dinan, T.G.; Ross, R.P., et al.
Dietary trans-10, cis-12-conjugated linoleic acid alters fatty acid
metabolism and microbiota composition in mice. *Br J Nutr* **2015**, *113*,
728-738, doi:10.1017/S0007114514004206.

13.   Cuciniello, R.; Luongo, D.; Ferramosca, A.; Lunetti, P.; Rotondi-Aufiero,
V.; Crispi, S.; Zara, V.; Maurano, F.; Filosa, S.; Bergamo, P. Conjugated
linoleic acid downregulates Alzheimer's hallmarks in aluminum mouse
model through an Nrf2-mediated adaptive response and increases brain
glucose transporter levels. *Free Radic Biol Med* **2022**, *191*, 48-58,
doi:10.1016/j.freeradbiomed.2022.08.027.

14.   Sisk, M.B.; Hausman, D.B.; Martin, R.J.; Azain, M.J. Dietary conjugated
linoleic acid reduces adiposity in lean but not obese Zucker rats. *J Nutr*
2001, *131*, 1668-1674, doi:10.1093/jn/131.6.1668.

15.   Li, S.L.; Ma, S.Y.; Xu, B.R.; Fan, Z.Y.; Li, M.J.; Cao, W.G.; Gou, K.M.
Effects of trans-10, cis-12-conjugated linoleic acid on mice are influenced
by the dietary fat content and the degree of murine obesity. *Eur J Lipid*
Sci Tech **2015**, *117*, 1908-1918, doi:10.1002/ejlt.201400568.

16.   den Hartigh, L.J. Conjugated Linoleic Acid Effects on Cancer, Obesity,
672   and Atherosclerosis: A Review of Pre-Clinical and Human Trials with
673   Current Perspectives. *Nutrients* **2019**, *11*, doi:10.3390/nu11020370.

674   17.   Flippo, K.H.; Potthoff, M.J. Metabolic Messengers: FGF21. *Nat Metab*
675   **2021**, *3*, 309-317, doi:10.1038/s42255-021-00354-2.

676   18.   Neville, O.B.; Fahey, A.G.; Mulligan, F.J. Comparison of milk and grass
677   composition from grazing Irish dairy herds with and without milk fat
678   depression. *Ir Vet J* **2023**, *76*, 5, doi:10.1186/s13620-023-00230-3.

679   19.   Yang, C.; Deng, X.; Lund, P.; Liu, H.; Ding, X.; Fu, Z.; Zhang, N.; Li, J.;
680   Dong, L. Rumen microbiota-host transcriptome interaction mediates the
681   protective effects of trans-10, cis-12 CLA on facilitating weaning
682   transition   of   lambs.   *Anim   Nutr*   **2023**,   *12*,   345-359,
683   doi:10.1016/j.aninu.2022.11.002.

684   20.   Subbaiah, P.V.; Gould, I.G.; Lal, S.; Aizezi, B. Incorporation profiles of
685   conjugated linoleic acid isomers in cell membranes and their positional
686   distribution in phospholipids. *Biochim Biophys Acta* **2011**, *1811*, 17-24,
687   doi:10.1016/j.bbalip.2010.09.004.

688   21.   Furuyashiki, T.; Narumiya, S. Stress responses: the contribution of
689   prostaglandin E(2) and its receptors. *Nat Rev Endocrinol* **2011**, *7*, 163-
690   175, doi:10.1038/nrendo.2010.194.

691   22.   Zeigerer, A.; Sekar, R.; Kleinert, M.; Nason, S.; Habegger, K.M.; Muller,
692   T.D. Glucagon's metabolic action in health and disease. *Compr Physiol*
693   **2021**, *11*, 1759-1783, doi:10.1002/cphy.c200013.

694   23.   Saper, C.B.; Lowell, B.B. The hypothalamus. *Curr Biol* **2014**, *24*, R1111-
695   1116, doi:10.1016/j.cub.2014.10.023.

24.   Cypess, A.M.; Weiner, L.S.; Roberts-Toler, C.; Franquet Elia, E.; Kessler,
S.H.; Kahn, P.A.; English, J.; Chatman, K.; Trauger, S.A.; Doria, A., et al.
Activation of human brown adipose tissue by a beta3-adrenergic receptor
agonist. *Cell Metab* **2015**, *21*, 33-38, doi:10.1016/j.cmet.2014.12.009.

25.   Contreras, C.; Gonzalez-Garcia, I.; Seoane-Collazo, P.; Martinez-Sanchez,
N.; Linares-Pose, L.; Rial-Pensado, E.; Ferno, J.; Tena-Sempere, M.;
Casals, N.; Dieguez, C., et al. Reduction of hypothalamic endoplasmic
reticulum stress activates browning of white fat and ameliorates obesity.
Diabetes **2017**, *66*, 87-99, doi:10.2337/db15-1547.

26.   Contreras, C.; Fondevila, M.F.; Lopez, M. Hypothalamic GRP78, a new
target against obesity? *Adipocyte* **2018**, *7*, 63-66,
doi:10.1080/21623945.2017.1405878.

27.   Martins, L.; Seoane-Collazo, P.; Contreras, C.; Gonzalez-Garcia, I.;
Martinez-Sanchez, N.; Gonzalez, F.; Zalvide, J.; Gallego, R.; Dieguez, C.;
Nogueiras, R., et al. A functional link between ampk and orexin mediates
the effect of bmp8b on energy balance. *Cell Rep* **2016**, *16*, 2231-2242,
doi:10.1016/j.celrep.2016.07.045.

28.   Chemelli, R.M.; Willie, J.T.; Sinton, C.M.; Elmquist, J.K.; Scammell, T.; Lee,
C.; Richardson, J.A.; Williams, S.C.; Xiong, Y.M.; Kisanuki, Y., et al.
Narcolepsy in orexin knockout mice: Molecular genetics of sleep
regulation. *Cell* **1999**, *98*, 437-451, doi:Doi 10.1016/S0092-8674(00)81973-
X.

29.   Ashwell, M.S.; Ceddia, R.P.; House, R.L.; Cassady, J.P.; Eisen, E.J.; Eling,
T.E.; Collins, J.B.; Grissom, S.F.; Odle, J. Trans-10, cis-12-conjugated
linoleic acid alters hepatic gene expression in a polygenic obese line of

mice displaying hepatic lipidosis. *J Nutr Biochem* **2010**, *21*, 848-855, doi:10.1016/j.jnutbio.2009.06.013.

30. Vyas, D.; Kadegowda, A.K.; Erdman, R.A. Dietary conjugated linoleic acid and hepatic steatosis: species-specific effects on liver and adipose lipid metabolism and gene expression. *J Nutr Metab* **2012**, *2012*, 932928, doi:10.1155/2012/932928.

31. Cordoba-Chacon, J.; Sugasini, D.; Yalagala, P.C.R.; Tummala, A.; White, Z.C.; Nagao, T.; Kineman, R.D.; Subbaiah, P.V. Tissue-dependent effects of cis-9,trans-11- and trans-10,cis-12-CLA isomers on glucose and lipid metabolism in adult male mice. *J Nutr Biochem* **2019**, *67*, 90-100, doi:10.1016/j.jnutbio.2019.01.020.

32. Laryea, G.; Schutz, G.; Muglia, L.J. Disrupting hypothalamic glucocorticoid receptors causes HPA axis hyperactivity and excess adiposity. *Mol Endocrinol* **2013**, *27*, 1655-1665, doi:10.1210/me.2013-1187.

33. Nicolaides, N.C.; Charmandari, E.; Kino, T.; Chrousos, G.P. Stress-related and circadian secretion and target tissue actions of glucocorticoids: Impact on health. *Front Endocrinol (Lausanne)* **2017**, *8*, 70, doi:10.3389/fendo.2017.00070.

34. Krilov, L.; Nguyen, A.; Miyazaki, T.; Unson, C.G.; Bouscarel, B. Glucagon receptor recycling: role of carboxyl terminus, beta-arrestins, and cytoskeleton. *Am J Physiol-Cell Ph* **2008**, *295*, C1230-C1237, doi:10.1152/ajpcell.00240.2008.

35. Charbonneau, A.; Unson, C.G.; Lavoie, J.M. High-fat diet-induced hepatic steatosis reduces glucagon receptor content in rat hepatocytes: potential

interaction with acute exercise. *J Physiol-London* **2007**, *579*, 255-267, doi:10.1113/jphysiol.2006.121954.

36. Sammons, M.F.; Lee, E.C.Y. Recent progress in the development of small-molecule glucagon receptor antagonists. *Bioorg Med Chem Lett* **2015**, *25*, 4057-4064, doi:10.1016/j.bmcl.2015.07.092.

37. Perry, R.J.; Zhang, D.; Guerra, M.T.; Brill, A.L.; Goedeke, L.; Nasiri, A.R.; Rabin-Court, A.; Wang, Y.; Peng, L.; Dufour, S., et al. Glucagon stimulates gluconeogenesis by INSP3R1-mediated hepatic lipolysis. *Nature* **2020**, *579*, 279-283, doi:10.1038/s41586-020-2074-6.

38. Hayashi, Y. Glucagon regulates lipolysis and fatty acid oxidation through inositol triphosphate receptor 1 in the liver. *J Diabetes Investig* **2021**, *12*, 32-34, doi:10.1111/jdi.13315.

39. Kleinert, M.; Sachs, S.; Habegger, K.M.; Hofmann, S.M.; Muller, T.D. Glucagon regulation of energy expenditure. *Int J Mol Sci* **2019**, *20*, doi:10.3390/ijms20215407.

40. Schlein, C.; Talukdar, S.; Heine, M.; Fischer, A.W.; Krott, L.M.; Nilsson, S.K.; Brenner, M.B.; Heeren, J.; Scheja, L. Fgf21 lowers plasma triglycerides by accelerating lipoprotein catabolism in white and brown adipose tissues. *Cell Metab* **2016**, *23*, 441-453, doi:10.1016/j.cmet.2016.01.006.

41. Bookout, A.L.; de Groot, M.H.; Owen, B.M.; Lee, S.; Gautron, L.; Lawrence, H.L.; Ding, X.; Elmquist, J.K.; Takahashi, J.S.; Mangelsdorf, D.J., et al. FGF21 regulates metabolism and circadian behavior by acting on the nervous system. *Nat Med* **2013**, *19*, 1147-1152, doi:10.1038/nm.3249.

771   42.   Holland, W.L.; Adams, A.C.; Brozinick, J.T.; Bui, H.H.; Miyauchi, Y.;

772   Kusminski, C.M.; Bauer, S.M.; Wade, M.; Singhal, E.; Cheng, C.C., et al.

773   An FGF21-adiponectin-ceramide axis controls energy expenditure and

774   insulin action in mice. *Cell Metab* **2013**, *17*, 790-797,

775   doi:10.1016/j.cmet.2013.03.019.

776   43.   Henriksen, E.J.; Teachey, M.K.; Taylor, Z.C.; Jacob, S.; Ptock, A.; Kramer,

777   K.; Hasselwander, O. Isomer-specific actions of conjugated linoleic acid

778   on muscle glucose transport in the obese Zucker rat. *Am J Physiol*

779   *Endocrinol Metab* **2003**, *285*, E98-E105, doi:10.1152/ajpendo.00013.2003.

780   44.   Tsuboyama-Kasaoka, N.; Takahashi, M.; Tanemura, K.; Kim, H.J.; Tange,

781   T.; Okuyama, H.; Kasai, M.; Ikemoto, S.; Ezaki, O. Conjugated linoleic

782   acid supplementation reduces adipose tissue by apoptosis and develops

783   lipodystrophy in mice. *Diabetes* **2000**, *49*, 1534-1542,

784   doi:10.2337/diabetes.49.9.1534.

785   45.   Clement, L.; Poirier, H.; Niot, I.; Bocher, V.; Guerre-Millo, M.; Krief, S.;

786   Staels, B.; Besnard, P. Dietary trans-10,cis-12 conjugated linoleic acid

787   induces hyperinsulinemia and fatty liver in the mouse. *J Lipid Res* **2002**,

788   *43*, 1400-1409, doi:10.1194/jlr.m20008-jlr200.