Thermoregulatory consequences of growing up during a heatwave

Authors: Elin Persson[1], Andreas Nord[1]

[1] Lund University, Department of Biology, Section for Evolutionary Ecology, Sölvegatan 37, SE-223
63 Lund, Sweden

* Author for correspondence: Elin Persson, elin.persson@biol.lu.se

ORCID numbers:

Elin Persson: 0000-0003-1824-1403

Andreas Nord: 0000-0001-6170-689X

Keywords: bird, development, evaporative cooling, heatwave, metabolic rate, thermoregulation

AUTHOR CONTRIBUTIONS: Andreas Nord conceived the idea. Elin Persson and Andreas Nord
performed the practical work. Elin Persson analysed the data and both authors interpreted the results.
Elin Persson wrote the first draft of the manuscript, which was edited by Andreas Nord. Both authors
approved submission and agreed to be accountable for all contents. Andreas Nord procured funding.

ACKNOWLEDGEMENTS: Lars Råberg and Fredrik Andreasson provided helpful comments that
improved a previous draft of the manuscript. We thank Camilla Björklöv and Agnieszka Czopek for
assistance in caring for the birds, Lars Fredriksson for all the help with building pens and chambers
and Cecilia Thomasson for providing equipment.

DATA AVAILABILITY: Data will be deposited in Figshare upon acceptance of the manuscript.

CONFLICT OF INTEREST: We have no conflicts of interest to disclose.

Abstract

Climate change is causing an increase in the frequency and intensity of heatwaves. In birds, changes in

environmental temperature during the developmental period can affect growth, metabolism, and

temperature tolerance of the offspring. We know little about whether such changes remain to

adulthood, which is important to understand the links between climate change, development, and

fitness. We investigated if phenotypic consequences of the thermal environment in early life remained

in adulthood by raising Japanese quail (*Coturnix japonica*) under simulated heatwave- or control

conditions from hatching until halfway through the growth period, and then in a common garden until

reproductively mature. We measured biometric and thermoregulatory (metabolic heat production

[MHP], evaporative water and heat loss [EWL, EHL] and body temperature) responses to variation in

ambient temperature at the end of the heatwave treatment and in adulthood. Developmental

temperature did not affect body mass or wing length. However, warm birds had lower MHP than

control birds at the end of the heat wave period and, in the warmest temperature studied (40°C), warm

birds had higher evaporative cooling capacity than control birds. None of the effects found at the end

of the heatwave period remained until adulthood. This implies that chicks exposed to higher

temperatures could be more prepared to counter heat stress as juveniles, but that they do not enjoy any

advantages of such developmental conditions when facing high temperatures as adults.

Introduction

Climate change, including increased global mean temperature and increased climatic instability, has

strongly impacted animal populations, for example, through changes in species distributions' (Chen et

al., 2011), mass mortality events (McKechnie et al., 2012) and population extinctions (Thomas et al.,

2006). A warmer climate is also associated with a range of sublethal consequences that can negatively

affect fitness (Conradie et al., 2019). For example, many insectivorous passerines in the Northern

Hemisphere have advanced breeding start to match a warming-related advancement of peak food

abundance, but still pay fitness costs since phenotypic plasticity in egg laying cannot fully compensate

for the earlier emergence of insect prey (Shipley et al., 2020; Both et al., 2009; Visser et al., 1998;

Hansson et al., 2014). Moreover, when warming is acute and pronounced, such as during a heatwave,

parents might have to trade-off workload against the risk of overheating (Cunningham et al., 2013; du

Plessis et al., 2012; Nilsson and Nord, 2018) with implications for both current and future reproduction

(Sharpe et al., 2019; Cunningham et al., 2021; Nord & Nilsson, 2019; Andreasson et al., 2020).

Understanding the ontogeny and evolution of animals' ability to cope with fluctuating temperatures is

thus fundamental to predict the consequences of climate change on wildlife (McKechnie and Wolf,

2019; Albright et al., 2017).

Changes in temperature during the developmental period before or after hatching can affect subsequent

temperature tolerance (reviewed by Nord and Giroud, 2020). Conceptually, it can be hypothesised that

programming effects of developmental temperature on adult thermoregulation fall on a scale between

constraining and priming: 1) if early-life temperature programs phenotypes that perform optimally in

that thermal environment, adults should maximise fitness when early- and later-life environments are

similar ("environmental matching hypothesis"; reviewed by Monaghan, 2008); 2) if deviations from

optimal developmental conditions act suppressively on the phenotype, then increasing distance from

optimality in early life should come at a fitness cost regardless of the adult thermal environment ("silver

spoon hypothesis"; Monaghan, 2008). In line with the environmental matching hypothesis, rats reared

in warmer conditions grew their tails longer and feet larger and reduced the amount of body fat, which

was interpreted as adaptions to facilitate heat loss (Demicka and Caputa, 1993). However, in wild

systems, exposure to warmer developmental temperatures have been shown to have negative effects on

nestling growth and fledging success; heat-stressed chicks weighed less and grew more slowly than

individuals reared in control conditions (Deaton et al., 1978; Andreasson et al., 2018; Rodriguez and

Barba, 2016; Cunningham et al., 2013; van de Ven et al., 2020; Andrew et al., 2017). These findings

conform more closely with the silver spoon hypothesis. However, other studies demonstrate positive

effects of warmer developmental temperatures on nestling growth and fledgling success (McCarty and

Winkler, 1999; Dawson et al., 2005). For example, tree swallow (*Tachycineta bicolor*) nestlings gained

more mass due to both higher temperatures and higher food abundance (which, in turn, was also

positively affected by higher temperatures) (McCarty and Winkler, 1999), and both nestling survival
and fledging success increased with higher temperatures in another study on the same species (Dawson
et al., 2005).

We know less about the physiological basis of any matching between developmental and adult thermal
environments (reviewed by Nord & Giroud, 2020). Studies on poultry suggest that precisely timed and
dosed thermal stimuli during embryonic or perinatal development can lead to epigenetic programming
that makes the birds better at withstanding matched temperature stressors, such as altered body
temperature (T_b) and metabolic rate, that last at least until adolescence (Nichelmann, 2004; reviewed by
e.g. Farag and Alagawany, 2018). There are few comparable studies in wild systems. However,
Andreasson et al. (2018) found that blue tit (*Cyanistes caeruleus*) nestlings from heated nest boxes had
higher T_b and grew more slowly than nestlings from control nest boxes, indicating a constraining role
of high ambient temperatures acting through a change in distribution of resources between growth and
thermoregulation. Yet, heat-challenged nestlings in the study by Andreasson et al. (2018) tended to
survive better to the next breeding season compared to control birds, providing mixed evidence for the
constraining or priming role of developmental temperature on adult phenotype. Furthermore, Page et al.
(2022) found that blue tits that were experimentally heated as embryos were more cold tolerant as mature
nestlings than nestlings in non-heated nests. The above discrepancies suggest we need to know more
about how morphological and physiological responses to developmental temperature affect the capacity
for thermal acclimation in a broader range of circumstances, particularly with regards to whether any of
the reported phenotypic changes remain in adulthood. To fill this knowledge gap, we must measure the
ontogeny and legacy of responses to variation in developmental temperature, including changes in
metabolic rate, evaporative cooling capacity and T_b, both in the juvenile stage and in adulthood.

Here, we investigated if development during a simulated heatwave affected offspring thermal, metabolic
and hygric physiology, and if any such effects remained in adulthood several weeks after the heatwave
had passed. We studied this in Japanese quail (*Coturnix japonica*) that completed the first half of their
somatic development under simulated heatwave-like conditions or under normal thermal conditions, and
the second half in a common garden. Epigenetic adaptation to temperature appears during critical phases
of development (Yahav, 2015). Even though precocial species like Japanese quail are covered with
feathers and can thermoregulate from an early age, they are typically not fully homeothermic until some
weeks after hatch (Jørgensen & Blix, 1988; Nord & Nilsson, 2021; reviewed by Price and Dzialowski,
2018). Accordingly, incipient thermogenesis develops from late incubation and over the first 10 or so
days post-hatch (Yahav, 2015), suggesting there could be lasting effects on the thermoregulatory
phenotype if chicks complete the first part of their post-embryonic life in heatwave-like conditions. If

developmental conditions prime heat acclimation, we predicted, under the environmental matching hypothesis, that heatwave birds would have lower T_b and correspondingly lower metabolic heat production rate (MHP) and a higher capacity for evaporative cooling during heat stress (Fig. 1A). However, if high developmental temperature constrains development, we predicted, under the silver spoon hypothesis, that heatwave birds would have higher T_b, higher MHP and lover evaporative cooling capacity during a heat challenge (Fig. 1B). If warm temperatures during the early developmental period permanently modifies temperature tolerance, as shown in studies on poultry (e.g., Arjona et al., 1988; Piestun et al., 2011), we also predicted that these phenotypic traits would remain in adulthood (Figs. 1A, B). If, however, developmental temperature-effects are reversible, we predict that the change in traits caused by thermal acclimation would return to pre-acclimation levels according to the phenotypic flexibility hypothesis (Fig. 1C). We also measured body mass and wing length development, because in line with Bergmann's rule, birds are known to grow larger when it is colder (Ashton, 2002) and smaller when it is warmer (Van Buskirk et al., 2010; Teplitsky et al., 2008). This is expected to decrease or increase dry heat loss rate due to a higher ratio between body surface area and volume. If a higher surface area to volume ratio is biologically meaningful to increase heat transfer rate in the warmth, we predicted that birds growing up under heatwave-like conditions would have lower body mass and longer wings in keeping with established biogeographic rules (Ryding et al., 2021). This study provides new insights into how the developmental period influences how birds cope with increasing temperatures in the short- and long term, by providing data on whether phenotypic effects are lasting, and under which circumstances such responses may be adaptive or maladaptive.

Methods

Species and husbandry

Japanese quail eggs (n=60) were bought from a commercial breeder (Sigvard Månsgård, Åstorp, Sweden) and were incubated in 37.9 ± 0.52°C at 50 % relative humidity using Brinsea OvaEasy 190 incubators (Brinsea, Weston-super-Mare, United Kingdom). The eggs were stored at room temperature (18-20°C) and were turned manually twice daily until put into the incubator (6 eggs per day). On day 15 of incubation, the eggs were moved from the incubation trays to a hatching tray contained inside the incubator. The eggs were checked for signs of hatching daily from day 17 onwards. The incubation period was 19 days (range: 17 to 21d). Forty-five of 60 eggs (75%) hatched.

After hatching, the chicks were left inside the incubator until completely dry or for a maximum of 12 hours. They were then transferred to open pens ($310 \times 120 \times 60$ cm) dressed with wood shavings. During the first 3 weeks after hatching, housing temperature was either 20° (henceforth "control treatment") or 30°C to simulate heatwave conditions (henceforth "warm treatment"). From then on, all chicks were

housed in a common garden at 20°C until the end of the experiment. These temperatures were near the upper and lower critical temperatures, respectively, in Japanese quail chicks (Ben-Hamo et al. 2010) and broadly similar to those used by others studying the thermal sensitivity of development in quail (Burness et al., 2013). Chicks were assigned randomly to the warm (n=25) or control (n=20) treatments. The quail had *ad libitum* access to pelleted feed, water, sandbathes and crushed seashells throughout the experiment. Mealworms and vegetables (kale, carrot, or lettuce) were each offered once weekly on alternate days. Until 3 weeks old, the birds were fed turkey starter (Kalkonfoder Start, Lantmännen, Stockholm, Sweden; 25.5% protein). After 3 weeks of age, the feed was switched to turkey grower (Kalkonfoder Tillväxt, Lantmännen, Stockholm, Sweden; 22.5% protein). Until 3 weeks old, all birds had access to a heat lamp providing a temperature of 37°C at floor level, but no heating source was provided from week 3 onwards. Photoperiod was 12:12 LD (lights on from 06:00 to 18:00 GMT+2) throughout the experiment.

Measurements of body size and body temperature

Morphometric measurements were taken starting on day 1 and then once weekly throughout the experiment. On day 1, birds were banded and weighed ($\pm$ 0.1 g). From day 7 onwards, wing length was also measured ($\pm$ 0.5 mm). When the birds were 13 to 16 days old, we implanted a temperature-sensitive PIT tag (LifeChip BioTherm, Destron Fearing, South St. Paul, MN, USA), 2.1×12 mm in size ($< 0.5\%$ of body weight), into the intraperitoneal cavity to measure T_b. Ventral skin surrounding the distal end of the sternum was sterilised (70% EtOH) and anaesthetized topically using 5% lidocaine (EMLA®, Aspen Pharma, Durban, South Africa). Thirty to 45 minutes later, the skin was sterilized again, and a sterile PIT tag was inserted immediately beneath the distal tip of the sternal keel using a sterile 12-gauge syringe. This positioned the PIT tag on top of the liver (Andreas Nord, pers. obs.). The incision was closed using UHU cyanoacrylate (Bolton Adhesives, Rotterdam, Nederland) and covered with antiseptic ointment (1% H_2O_2, LHP, Bioglan AB, Malmö, Sweden).

Measurements of metabolic heat production and evaporative water loss

Metabolic heat production (MHP) and evaporative water loss (EWL) were measured using flow-through respirometry in a climate test chamber (Weiss Umwelttechnik C180, Reiskirchen, Germany). Measurements were taken, firstly, at the end of the heatwave treatment when the birds were 3 weeks old and halfway through the somatic growth phase. Measurements were taken again at 8 weeks, when the birds had reached asymptotic size and started reproducing (i.e., when they were young adults). During the first experimental period, we used 3.3 L glass respirometry chambers, and for the second experimental period 8.0 L glass chambers. The chambers were ventilated with dry (drierite; Sigma-Aldrich, Stockholm, Sweden) atmospheric air at $1.873 - 2.638$ L/min (mean $\pm$ SE: 2.149 ± 0.013 L/min;

standard temperature and pressure, dry, STPD) during the 3 week measurements and 3.653 – 4.751 L/min (mean ± SE: 4.179 ± 0.013 L/min, STPD) during the 8 week measurements as measured with a FB8 mass flow meter (Sable Systems, Las Vegas, NV, USA). We sub-sampled 308 – 388 ml/min (mean ± SE: 333 ± 1 ml/min, STPD) from the main gas stream using a SS-4 sub-sampler (Sable Systems) for gas analyses. Oxygen was measured using a FC-10 analyser (Sable Systems) and water vapor was measured with a RH-300 water vapour meter (Sable Systems). Water vapour and carbon dioxide were removed from the airstream before measuring oxygen using drierite and ascarite (II) (Acros Organics, Geel, Belgium). The oxygen analyser was zero calibrated using 100% N_2 and span calibrated to 20.95% O_2 using dry (drierite) room air. The RH-300 was zero calibrated using dry room air, and span calibrated by exploiting the dilution of oxygen by water vapour in moist air. Thus, by recording barometric pressure (BNP, kPa) and O_2 content in dry (O_{2dry}) and wet (O_{2wet}) air, it is possible to calculate water vapour pressure (WVP, kPa):

$$WVP = BP \times \left(O_{2_{dry}} - O_{2_{wet}}\right) / O_{2_{dry}} \qquad \text{(eq. 1)}$$

Birds were placed on a metal grid platform over a reservoir of mineral oil to avoid that evaporation from faeces affected the measurements of EWL (Lasiewski et al., 1966). Air temperature in the chambers were measured using thermocouples (36-gauge type T, copper-constantan) connected to a thermocouple box (TC-2000, Sable Systems) at a height where the temperature reading was not affected by contact with the bird. T_b was measured at 1 Hz throughout the experiment by placing custom-made antennas connected to a RM310 multiplexer board (Biomark, Boise, ID, USA) below the respirometry chambers. Measurements were performed with the climate test chamber set at 10, 20, 30 and 40°C. The corresponding temperatures in the respirometer chambers were 12 ± 0.14, 21 ± 0.07, 31 ± 0.05 and 40 ± 0.07°C.

All experiments took place during daytime. Birds were placed in a climate chamber illuminated by a light bulb at least 30 minutes before the start of data collection for thermal acclimation to the first measurement temperature (which was always 10°C). During the first experimental period, a measurement cycle started with a 10-minute baseline followed by 10 minute of sequential data collection for each of four birds and ended with a 20-minute baseline. During the second experimental period (i.e., when birds where 8 weeks old), 2 or 3 birds were measured simultaneously. Measurements started with a 15-minute baseline followed by 10 minute of data collection for each bird and ended with a 20-minute baseline. Temperature in the climate chamber was increased acutely to the next target temperature at the start of the last baseline measurement in a cycle, the new target being reached within 10 – 20 min. A bird was removed from the experiment if it showed signs of distress that did not pass in 5 minutes, or if T_b rose > 44°C at the same time as it showed signs of distress.

Data analyses

Missing data

Of the 45 birds that hatched, 39 (control: 17; warm: 22) were subsequently used in the first experimental

period and 36 (control: 16; warm: 20) in the second experimental period (Table 1). Four birds died of

natural causes or were euthanized before any measurements took place, and an additional 3 died between

3 and 8 weeks of age. Furthermore, 1 bird was never considered for any measurements because it showed

abnormally slow growth rate and was subsequently subject to severe pecking by the other birds. One

bird was left out of the thermal and metabolic experiments because it could not be fitted into the

measurement schedule (see above) but was included in the analyses of morphological characteristics.

Five birds were removed from the experiment before measurements in 40°C at 3 weeks of age due to

signs of stress (see above). Data from 8 birds from 6 sessions (of 100 total) were excluded from the data

set because gas concentrations were not at steady state, typically because the birds were non-resting and

so did not conform to the criteria for measuring resting metabolic rate (IUPS Thermal Commission,

2003). An overview of sample sizes per ages and response variable is provided in Table 1.

Calculations

Oxygen consumption (ml / min) was calculated from the most stable 2 min period of the 10 min

recording using eq. 11.1 in Lighton (2008), and was converted to metabolic heat production (MHP, in

W) assuming 1 ml of O_2 = 20 J (Kleiber, 1961). Evaporative water loss (EWL, in mg / min) was

calculated from the same period using eq. 11.9 in Lighton (2008) and was converted to EHL (in W),

assuming 2.406 J is required to evaporate 1 ml of H_2O (Wallace and Hobbs, 2006). We then calculated

evaporative cooling efficiency as the ratio between EHL and MHP (Lasiewski et al., 1966). Body

temperature was calculated as the mean of the 2 min period for which MHP and EWL were defined.

Statistical analyses

Statistical analyses were made in R (version 4.0.3; http://www.R-project.org/) using linear mixed

models (lmer function in lme4; Bates et al., 2015). To test if the treatment affected size and growth, we

used body mass or wing length as dependent variables, treatment as a factor, and age and age^2 as co-

variates. The original model also included the interactions treatment × age and treatment × age^2. Bird

ID was used as a random factor to account for repeated measurements on the same individual. For the

thermal and metabolic models, MHP, EWL, EHL/MHP, and T_b were used as response variables. To test

the effects of measurement temperature on these metrics, we fitted separate models at 3 and 8 weeks of

age with treatment, temperature, and treatment × temperature as factors, body mass as a continuous

covariate and bird ID as random intercept (to account for repeated measurements). MHP, EWL and

EHL/MHP were log-transformed before the analyses to meet parametric assumptions.

P-values for both fixed and random factors were assessed using likelihood ratio tests. The interaction

was removed from the model when non-significant (i.e., *P* > 0.05), but all main effects were retained. If

interactions were significant, a post hoc test (using the pairs() function in the emmeans package; Lenth,

2016) was performed between treatments within temperatures. Model estimates where calculated using

the emmean() function in the emmeans package (Lenth, 2016), and were back-transformed from the log

scale when applicable. Figures show raw data.

Results

Body mass and wing length

Both body mass and wing length increased non-linearly with age, with asymptotic phases for both traits

reached between 5 and 6 weeks of age. Neither growth curve differed between the treatments. Nor was

there any difference in the body mass (Fig. 2A) or wing length (Fig. 2B) intercepts between warm- and

control birds (Table 2). Birds had reached 58% of their adult (i.e., 8 week) body mass, and 81 % of adult

wing length at 3 weeks of age (Table 2; Fig. 2).

Effects of air temperature on thermal and metabolic responses

There was no interaction between treatment and temperature on log(MHP) at any age (Table 3). At 3

weeks, control birds had higher log(MHP) (2.09 ± 0.058 W) than warm birds (1.94 ± 0.051 W) (Table

3; Fig. 3A), but there was no effect of treatment on log(MHP) at 8 weeks (Table 3; Fig. 3A). Log(MHP)

increased with increasing body mass both at 3 and 8 weeks (Table 3).

Log(EWL) was not affected by the interaction between treatment and temperature, nor by treatment at

any age (Table 3). At 3 and 8 weeks, log(EWL) increased with temperature (Table 3; Fig. 3B).

Log(EWL) was not affected by body mass at 3 weeks (Table 3), but it increased with body mass at 8

weeks (Table 3).

The log-transformed ratio between EHL and MHP (i.e., log(EHL/MHP)), which measures how large a

proportion of MHP that the birds lost by evaporation, was affected by the interaction between treatment

and temperature at 3 weeks. Specifically warm birds lost a larger proportion of MHP (0.618 ± 0.022)

287 than control birds (0.523 ± 0.027) in 40°C (Table 3; Fig. 3C), but there was no difference between the

288 treatments in any other temperature (Table 3). At 8 weeks, log(EHL/MHP) was not influenced by

289 treatment, neither as a main effect or when interacting with temperature (Table 3), but it increased

290 significantly between all experimental temperatures (Table 3; Fig. 3C). At 3 weeks, log(EHL/MHP)

291 decreased with increasing body mass (Table 3) but this effect had disappeared by week 8 (Table 3).

293 T_b did not differ between treatments at any age, neither when considering the main effect nor its

294 interaction with air temperature (Table 3). However, T_b increased with temperature across treatments

295 for both ages (Table 3; Fig. 4). There was no effect of body mass on T_b at 3 weeks (Table 3). At 8 weeks,

296 T_b increased with increasing body mass (Table 3).

Discussion

299 By raising Japanese quail in heatwave-like conditions, we found that birds growing up in warm post-

300 hatch environments were more equipped to counter high air temperature as juveniles than control birds.

301 In line with our predictions, warm-reared birds had lower metabolic rate than control birds, and so could

302 dissipate more of their metabolic heat production (MHP) by evaporation compared to control birds when

303 exposed to hot air (i.e., the ratio between evaporative heat loss (EHL) and MHP was higher). Thus, birds

304 growing up warm are more likely to handle heat better, because evaporative cooling capacity is crucial

305 for heat tolerance (McKechnie and Wolf, 2019; McKechnie et al., 2016). Our results contrast a recent

306 study on zebra finches (*Taeniopygia guttata*) that found no effects of temperature on MHP or

307 evaporative water loss (EWL) close to fledging (Ton et al., 2021). However, the zebra finches studied

308 by Ton and colleagues had lower T_b than control birds both at fledging and as juveniles more than a

309 month after the birds had fledged and the experimental heatwave had ended. (Ton et al., 2021). In

310 contrast to this, we found no effects on T_b at any age, and there was also no difference in MHP or

311 evaporative cooling capacity at 8 weeks, when the birds had been kept in common garden for more than

312 one month. This indicates that evaporative cooling capacity develops faster when birds grow up in

313 warmer conditions, but that the temperature stimulus in this study did not cause permanent effects on

314 thermal and hygric physiology.

316 Even though a smaller body size could be expected according to biogeographical rules, we found no

317 effect of developmental temperature on body mass and wing length in our study. It is possible that

318 temperature-dependent constraints on development are not much of an issue when there is unrestricted

319 access to food and water. This is different from the effects of high temperature in the wild, where

320 heatwave-conditions are often associated with reduced food availability and drought (Kingsolver and

Woods, 1997; Smith, 1982; reviewed by Cunningham et al., 2021). Moreover, while many studies have shown that high temperatures cause decreases in body size (e.g. Salewski et al., 2010, Andrew et al., 2017, Andreasson et al., 2018), there are several exceptions to this pattern (Salewski et al., 2010; Siepielski et al., 2019; Ton et al., 2021). Thus, selection for small body size due to warmer temperatures might not be a general response (Siepielski et al., 2019). Nonetheless, Burness et al. (2013) and Andrew et al. (2017) report on reduced size in heat-challenged captive birds with *ad libitum* access to food and water. Moreover, broilers raised for meat production gain less body mass and eat less food, and have higher mortality, when exposed to both acute (Zaboli et al., 2017; Arjona et al., 1988) and chronic (Imik et al., 2012; Sohail et al., 2012; Ranjan et al., 2019) heat stress. Perhaps the difference between the control and warm treatment in our study was too small, or the warm temperature too low, to cause any irreversible changes on body size.

Alternatively, it is possible that other selection pressures for large body size, such as competitive ability in males or egg size in females, were stronger than selection for any size-related increase in dry heat flux in the warmth; demands for which would be more easily accommodated by physiological flexibility within a generation. Future studies should address if developmental temperature affected size of the appendages. According to Allen's rule (Allen, 1877), appendages such as the legs and bills of birds or the tails and ears of mammals, should be longer in warmer temperatures (cf. Demicka & Caputa, 1993). Since bird bills and legs are both richly vascularised and uninsulated, changes in local circulation can quickly either increase or decrease heat flux (Scholander et al., 1950; Tattersall et al., 2009). Thus, long appendages are likely aiding thermoregulation by the whole bird much more than any change in size per se.

The environmental matching hypothesis predicts that animals growing up in warmer environments should be better at handling warm temperatures as adults, which is shown by several studies (e.g., Demicka and Caputa, 1993; Stawski and Geiser, 2020; reviewed by Nord and Giroud, 2020). In other studies, silver spoon effects appear to be more common, reflected by reduced thermoregulatory performance or constrained growth after development in thermally challenging conditions (reviewed by Andreasson et al., 2020). We found support for neither hypothesis. However, it is important to note that we did not measure maximum heat and cold tolerance, which might be more telling of thermal acclimation than submaximal physiological rates. Such measurements could also be more informative when extrapolating laboratory results to free-ranging animals because, in the wild, temperatures vary by several tens of degrees between different microhabitats, e.g., in relation to vegetation and topography (Carroll et al., 2016; Cunningham et al., 2015; Cunningham et al., 2021), suggesting knowledge of heat

tolerances will inform on which animals are better able to exploit resources spread across the thermal
landscape.

Conclusion

We show that warm temperatures early in life can make chicks more prepared to counter heat stress.
However, we found no support for environmental matching of juvenile and adult environments, because
reproductively mature warm-origin birds that had spent the latter half of their growth and maturation
period in a common garden thermoregulated in the same way as control birds. Thus, a mild or short-
duration heatwave during the developmental period does not seem to be constraining for growth and
does not seem to have any obvious consequences for how animals respond to cold or warm temperatures
as adults. Future studies should address if a longer heatwave period, or a more severe heat treatment,
cause longer-term or irreversible effects. Such studies should also address if there are any differences
between appendages of birds subjected to different temperatures during development.

Funding

Ethics

Ethical approval for the experiment was granted by the Malmö/Lund Animal Ethics Committee (permit
no. 9246-19).

References

Albright, T. P., Mutiibwa, D., Gerson, A. R., Smith, E. K., Talbot, W. A., O'Neill, J. J., McKechnie, A.
E. and Wolf, B. O. (2017). Mapping evaporative water loss in desert passerines reveals an expanding
threat of lethal dehydration. *Proceedings of the National Academy of Science of the United States of*
America, 114, 2283-2288. doi: 10.1073/pnas.1613625114.

Allen, J. A. (1877). The influence of physical conditions in the genesis of species. *Radical Review*, i,
108–140.

Andreasson, F., Nilsson, J-Å. and Nord, A. (2020). Avian reproduction in a warming world. *Frontiers*
in Ecology and Evolution, 8, Article 576331. doi: 10.3389/fevo.2020.576331.

Andreasson, F., Nord, A. and Nilsson, J-Å. (2018). Experimentally increased nest temperature affects

body temperature, growth and apparent survival in blue tit nestlings. *Journal of Avian Biology*, 49,

Article e01620. doi: 10.1111/jav.01620.

Andrew, S. C., Hurley, L. L., Mariette, M. M. and Griffith, S. C. (2017). Higher temperatures during

development reduce body size in the zebra finch in the laboratory and in the wild. *Journal of*

Evolutionary Biology, 30, 2156-2164. doi: 10.1111/jeb.13181.

Arjona, A. A., Denbow, D. M. and Weaver, W. D. (1988). Effect of heat stress early in life on mortality

of broilers exposed to high environmental temperatures just prior to marketing. *Poultry science*, 67, 226-

231. doi: 10.3382/ps.0670226.

Ashton, K. G. (2002). Patterns of within-species body size variation of birds: strong evidence for

Bergmann's rule. *Global ecology and Biogeography*, 11, 505-523. doi: 10.1046/j.1466-

822X.2002.00313.x.

Bates, D., Mächler, M., Bolker, B. and Walker, S. (2015). Fitting linear mixed-effects models using

lme4. *Journal of Statistical Software*, 67, 1-48. doi: 10.18637/jss.v067.i01.

Ben-Hamo, Pinshow, B., McCue, M. D., McWilliams, S. R. and Bauchinger, U. (2010). Fasting triggers

hypothermia, and ambient temperature modulates its depth in Japanese quail Coturnix japonica.

Comparative Biochemistry and Physiology A, 156, 84-91. doi: 10.1016/j.cbpa.2009.12.020.

Both, C., van Asch, M., Bijlsma, R. G., van den Burg, A. B., & Visser, M. E. (2009). Climate change

and unequal phenological changes across four trophic levels: constraints or adaptations? *Journal of*

Animal Ecology, *78*, 73-83. doi: 10.1111/j.1365-2656.2008.01458.x.

Burness, G., Huard, J. R. Malcom, E. and Tattersall, G. J. (2013). Post-hatch heat warms adult beaks:

irreversible physiological plasticity in Japanese quail. *Proceedings of the Royal Society B*, 280, Article

20131436. doi: 10.1098/rspb.2013.1436.

Carroll, J. M., Davis, C. A., Fuhlendorf, S. D. and Elmore, R. D. (2016). Landscape pattern is critical

for the moderations of thermal extremes. *Ecosphere*, 7, Article e01403. doi: 10.1002/ecs2.1403.

Chen, I-C., Hill, J. K., Ohlemüller, R., Roy, D. B. and Thomas, C. D. (2011). Rapid range shifts of

species associated with high levels of climate warming. *Science*, 333, 1024-1026. doi:

10.1126/science.1206432.

Conradie, S. R., Woodborne, S. M., Cunningham, S. J., & McKechnie, A. E. (2019). Chronic, sublethal

effects of high temperatures will cause severe declines in southern African arid-zone birds during the

21st century. *Proceedings of the National Academy of Sciences of the United States of America*, 116,

14065-14070. doi: 10.1073/pnas.1821312116.

Cunningham, S. J., Gardner, J., L. and Martin, R. O. (2021). Opportunity costs and the response of birds and mammals to climate warming. *Frontiers in Ecology and the Environment*, 19, 300-307. doi: 10.1002/fee.2324.

Cunningham, S. J., Martin, R. O. and Hockey, P. A. R. (2015). Can behaviour buffer the impacts of climate change on an arid-zone bird?. *Ostrich*, 86, 119-126. doi: 10.2989/00306525.2015.1016469.

Cunningham, S. J., Martin, R. O., Hojem, C. L. and Hockey, P. A. R. (2013). Temperatures in excess of critical thresholds threaten nestling growth and survival in a rapidly-warming arid savanna: A study of common fiscals. *PLoS One*, 8, Article e74613. doi: 10.1371/journal.pone.0074613.

Dawson, R. D., Lawrie, C. C. and O'Brien, E. L. (2005). The importance of microclimate variation in determining size, growth and survival of avian offspring: experimental evidence from a cavity nesting passerine. *Oecologia*, 144, 499-507. doi: 10.1007/s00442-005-0075-7.

Deaton, J. W., Reece, F. N. and McNaughton, J. L. (1978). Effect of temperature during growth period on broiler performance. *Poultry Science*, 54, 1070-1074. doi: 10.3382/ps.0571070.

Demicka, A. and Caputa, M. (1993). Effect of warm rearing on the development of thermolytic effectors in rats. *Journal of Thermal Biology*, 18, 251-256. doi: 10.1016/0306-4565(93)90010-Q.

du Plessis, K. L., Martin, R. O., Hockey, P. A. R., Cunningham, S. J and Ridley, A. R. (2012). The costs of keeping cool in a warming world: implications of high temperatures for foraging, thermoregulation and body condition of an arid-zone bird. *Global Change Biology*, 18, 3063-3070. doi: 10.1111/j.1365-2486.2012.02778.x.

Farag, M. R. and Alagawany, M. (2018). Physiological alterations of poultry to the high environmental temperature. *Journal of Thermal Biology*, 76, 101-106. doi: 10.1016/j.jtherbio.2018.07.012.

Hansson, L. A., Ekvall, M. K., Ekvall, M. T., Ahlgren, J., Holm, W. S., Dessborn, L., & Bronmark, C. (2014). Experimental evidence for a mismatch between insect emergence and waterfowl hatching under increased spring temperatures. *Ecosphere*, 5(9), Article 120. doi: 10.1890/es14-00133.1.

Imik, H., Ozlu, H., Gumus, R., Atasever, M. A., Urgar, S. and Atasever, M. (2012). Effects of ascorbic acid and alpha-lipoic acid on performance and meat quality of broilers subjected to heat stress. *British Poultry Science*, 53, 800-808. doi: 10.1080/00071668.2012.740615.

IUPS Thermal Commission. (2003). Glossary of terms for thermal physiology. *Journal of Thermal Biology*, 28, 75-106. doi: 10.1016/S0306-4565(02)00055-4.

Jørgensen, E. and Blix, A. S. (1988). Energy-conservation by restricted body cooling in cold-exposed willow ptarmigan chicks. *Ornis Scandinavica,* 19, 17-20. doi: 10.2307/3676521.

Kingsolver, J. G. and Woods, H. A. (1997). Thermal sensitivity of growth and feeding in *Manduca sexta*

caterpillars. *Physiological Zoology,* 70, 631-638. doi: 10.1086/515872.

Kleiber, M. (1961). *The fire of life - An introduction to animal energetics*. New York. John Wiley &

Sons.

Lasiewski, R. C., Acosta, A. L. and Bernstein, M. H. (1966). Evaporative water loss in birds—I.

Characteristics of the open flow method of determination, and their relation to estimates of

thermoregulatory ability. *Comparative Biochemistry and Physiology*, 19, 445-457. doi: 10.1016/0010-

406X(66)90153-8.

Lenth, R. V. (2016). Least-squares means: the R package ismeans. *Journal of Statistical Software*, 69,

1-33. doi: 10.18637/jss.v069.i01.

Lighton, J. R. B. (2008). *Measuring metabolic rates: a manual for scientists*. New York. Oxford

University Press.

McCarty, J. P. and Winkler, D. W. (1999). Relative importance of environmental variables in

determining the growth of nestling tree swallows *Tachycineta bicolor*. *IBIS*, 141, 286-296. doi:

10.1111/j.1474-919X.1999.tb07551.x.

McKechnie, A. E., Hockey, P. A. R., and Wolf, B. O. (2012). Feeling the heat: Australian landbirds and

climate change. *Emu*, 112, 1–7. doi: 10.1071/MUv112n2_ED.

McKechnie, A. E., Whitfeld, M. C., Smit, B., Gerson, A. R., Smith, E. K., Talbot, W. A., McWhorter,

T. J. and Wolf, B. O. (2016). Avian thermoregulation in the heat: efficient evaporative cooling allows

for extreme heat tolerance in four southern hemisphere columbids. *Journal of Experimental Biology*,

219, 2145-2155. doi: 10.1242/jeb.138776.

McKechnie, A. E. and Wolf, B. O. (2019). The physiology of heat tolerance in small endotherms.

Physiology, 34, 302-313. doi: 10.1152/physiol.00011.2019.

Monaghan, P. (2008). Early growth conditions, phenotypic development and environmental change.

Philosophical Transactions of the Royal Society B, 363, 1635–1645. doi: 10.1098/rstb.2007.0011.

Nichelmann, M. (2004). Perinatal epigenetic temperature adaptation an avian species: comparison of

turkey   and   Muscovy   duck.   *Journal   of   thermal   biology*,   29,   613-619.   doi:

10.1016/j.jtherbio.2004.08.032.

Nilsson, J-Å. and Nord, A. (2018). Testing the heat dissipation limit theory in a breeding passerine.

Proceedings of the Royal Society B, 285, Article 20180652. doi: 10.1098/rspb.2018.0652.

Nord, A. and Giroud, S. (2020). Lifelong effects of thermal challenges during development in birds and

mammals. *Frontiers in Physiology*, 11, Article 419. doi: 10.3389/fphys.2020.00419.

Nord, A. and Nilsson, J-Å. (2019). Heat dissipation rate constrains reproductive investment in a wild

bird. *Functional Ecology*, 33, 250-259. doi: 10.1111/1365-2435.13243.

Nord, A. and Nilsson, J-Å. (2021). Low incubation temperature slows the development of cold tolerance

in a precocial bird. *The Journal of Experimental Biology,* 224, Article jeb237743. doi:

10.1242/jeb.237743.

Page, J. L., Nord, A., Dominoni, D. M, and McCafferty, D. J. (2022). Experimental warming during

incubation improves cold tolerance of blue tit (Cyanistes caeruleus) chicks. *Journal of Experimental*

Biology, 225, Article jeb243933. doi: 10.1242/jeb.243933.

Piestun, Y., Halevy, O., Shinder, D., Ruzal, M., Druyan, S., & Yahav, S. (2011). Thermal manipulations

during broiler embryogenesis improves post-hatch performance under hot conditions. *Journal of*

Thermal Biology, 36, 469-474. doi: 10.1016/j.jtherbio.2011.08.003.

Price, E. R., & Dzialowski, E. M. (2018). Development of endothermy in birds: patterns and

mechanisms. *Journal of Comparative Physiology B-Biochemical Systems and Environmental*

Physiology, 188, 373-391. doi: 10.1007/s00360-017-1135-0.

Ranjan, A., Sinha, R., Devi, I., Rahim, A. and Tiwari, S. (2019). Effect of heat stress on poultry

production and their managemental approaches. *International Journal of Current Microbiology and*

Applied Sciences, 8, 1548-1555. doi: 10.20546/ijcmas.2019.802.181.

Rodriguez, S. and Barba, E. (2016). Nestling growth is impaired by heat stress: an experimental study

in a Mediterranean great tit population. *Zoological Studies*, 55, Article 40. doi: 10.6620/ZS.2016.55-40.

Ryding, S., Klaassen, M., Tattersall, G. J., Gardner, J. L., & Symonds, M. R. E. (2021). Shape-shifting:

changing animal morphologies as a response to climatic warming. *Trends in Ecology & Evolution*, 36,

1036-1048. doi: 10.1016/j.tree.2021.10.012.

Salewski, V., Hochachka, W. M. and Fiedler, W. (2010). Global warming and Bergmann's rule: do

central European passerines adjust their body size to rising temperatures?. *Oecologia*, 162, 247-260.

doi: 10.1007/s00442-009-1446-2.

Scholander, P. F, Walters, V., Hock, R. and Irving, L. (1950). Body insulation of some arctic and tropical

mammals and birds. *Biological Bulletin*, 99, 225-236. doi: 10.2307/1538740.

Sharpe, L., Cale, B. and Gardner, J. L. (2019). Weighing the cost: the impact of serial heatwaves on

bodymass in a small Australian passerine. *Journal of Avian Biology*, 50, Article e02355. doi:

10.1111/jav.02355.

Shipley, J. R., Twining, C. W., Taff, C. C., Vitousek, M. N., Flack, A., and Winkler, D. W. (2020). Birds

advancing lay dates with warming springs face greater risk of chick mortality. *Proceedings of the*

National Academy of Sciences of the United States of America, *117*, 25590-25594. doi: 10.1073/pnas.2009864117.

Siepielski, A. M., Morrissey, M. B., Carlson, S. M., Francis, C. D., Kingsolver, J. G., Whitney, K. D. and Kruuk, L. E. B. (2019). No evidence that warmer temperatures are associated with selection for smaller body sizes. *Proceedings of Royal Society B*, 286, Article 20191332. doi: 10.1098/rspb.2019.1332.

Smith, K. G. (1982). Drought-induced changes in avian community structure along a montane sere. *Ecology,* 63, 952-961. doi: 10.2307/1937235.

Sohail, M.U., Ijas, A., Yousaf, M.S., Ashraf, K., Zaneb, H., Aleem, M. and Rehman, H. (2012). Alleviation of cyclic heat stress in broilers by dietary supplementation of mannan-oligosaccharide and Lactobacillus-based Probiotics: Dynamics of cortisol, thyroid hormones, cholesterol, C-reactive protein, and humoral immunity. *Poultry Science*, 89, 1934-1938. doi: 10.3382/ps.2010-0075.

Stawski, C. and Geiser, F. (2020). Growing up in a changing climate: how temperature affects the development of morphological, behavioural and physiological traits of a marsupial mammal. *Frontiers in Physiology*, 11, Article, 49. doi: 10.3389/fphys.2020.00049.

Tattersall, G. J., Andrade, D. V. and Abe, A. S. (2009). Heat Exchange from the Toucan Bill Reveals a Controllable Vascular Thermal Radiator. *Science*, 325, 468-470. doi: 10.1126/science.1175553.

Teplitsky, C., Mills, J. A., Alho, J. S. Yrrall, J. W. and Merilä, J. (2008). Bergmann's rule and climate change revisited: Disentangling environmental and genetic responses in a wild bird population. *Proceedings of the National Academy of Sciences of the United States of America*, 105, 13492-13496. doi: 10.1073/pnas.0800999105.

Thomas, C. D., Franco, A. M. A. and Hill, J. K. (2006). Range retractions and extinction in the face of climate warming. *Trends in Ecology and Evolution*, 21, 415-416. doi: 10.1016/j.tree.2006.05.012.

Ton, R., Stier, A., Cooper, C. E. and Griffith, S. C. (2021). Effects of heat waves during post-natal development on mitochondrial and whole body physiology: an experimental study in Zebra finches. *Frontiers in Physiology*, 12, Article 661670. doi: 10.3389/fphys.2021.661670.

Van Buskirk, J., Mulvihill, R. S. and Leberman, R. C. (2010). Declining body sizes in North American birds associated with climate change. *Oikos* 199, 1047-1055. doi: 10.1111/j.1600-0706.2009.18349.x.

van de Ven, T. M. F. N., McKechnie, A. E., Er, S. and Cunningham, S. J. (2020). High temperatures are associated with substantial reductions in breeding success and offspring quality in an arid-zone bird. *Oecologia*, 193, 225–235. doi: 10.1007/s00442-020-04644-6.

Visser, M. E., van Noordwijk, A. J., Tinbergen, J. M., & Lessells, C. M. (1998). Warmer springs lead

to mistimed reproduction in great tits (Parus major). *Proceedings of the Royal Society B-Biological*

Sciences, *265*, 1867-1870. doi: 10.1098/rspb.1998.0514.

Wallace, J. M. and Hobbs, P. V. (2006). *Atmospheric science*. Cambridge, Academic Press Inc.

Yahav, S. (2015). Chapter 37 - Regulation of Body Temperature: Strategies and Mechanisms. In Scanes,

C. G. (ed.) *Sturkie's Avian Physiology (Sixth Edition)*. London. Academic Press.

Zaboli, G-R., Rahimi, S., Shariatmadari, F., Torshizi, M. A. K., Baghbanzadeh, A. and Mehri, M.

(2017). Thermal manipulation during pre and post-hatch on thermotolerance of male broiler chickens

exposed to chronic heat stress. *Poultry science*, 96, 478-485. doi: 10.3382/ps/pew344.

TABLES

Table 1. The number of Japanese quail used when studying the thermo-physiological responses

(metabolic heat production (MHP), evaporative water loss (EWL), body temperature (T_b)) to

increasing air temperature at two different ages. The birds were raised in 20°C (control) or 30°C

(warm) for the first 3 weeks of life, and in 20°C thereafter. The same birds were measured at both

ages and at all temperatures, but some experiments were excluded from the final data set for

reasons stated in the main text.

		Sample size			
	Total	*10°C*	*20°C*	*30°C*	*40°C*
<u>3 weeks</u>	39				
Control		17	17	17	15
Warm		22	22	22	19
<u>8 weeks</u>	36				
Control		16	15	16	15
Warm		18	19	19	18

562   **Table 2.** Estimates, test statistics, degrees of freedom, P-values and standard deviations (σ) for the

563   random factor for models explaining the effects of developing in either control (20°C) or warm (30°C)

564   temperature conditions on body mass and wing length in Japanese quail.

Model	Estimate ± SE	LR	Df	P	σ_{ID} | σ_{total}
<u>Body mass (g)</u>					
Treatment		0.19	1	0.6601	
Age	35.07 ± 0.720	476.71	1	<0.0001 ***	
Age^2	-2.75 ± 0.078	225.62	1	<0.0001***	
Treatment:Age		0.30	1	0.5870	
Treatment:Age^2		0.39	1	0.5352	
Random factor bird ID		67.01	1	<0.0001 ***	11.44|25.69
<u>Wing length (mm)</u>					
Treatment		2.07	1	0.1500	
Age	58.40 ± 1.497	560.34	1	<0.0001 ***	
Age^2	-2.83 ± 0.145	429.19	1	<0.0001 ***	
Treatment:Age		0.01	1	0.9417	
Treatment:Age^2		0.00	1	0.9776	
Random factor bird ID		0.08	1	0.7844	0.60|5.99

Table 3. Estimates, test statistics, degrees of freedom, P-values and standard deviations (σ) for the

random factor for models explaining the effects of developing in either control (20°C) or warm (30°C)

temperature conditions on treatment at 3- and 8 weeks on metabolic heat production, evaporative

water loss, evaporative heat loss/metabolic heat production and body temperature in Japanese

quail. All dependent variables were log-transformed in the models, but back-transformed model

estimates are shown in the table. Statistics for main effects is not provided when the interaction

between treatment and temperature was significant. Different letters within brackets represent

significant (P < 0.05) post hoc comparisons. Abbreviations: MHP: Metabolic heat production; EWL:

Evaporative water loss; EHL/MHP: Evaporative heat loss/Metabolic heat production; T_b: Body

temperature.

Model	Estimate ± SE	LR	Df	P	σ_{ID}|σ_{total}
Log(MHP (W)) – 3 weeks					
Treatment		3.88	1	0.0460 *	
Control [A]	2.03 ± 0.060				
Warm [B]	1.88 ± 0.049				
Temperature		187.06	3	<0.0001 ***	
10°C [A]	2.51 ± 0.061				
20°C [B]	2.01 ± 0.049				
30°C [C]	1.51 ± 0.037				
40°C [B]	1.92 ± 0.049				
Treatment:Temperature		4.07	3	0.2536	
Body mass	1.004 ± 1.001	8.14	1	0.0043 **	
Random factor bird ID		45.15	1	<0.0001 ***	1.121.12|2.23
Log(MHP (W)) – 8 weeks					
Treatment		1.21	1	0.2711	
Temperature		75.27	3	<0.0001 ***	
10°C [A]	2.93 ± 0.097				
20°C [BD]	2.35 ± 0.078				
30°C [CD]	2.01 ± 0.066				
40°C [D]	2.19 ± 0.073				
Treatment:Temperature		2.02	3	0.5677	
Body mass	1.002 ± 1.001	5.42	1	0.0199 *	
Random factor bird ID		14.91	1	0.0001 ***	1.12|2.29
Log(EWL (g h^{-1})) – 3 weeks					
Treatment		0.00	1	0.9516	
Temperature		398.38	3	<0.0001 ***	
10°C [A]	0.500 ± 0.014				
20°C [B]	0.583 ± 0.016				
30°C [C]	0.710 ± 0.020				
40°C [D]	1.632 ± 0.047				
Treatment:Temperature		3.04	3	0.3856	
Body mass		0.68	1	0.4085	
Random factor bird ID		83.35	1	<0.0001 ***	1.16|2.25
Log(EWL (g h^{-1})) – 8 weeks					
Treatment		2.01	1	0.1559	
Temperature		321.04	3	<0.0001 ***	
10°C [A]	0.556 ± 0.019				
20°C [AB]	0.575 ± 0.020				
30°C [B]	0.628 ± 0.021				
40°C [D]	2.036 ± 0.070				
Treatment:Temperature		1.35	3	0.7173	
Body mass	1.002 ± 1.001	5.28	1	0.0216 *	
Random factor bird ID		25.56	1	<0.0001 ***	1.14|2.30
Log(EHL/MHP) – 3 weeks					

Treatment:Temperature		8.36	3	0.0391 *	
10°C					
Control [A]	0.134 ± 0.005				
Warm [A]	0.134 ± 0.004				
20°C					
Control [A]	0.187 ± 0.007				
Warm [A]	0.202 ± 0.007				
30°C					
Control [A]	0.299 ± 0.011				
Warm [A]	0.328 ± 0.011				
40°C					
Control [A]	0.518 ± 0.020				
Warm [B]	0.612 ± 0.021				
Body mass	0.998 ± 1.001	4.57	1	0.0325 *	
Random factor bird ID		17.77	1	<0.0001 ***	1.09|2.23
Log(EHL/MHP) – 8 weeks					
Treatment		0.14	1	0.7130	
Temperature		318.56	3	<0.0001 ***	
10°C [A]	0.127 ± 0.005				
20°C [B]	0.163 ± 0.006				
30°C [C]	0.209 ± 0.007				
40°C [D]	0.623 ± 0.022				
Treatment:Temperature		0.91	3	0.8238	
Body mass		0.02	1	0.8748	
Random factor bird ID		3.67	1	<0.0554	1.09|2.30
T_b (°C) – 3 weeks					
Treatment		0.61	1	0.4359	
Temperature		218.04	3	<0.0001 ***	
10°C [A]	42.68 ± 0.061				
20°C [B]	42.31 ± 0.061				
30°C [C]	42.00 ± 0.061				
40°C [D]	43.61 ± 0.064				
Treatment:Temperature		4.17	3	0.2491	
Body mass		0.01	1	0.9046	
Random factor bird ID		30.35	1	<0.0001 ***	1.23|2.61
T_b (°C) – 8 weeks					
Treatment		0.37	1	0.5409	
Temperature		134.80	3	<0.0001 ***	
10°C [A]	43.07 ± 0.077				
20°C [B]	42.66 ± 0.077				
30°C [B]	42.57 ± 0.076				
40°C [C]	43.69 ± 0.078				
Treatment:Temperature		2.97	3	0.3960	
Body mass	0.005 ± 0.003	4.73	1	0.0297 *	
Random factor bird ID		43.69	1	<0.0001 ***	1.40|2.75

Figure 1. Three hypothetical scenarios for thermo-physiological changes in response to a

developmental heatwave. A) Under the Environmental matching hypothesis, a developmental

heatwave leads to a non-reversible reduction in metabolic heat production and improvement in

evaporative cooling capacity, such that individuals developing under heatwave-like conditions are

more heat tolerant as adults. B) Under the Silver spoon hypothesis, a developmental heatwave

impairs normal growth and maturation of the thermoregulatory system which lowers heat tolerance

both in juveniles and adults. C) Under the Phenotypic flexibility hypothesis, a developmental

heatwave triggers reduced metabolic heat production and improved evaporative cooling capacity.

However, these effects are reversible so that the developmental thermal environment does not

affect heat tolerance in adulthood.

Figure 2. Effects on body mass and wing length when Japanese quail developed in either control

(20°C) or warm (30°C) temperature conditions. The figure shows mean ± SE (A) body mass and (B)

wing length of Japanese quail during the 8-week experimental period. The grey curve is LOESS (locally

estimated scatterplot smoothing) ± 95% confidence interval. The dashed lines represent the timing of

measurements of metabolic rate and evaporative water loss. Sample sizes are stated in the Methods.

Transparent points show raw data.

Figure 3. Effects on metabolic heat production (MHP), evaporative water loss (EWL) and the ratio

between evaporative heat loss (EHL) and MHP when Japanese quail developed in either control

(20°C) or warm (30°C) temperature conditions. The figure shows mean ± SE (A) MHP, (B) EWL and

(C) EHL/MHP of Japanese quail in the different test temperatures (10°C, 20°C, 30°C and 40°C) at 3

weeks and 8 weeks of age. Sample sizes are stated in the Methods. Transparent points show raw

data.

Figure 4. Effects on body temperature (T_b) when Japanese quail developed in either control (20°C) or warm (30°C) temperature conditions. The figure shows mean ± SE T_b of Japanese quail in the different test temperatures (10°C, 20°C, 30°C and 40°C) at 3 weeks and 8 weeks of age. Sample sizes are stated in the Methods. Transparent points show raw data.