Mutualism at the leading edge: Insights into the eco-evolutionary dynamics of host-symbiont communities during range expansion

Maria M Martignoni[1*], Rebecca C Tyson[2†], Oren Kolodny[1†], Jimmy Garnier[3†]

[1*]Department of Ecology, Evolution and Behavior, A. Silberman Institute of Life Sciences, Faculty of Sciences, Hebrew University of Jerusalem, Israel.
[2]Department of Mathematics, University of British Columbia, British Columbia, Canada.
[3]Laboratory of Mathematics, CNRS, Université Savoie-Mont Blanc, Université Grenoble Alpes, Chambery, France.

*Corresponding author(s). E-mail(s):
maria.martignonimseya@mail.huji.ac.il;
Contributing authors: rebecca.tyson@ubc.ca; orenkolodny@gmail.com;
jimmy.garnier@univ-smb.fr;
[†]These authors contributed equally to this work.

Abstract

The evolution of mutualism between host and symbiont communities plays an essential role in maintaining ecosystem function and should therefore have a profound effect on their dynamics of range expansion. In particular, the presence of mutualistic symbionts at the leading edge of a host-symbiont community should enhance its propagation in space. We develop a theoretical framework that captures the eco-evolutionary dynamics of host-symbiont communities, to investigate how the evolution of resource exchange may shape community structure during range expansion. We consider a community which includes symbionts that are mutualistic or parasitic to various degrees, where parasitic symbionts receive the same amount of resource from the host as mutualistic symbionts, but at lower cost. The selective advantage of parasitic over mutualistic symbionts is strengthened with resource availability (i.e. with host density), promoting mutualism at the range edges, where host density is low, and parasitism in the population core,

where host density is higher. This spatial selection also influences the speed of spread. We find that the host growth rate (which depends on the average benefit provided by the symbionts) is maximal at the range edges, where symbionts are more mutualistic, and that host-symbiont communities with high symbiont density at their core (e.g. resulting from more mutualistic hosts) spread faster into new territories. These results indicate that the expansion of host-symbiont communities is pulled by the hosts but pushed by the symbionts, in a unique push-pull dynamics of which both, host and symbionts, are active and tightly-linked players.

Keywords: eco-evolutionary dynamics, mathematical model, mutualism, range expansion, arbuscular mycorrhizal fungi, host-microbial communities

MSC Classification: 92D40: Ecology , 92D15: Problems related to evolution , 92D25: Population dynamics (general).

Acknowledgments. MMM acknowledges the Azrieli Foundation. MMM and OK acknowledge the Israel Science Foundation (ISF) (grant number 1860/20), the Gordon and Betty Moore Foundation, and the United States-Israel Binational Science Foundation (BSF). RCT acknowledges the Natural Sciences and Engineering Research Council (NSERC) of Canada Discovery Grants Program, grant number RGPIN-2022-03589, and the University of British Columbia Okanagan Institute for Biodiversity, Resilience, and Ecosystem Services. JG acknowledges ModEcoEvo project funded by the Université Savoie Mont-Blanc.

Statements and declarations

- The authors have no competing interests to declare that are relevant to the content of this article.
- All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this manuscript.
- The authors have no financial or proprietary interests in any material discussed in this article.
- All authors agree with the present manuscript
- For their simulations, authors used Matlab R2022. All the codes developed for the manuscript are available on GitHub https://github.com/garnieji/AMF_Evolution
- Authors' contributions: MMM: Conceptualization, Methodology, Investigation, Writing - Original Draft, Writing - Review & Editing, Visualization. RT: Conceptualization, Methodology, Investigation, Data curation, Writing - Review & Editing, Visualization, Funding acquisition. OK: Writing - Review & Editing, Funding acquisition. JG: Conceptualization, Methodology, Software, Formal analysis, Investigation, Data curation, Writing - Review & Editing, Visualization, Funding acquisition.

Introduction

Climate change and anthropogenic disturbance are inducing species to engage in range expansion toward more suitable habitat at an unprecedented rate (Chen et al, 2011). Understanding the mechanisms governing population spread is therefore a key priority in conservation biology. Nonetheless, predicting the specific contours of a species' range expansion remains challenging (Fournier et al, 2019). Spatial spread is determined by processes acting at the edge of a species' range, where population densities are lower in comparison to those in the core of the range (Chuang and Peterson, 2016). Low density populations experience unique interaction structures and selective pressures, such as decreased intraspecific competition, or altered reproductive success (Phillips, 2009; Huang and Peng, 2016)), which together favour the evolution of certain traits with respect to others and can affect population spread in an unexpected manner.

In this regard, the range expansion of a host population associated with a mutualistic symbiont community, represents a particularly challenging study system. The term 'host-associated community' refers to a community of small organisms with a short generation time (the 'symbionts') associated with a large host (Bronstein, 2015). Host-associated communities may encompass a collection of parasitic, commensal, and/or mutualistic organisms where, in order to be defined as 'mutualistic', the associated symbionts should have an overall positive effect on host fitness, at least to some extent (Bronstein, 2015). If the fitness of a host species is affected by the presence of its mutualistic symbionts, it follows that the spatial spread of the host population will be intrinsically coupled to the density and characteristics of its associated symbionts at the range edge. Additionally, lower population densities of the host at the range edges of an expanding population may lead to the the selection of symbionts with mutualistic traits that differ from those observed in the population core, further affecting the expansion dynamics (Koella, 2000; Van Dyken et al, 2013). Mutualism thus adds a further degree of complexity to the evaluation of the speed of spread of hosts and symbionts.

Host-associated mutualistic communities are widespread in nature, and many have well-established ecological and economic significance. Examples include the crucial role played by mycorrhizal fungi in plant growth and agricultural productivity (Smith and Read, 2010), or the fundamental function of zooxanthellae in supporting coral reef ecosystems and fisheries (Muller-Parker et al, 2015). Only recent studies, however, have investigated how mutualistic interactions can affect the range limits of a host population and its symbionts (Afkhami et al, 2014; Araújo and Rozenfeld, 2014; Stanton-Geddes and Anderson, 2011; Dickie et al, 2017; Godsoe et al, 2017; Fournier et al, 2019; Benning and Moeller, 2021; Paquette and Hargreaves, 2021; Stephan et al, 2021). In this context, much greater attention has been given to the influence of abiotic factors (Chen et al, 2011; Spence and Tingley, 2020), or to the negative impacts of biotic factors such as predation, competition, or pathogen transmission (Phillips et al, 2010; Brown and Vellend, 2014; Wan et al, 2016). Theoretical approaches have also been scarce (Case et al, 2005; Brooker et al, 2007; Mack, 2012; Kubisch et al, 2014; Shaw, 2022), with very few mathematical models allowing treatment of hosts and their multiple symbionts as separate entities capable of influencing each others' population dynamics and evolutionary trajectories.

Even less is known about how the dynamics of interactions between hosts and symbionts are shaped by spatial dispersal (Mony et al, 2022; Hu et al, 2022), and whether the community of symbionts in a core population should be expected to differ from what is observed at the edges of the expansion range (Doebeli and Knowlton, 1998). Empirical work on host-microbial symbioses has revealed changes in microbial diversity at the range edge (Lankau and Keymer, 2016; Fowler et al, 2023). This variation could lead to a decrease in host fitness, or could help the host to better adapt to new environmental challenges, depending on the microbial composition observed in the edge community. These studies, however, do not address the question of whether range expansion favours selection for specific mutualistic traits in the symbionts, affecting the structure of the symbiotic community. Currently, only a few studies of protective plant-ant mutualisms have tackled this issue, and have identified variation in the mutualistic investment of ants at the range edge (Léotard et al, 2009; Vittecoq et al, 2012).

To address these questions, we develop a theoretical framework considering the mutualistic interactions of a population of plants (the 'host') and a community of host-associated arbuscular mycorrhizal fungi (the 'symbionts'). We consider host and symbionts to interact by exchanging resources necessary for their growth, where access to the hosts resource is the same for all symbionts, regardless of the benefit they provide in return. We further consider symbionts to have the ability to disperse independently from the host, and to transmit horizontally within the host population.

Host control, intended as any mechanism by which hosts can actively select for more beneficial symbionts in the community, has been a popular explanation for the persistence of mutualism in the face of cheaters (Hoeksema and Kummel, 2003; Bever, 2015; Bachelot and Lee, 2018; Christian and Bever, 2018). Host species have been found to have some influence over the composition and persistence of their symbionts, e.g. through gene regulation (Wier et al, 2010; Davenport et al, 2015), through secretion of compounds such as amino acids, sugars, and organic acids (Yuan et al, 2015; Frenkel and Ribbeck, 2017), or through preferential allocation of resources (Bever et al, 2009; Kiers et al, 2011). The universality of host control mechanisms, however, and the extent to which these mechanisms apply in natural contexts, have both been questioned in numerous studies, particularly in the context of the mycorrhizal symbiosis (Cameron et al, 2008; Walder et al, 2012; Walder and Van Der Heijden, 2015; Zhang et al, 2015; van der Heijden and Walder, 2016).

In the absence of host control mechanisms, vertical transmission of symbionts has also been found to favour mutualism (Ewald, 1987; Sachs et al, 2004). Indeed the faithful transmission of symbionts from parent to offspring may promote the evolution of more mutualistic traits when the reproductive success of a host and its symbionts is linked (Sachs et al, 2011; Frederickson, 2013). Symbionts, however, are often horizontally or environmentally inherited (Wilkinson and Sherratt, 2001; Vandenkoornhuyse et al, 2015; Shade et al, 2017). In this case, a symbiont may remain parasitic rather than engage in a mutualistic interaction (Ferdy and Godelle, 2005; Drew et al, 2021a), and little is known about how mutualism can persist when the fates of the symbionts and the host are decoupled. We have therefore developed a model that allows us to

investigate mutualism persistence in the absence of host control and when symbionts are shared among hosts.

Evolutionary changes in symbionts have been proposed to play a key role in structuring microbial communities, and represent an important factor to consider in elucidating the interaction dynamics of a host and its symbionts (Gómez et al, 2016; Miller et al, 2018). We therefore assume that the mutualistic investment of each symbiont (quantified by the rate at which the symbiont provides a benefit to the host) can change in time through a diversification process, e.g. genetic or phenotypic mutation. We thus develop a theoretical framework in which the evolutionary dynamics of the symbionts and their ecological interactions with the host occur on the same timescale. Specifically, we ask the following questions:

(i) How do eco-evolutionary dynamics shape the structure of host-associated mutualistic communities?

(ii) How does symbiont investment,in the mutualistic relationship with the host, differ between the population core and the edge of the expansion range?

(iii) How does resource supply of a host population toward its symbionts affect the spread of both host and symbiont populations?

Our analysis provides insights into the eco-evolutionary dynamics and linked dynamics of expansion of a host and its symbionts. To make our work more concrete, we select the plant-mycorrhizal mutualism as our motivating example, but our model and results are applicable to multiple ecological systems (Palmer et al, 2003; Robinson et al, 2010).

Models and Methods

We present an eco-evolutionary model describing the dynamics of a host population of plants supporting a host-dependent community of arbuscular mycorrhizal fungi (AMF), where the associated AMF compete amongst each other for space (Thonar et al, 2014; Engelmoer et al, 2014). The plant and AMF densities correspond to the amount of biomass (leaves, roots, hyphae, etc) present at each time and location and are described by the variables $p(x,t)$ and $m(x,t,\alpha)$ respectively. The AMF community is composed of fungi that differ in their ability α to deliver resource to host plants, as explained in detail below.

Individuals within the plant population and AMF community are each characterised by a trait quantifying their mutualistic investment, i.e. the rate at which each delivers resource to the partner: Each plant delivers carbon to the AMF at fixed density-dependent rate β, while each associated AMF delivers phosphorus to the plant at density-dependent rate α. The mutualistic investment of the fungi is subject to evolution, e.g. α can evolve through mutation or recombination during spore production and spread, or during root colonization (Vandenkoornhuyse et al, 2001; den Bakker et al, 2010). Thus, a plant-dependent community of AMF includes fungi that differ in their mutualistic investment α. When the mutation rate is high, or mutations with large effects on traits are rare (as it is commonly observed in microbial communities (LeClerc et al, 1996; Oliver et al, 2000; Trindade et al, 2010)), diversification of the trait α can be modeled as a diffusion process with a diffusion or *diversification rate* d_m, corresponding to the product of the mutation rate and half the variance associated

with mutational effects (Fleming, 1979). In addition to trait evolution, we consider the plant population and AMF community to undergo spatial expansion through the random dispersal of seeds or spores, respectively, and we quantify their dispersal ability by the diffusion coefficients D_p (for the plant) and D_m (for the associated AMF).

We thus obtain the following model for a population of plants with biomass density $p(x,t)$ interacting with a community of AMF with total biomass density $M(t,x)$, composed of multiple AMF whose biomass density $m(t,x,\alpha)$ depends on their mutualistic investment α:

$$\partial_t p = D_p \partial_x^2 p + p \left(r_p + \frac{Q \overline{\alpha}(t,x) M(t,x)}{p+d} - \beta M(t,x) - \mu_p p \right), \ x \in \mathbb{R}, \ t > 0, \qquad (1a)$$

$$\partial_t m = D_m \partial_x^2 m + d_m \partial_\alpha^2 m + m \left(\beta p - \frac{\alpha p}{p+d} - \mu_m M(t,x) \right), \ \alpha \in (\alpha_{\min}, \alpha_{\max}). \ (1b)$$

Note that p, m, and M have been converted into dimensionless quantities with arbitrary units. The parameter $\alpha_{\min}$ is the minimal mutualistic investment of the AMF (which can be zero) and $\alpha_{\max} > 0$ is the maximal mutualistic investment. Each AMF may be considered parasitic (if the cost of the symbiont for the plant is higher than the benefit it provides) or mutualistic (if the benefit provided is higher than the cost), as we will explain in detail below (see Eq. (3)). The quantity $\overline{\alpha}(t,x)$ is the mean mutualistic investment of the AMF community at location x and time t, and is defined as

$$\overline{\alpha}(t,x) = \int_{\alpha_{\min}}^{\alpha_{\max}} \alpha \, \frac{m(t,x,\alpha)}{M(t,x)} \, d\alpha \ \ \text{and} \ \ M(t,x) = \int_{\alpha_{\min}}^{\alpha_{\max}} m(t,x,\alpha) \, d\alpha \,. \qquad (2)$$

To ensure that mutations have no effect on the total population biomass of the symbiont community, Eqs. (1) have no-flux boundary conditions on the boundary of the trait domain $(\alpha_{\min}, \alpha_{\max})$, that is,

$$\partial_\alpha m(t,x,\alpha_{\min}) = \partial_\alpha m(t,x,\alpha_{\max}) = 0, \ \ \text{for all } t > 0, x \in \mathbb{R} \,.$$

The delivery of resource from the plant to the AMF depends linearly on plant and AMF densities (see fourth and third terms in Eqs. (1a) and (1b) respectively). Resource delivery from the AMF to the plant increases linearly with increasing fungal density, and nearly linearly with increasing plant density when plant density is low (relative to parameter d) (see third and forth terms in Eqs. (1a) and (1b) respectively). When plant density is large (relative to parameter d) resource delivery tends toward a dependency on fungal density only, as we assume host availability to not be a factor affecting the resource delivery capacity of associated symbionts (see Martignoni et al (2020) for more details about the underlying biological assumptions underlying these functional responses for the benefit and cost of mutualism).

The nutrients (e.g., phosphorus) received by the plant from the AMF, and the carbon received by the AMF from the plant are converted into plant and AMF biomass respectively. The parameter Q represents the efficiency of these conversions (see Martignoni et al (2020)). Parameters μ_p and μ_m correspond to the density-dependent rates at which resources are directed to the maintenance of the existing plant and AMF biomass, respectively. These costs encompass, for example, energetic costs and the

cost of direct competition. Note that the maintenance term of the AMF community depends on the total AMF density $M(x,t)$ (see last term of Eq. (1b)), and therefore includes direct competition between symbionts.

The eco-evolutionary dynamics of host-associated communities:

We first consider the model of Eq. (1) in the absence of spatial spread (i.e., $D_m = D_p = 0$). We aim to understand how a symbiotic community can emerge from natural selection and persist in association with a population of host plants. For this purpose, we investigate the coupled dynamics of a community of AMF associated with a population of host plants, and we consider the role that evolutionary changes in the mutualistic investment of the symbionts (α) may play in the establishment and persistence of mutualistic communities. Symbionts with low mutualistic investment reduce the growth rate of the host. We will refer to these symbionts as 'parasitic', and we will define them as AMF characterized by $\alpha \leqslant \alpha_c$, where α_c is the threshold below which an obligate mutualistic plant can not survive in the presence of that single symbiont (see SI.A for mathematical details). It is defined by

$$\alpha_c = \frac{\beta\,d}{Q} \, \frac{(Q+1) - \sqrt{(Q-1)^2 - 4\dfrac{Q\mu_m\mu_p}{\beta^2}}}{2} \, . \tag{3}$$

Symbionts with large mutualistic investment ($\alpha \geqslant \alpha_c$) enhance the host growth rate, which allows the host-associated community to survive. We will refer to these symbionts as 'mutualistic'. We investigate the mechanisms leading to the coexistence of mutualistic and parasitic symbionts in a community supported by a host population. In addition, we study how the mutualistic investment (β) of host plants toward their associated AMF affects the distribution of mutualistic traits (α) among the AMF community and the proportion of parasitic and mutualistic symbionts in the community.

Mutualistic investment at the range edges:

We use Eqs. (1) to understand how dispersal structures the average mutualistic investment α along a propagating wave. We compare the proportion of parasitic and mutualistic symbionts and their average mutualistic investment $\overline{\alpha}$ at the edge and in the core of an expanding plant-AMF community. By computing the plant and AMF growth rates along the wave front, we also gain insights into whether the expansion of host-associated communities can be defined as a pushed wave (i.e. driven by a growth rate that is highest some distance behind the wave front) or pulled wave (i.e. driven by a growth rate that is highest at the leading edge).

The spread of a host population and its symbionts:

We aim to understand how mutualistic interactions between a plant and its associated AMF affect the speed of spread of a host and its associated symbionts in a homogeneous landscape. We first compute the speed of spread of a host-symbiont community for different values of the rate β at which the plant provides carbon to its AMF. We

<div align="center">(a)   (b)</div>

Fig. 1 Host-symbiont community dynamics in the absence of dispersal. Panel (a), Plant density at equilibrium (black) and total density of the AMF community (blue) as functions of the plant carbon supply rate β. The mean mutualistic trait $\overline{\alpha}$ of the AMF community is shown in red and the mutualism/parasitism threshold α_c is shown in grey. Black asterisks correspond to the simulation results in panel (b). Panel (b), Trait distribution in the AMF community as a function of AMF phosphorus supply rate α and for various values of plant carbon supply rate β. For both panels, symbols correspond to equilibria of the model (1) without spatial spread ($D_p = D_m = 0$) obtained from numerical simulations, while curves correspond to theoretical approximations defined in SI.A. Parameter values used for the simulations are: $d_m = 0.01$, $Q = 6$, $\mu_p = \mu_m = 0.3$, $\alpha_{\min} = 0$ and $\alpha_{\max} = 5$.

further investigate how the speed of spread is affected by the difference between the dispersal abilities of the plant and the AMF. More precisely, we vary the ratio between the dispersal ability of the plant population (D_p) and that of the AMF community (D_m). Finally, we test how results differ when the plant is an obligate ($r_p = 0$) versus a facultative mutualist ($r_p > 0$).

Results

The eco-evolutionary dynamics of host-associated communities

Our model predicts the establishment and persistence of a mutualistic host-associated community (Fig. 1(a)). The emerging community is composed of a combination of symbionts that are parasitic or mutualistic to various degrees, where the mutualistic investment of the symbionts does not converge to a specific value but is distributed within the community across the range $\alpha_{\min} - \alpha_{\max}$ (Fig. 1(b)).

Mutualistic symbionts receive the same benefit from the hosts as parasitic ones (proportional to $\beta\,p$) but at higher cost (proportional to $\alpha\,p/(p+d)$, where α is lower for parasitic symbionts) as described by Eq. (1b). Parasites thus benefit from higher fitness and reach higher densities than mutualists, which confers a competitive advantage to parasitic symbionts. As a result, the trait distribution of AMF has its maximum at $\alpha_{\min}$ and decreases with increasing α (Fig. 1(b) and SI.B for mathematical details). The selection strength favouring parasitic symbionts depends linearly on plant density p. However, host growth benefits from high mutualistic investment by the symbionts (see Eq. 1a), and so host density decreases with decreasing mean mutualistic investment $\overline{\alpha}$

<div align="center">8</div>

(see Box 1). Thus, in the absence of mutations, mutualistic symbionts are competitively excluded by parasitic symbionts, with a consequent reduction in plant growth and collapse of the whole community (see SI.A for more details). Mutations cause parasites to evolve into better mutualists, driving a diversification in α toward higher mutualistic investment. Even when diversification occurs at an extremely low rate (Fig. B1a in SI.B), this process can balance selection due to competition between symbionts (which selects for low values of α) to produce a mean mutualistic investment of the AMF community large enough to support plant growth (Fig.1(a)).

From the perspective of the symbionts, the intertwined effects of mutation and selection generate an average mutualistic investment $\overline{\alpha}$ that is decreasing with selection strength and thus decreasing with respect to plant density. More specifically, the resulting balance between selection and mutation produces a distribution of AMF density as a function of plant density p that can be approximated by the Airy function truncated at α_{max}, with mean $\overline{\alpha}$ satisfying

$$\overline{\alpha} = z_0 \left(\frac{(p+d)d_m}{p} \right)^{1/3}, \tag{4}$$

where z_0 is a positive constant which depends only on the Airy function that solves the dimensionless problem $\mathrm{Ai}''(z) - z\,\mathrm{Ai}(z) = 0$ on $\mathbb{R}$ (see SI.B.3 for more details and Fig. 1b). In addition, the mean mutualistic investment $\overline{\alpha}$ is increasing with the diversification rate d_m (Fig. B1).

From the perspective of the plants, the ecological interaction between the plants and the AMF community produces dynamics with antagonistic effects. Host plants benefit from the community at rate $Q\overline{\alpha}M/(p+d)$, which is increasing with the average mutualistic investment $\overline{\alpha}$. At ecological equilibrium, plant density satisfies

$$p = d\left(\frac{\overline{\alpha}}{\alpha_c} - 1 \right), \tag{5}$$

where α_c is the parasitic/mutualistic threshold defined by (3) (see SI. B).

The antagonistic outcomes of the evolutionary process described by (4), and the ecological interactions between the host plants and symbiont community described by (5), produce an eco-evolutionary equilibrium. The eco-evolutionary feedback between plant density and the mean mutualistic trait is the key to the persistence of a community where parasites and mutualists coexist (see Box 1). More precisely, the community at equilibrium persists despite strong competition between symbionts, thanks to parasitic symbionts evolving into more mutualistic symbionts through mutation (Fig. B3 and SI.B.5).

Box 1: Forces stabilizing the eco-evolutionary dynamics of host-symbiont communities

Diagram illustrating the evolutionary and ecological forces stabilizing the population dynamics of host-symbiont communities. Ecological interactions between hosts and symbionts tend to increase host density in a way that is proportional to the mean mutualistic investment $\overline{\alpha}$ of its symbionts (see Eq.(5)). Evolutionary forces, on the other hand, tend to favor parasitic symbionts, which have a greater fitness advantage over mutualistic symbionts. This advantage increases with increasing resource availability, and thus with increasing host density (see Eq.(4)). If the community is highly parasitic, $\overline{\alpha}$ assumes a lower value, which causes a reduction in host density and a decrease in the fitness advantage of parasitic symbionts, and the community becomes on average more mutualistic. If the community is more mutualistic, host density increases and so does the fitness advantage of parasitic symbionts. The balance between evolutionary forces (selecting for more parasitic traits) and ecological forces (selecting for more mutualistic traits) stabilizes host density around a fixed value.

Carbon supply to the symbionts (parameter β) represents the mutualistic investment of the host plants. It determines the growth of the associated AMF and, in turn, their own growth. Plant density at equilibrium is therefore affected by β (see Eq. (5)). Since the plant density affects community structure through the evolutionary dynamics of symbionts (see Eq. (4) and SI.B), the carbon supply rate β is also an indirect determinant of community structure. In particular, a low rate of carbon supply β from the host to the symbionts does not provide enough resource for the AMF community to grow and support host growth, while a large β leads to a high cost of mutualism for the plant population itself, and thus directly reduces its growth. Plant density is therefore maximised at an intermediate value of β (Fig. 1(a)).

Mutualistic investment at the range edges

In this section we focus on the specific contribution of parasitic and mutualistic symbionts in shaping the expansion dynamics of the community. We observe that coexistence of parasitic and mutualistic symbionts is dynamically stable, consistent with the non-spatial eco-evolutionary dynamics described above. In the core of an expanding population selection favors parasitic symbionts and allows them to grow larger than mutualistic symbionts (Fig. 2(a)). As plant density decreases toward the expansion edges, the strength of selection against mutualistic symbionts is reduced, and thus the mean mutualistic investment of the community increases (Fig. 2(a)). Thus the resulting effect on the distribution of α across the AMF community is similar to the one observed when varying β in the absence of dispersal, but with position along the wavefront, x, playing the role of β (cfr. Fig. 1(b) and Fig. 2(b)). Indeed, the positive dependence of the selection strength on plant density combined with the low plant density at the leading edge of the traveling wave explains the decrease in the proportion of parasitic symbionts toward the traveling front (Fig. 2(c) and SI.B.4).

It is evolution among the parasitic symbionts in the population core that gives rise to the mutualistic population at the range edge. Indeed, parasitic symbionts are the most likely common ancestors of mutualistic symbionts (Fig. B3 and Fig. C5). A mutualistic community at the leading edge provides a large growth rate to the host plants, leading to a rapid increase in host density (Fig. 2(d)). As plant density increases, the growth rate of the symbionts will also increase, given that the benefit provided by the host to the symbionts increases linearly with host density (see third term in Eq. (1b)). However, also the strength of selection favoring parasites increases with increasing host density. As the symbiotic community becomes more parasitic behind the wave front, the host growth rate decreases, causing in turn a decrease in the growth rate of the whole symbiotic community. Thus, the maximal symbiont growth rate is located just behind the maximal growth rate of the host population, where host density is large enough to support symbiont growth, but low enough to limit the growth of parasites. As the host population colonizes first, followed by its associated symbiotic community, the expansion is pulled by the host population, which benefits from the high proportion of mutualists at the range edge, but is pushed by the symbionts, which generate mutualists from mutation within the population core (Fig. C5).

The expansion dynamics of host-associated communities is therefore the result of a unique interaction of pushed and pulled waves. That is, the speed of spread depends on both the characteristics of the wave core as well as its leading edge. We have derived approximations for the speed of spread in SI.C, shown in Fig. 3 (solid curves), which combine information on the mean trait of the population in the core with the growth dynamics at the leading edge of the wave. Our approximations capture the general shape of the speed function, but under-estimate the speed of spread. This difference is partly due to spatial heterogeneity in the mean mutualistic investment of the symbiont community $\overline{\alpha}$, which is not embedded our approximation (i.e., in our analytical approximation $\overline{\alpha}$ is considered to be constant, while effectively $\overline{\alpha}$ increases in space x, see Fig. 2).

Fig. 2 Panel (a), Travelling wave profile at steady state. Plot of AMF density for increasing values of α (green to blue curves), total AMF density with respect to α (black curve), plant density (dashed black curve), and mean AMF trait $\overline{\alpha}$ (thick solid red curve). The crosses (and vertical dotted lines) indicate the positions along the travelling wave corresponding to the curves in panel (b). Panel (b), The distribution of the trait α at four different positions along the travelling wavefront. For both plots, $\beta = 0.4$. Panel (c), Cumulative densities of parasitic AMF ($\alpha \leqslant \alpha_c$, green area) and mutualistic AMF ($\alpha > \alpha_c$, blue area) inside the traveling front (black curve). Red curve corresponds to the proportion of parasitic AMF along the traveling wave. Panel (d), Growth rate of the plant (dashed red curve) and AMF (solid red curve) as a function of position along the travelling wave (plant - dashed black curve, AMF - solid black curve). Parameter values used for the simulations are: $D_m = D_p = 0.1$, $d_m = 0.01$, $Q = 6$, $\mu_p = \mu_m = 0.3$, $\beta = 0.72$, and $\alpha_{\min} = 0$ and $\alpha_{\max} = 5$.

The spread of a host population and its symbionts

Resource supply from the hosts to the symbionts determines the speed of spread of the whole community (Fig. 3). The speed of spread of host-symbiont communities reflects a similar broad unimodal dependence already observed for the plant and AMF densities (cfr. Fig. 1(a) and 3(a)). We find that the speed of spread is maximal when conditions in the core of the population maximise the density of the symbiont community rather than the density of the plants (i.e., the maximal speed of spread is obtained around $\beta = 0.86$, a value close to the one which maximizes symbiont density at equilibrium, $\beta = 0.72$). Indeed, as discussed in the previous section, the wave of expansion is pushed by the symbionts (Fig. 2(d) and Fig. C5). Thus, large carbon supply from the host,

(a)   (b)

Fig. 3 Panel (a), Speed of spread of AMF and plants with different relative dispersal abilities described by the diffusivity ratio (D_p/D_m): slow plant (blue), slow AMF (orange), and both identical (red); Panel (b) Difference between the speed of spread of a plant with and without an AMF community for various plants with different carbon supply rate (β) and different degree of dependence on the symbiont: obligate mutualistic plants (blue, $r_p = 0$), facultative mutualistic plants with various intrinsic growth rates, $r_p = 0.1$ (red), $rp = 0.2$ (orange) and $r_p = 0.8$ (purple). For the simulations, the parameters are: $d_m = 0.01$, $Q = 6$, $\mu_p = \mu_m = 0.3$, and $\alpha_{\min} = 0$ and $\alpha_{\max} = 5$.

despite not resulting in maximal plant density, may accelerate range expansion of a given host-symbiont community. Any difference in the dispersal abilities of the host and the symbionts decreases the speed of spread (compare the blue and yellow curves with the red curve of Fig. 3(a)), although the carbon supply rate that maximizes the speed of spread remains essentially the same (compare the location of the peak in the blue, yellow, and red curves of Fig. 3(a)).

As the wave of expansion is pulled by the plants (Fig. 2(d) and Fig. C5), the speed of spread increases if the plants can grow in the absence of the symbiont community (Fig. C4(b)). However, the speed advantage provided by the symbiont community depends on the obligate degree of the plant. When considering the difference between the speed of spread in the absence and presence of symbionts, we show that symbionts can induce either a positive or negative effect on the speed of spread (Fig. 3(b)). When the plants have a small intrinsic growth rate and need the symbiont community for growth, the symbionts enhance the speed of spread. For plants with a large intrinsic growth rate, however, the presence of symbionts reduces the speed of spread. The speed load induced by the symbionts depends on the cost of the symbionts to the host, which corresponds to the carbon supply β. In conclusion, strategies that decrease the dependency of hosts on their symbionts (such as a transition from obligate to facultative mutualism) or, alternatively, strategies that increase the symbiont density in the core of the population (such as selection for a carbon supply rate that maximizes symbiont density), enhance the speed of spread of the whole host-symbiont community.

Discussion

The Eco-evolutionary dynamics of host-associated communities

There have been several calls for investigation of the dynamics that emerge when evolutionary and ecological changes in symbiotic communities occur at the same timescale (Koskella et al, 2017; Fitzpatrick et al, 2020; Drew et al, 2021b). Evolutionary and ecological forces are known to drive the spread of pathogens in host populations (Restif, 2009; Day et al, 2020). However, our understanding of eco-evolutionary dynamics involving mutualistic symbionts remains limited. Here we present a theoretical framework that allows us to investigate, under minimal assumptions, the linked population dynamics, spread, and evolution of a host population and its associated mutualistic and parasitic symbionts. We show how mutualism can emerge from a parasitic community of symbionts, and stably persist in a community in which parasites and mutualists coexist and rapidly evolve along the parasitism-mutualism continuum (Lin and Koskella, 2015; Rogalski et al, 2021). Our framework accounts for unique characteristics of host-associated communities, such as their short generation time and horizontal modes of transmission, and constitutes the basis for further investigation of the eco-evolutionary dynamics of host-symbiont communities.

In our model, coexistence of parasites and mutualists occurs thanks to two key features: (1) a mutation-selection balance acting on the symbionts at the individual scale, and (2) a positive eco-evolutionary feedback acting at the host-community level. At the individual scale, parasitic symbionts benefit from a higher fitness than mutualistic symbionts, because they receive the same benefit from the host at a smaller cost. However, the average mutualistic investment of the whole community, and not the individual contribution of each symbiont, determines whether a community will establish and be mutualistic as a whole, as observed in previous modeling work (Archetti and Scheuring, 2011, 2013; Martignoni et al, 2020). Thus, in our context, the evolution of mutualism looks similar to the evolution of altruism. Even though the emergence of altruism is a fraternal transition (i.e. is arising from a division of labour among individuals of the same species), whereas the transition from parasitism to mutualism is egalitarian (i.e. resulting from the association of different species to complement their function), both result from the conflict between two levels of selection: the individual-level, favouring parasites, and the group-level, favoring altruists or mutualists (Wilson and Sober, 1989; Van Baalen and Rand, 1998; Simon et al, 2013).

Previous work has shown a strong shift from mutualism to parasitism with increasing availability of resources, with more beneficial symbionts dominating the community when host productivity is low, and parasites dominating when host productivity is high (Schwartz and Hoeksema, 1998; Hochberg et al, 2000; Hochberg and van Baalen, 2000; Neuhauser and Fargione, 2004). Similarly, in our simulations the selective advantage of parasites increases with the amount of resource available (i.e. with host density). Thus, high host density promotes the establishment of parasitic communities and, conversely, low host density reduces the selective advantage of parasitic symbionts, causing the community to become more mutualistic on average. Fluctuations in selection pressures due to, for instance, nonlinear public goods Archetti and Scheuring (2013), genotype-environment interactions (Parker,

1995), negative frequency-dependent selection (Bever, 1999; Brown and Tellier, 2011) or selection mosaics (Thompson, 2005) are known to maintain variation in partner quality (Mitchell-Olds et al, 2007). Here we show that fluctuation in selection pressures, due to variation in host density, can induce a shift from parasitism to mutualism in a symbiotic community.

Interesting future directions could consider how the inclusion of additional mechanisms, such as host control and different symbiont transmission modes, may improve the quality of the mutualistic interactions between hosts and symbionts. For example, host phenotypic plasticity in its mutualistic investment toward the symbionts may facilitate the evolution of more mutualistic traits (Koskella and Bergelson, 2020; Hou et al, 2021), even though the evolution of mutualism among hosts occurs after the emergence of mutualism among symbionts (Ledru et al, 2022). Strict vertical transmission of symbionts could also enhance evolution toward stronger mutualistic communities (Ewald, 1987; Sachs et al, 2004). Competition between hosts can also shape community structure: More mutualistic symbiotic communities should provide a fitness advantage to their host, which could drive the selection for communities with a lower proportion of parasites (Hartnett et al, 1993; Jones et al, 2012). Extension of our current model would allow us to investigate the evolutionary potential of these additional mechanisms.

Mutualistic investment at the range edges

Range expansion can result from various mechanisms involving growth and dispersal. Generally, we distinguish between two types of expansions: those that are pushed *versus* those that are pulled. In a pulled propagating wave, the population expands its range thanks to a higher growth rate at the population edges, while in a pushed wave population expansion is driven by a higher growth rate in the core (but see (Gandhi et al, 2016; Miller et al, 2020; Erm and Phillips, 2020)). Here, host-symbiont communities follow a combined expansion dynamic in which the wave of expansion is pulled by the hosts and pushed by the symbionts.

When host-symbiont communities expand their range, the selective pressure on symbionts becomes a function of space, due to changes in host density along the travelling wave, with strong selection favouring parasitic symbionts in the population core (where host density is large) but not at the population edges. The low density of plants at the population edges promotes the formation of a more mutualistic community, causing plant growth rate to be maximal at the wave front. Thus, the host plants pull the expansion wave by providing new resources to their obligate symbionts. In contrast to the pulling dynamics of the plants, the wave of expansion is pushed by the symbionts from the population core. The symbiotic community in the population core is more parasitic than that at the wave front, due to higher resource availability. As discussed in the previous section, parasitic symbionts are the common ancestors of good mutualists at the wave front. Thus, these symbionts are pushed from the population core into the front and closely follow their hosts in their expansion dynamics.

Spatial self-organization has been found to increase the abundance of mutualistic symbionts during range expansion in theoretical models (Momeni et al, 2013; Van Dyken et al, 2013) and synthetic microbial communities (Pande et al, 2016; Amor

et al, 2017; Rodríguez Amor and Dal Bello, 2019), with cheaters or parasitic symbionts lagging behind the mutualists at the leading edge. Emerging evidence also indicates that under stressful conditions plants may actively 'cry for help' (i.e., by secreting chemical compounds) and recruit beneficial symbionts (Schuman et al, 2015; Rizaludin et al, 2021). In our work, spatial selection occurs in the absence of built-in mechanisms. Instead, it is spontaneously driven by lower host density at the leading edges, which reduces the selection strength against mutualists in this region.

Invasion by alien organisms is often studied from the perspective of a single population, despite the fact that the success of an invasion might require the simultaneous successful spread of the focal population and its invisible symbionts (Dickie et al, 2017). Eradication and restoration strategies should therefore consider both hosts and symbionts as active and tightly-linked players in invasion dynamics. As conservation strategies to limit pulled and pushed invasions differ (Gandhi et al, 2016), the combined push-pull dynamics of linked host-symbiont invasions may require novel containment strategies (Taylor and Hastings, 2004). For example, limiting the invasive spread of species expanding as a pulled wave is currently accomplished by eradicating invaders at the population edges. However, this strategy may be inefficient if expansion and evolution of the symbionts in the population core facilitates the quick restoration of the population at the range edges. Our general framework may be adapted to host-symbiont populations expanding over a heterogeneous landscape, such as a fragmented or disturbed habitat (Willing et al, 2021), and to consider range overlaps with native populations (Dickie et al, 2017).

The spread of a host population and its symbionts

Currently, the effect of mutualistic interactions between hosts and symbionts on their range expansion is unclear. On the one hand, the growth benefits provided by mutualism can increase the speed of spread of the host, while on the other hand the absence of mutualistic partners at the range edge may produce an Allee effect that limits the rate of spread into new territories (Stanton-Geddes and Anderson, 2011; Kubisch et al, 2014; Fowler et al, 2023). Our framework allows us to disentangle these two tendencies and study how resource exchange (determining fitness of hosts and symbionts along the expansion range) and differences in host and symbiont dispersal abilities can affect the speed of spread of the community.

The 'enemy release hypothesis' postulates that species can escape their enemies (such as predators, pathogens, parasites, or herbivores) when introduced into a new range (Torchin et al, 2003; Colautti et al, 2004). A similar effect can occur at the edge of an expanding population, if enemies disperse more slowly than their hosts (Fagan et al, 2002; Nomikou et al, 2003). In contrast, if the linked species is beneficial, an expanding host population may also escape its mutualistic symbionts (Dickie et al, 2017; Shaw, 2022), to the detriment of its speed of spread. A similar behaviour occurs in cooperative systems, where the phenotype with the slowest speed of spread limits the spreading speed of the community Li et al (2005). In agreement with these results, we found that differences in the dispersal abilities of hosts and symbionts may reduce expansion rates. Thus, symbiont inoculation at the leading edges can also contribute to accelerating range expansion if symbionts have limited dispersal ability.We also found

that the optimal carbon supply rate, i.e., the one that maximizes AMF density at equilibrium, corresponds to the rate that maximizes the speed of spread of the whole host-symbiont community. Thus, hosts investment in their mutualistic relationship with symbionts may be considered an evolutionary strategy to increase the colonization speed of the host-symbiont community, and might also compensate for an eventual decrease in the speed of spread due to a lack of symbionts at the leading edge.

Our results also show that the speed of spread of host-symbiont communities can be maximized with reduced host dependence on the mutualism (from obligate to facultative). It follows that the evolution of reduced dependence of a host on its symbionts at the range edges of an expanding population can potentially facilitate plant invasion (Seifert et al, 2009). Note, however, that we only included one axis of benefit to the hosts, that is, one symbiont-supplied resource. Mutualistic interactions can be multidimensional; AMF, in particular, have been found to provide a suite of benefits to their hosts, such as protection against pathogens or abiotic factors (Smith and Read, 2010; Fowler et al, 2023). The inclusion of these other benefit axes might reduce the benefit of a host reducing its dependence on symbionts.

Our model could be extended to consider the evolution of obligate mutualism from facultative mutualism. Indeed, hosts that are facultative mutualists may reach higher densities in the absence of symbionts. Subsequent colonization of these hosts with parasites, and the evolution of some mutualistic traits in these parasites, may be sufficient to drive a transition from parasitism to mutualism in the community. Future studies are needed to consider how, once mutualism has emerged, host adaptation may lead to a strategy increasing investment into the mutualistic relationship with symbionts, rather than into self-growth (Ledru et al, 2022). Similarly, the model could be extended to consider the evolution of microorganisms from free-living to host-associated states (Drew et al, 2021b).

Finally, it might therefore be interesting to investigate the co-evolution between the mutualistic investment of symbionts and their dispersal ability. Indeed, in some circumstances dispersal ability have been found to evolve during range expansion (Urban et al, 2007; Brown and Vellend, 2014). A recent study (Ledru et al, 2022) has shown that the co-evolution between mutualistic effort and dispersal results in mutualistic symbionts that disperse locally (i.e., have low dispersal ability), while parasitic symbionts disperse over longer distances (i.e., have high dispersal ability). Such negative correlation between dispersal ability and mutualistic quality has also been observed in the context of altruism (Koella, 2000; Le Galliard et al, 2005; Hochberg et al, 2008; Eldakar et al, 2010; Purcell et al, 2012; Mullon et al, 2018), and between local interactions and evolution of avirulence (Boots and Mealor, 2007). In a range expansion context, this correlation might be reversed because the mutualistic symbiont has an evolutionary advantage at the leading edge.

Conclusion and future work

Current ecological and evolutionary theory fails to explain the complex dynamics structuring stable host-symbiont communities (Koskella et al, 2017). This gap is at

least partly due to unique characteristics of these communities, such as rapid evolution, interdependent fitness, and horizontal transmission of symbionts, which make study and analysis challenging. Here we present a theoretical framework that contributes to our understanding of host-symbiont linked population dynamics, evolution, and spread, with possible implications for microbiome research (Koskella et al, 2017; Fitzpatrick et al, 2020) and for the management of linked plant-microbial invasions (Dickie et al, 2017).

Appendix A   No evolution no space, competitive exclusion and parasitic threshold

We first look at the dynamical system without spacial component and evolution of the trait α.

Monomorphic population of AMF.

We first look at a population of symbionts that are monomoprhic, that is all the individuals share the same trait α. In this situation, the system is reduce to the following differential equations for the biomass of plant P and the biomass of the monomorphic symbionts M:

$$P'(t) = P \left(\frac{Q\,\alpha M(t)}{P(t) + d} - \beta\,M(t) - \mu_p P(t) \right), \ t > 0 \tag{A1a}$$

$$M'(t) = M(t) \left(\beta\,P(t) - \frac{\alpha P(t)}{P(t) + d} - \mu_m M(t) \right). \tag{A1b}$$

From the analysis of Martignoni et al (2020), the system admits the extinction steady state $(0, 0)$, which might be stable if $d < \alpha/\beta$ and a stable positive steady state (P, M) if the following properties are satisfied:

- $Q > 1$
- $\alpha \geqslant \alpha_c(\beta)$ where $\alpha_c(\beta)$ satisfies

$$\alpha_c(\beta) = \frac{\beta\,d}{Q} \frac{(Q + 1) - \sqrt{(Q - 1)^2 - 4\dfrac{Q\mu_m\mu_p}{\beta^2}}}{2} \tag{A2}$$

- $\mu_p\mu_m \leqslant \beta^2(Q - 1)^2/(4Q)$.

The assumption on α shows that the mutualistic investment of the symbiont needs to be large enough to persist with the plant, that is $\alpha \geqslant \alpha_c$ where the threshold satisfies

$$\alpha_c = \frac{\beta\,d}{Q} \frac{(Q + 1) - \sqrt{(Q - 1)^2 - 4\dfrac{Q\mu_m\mu_p}{\beta^2}}}{2} \geqslant \frac{\beta\,d}{Q} \tag{A3}$$

which means that the symbiont provides a positive benefit to the plant when the plant is at low density. In addition, the analysis first points out that if the monomorphic population of AMF have a low mutualistic quality α below the threshold α_c, then the system goes extinct. As such, we say the a symbionts is *parasitic* if its mutualsitic investment α is below α_c, while it is *mutualistic* if $\alpha > \alpha_c$.

Polymorphic population of AMF and competitive exclusion

On the other hand, if the symbiont population is initially polymorphic, with an initial trait density $m(t, \alpha)$ with $\alpha \in [\alpha_{\min}, \alpha_{\max}]$, then the population will converges toward

a monomorphic population with mutualistic quality $\alpha_{\min}$. If this quality is below the critical value α_c, the population will go extinct.

The population densities (P, m) satisfies in this case the following model:

$$P'(t) = P\left(\frac{Q\,\overline{\alpha}(t)M(t)}{P(t) + d} - \beta\,M(t) - \mu_p P(t)\right),\ t > 0 \tag{A4a}$$

$$\partial_t m(t, \alpha) = m\left(\beta\,P(t) - \frac{\alpha\,P(t)}{P(t) + d} - \mu_m M(t)\right),\ \alpha \in (\alpha_{\min}, \alpha_{\max}) \tag{A4b}$$

where the quantity $\overline{\alpha}(t)$ is the mean AMF trait in the community at location x and $M(t)$ is the total biomass of AMF, defined by

$$\overline{\alpha}(t) = \int_{\alpha_{\min}}^{\alpha_{\max}} \alpha\,\frac{m(t, \alpha)}{M(t)}\,d\alpha \ \ \text{and}\ \ M(t) = \int_{\alpha_{\min}}^{\alpha_{\max}} m(t, \alpha)\,d\alpha \tag{A5}$$

As the result, the total biomass of AMF satisfies the following equation

$$M'(t) = M\left(\beta\,P(t) - \frac{\overline{\alpha}(t)\,P(t)}{P(t) + d} - \mu_m M(t)\right) \tag{A6}$$

and the trait distribution $\phi_m(t, \alpha) = m(t, \alpha)/M(t)$ is described by the following model

$$\partial_t \phi_m(t, \alpha) = \phi_m(t, \alpha)\big(\overline{\alpha}(t) - \alpha\big)\frac{P(t)}{P(t) + d},\ \alpha \in (\alpha_{min}, \alpha_{max}),\ t > 0. \tag{A7}$$

From the definition of the mean trait $\overline{\alpha}$, it satisfies $\overline{\alpha}(t) \in [\alpha_{\min}, \alpha_{\max}]$ for all time $t > 0$ and for all $\alpha < \overline{\alpha}(t)$, $\phi_m(t, \alpha)$ is decreasing over time, while $\phi_m(t, \alpha)$ is increasing for $\alpha \leqslant \overline{\alpha}(t)$. Consequently, $\overline{\alpha}(t)$ is decreasing over time, and it converges to $\alpha_{\min}$, while ϕ_m converges toward the dirac mass at $\alpha = \alpha_{rmmin}$.

If the minimal value of the trait $\alpha_{\min}$ is below the persistence threshold α_c then the population goes extinct. So in absence of diversification of trait α, the population experience a competitive exclusion that can expose the population to extinction.

Appendix B   Evolution, no dispersal, analysis

To understand the co-evolution of a plant with its AMF community, we first look at the diversification of AMF for a given plant at a fixed location. In this situation, the plant with biomass $P(t)$ is able to provide carbon for the AMF with trait α and density $m(t, \alpha)$, at a rate β.

The model (1) then becomes

$$P'(t) = P(t)\left(\frac{Q\,\overline{\alpha}(t)\,M(t)}{P(t) + d} - \beta M(t) - \mu_p P(t)\right),\ t > 0 \tag{B8a}$$

$$\partial_t m(t, \alpha) = d_m \partial_\alpha^2 m(t, \alpha) + m(t, \alpha) \left(P(t) \left(\beta - \frac{\alpha}{P(t) + d} \right) - \mu_m M(t) \right), \alpha \in (\alpha_{min}, \alpha_{max})$$

$$(\text{B8b})$$

We here aim to understand the AMF community that emerges from the interaction with the plant and the evolutionary process. More precisely, we aim to characterize the steady states of this model (B8). We first prove the existence and some properties of the non trivial steady state and then we provide an approximation which corresponds to special case where $\alpha_{\max} = \infty$.

Let us first write the model (B8) with different variables. First, if we integrate over α the equation for m, we obtain the following system of equations for the total biomass of AMF M and plant P:

$$\begin{aligned} P'(t) &= P(t) \left(\frac{Q \, \overline{\alpha}(t) \, M(t)}{P(t) + d} - \beta M(t) - \mu_p P(t) \right) \\ M'(t) &= M(t) \left(P(t) \left(\beta - \frac{\overline{\alpha}(t)}{P(t) + d} \right) - \mu_m M(t) \right) \end{aligned} \quad , \ t > 0. \qquad (\text{B9})$$

Then, let us write an equation for the trait distribution of AMF $\phi_m(t, \alpha) = m(t, \alpha)/M(t)$. Using equation for m and M, we obtain

$$\begin{cases} \partial_t \phi_m(t, \alpha) = d_m \partial_\alpha^2 \phi_m(t, \alpha) + \phi_m(t, \alpha) \big(\overline{\alpha}(t) - \alpha \big) \dfrac{P(t)}{P(t) + d}, \ \alpha \in (\alpha_{min}, \alpha_{max}), \ t > 0 \\ \partial_\alpha \phi_m(t, \alpha_{min}) = \partial_\alpha \phi_m(t, \alpha_{max}) = 0, \ t > 0 \end{cases}$$

$$(\text{B10})$$

B.1 Steady state.

A non trivial steady state $(P, m(\alpha))$ of the model (1) without spatial dispersal ($D_p = D_m = 0$, should satisfy the two following problems. First, the biomass of the plant P and the total biomass of AMF $M = \int m(\alpha) d\alpha$ solve the following system

$$\begin{aligned} 0 &= \frac{Q \, \overline{\alpha} \, M}{P + d} - \beta M - \mu_p P \\ 0 &= P \left(\beta - \frac{\overline{\alpha}}{P + d} \right) - \mu_m M \end{aligned} \qquad (\text{B11})$$

where $\overline{\alpha}$ is the mean trait of the AMF distribution at steady state. And the trait distribution at equilibrium $\phi_m(\alpha) = m(\alpha)/M$ satisfies the following elliptic problem

$$
\begin{cases}
-d_m \phi_m'' + \alpha \dfrac{P}{P+d} \phi_m = \overline{\alpha} \dfrac{P}{P+d} \phi_m, & \text{in } (\alpha_{min}, \alpha_{max}) \\
\phi_m'(\alpha_{min}) = \phi_m'(\alpha_{max}) = 0 \\
\phi_m > 0
\end{cases}
\tag{B12}
$$

Proposition 1. *If, the diversification rate $d_m > 0$ and the following inequality holds true*

$$
\frac{\alpha_{min} + \alpha_{max}}{2} > \alpha_c(\beta) \quad \text{where} \quad \alpha_c(\beta) = \frac{\beta \, d}{Q} \frac{(Q+1) - \sqrt{(Q-1)^2 - 4\dfrac{Q\mu_m\mu_p}{\beta^2}}}{2} \tag{B13}
$$

then, there exists a steady state of the model (B8).

In addition, the mean trait $\overline{\alpha}$ at equilibrium satisfies the following property

$$
\alpha_c(\beta) \leqslant \overline{\alpha} \leqslant \frac{\alpha_{\min} + \alpha_{\max}}{2} \tag{B14}
$$

Proof of Proposition 1. If a non trivial steady state $(P, m(\alpha))$ of the system (B8) exists then the total biomass (P, M) should satisfy the following system, which depends on $\overline{\alpha}$,

$$
\begin{aligned}
0 &= \frac{Q\,\overline{\alpha}\,M}{P+d} - \beta M - \mu_p P \\
0 &= P\left(\beta - \frac{\overline{\alpha}}{P+d}\right) - \mu_m M
\end{aligned}
\tag{B15}
$$

From Martignoni et al (2020), we know that the system admits at least a positive solution, if the following conditions holds true

$$
Q > 1 \quad \text{and} \quad \frac{\sqrt{4Q\mu_m\mu_p}}{|Q-1|} < \beta < Q\frac{\overline{\alpha}}{d} \tag{B16}
$$

Under this condition, the following equilibrium exists

$$
\begin{aligned}
P(\overline{\alpha}) &= d\left(\frac{\overline{\alpha}}{\alpha_c} - 1\right), \quad \text{with} \quad \alpha_c(\beta) = \frac{\beta\,d}{Q} \frac{(Q+1) - \sqrt{(Q-1)^2 - 4\dfrac{Q\mu_m\mu_p}{\beta^2}}}{2} \\
M(\overline{\alpha}) &= \frac{P(\overline{\alpha})}{\mu_m}\left(\beta - \frac{\overline{\alpha}}{P+d}\right)
\end{aligned}
\tag{B17}
$$

In addition, we can show from (B10) that $(\overline{\alpha}, \phi_m)$ solves the following problem, which depends on P

$$-d_m \phi_m''(\alpha) + \alpha \frac{P}{P+d} \phi_m(\alpha) = \overline{\alpha} \frac{P}{P+d} \phi_m(\alpha), \ \alpha \in (\alpha_{min}, \alpha_{max}),$$

$$\phi_m'(\alpha_{min}) = \phi_m'(\alpha_{max}) = 0.$$

(B18)

However, we know that for any positive P, the Neumann principal eigenproblem

$$\begin{cases} -d_m \phi'' + \alpha \dfrac{P}{P+d} \phi = \lambda \phi, & \text{in} \ \ (\alpha_{min}, \alpha_{max}) \\ \phi'(\alpha_{min}) = \phi'(\alpha_{max}) = 0 \\ \phi > 0 \end{cases}$$

(B19)

admits a principal eigenpair (λ_P, ϕ_P), normalized by $\int \phi_P^2 = 1$ and we have the variational formula

$$\lambda_P = \min \left\{ Q_P(\phi) := \int_{\alpha_{min}}^{\alpha_{max}} \left(d_m \phi'^2 + \alpha \frac{P}{P+d} \phi^2 \right) d\alpha, \ \text{for} \ \phi \in E \right\}, \quad (B20)$$

where

$$E = \left\{ \phi \in H^1(\alpha_{min}, \alpha_{max}) \ \text{and} \ \int \phi^2 = 1 \right\}. \quad (B21)$$

If the parameter, $d_m > 0$, the eigenpair (λ_P, ϕ_P) satisfies the following properties
(i) for all $P > 0$, we have

$$\frac{P}{P+d} \alpha_{min} < \lambda_P < \frac{P}{P+d} \frac{\alpha_{min} + \alpha_{max}}{2} \quad (B22)$$

(ii) the function $P \mapsto \lambda_P$ is increasing and concave in $(0, \infty)$, and

$$\lim_{P \to 0} \lambda_P = 0, \ \lambda_P \underset{P \to 0}{\sim} \frac{P}{P+d} \frac{\alpha_{min} + \alpha_{max}}{2} \ \text{and} \ \lim_{P \to \infty} \lambda_P = \lambda_{d_m} < \frac{\alpha_{min} + \alpha_{max}}{2}$$

(B23)

where $\lambda_{d_m} > 0$ is the principal eigenvalue of the problem when $P/(P+d) = 1$.
(iii) the eigenfunction ϕ is is decreasing on $[\alpha_{\min}, \alpha_{\max}]$ and it is strictly concave on $[\alpha_{\min}, \lambda_P)$ and strictly convex on $(\lambda_P, \alpha_{\max}]$.
Since $\overline{\alpha} = \dfrac{P+d}{d} \alpha_c(\beta)$ from (B17), we know that there exists a unique positive P such that $\lambda_P = \overline{\alpha} P/(P+d)$ if and only if the following inequality holds true

$$\frac{\alpha_{min} + \alpha_{max}}{2} > \alpha_c(\beta). \quad (B24)$$

This inequality ensures the existence of a unique steady state $(P, M, \overline{\alpha}, \phi_m)$ which satisfies simultaneously (B17) and (B12).

Moreover, this inequality shows that when $\beta \to \beta_{max}$ such that inequality (B24) is an equality, then the mean trait in the AMF population converges towards $\dfrac{\alpha_{min} + \alpha_{max}}{2}$ and thus the distribution ϕ_m converges towards the uniform distribution $\phi_m(\alpha) = 1/(\alpha_{max} - \alpha_{min})$ for all $\alpha \in (\alpha_{min}, \alpha_{max})$.

So, if $\beta = \beta_{max}$ and competition between AMF all sharing one plant, we expect a uniform distribution of AMF biomass across the range of α values (Fig. 1b).

In addition, if the plant density is low $P \to 0$ then the mean AMF trait should be close to $\frac{(\alpha_{min} + \alpha_{max})}{2}$ and thus the AMF distribution should be uniform over α. As a result, when the plant density is low, we expect the AMF distribution to be uniform

Moreover, the presence of mutation through the diffusion operator is crucial for the existence of a non trivial positive steady state. In absence of mutation, the competitive exclusion drives the population toward extinction.

B.2 Approximation of the steady states

We aim to find some approximations for $\overline{\alpha}$, P, M, and m. First, we assume that

$$\alpha_{min} = 0 \quad \text{and} \quad \alpha_{max} = \infty. \tag{B25}$$

Exact solution for the eigenpair problem.

In this situation, we can compute an explicit solution of (B19) using the Airy function Ai, which satisfies $\mathrm{Ai}''(z) - z\,\mathrm{Ai}(z) = 0$ for all $z \in \mathbb{R}$. Indeed, the following function ϕ is a positive solution of (B19):

$$\phi(\alpha) = \frac{\mathrm{Ai}\left(\left(\dfrac{P}{(P+d)d_m}\right)^{1/3}\alpha - z_0\right)}{\displaystyle\int_{-z_0}^{\infty}\left(\dfrac{P}{(P+d)d_m}\right)^{-1/3}A_i(z)dz} \quad \text{and} \quad \lambda = z_0\, d_m^{1/3}\left(\frac{P}{(P+d)}\right)^{2/3} \tag{B26}$$

where $-z_0$ is the maximal value such that Ai achieves a maximum at $-z_0$ and the mean trait $\overline{\alpha}$ and P solves the following system:

$$\begin{cases} P = d\left(\dfrac{\overline{\alpha}}{\alpha_c} - 1\right) \\[2mm] \overline{\alpha}\dfrac{P}{P+d} = \lambda = z_0\, d_m^{1/3}\left(\dfrac{P}{(P+d)}\right)^{2/3} \end{cases} \tag{B27}$$

The following system can be reduced to $P = d(\overline{\alpha}/\alpha_c - 1)$ and the mean trait $\overline{\alpha}$ is the real positive root of the following polynomial function $A(X) = X^3 - \alpha_c(\beta)X^2 - z_0^3\, d_m$.

B.3 Effect of the diversification rate d_m and plant biomass

We can write the previous system (B27) in the following form

$$\overline{\alpha} = \frac{P+d}{d}\alpha_c(\beta) \tag{B28}$$

$$\overline{\alpha} = z_0 \left(\frac{d_m}{P}(P+d) \right)^{1/3}. \tag{B29}$$

First, the mean trait $\overline{\alpha}$ is increasing with the diversification rate d_m. As a result, the biomass of the plant as well as the total biomass of AMF is also increasing with the diversification rate d_m (Fig. B1. Secondly, the plant biomass P has antagonistic effects on the mean trait $\overline{\alpha}$. The ecological dynamics, described by (B28), increases the mean trait with the biomass of the plant P. A more mutualsitic community will support a larger plant (see Martignoni et al, 2020). While, the evolutionary dynamics described by (B29), decreases the mean trait when the bimoass of the plant increases. Indeed, in the evolutionary dynamics, the strength of selection, tha is $P/(P+d)$, is driven by the biomass of the plant. If the biomass of the plant is large, selection is strong and it drives the trait distribution toward small values of α. Conversely, when the biomass of the plant is small, selection is weak and the mutations drive the trait distribution toward larger α. The equilibrium results from the balance between the ecological forces which promotes mutualism and evolutionary forces enhancing parasitism.

B.4 Proportion of parasitic symbionts

In previous section, we introduce the critical value α_c of the trait, such that if the plant is in association with a single symbiont with trait $\alpha \leqslant \alpha_c$, then the plant-symbiont system goes extinct. Conversely, the system survives if $\alpha > \alpha_c$. The critical value $\alpha_c(\beta)$ does depend on the mutualistic investment of the plant β.

In a polymorphic community, where the trait can range from $\alpha_{\min}$ to $\alpha_{\max}$ such that $\alpha_{\min} < \alpha_c < \alpha_{\max}$, the symbionts can be split in two groups: parasitic symbionts ($\alpha < \alpha_c$) and mutualistic symbionts ($\alpha \geqslant \alpha_c$). We investigate the proportion of parasitic and mutualistic symbiont in the community for different plant (Fig. B2). The proportion is defined by

$$prop = \frac{\displaystyle\int_{\alpha_{\min}}^{\alpha_c} m(\alpha)d\alpha}{M} = \int_{\alpha_{\min}}^{\alpha_c} \phi_m(\alpha)d\alpha \tag{B30}$$

With the approximation of ϕ_m defined in (B19), we obtain

$$prop = \frac{\displaystyle\int_{-z_0}^{\left(\frac{P}{(P+d)d_m}\right)^{1/3}\alpha_c - z_0} \mathrm{Ai}(\alpha)d\alpha}{\displaystyle\int_{-z_0}^{\infty} \mathrm{Ai}(\alpha)d\alpha} = \frac{\displaystyle\int_{-z_0}^{z_0\left(\frac{\alpha_c}{\overline{\alpha}}-1\right)} \mathrm{Ai}(\alpha)d\alpha}{\displaystyle\int_{-z_0}^{\infty} \mathrm{Ai}(\alpha)d\alpha} \tag{B31}$$

(a)

(b) $\beta = 0.43$ (c) $\beta = 0.72$

Fig. B1 Effect of the diversification rate d_m on the biomass of the plant (black curves and triangles) and the total biomass of the AMF (blue curves and dots), and the mean mutualistic investment $\overline{\alpha}$ (red curves and dots) at eco-evolutionary equilibrium. For the simulations, the parameters are: $Q = 6$, $\mu_p = \mu_m = 0.3$, and $\alpha_{\min} = 0$ and $\alpha_{\max} = 5$.

On the one hand the proportion of parasitic symbionts increases with the plant biomass if the parasitic threshold α_c remains constant. And the proportion is decreasing with the diversification rate. On the other hand, the proportion of parasitic symbionts depends on the ratio between the critical threshold $\alpha_c(\beta)$ and the mean trait $\overline{\alpha}$. Since $\overline{\alpha}$ is increasing with respect to the diversification rate d_m, the proportion of parasits in the community decreases with the diversification rate, as expected by the previous results (Fig. B2(a-b)).

At equilibrium, the expression of α_c and the relationship between P and $\overline{\alpha}$ (B28) provide the following expression:

$$prop = \frac{\displaystyle\int_{-z_0}^{-z_0 \frac{P}{P+d}} \mathrm{Ai}(\alpha)d\alpha}{\displaystyle\int_{-z_0}^{\infty} \mathrm{Ai}(\alpha)d\alpha} \tag{B32}$$

In this case, the porportion of parasitic symbionts depends on the bimoass of the plant at equilibrium. The effect of the mutualistic investment of the plant on the proportion is thus not monotonic because, the biomass of the plant is not monotonic with the parameter β. However, we have shown that the plant biomass at equilibrium is increasing and then decreasing with respect to β. The proportion of parasitic symbionts is decreasing and then increasing with respect to the mutualistic investment of the plant β (Fig. B2).

B.5 Lineages among the symbionts community

We aim to understand the lineages of individuals present at time t in the community at equilibrium. We first track the offspring of individuals, which are present initially, using the inside dynamics approach developed by Roques et al (2012). Individuals are labeled, and transmit their label to their offspring. Since individuals only differ by their label and their trait, each label $k \in \mathbb{N}$ corresponds to a neutral fraction of density v^k inside the community at equilibrium $m(\alpha)$. Initially we assume that

$$m(\alpha) = \sum_{k \geqslant 1} v^k(0, \alpha), \text{ for all } \alpha \in (\alpha_{\min}, \alpha_{\max}). \tag{B33}$$

Their dynamics is described by

$$\partial_t v^k(t, \alpha) = d_m \partial_\alpha^2 v^k(t, \alpha) + v^k(t, \alpha)\left(P\left(\beta - \frac{\alpha}{P+d}\right) - \mu_m M\right) \tag{B34}$$

where $(P, M, \overline{\alpha})$ are the equilibrium satisfying (B11)-(B12). From Cloez and Gabriel (2020), the solution v^k converges uniformly in space toward the following quantity $p[v_0^k]m(\alpha)$ where the positive scalar $p[v_0]$ is defined by

$$p[v_0] = \frac{\displaystyle\int_{\alpha_{\min}}^{\alpha_{\max}} v_0(\alpha)m(\alpha)d\alpha}{\displaystyle\int_{\alpha_{\min}}^{\alpha_{\max}} m^2(\alpha)d\alpha} \text{ and } v_0(\alpha) = v(0, \alpha) \leqslant m(\alpha). \tag{B35}$$

Using this forward approach, we are able to define a backward *ancestral process* Y_s, for each individual with trait α at time t (see Forien et al (2022) for more details on the derivation). More precisely, the ancestral process at time s aims to describe the

Fig. B2 Proportion of parasitic symbionts in the symbionts community associated with an obligate (panel (a-c) $r_p = 0$) or facultative (panel (b-d) $r_p = 0.2$) mutualistic plant with various carbon supply ability β. Blue dots corresponds to the proportion at equilibrium from the model (1) and the blue curve corresponds to the approximation defined by (B26). Biomass of the plant (black curve and triangles) and mean mutualistic investment of AMF in the community (red curve and dots). Grey curve corresponds to the mutualistic threshold α_c. Dashed black line in panel (d) is the approximation for the obligate mutualistic plant biomass. For the simulations, the parameters are: $d_m = 0.01$, $Q = 6$, $\mu_p = \mu_m = 0.3$, $\beta = 0.43$ and $\alpha_{\min} = 0$ and $\alpha_{\max} = 5$.

trait of the ancestor (alive at time $t - s$) of an individual sampled randomly among individuals with trait α at time t. The ancestral process Y_s is related to the previous model through the following relationship for all $(t, \alpha) \in (0, \infty) \times (\alpha_{\min}, \alpha_{\max})$ and $k \in \mathbb{N}$:

$$\frac{v^k(t, \alpha)}{m(\alpha)} = \mathbb{E}_\alpha \left[\frac{v^k(s, Y_{t-s})}{m(Y_{t-s})} \right], \ 0 \leqslant s \leqslant t. \tag{B36}$$

This equality means that the probability of sampling an individual of type k among individual with α at time t is equal to the probability of drawing an individual of type

k among individuals with trait y in the past, at the time $t - s$ (right hand side). In particular, the previous long time behavior property (B.5), shows that the ancestral process converges as $s \to \infty$ towards a random variable Y_∞, which admits the density

$$\alpha \mapsto \frac{m^2(\alpha)}{\displaystyle\int_{\alpha_{\min}}^{\alpha_{\max}} m^2(\alpha)d\alpha} \tag{B37}$$

In particular, we show that the most likely common ancestor is a parasitic symbiont with trait $\alpha = \alpha_{\min}$. In addition, on average the trait of the common ancestor is smaller than the mean trait of the actual community (see red and orange curves in Fig. B3). In addition, for small β, the mean trait of the common ancestor is mutualistic while for large β, it is parasitic (see orange curve in Fig. B3). As a result, we show that the equilibrium is most likely produced by parasitic individuals that mutate and generate the equilibrium distribution. So the mutualism persist in the population thanks to more parasitic individuals. This behavior persists even for facultative mutualistic plants (Fig. B3c).

Appendix C   Spreading with and without evolution

C.1   Spreading without evolution

First we investigate the spreading properties of the solution of our problem without variation in trait α. So we assume that AMF community is monomorphic: there is only one trait $\boldsymbol{\alpha}$ in the AMF population. We assume that $\boldsymbol{\alpha}$ is a real positive constant value between ($\alpha_{\min}$ and $(\alpha_{\max} + \alpha_{\min})/2$, so that the following system admits a constant positive steady state. We investigate the spreading speed of solutions of the following system of equations

$$\partial_t P(t, x) = D_p \partial_x^2 P + P\left(\frac{Q\,\boldsymbol{\alpha}M(t,x)}{P + d} - \beta\,M(t,x) - \mu_p P(t,x)\right), \ x \in \mathbb{R}, \ t > 0 \tag{C38}$$

$$\partial_t M(t, x) = D_m \partial_x^2 M + M\left(\beta\,P - \frac{\boldsymbol{\alpha}\,P}{P + d} - \mu_m M(t,x)\right). \tag{C39}$$

Under, the assumption of Proposition 1, there exists a positive steady state of the model. We aim to investigate the traveling wave solutions of this model, that are solutions of the form $P(x - ct)$ and $M(x - ct)$ where c is the spreading speed of the solution and P and M are profiles that connect the positive stationary state (P^*, M^*) to the trivial state $(0, 0)$. Let us remind tha the positive steady state satisfies

$$P^* = d\left(\frac{\boldsymbol{\alpha}}{\alpha_c} - 1\right) \tag{C40}$$

$$M^* = \frac{P^*}{\mu_m}\left(\beta - \frac{\alpha_c}{d}\right) \tag{C41}$$

Obligate mutualistic plant   Facultative mutualistic plant

(a) $(r_p = 0)$ (b) $(r_p = 0.2)$

(c) $(r_p = 0, \ \beta = 0.4)$ (d) $(r_p = 0.2, \ \beta = 0.4)$

Fig. B3 Panel (a-b), Probability of a parasitic common ancestor (dark blue curve and triangles) and mean mutualistic investment of the common ancestor (orange curve and diamonds) compared respectively with the proportion of parasitic symbiont in the AMF community at equilibrium without spatial spread (light blue curve and dots) and the mean mutualistic investment of the comunity (red curve and dots) associated with obligate mutualistic plant (panel (a)) and facultative mutualistic plant (panel c). Distribution of the mutualistic investment for the common ancestor (dark blue curve and triangles) and the AMF community (red curve and dots) with obligate mutualistic plant (panel (b)) and facultative mutualistic plant (panel d). The curves correspond to the approximations, while the markers correspond to the outcome of the model. For the simulations, the parameters are: $d_m = 0.01$, $D_p = D_m = 0.1$, $Q = 6$, $\mu_p = \mu_m = 0.3$, $\beta = 0.72$ and $\alpha_{\min} = 0$ and $\alpha_{\max} = 5$.

where $\alpha_c(\beta)$ is the mutualistic/parasitic threshold defined in (3) and it depends on the parameter β.

To investigate the spreading speed problem, we look at the following equations

$$\partial_t p(t, x) = D_p \partial_x^2 p + f_p p^2 (1 - p/P^*), \ x \in \mathbb{R}, \ t > 0 \qquad \text{(C42)}$$

$$\partial_t m(t, x) = D_m \partial_x^2 m + f_m m^2 (1 - m/M^*), \qquad \text{(C43)}$$

where f_p and f_m are positive constants. We know (Turchin, 1998) that these equations, which are linked only through the positive steady state (P^*, M^*), admit a unique traveling wave solutions, that move at a speed c_p and c_m respectively. The speed of spread are given by the following formula

$$c_p = \sqrt{\frac{D_p \, f_p \, P^*}{2}} \tag{C44}$$

$$c_m = \sqrt{\frac{D_m \, f_m \, M^*}{2}} \tag{C45}$$

For appropriate constant f_p and f_m, the associated solutions p and m should be sub-solutions of the initial problem (C38). More precisely, we choose f_p and f_m as follows. Let γ be a positive constant that we will determine later, and define

$$f_p(\gamma) = \left(\frac{Q \, \boldsymbol{\alpha}}{d} - \beta\right) \gamma - \mu_p \tag{C46}$$

$$f_m(\gamma) = \left(\beta - \frac{\boldsymbol{\alpha}}{d}\right) \frac{1}{\gamma} - \mu_m \tag{C47}$$

We choose γ such that $c_p(\gamma) = c_m(\gamma)$, so that the solutions p and m will spread at the same speed

$$c^* = c_p(\gamma^*) = c_m(\gamma^*). \tag{C48}$$

The parameter γ^* should be the positive root of the following polynomial function

$$G(X) = \left(\frac{Q \, \boldsymbol{\alpha}}{d} - \beta\right) X^2 + \left(\mu_m \frac{D_m \, M^*}{D_p \, P^*} - \mu_p\right) X - \frac{D_m \, M^*}{D_p \, P^*} \left(\beta - \frac{\boldsymbol{\alpha}}{d}\right) \tag{C49}$$

Since p and m are sub-solutions of the initial system, the solution (P, M) should travel at a speed $c > c^*$ (Fig. C4).

C.2 Spreading with evolution

Speed of spread

We now look at the spreading sped of the solution with evolution. In this case, we know that the biomass satisfy the following problem

$$\partial_t P(t,x) = D_p \partial_x^2 P + P \left(\frac{Q \, \overline{\alpha}(t,x) M(t,x)}{P + d} - \beta \, M(t,x) - \mu_p P(t,x)\right), \ x \in \mathbb{R}, \ t > 0 \tag{C50}$$

$$\partial_t M(t,x) = D_m \partial_x^2 M + + M \left(\beta \, P - \frac{\overline{\alpha}(t,x) \, P}{P + d} - \mu_m M(t,x)\right). \tag{C51}$$

where $\overline{\alpha}(t,x)$ is increasing and satisfies the following property $\overline{\alpha}(t,x) \to \overline{\alpha}$ as $t \to +\infty$ locally uniformly in space, where $\overline{\alpha}$ solves the steady state problem (B11)- (B12). From numerical simulation we see that $\overline{\alpha}(t,x)$ is increasing with respect to the space

<div style="text-align:center">(a)   (b)</div>

Fig. C4 Panel (a), Effect of the carbon supply rate β on the speed of spread a plant associated with one AMF with various fixed trait α (panel (a)): blue curve ($\alpha = 0.4$), red curve ($\alpha = 0.7$) and orange curve ($\alpha = 0.9$); Panel (b), Difference between the speed of spread of a plant with and without the monomorphic AMF community for various plants with different carbon supply rate and different degree of dependence on the symbiont: obligate mutualistic plants (blue, $r_p = 0$), facultative mutualistic plants with various intrinsic growth rates, $r_p = 0.1$ (red), $r_p = 0.2$ (orange) and $r_p = 0.8$ (purple). The markers correspond to the speed of the model (1) without evolution $d_m = 0$) and the curves correspond to the approximation speed (C48). For the simulations, the parameters are: $D_p = D_m = 0.1$, $Q = 6$, $\mu_p = \mu_m = 0.3$.

variable x. Under these conditions, we can use the previous sub-solutions to show that the spreading speed

$$c^* \geqslant c_p(\gamma^*(\overline{\alpha})). \tag{C52}$$

Fixation probability along the range expansion

Here, we aim to understand how the propagation occurs in space. We first assume that the population of host and symbionts are represented by traveling waves of profile $\mathbf{P}(x)$ and $\mathbf{m}(x, \alpha)$ respectively, which satisfy the following problem

$$D_p \mathbf{P}''(z) + c\mathbf{P}(z) \left(\frac{Q\,\overline{\alpha}(z)\mathbf{M}(z)}{\mathbf{P}+d} - \beta\,\mathbf{M}(z) - \mu_p\mathbf{P}(z) \right) = 0, \ z \in \mathbb{R} \tag{C53}$$

$$\begin{cases} D_m \partial_z^2 \mathbf{m}(z, \alpha) + d_m \partial_\alpha^2 \mathbf{m}(z, \alpha) + c\partial_z \mathbf{m}(z, \alpha) + \mathbf{m} \left(\beta\,\mathbf{P} - \dfrac{\overline{\alpha}(z)\,\mathbf{P}}{\mathbf{P}+d} - \mu_m\mathbf{M}(z) \right) = 0 \\ \partial_\alpha \mathbf{m}(z, \alpha_{\min}) = \partial_\alpha \mathbf{m}(z, \alpha_{\max}) = 0, \ z \in \mathbb{R} \end{cases} \tag{C54}$$

where

$$\overline{\alpha}(z) = \int_{\alpha_{\min}}^{\alpha_{\max}} \alpha\,\frac{\mathbf{m}(z, \alpha)}{\mathbf{M}(z)}\,d\alpha \ \ \text{and} \ \ \mathbf{M}(z) = \int_{\alpha_{\min}}^{\alpha_{\max}} \mathbf{m}(z, \alpha)\,d\alpha. \tag{C55}$$

The traveling wave profile satisfy the following boundary conditions

$$\begin{cases} \mathbf{P}(-\infty) = P^* \\ \mathbf{m}(-\infty, \alpha) = M^*\phi(\alpha) \end{cases} \text{and} \ \begin{cases} \mathbf{P}(\infty) = 0 \\ \mathbf{m}(\infty, \alpha) = 0 \end{cases} \text{for all } \alpha \in (\alpha_{\min}, \alpha_{\max}) \tag{C56}$$

where (P^*, M^*, ϕ) solves the stationnary problem (B11) and (B12)

We use the inside dynamics approach presented in the previous section B.5 and introduced by (Roques et al, 2012). We here label the individuals – hosts or symbionts – according to their initial position in the front, and they transmit their label to their offspring. Since they only differ by their label and their trait for the symbionts, each label $k \in \mathbb{N}$ corresponds to a neutral fraction of density v_p^k inside the host population $\mathbf{P}(x - ct)$ and v_m^k inside the symbiont population $\mathbf{M}(x - ct, \alpha)$. Initially, the fractions satisfy

$$\mathbf{P}(z) = \sum_{k \in \mathbb{N}} v_p^k(0, z) \quad \text{and} \quad \mathbf{m}(z, \alpha) = \sum_{k \in \mathbb{N}} v_m^k(0, z, \alpha) \tag{C57}$$

Their dynamics in the moving frame of speed c, is described by

$$\partial_t v_p^k(t, z) = D_p \partial_z^2 v_p^k + c \partial_z v_p^k + v_p^k \left(\frac{Q \, \overline{\alpha}(z) \mathbf{M}(z)}{\mathbf{P}(z) + d} - \beta \, \mathbf{M}(z) - \mu_p \mathbf{P}(z) \right), \; z \in \mathbb{R}, \; t > 0 \tag{C58}$$

$$\partial_t v_m^k(t, z, \alpha) = D_m \partial_z^2 v_m^k + c \partial_z v_m^k + d_m \partial_\alpha^2 v_m^k + v_m^k \left(\beta \, \mathbf{P}(z) - \frac{\overline{\alpha}(z) \, \mathbf{P}(z)}{\mathbf{P}(z) + d} - \mu_m \mathbf{M}(z) \right), \; \alpha \in (\alpha_{\min}, \alpha_{\max}) \tag{C59}$$

$$\partial_\alpha v_m^k(z, \alpha_{\min}) = \partial_\alpha v_m^k(z, \alpha_{\max}) = 0, \; z \in \mathbb{R} \tag{C60}$$

Since we are interested in the effect of initial space location, we can integrate the equation in v_m^k and we obtain that the quantity $\Upsilon_m^k(z) = \int v_m^k(z, \alpha) d\alpha$ satisfies the following problem

$$\partial_t \Upsilon_m^k(t, z, \alpha) = D_m \partial_z^2 \Upsilon_m^k + c \partial_z \Upsilon_m^k + \Upsilon_m^k \left(\beta \, \mathbf{P}(z) - \frac{\overline{\alpha}(z) \, \mathbf{P}(z)}{\mathbf{P}(z) + d} - \mu_m \mathbf{M}(z) \right), \; \alpha \in (\alpha_{\min}, \alpha_{\max}) \tag{C61}$$

Formally, the solutions v_p^k and Υ_m^k converge uniformly in space toward $p_p^k \mathbf{P}(z)$ and $p_m^k \mathbf{M}(z)$ respectively, where the positive constant p_p^k and p_m^k satisfies

$$p_p^k = \frac{\displaystyle\int_{\mathbb{R}} v_p^k(0, z) e^{\frac{cz}{D_p}} \mathbf{P}(z) dz}{\displaystyle\int_{\mathbb{R}} e^{\frac{cz}{D_p}} \mathbf{P}^2(z) dz} \quad \text{and} \quad p_m^k = \frac{\displaystyle\int_{\mathbb{R}} v_m^k(0, z) e^{\frac{cz}{D_m}} \mathbf{M}(z) dz}{\displaystyle\int_{\mathbb{R}} e^{\frac{cz}{D_m}} \mathbf{M}^2(z) dz} \tag{C62}$$

As a result, the probability that individuals from respectively the host population of symbiont population, starting in location x, fix their label in the population is given respectively by $PF_p(x)$ and $PF_m(x)$, which are defined by

$$PF_p(x) = \frac{e^{\frac{cx}{D_p}} \mathbf{P}^2(x)}{\displaystyle\int_{\mathbb{R}} e^{\frac{cz}{D_p}} \mathbf{P}^2(z) dz} \quad \text{and} \quad PF_m(x) = \frac{e^{\frac{cx}{D_m}} \mathbf{M}^2(x) dz}{\displaystyle\int_{\mathbb{R}} e^{\frac{cz}{D_m}} \mathbf{M}^2(z) dz} \tag{C63}$$

Since the traveling profiles $\mathbf{P}$ and $\mathbf{M}$ decay at different rates (Fig. C5), the fixation probability attains is maximum at different location. In particular, we see that the

Fig. C5 Fixation probability of plant (dashed curves) and symbionts (plain curves) with respect to space location: red curves are numerical simulation from Eq. (C58)-(C61) and gray curves are numerical simulation from Eq. (C63) . We also plot the densities of plant (black dashed curve) and AMF symbiont densities (dots). The color of the dots represent the value of the mean mutualistic trait $\overline{\alpha}(x)$ at the specific location x (green parasitic symbiont to blue mutualistic symbionts). For the simulations, the parameters are: $d_m = 0.01$, $D_p = D_m = 0.1$, $Q = 6$, $\mu_p = \mu_m = 0.3$, $\beta = 0.72$ and $\alpha_{\min} = 0$ and $\alpha_{\max} = 5$.

plant at the leading edge have a higher probability of fixation than the symbiont at the same location. Thus plant at the leading edge mainly come from this edge, while the symbiont come more from the core of the population (Fig. C5).

References

Afkhami ME, McIntyre PJ, Strauss SY (2014) Mutualist-mediated effects on species' range limits across large geographic scales. Ecology letters 17(10):1265–1273

Amor DR, Montañez R, Duran-Nebreda S, et al (2017) Spatial dynamics of synthetic microbial mutualists and their parasites. PLOS Computational Biology 13(8):e1005689

Araújo MB, Rozenfeld A (2014) The geographic scaling of biotic interactions. Ecography 37(5):406–415

Archetti M, Scheuring I (2011) Coexistence of cooperation and defection in public goods games. Evolution 65(4):1140–1148

Archetti M, Scheuring I (2013) Trading public goods stabilizes interspecific mutualism. Journal of Theoretical Biology 318:58–67. https://doi.org/10.1016/j.jtbi.2012.10.022, URL https://www.sciencedirect.com/science/article/pii/S0022519312005619

Bachelot B, Lee CT (2018) Dynamic preferential allocation to arbuscular mycorrhizal fungi explains fungal succession and coexistence. Ecology 99(2):372–384

den Bakker HC, VanKuren NW, Morton JB, et al (2010) Clonality and Recombination in the Life History of an Asexual Arbuscular Mycorrhizal Fungus. Molecular Biology and Evolution 27(11):2474–2486. https://doi.org/10.1093/molbev/msq155, URL https://doi.org/10.1093/molbev/msq155

Benning JW, Moeller DA (2021) Microbes, mutualism, and range margins: testing the fitness consequences of soil microbial communities across and beyond a native plant's range. New Phytologist 229(5):2886–2900

Bever J (1999) Dynamics within mutualism and the maintenance of diversity: inference from a model of interguild frequency dependence. Ecology Letters 2(1):52–61. https://doi.org/10.1046/j.1461-0248.1999.21050.x

Bever JD (2015) Preferential allocation, physio-evolutionary feedbacks, and the stability and environmental patterns of mutualism between plants and their root symbionts. New Phytologist 205(4):1503–1514

Bever JD, Richardson SC, Lawrence BM, et al (2009) Preferential allocation to beneficial symbiont with spatial structure maintains mycorrhizal mutualism. Ecology letters 12(1):13–21

Boots M, Mealor M (2007) Local interactions select for lower pathogen infectivity. Science 315(5816):1284–1286. https://doi.org/10.1126/science.1137126

Bronstein JL (2015) Mutualism. Oxford University Press, USA

Brooker RW, Travis JM, Clark EJ, et al (2007) Modelling species' range shifts in a changing climate: the impacts of biotic interactions, dispersal distance and the rate of climate change. Journal of Theoretical Biology 245(1):59–65

Brown CD, Vellend M (2014) Non-climatic constraints on upper elevational plant range expansion under climate change. Proceedings of the Royal Society B: Biological Sciences 281(1794):20141779

Brown JKM, Tellier A (2011) Plant-parasite coevolution: Bridging the gap between genetics and ecology. Annual Review of Phytopathology 49(1):345–367. https://doi.org/10.1146/annurev-phyto-072910-095301

Cameron DD, Johnson I, Read DJ, et al (2008) Giving and receiving: measuring the carbon cost of mycorrhizas in the green orchid, goodyera repens. New Phytologist

180(1):176–184

Case TJ, Holt RD, McPeek MA, et al (2005) The community context of species' borders: ecological and evolutionary perspectives. Oikos 108(1):28–46

Chen IC, Hill JK, Ohlemüller R, et al (2011) Rapid range shifts of species associated with high levels of climate warming. Science 333(6045):1024–1026

Christian N, Bever JD (2018) Carbon allocation and competition maintain variation in plant root mutualisms. Ecology and Evolution 8(11):5792–5800

Chuang A, Peterson CR (2016) Expanding population edges: theories, traits, and trade-offs. Global change biology 22(2):494–512

Cloez B, Gabriel P (2020) On an irreducibility type condition for the ergodicity of nonconservative semigroups. Comptes Rendus Mathématique 358(6):733–742

Colautti RI, Ricciardi A, Grigorovich IA, et al (2004) Is invasion success explained by the enemy release hypothesis? Ecology letters 7(8):721–733

Davenport ER, Cusanovich DA, Michelini K, et al (2015) Genome-wide association studies of the human gut microbiota. PloS one 10(11):e0140301

Day T, Gandon S, Lion S, et al (2020) On the evolutionary epidemiology of sars-cov-2. Current Biology 30(15):R849–R857

Dickie IA, Bufford JL, Cobb RC, et al (2017) The emerging science of linked plant–fungal invasions. New Phytologist 215(4):1314–1332

Doebeli M, Knowlton N (1998) The evolution of interspecific mutualisms. Proceedings of the National Academy of Sciences 95(15):8676–8680

Drew GC, Stevens EJ, King KC (2021a) Microbial evolution and transitions along the parasite–mutualist continuum. Nature Reviews Microbiology 19:1–16. https://doi.org/10.1038/s41579-021-00550-7

Drew GC, Stevens EJ, King KC (2021b) Microbial evolution and transitions along the parasite–mutualist continuum. Nature Reviews Microbiology 19(10):623–638

Eldakar OT, Dlugos MJ, Holt GP, et al (2010) Population structure influences sexual conflict in wild populations of water striders. Behaviour 147(12):1615–1631. https://doi.org/10.1163/000579510X510520, URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3774177/

Engelmoer DJ, Behm JE, Toby Kiers E (2014) Intense competition between arbuscular mycorrhizal mutualists in an in vitro root microbiome negatively affects total fungal abundance. Molecular ecology 23(6):1584–1593

Erm P, Phillips BL (2020) Evolution transforms pushed waves into pulled waves. The American Naturalist 195(3):E87–E99

Ewald PW (1987) Transmission modes and evolution of the parasitism-mutualism continuum. Annals of the New York Academy of Sciences 503:295–306

Fagan WF, Lewis MA, Neubert MG, et al (2002) Invasion theory and biological control. Ecology letters 5(1):148–157

Ferdy JB, Godelle B (2005) Diversification of transmission modes and the evolution of mutualism. The American Naturalist 166(5):613–627

Fitzpatrick CR, Salas-González I, Conway JM, et al (2020) The plant microbiome: from ecology to reductionism and beyond. Annual Review of Microbiology 74:81–100

Fleming WH (1979) Equilibrium distributions of continuous polygenic traits. SIAM J Appl Math 36(1):148–168. https://doi.org/10.1137/0136014

Forien R, Garnier J, Patout F (2022) Ancestral lineages in mutation selection equilibria with moving optimum. Bull Math Biol 84(93)

Fournier A, Penone C, Pennino MG, et al (2019) Predicting future invaders and future invasions. Proceedings of the National Academy of Sciences 116(16):7905–7910

Fowler JC, Donald ML, Bronstein JL, et al (2023) The geographic footprint of mutualism: How mutualists influence species' range limits. Ecological Monographs 93(1):e1558

Frederickson ME (2013) Rethinking mutualism stability: cheaters and the evolution of sanctions. The Quarterly review of biology 88(4):269–295

Frenkel ES, Ribbeck K (2017) Salivary mucins promote the coexistence of competing oral bacterial species. The ISME Journal 11(5):1286–1290

Gandhi SR, Yurtsev EA, Korolev KS, et al (2016) Range expansions transition from pulled to pushed waves as growth becomes more cooperative in an experimental microbial population. Proceedings of the National Academy of Sciences 113(25):6922–6927

Godsoe W, Holland NJ, Cosner C, et al (2017) Interspecific interactions and range limits: Contrasts among interaction types. Theoretical Ecology 10:167–179

Gómez P, Paterson S, De Meester L, et al (2016) Local adaptation of a bacterium is as important as its presence in structuring a natural microbial community. Nature communications 7(1):12453

Hartnett D, Hetrick B, Wilson G, et al (1993) Mycorrhizal influence on intra-and interspecific neighbour interactions among co-occurring prairie grasses. Journal of

Ecology pp 787–795

van der Heijden MG, Walder F (2016) Reply to 'misconceptions on the application of biological market theory to the mycorrhizal symbiosis'. Nature Plants 2(5):1–1

Hochberg M, Gomulkiewicz R, Holt R, et al (2000) Weak sinks could cradle mutualistic symbioses-strong sources should harbour parasitic symbioses. Journal of Evolutionary biology 13(2):213–222

Hochberg ME, van Baalen M (2000) A geographical perspective of virulence. Evolutionary biology of host-parasite relationships: theory meets reality Elsevier pp 81–96

Hochberg ME, Rankin DJ, Taborsky M (2008) The coevolution of cooperation and dispersal in social groups and its implications for the emergence of multicellularity. BMC Evolutionary Biology 8(1):1–14. https://doi.org/10.1186/1471-2148-8-238

Hoeksema JD, Kummel M (2003) Ecological persistence of the plant-mycorrhizal mutualism: a hypothesis from species coexistence theory. The american naturalist 162(S4):S40–S50

Hou S, Wolinska KW, Hacquard S (2021) Microbiota-root-shoot-environment axis and stress tolerance in plants. Current Opinion in Plant Biology 62:102028

Hu J, Vandenkoornhuyse P, Khalfallah F, et al (2022) Ecological corridors homogenize plant root endospheric mycobiota. New Phytologist 237(4):1347–1362. https://doi.org/10.1111/nph.18606

Huang F, Peng S (2016) Intraspecific competitive ability declines towards the edge of the expanding range of the invasive vine mikania micrantha. Oecologia 181:115–123

Jones EI, Bronstein JL, Ferrière R (2012) The fundamental role of competition in the ecology and evolution of mutualisms. Annals of the New York Academy of Sciences 1256(1):66–88

Kiers ET, Duhamel M, Beesetty Y, et al (2011) Reciprocal rewards stabilize cooperation in the mycorrhizal symbiosis. science 333(6044):880–882

Koella JC (2000) The spatial spread of altruism versus the evolutionary response of egoists. Proceedings of the Royal Society of London Series B: Biological Sciences 267(1456):1979–1985. https://doi.org/10.1098/rspb.2000.1239

Koskella B, Bergelson J (2020) The study of host–microbiome (co) evolution across levels of selection. Philosophical Transactions of the Royal Society B 375(1808):20190604

Koskella B, Hall LJ, Metcalf CJE (2017) The microbiome beyond the horizon of ecological and evolutionary theory. Nature ecology & evolution 1(11):1606–1615

Kubisch A, Holt RD, Poethke HJ, et al (2014) Where am I and why? Synthesizing range biology and the eco-evolutionary dynamics of dispersal. Oikos 123(1):5–22

Lankau RA, Keymer DP (2016) Ectomycorrhizal fungal richness declines towards the host species' range edge. Molecular Ecology 25(13):3224–3241

Le Galliard JF, Ferriere R, Dieckmann U (2005) Adaptive evolution of social traits: origin, trajectories, and correlations of altruism and mobility. The American Naturalist 165(2):206–224. https://doi.org/10.1086/427090

LeClerc JE, Li B, Payne WL, et al (1996) High mutation frequencies among escherichia coli and salmonella pathogens. Science 274(5290):1208–1211

Ledru L, Garnier J, Rhor M, et al (2022) Mutualists construct the ecological conditions that trigger the transition from parasitism. Peer Community Journal 2:e41. https://doi.org/10.24072/pcjournal.139, URL https://peercommunityjournal.org/articles/10.24072/pcjournal.139/

Léotard G, Debout G, Dalecky A, et al (2009) Range expansion drives dispersal evolution in an equatorial three-species symbiosis. PLoS One 4(4):e5377

Li B, Weinberger HF, Lewis MA (2005) Spreading speeds as slowest wave speeds for cooperative systems. Mathematical Biosciences 196(1):82–98. https://doi.org/10.1016/j.mbs.2005.03.008

Lin D, Koskella B (2015) Friend and foe: factors influencing the movement of the bacterium helicobacter pylori along the parasitism–mutualism continuum. Evolutionary Applications 8(1):9–22

Mack KM (2012) Selective feedback between dispersal distance and the stability of mutualism. Oikos 121(3):442–448

Martignoni MM, Hart MM, Garnier J, et al (2020) Parasitism within mutualist guilds explains the maintenance of diversity in multi-species mutualisms. Theoretical Ecology 13:615–627

Miller ET, Svanbäck R, Bohannan BJ (2018) Microbiomes as metacommunities: understanding host-associated microbes through metacommunity ecology. Trends in Ecology & Evolution 33(12):926–935

Miller TE, Angert AL, Brown CD, et al (2020) Eco-evolutionary dynamics of range expansion. Ecology 101(10):e03139

Mitchell-Olds T, Willis J, Goldstein D (2007) Which evolutionary processes influence natural genetic variation for phenotypic traits? Nat Rev Genet 8:845–856. https://doi.org/10.1038/nrg2207

Momeni B, Waite AJ, Shou W (2013) Spatial self-organization favors heterotypic cooperation over cheating. Elife 2:e00960

Mony C, Uroy L, Khalfallah F, et al (2022) Landscape connectivity for the invisibles. Ecography 2022(8). https://doi.org/10.1111/ecog.06041

Muller-Parker G, D'elia CF, Cook CB (2015) Interactions between corals and their symbiotic algae. Coral reefs in the Anthropocene pp 99–116

Mullon C, Keller L, Lehmann L (2018) Social polymorphism is favoured by the co-evolution of dispersal with social behaviour. Nature ecology & evolution 2(1):132–140. https://doi.org/10.1038/s41559-017-0397-y

Neuhauser C, Fargione JE (2004) A mutualism–parasitism continuum model and its application to plant–mycorrhizae interactions. Ecological modelling 177(3-4):337–352

Nomikou M, Janssen A, Sabelis MW (2003) Herbivore host plant selection: white-fly learns to avoid host plants that harbour predators of her offspring. Oecologia 136:484–488

Oliver A, Cantón R, Campo P, et al (2000) High frequency of hypermutable pseudomonas aeruginosa in cystic fibrosis lung infection. Science 288(5469):1251–1253

Palmer TM, Stanton ML, Young TP (2003) Competition and coexistence: exploring mechanisms that restrict and maintain diversity within mutualist guilds. the american naturalist 162(S4):S63–S79

Pande S, Kaftan F, Lang S, et al (2016) Privatization of cooperative benefits stabilizes mutualistic cross-feeding interactions in spatially structured environments. The ISME Journal 10(6):1413–1423

Paquette A, Hargreaves AL (2021) Biotic interactions are more often important at species' warm versus cool range edges. Ecology Letters 24(11):2427–2438

Parker MP (1995) Plant fitness variation caused by different mutualist genotypes. Ecology 76(5):1525–1535. https://doi.org/10.2307/1938154

Phillips BL (2009) The evolution of growth rates on an expanding range edge. Biology Letters 5(6):802–804

Phillips BL, Kelehear C, Pizzatto L, et al (2010) Parasites and pathogens lag behind their host during periods of host range advance. Ecology 91(3):872–881

Purcell J, Brelsford A, Avilés L (2012) Co-evolution between sociality and dispersal: the role of synergistic cooperative benefits. Journal of Theoretical Biology 312:44–54. https://doi.org/10.1016/j.jtbi.2012.07.016

Restif O (2009) Evolutionary epidemiology 20 years on: challenges and prospects. Infection, Genetics and Evolution 9(1):108–123

Rizaludin MS, Stopnisek N, Raaijmakers JM, et al (2021) The chemistry of stress: understanding the 'cry for help'of plant roots. Metabolites 11(6):357

Robinson CJ, Bohannan BJ, Young VB (2010) From structure to function: the ecology of host-associated microbial communities. Microbiology and Molecular Biology Reviews 74(3):453–476

Rodríguez Amor D, Dal Bello M (2019) Bottom-up approaches to synthetic cooperation in microbial communities. Life 9(1):22

Rogalski MA, Stewart Merrill T, Gowler CD, et al (2021) Context-dependent host-symbiont interactions: Shifts along the parasitism-mutualism continuum. The American Naturalist 198(5):563–575

Roques L, Garnier J, Hamel F, et al (2012) Allee effect promotes diversity in traveling waves of colonization. Proc Natl Acad Sci USA 109:8828–8833

Sachs JL, Mueller UG, Wilcox TP, et al (2004) The evolution of cooperation. The Quarterly review of biology 79(2):135–160

Sachs JL, Skophammer RG, Regus JU (2011) Evolutionary transitions in bacterial symbiosis. Proceedings of the National Academy of Sciences 108(supplement_2):10800–10807

Schuman MC, Allmann S, Baldwin IT (2015) Plant defense phenotypes determine the consequences of volatile emission for individuals and neighbors. Elife 4:e04490

Schwartz MW, Hoeksema JD (1998) Specialization and resource trade: biological markets as a model of mutualisms. Ecology 79(3):1029–1038

Seifert EK, Bever JD, Maron JL (2009) Evidence for the evolution of reduced mycorrhizal dependence during plant invasion. Ecology 90(4):1055–1062

Shade A, Jacques MA, Barret M (2017) Ecological patterns of seed microbiome diversity, transmission, and assembly. Current opinion in microbiology 37:15–22

Shaw AK (2022) How to outrun your parasites (or mutualists): symbiont transmission mode is key. Oikos 2022(11):e09374

Simon B, Fletcher JA, Doebeli M (2013) Towards a general theory of group selection. Evolution 67(6):1561–1572. https://doi.org/10.1111/j.1558-5646.2012.01835.x

Smith SE, Read DJ (2010) Mycorrhizal symbiosis. Academic press

Spence AR, Tingley MW (2020) The challenge of novel abiotic conditions for species undergoing climate-induced range shifts. Ecography 43(11):1571–1590

Stanton-Geddes J, Anderson CG (2011) Does a facultative mutualism limit species range expansion? Oecologia 167:149–155

Stephan P, Bramon Mora B, Alexander JM (2021) Positive species interactions shape species' range limits. Oikos 130(10):1611–1625

Taylor CM, Hastings A (2004) Finding optimal control strategies for invasive species: a density-structured model for spartina alterniflora. Journal of Applied Ecology 41(6):1049–1057. https://doi.org/10.1111/j.0021-8901.2004.00979.x

Thompson JN (2005) The geographic mosaic of coevolution (interspecific interactions). University of Chicago Press

Thonar C, Frossard E, Šmilauer P, et al (2014) Competition and facilitation in synthetic communities of arbuscular mycorrhizal fungi. Molecular Ecology 23(3):733–746

Torchin ME, Lafferty KD, Dobson AP, et al (2003) Introduced species and their missing parasites. Nature 421(6923):628–630

Trindade S, Perfeito L, Gordo I (2010) Rate and effects of spontaneous mutations that affect fitness in mutator escherichia coli. Philosophical Transactions of the Royal Society B: Biological Sciences 365(1544):1177–1186

Turchin P (1998) Quantitative analysis of movement : measuring and modeling population redistribution in animals and plants

Urban MC, Phillips BL, Skelly DK, et al (2007) The cane toad's (*Chaunus [Bufo] marinus*) increasing ability to invade australia is revealed by a dynamically updated range model. Proceedings of the Royal Society B: Biological Sciences 274(1616):1413–1419. https://doi.org/10.1098/rspb.2007.0114

Van Baalen M, Rand DA (1998) The unit of selection in viscous populations and the evolution of altruism. Journal of theoretical biology 193(4):631–648. https://doi.org/10.1006/jtbi.1998.0730

Van Dyken JD, Müller MJI, Mack KM, et al (2013) Spatial population expansion promotes the evolution of cooperation in an experimental prisoner's dilemma. Current Biology 23(10):919–923

Vandenkoornhuyse P, Leyval C, Bonnin I (2001) High genetic diversity in arbuscular mycorrhizal fungi: evidence for recombination events. Heredity 87. https://doi.org/10.1046/j.1365-2540.2001.00941.x, URL https://doi.org/10.1046/j.1365-2540.2001.00941.x

Vandenkoornhuyse P, Quaiser A, Duhamel M, et al (2015) The importance of the microbiome of the plant holobiont. New Phytologist 206(4):1196–1206

Vittecoq M, Djiéto-Lordon C, McKey D, et al (2012) Range expansion induces variation in a behavioural trait in an ant–plant mutualism. Acta oecologica 38:84–88

Walder F, Van Der Heijden MG (2015) Regulation of resource exchange in the arbuscular mycorrhizal symbiosis. Nature plants 1(11):1–7

Walder F, Niemann H, Natarajan M, et al (2012) Mycorrhizal networks: common goods of plants shared under unequal terms of trade. Plant physiology 159(2):789–797

Wan JS, Fazlioglu F, Bonser SP (2016) Populations evolving towards failure: costs of adaptation under competition at the range edge of an invasive perennial plant. Plant Ecology & Diversity 9(4):349–358

Wier AM, Nyholm SV, Mandel MJ, et al (2010) Transcriptional patterns in both host and bacterium underlie a daily rhythm of anatomical and metabolic change in a beneficial symbiosis. Proceedings of the National Academy of Sciences 107(5):2259–2264

Wilkinson DM, Sherratt TN (2001) Horizontally acquired mutualisms, an unsolved problem in ecology? Oikos 92(2):377–384

Willing CE, Pierroz G, Guzman A, et al (2021) Keep your friends close: Host compartmentalisation of microbial communities facilitates decoupling from effects of habitat fragmentation. Ecology Letters 24(12):2674–2686

Wilson DS, Sober E (1989) Reviving the superorganism. Journal of theoretical Biology 136(3):337–356. https://doi.org/10.1016/s0022-5193(89)80169-9

Yuan J, Zhang N, Huang Q, et al (2015) Organic acids from root exudates of banana help root colonization of pgpr strain bacillus amyloliquefaciens njn-6. Scientific reports 5(1):1–8

Zhang H, Ziegler W, Han X, et al (2015) Plant carbon limitation does not reduce nitrogen transfer from arbuscular mycorrhizal fungi to plantago lanceolata. Plant and Soil 396:369–380