Metagenomics uncovers dietary adaptations for chitin digestion in the gut microbiota of convergent myrmecophagous mammals

Sophie Teullet[a,#], Marie-Ka Tilak[a], Amandine Magdeleine[a], Roxane Schaub[b,c], Nora M. Weyer[d], Wendy Panaino[d,e], Andrea Fuller[d], W. J. Loughry[f], Nico L. Avenant[g], Benoit de Thoisy[h,i], Guillaume Borrel[j] and Frédéric Delsuc[a]

[a] Institut des Sciences de l'Evolution de Montpellier (ISEM), Univ Montpellier, CNRS, IRD, Montpellier, France

[b] CIC AG/Inserm 1424, Centre Hospitalier de Cayenne Andrée Rosemon, Cayenne, French Guiana

[c] Tropical Biome and immunopathology, Université de Guyane, Labex CEBA, DFR Santé, Cayenne, French Guiana

[d] Brain Function Research Group, School of Physiology, University of the Witwatersrand, Johannesburg, South Africa

[e] Centre for African Ecology, School of Animals, Plant, and Environmental Sciences, University of the Witwatersrand, Johannesburg, South Africa

[f] Department of Biology, Valdosta State University, Valdosta, GA, USA

[g] National Museum and Centre for Environmental Management, University of the Free State, Bloemfontein, South Africa

[h] Institut Pasteur de la Guyane, Cayenne, French Guiana, France

[i] Kwata NGO, Cayenne, French Guiana, France

[j] Institut Pasteur, Université Paris Cité, UMR CNRS 6047, Evolutionary Biology of the Microbial Cell, Paris, France

Running title: Ant-eating mammal metagenomics

[#]Address correspondence to Sophie Teullet, sophie.teullet@umontpellier.fr

Abstract

In mammals, myrmecophagy (ant and termite consumption) represents a striking example of dietary convergence. This trait evolved independently at least five times in placentals with myrmecophagous species comprising aardvarks, anteaters, some armadillos, pangolins, and aardwolves. The gut microbiome plays an important role in dietary adaptation, and previous analyses of 16S rRNA metabarcoding data have revealed convergence in the composition of the gut microbiota among some myrmecophagous species. However, the functions performed by these gut bacterial symbionts and their potential role in the digestion of prey chitinous exoskeletons remain open questions. Using long- and short-read sequencing of fecal samples, we generated 29 gut metagenomes from nine myrmecophagous and closely related insectivorous species sampled in French Guiana, South Africa, and the USA. From these, we reconstructed 314 high-quality bacterial genome bins of which 132 carried chitinase genes, highlighting their potential role in insect prey digestion. These chitinolytic bacteria belonged mainly to the family Lachnospiraceae, and some were likely convergently recruited in the different myrmecophagous species as they were detected in several host orders (i.e., *Enterococcus faecalis*, *Blautia* sp), suggesting that they could be directly involved in the adaptation to myrmecophagy. Others were found to be more host-specific, possibly reflecting phylogenetic constraints and environmental influences. Overall, our results highlight the potential role of the gut microbiome in chitin digestion in myrmecophagous mammals and provide the basis for future comparative studies performed at the mammalian scale to further unravel the mechanisms underlying the convergent adaptation to myrmecophagy.

Importance

Myrmecophagous mammals are specialized in the consumption of ants and/or termites. They do not share a direct common ancestor and evolved convergently in five distinct placental orders raising questions about the underlying adaptive mechanisms involved and the relative contribution of natural selection and phylogenetic constraints. Understanding how these species digest their prey can help answer these questions. More specifically, the role of their gut microbial symbionts in the digestion of the insect chitinous exoskeleton has not been investigated in all myrmecophagous orders. We generated 29 new gut metagenomes from nine myrmecophagous species to reconstruct more than 300 bacterial genomes in which we identified chitin-degrading enzymes. Studying the distribution of these chitinolytic bacteria among hosts revealed both shared and specific bacteria between ant-eating species. Overall, our results highlight the potential role of gut symbionts in the convergent dietary adaptation of myrmecophagous mammals and the evolutionary mechanisms shaping their gut microbiota.

Keywords

convergent evolution; myrmecophagy; mammals; gut microbiota; chitinases; metagenomics;
genome assembly.

Introduction

In mammals, the gut microbiota has played a major role in dietary diversification, enabling
transitions to novel carbon sources (1, 2). Several factors such as host diet, physiology,
genetics, and phylogeny, but also environmental factors, shape the taxonomic composition
and functional structure of the gut microbiota (3). Many studies have focused on convergent
dietary adaptations and the effects of host diet, investigating how host phylogeny shapes the
gut microbiota (4–10). A striking example of convergent dietary specialization is the
adaptation to myrmecophagy in placental mammals. Myrmecophagous species are
characterized by a diet composed of at least 90% of ants and/or termites (11). A total of 22
placental species evolved toward this diet and are found in five independent orders:
Tubulidentata (aardvarks), Pilosa (anteaters), Cingulata (tolypeutine armadillos), Pholidota
(pangolins), and Carnivora (aardwolves) (12–14). Their divergent evolutionary histories raise
the question of how these species convergently adapted to the myrmecophagous diet and
whether similar adaptive mechanisms were involved between the different species.
Myrmecophagous species need to digest the chitinous exoskeleton of their prey to extract
nutrients. Mammals carry chitinase genes in their genomes, which participate in chitin
digestion (15–18), but their gut microbiota might also have played an important role in
facilitating prey digestion (19–21).

92   Indeed, chitinolytic bacteria are present in a diversity of environments (i.e., soils,
sediments, aquatic environments) where they ensure chitin degradation and play an
important role in the carbon cycle (22–25). They also have been identified in the digestive
tract of mammals where they could participate in prey digestion (26–30). Taxonomic analyses
based on 16S rRNA sequences have highlighted similarities in the gut microbiota composition
of myrmecophagous species (19) compared to their non-myrmecophagous sister species. For
instance, genera such as *Blautia* (Lachnospiraceae), *Streptococcus* (Streptococcaceae),
Peptococcus (Peptococcaceae), or *Eubacterium* (Lachnospiraceae) were found to be
significantly more abundant in the gut microbiota of myrmecophagous species than in their
sister species. Focusing on the gut archaeome of placentals, *Methanobrevibacter* has been
found almost only in myrmecophagous species (10). Yet, little is known regarding the role
these symbionts play in the digestion of the chitinous exoskeleton of their prey.

Microbial chitinases and N-acetylglucosaminidases are carbohydrate-active enzymes
(CAZymes) (31) which ensure the degradation of chitin into chitosan and are mostly found in
the Glycosyl Hydrolases (GH) families 18, 19, and 20 (23, 32). Since bacterial chitinases mainly
belong to the GH18 family (32–35), they represent relevant candidates to assess the potential
of the gut microbiota to digest chitin in our focal species. The GH18 enzyme family comprises

chitinases and chitin-binding proteins and within this family, bacterial chitinases are classified
into three subfamilies (A, B, and C) based on sequence homology (32–34). Symbiotic chitin-
degrading bacteria have been identified, as well as their chitinase genes, in the Malayan
pangolin and the giant anteater (20, 21) and in other mammals having a chitin-rich diet (36,
37). This suggests that other myrmecophagous species might also carry chitinolytic gut
bacteria that could participate in prey digestion. In the specific example of dietary
convergence, one question resides in understanding whether these potential microbial
mechanisms of chitin exoskeleton digestion have converged among the five different
placental orders of myrmecophagous mammals. The different myrmecophagous mammal
lineages diverged millions of years ago and evolved in diverse environments. Chitinolytic
microbes might thus have been independently recruited to ensure chitin digestion, but
whether similar bacteria carrying similar functions are involved still needs further
investigation. Assessing whether chitinolytic bacteria present in myrmecophagous mammals
are taxonomically and functionally similar will shed light on their origin.

To investigate the chitin-degrading potential of the gut microbiota of
myrmecophagous mammals, we took advantage of recent advances in metagenomics to
reconstruct high-quality bacterial genomes and identify Glycoside Hydrolases 18 (GH18) as
this enzyme family comprises most bacterial chitinases. By combining Nanopore long-read
and Illumina short-read shotgun metagenomic sequencing, we generated 29 new gut
metagenomes from field-collected fecal samples of nine myrmecophagous and insectivorous
species representatives of the five myrmecophagous placental orders. From these, we
reconstructed 314 high-quality bacterial genomes harboring a diversity of GH18 chitinases.
Identification of GH18 sequences in the same bacterial genomes revealed a potential role of
gut symbionts in prey digestion through putative complex metabolic pathways. Both
generalist and host-specific bacteria were identified in the different myrmecophagous
species, potentially reflecting the divergent evolutionary histories of their hosts, and raising
questions about the evolutionary forces (historical contingency and determinism) at play in
shaping the gut microbiome of convergently evolved myrmecophagous mammals.

Results

Potential role of the gut microbiota in chitin digestion

Combining long-read and short-read assemblies, we were able to reconstruct 314
dereplicated high-quality genome bins (156 and 158 from each dataset respectively; see Table
S1F) and highlighted the usefulness of using the two methods (see supplementary results part
1 available via Zenodo). The following analyses were done on this set of high-quality selected
bins to assess the presence of chitinolytic bacteria in the gut microbiota of myrmecophagous
mammals.

All selected bins were taxonomically assigned to bacterial genomes and their
phylogeny is presented in Fig. S1. Overall, 58.3 % of the bins (n = 183) were not assigned at

the species level, 29.3% (n = 92) at the genus level, and 6.7% (n = 21) at the family level.
Taxonomically assigned bins belonged mainly to the Lachnospiraceae (n = 63),
Burkholderiaceae (n = 24), Acutalibacteraceae (n = 18), Ruminococcaceae (n = 12), and
Bacteroidaceae (n = 11) families (Table S1F). The 314 selected bins were integrated into a
phylogeny of prokaryote reference genomes to confirm these results (Fig. 1A). All selected
bins were well placed in this phylogeny, suggesting they are similar to known bacterial
genomes and not distantly related, even for genome bins for which taxonomic assignment
failed. Some of the reconstructed bacterial genome bins clustered together within the
Firmicutes (Fig. 1B), Bacteroidetes, and Proteobacteria (Figs. S2 A and B), in clades not
including reference genomes, suggesting that these genome bins could be specific to
myrmecophagous species (here called myrmecophagous-specific clades). Within the
Firmicutes, six clades containing more than two bins reconstructed from myrmecophagous
samples can be defined within the Lachnospiraceae family with two clades including
respectively 12 and 11 genome bins (Fig. 1B). In the first clade, six bins had no taxonomic
assignment below the family level and could represent bacterial genomes not yet described,
two belonged to the *Acetatifactor* genus, and four to the *CAG-510* genus. In the other clade,
four bins were not taxonomically assigned below the family level, three bins belonged to the
CAG-590 genus, one was assigned to the *CAG-127* genus and two were assigned at the species
level (*Frisingiococcus caecimuris* and *Acetivibrio ethanolgignens*) (Fig. 1B). These bins were all
reconstructed from xenarthran gut metagenomes (Fig. 1B). Within the Acutalibacteraceae
family, one clade contained 18 genome bins reconstructed mostly from xenarthran gut
metagenomes and the three aardvark samples (Fig. 1B). Eleven had no taxonomic assignment
below the family level, four belonged to the *Eubacterium* genus, and three to the *UBA1227*,
UBA1691, and *UBA6857* genera (Fig. 1B). Other clades containing only genome bins from
myrmecophagous species were found within the Ruminococcaceae, Oscillospiraceae,
Anaerovoracaceae, and Erysipelotrichaceae families.

Kingdom
- Archaea
- Eukaryota
- Bacteria

Phyla
- Actinobacteriota
- Bacteroidota
- Campylobacter
- Desulfobacterota
- Firmicutes
- Fusobacteriota
- Gemmatimonada
- None
- Proteobacteria
- Verrucomicrobiota

Family
- Acutalibacteraceae
- Anaeroboracaceae
- Clostridiaceae
- Eggerthellaceae
- Erysipelotrichaceae
- Fusobacteriaceae
- Lachnospiraceae
- None
- Oscillospiraceae
- Other
- Ruminococcaceae

Fig 1 Phylogenetic position of the 314 high-quality selected bins reconstructed from 29 gut metagenomes of the nine focal myrmecophagous species within a reference prokaryotic phylogeny. A: Phylogeny of the 314 selected bins (red branches) with 2496 prokaryote reference genomes. Circles respectively indicate (from inner to outer circles): the bacterial

phyla and kingdom to which these genome bins were assigned based on the Genome
Taxonomy Database release 7 (38). Clades, where a subtree was defined, are highlighted in
blue for the Firmicutes (Fig. 1B), green for the Bacteroidetes, and pink for the Proteobacteria
(Figs. S2 A and B, respectively). **B**: Subtree within Fimircutes showing myrmecophagous-
specific clades (blue highlights; dark blue corresponds to the three clades mentioned in the
results, light blue to the other clades). The outer circle indicates the bacterial family to which
these genome bins were assigned based on the Genome Taxonomy Database. Bins' names of
the myrmecophagous-specific clades are indicated at leaves of the phylogenetic tree together
with the genus to which they were assigned to.

To understand whether these bacteria could potentially degrade chitin, these high-quality
genome bins were then used to search for GH18 enzymes. This resulted in 132 bins containing
at least one GH18 sequence (between 0 and 17 bins per sample, 4.93 on average). Having
complete genome bins enabled us to identify several GH18 genes in the same bin (between
0 and 17 per bin, 1.26 on average; Table S1F). In total, 394 GH18 sequences were identified
(Fig. 2) with 237 sequences presenting an active chitinolytic site (DXXDXDXE) (Fig. 2) and
distributed among 82 bins (here called chitinolytic bins). These chitinolytic sequences are
found in genome bins belonging mainly to the Lachnospiraceae (n = 183 sequences; e.g.,
Blautia, *Acetatifactor*, *Roseburia*, *Clostridium* genera), Acutalibacteraceae (n = 76; e.g.,
Eubacterium genus), and Ruminoccocaceae (n = 23; e.g., *Ruminococcus*,
Acetanaerobacterium genera) bacterial families (Fig. 2). Fifty-three sequences not presenting
an active site were found in a clade with sequences similar to Lysin Motif (LysM) domain-
containing proteins (Fig. 2), which is a 40 amino acid domain involved in peptidoglycan and
chitin-binding (39). Sixty-two were placed in a clade with sequences similar to src Homology-
3 (SH3) domain-containing proteins (Fig. 2), which is a 50 amino acid domain found in
intracellular and membrane proteins involved in the binding of ligands (40). These two clades
were used to root the tree. Finally, 278 GH18 sequences formed a clade with sequences
similar to known bacterial chitinases (Fig. 2). The majority had an active chitinolytic site. These
sequences were identified in bins reconstructed from different host species, representative
of the five myrmecophagous placental orders. They represent a diversity of chitinase genes,
as they are distributed in distinct clades (Fig. 2).
210

Fig 2 Phylogeny of the 394 GH18 sequences identified in 132 high-quality selected bins reconstructed from 29 gut metagenomes of the nine focal myrmecophagous species and relatives. Red branches indicate the 237 sequences having an active chitinolytic site (DXXDXDXE). Circles respectively indicate (from inner to outer circles): the bacterial family and phyla of the bin the sequence was retrieved from. Colored sequence names indicate the host species. Colored circles at certain nodes indicate enzymes to which sequences are similar when blasting them against the NCBI non-redundant protein database. Sequence names are indicated at leaves of the tree and begin with the genus to which the bin they were identified in was assigned to.

Distribution of chitinolytic bacteria among myrmecophagous mammals

Numbers of shared and specific selected bins in the nine host species were computed based on a detection threshold of 0.25 (the percentage of the reference covered by at least one read) of the selected bins across samples (Fig. 3, Table S2, and detection table available via Zenodo). According to this threshold, five selected bins were not considered to be detected in any sample. Between 6 and 124 selected bins were detected in each sample (52.62 on average) and 252 selected genomes were shared and detected in samples other than the ones they were reconstructed from (Fig. 3). Selected bins were detected in 4.86 different samples on average (between 0 and 18). Selected bins were shared between individuals of the same species but with some intra-specific variability (Fig. 3). Conversely, 57 bacterial genomes were detected in only one sample (Fig. 3, Table S2) and 194 bins were detected in more than one host species (between 0 and 7 host species; 2.73 on average), between closely related species

or distantly related species (Fig. 3, Table S2). On the other hand, 115 selected bins were
specific to a particular host species, meaning they were found in only one of the nine species
studied here (Fig. 3, Table S2). Selected bins were detected in 1.80 different host orders on
average (between 0 and 5) for a total of 179 bins detected in more than one host order.
Finally, 130 bins were detected in only one host order.
There were more shared selected bins between host species carrying GH18 genes than
shared bins with no GH18 gene (Fig. S3). Among the 194 bins shared between host species,
110 had at least one GH18 gene. These bins belonged to 15 different bacterial families, mainly
the Lachnospiraceae (n = 60; e.g., *Blautia*, *Acetatifactor*, *Roseburia* genera) and
Acutalibacteraceae (n = 10; e.g., *Eubacterium* genus) (Fig. S3). The 84 shared selected bins
without chitinases belonged to a more diverse range of bacterial families (n = 33), mainly
Burkholderiaceae (n = 8; e.g., *CAG-521*, *Sutterella*, *Bordetella* genera) and Bacteroidaceae (n
= 7; e.g., *Bacteroidetes*, *Prevotella* genera) (Fig. S3). On the contrary, among bins found in only
one host species, there were fewer chitinase-carrying bins. Among the 115 host-species
specific bins, 21 had at least one GH18 gene. They were found in 12 different bacterial families
(Fig. S3). The 94 specific bins with no chitinase genes belonged to 37 different bacterial
families, mainly Burkholderiaceae (n = 16) (Fig. S3). A similar pattern was observed at the host
order level (see supplementary results part 2 available via Zenodo).
Three high-quality selected bins carrying GH18 were shared between species
belonging to the five different myrmecophagous orders (Table S1F): one bin of *Bacteroides*
fragilis (mean absolute abundance across samples 2.39, see abundance table available via
Zenodo) and two bins of Enterobacteriaceae (mean absolute abundances across samples 0.58
and 3.66). Two of these three bins shared across orders were a bit more abundant, on
average, than the mean absolute abundance of selected bins across samples of 0.89 (the
minimal abundance of a bin across samples was 2.82e-6 and the maximal absolute abundance
was 198). Seven selected bins carrying GH18 were shared among four myrmecophagous
orders (Table S1F). Two bins belonged to *Enteroccocus faecalis* and were reconstructed from
pangolin samples, along with bins belonging to Lachnospiraceae bacteria (notably one from
Blautia sp., one from *Faecalimonas* sp. and one from an unknown genus), one from
Bacteroides sp. (Bacteroidaceae) and one from *Emergencia timonensis* (Anaerovoracaceae)
with mean absolute abundances across samples ranging from 0.21 (*E. timonensis*) to 4.74
(*Bacteroides sp*), the latter being found to be abundant in several samples (for example: one
D. novemcinctus (12.2), one *T. tetradactyla* (21.2), one *O. afer* (41.2)). One bin of *Enteroccus*
faecalis also had a mean absolute abundance (3.43) above the mean absolute abundance of
selected bins and was found to be abundant in several samples as well (one *D. sp. nov* FG
(33.8), three pangolin samples (10.8, 13.3 and 21.8), and one *T. tetradactyla* (15.3)).
At least one chitinolytic bin (a high-quality selected bin carrying at least one GH18
sequence with an active chitinolytic site) was detected in the gut microbiota of each host
species (Fig. 4, Table S2). Consistent with our previous results (Fig. 2) these bins mainly
belonged to the Firmicutes (Fig. 4). More chitinolytic bins were detected in gut metagenomes
of xenarthran species compared to the aardvark, the southern aardwolf, and the ground

pangolin, except for *Dasypus kappleri* for which we only had one sample (Fig. 4; Table S3). When compared to the total number of genome bins detected per species, xenarthran species had higher proportions of chitinolytic genome bins (Table S3).

Fig 3 Detection of the 314 high-quality bacterial genomes (lines) **in the 29 gut metagenomes** (columns) **of the nine focal species.** Each square indicates the detection of a genome bin in a sample as estimated by anvi'o v7 (41). Names of bins are indicated on the left with red indicating chitinolytic bins (Table S2). The names begin with the genus to which the bin was assigned to. Asterisks (*****) indicate bins detected in at least one soil sample (detection > 0.25) (Fig. S4, Table S2, and detection table available via Zenodo). Phylogenetic relationships of host species distinguished by different color strips are represented at the bottom of the graph. Columns on the right indicate (from left to right): the number of GH18 sequences identified in each bin (from 0 to 17), the bin's taxonomic phylum, class, order, and family. The phylogeny of the 314 selected bins inferred with PhyloPhlAn v3.0.58 (42) is also represented on the right of the graph (see Fig. S1). Silhouettes were downloaded from phylopic.org.

Fig 4 Distribution of chitinolytic selected bins (red links) **among the nine focal myrmecophagous species and relatives.** Phylogenies of the 314 high-quality selected bins (Fig. S1) and of the nine host species (downloaded from timetree.org) are represented respectively on the left and the right of the graph. Links illustrate, for each bin, in which host species the bin was detected (detection threshold > 0.25). Red links indicate bins in which at least one GH18 sequence with an active chitinolytic site (DXXDXDXE) was found (chitinolytic bins). The size of the circles at the tips of the host phylogeny is proportional to the number of samples (n = 1 for *D. kap*; n = 2 for *D. nov, C. uni* and *M. tri*; n = 3 for *T. tet* and *O. afe*; n = 4 for *D. sp. nov* FG; n = 6 for *P. cri* and *S. tem*). Bins' names are indicated at the tip of the bins' phylogeny and main bacterial phyla are indicated by colored vertical bars. This graph was done with the cophylo R package within the phytools suite (43). Silhouettes were downloaded from phylopic.org.

Discussion

Potential role of the gut microbiota in prey digestion in myrmecophagous mammals

The gut microbiota plays an important role in host digestion, which has led to several cases of gut microbiome convergence in distantly related species that share similar diets (5, 19, 44). In animals with a chitin-rich diet, symbiotic chitinolytic bacteria could be involved in prey

digestion, as they have been identified in species as diverse as nine-banded armadillos (45),
insectivorous bats (28), insectivorous monkeys (30), or crustacean-eating whales (46). Here,
we focused on reconstructing high-quality bacterial genome bins from gut metagenomes of
species representative of the five myrmecophagous orders and identified sequences
belonging to the main bacterial chitinases (i.e., GH18) (32, 33, 47). This allowed us to assess
the potential of chitin degradation of gut symbionts in myrmecophagous placentals and
better understand their convergent adaptation to this highly specialized diet.

Among the high-quality bacterial genome bins reconstructed, several clustered
together in myrmecophagous-specific clades. These clades were found mainly within the
Lachnospiraceae and Actualibacteraceae families within Firmicutes but also within the
Oscillospiraceae and Ruminococcaceae. Several of these genomes had no taxonomic
assignment below the family level, suggesting they could represent bacterial taxa not
described yet and could be linked with the adaptation to myrmecophagy. Most of these
bacterial taxa (i.e., Lachnospiraceae like *Blautia* or *Roseburia* genera or Ruminococcaceae)
were previously found to be significantly more abundant in the gut microbiota of the
aardwolf, giant anteater, and southern tamandua when compared to their non-
myrmecophagous sister-species based on 16S rRNA metabarcoding data (19), which suggests
that their presence may be associated with dietary adaptations. Besides, chitin-degrading
metabolic pathways were identified in the gut metagenome of the Malayan pangolin (*Manis
javanica*) as well as genes encoding chitinases and chitin-binding proteins belonging to the
GH18 and GH19 enzyme families (20). This allowed the identification of chitinolytic symbionts
such as *Enterococcus faecalis* (Enterococcaceae), *Clostridium paraputrificum* (Clostridiaceae),
and *Bacteroides fragilis* (Bacteroidaceae) (20), whose genomes were also reconstructed here
from Temminck's pangolin (*Smutsia temminckii*) gut metagenomes, confirming their probable
role in prey digestion in pangolins.

Numerous GH18 genes were identified within the high-quality bacterial genomes we
reconstructed here. We were able to identify 132 genome bins carrying at least one GH18
gene among which 83 had at least one GH18 sequence with an active chitinolytic site
(DXXDXDXE) (48, 49), here called chitinolytic bins, and representing bacteria that might thus
play a role in insect prey digestion. Among genome bins found in the three myrmecophagous-
specific clades mentioned previously, almost all had at least one GH18 sequence. This further
suggests that these genome bins could represent bacterial taxa playing a significant role in
chitin digestion in myrmecophagous species. Studying the distribution of these bacteria in
other insectivorous mammals will allow further understanding their role in the adaptation
toward these specialized diets.

In addition, some of the bacterial taxa we identified here as being potentially involved
in chitin digestion in myrmecophagous species belonged to or were closely related to
bacterial taxa known for their chitinolytic properties. For example, *Clostridium* species are
known for their chitinolytic activity (50–52). Chitinases have also been identified and studied
in *Enterococcus faecalis* (53, 54). More generally, Lachnospiraceae, for which we
reconstructed the most genomes, are known to degrade complex polysaccharides (55) and

our results suggest they could be involved in chitin hydrolysis. Moreover, several genomes of
Ruminococcaceae bacteria reconstructed here encode a GH18. Within this family,
Ruminococcus species have been suggested to play a role in chitin digestion in the gut
microbiota of insectivorous mammals (56). Several GH18 sequences were found in the same
bacterial genomes, suggesting that complex chitin-degrading pathways are present in the gut
microbiota of the myrmecophagous species, consistent with previous analyses of the
chitinolytic properties of certain bacteria. Indeed, several enzymes are involved in the
different steps of chitin hydrolysis (23). For example, two types of chitin-binding and chitinase
enzymes have been described in *E. faecalis* (54). This bacteria also carries endo-β-N-
acetylglucosaminidases (57) which are part of the chitin hydrolysis pathway and degrade
chitin into chitosan (23). This further highlights the complex chitinolytic machinery of these
bacteria.

Finally, some sequences do not have the ability to hydrolyze chitin because no active
site was identified, but could still bind chitin. For example, the Lysin-motif (LysM) domain of
certain proteins can bind peptidoglycan (39). It is present in certain eukaryotes chitinases (i.e.,
algae, nematodes) and participates in the recognition of symbiotic rhizobial bacteria by
leguminous plants and is thus thought to bind chitin (39, 58). The SH3 domain is also known
to have binding properties and is involved in protein interactions (40). Therefore, sequences
similar to LysM and SH3 domain-containing proteins could still be part of the chitin hydrolysis
process by influencing molecular interactions. In addition, enzymes carrying the
carbohydrate-binding module 37 (CBM37), which is known to have binding properties notably
to chitin, could also be involved in the chitin degrading process. The presence of other
enzymes potentially involved in insect digestion could be investigated, such as trehalases,
enzymes that break down trehalose, a sugar found in insect blood. The gut microbiota of the
Malayan pangolin and giant anteater were found to be significantly enriched in these enzymes
when compared to non-myrmecophagous species (21). The gut microbiota also plays a role
in the detoxification of ingested compounds (26, 44), such as insect toxins. For example,
metabolic pathway analyses have revealed the potential of microbial symbionts to detoxify
formic acid in some myrmecophagous species (21).

Shared and species-specific gut bacteria among myrmecophagous mammals

Morphological and genomic comparative studies have shown that the convergent adaptation
to myrmecophagy in mammals involved different mechanisms between the different species
and that phylogenetic constraints played a central role (15, 18, 59–61). As diet is one of the
main factors shaping the evolution of the gut microbiota (e.g., (5)), this raises the question of
whether the same bacterial symbionts with similar functions were convergently recruited
between the different ant-eating species, or whether it was different bacteria with similar
functions (functional convergence).

Analysis of the distribution of selected genome bins across samples revealed both
shared and host-specific bacteria among myrmecophagous species. Selected genomes were
mainly shared between closely related species and mainly among Xenarthra but less so

between distantly related orders such as pangolins (Pholidota) and anteaters (Pilosa),
highlighting the influence of host evolutionary history. In mammals, phylosymbiosis is
particularly strong (9, 62) in part because of mammalian-specific traits such as viviparity or
parental care (63) but also limited microbial dispersal abilities (64) facilitating the vertical
transmission of microbes, which may explain these results. Some bacterial genomes were
shared between distantly related host species and those shared across more than one order
mainly belong to the family Lachnospiraceae. One selected genome of *Bacteroides fragilis* and
two of Enterobacteriaceae were shared between species belonging to the five different
myrmecophagous orders and carry GH18. Similarly, seven selected genome bins were shared
among four host orders and belong to *Enteroccocus faecalis*, the Lachnospiraceae (i.e., *Blautia*
sp., *Faecalimonas* sp.), *Bacteroides* sp. (Bacteroidaceae) and *Emergencia timonensis*
(Anaerovoracaceae). These shared high-quality selected genome bins carrying GH18 may thus
participate in the adaptation to myrmecophagy and highlight the influence of the host diet,
consistent with previous studies showing that it is an important factor in shaping the
mammalian gut microbiota (e.g., (1, 4, 5, 8)). Each host species appears to carry chitinolytic
bacteria in its gut microbiota, confirming their potential role in adapting to an insect-based
diet. However, differences in the distribution of these chitinolytic selected bins could highlight
divergent microbiota adaptations, especially in xenarthran species, which carry more
chitinolytic bacteria than aardvark, ground pangolin, and southern aardwolf. Some bins were
found to be abundant in some samples raising questions on how the gut microbiota is
involved in prey digestion. Having few highly abundant chitinolytic bacterial species producing
large amounts of chitinases might be sufficient for a specific host to digest its prey, whereas
having diverse chitinolytic bacterial species might have been selected for in other host
species.

The distribution of high-quality selected genome bins across samples and species
revealed how shared and host-specific bacteria may reflect the divergent evolutionary
histories of host species and the effect of adapting to a similar diet despite phylogenetic
constraints. This relates to the evolution of the mammalian gut microbiota. From a
mammalian ancestor that probably had an insectivore-like gut microbiota (62), its
composition might have changed as placentals diversified and occupied new niches,
constraining its composition in certain species. Chitinolytic bacteria shared by all
myrmecophagous orders may represent ancient bacterial lineages inherited from a common
ancestor, while bacteria found in only certain host species may reflect more recent
adaptations with bacteria acquired in specific lineages.

The environment and biogeography also influence host-associated microbial
communities, which could explain the distribution pattern of selected bins across samples,
with large differences observed between South Africa and South America. Anteaters and
armadillos (Xernarthra) diverged anciently from aardvarks, pangolins, and aardwolves (~80
Ma) (14), and these lineages have evolved in very distinct environments and biogeographic
contexts for most of their evolutionary history. Soil samples from the fecal sampling sites have
been collected for the South African samples. Some selected bins reconstructed from gut

metagenomes of the ground pangolin, southern aardwolf, and aardvark belonged to bacterial
taxa that were indeed not expected to be found in the gut but rather in the environment (e.g.,
Sphingobacteriaceae, Xanthomonadaceae, Burkholdericaeae). The distribution of selected
genome bins reconstructed from gut metagenomes of these species was examined in soil
samples and showed that 91 selected bins were also detected in at least one soil sample
(detection > 0.25; Fig. S4, Table S2). For example, among the Proteobacteria selected genome
bins found in myrmecophagous-specific clades (Fig. S2B), some belonged to Burkholderiaceae
bacteria and were detected in at least one soil sample. This may not necessarily reflect
environmental contamination, as the center of the feces was specifically sampled to minimize
contamination. Rather, it may reflect the fact that these species ingest soil and thus
environmental microbes while foraging (11). This could be beneficial to the host (65), as it
may compensate for the lack of mastication and/or help to deal with toxins that might be
present in prey (11). Environmental acquisition of microbes has also been suggested in lemurs
(66) and wild echidnas (67), species that ingest soil during foraging and in which soil bacteria
have been identified in their gut microbiota. These environmental bacteria could therefore
represent transient bacteria in the gut, reflecting what is ingested by the host. They could also
represent resident bacteria that have been recruited from the environment into the gut
microbiota of myrmecophagous species. Indeed, some beneficial microorganisms could be
acquired from soil microbiomes (68), which could allow the host to better adapt to its
environment through horizontal gene transfer of beneficial genes (69). Moreover, it has also
been suggested that environmental bacteria could be a source of chitinase genes to digest
prey (22). Thus, their recruitment into the gut microbiota may have been selected to
participate in the host digestion and further contribute to the adaptation to its environment.
This would highlight the potential influence of the environment on the gut microbiota of
myrmecophagous mammals.

Role of the holobiont in prey digestion in myrmecophagous mammals

The influence of the microbiota on host evolution and their coevolution is increasingly
recognized as studies of the diverse microbiota of captive and wild animals proliferate. Many
scientists now recognize the term holobiont to describe the host and all its associated
symbionts, which would be the unit of natural selection as defined by the hologenome theory
of evolution (70). Integrative studies (e.g., (21, 71)) and initiatives such as the "Earth
Hologenome Initiative'' (http://www.earthhologenome.org/) to study hologenomic
adaptations are now becoming more common. In the specific case of myrmecophagous
mammals, a hologenomic approach would help to better understand the adaptive
mechanisms involved. Indeed, if chitinolytic bacteria can digest chitin, prey digestion can also
be ensured by host-produced chitinases. In mammals, from a placental ancestor that probably
carried five functional chitinase paralogs (*CHIA*), some of these paralogs were subsequently
lost during placental diversification in non-insectivorous species, leading to a positive
correlation between the number of functional chitinase paralogs and the proportion of
invertebrates in the diet (15); a correlation also observed in primates (72). Among

myrmecophagous species, this chitinase gene repertoire has evolved differently to ensure
chitin digestion, reflecting phylogenetic constraints (15, 18). Thus, both endogenous and
microbial chitinases may be involved in prey digestion, raising the question of the relative
contribution of the host and its symbionts in providing the same function. For example,
myrmecophagous mammals with only one functional *CHIA* paralog (*CHIA5*), such as
aardwolves and pangolins, may compensate for this by overexpressing their only functional
paralog or by relying more on their chitinolytic symbionts to digest their prey. In pangolins,
CHIA5 has been found to be overexpressed in all digestive organs (18). In the aardwolf, which
recently diverged from the other Hyaenidae species (~10 Ma) (73), and does not present
strong morphological adaptations to myrmecophagy, no host chitinases have been found to
be expressed in its salivary glands (Allio *et al*, 2023) and expression of these enzymes in other
digestive organs needs further investigation. Therefore, this species could rely more on its gut
microbiota to ensure prey digestion. Based on our results, fewer chitinolytic bacteria were
found in the gut microbiota of the southern aardwolf compared to other species but they
could still ensure chitin digestion, for instance, by overexpressing their chitinases. The
combination of the host genomic and transcriptomic data with metagenomic and
metatranscriptomic data of its microbiota would allow answering this type of question by
comparing host gene repertoires in light of the chitinolytic abilities of the gut microbiota. This
will shed light on the adaptation to chitin digestion in placentals and, more generally, on the
role of the holobiont in the adaptation to a specific function.
492
493

Materials and methods

1.   Sampling
Thirty-three fecal samples were collected from nine species representative of the five
myrmecophagous placental orders (Table 1). For armadillos and anteaters provided by the
JAGUARS collection (Cayenne, French Guiana), fecal samples from roadkill and deceased zoo
animals were obtained after unfreezing the specimens and dissecting the lower part of the
digestive tract in the lab facilities provided by Institut Pasteur de la Guyane (Cayenne, French
Guiana). Roadkill armadillos collected in the USA were also dissected to sample feces. For the
aardvark, ground pangolin, and southern aardwolf, fresh fecal samples were collected directly
in the field during fieldwork sessions conducted in Tswalu Kalahari and Tussen-die-Riviere
reserves (South Africa). The inner part of the feces was sampled with a sterile scalpel blade
to avoid soil contamination. In the South African reserves, eight soil samples were also
collected near feces sampling sites to serve as a control for potential environmental
contamination (Table S4). All fecal and soil samples were stored at -20°C in 96% ethanol
before DNA extraction.
509
510

Table 1. Detailed sample information for the 33 fecal samples collected.

Sample name	Species name	Common name	Order	Diet	Sex	Age	Wild / Captive	Country of origin	Location	Accession number Illumina	ONT
CAB M2808	Cabassous unicinctus	Southern naked-tailed armadillo	Cingulata	Myrmecophagous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925023	SRR23925022
CAB M2982	Cabassous unicinctus	Southern naked-tailed armadillo	Cingulata	Myrmecophagous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925011	SRR23925000
CAB M3141	Cabassous unicinctus	Southern naked-tailed armadillo	Cingulata	Myrmecophagous	Female	Adult	Wild	French Guiana	coll. JAGUARS	SRR23924989	SRR23924978
DASY M1746	Dasypus novemcinctus sp FG	Guianan long-nosed armadillo	Cingulata	Omnivorous	Male	Adult	Wild	French Guiana	coll. JAGUARS	NA	SRR23924967
DASY M1777	Dasypus novemcinctus sp FG	Guianan long-nosed armadillo	Cingulata	Omnivorous	Male	Juvenile	Wild	French Guiana	coll. JAGUARS	SRR23924962	SRR23924961
DASY M2265	Dasypus novemcinctus sp FG	Guianan long-nosed armadillo	Cingulata	Omnivorous	NA	Adult	Wild	French Guiana	coll. JAGUARS	SRR23924960	SRR23925021
DASY M2886	Dasypus novemcinctus sp FG	Guianan long-nosed armadillo	Cingulata	Omnivorous	Female	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925020	SRR23925019
DASY M3021	Dasypus novemcinctus sp FG	Guianan long-nosed armadillo	Cingulata	Omnivorous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925018	SRR23925017
DASY VLD165	Dasypus novemcinctus	Nine-banded armadillo	Cingulata	Omnivorous	Female	Adult	Wild	USA	Valdosta (GA)	SRR23925016	SRR23925015
DASY VLD168	Dasypus novemcinctus	Nine-banded armadillo	Cingulata	Omnivorous	Male	Adult	Wild	USA	Valdosta (GA)	NA	SRR23925014
DASY VLD172	Dasypus novemcinctus	Nine-banded armadillo	Cingulata	Omnivorous	Female	Adult	Wild	USA	Valdosta (GA)	SRR23925013	SRR23925012
KAP M2887	Dasypus kappleri	Greater long-nosed armadillo	Cingulata	Insectivorous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925010	SRR23925009
MYR M6293	Myrmecophaga tridactyla	Giant anteater	Pilosa	Myrmecophagous	Female	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925008	SRR23925007
MYR M6296	Myrmecophaga tridactyla	Giant anteater	Pilosa	Myrmecophagous	NA	Juvenile	Wild	French Guiana	coll. JAGUARS	SRR23925006	SRR23925005
MYR ZOO	Myrmecophaga tridactyla	Giant anteater	Pilosa	Myrmecophagous	Female	Adulte	Captive	France	Montpellier zoo	SRR23925004	SRR23925003
ORY T8217	Orycteropus afer	Aardvark	Tubulidentata	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23925002	SRR23925001
ORY T8513	Orycteropus afer	Aardvark	Tubulidentata	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924999	SRR23924998
ORY T8586	Orycteropus afer	Aardvark	Tubulidentata	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924997	SRR23924996
PAN T8471	Smutsia temminckii	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924995	SRR23924994
PAN T8476	Smutsia temminckii	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924993	SRR23924992
PAN T8482	Smutsia temminckii	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924991	SRR23924990
PAN T8488	Smutsia temminckii	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924988	SRR23924987
PAN T8489	Smutsia temminckii	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924986	SRR23924985
PAN T8626	Smutsia temminckii	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924984	SRR23924983
PRO TDR7	Proteles cristatus	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924982	SRR23924981
PRO TDR17	Proteles cristatus	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924980	SRR23924979
PRO TDR22	Proteles cristatus	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924977	SRR23924976
PRO TDR48	Proteles cristatus	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924975	SRR23924974
PRO TDR82	Proteles cristatus	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924973	SRR23924972
PRO TDR87	Proteles cristatus	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924971	SRR23924970
TAM M3076	Tamandua tetradactyla	Southern tamandua	Pilosa	Myrmecophagous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23924969	SRR23924968
TAM M6331	Tamandua tetradactyla	Southern tamandua	Pilosa	Myrmecophagous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23924966	SRR23924965
TAM M6584	Tamandua tetradactyla	Southern tamandua	Pilosa	Myrmecophagous	Female	Adult	Wild	French Guiana	coll. JAGUARS	SRR23924964	SRR23924963

N.B.: Diet was determined based on field observations (i.e., dissections) and the literature (11, 74-78).

2. DNA extraction

Whole DNA was extracted from fecal samples following the optimized protocol of (79) using the Genomic DNA from soil kit (NucleoSpin, Macherey-Nagel). Two successive extractions were done and a purification step was added to retrieve high molecular weight DNA suitable for long-read sequencing (79). The same kit was used to extract DNA from soil samples. Before the extraction, samples were incubated with 700 µL of the lysis buffer (SL1) (79) and 30 µl of proteinase K at 56°C for 30 minutes.

3. Library preparation and DNA sequencing

3.1. Long-read sequencing

Long-read libraries were constructed using the SKQ-LSK109 and SKQ-LSK110 library preparation kits (ONT, Oxford Nanopore Technologies, https://nanoporetech.com/). Shotgun metagenomic sequencing was done using the MinION, MK1C, and GridION devices using one R9 flowcell per sample. Between 150 and 988 ng of DNA were loaded per flowcell, which were run for 48 to 72 hours (Table S1A). For samples sequenced on the GridION, super-accurate basecalling was performed with Guppy v5+ (Qscore = 10) whereas on the MinION and MK1C fast basecalling was used (Qscore = 7). Sequencing output statistics were checked using PycoQC v2.5.2 (80) with a minimum quality score set to 7 or 10 depending on the sequencing device used (Table S1A). Rebasecalling of samples sequenced on the MinION and MK1C was done on a GPU machine with Guppy v5.0.16 (super accurate mode, Qscore = 10, config_file = dna_r9.4.1_450bps_sup.cfg) (ONT).

3.2. Short-read sequencing

Short-read shotgun metagenomic Illumina sequencing was done for all samples to generate data for long-read assemblies polishing. Either the first or second extraction was used for sequencing as they have previously been shown to be both suitable for short-read sequencing

(79). Library preparation and Illumina sequencing on a NovaSeq instrument were outsourced
to Novogene Europe (Cambridge, UK) to generate metagenomes using 50 million of 150 bp
paired-end reads each (15 Gb of raw data per sample) (Table S1A). Two samples of two
Dasypus species (DASY M1746 and DASY VLD168) could not be sequenced using short reads
due to low quantity of starting material and were not included in the following analyses;
resulting in 31 samples being analyzed. Short-read metagenomic sequencing of soil samples
was performed following the same protocol.

4. Data filtering

4.1. Long-read data

Sequencing adapters were removed with Porechop v0.2.4 (81) used with default parameters.
Reads shorter than 200 bp were removed using Filtlong v0.2.1 (82). No quality filtering was
performed at this stage as it was previously done during basecalling (Qscore > 10).

Host mitogenomes were downloaded for the nine myrmecophagous species of our
dataset from the National Center for Biotechnology Information (NCBI) Genbank database
(Table S1A). Long-read metagenomes were mapped to the host mitogenome with Minimap2
v2.17 (83, 84) with the ONT preset (*-ax map-ont*) for Oxford Nanopore reads. This allowed us
to confirm the host species. Host mitogenomes were then assembled from the mapping reads
with Flye v2.8.3 (85) with default parameters for ONT raw reads (*--nano-raw*). Mapping of
host mitogenomic reads on host mitogenomes was visually checked with Geneious Prime
2022.0.2 (86), and mitochondrial reads were removed from the metagenomes for
downstream analyses.

The same was done to remove host nuclear reads by mapping long-read
metagenomes against the host genome using Minimap2 v2.17 (83, 84) using the Nanopore
read option (*-ax map-ont*) (Table S1A). The *Dasypus novemcinctus* RefSeq genome assembly
(Dasnov 3.0; GCF_000208655.1) was downloaded from Genbank. The *Orycteropus afer*
genome was downloaded from the DNA Zoo database (87) (HiC assembly based on the draft
assembly of Di Palma *et al*, unpublished). *Myrmecophaga tridactyla*, *Tamandua tetradactyla*,
Smutsia gigantea, and *Proteles cristatus* genomes have been previously reconstructed using
long (ONT) and short (Illumina) reads assembled with MaSuRCA v3.2.9 (88) (see (89) for a
detailed description of the hybrid assembly process). For *Cabassous unicinctus*, we used the
Discovar draft genome assembly generated by (90). For species lacking an available reference
genome, the genome of the closest relative in our dataset was used: *Smutsia gigantea* for *S.*
temminckii and *Dasypus novemcinctus* for *D. kappleri* and *Dasypus sp. nov.* FG. Host reads
were removed from the metagenomes for downstream analyses. One southern naked-tailed
armadillo sample (CAB M3141) and one giant anteater sample (MYR M5295) presented a high
proportion of host reads (> 95%) and were excluded from the following analyses. The final
dataset thus included 29 samples.

Finally, contaminant human reads were removed following the same approach and
using the telomere-to-telomere human genome assembly (GCA009914755.3) as reference
(Table S1A).

 4.2.   **Short-read data**

Illumina sequencing adapters were removed using FASTP v0.20.0 with default quality filtering parameters (91). Host and human reads were removed using the same approach as for long-read metagenomes but the mapping was done with bowtie2 v2.3.5 (92) with default parameters (Table S1A).

Samtools v1.7 (93) was used to manipulate long- and short-read mapping files.

5.   Metagenome assembly

Long-read metagenomes were assembled for each sample using metaFlye v2.9 (94) with default parameters for ONT reads basecalled with Guppy v5+ (*--nano-hq*, recommended mode when < 5% of sequencing errors is expected) and the strain mode (*--keep-haplotypes*), which prevents closely related strains represented by different paths in the assembly graph to be collapsed. Long-read metagenome assemblies were then polished with short-reads using Pilon v1.24 (95) with default parameters. Short-read metagenomes were assembled using metaSPAdes v3.11.0 (96) and MEGAHIT v1.1.2 (97) with default parameters.

Assembly statistics were computed with anvi'o v7 (41) (Tables S1B and C). Anvi'o first generates a contig database in which k-mer frequencies for each contig are computed (k = 4). Prodigal v2.6.3 (98) is then used to predict archaeal and bacterial open reading frames and estimate the number of genes. Finally, microbial single-copy core genes are searched using HMMER (99) with the hmmscan v3.2.1 program (default in anvi'o).

6.   Reconstruction of bacterial genomes

6.1.   Genome binning and bins selection

Prior to binning, reads were mapped to metagenome assemblies to obtain the coverage information needed for binning. This was done using Bowtie2 v2.2.9 (92) for short reads and Minimap2 v2.17 (83, 84) for long reads. Genome binning was conducted on single long-read polished assemblies and short-read assemblies separately (trimmed to keep contigs longer than 1000 bp) using metaBAT2 v2.15 (100) with default parameters (minimum size of contigs set to 2500 bp and bin minimal length set to 200 kb).

Completeness and redundancy of bins were estimated using anvi'o v7 (Tables S1D and E), which relies on the detection of single-copy core genes (SCG). Bins with more than 90% completeness and less than 5% redundancy were selected for downstream analyses. Based on these criteria, 239 genome bins were selected from the long-read assemblies and 254 from the short-read assemblies. Genome bins statistics were computed using anvi'o v7 as done for the metagenome assemblies (Tables S1D and E).

6.2.   Genome bins dereplication

To remove potentially redundant bins, a dereplication step was performed on the set of selected bins using dRep v3.3.0 (101). dRep first filters genomes based on their length (> 50 000 by default) and their completeness and redundancy as computed with CheckM v1.1.11 (102) (> 75% completeness, < 25% redundancy by default). dRep then clusters genomes based

on an Average Nucleotide Identity (ANI) threshold of 90% using the Mash algorithm v2.3
(103). Genomes having at least 90% ANI are then clustered using fastANI v1.33 (104), here
with a secondary ANI threshold of 98% (recommended to avoid mis-mapping of metagenomic
reads against bins, for example, to study their distribution across metagenomes). By default,
at least 10% of the genome is compared (minimum alignment fraction). Two bins did not pass
dRep quality filtering in each dataset and were not included in the analysis. In the long-read
dataset, 38 genomes were removed by dRep with an ANI of 98%, resulting in 201 unique
genome bins. In the short-read dataset, 48 genomes were removed, resulting in 206 unique
genome bins. Selected genome bin statistics were compared between bins reconstructed
from long-read and short-read assemblies using anvi'o v7 (see supplementary results part 1
available via Zenodo).

To determine whether similar genome bins were reconstructed using long-read
polished assemblies (n = 201) or short-read assemblies (n = 206), dRep was also run on the
set of all selected bins (n = 407) with 98% ANI to remove redundant genome bins (93 genomes
removed, resulting in 314 non-redundant high-quality bins). To capture the majority of the
bacterial taxa present in our samples, we decided to combine high-quality selected bins
reconstructed from both long- and short-read assemblies dereplicated with an ANI of 98% in
our final dataset for downstream analyses (n = 314; Table S1F) (see supplementary results
part 1 and dRep output results available via Zenodo).

6.3.   Taxonomic assignment of selected bins
The taxonomy of selected genome bins reconstructed from long- and short-read assemblies
was assessed using anvi'o v7 which uses 22 single-copy core genes (SCGs) and the taxonomy
of the genomes defined by the Genome Taxonomy Database release 7 (GTDB) (38) from which
these genes have been extracted (Table S1F).

The 314 selected bins were placed in a phylogeny of 2566 reference prokaryotic
genomes downloaded from Genbank using PhyloPhlAn v3.0.58 (42). PhyloPhlAn first searches
universal prokaryotic marker genes (option *-d* phylophlan, a database comprising 400
markers) (105), here using Diamond v2.0.6 (106). Marker genes were then aligned using
MAFFT v7.475 (107), and cleaned using trimAl v1.4 (108). Next, marker gene alignments were
concatenated. Finally, the phylogeny was inferred using IQ-TREE v2.0.3 (109) under the LG
model. PhyloPhlAn was run with the *–diversity* high and *–fast* options, which together set a
range of parameters for the reconstruction of high-ranked taxonomic level phylogenies (42).
By default, at least 100 markers of the PhyloPhlAn database must be present in a genome for
it to be included in the analysis (*--min_num_markers* 100) and each marker should be found
in at least four genomes to be included in the analysis (*--min_num_entries* 4). Seventy of the
reference genomes had less than 100 markers and were excluded from the analysis resulting
in a total of 2810 genomes in the final reconstructed tree. The phylogeny was rooted with
Archaea so the Bacteria were monophyletic. Using the same PhyloPhlAn parameters, a
phylogeny of the set of the 314 selected bacterial genome bins was also inferred (Fig. S1). This
phylogeny was rooted according to (110), placing the root of the bacterial tree separating

Terrabacteria (e.g., Firmicutes, Actinobacteriota) and Gracilicutes (e.g., Proteobacteria,
Desulfobacterota, Campylobacterota, Fibrobacteres-Chlorobi-Bacteroidetes group).

6.4. **Distribution of selected bins in gut metagenomes**
Short-read metagenomes were mapped against the 314 selected bins using bowtie2 v2.3.5
with local sensitive alignment parameters. The distribution of bins across samples was
explored using anvi'o v7. Anvi'o enables us to compute the percentage of mapped reads on
each bin for each metagenome and computes statistics such as the coverage, percentage of
recruited reads, abundance, and detection (horizontal coverage) of bins across metagenomes
(samples). The distribution across samples was studied using the anvi'o interface by
visualizing the detection of selected genome bins across samples. A detection threshold of
0.25 % of the reference covered by at least one read was chosen to consider a bin detected
in a sample. The number of bins shared among the different samples and host species and
bins specific to a sample or host species were calculated. Applying the same approach, the
distribution of selected bins reconstructed from the aardvark, ground pangolin, and southern
aardwolf samples (n = 140) was explored in the eight soil samples collected in South Africa
near feces sampling sites (Fig. S4).

7. Identification of bacterial chitinase genes
Chitinase genes were searched in selected genome bins. More specifically, sequences of
enzymes belonging to the Glycoside Hydrolase 18 (GH18) family (comprising chitinases and
chitin-binding proteins) as determined by the classification of Carbohydrate Active enzymes
(CAZymes) (31) were scanned using dbCAN2 (111) with default parameters. dbCAN2 is
designed to annotate CAZymes sequences in genomes using an HMMER search (99) against
the dbCAN HMM database containing HMM models for each CAZyme family. Proteins were
predicted using Prodigal (98).
The amino acid sequences of 420 identified GH18 genes were imported in Geneious
Prime 2022.0.2. Sequences were aligned using MAFFT v7.450 with default parameters. The
alignment was cleaned by removing sites not present in at least 50% of the sequences,
resulting in an alignment of 390 amino acids. The chitinase gene tree was inferred with RAxML
v8.2.11 (112) under the LG+G model and the rapid hill-climbing algorithm for topology
rearrangements. After removing 26 sequences that did not align correctly and had very long
branches, the final alignment included 394 sequences. The phylogeny was rooted with the
clades containing sequences without chitinolytic sites as they were divergent from the other
sequences. The conserved chitinolytic site (DXXDXDXE) (48, 49), typical of chitin-degrading
enzymes, was searched in the aligned sequences. BLAST searches against the NCBI non-
redundant protein database were conducted to assess whether identified GH18 genes were
similar to known microbial chitinases.

8. Data availability

Raw RNAseq short Illumina and long ONT reads have been submitted to the Short Read Archive (SRA) of the National Center for Biotechnology Information (NCBI) and are available under BioProject number PRJNA942254. Metagenome assemblies, genome bins (all and selected), phylogenetic datasets (alignments and corresponding trees), and other supplementary materials and results are available from zenodo.org (https://doi.org/: 10.5281/zenodo.7995394).

Acknowledgments

We are indebted to Lionel Hautier, Rémi Allio, Fabien Condamine, Sérgio Ferreira-Cardoso, Pierre-Henri Fabre, Quentin Martinez, Mathilde Barthe, Aude Caizergues, Nathalie Delsuc, Jessica Briner, Tshediso Putsane, Baptiste Chenet, Edith Guilloton, Solène Lefort, Margo Traimond, Michel Blanc, Bertrand Goguillon, Antoine Baglan for their help with sampling. Fieldwork sessions at Tswalu Kalahari Reserve (South Africa) were conducted under the auspices of the Tswalu Foundation thanks to Duncan MacFadyen (Oppenheimer Generations Research and Conservation), Dylan Smith, and Gus van Dyk. We thank two anonymous reviewers for their helpful comments. Computational analyses benefited from the Montpellier Bioinformatics Biodiversity (MBB) platform. This work has been supported by grants from the European Research Council (ConvergeAnt project: ERC-2015-CoG-683257) and Investissements d'Avenir of the Agence Nationale de la Recherche (CEBA: ANR-10-LABX-25-01; CEMEB: ANR-10-LABX-0004). The JAGUARS collection, managed by Kwata NGO is supported by the Collectivité Territoriale de Guyane and the Direction Générale des Territoires et de la Mer de Guyane. This is contribution ISEM 2023-XXX-SUD of the Institut des Sciences de l'Evolution de Montpellier.

References

1. Ley RE, Lozupone CA, Hamady M, Knight R, Gordon JI. 2008. Worlds within worlds: evolution of the vertebrate gut microbiota. 10. Nature Reviews Microbiology 6:776–788.
2. Moeller AH, Sanders JG. 2020. Roles of the gut microbiota in the adaptive evolution of mammalian species. Philosophical Transactions of the Royal Society B: Biological Sciences 375:20190597.
3. Alessandri G, Rizzo SM, Ossiprandi MC, van Sinderen D, Ventura M. 2022. Creating an atlas to visualize the biodiversity of the mammalian gut microbiota. Current Opinion in Biotechnology 73:28–33.
4. Ley RE, Hamady M, Lozupone C, Turnbaugh PJ, Ramey RR, Bircher JS, Schlegel ML, Tucker TA, Schrenzel MD, Knight R, Gordon JI. 2008. Evolution of mammals and their gut microbes. Science 320:1647–1651.
5. Muegge BD, Kuczynski J, Knights D, Clemente JC, González A, Fontana L, Henrissat B,

Knight R, Gordon JI. 2011. Diet drives convergence in gut microbiome functions across
mammalian phylogeny and within humans. Science 332:970–974.

6. Groussin M, Mazel F, Sanders JG, Smillie CS, Lavergne S, Thuiller W, Alm EJ. 2017.
Unraveling the processes shaping mammalian gut microbiomes over evolutionary time.
1. Nature Communications 8:14319.

7. Nishida AH, Ochman H. 2018. Rates of gut microbiome divergence in mammals. Molecular
Ecology 27:1884–1897.

8. Youngblut ND, Reischer GH, Walters W, Schuster N, Walzer C, Stalder G, Ley RE,
Farnleitner AH. 2019. Host diet and evolutionary history explain different aspects of gut
microbiome diversity among vertebrate clades. 1. Nature Communications 10:2200.

9. Song SJ, Sanders JG, Delsuc F, Metcalf J, Amato K, Taylor MW, Mazel F, Lutz HL, Winker K,
Graves GR, Humphrey G, Gilbert JA, Hackett SJ, White KP, Skeen HR, Kurtis SM, Withrow
J, Braile T, Miller M, McCracken KG, Maley JM, Ezenwa VO, Williams A, Blanton JM,
McKenzie VJ, Knight R. 2020. Comparative analyses of vertebrate gut microbiomes reveal
convergence between birds and bats. mBio 11:e02901-19.

10. Thomas CM, Desmond-Le Quéméner E, Gribaldo S, Borrel G. 2022. Factors shaping the
abundance and diversity of the gut archaeome across the animal kingdom. 1. Nat
Commun 13:3358.

11. Redford KH. 1987. Ants and termites as food, p. 349–399. In Genoways, HH (ed.), Current
Mammalogy. Springer US, Boston, MA.

12. Delsuc F, Scally M, Madsen O, Stanhope MJ, de Jong WW, Catzeflis FM, Springer MS,
Douzery EJP. 2002. Molecular phylogeny of living xenarthrans and the impact of character
and taxon sampling on the placental tree rooting. Mol Biol Evol 19:1656–1671.

13. Meredith RW, Janečka JE, Gatesy J, Ryder OA, Fisher CA, Teeling EC, Goodbla A, Eizirik E,
Simão TLL, Stadler T, Rabosky DL, Honeycutt RL, Flynn JJ, Ingram CM, Steiner C, Williams
TL, Robinson TJ, Burk-Herrick A, Westerman M, Ayoub NA, Springer MS, Murphy WJ.
2011. Impacts of the cretaceous terrestrial revolution and KPg extinction on mammal
diversification. Science 334:521–524.

14. Álvarez-Carretero S, Tamuri AU, Battini M, Nascimento FF, Carlisle E, Asher RJ, Yang Z,
Donoghue PCJ, dos Reis M. 2021. A species-level timeline of mammal evolution
integrating phylogenomic Data. Nature 602: 263–267.

15. Emerling CA, Delsuc F, Nachman MW. 2018. Chitinase genes (*CHIAs*) provide genomic
footprints of a post-Cretaceous dietary radiation in placental mammals. Science Advances
4:eaar6478.

16. Ma J-E, Li L-M, Jiang H-Y, Zhang X-J, Li J, Li G-Y, Yuan L-H, Wu J, Chen J-P. 2017.
Transcriptomic analysis identifies genes and pathways related to myrmecophagy in the
Malayan pangolin (*Manis javanica*). PeerJ 5:e4140.

17. Ma J-E, Jiang H-Y, Li L-M, Zhang X-J, Li H-M, Li G-Y, Mo D-Y, Chen J-P. 2019. SMRT
sequencing of the full-length transcriptome of the Sunda pangolin (*Manis javanica*). Gene
692:208–216.

18. Allio R, Teullet S, Lutgen D, Magdeleine A, Koual R, Tilak M-K, Thoisy B de, Emerling CA,

773  Lefébure T, Delsuc F. 2023. Comparative transcriptomics reveals divergent paths of
774  chitinase evolution underlying dietary convergence in ant-eating mammals. bioRxiv
775  https://doi.org/10.1101/2022.11.29.518312.

776 19. Delsuc F, Metcalf JL, Wegener Parfrey L, Song SJ, González A, Knight R. 2014. Convergence
777  of gut microbiomes in myrmecophagous mammals. Mol Ecol 23:1301–1317.

778 20. Ma J-E, Jiang H-Y, Li L-M, Zhang X-J, Li G-Y, Li H-M, Jin X-J, Chen J-P. 2018. The fecal
779  metagenomics of Malayan pangolins identifies an extensive adaptation to
780  myrmecophagy. Front Microbiol 9:2793.

781 21. Cheng S-C, Liu C-B, Yao X-Q, Hu J-Y, Yin T-T, Lim BK, Chen W, Wang G-D, Zhang C-L, Irwin
782  DM, Zhang Z-G, Zhang Y-P, Yu L. 2022. Hologenomic insights into mammalian adaptations
783  to myrmecophagy. National Science Review nwac174.

784 22. Gooday GW. 1990. The ecology of chitin degradation, p. 387–430. In Marshall, KC (ed.),
785  Advances in Microbial Ecology. Springer US, Boston, MA.

786 23. Beier S, Bertilsson S. 2013. Bacterial chitin degradation—mechanisms and
787  ecophysiological strategies. Front Microbiol 4.

788 24. Swiontek Brzezinska M, Jankiewicz U, Burkowska A, Walczak M. 2014. Chitinolytic
789  microorganisms and their possible application in environmental protection. Curr
790  Microbiol 68:71–81.

791 25. Raimundo I, Silva R, Meunier L, Valente SM, Lago-Lestón A, Keller-Costa T, Costa R. 2021.
792  Functional metagenomics reveals differential chitin degradation and utilization features
793  across free-living and host-associated marine microbiomes. Microbiome 9:43.

794 26. Stevens CE, Hume ID. 1998. Contributions of microbes in vertebrate gastrointestinal tract
795  to production and conservation of nutrients. Physiological Reviews 78:393–427.

796 27. Šimůnek J, Hodrová B, Bartoňová H, Kopečný J. 2001. Chitinolytic bacteria of the mammal
797  digestive tract. Folia Microbiol 46:76–78.

798 28. Whitaker JO Jr, Dannelly HK, Prentice DA. 2004. Chitinase in insectivorous bats. Journal of
799  Mammalogy 85:15–18.

800 29. Irulan A, Nathan PT, Priya YS, Marimuthu S, Elangovan V. 2011. Isolation and
801  characterization of chitinase producing gut microflora of insectivorous bats. Trends in
802  Biosciences 4:8–11.

803 30. Macdonald C, Barden S, Foley S. 2014. Isolation and characterization of chitin-degrading
804  micro-organisms from the faeces of Goeldi's monkey, *Callimico goeldii*. Journal of Applied
805  Microbiology 116:52–59.

806 31. Drula E, Garron M-L, Dogan S, Lombard V, Henrissat B, Terrapon N. 2022. The
807  carbohydrate-active enzyme database: functions and literature. Nucleic Acids Research
808  50:D571–D577.

809 32. Deeba F, Shakir H, Irfan M, Qazi J. 2016. Chitinase production in organisms: A review.
810  Punjab University Journal of Zoology 31:101–106.

811 33. Adrangi S, Faramarzi MA. 2013. From bacteria to human: A journey into the world of
812  chitinases. Biotechnology Advances 31:1786–1795.

813 34. Yan Q, Fong SS. 2015. Bacterial chitinase: nature and perspectives for sustainable

bioproduction. Bioresources and Bioprocessing 2:31.

35. Rathore AS, Gupta RD. 2015. Chitinases from bacteria to human: properties, applications,
and future perspectives. Enzyme Research 2015:791907.

36. Sanders JG, Beichman AC, Roman J, Scott JJ, Emerson D, McCarthy JJ, Girguis PR. 2015.
Baleen whales host a unique gut microbiome with similarities to both carnivores and
herbivores. 1. Nature Communications 6:8285.

37. Kohl KD, Dieppa-Colón E, Goyco-Blas J, Peralta-Martínez K, Scafidi L, Shah S, Zawacki E,
Barts N, Ahn Y, Hedayati S, Secor SM, Rowe MP. 2022. Gut microbial ecology of five
species of sympatric desert rodents in relation to herbivorous and insectivorous feeding
strategies. Integrative and Comparative Biology icac045.

38. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. 2021. GTDB:
an ongoing census of bacterial and archaeal diversity through a phylogenetically
consistent, rank normalized and complete genome-based taxonomy. Nucleic Acids
Research 50:D785–D794.

39. Buist G, Steen A, Kok J, Kuipers OP. 2008. LysM, a widely distributed protein motif for
binding to (peptido)glycans. Mol Microbiol 68:838–847.

40. Musacchio A, Wilmanns M, Saraster M. 1994. Structure and function of the SH3 domain.
Progress in Biophysics and Molecular Biology 61:283–297.

41. Eren AM, Kiefl E, Shaiber A, Veseli I, Miller SE, Schechter MS, Fink I, Pan JN, Yousef M,
Fogarty EC, Trigodet F, Watson AR, Esen ÖC, Moore RM, Clayssen Q, Lee MD, Kivenson V,
Graham ED, Merrill BD, Karkman A, Blankenberg D, Eppley JM, Sjödin A, Scott JJ, Vázquez-
Campos X, McKay LJ, McDaniel EA, Stevens SLR, Anderson RE, Fuessel J, Fernandez-Guerra
A, Maignien L, Delmont TO, Willis AD. 2021. Community-led, integrated, reproducible
multi-omics with anvi'o. Nat Microbiol 6:3–6.

42. Asnicar F, Thomas AM, Beghini F, Mengoni C, Manara S, Manghi P, Zhu Q, Bolzan M,
Cumbo F, May U, Sanders JG, Zolfo M, Kopylova E, Pasolli E, Knight R, Mirarab S,
Huttenhower C, Segata N. 2020. Precise phylogenetic analysis of microbial isolates and
genomes from metagenomes using PhyloPhlAn 3.0. 1. Nature Communications 11:2500.

43. Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and other
things). Methods in Ecology and Evolution 3:217–223.

44. Suzuki TA. 2017. Links between natural variation in the microbiome and host fitness in
wild mammals. Integrative and Comparative Biology 57:756–769.

45. Smith S. 1996. Isolation and characterization of chitinases from *Dasypus novemcinctus*,
the nine-banded armadillo. MSU Graduate Theses 184.

46. Olsen MA, Blix AS, Utsi T, Sørmo W, Mathiesen SD. 1999. Chitinolytic bacteria in the minke
whale forestomach. Can J Microbiol 46:85–94.

47. Wardman JF, Bains RK, Rahfeld P, Withers SG. 2022. Carbohydrate-active enzymes
(CAZymes) in the gut microbiome. Nat Rev Microbiol 20:542–556.

48. Olland AM, Strand J, Presman E, Czerwinski R, Joseph-McCarthy D, Krykbaev R,
Schlingmann G, Chopra R, Lin L, Fleming M, Kriz R, Stahl M, Somers W, Fitz L, Mosyak L.
2009. Triad of polar residues implicated in pH specificity of acidic mammalian chitinase.

Protein Sci 18:569–578.

49. Hamid R, Khan MA, Ahmad M, Ahmad MM, Abdin MZ, Musarrat J, Javed S. 2013. Chitinases: an update. J Pharm Bioallied Sci 5:21–29.

50. Pel R, Hessels G, Aalfs H, Gottschal JC. 1989. Chitin degradation by *Clostridium* sp. strain 9.1 in mixed cultures with saccharolytic and sulfate-reducing bacteria. FEMS Microbiology Letters 62:191–200.

51. Simůnek J, Kopecný J, Hodrová B, Bartonová H. 2002. Identification and characterization of *Clostridium paraputrificum*, a chitinolytic bacterium of human digestive tract. Folia Microbiol (Praha) 47:559–564.

52. Sebaihia M, Peck MW, Minton NP, Thomson NR, Holden MTG, Mitchell WJ, Carter AT, Bentley SD, Mason DR, Crossman L, Paul CJ, Ivens A, Wells-Bennik MHJ, Davis IJ, Cerdeño-Tárraga AM, Churcher C, Quail MA, Chillingworth T, Feltwell T, Fraser A, Goodhead I, Hance Z, Jagels K, Larke N, Maddison M, Moule S, Mungall K, Norbertczak H, Rabbinowitsch E, Sanders M, Simmonds M, White B, Whithead S, Parkhill J. 2007. Genome sequence of a proteolytic (Group I) *Clostridium botulinum* strain Hall A and comparative analysis of the clostridial genomes. Genome Res 17:1082–1092.

53. Leisner JJ, Larsen MH, Ingmer H, Petersen BO, Duus JØ, Palcic MM. 2009. Cloning and comparison of phylogenetically related chitinases from Listeria monocytogenes EGD and *Enterococcus faecalis* V583. J Appl Microbiol 107:2080–2087.

54. Vaaje-Kolstad G, Bøhle LA, Gåseidnes S, Dalhus B, Bjørås M, Mathiesen G, Eijsink VGH. 2012. Characterization of the chitinolytic machinery of *Enterococcus faecalis* V583 and high-resolution structure of its oxidative CBM33 enzyme. Journal of Molecular Biology 416:239–254.

55. Vacca M, Celano G, Calabrese FM, Portincasa P, Gobbetti M, De Angelis M. 2020. The controversial role of human gut Lachnospiraceae. Microorganisms 8:573.

56. La Reau AJ, Meier-Kolthoff JP, Suen G. 2016. Sequence-based analysis of the genus *Ruminococcus* resolves its phylogeny and reveals strong host association. Microb Genom 2:e000099.

57. Bøhle LA, Mathiesen G, Vaaje-Kolstad G, Eijsink VGH. 2011. An endo-β-N-acetylglucosaminidase from *Enterococcus faecalis* V583 responsible for the hydrolysis of high-mannose and hybrid-type N-linked glycans. FEMS Microbiology Letters 325:123–129.

58. Spaink HP. 2004. Specific recognition of bacteria by plant LysM domain receptor kinases. Trends in Microbiology 12:201–204.

59. Ferreira-Cardoso S, Delsuc F, Hautier L. 2019. Evolutionary tinkering of the mandibular canal linked to convergent regression of teeth in placental mammals. Current Biology 29:468-475.e3.

60. Ferreira-Cardoso S, Fabre P-H, Thoisy B de, Delsuc F, Hautier L. 2020. Comparative masticatory myology in anteaters and its implications for interpreting morphological convergence in myrmecophagous placentals. PeerJ 8:e9690.

61. Ferreira-Cardoso S, Claude J, Goswami A, Delsuc F, Hautier L. 2022. Flexible conservatism

in the skull modularity of convergently evolved myrmecophagous placental mammals.
BMC Ecology and Evolution 22:87.

62. Perez-Lamarque B, Sommeria-Klein G, Duret L, Morlon H. 2023. Phylogenetic comparative
approach reveals evolutionary conservatism, ancestral composition, and integration of
vertebrate gut microbiota. bioRxiv https://doi.org/10.1101/2023.01.03.522549.

63. Mallott EK, Amato KR. 2021. Host specificity of the gut microbiome. Nat Rev Microbiol
19:639–653.

64. Mazel F, Davis K, Loudon A, Kwong W, Groussin M, Wegener Parfrey L. 2018. Is host
filtering the main driver of phylosymbiosis across the tree of life? mSystems 3:e00097-18.

65. Gull JM, Stahl M, Osmann C, Ortmann S, Kreuzer M, Hatt J-M, Clauss M. 2015. Digestive
physiology of captive giant anteaters (*Myrmecophaga tridactyla*): determinants of faecal
dry matter content. Journal of Animal Physiology and Animal Nutrition 99:565–576.

66. Bornbusch SL, Greene LK, Rahobilalaina S, Calkins S, Rothman RS, Clarke TA, LaFleur M,
Drea CM. 2022. Gut microbiota of ring-tailed lemurs (*Lemur catta*) vary across natural and
captive populations and correlate with environmental microbiota. Animal Microbiome
4:29.

67. Perry T, West E, Eisenhofer R, Stenhouse A, Wilson I, Laming B, Rismiller P, Shaw M,
Grützner F. 2022. Characterising the gut microbiomes in wild and captive short-beaked
echidnas reveals diet-associated changes. Frontiers in Microbiology 13:687115.

68. Banerjee S, van der Heijden MGA. 2023. Soil microbiomes and one health. 1. Nat Rev
Microbiol 21:6–20.

69. Hehemann J-H, Correc G, Barbeyron T, Helbert W, Czjzek M, Michel G. 2010. Transfer of
carbohydrate-active enzymes from marine bacteria to Japanese gut microbiota. 7290.
Nature 464:908–912.

70. Zilber-Rosenberg I, Rosenberg E. 2008. Role of microorganisms in the evolution of animals
and plants: the hologenome theory of evolution. FEMS Microbiology Reviews 32:723–
735.

71. Zepeda Mendoza ML, Xiong Z, Escalera-Zamudio M, Runge AK, Thézé J, Streicker D, Frank
HK, Loza-Rubio E, Liu S, Ryder OA, Samaniego Castruita JA, Katzourakis A, Pacheco G,
Taboada B, Löber U, Pybus OG, Li Y, Rojas-Anaya E, Bohmann K, Carmona Baez A, Arias
CF, Liu S, Greenwood AD, Bertelsen MF, White NE, Bunce M, Zhang G, Sicheritz-Pontén T,
Gilbert MPT. 2018. Hologenomic adaptations underlying the evolution of sanguivory in
the common vampire bat. 4. Nature Ecology & Evolution 2:659–668.

72. Janiak MC, Chaney ME, Tosi AJ. 2018. Evolution of acidic mammalian chitinase genes
(*CHIA*) is related to body mass and insectivory in Primates. Molecular Biology and
Evolution 35:607–622.

73. Eizirik E, Murphy WJ, Koepfli K-P, Johnson WE, Dragoo JW, Wayne RK, O'Brien SJ. 2010.
Pattern and timing of diversification of the mammalian order Carnivora inferred from
multiple nuclear gene sequences. Molecular Phylogenetics and Evolution 56:49–63.

74. Redford KH. 1985. Food habits of armadillos (Xenarthra, Dasypodidae), p. 429–438. In
G.G. Montgomery (ed.), The evolution and ecology of armadillos, sloths, and

vermilinguas. Smithsonian Institution Press, Washington, DC.

75. Redford KH. 1986. Dietary specialization and variation in two mammalian
myrmecophages (variation in mammalian myrmecophagy). Revista Chilena de Historia
Natural 59:201–208.

76. Anacleto TC da S. 2007. Food habits of four armadillo species in the Cerrado Area, Mato
Grosso, Brazil. Zoological Studies 46:529–537.

77. Vaz VC, Santori RT, Jansen AM, Delciellos AC, D'Andrea PS. 2012. Notes on food habits of
armadillos (Cingulata, Dasypodidae) and anteaters (Pilosa, Myrmecophagidae) at Serra
Da Capivara National Park (Piauí State, Brazil). eden 13:84–89.

78. Aya-Cuero C, Chacón-Pacheco J, Anacleto TCS. 2019. *Dasypus kappleri* (Cingulata:
Dasypodidae). mmsp 51:51–60.

79. Magdeleine A, Tilak M-K, Teullet S, Delsuc F. 2021. High molecular weight bacterial DNA
extraction from field-collected fecal samples preserved in ethanol for long-read
sequencing. protocol.io https://doi.org/10.17504/protocols.io.bvyin7ue.

80. Leger A, Leonardi T. 2019. pycoQC, interactive quality control for Oxford Nanopore
Sequencing. JOSS 4:1236.

81. Wick RR, Judd LM, Gorrie CL, Holt KEY 2017. 2017. Completing bacterial genome
assemblies with multiplex MinION sequencing. Microbial Genomics 3:e000132.

82. Wick RR. 2021. Filtlong (0.2.1). https://github.com/rrwick/Filtlong.

83. Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics
34:3094–3100.

84. Li H. 2021. New strategies to improve minimap2 alignment accuracy. Bioinformatics
37:4572–4574.

85. Kolmogorov M, Yuan J, Lin Y, Pevzner PA. 2019. Assembly of long, error-prone reads using
repeat graphs. 5. Nat Biotechnol 37:540–546.

86. Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A,
Markowitz S, Duran C, Thierer T, Ashton B, Meintjes P, Drummond A. 2012. Geneious
Basic: An integrated and extendable desktop software platform for the organization and
analysis of sequence data. Bioinformatics 28:1647–1649.

87. Dudchenko O, Batra SS, Omer AD, Nyquist SK, Hoeger M, Durand NC, Shamim MS, Machol
I, Lander ES, Aiden AP, Aiden EL. 2017. De novo assembly of the *Aedes aegypti* genome
using Hi-C yields chromosome-length scaffolds. Science 356:92–95.

88. Zimin AV, Puiu D, Luo M-C, Zhu T, Koren S, Marçais G, Yorke JA, Dvořák J, Salzberg SL.
2017. Hybrid assembly of the large and highly repetitive genome of *Aegilops tauschii*, a
progenitor of bread wheat, with the MaSuRCA mega-reads algorithm. Genome Res
27:787–792.

89. Allio R. 2021. Phylogenomics and comparative genomics in ant-eating mammals. Doctoral
dissertation. University of Montpellier.

90. Emerling CA, Gibb GC, Tilak M-K, Hughes JJ, Kuch M, Duggan AT, Poinar HN, Nachman
MW, Delsuc F. 2022. Genomic data suggest parallel dental vestigialization within the
xenarthran radiation. bioRxiv https://doi.org/10.1101/2022.12.09.519446.

91. Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34:i884–i890.

92. Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. 4. Nature
Methods 9:357–359.

93. Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,
1000 Genome Project Data Processing Subgroup. 2009. The Sequence Alignment/Map
format and SAMtools. Bioinformatics 25:2078–2079.

94. Kolmogorov M, Bickhart DM, Behsaz B, Gurevich A, Rayko M, Shin SB, Kuhn K, Yuan J,
Polevikov E, Smith TPL, Pevzner PA. 2020. metaFlye: scalable long-read metagenome
assembly using repeat graphs. 11. Nat Methods 17:1103–1110.

95. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, Cuomo CA, Zeng Q,
Wortman J, Young SK, Earl AM. 2014. Pilon: an integrated tool for comprehensive
microbial variant detection and genome assembly improvement. PLOS ONE 9:e112963.

96. Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. 2017. metaSPAdes: a new versatile
metagenomic assembler. Genome Res 27:824–834.

97. Li D, Liu C-M, Luo R, Sadakane K, Lam T-W. 2015. MEGAHIT: an ultra-fast single-node
solution for large and complex metagenomics assembly via succinct de Bruijn graph.
Bioinformatics 31:1674–1676.

98. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal:
prokaryotic gene recognition and translation initiation site identification. BMC
Bioinformatics 11:119.

99. Eddy SR. 2011. Accelerated Profile HMM Searches. PLOS Computational Biology
7:e1002195.

100.   Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. 2019. MetaBAT 2: an adaptive
binning algorithm for robust and efficient genome reconstruction from metagenome
assemblies. PeerJ 7:e7359.

101.   Olm MR, Brown CT, Brooks B, Banfield JF. 2017. dRep: a tool for fast and accurate
genomic comparisons that enables improved genome recovery from metagenomes
through de-replication. 12. The ISME Journal 11:2864–2868.

102.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM:
assessing the quality of microbial genomes recovered from isolates, single cells, and
metagenomes. Genome Res 25:1043–1055.

103.   Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM.
2016. Mash: fast genome and metagenome distance estimation using MinHash. Genome
Biology 17:132.

104.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. 2018. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. 1.
Nat Commun 9:5114.

105.   Segata N, Börnigen D, Morgan XC, Huttenhower C. 2013. PhyloPhlAn is a new method
for improved phylogenetic and taxonomic placement of microbes. 1. Nature
Communications 4:2304.

106.   Buchfink B, Xie C, Huson DH. 2015. Fast and sensitive protein alignment using
1020   DIAMOND. 1. Nature Methods 12:59–60.

1021   107.   Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7:
1022   improvements in performance and usability. Mol Biol Evol 30:772–780.

1023   108.   Capella-Gutierrez S, Silla-Martinez JM, Gabaldon T. 2009. trimAl: a tool for automated
1024   alignment trimming in large-scale phylogenetic analyses. Bioinformatics 25:1972–1973.

1025   109.   Nguyen L-T, Schmidt HA, von Haeseler A, Minh BQ. 2015. IQ-TREE: a fast and effective
1026   stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol
1027   32:268–274.

1028   110.   Coleman GA, Davín AA, Mahendrarajah TA, Szánthó LL, Spang A, Hugenholtz P, Szöllősi
1029   GJ, Williams TA. 2021. A rooted phylogeny resolves early bacterial evolution. Science
1030   372:eabe0511.

1031   111.   Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Busk PK, Xu Y, Yin Y. 2018.
1032   dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. Nucleic
1033   Acids Research 46:W95–W101.

1034   112.   Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-
1035   analysis of large phylogenies. Bioinformatics 30:1312–1313.

1036

Tree scale: 1

Family
- Acutalibacteraceae
- Anaerovoracaceae
- Bacteroidaceae
- Enterobacteriaceae
- Erysipelotrichaceae
- Lachnospiraceae
- None
- Oscillospiraceae
- Other
- Ruminococcaceae
- Sphingobacteriaceae
- Xanthomonadaceae

Phyla
- Firmicutes
- Proteobacteria
- Actinobacteriota
- Bacteroidota
- None
- Verrucomicrobiota
- Gammaproteobacteria

Host species
- *Myrmecophaga tridactyla*
- *Tamandua tetradactyla*
- *Cabassous unicinctus*
- *Dasypus novemcinctus sp* FG
- *Dasypus novemcinctus*
- *Orycteropus afer*
- *Smutsia temminckii*
- *Proteles cristatus*

SH3 domain-containing protein

Lysin-motif domain-containing protein

GH18, chitinases

Phyla
Family

Firmicutes

Actinobacteriota

Bacteroidota

Proteobacteria

Dasypus kappleri

Dasypus sp. nov. FG

Dasypus novemcinctus

Cabassous unicinctus

Myrmecophaga tridactyla

Tamandua tetradactyla

Orycteropus afer

Proteles cristatus

Smutsia temminckii

Sample name	Species name	Common name	Order	Diet	Sex	Age	Wild / Captive	Country of origin	Location	Accession number Illumina	ONT
CAB M2809	*Cabassous unicinctus*	Southern naked-tailed armadillo	Cingulata	Myrmecophagous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925023	SRR23925022
CAB M2962	*Cabassous unicinctus*	Southern naked-tailed armadillo	Cingulata	Myrmecophagous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925011	SRR23925000
CAB M3141	*Cabassous unicinctus*	Southern naked-tailed armadillo	Cingulata	Myrmecophagous	Female	Adult	Wild	French Guiana	coll. JAGUARS	SRR23924989	SRR23924978
DASY M1746	*Dasypus novemcinctus sp FG*	Guianan long-nosed armadillo	Cingulata	Omnivorous	Male	Adult	Wild	French Guiana	coll. JAGUARS	NA	SRR23924967
DASY M1777	*Dasypus novemcinctus sp FG*	Guianan long-nosed armadillo	Cingulata	Omnivorous	Male	Juvenile	Wild	French Guiana	coll. JAGUARS	SRR23924962	SRR23924961
DASY M2255	*Dasypus novemcinctus sp FG*	Guianan long-nosed armadillo	Cingulata	Omnivorous	NA	Adult	Wild	French Guiana	coll. JAGUARS	SRR23924960	SRR23925021
DASY M2865	*Dasypus novemcinctus sp FG*	Guianan long-nosed armadillo	Cingulata	Omnivorous	Female	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925020	SRR23925019
DASY M3021	*Dasypus novemcinctus sp FG*	Guianan long-nosed armadillo	Cingulata	Omnivorous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925018	SRR23925017
DASY VLD165	*Dasypus novemcinctus*	Nine-banded armadillo	Cingulata	Omnivorous	Female	Adult	Wild	USA	Valdosta (GA)	SRR23925016	SRR23925015
DASY VLD168	*Dasypus novemcinctus*	Nine-banded armadillo	Cingulata	Omnivorous	Male	Adult	Wild	USA	Valdosta (GA)	NA	SRR23925014
DASY VLD172	*Dasypus novemcinctus*	Nine-banded armadillo	Cingulata	Omnivorous	Female	Adult	Wild	USA	Valdosta (GA)	SRR23925013	SRR23925012
KAP M2867	*Dasypus kappleri*	Greater long-nosed armadillo	Cingulata	Insectivorous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925010	SRR23925009
MYR M5293	*Myrmecophaga tridactyla*	Giant anteater	Pilosa	Myrmecophagous	Female	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925008	SRR23925007
MYR M5295	*Myrmecophaga tridactyla*	Giant anteater	Pilosa	Myrmecophagous	NA	Juvenile	Wild	French Guiana	coll. JAGUARS	SRR23925006	SRR23925005
MYR ZOO	*Myrmecophaga tridactyla*	Giant anteater	Pilosa	Myrmecophagous	Female	Adulte	Captive	France	Montpellier zoo	SRR23925004	SRR23925003
ORY TS217	*Orycteropus afer*	Aardvark	Tubulidentata	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23925002	SRR23925001
ORY TS513	*Orycteropus afer*	Aardvark	Tubulidentata	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924999	SRR23924998
ORY TS566	*Orycteropus afer*	Aardvark	Tubulidentata	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924997	SRR23924996
PAN TS471	*Smutsia temminckii*	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924995	SRR23924994
PAN TS475	*Smutsia temminckii*	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924993	SRR23924992
PAN TS482	*Smutsia temminckii*	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924991	SRR23924990
PAN TS488	*Smutsia temminckii*	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924988	SRR23924987
PAN TS489	*Smutsia temminckii*	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924986	SRR23924985
PAN TS525	*Smutsia temminckii*	Ground pangolin	Pholidota	Myrmecophagous	NA	NA	Wild	South Africa	Tswalu Kalahari reserve	SRR23924984	SRR23924983
PRO TDR7	*Proteles cristatus*	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924982	SRR23924981
PRO TDR17	*Proteles cristatus*	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924980	SRR23924979
PRO TDR22	*Proteles cristatus*	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924977	SRR23924976
PRO TDR49	*Proteles cristatus*	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924975	SRR23924974
PRO TDR62	*Proteles cristatus*	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924973	SRR23924972
PRO TDR67	*Proteles cristatus*	Southern aardwolf	Carnivora	Myrmecophagous	NA	NA	Wild	South Africa	Tussen-die-Riviere reserve	SRR23924971	SRR23924970
TAM M3075	*Tamandua tetradactyla*	Southern tamandua	Pilosa	Myrmecophagous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23924969	SRR23924968
TAM M5331	*Tamandua tetradactyla*	Southern tamandua	Pilosa	Myrmecophagous	Male	Adult	Wild	French Guiana	coll. JAGUARS	SRR23925966	SRR23924965
TAM M5584	*Tamandua tetradactyla*	Southern tamandua	Pilosa	Myrmecophagous	Female	Adult	Wild	French Guiana	coll. JAGUARS	SRR23924964	SRR23924963

N.B.: Diet was determined based on field observations (i.e., dissections) and the literature (11, 74-78).