Traditionally fermented foods still as a critical avenue impacting host gut antibiotic

resistome

Yutong Li[1], Siying Fu[1], Matthias S. Klein[1], Hua Wang[1*]

[1]Department of Food Science and Technology, The Ohio State University. 2015 Fyffe Court,

Columbus, OH 43210, USA

*Corresponding author: wang.707@osu.edu

1   **Abstract**

Disrupted gut microbiota as a critical risk factor for many noncommunicable diseases is largely

driven by gut microbiota-impacting drugs, especially orally administered and biliary excreted

antibiotics. Fermented food consumption has been encouraged to replenish disrupted gut

microbiota; however, its overall impact on host gut health remains to be elucidated. This study

examined traditionally fermented foods and the gut microbiota of consumers of fermented foods

for antibiotic resistance. Dietary intervention with fermented foods was found to lead to a surge

in the antibiotic resistome in the gut microbiota of most human subjects. The antibiotic resistome

was further illustrated in traditionally fermented food samples, and antibiotic-resistant bacteria

were isolated from approximately 90% of the retail kimchi and all artisan cheeses assessed in

this pilot screening. The identified isolates included pathogens of importance in nosocomial

infections such as *Klebsiella pneumoniae* and *Enterococcus*, as well as commensals and lactic

acid bacteria, and some isolates exhibited extremely high minimum inhibitory concentration

(MIC) against antibiotics of clinical significance. Exposing fermented food microbiota to

representative antibiotics further led to a boost in the corresponding antibiotic and multidrug

resistance gene pools and disturbed microbiota. The results revealed an underestimated public

health risk associated with fermented food interventions, particularly in susceptible populations

with gastrointestinal tract symptoms and compromised immune functions seeking gut microbiota

rescue. These findings call for more comprehensive assessments of the benefits and potential

safety challenges associated with traditionally fermented foods and strategic movements to

mitigate unnecessary damage to the host gut microbiota and improve food safety.

<u>Key words</u>

Gut microbiota; traditionally fermented foods; dietary intervention; antibiotic resistance;

resistome; opportunistic pathogens; kimchi; artisan cheese

Introduction

The rapid rise in antibiotic resistance (AR) negates the effective treatment of bacterial infections, shaking the foundation of modern medicine. AR bacteria in host gastrointestinal tract further indirectly contribute to gut microbiota dysbiosis resulting from even short-term antibiotic treatment.[1] Gut microbiota destruction has been recognized as a critical, shared risk factor for a growing list of noncommunicable "modern" diseases, ranging from malfunction of the host immune system, type-II diabetes, brain/neurological disorders, *Clostridium difficile* infections, to certain cardiovascular diseases and cancers.[2-10] Both AR and noncommunicable diseases are among the top global public health threats in the 21[st] century.[11]

For decades, the broad applications of antibiotics have been blamed for the rapid surge of AR and, more recently, for disrupted host gut microbiota and associated "modern" diseases. But despite limiting the uses of antibiotics has been the primary control strategy worldwide, even essential antibiotic applications for infection prevention and treatment still cause irreversible damages in host gut microbiota.[12,13] AR is a complicated issue with multiple risk factors.[14,15] Particularly, gut impacting antibiotics, i.e., the mainstream oral administration of antibiotics, and using drugs with primary biliary instead of renal excretion, rather than the application of antibiotics, have been the key and shared driver for the rapid surge of the antibiotic resistance gene (ARG) pool, massively disrupted gut microbiota, and the rise of pathogens in gut microbiota.[16,1] This paradigm-changing discovery is further supported by clinical observations on the trends of AR worldwide.[17,1] The finding is also applicable to non-antibiotic drugs with impact on gut microbiota.[18] According to CDC, over 200 million outpatient oral antibiotic prescriptions were given in the US annually.[19] Assuming even distribution, this is equivalent to

over 60% of the U.S. population alone affected every year, contributing to the global rising trends of the aforementioned diseases.

Given its impact on host health, various approaches to replenish damaged gut microbiota have been attempted. Fecal microbiota transplant (FMT) is gaining approval for treating *C. difficile* infections,[20,21] but the procedure was associated with acquired infections, including death by ESBL *E. coli*.[22] It further led to a surge in resistome in both human and gnotobiotic pig recipients after FMT procedures.[23] Autologous fecal microbiota transplantation (auto-FMT) facilitated the restoration of the gut microbiota and immune functions in patients.[24] Yet, re-assessing the gut microbiota profiles and resistome illustrated that auto-FMT using own fecal microbiota banked before disruptive drug therapy, with the intention of avoiding the introduction of infectious agents from other donor(s), still resulted in the rise of resistome and opportunistic pathogens in multiple recipients even without further antibiotic treatment (Wang et al., unpublished data. Communicated with authors of MKSCC with consensus on the above conclusion in January 2021).

Fermented foods have emerged in recent years as a popular alternative to repair disrupted gut microbiota.[25,26] Wastyk et al.[26] reported that consuming diets high in fermented foods increased diversity of the gut microbiota and improved immune function of the human recipients, while the same effect was not observed in subjects with diets high in plant fibers. Fermented vegetables, followed by kombucha and fermented dairy products (yogurt and kefir), were among the most consumed categories of fermented foods by this group of subjects (Supplemental Data STable 1).

However, fermented foods that are rich in viable microbes are susceptible to AR. Before 2007, fermented dairy foods had been the most significant foodborne avenue transmitting AR to

consumers.[27] A gram of retail cheese contained up to 10^8 copies of AR genes,[28] and the *Bifidobacterium* strain supplemented to yogurt products worldwide used to contain a tetracycline (Tet) resistance-encoding gene (personal communication, 2007 ASM General Meeting). Various bacteria isolated from fermented foods with mobile AR genes were able to cause acquired resistance in human commensals and pathogens via horizontal gene transfer mechanisms.[27,29,30] Although successful mitigation of the AR gene pool was quickly achieved in mainstream fermented dairy products by 2011, primarily by removal of problematic fermentation starter cultures and probiotic strains from the product lines,[31,15] other traditionally fermented products may still be prone to AR.[32-48] Unlike mainstream modern dairy fermentation, which uses pasteurized milk and commercial starter cultures carefully screened by major culture companies, traditionally fermented vegetables and artisan cheeses, for instance, still rely on microbiota associated with raw materials, environment, or "mother" cultures other than carefully screened commercial starters.[31-33,40,43,47,49]

The fermented food and beverage market is expected to grow by 5.6% in the next 10 years and will exceed US$ 989.2 billion by 2032 amid escalating demand for healthy and nutritious foods.[50] The COVID-19 pandemic has further facilitated the explosive growth of sales of fermented products such as sauerkraut and kimchi.[51] Given the booming market and increased consumption, re-assessing potential AR risks associated with traditionally fermented food products has become an urgent food safety and public health need, especially for susceptible populations with compromised digestive tract and immune systems. Therefore, this study investigated the incidence of antibiotic resistome in representative fermented foods and the impact of fermented food interventions on the human gut resistome.

Results

Impact of food intervention on fecal microbiota resistome. Ten-week dietary interventions with diets high in fermented foods or plant fibers led to 11 increased and 6 decreased, as well as 10 increased and 5 decreased gut antibiotic resistomes in human subjects, respectively, but the impact of the two types of diets was distinctive (Fig 1 & Fig 2). Fermented foods intervention led to significant changes in fecal antibiotic resistome of subjects with a p-value of 0.03, and the mean of resistome increased from 0.36 to 0.42 copies/16S rRNA gene (Fig 2A). Intervention with high plant fiber diets caused insignificant changes in the host gut resistome, with a p-value of 0.94. The mean resistome remained at 0.38 copies/16S rRNA gene in subjects before and after dietary intervention (Fig 2B). The summarized resistome data are illustrated in Supplemental Data STable 2. For details on the resistome data, see the Supplemental Data at the end (STables 7 and 8).

[Insert Fig 1 here]

[Insert Fig 2 here]

[Insert Table 1 here]

Antibiotic resistome of traditionally fermented foods. Results of the pilot survey on retail fermented foods supported the aforementioned impact on the host gut resistome. Multidrug resistance, bacitracin, and macrolide-lincosamide-streptogramin (MLS) genes were most abundant in pooled kimchi samples (K1-K4) recovered from brain heart infusion (BHI) plates (Fig 3A and Table 1), and they were also among the most abundant AR genes of all five individual kimchi microbiota (K7-K11) assessed directly, despite the exact abundance of the top AR genes varied among these samples (Table 1 & Fig 4A). Likewise, even though AR isolates were identified in eight kimchi samples (K5, K7-K13, Table 2), the overall abundance of the

resistome of the assessed samples also varied, ranging from 0.372 (K9) to 0.029 (K11) copies of

AR genes/16S rRNA (Fig 4A).

[Insert Fig 3 about here]

[Insert Fig 4 about here]

[Insert Table 2 here]

Fig 4B further compares the resistome outcomes of kimchi sample K7 using total DNA extracted

from kimchi microbiota directly (0.048 copies of AR genes/16S) and total DNA of kimchi

microbiota recovered from BHI agar plates (2.909 copies of AR genes/16S). The results

indicated that with specific medium and incubation conditions, even without antibiotic selective

pressure, certain bacteria in the kimchi microbiota might have been selectively enriched, leading

to a drastic difference in the detected antibiotic resistome. Likewise, the top genera of pooled

kimchi microbiota recovered from BHI plates (Supplemental Data STable 3) were affected by

cultivation conditions as well, as the illustrated resistome (Fig 4B).

Indeed, Table 3 illustrates that the top genera of each individual kimchi microbiota (K7 to K11)

assessed using total DNA extracted directly were dominated by lactic acid bacteria, which are

well-recognized for driving kimchi fermentation. The data further demonstrated that the

antibiotic resistome and AR bacteria were still prevalent in these successfully fermented kimchi

products.

[Insert Table 3 here]

The resistome results of pooled artisan cheese samples C1 to C4 (Fig 3B) and individual cheese

samples C5 to C8 (Fig 4C), all using microbiota recovered from BHI agar plates, further

illustrated the prevalence of various AR genes in these products. Multidrug, MLS, and bacitracin were among the top AR genes of the pooled samples C1-C4, while multidrug, bacitracin, tetracycline, and MLS were the most abundant among individual samples C5 to C8 (Table 4). Notably, the abundance of the resistome of cheese microbiota recovered from BHI plates may also be affected by the medium and culture conditions, deviating from those of the original samples. Supplementary STable 5 further illustrated that *Staphylococcus* (including *Mammaliicoccus,* also known as *Staphylococcus*) was the dominant genus of individual artisan cheese microbiota recovered from BHI agar plates.

[Insert Table 4 here]

Representative AR bacteria from traditionally fermented foods. Various AR bacteria were isolated from traditionally fermented products. Under the very limited cultivation conditions applied in this study, AR bacteria were identified in eight out of nine kimchi and four out of four cheese samples assessed by culture recovery (Tables 2 and 5).

[Insert Table 5 here]

Identified AR isolates from kimchi products ranged from important human pathogens and opportunistic pathogens, such as *Klebsiella pneumoniae, Serratia marcescens* etc., which are recognized agents for nonsocomial infections,[52,53] to organisms still considered commensals, as well as lactic acid bacteria driving food fermentations. For instance, isolates of several species of *Latilactobacillus, Leuconostoc* and *Weissella* were among those highly resistant to the four antibiotics screened for, namely tetracycline (Tet), erythromycin (Ery), ampicillin (Amp), and vancomycin (Van). Some isolates exhibited multidrug resistance.

The AR isolates identified from cheese products were mostly *Staphylococcus* spp. resistant to tetracycline, and one isolate was resistant to Van. An isolate of *Enterococcus* sp. was highly resistant to all the four antibiotics assessed.

Antibiotic exposure further shaped the profiles of fermented food originated microbiota. As illustrated in Fig. 5 and Supplementary STable 5, the cheese microbiota harvested from BHI plates without antibiotics was dominated by *Mammaliicoccus* sp. (formerly *Staphylococcus*), *Staphyloccocus, Camobacterium, Psychrobacter* and *Glutamicibacter (*belonging to Micrococcaceae) for cheese sample C5. However, in the presence of Tet or Amp, the dominant bacteria were *Stenotrophomonas*, *Pseudomonas*, *Burkholderia*, *Alcanivorax* and *Streptococcus*. Likewise, while the dominant cultures for cheese C7 recovered on BHI were *Staphylococcus* and *Lactococcus*, the top-ranked bacteria of cheese-originated microbiota shifted to *Pseudomonas, Burkholderia*, *Staphylococcus, Streptococcus* and *Leuconostoc* on BHI-Amp and *Staphylococcus* on BHI-Tet. In the case of C7 and pooled sample C1-C4, the cheese starter culture *Lactococcus* was eliminated when the cheese microbiota was exposed to either of the two antibiotics.

[Insert Fig 5 about here]

Although antibiotic-containing agar plates were used to screen for AR bacteria, the above results also indicated how the food-originated microbiota might impact the gut microbiota when the hosts receive antibiotic treatment. *Stenotrophomonas*, *Pseudomonas*, *Burkholderia,* and *Staphylococcus* are opportunistic pathogens associated with antibiotic-resistant hospital-acquired infections, and most *Streptococcus* species are pathogenic.

Microbial profiles by metagenomics of dominant cheese bacteria recovered on BHI plates included *Staphylococcus, Enterococcus* and *Lactocaseibacillus* (formerly *Lactobacillus*) for

cheese sample C6, and *Enterococcus, Tetragenococcus, Staphylococcus, Lactococcus* and *Camobacterium* for sample C8 (Fig. 5 and Supplementary STable 5). In agreement with this, some of the mentioned antibiotic-resistant bacteria, such as multidrug-resistant *Enterococcus* sp. and *Staphylococcus* were also confirmed in these samples (Table 5).

Discussion and Conclusion

Although unexpected for many, the susceptibility of fermented foods to AR has been well-recognized at least in the food microbiology community. Since the first systematic demonstration of the problem in the early 2000s with a broad spectrum of AR isolates, including starter cultures, opportunistic pathogens, and commensals of mainstream fermented dairy products being identified and characterized,[27] AR bacteria have been isolated from various fermented foods worldwide.[54-56] For instance, *Pantoea agglomerans* (formerly *Enterobacter agglomerans* or *Erwinia herbicola*), an opportunistic pathogen that causes a wide range of opportunistic infections, especially in immunocompromised patients, was isolated from kimchi in South Korea. The genome of *P. agglomerans* isolate KM1 contains 13 genes conferring resistance to clinically important antibiotics, and the strain exhibited immunostimulatory properties *in vitro*, including the production of pro-inflammatory and anti-inflammatory cytokines in stimulated cells.[57]

This study, however, using a combination of approaches, for the first time systematically reveals the impact of dietary intervention by fermented foods on antibiotic resistome in human gut microbiome, backed by data on the prevalence and abundance of antibiotic resistome in popular fermented foods. The original human study on dietary intervention by Wastyk et al.[26] had a total of 36 healthy adult subjects who completed the study; half received fermented foods, and the other half received dietary fiber intervention. The sample size was small, providing the variables

for human subjects and activities. Among these, the fecal microbiome outcomes of 32 subjects with required data points, including 17 subjects with fermented food intervention and 15 subjects as control (high plant fiber diet), were assessed and reported for the impact of dietary intervention on the host gut resistome in this study.

Although the sample size of the human study was rather limited, the illustration of antibiotic resistome in kimchi and artisan cheese microbiome, along with the high prevalence of confirmed AR bacteria in approximately 90% of the kimchi and all artisan cheeses assessed in this pilot study, supports the conclusion that consuming these traditionally fermented products would result in an increase in the gut antibiotic resistome in consumers. Since only limited conditions were applied in cultivating AR bacteria, and resistance against only four antibiotics was assessed in this pilot study, the recovered isolates represented only a small percentage of the AR microbiota of the fermented foods. The results of this pilot study thus call for more extensive investigations on the food safety risks associated with traditionally fermented foods and for effective mitigation strategies to protect public health.

It is worth noting that even in subjects who consumed only yogurt and kombucha (Subject 8020) or yogurt, kombucha, and kefir (Subject 8014) (STables 2 and 6), the gut resistome of the subjects still increased by 11% and 17%, respectively, after the 10-week intervention. The data suggest that consuming these products at least did not reduce gut antibiotic resistance in the subjects.

The identification of AR bacteria from these products, ranging from human and plant pathogens to lactic acid bacteria that drive food fermentation, especially those exhibiting multidrug resistance and with extremely high MICs against antibiotics of clinical significance, is

particularly concerning. Multidrug-resistant *Klebsiella pneumoniae, Serratia marcescens,* as well

as *Stenotrophomonas*, *Pseudomonas*, *Burkholderia, Staphylococcus* and *Streptococcus* spp. are

also known to cause hospital-acquired infections, including some of the most troublesome cases

in healthcare facilities. Thus, data from this study also provide an alternative interpretation of

the route of dissemination and potential origins of infections in patients.

Balanced gut microbiota as well as effective mucosal, intestinal epithelial, and gut vascular

barriers as part of an integral and functional gastrointestinal tract system are essential for host

health.[58] Physical damage to the intestinal barriers has various health consequences. Intestinal

tract problems such as inflammatory bowel diseases, enteric infections, or overgrowth by

pathogens such as EHEC and Shiga toxin-producing *E. coli*, *Shigella*, *Salmonella*,

Campylobacter, *C. difficile*, enteric viruses, and certain drug treatments such as the antibiotic

clindamycin, and even alcohol consumption can cause acute or chronic intestinal barrier

damages. Once the intestinal barrier collapses, intestinal microorganisms and their metabolites

can enter the bloodstream, reach other organs, and cause further health damage. Once they enter

the bloodstream, even commensals and probiotics can also be problematic. For instance, the

popular probiotic strains *Bifidobacterium* and *L. rhamnosus* GG were found causative to

bacteremia and sepsis in high-risk populations, with strain confirmation by metagenomic

sequencing.[59-68] While bacteremia and sepsis by commercial probiotic strains absence of AR

genes are still controllable by mainstream antibiotics such as ampicillin, introducing the broad

spectrum of AR bacteria highly resistant to clinically important antibiotics associated with

traditionally fermented foods presents an underestimated and potentially serious public health

risk, especially to targeted populations suffering from various gut symptoms associated with

"leaky gut" as well as compromised immune functions. Bacteremia and sepsis risk, antibiotic

therapy failures, and further gut microbiota destruction after antibiotic treatment may

significantly increase in targeted susceptible consumer populations who intend to repair the

damaged gut microbiome by enhancing fermented food consumption.

With the recipients of over 200 million oral antibiotic prescriptions added to the susceptible

population suffering from broad gut microbiota destruction annually in the US alone,[19] the

impacted patient population worldwide is astonishing. While various gut microbiota-replenishing

approaches, including fermented food intervention, have been practiced desperately, they all

have introduced additional public health risks to patients. The results from this study thus further

call for strategic efforts to tackle the key and shared driver of the antibiotic resistome surge and

gut microbiota disruption in humans and animals by mitigating the applications of gut-impacting

antibiotics (oral administration and biliary excretion) and offering alternatives to significantly

minimize unnecessary damages to the gut microbiome for productive outcomes.

14   **Methods and Materials**

Host gut resistome assessment. In the original study by Wastyk et al.[26], 39 healthy adult

participants were recruited, with 18 randomly assigned to the dietary intervention group by

fermented foods, and the remaining 21 were assigned to the plant fiber intervention group. Three

participants in the fiber intervention group dropped out of the study; thus, both groups had 18

participants who completed the entire study (25 female,11 male, with an average age of 52 ± 11

years). Fecal shotgun metagenomics data of these participants were collected at four checkpoints,

(Week -2, Week 0, Week 8, and Week 10) throughout the study and deposited in the NCBI

BioProject database (ID number: PRJNA743361). After initial screening for data availability,

three participants in the fiber group and one participant in the fermented food group were

excluded from our study because of missing Week 10 endpoint shotgun sequencing data in the

database provided by the original research team. Raw sequencing data were retrieved from the

Sequence Read Archive (. sra) format, and converted to. fastq using the NCBI SRA Toolkit

(version 3.0.0) and further processed on a high-performance supercomputer at the Ohio

Supercomputer Center.

For quality control, the FASTP tool (version 0.22.0) was used (parameters: -q 20 -u 20 -n 2 -l 80)

to clean and trim the raw sequences before further annotation and analysis, as described by Chen

et al.[69]. The cleaned sequences were then processed using the ARGs-OAP pipeline (Online

Analysis Pipeline for Anti-biotic Resistance Genes, version 2.5).[70] The resistome analysis criteria

were set as hit length of 50%, e-value of 1e-07 and identity of 80%. The generated data were

further analyzed and plotted using Tidyverse packages, including ggplot2 (version 3.3.6), dplyr

(version 1.0.10), ggstatsplot package (version 0.10.0), and car package (Companion to Applied

Regression, version 3.1-1) on R version 4.2.1.[71-75]

Data transformation and analysis. The sum of the total ARG was calculated by adding up the

relative abundances of all types of ARG annotated by ARGs-OAP. The percentage of total

resistome change for each subject was calculated using the following equation: ($\sum$resistome

reads after dietary intervention- $\sum$resistome reads before dietary intervention)/ $\sum$resistome reads

before dietary intervention $\times$ 100%. The resistome data from Week -2 and Week 0 were

averaged as the baseline to ensure that all subjects had baseline data before dietary intervention

and to reduce natural variation in gut microbiota. Normality and variance of the data sets were

verified using the Shapiro-Wilk normality test and Levene's test. Violin-boxplots with Student's

t-test p-values ($\alpha<0.05$) were generated using the ggstatsplot package.[74] Other plots were

constructed using the ggplot2 package.[72] Summary statistics such as mean, median, standard

deviation, and significance of dietary intervention were calculated using the dplyr package.[75]

Source of fermented food samples. Representative kimchi samples of different brands were

purchased from seven retail stores, one chain restaurant, and two local Japanese and Korean

restaurants in Columbus, OH, between 2021 and 2022. Artisan cheese products were purchased

from a national grocery chain store in Columbus, OH, with four purchased in 2021 and four in

2022. Sample designation and brief descriptions are illustrated in Supplementary Data STable 1.

Recovery and assessment of viable bacteria from fermented food samples. Five grams of each

food sample were stomached in 45 ml 0.1% peptone water using a Seward stomacher 80 (Seward,

UK). The juice was assessed for total bacteria and AR bacteria using Brain Heart Infusion (BHI),

Luria-Bertani (LB), or De Man, Rogosa, and Sharpe (MRS) agar plates with cycloheximide or

nystatin as a mold inhibitor, with or without the corresponding antibiotic (32 µg/mL Amp; 32 or

64 µg/mL Tet; 2 µg/mL Ery for BHI or MRS). The plates were incubated at 30 °C under aerobic

or anaerobic conditions. Single colonies recovered from antibiotic-containing agar plates were

picked based on their morphology and identified by Sanger sequencing of 16S rRNA gene PCR

amplicons.[27] The MIC of recovered AR isolates against four commonly used antibiotics was

determined by a microdilution procedure,[76] but using the corresponding recovery media instead.

Antibiotic resistome of representative fermented food items. Twenty-five grams of food sample

was mixed with 25 ml 0.1% peptone water and stomached as described above. Large particles of

food debris were removed using sterile sieves. The liquid phase was centrifuged at 1711 relative

centrifugal force using Multifuge X1R (Thermo Scientific, USA, MA) for 5 min. The pellet was

washed once with 0.1% peptone water and subjected to total DNA extraction using a QIAamp

PowerFecal Pro DNA Kit (Qiagen, Hilden, Germany) following the manufacturer's instructions.

Alternatively, the culture-recovered microbiota was scraped from agar plates with 3mL of 0.1% peptone water and subjected to DNA extraction using the QIAamp kit.

The DNA samples were subjected to shotgun metagenomic sequencing (2X155bp) with an average depth of 16.7 to 20 million paired-end reads for each sample, using an Illumina NextSeq2000 Sequencing System. The raw sequences were cleaned (parameters: -q 20 -u 40 -n 2 -l 80) and analyzed using the same metagenomic tools and parameters as described above to obtain the relative abundance of ARGs normalized to the number of 16S rRNA genes. The sequencing data were further assessed for microbiota profiles through the Kraken2 pipeline with a standard database and minimum-base-quality 20, as described previously.[77] the food microbiota profile of Kimchi #7 was assessed by 16S rRNA sequencing using an Illumina MiSeq system.

All local metagenomic data analyses were conducted using a high-performance supercomputer at the Ohio Supercomputer Center.

Acknowledgement

The study was possible with funding from the OSU Women and Philanthropy. YL and SF were funded by OSU FST Graduate Research Associateship for HW. YL is also the recipient of the 2022 Elwood Caldwell graduate fellowship of the Institute of Food Technologists (IFT). We thank Dr. Erin DiCaprio of UC Davis for sharing information on the retail trends in fermented foods. We also thank our collaborators who supported and enabled this research.

We also thank Dr. Justin Sonnenburg's team for properly organizing and depositing the human study data published in *Cell* (2021) in the public domain, which enabled our data mining and reassessment.

Part of the data was presented at the 2022 ASM Microbe Satellite/Sino-Micro annual meeting (Washington, DC) and USDA multistate NC1206 project meeting on antibiotic resistance (Ames, IA).

Data availability statement

The metagenomic sequencing data were deposited in the NCBI BioProject database (ID number: PRJNA944373).

References

1. Zhou Y, Li Y, Zhang L, Wu Z, Huang Y, Yan H, Zhong J, Wang LJ, Abdullah HM, Wang HH. Antibiotic administration routes and oral exposure to antibiotic resistant bacteria as key drivers for gut microbiota disruption and resistome in poultry. Front Microbiol. 2020;11:1319. doi: 10.3389/fmicb.2020.01319.

2. Sekirov I, Russell SL, Antunes LC, Finlay BB. Gut microbiota in health and disease. Physiol Rev. 2010;90:859-904.

3. Chung H, Pamp SJ, Hill JA, Surana NK, Edelman SM, Troy EB, Reading NC, Villablanca EJ, Wang S, Mora JR, et al. Gut immune maturation depends on colonization

with a host-specific microbiota. Cell. 2012;149:1578-1593. doi: 10.1016/j.cell.2012.04.037.

4. Foster JA, McVey Neufeld KA. Gut-brain axis: how the microbiome influences anxiety and depression. Trends Neurosci. 2013;36:305-312. doi: 10.1016/j.tins.2013.01.005.

5. Blaser MJ. Antibiotic use and its consequences for the normal microbiome. Science. 2016;352:544-545.

6. Lynch SV, Pedersen O. The human intestinal microbiome in health and disease. N Engl J Med. 2016;375:2369-2379. doi: 10.1056/NEJMra1600266.

7. Smits WK, Lyras D, Lacy DB, Wilcox MH, Kuijper EJ. Clostridium difficile infection. Nat Rev Dis Primers. 2016;2:16020.

8. Meng C, Bai C, Brown TD, Hood LE, Tian Q. Human gut microbiota and gastrointestinal cancer. Genomics Proteomics Bioinformatics. 2018;16:33-49.

9. Gurung M, Li Z, You H, Rodrigues R, Jump DB, Morgun A, Shulzhenko N. Role of gut microbiota in type 2 diabetes pathophysiology. EbioMedicine. 2020;51:102590.

10. Witkowski M, Weeks TL, Hazen SL. Gut microbiota and cardiovascular disease. Circ Res. 2020;127:553-570.

11. World Health Organization (WHO). World Health Organization (WHO). Ten threats to global health in 2019. Geneva (Switzerland): World Health Organization; 2019 Jan 16 [accessed 2023 Mar 24]. https://www.who.int/news-room/spotlight/ten-threats-to-global-health-in-2019.

12. Zimmermann P, Curtis N. The effect of antibiotics on the composition of the intestinal microbiota-a systematic review. J Infect. 2019 Dec;79(6):471-489. doi: 10.1016/j.jinf.2019.08.008. Epub 2019 Aug 22. PMID: 31445907.

13. Wang HH. The impact of food and drug, and the mainstream rescue strategies on host gut microbiota and resistome. Presented at: 2022 Multistate AR meeting; Oct 27, 2022; Ames, IA, USA.

14. Wang HH. Commensal bacteria, microbial ecosystems and horizontal gene transmission: adjusting our focus for strategic breakthroughs against antibiotic resistance. In: Jaykus L, Wang HH, Schlesinger L., editors. Foodborne Microbes: Shaping the Host Ecosystems. Washington (DC): ASM Press; 2009. p. 267-281.

15. Wang HH, Zhou Y, Zhang L. Antibiotic resistance, gut microbiota and health. In: Doyle MP, Diez-Gonzalez F, Hill C, editors. Food Microbiology: Fundamentals and Frontiers. 5th ed. Washington (DC): ASM Press; 2019. p.903-926.

16. Zhang L, Huang Y, Zhou Y, Buckley T, Wang HH. Antibiotic administration routes significantly influence the levels of antibiotic resistance in gut microbiota. Antimicrob Agents Chemother. 2013;57:3659-3666. doi: 10.1128/AAC.00670-13.

17. Luber AD, Jacobs RA, Jordan M, Guglielmo BJ, Shay DK, Jarvis WR, Montecalvo M. Relative importance of oral versus intravenous vancomycin exposure in the development of vancomycin-resistant enterococci [with Reply]. J Infect Dis. 1996;173:1292-1294.

18. Weersma RK, Zhernakova A, Fu J. Interaction between drugs and the gut microbiome. Gut. 2020 Aug;69(8):1510-1519. doi: 10.1136/gutjnl-2019-319350. PMID: 32265294.

19. U.S. Centers for Disease Control and Prevention. Outpatient antibiotic prescriptions — United States, 2021. Atlanta (GA): CDC; 2021. [accessed 2022 Dec 02]. https://www.cdc.gov/antibiotic-use/data/report-2021.html.

20. U.S. Food and Drug Administration. FDA approves first fecal microbiota product. Silver

Spring (MD): FDA; 2021 Jun 28 [cited 2022 Dec 28]. https://www.fda.gov/news-

events/press-announcements/fda-approves-first-fecal-microbiota-product.

21. Therapeutic Goods Administration. Faecal microbiota transplant products regulation.

Canberra (AU): TGA; 2021 Sep 29 [accessed 2022 Mar 8].

https://www.tga.gov.au/products/biologicals-blood-and-tissues-and-advanced-

therapies/biologicals/faecal-microbiota-transplant-products-regulation.

22. U.S. Food and Drug Administration. Fecal microbiota transplantation: safety

communication - risk of serious adverse reactions due to transmission of multi-drug

resistant organisms. Silver Spring (MD): FDA; 2019 Jun 13 [accessed 2022 Oct 04].

https://www.fda.gov/safety/medical-product-safety-information/fecal-microbiota-

transplantation-safety-communication-risk-serious-adverse-reactions-due

23. Liu H, Wang HH. Impact of microbiota transplant on resistome of gut microbiota in

gnotobiotic piglets and human subjects. Front Microbiol. 2020;11:932. doi:

10.3389/fmicb.2020.00932.

24. Schluter J, Peled JU, Taylor BP, Markey KA, Smith M, Taur Y, Niehus R, Staffas A, Dai

A, Fontana E, et al. The gut microbiota is associated with immune cell dynamics in

humans. Nature. 2020;588(7837):303-307. doi: 10.1038/s41586-020-2971-8.

25. Taylor BC, Lejzerowicz F, Poirel M, Shaffer JP, Jiang L, Aksenov A, Litwin N,

Humphrey G, Martino C, Miller-Montgomery S, et al. Consumption of fermented foods

is associated with systematic differences in the gut microbiome and metabolome.

mSystems. 2020;5(2):e00901-19. doi: 10.1128/mSystems.00901-19.

26. Wastyk HC, Fragiadakis GK, Perelman D, Dahan D, Merrill BD, Feiqiao BY, Topf M, Gonzalez CG, Van Treuren W, Han S, et al. Gut-microbiota-targeted diets modulate human immune status. Cell. 2021;184(16):4137-4153.e13. doi: 10.1016/j.cell.2021.06.034.

27. Wang HH, Manuzon M, Lehman M, Wan K, Luo H, Wittum TE, Yousef A, Bakaletz LO. Food commensal microbes as a potentially important avenue in transmitting antibiotic resistance genes. FEMS Microbiol Lett. 2006;254(2):226-231. doi: 10.1111/j.1574-6968.2005.00043.x.

28. Manuzon MY, Hanna SE, Luo H, Yu Z, Harper WJ, Wang HH. Quantitative assessment of the tetracycline resistance gene pool in cheese samples by real-time TaqMan PCR. Appl Environ Microbiol. 2007;73(5):1676-1677. doi: 10.1128/AEM.02591-06.

29. Li X, Alvarez V, Harper WJ, Wang HH. Persistent, toxin-antitoxin system-independent, tetracycline resistance-encoding plasmid from a dairy *Enterococcus faecium* isolate. Appl Environ Microbiol. 2011;77(20):7096-7103. doi: 10.1128/AEM.05430-11.

30. Jahan M, Holley RA. Transfer of antibiotic resistance from *Enterococcus faecium* of fermented meat origin to *Listeria monocytogenes* and *Listeria innocua*. Lett Appl Microbiol. 2016;62(4):304-310. doi: 10.1111/lam.12553.

31. Li X, Li Y, Alvarez V, Harper WJ, Wang HH. Effective antibiotic resistance mitigation during cheese fermentation. Appl Environ Microbiol. 2011;77(20):7171-7175.

32. Belletti N, Gatti M, Bottari B, Neviani E, Tabanelli G, Gardini F. Antibiotic resistance of lactobacilli isolated from two Italian hard cheeses. J Food Prot. 2009;72(10):2162-2169.

33. Comunian R, Daga E, Dupré I, Paba A, Devirgiliis C, Piccioni V, Perozzi G, Zonenschain D, Rebecchi A, Morelli L, et al. Susceptibility to tetracycline and

erythromycin of Lactobacillus paracasei strains isolated from traditional Italian fermented foods. Int J Food Microbiol. 2010;138(1-2):151-156.

34. Karasu N, Şimşek Ö, Çon AH. Technological and probiotic characteristics of Lactobacillus plantarum strains isolated from traditionally produced fermented vegetables. Ann Microbiol. 2010;60(2):227-234.

35. Nawaz M, Wang J, Zhou A, Ma C, Wu X, Moore JE, Cherie Millar B, Xu J. Characterization and transfer of antibiotic resistance in lactic acid bacteria from fermented food products. Curr Microbiol. 2011;62:1081-1089.

36. Pan L, Hu X, Wang X. Assessment of antibiotic resistance of lactic acid bacteria in Chinese fermented foods. Food Control. 2011;22(8):1316-1321.

37. Zhou N, Zhang JX, Fan MT, Wang J, Guo G, Wei XY. Antibiotic resistance of lactic acid bacteria isolated from Chinese yogurts. J Dairy Sci. 2012;95(9):4775-4783.

38. Rebecchi A, Pisacane V, Callegari ML, Puglisi E, Morelli L. Ecology of antibiotic resistant coagulase-negative staphylococci isolated from the production chain of a typical Italian salami. Food Control. 2015;53:14-22.

39. Zielińska D, Rzepkowska A, Radawska A, Zieliński K. In vitro screening of selected probiotic properties of Lactobacillus strains isolated from traditional fermented cabbage and cucumber. Curr Microbiol. 2015;70(2):183-194.

40. Nunes RSC, de Souza CP, Pereira KS, Del Aguila EM, Paschoalin VMF. Identification and molecular phylogeny of coagulase-negative staphylococci isolates from Minas Frescal cheese in southeastern Brazil: Superantigenic toxin production and antibiotic resistance. J Dairy Sci. 2016;99(4):2641-2653.

41. Park S, Ji Y, Park H, Lee K, Park H, Beck BR, Shin H, Holzapfel WH. Evaluation of functional properties of lactobacilli isolated from Korean white kimchi. Food Control. 2016;69:5-12.

42. Guo H, Pan L, Li L, Lu J, Kwok L, Menghe B, Zhang H, Zhang W. Characterization of antibiotic resistance genes from Lactobacillus isolated from traditional dairy products. J Food Sci. 2017;82(3):724-730.

43. Erginkaya ZERRIN, Turhan EU, Tatlı D. Determination of antibiotic resistance of lactic acid bacteria isolated from traditional Turkish fermented dairy products. Iranian J Vet Res. 2018;19(1):53.

44. Touret T, Oliveira M, Semedo-Lemsaddek T. Putative probiotic lactic acid bacteria isolated from sauerkraut fermentations. PloS One. 2018;13(9):e0203501.

45. Wang J, Li M, Wang J, Liu M, Yang KU, Zhang JI, Fan M, Wei X. Antibiotic resistance of coagulase-negative staphylococci and lactic acid bacteria isolated from naturally fermented Chinese cured beef. J Food Prot. 2018;81(12):2054-2063.

46. Li Y, Li L, Kromann S, Chen M, Shi L, Meng H. Antibiotic resistance of Lactobacillus spp. and Streptococcus thermophilus isolated from Chinese fermented milk products. Foodborne Pathog Dis. 2019;16(3):221-228.

47. Leech J, Cabrera-Rubio R, Walsh AM, Macori G, Walsh CJ, Barton W, Finnegan L, Crispie F, O'Sullivan O, Claesson MJ, Cotter PD. Fermented-food metagenomics reveals substrate-associated differences in taxonomy and health-associated and antibiotic resistance determinants. mSystems. 2020;5(6):e00522-20.

48. Yasir M, Al-Zahrani IA, Bibi F, Abd El Ghany M, Azhar EI. New insights of bacterial communities in fermented vegetables from shotgun metagenomics and identification of antibiotic resistance genes and probiotic bacteria. Food Res Int. 2022;157:111190.

49. Wang HH, Schaffner DW. Antibiotic resistance: how much do we know and where do we go from here? Appl Environ Microbiol. 2011;77(20):7093-7095.

50. Future Market Insights. Fermented foods and beverages market. Future Market Insights; 2022 [accessed 2022 Aug 30]. Available from: https://www.futuremarketinsights.com/reports/fermented-foods-and-beverages-market.

51. Manskar N, Raskin S. Here's why sauerkraut and kimchi sales surged during the coronavirus. New York Post; 2020 Apr 18 [accessed March 26, 2021]. Available from: https://nypost.com/2020/04/17/heres-why-sauerkraut-and-kimchi-sales-are-surging-thanks-to-coronavirus/.

52. Farmer JJ. *Enterobacteriaceae*: introduction and identification. In: Manual of Clinical Microbiology 8th edition. Vol 1. 2003:636-653.

53. Mahlen SD. *Serratia* infections: from military experiments to current practice. Clin Microbiol Rev. 2011;24(4):755-791.

54. Muñoz MD, Benomar N, Lerma LL, Gálvez A, Abriouel H. Antibiotic resistance of *Lactobacillus pentosus* and *Leuconostoc pseudomesenteroides* isolated from naturally-fermented Aloreña table olives throughout fermentation process. Int J Food Microbiol. 2014;172:110-118.

55. Fraqueza MJ. Antibiotic resistance of lactic acid bacteria isolated from dry-fermented sausages. Int J Food Microbiol. 2015;212:76-88.

56. Kim T, Heo S, Na HE, Lee G, Kim JH, Kwak MS, Sung MH, Jeong DW. Bacterial community of Galchi-Baechu kimchi based on culture-dependent investigations and selection of starter candidates. J Microbiol Biotechnol. 2022;32(3):309-321.

57. Guevarra RB, Magez S, Peeters E, Chung MS, Kim KH, Radwanska M. Comprehensive genomic analysis reveals virulence factors and antibiotic resistance genes in *Pantoea agglomerans* KM1, a potential opportunistic pathogen. PLoS One. 2021;16(1):e0239792.

58. Gao W, Zhu Y, Ye J, Chu H. Gut non-bacterial microbiota contributing to alcohol-associated liver disease. Gut Microbes. 2022;13(1):1984122.

59. Land MH, Rouster-Stevens K, Woods CR, Cannon ML, Cnota J, Shetty AK. Lactobacillus sepsis associated with probiotic therapy. Pediatrics. 2005;115(1):178-181. doi:10.1542/peds.2004-2137.

60. Boyle RJ, Robins-Browne RM, Tang ML. Probiotic use in clinical practice: what are the risks?. Am J Clin Nutr. 2006;83(6):1256-1264.

61. Ohishi A A, Takahashi S, Ito Y, Ohishi Y, Tsukamoto K, Nanba Y, Ito N, Kakiuchi S, Saitoh A, Morotomi M, et al. Bifidobacterium septicemia associated with postoperative probiotic therapy in a neonate with omphalocele. J Pediatr. 2010;156(4):679-681. doi:10.1016/j.jpeds.2009.11.041.

62. Kochan P, Chmielarczyk A, Szymaniak L, Brykczynski M, Galant K, Zych A, Pakosz K, Giedrys-Kalemba S, Lenouvel E, Heczko PB. Lactobacillus rhamnosus administration causes sepsis in a cardiosurgical patient--is the time right to revise probiotic safety guidelines? Clin Microbiol Infect. 2011;17(10):1589-1592. doi:10.1111/j.1469-0691.2011.03614.x

63. Bertelli C, Pillonel T, Torregrossa A, Prod'hom G, Fischer CJ, Greub G, Giannoni E. Bifidobacterium longum bacteremia in preterm infants receiving probiotics. Clin Infect Dis. 2015;60(6):924-927. doi:10.1093/cid/ciu946

64. Angurana SK, Bansal A, Singhi S, Aggarwal R, Jayashree M, Salaria M, Mangat NK. Evaluation of effect of probiotics on cytokine levels in critically ill children with severe sepsis: a double-blind, placebo-controlled trial. Crit Care Med. 2018;46(10):1656-1664.

65. Chiang MC, Chen CL, Feng Y, Chen CC, Lien R, Chiu CH. Lactobacillus rhamnosus sepsis associated with probiotic therapy in an extremely preterm infant: Pathogenesis and a review for clinicians. J Microbiol Immunol Infect. 2021;54(4):575-580. doi:10.1016/j.jmii.2020.03.029.

66. Aydoğan S, Dilli D, Özyazici A, Aydin N, Şimşek H, Orun UA, Aksoy ÖN. Lactobacillus rhamnosus sepsis associated with probiotic therapy in a term infant with congenital heart disease. Fetal Pediatr Pathol. 2022;41(5):823-827. doi:10.1080/15513815.2021.1966144.

67. Kulkarni T, Majarikar S, Deshmukh M, Ananthan A, Balasubramanian H, Keil A, Patole S. Probiotic sepsis in preterm neonates-a systematic review. Eur J Pediatr. 2022;181(6):2249-2262. doi:10.1007/s00431-022-04452-5.

68. Colautti A, Arnoldi M, Comi G, Iacumin L. Antibiotic resistance and virulence factors in lactobacilli: something to carefully consider. Food Microbiol. 2022;103:103934.

69. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics. 2018 Sep 1;34(17):i884-i890.

70. Yin X, Jiang XT, Chai B, Li L, Yang Y, Cole JR, Tiedje JM, Zhang T. ARGs-OAP v2.0 with an expanded SARG database and Hidden Markov Models for enhancement

characterization and quantification of antibiotic resistance genes in environmental metagenomes. Bioinformatics. 2018 Jul 1;34(13):2263-70.

71. Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes A, Henry L, Hester J, et al. Welcome to the Tidyverse. J Open Source Softw. 2019;4(43):1686.

72. Wickham H. Data analysis. In: ggplot2. Springer, Cham; 2016:189-201.

73. Fox J, Weisberg S. An R companion to applied regression. Sage publications; 2018.

74. Patil I. Visualizations with statistical details: The'ggstatsplot'approach. J Open Source Softw. 2021;6(61):3167.

75. Wickham H, François R, Henry L, Müller K. dplyr: A Grammar of Data Manipulation. https://dplyr.tidyverse.org, https://github.com/tidyverse/dplyr; 2022.

76. Stock I, Grueger T, Wiedemann B. Natural antibiotic susceptibility of strains of *Serratia marcescens* and the *S. liquefaciens* complex: S. *liquefaciens sensu stricto*, *S. proteamaculans* and *S. grimesii*. Int J Antimicrob Agents. 2003;22(1):35-47.

77. Wood DE, Lu J, Langmead B. Improved metagenomic analysis with Kraken 2. Genome Biol. 2019;20(1):1-13.

1 **Tables**

2 Table 1. Most abundant AR genes in kimchi samples[1].

Sample #	Rank				
	1	2	3	4	5
K1-4 (BHI, pooled)	Multidrug 0.3189*	Bacitracin 0.0901	Unclassified 0.0780	MLS** 0.0684	Fosfomycin 0.0465
K1-4 (BHI-Tet, pooled)	Multidrug 0.2556	Tetracycline 0.1059	Unclassified 0.0777	Bacitracin 0.0485	MLS 0.0422
K1-4 (BHI-Amp, pooled)	Multidrug 0.3235	Bacitracin 0.1795	β-lactam 0.0680	MLS 0.0557	Fosfomycin 0.0498
K7 (BHI)	Multidrug 1.7085	Unclassified 0.3728	Bacitracin 0.1609	MLS 0.1278	Fosmidomycin 0.1216
K7 (BHI)	unclassified 0.0248	Multidrug 0.0118	Bacitracin 0.0050	MLS 0.0023	fosmidomycin 0.0012
K8 (BHI)	Multidrug 0.0681	Unclassified 0.0469	Bacitracin 0.0136	MLS 0.0047	Tetracycline 0.0006
K9 (BHI)	MLS 0.2557	Multidrug 0.0577	Unclassified 0.0390	Bacitracin 0.0191	Vancomycin 0.0001
K10 (BHI)	Bacitracin 0.0880	Unclassified 0.0669	Multidrug 0.0182	MLS 0.0009	Fosfomycin 0.0005
K11 (BHI)	Unclassified 0.0191	Bacitracin 0.0063	Multidrug 0.0026	Tetracycline 0.0002	MLS* 0.0002

5 *Numerical readings of AR: total ARG copies per 16S rRNA gene.

6 **MLS: Macrolide-Lincosamide-Streptogramin.

1   Table 2. Summary of identified AR colonies from kimchi products purchased in 2022[*].

Isolate	Identity	Food source	Recovered from	Resistance with MIC* (µg/mL)
K5-1	*Latilactobacillus (Lactobacillus) sakei*	Kimchi 5	MRS-tet	Tet (>64)
K5-2	*Bacillus (licheniformis/paralicheni formis)*	Kimchi 5	BHI-ery	Amp (>64) Ery (>16)
K5-3	*Weissella sp.*	Kimchi 5	MRS-ery	Amp (>256) Ery (>16) Tet (>128)
K7-5	*Serratia quinivorans*	Kimchi 7	BHI/BHI-amp	Amp (>64) Van (>1024)
K7-6	*Klebsiella pneumoniae*	Kimchi 7	BHI/BHI-amp	Amp (>256) Van (>1024)
K7-8	*Enterobacter sp.*	Kimchi 7	BHI/BHI-amp	Amp (>256) Van (>1024)
K7-9	*Rahnella aquatilis*	Kimchi 7	BHI/BHI-amp	Amp (>256) Ery (>100) Van(>1024)
K8-6	*Bacillus licheniformis*	Kimchi 8	BHI-ery	Amp (>64) Ery (>16)
K9-1	*Latilactobacillus (Lactobacillus) curvatus*	Kimchi 9	MRS-tet	Tet (>64)
K9-2	*Bacillus (licheniformis/paralicheni formis)*	Kimchi 9	BHI-ery	Amp (>64) Ery (>16)
K9-3	*Latilactobacillus (Lactobacillus) sakei*	Kimchi 9	MRS-ery	Ery (>16)
K10-1	*Levilactobacillus (Lactobacillus) brevis*	Kimchi 10	BHI-tet	Tet (>64) Van (>512)
K10-2	*Levilactobacillus (Lactobacillus) brevis*	Kimchi 10	BHI-tet	Tet (>64) Van (>512)
K11-3	*Rahnella aquatilis*	Kimchi 11	MRS-amp	Amp (>256)
K12-1	*Leuconostoc citreum*	Kimchi 12	BHI-tet	Tet (>128)
K12-6	*Latilactobacillus (Lactobacillus) sakei*	Kimchi 12	BHI-tet	Tet (>32)
K13-1	*Serratia marcescens*	Kimchi 13	BHI-tet	Amp (>256) Ery (>200) Tet (>128) Van (>1024)
K13-2	*Oceanobacillis jeddahense*	Kimchi 13		Tet (>32) Ery (>16)

2   *Tested MIC for representative antibiotics, including vancomycin (Van), ampicillin (Amp),
3   erythromycin (Ery), and tetracycline (Tet).
4

1   Table 3. Top 5 genera classified of microbiota directly of 5 kimchi samples purchased in 2022.

Sample	Top genera	Percentage
Kimchi7	*Leuconostoc*	94.87%
	Other	3.17%
	Rahnella	0.85%
	Weissella	0.48%
	Levilactobacillus	0.36%
	Xanthomonas	0.27%
Kimchi8	*Latilactobacillus*	55.98%
	Leuconostoc	28.90%
	Lactiplantibacillus	7.49%
	Other	5.46%
	Lactococcus	1.59%
	Levilactobacillus	0.58%
Kimchi9	*Latilactobacillus*	81.60%
	Leuconostoc	11.79%
	Other	3.45%
	Weissella	2.02%
	Lactiplantibacillus	0.59%
	Levilactobacillus	0.55%
Kimchi10	*Lactiplantibacillus*	54.60%
	Leuconostoc	32.89%
	Levilactobacillus	9.80%
	Other	1.86%
	Weissella	0.55%
	Latilactobacillus	0.30%
Kimchi11	*Weissella*	58.85%
	Leuconostoc	39.19%
	Other	1.41%
	Latilactobacillus	0.23%
	Lactiplantibacillus	0.19%
	Stenotrophomonas	0.13%

1 Table 4. Most abundant AR genes in BHI plates recovered microbiota of artisan cheese samples[1].

Sample #	Rank				
	1	2	3	4	5
C1-4 (BHI, pooled)	Multidrug 0.4704*	MLS** 0.0783	Bacitracin 0.0353	Fosfomycin 0.0265	Tetracycline 0.0105
C1-4 (BHI-Tet, pooled)	Tetracycline 0.5377	Multidrug 0.2359	Unclassified 0.0876	Bacitracin 0.0744	MLS 0.0295
C1-4 (BHI-Amp, pooled)	Multidrug 0.0308	Tetracycline 0.0093	MLS 0.0084	Bacitracin 0.0053	Unclassified 0.0033
C5 (BHI)	Multidrug 0.1343	Fosfomycin 0.1123	Unclassified 0.0709	β-lactam 0.0617	MLS 0.0555
C5 (BHI-Tet)	Multidrug 0.0033	β-lactam 0.0031	Tetracycline 0.0014	Bacitracin 0.0012	unclassified 0.0012
C5 (BHI-Amp)	Multidrug 0.0040	β-lactam 0.0038	unclassified 0.0015	bacitracin 0.0014	MLS 0.0013
C6 (BHI)	Multidrug 0.6202	Tetracycline 0.4674	Bacitracin 0.2046	Unclassified 0.1750	MLS 0.0694
C7 (BHI)	Multidrug 0.3538	Tetracycline 0.2740	Unclassified 0.1442	Bacitracin 0.1308	MLS 0.0557
C7 (BHI-Tet)	Multidrug 0.2558	Tetracycline 0.2215	Unclassified 0.1183	Bacitracin 0.1176	Quinolone 0.0167
C7 (BHI-Amp)	Multidrug 0.0394	Bacitracin 0.0085	Tetracycline 0.0078	Unclassified 0.0075	β-lactam 0.0035
C8 (BHI)	Multidrug 0.2605	Unclassified 0.0737	Vancomycin 0.0671	Bacitracin 0.0472	MLS 0.0345

3 *Numerical readings of total ARG copies per 16S rRNA gene.

4 **MLS: Macrolide-Lincosamide-Streptogramin.

1   Table 5. Summary of identified AR colonies from artisan cheese products purchased in 2022[*].
2

Isolate	Identity	Food source	Recovered from	Resistance with MIC* (µg/mL)
C5-3	*Staphylococcus equorum*	Cheese 5	BHI-ery	Ery (>16)
C5-9	*Mammaliicoccus vitulinus*	Cheese 5	BHI-ery	Ery (>16)
C6-2	*Staphylococcus xylosus*	Cheese 6	BHI-tet	Tet (>64)
C6-4	*Staphylococcus saprophyticus*	Cheese 6	BHI-tet	Tet (>64)
C6-10	*Staphylococcus xylosus*	Cheese 6	BHI-tet	Tet (>64) Van (>64)
C6-11	*Enterococcus faecalis*	Cheese 6	MRS-ery	Amp (>128) Tet (>128) Ery (>16) Van (>64)
C7-1	*Staphylococcus xylosus*	Cheese 7	BHI-tet	Tet (>32)
C7-5	*Staphylococcus saprophyticus*	Cheese 7	BHI-tet	Tet (>64)
C8-11	*Staphylococcus equorum*	Cheese 8	BHI-ery	Ery (>16) Van (>16)

3   *Tested MIC for representative antibiotics, including vancomycin, ampicillin, erythromycin, and
4   tetracycline.
5

1 **Figures**

2 Fig 1. Impact of 10-week food intervention on the change of antibiotic resistome of fecal
3 microbiota of human subjects in percentage comparing to the baseline. Green: by foods rich in
4 fermented products; pink: by foods rich in plant fibers.

1 Fig 2. Statistical significance analysis of the impact of 10-week food intervention on the change
2 of antibiotic resistome of fecal microbiota of human subjects in percentage comparing to the
3 baseline by diets rich in A) fermented foods; B) plant fibers.

4 A)

$t_{Student}(16) = -2.43, p = 0.03, \hat{g}_{Hedges} = -0.56, CI_{95\%} [-1.05, -0.06], n_{pairs} = 17$

$\log_e(BF_{01}) = -0.87, \hat{\delta}_{difference}^{posterior} = -0.05, CI_{95\%}^{ETI} [-0.10, -1.19e-03], r_{Cauchy}^{JZS} = 0.71$

6 B)

$t_{Student}(14) = -0.07, p = 0.94, \hat{g}_{Hedges} = -0.02, CI_{95\%} [-0.50, 0.46], n_{pairs} = 15$

$\log_e(BF_{01}) = 1.34, \hat{\delta}_{difference}^{posterior} = -1.05e-03, CI_{95\%}^{ETI} [-0.04, 0.04], r_{Cauchy}^{JZS} = 0.71$

1   Fig 3. Antibiotic resistome of pooled traditionally fermented food microbiota recovered from
2   agar plates. A) 4 kimchi samples (K1-K4) purchased in 2021. B) 4 artisan cheeses (C1-C4)
3   purchased in 2021.

5   A).   B).

Fig 4. Resistome assessment of individual fermented foods purchased in 2022. A) resistome of
direct kimchi microbiota of samples; B) resistome of direct kimchi microbiota (left bar) and BHI
recovered microbiota (right bar) of kimchi sample K7; C). resistome of BHI recovered cheese
microbiota.

A).   B).

C).

1   Fig 5. Detected cheese microbiota recovered from BHI agar plates with and without antibiotics.

1   **Supplemental tables**

2   STable 1. Fermented food samples assessed in this study.

Sample #	Description	Year of Purchase	Plate recovered microbiota extraction shotgun	Direct sample extraction shotgun	AR isolate screening	AR isolate illustrated
Kimchi 1 (K1)	From Korean market A	2021	+ (mixed)			
Kimchi 2 (K2)	From Korean restaurant A	2021	+ (mixed)			
Kimchi 3 (K3)	From Japanese restaurant A	2021	+ (mixed)			
Kimchi 4 (K4)	From Korean restaurant B	2021	+ (mixed)			
Kimchi 5 (K5)	From Korean restaurant B	2022			+	+
Kimchi 6 (K6)	Bottled brand A	2022			+	
Kimchi 7 (K7)	From a Korean chain restaurant C	2022	+	+	+	+
Kimchi 8 (K8)	Bottled brand B	2022		+	+	+
Kimchi 9 (K9)	Bottled brand C	2022		+	+	+
Kimchi 10 (K10)	Bagged brand D	2022		+	+	+
Kimchi 11 (K11)	From Japanese market B	2022		+	+	+
Kimchi 12 (K12)	Bottled brand E	2022			+	+
Kimchi 13 (K13)	From Korean Market A	2022			+	+
Cheese 1 (C1)	X(i)* Blue, pasteurized milk (Germany)	2021	+ (mixed)			
Cheese 2 (C2)	X(ii)*, raw cow's milk (Aged > 60 days) (France)	2021	+ (mixed)			
Cheese 3 (C3)	X(iii)* Blue, pasteurized cow and sheep milk (France)	2021	+ (mixed)			
Cheese 4 (C4)	X(iv)* Blue, organic pasteurized milk	2021	+ (mixed)			

	(Unknown)					
Cheese 5 (C5)	C(i)*, pasteurized buffalo milk (Italy)	2022	+		+	+
Cheese 6 (C6)	C(ii)*, raw cow's milk (Aged > 60 days) (United States)	2022	+		+	+
Cheese 7 (C7)	C(iii)*, raw cow's milk (Aged > 60 days) (United States)	2022	+		+	+
Cheese 8 (C8)	C(iv)*, pasteurized part-skim cow's milk (Switzerland)	2022	+		+	+

1   *Cheese products were coded to protect the identities of individual producers.

1   STable 2. Gut antibiotic resistome of subjects with dietary intervention[1].

	Total ARG copies/16S gene				
Subject ID	"Week-02"	Week0	Week08	Week10	Intervention by
8004	0.272186	0.284192	0.300493	0.239926	Fermented foods
8008	0.479228	0.508536	0.404436	0.527567	Fermented foods
8010	0.438165	0.460143	0.377093	0.408678	Fermented foods
8011	0.283978	0.241384		0.345206	Fermented foods
8014		0.367376	0.420609	0.411247	Fermented foods
8016	0.608693	0.498886	0.611231	0.591077	Fermented foods
8020	0.251731	0.287428		0.311607	Fermented foods
8021	0.357738	0.428821	3.77E-01	0.585925	Fermented foods
8024	0.345131	0.372126	0.364187	0.510634	Fermented foods
8025	0.188706	0.322113	0.308409	0.346233	Fermented foods
8026		0.33126	0.515311	0.446267	Fermented foods
8027	0.414694	0.332016	0.455848	0.347911	Fermented foods
8028	0.265774			0.351583	Fermented foods
8030	0.407724	0.514995	0.598983	0.448453	Fermented foods
8032	0.383879	0.416004	0.468486	0.718217	Fermented foods
8033	0.290082	0.272963	0.236222	0.23334	Fermented foods
8034	0.297562	0.364338	0.363455	0.287874	Fermented foods
8001		0.376419	0.316027	0.310167	High plant fibers
8002	0.274209	0.289648	0.264493	0.265522	High plant fibers
8003	0.624062	0.481308	0.447535	0.394988	High plant fibers
8006	0.219233	0.219967	0.233956	0.253023	High plant fibers
8009	0.182759	0.449417	0.415087	0.433871	High plant fibers
8017	0.765387	0.246743		0.342993	High plant fibers
8018	0.259782		0.241797	0.279882	High plant fibers
8022	0.364975		0.534551	0.44805	High plant fibers
8023		0.467738	0.385584	0.507301	High plant fibers
8029	0.397652	0.313326	0.326775	0.281955	High plant fibers
8036	0.381629	0.421726		0.45393	High plant fibers
8037	0.354639	0.320164	0.283361	0.384654	High plant fibers
8038	0.338513	0.269464	0.420975	0.311128	High plant fibers
8039	0.409468	0.552285	0.551152	0.555141	High plant fibers
8041	0.499627	0.501698	0.510558	0.526228	High plant fibers

2   [1]Numerical readings in total ARG copies per 16S gene.

STable 3. Top 5 genera classified of microbiota recovered from BHI agar plates of pooled 4 kimchi samples purchased in 2021.

Sample	Top genera	Percentage
Mix_Kimchi_BHI	*Bacillus*	58.29%
	Serratia	13.42%
	Pseudomonas	12.11%
	Leuconostoc	9.92%
	Other	4.72%
	Lelliottia	1.55%
Mix_Kimchi_BHI_amp	*Pseudomonas*	46.63%
	Bacillus	41.71%
	Brucella	5.92%
	Enterobacter	2.71%
	Other	1.70%
	Rahnella	1.34%
Mix_Kimchi_BHI_tet	*Latilactobacillus*	34.33%
	Bacillus	33.42%
	Serratia	18.88%
	Stenotrophomonas	5.65%
	Lactococcus	4.14%
	Other	3.57%

1 STable 4: Top 5 genera classified of microbiota recovered from BHI agar plates of pooled 4
2 cheese samples purchased in 2021.

Sample	Top genera	Percentage
Mix_Cheese_BHI	*Lactococcus*	47.32%
	Lacticaseibacillus	25.26%
	Staphylococcus	23.19%
	Other	2.04%
	Enterococcus	1.09%
	Streptococcus	1.09%
Mix_Cheese_BHI_tet	*Staphylococcus*	52.84%
	Bacillus	31.53%
	Streptococcus	12.24%
	Microbacterium	1.92%
	Other	1.29%
	Bacteroides	0.17%

STable 5: Top 5 genera classified of BHI-recovered microbiota of individual cheese samples
purchased in 2022.

Sample	Top genera	Percentage
Cheese5-BHI	*Mammaliicoccus*	64.41%
	Staphylococcus	18.62%
	Carnobacterium	5.26%
	Other	4.49%
	Glutamicibacter	3.73%
	Psychrobacter	3.51%
Cheese5-BHI-amp	Other	68.49%
	Stenotrophomonas	9.89%
	Pseudomonas	7.20%
	Burkholderia	5.91%
	Alcanivorax	5.31%
	Streptococcus	3.19%
Cheese5-BHI-tet	Other	64.56%
	Stenotrophomonas	11.03%
	Pseudomonas	7.46%
	Burkholderia	6.17%
	Alcanivorax	5.66%
	Streptococcus	5.14%
Cheese6-BHI	*Staphylococcus*	64.68%
	Enterococcus	22.47%
	Lacticaseibacillus	12.22%
	Other	0.29%
	Streptococcus	0.28%
	Leuconostoc	0.07%
Cheese7-BHI	*Staphylococcus*	73.00%
	Lactococcus	26.05%
	Leuconostoc	0.40%
	Other	0.32%
	Lacticaseibacillus	0.13%
	Streptococcus	0.11%
Cheese7-BHI-amp	Other	50.11%
	Pseudomonas	30.32%
	Burkholderia	5.63%
	Staphylococcus	5.18%
	Streptococcus	5.16%
	Leuconostoc	3.60%
Cheese7-BHI-tet	*Staphylococcus*	99.71%
	Other	0.10%
	Enterococcus	0.09%
	Leuconostoc	0.05%

	Mammaliicoccus	0.03%
	Lactococcus	0.02%
Cheese8-BHI	*Enterococcus*	44.89%
	Tetragenococcus	16.18%
	Staphylococcus	13.08%
	Lactococcus	12.68%
	Other	11.26%
	Carnobacterium	1.91%