Myocardial IGF2R is a critical mediator of inflammation and fibrosis after ischemia-reperfusion injury

Zhichao Wu[1], Wei Huang[1], Xingyu He[1], Suchandrima Dutta[2], Christian Paul[1], Guo-Chang Fan[3], Onur Kanisicak[1], Meifeng Xu[1], Jialiang Liang[1,*], Yigang Wang[1,*]

1 Department of Pathology and Laboratory Medicine, College of Medicine, University of Cincinnati, Cincinnati, OH, USA, 45267

2 Department of Internal Medicine, College of Medicine, University of Cincinnati, Cincinnati, OH 45267

3 Department of Pharmacology and Systems Physiology, College of Medicine, University of Cincinnati, Cincinnati, OH, USA, 45267

* Correspondence to: Jialiang Liang (liangjl@ucmail.uc.edu), Department of Pathology and Laboratory Medicine, College of Medicine, University of Cincinnati, 231 Albert Sabin Way, Cincinnati, Ohio, 45267, USA. Yigang Wang (yigang.wang@uc.edu), Department of Pathology and Laboratory Medicine, College of Medicine, University of Cincinnati, 231 Albert Sabin Way, Cincinnati, Ohio, 45267, USA.

ABSTRACT

Ischemia-reperfusion (I/R) injury is a common occurrence in various surgical procedures used to treat heart diseases. However, the role of insulin-like growth factor 2 receptor (IGF2R) during the process of myocardial I/R remains unclear. Therefore, this study aims to investigate the expression, distribution, and functionality of IGF2R in various I/R-associated models (such as reoxygenation, revascularization, and heart transplant). Loss-of-function studies (including myocardial conditional knockout and CRISPR interference) were performed to clarify the role of IGF2R in I/R injuries. Following hypoxia, IGF2R expression increased, but this effect was reversed upon restoration of oxygen levels. Loss of myocardial IGF2R was found to enhance the cardiac contractile functions, and reduced cell infiltration or cardiac fibrosis of I/R mouse models compared to the genotype control. CRISPR-inhibition of IGF2R decreased cell apoptotic death under hypoxia. RNA sequencing analysis indicated that myocardial IGF2R played a critical role in regulating the inflammatory response, innate immune response, and apoptotic process following I/R. Integrated analysis of the mRNA profiling, pulldown assays, and mass spectrometry identified granulocyte-specific factors as potential targets of myocardial IGF2R in the injured heart. In conclusion, myocardial IGF2R emerges as a promising therapeutic target to ameliorate inflammation or fibrosis following I/R injuries.

INTRODUCTION

Ischemia-reperfusion (I/R) injury is a well-known phenomenon that occurs in various surgical procedures to treat heart diseases, such as percutaneous coronary intervention for acute myocardial infarction, heart transplantation, and coronary artery bypass grafting.[1] While several pharmacological interventions have been investigated to improve I/R injury, no single therapy can effectively target the multiple pathways involved in this complex process.[2] Eventually, the activation of irreversible cell death pathways after I/R can contribute to the development of heart failure. Understanding the molecular and cellular mechanisms underlying I/R injury is critical for refining the existing surgical approaches to prevent or mitigate their adverse effects on the heart.

Currently, most of the drugs being evaluated for their cardioprotective effects are derived from the cardioprotective signal transduction pathways.[3, 4] The activation of prosurvival pathways (such as PI3K/AKT, JAK/STAT, and ERK) modulated by the insulin-like growth factor (IGF) system is essential for cardiac homeostasis including cell growth or death and regulation of energy metabolism.[5-7] Previous

studies have established the homology between insulin, IGF1, and IGF2, as well as

elucidated the interactions among the insulin receptor (INSR), IGF1 receptor (IGF1R),

and IGF2 receptor (IGF2R).[8-10] As the administration of insulin, IGF1, IGF2, and

other growth factors gains increasing interest as a potential treatment for myocardial

I/R,[11-13] investigating the receptors of these factors could provide valuable insights

into the role of the IGF system in cardioprotection.

It is well-known that canonical downstream of IGF1R such as PI3K/AKT can

maintain outer mitochondrial membrane potential and inhibit glycogen synthase

kinase-3β, thereby protecting cells against necrosis triggered by I/R.[14-16] Recent

research has shown that inhibiting IGF2R signaling promotes cardiomyocyte

proliferation and reduces hypoxia- or TNFα-induced cell death,[17-19] while the role of

IGF2R signaling in I/R remains largely unknown. With broad expression across many

cell types, IGF2R is a versatile receptor that can bind multiple ligands (including

IGF2 and M6P-modified proteins) and plays a critical role in intracellular transport,

shuttling between the trans-Golgi network, endosomes, and plasma membrane.[20, 21]

Therefore, disruption of the Golgi-to-lysosome transport by loss of IGF2R can lead to

reduced lysosomal activity, resulting in the accumulation of misfolded proteins and

increased production of reactive oxygen species due to dysfunction in both autophagy

and mitophagy pathways.[22, 23] Given its multifunctional nature, the effects of IGF2R

on cell viability appear to be cell type-specific and context-dependent under stress

conditions. In this study, the loss-of-function models were used to investigate the role

of IGF2R in myocardial I/R injuries.

RESULTS

Dynamic expression of IGF2R during myocardial I/R

The expression of IGF2R in mouse hearts exhibited an initial increase

following myocardial ischemia (MI) and reperfusion as compared to the sham

operation but subsequently decreased to baseline levels after a 6-hour reperfusion

period (**Fig.1A**). In addition to cardiomyocytes, IGF2R was expressed in non-CM

cells located in the interstitial space of reperfused heart (**Fig.1B**). The single-cell

RNA-seq dataset (Tabula Muris)[24] also provides evidence that IGF2R is expressed

constitutively across multiple cell types within the heart (**Fig.1C**). To investigate the

precise role of myocardial IGF2R in I/R, we generated conditional knockout (cKO)

models by crossing Cre-inducible *Igf2r*[fl/fl] mice with *Myh6*[MerCreMer] mice and treating

the resultant offspring with tamoxifen (**Fig.1D**), while *Igf2r^{fl/fl}* litters were used as the

control group (Ctrl). IGF2R gene expression was significantly decreased in cKO as

confirmed by PCR and western blotting (**Fig.1E-F**).

Loss of myocardial IGF2R improves heart function of I/R models

Subsequently, two different types of IR injury were employed to assess the

effect of cKO on heart function. In a form of warm I/R (**Fig.2A**), the heart function of

Ctrl or cKO mice was analyzed by echocardiography after LAD ligation surgery. At 1

week after I/R, there was no difference in the ejection fraction (EF) between Ctrl and

cKO (**Fig.2B**). The heart function of cKO mice demonstrated an improvement at 4

weeks post-I/R when compared to Ctrl mice, with increases in EF, fractional

shortening, and interventricular septal (**Fig.2C-D**). In addition, heterotopic cardiac

transplant models were established to mimic cold I/R injury (**Fig.2E**). Following

transplantation within 12-hour preservation in cold HTK solutions, there was no

difference in re-beating time between the control group and the cKO donor hearts

(**Fig. 2F**). Although the re-beating time of cKO hearts was shorter than that of the

control group after 24-hour preservation in cold HTK solutions, both grafts had a

dramatically decreased median survival time (**Fig. 2G**).

All Ctrl and cKO donor hearts survived after 6-hour preservation without a

significant difference, but ~60% of cKO hearts exhibited a significantly longer

median survival time than Ctrl hearts after 12-hour preservation (**Fig.2G**), suggesting

a potential protective effect of cKO against ischemic injury. Therefore, on day one

post-surgery, a catheter-based approach was used to evaluate the left ventricular (LV)

pressure and contractility of Ctrl or cKO hearts that underwent 12-hour preservation

(**Fig.2H-I**). The LV systolic pressure and heart rate were increased in cKO donor

hearts as compared with the control, while there was no significant difference in the

LV end-diastolic pressures. The +dp/dt max and the -dp/dt max were rapidly raised in

cKO donor hearts, reflecting an improved contractile function. Furthermore,

recipients with cKO hearts demonstrated a significant decrease in serum levels of

intracellular enzymes, including creatine kinase and lactate dehydrogenase, when

compared to the control group (**Fig.2J**).

Deletion of myocardial IGF2R alters transcriptional profiles in response to I/R

RNA sequencing (RNA-seq) was performed to analyze the transcriptome of

cKO and Ctrl hearts at baseline and after 12-hour preservation and transplantation

(I/R). Principal component analysis of the normalized datasets showed that the

transcriptional profiles of the uninjured cKO and Ctrl hearts were similar, while the

cKO hearts exhibited distinct expression patterns compared to the other samples

following I/R (**Fig.3A**). To gain a deeper understanding of the underlying mechanism

of IGF2R, we analyzed the differentially expressed genes in cKO and Ctrl hearts

subjected to the I/R condition.

We identified 3,909 significant genes (*padj*<0.05) differentially expressed in

cKO and Ctrl hearts following I/R, and used hierarchical clustering analysis to group

these genes into distinct clusters (**Fig.3B**). After I/R, clusters a and b showed up-

regulated gene expression in cKO hearts, while clusters c and d exhibited up-

regulated gene expression in Ctrl hearts. At baseline, both cKO and Ctrl hearts

displayed similar patterns of up-regulated gene expression in clusters b and c, and

down-regulated gene expression in clusters a and d. Gene ontology (GO) analysis was

used to identify the over-representation of GO terms (biological processes) in the gene

lists of different clusters (**Fig.3C**). Genes involved in DNA repair, cell cycle

regulation, and autophagy were predominantly represented in clusters a and b, while

clusters c and d were enriched for genes related to inflammatory response, innate

immune response, and apoptotic process. The top 20 GO terms showed that the cKO

of IGF2R significantly attenuated innate immune responses, including cytokine

production, leukocyte migration, and cell adhesion, while promoting cell cycle

progression and self-healing processes. (**Fig.3D**).

We also performed Gene Set Enrichment Analysis (GSEA) on a ranked list of

significant genes to identify enriched gene sets and biological pathways associated

with the observed gene expression changes (**Fig.3E**). The results of the gene set

distribution and enrichment score using GSEA confirmed that the Ctrl hearts were

enriched for GO terms related to inflammatory response, innate immune response,

and chemotaxis, whereas the cKO hearts showed enrichment for the mitotic cell cycle

process except for autophagy. The cKO of IGF2R resulted in the activation of genes

involved in cell growth pathways, which supports the established concept that IGF2R

functions as a growth-inhibitory gene.[21, 25].

Identification of IGF2R-associated proteins from reperfused myocardium

Although the innate immune response is known to play a critical role in the

pathogenesis of myocardial I/R injury,[26, 27] the precise mechanism by which IGF2R

participates in this process is still unknown. Therefore, we attempted to identify the

IGF2R targets by proteomic analysis of lysed tissues obtained from heart grafts using

pull-down assays on day 1 post-I/R (**Fig.4A**). To this end, we selectively excised

protein bands from the control sample, which were missing from the cKO sample at

equivalent molecular weights, and performed mass spectrometry analysis. The lists of

identified proteins were compared with the RNA-seq results of the down-regulated

(log2fc<-1, padj<0.05) or up-regulated (log2fc>1, padj<0.05) genes (**Fig.4B**). Notably,

the consistent down-regulation of *Actb*, *Ngp*, *Chil3 (Chi3l3)*, *S100a9*, and *Ltf* gene in

cKO heart tissue at both transcriptomic and translational levels highlighted their

potential role as therapeutic targets of IGF2R. The single-cell RNA-seq dataset

(Tabula Muris)[24] also reveals that these genes are expressed in granulocytes,

granulocytopoietic cells, and monocytes in bone marrow, but not in the native heart

(**Fig.4C**). At 4 weeks post I/R injury, the Ctrl heart exhibited a substantial increase in

cell infiltration with visible nuclei, while the cKO heart demonstrated a significant

reduction in cell infiltration (**Fig.4D**). Furthermore, Masson's trichrome staining

showed that the fibrosis area in the cKO heart was smaller than that in the Ctrl heart

(**Fig.4E**).

Hypoxia-induced activation of IGF2R leads to apoptotic cell death

IGF2R was expressed in H9C2 cardiomyoblasts, and *in vitro* model of

hypoxia-reoxygenation was used to mimic I/R injury. As compared to the normoxic

condition, the IGF2R protein level was increased after hypoxia for 3 hours but slightly

decreased after reoxygenation (**Fig.5A-B**). It is known that IGF2R shuttles constantly

between intracellular organelles (~90% of the total IGF2R protein) and the cell

surface membrane.[9, 25] Labeling with wheat germ agglutinin indicated that IGF2R was

primarily localized in the intracellular organelles of H9C2 cells (**Fig.5C**).

Interestingly, under hypoxia, the intracellular IGF2R appeared to diffuse from the

perinuclear region toward the plasma membrane, resulting in a dispersed cytoplasmic

distribution. Following hypoxia-reoxygenation, the immunoblot analysis of protein

fractions revealed that IGF2R was highly expressed in the intracellular membrane

fraction, indicating membrane attachment rather than cytoplasmic solubility (**Fig.5D**).

A doxycycline (Dox)-inducible CRISPR-interference (CRISPRi) system

(pHAGE-TRE-dCas9-KRAB)[28] was established to determine whether the increased

IGF2R protein level was due to its promoter transcription activation (**Fig.5E**). The

IGF2R protein expression in H9C2 cells was decreased by Dox-induced CRISPRi
under normal or hypoxia-reoxygenation conditions (**Fig.5F**). Hypoxia caused a
significant increase in the level of cleaved caspase3 (active form), which was
attenuated by Dox treatment. Although the cleaved caspase3 level progressively
decreased after reoxygenation (to the normoxic level), CRISPRi of IGF2R did not
affect cell death during this period.

DISCUSSION

193   Although IGF2R is known to be multifaceted, its role in the myocardial I/R
response remains poorly characterized. Here, we investigated the role of IGF2R in
different models of myocardial I/R injuries. We observed the dynamic expression of
IGF2R during myocardial I/R and found that cardiac-specific depletion of IGF2R
improved heart dysfunction and reduced cell death. Mechanistic studies revealed that
IGF2R plays a role in the inflammatory response and fibrosis by interacting with
immune cell-derived proteins. Our findings suggest that IGF2R could be an important
therapeutic target for I/R injury and warrant further exploration of its function in
cardiac myocytes.

202   Our results showed that IGF2R expression was upregulated in the ischemic
heart but returned to baseline levels following reperfusion. The dynamic expression of
IGF2R was also observed in renal macrophages during the I/R processes.[29] IGF2R is
expressed widely in both cardiac and non-cardiac cells, but whether its expression is
consistent or variable across different cell types in response to injuries remains to be
determined. To this end, single-cell RNA sequencing can be used to characterize the
expression of IGF2R across different cell types and heart disease models in future
studies. Further research is also needed to fully elucidate the potential transcriptional
or epigenetic mechanisms (such as DNA methylation, histone modifications,
microRNAs, long non-coding RNAs, and various transcription factors)[30-34] controlling
IGF2R expression in the context of I/R.

213   Our loss-of-function studies provided novel evidence that myocardial IGF2R
played a role in the immune responses following reperfusion or heart transplantation.
Depleting cardiac IGF2R expression resulted in a reduction of cell infiltration and
fibrosis in the heart tissue after reperfusion injury. The integrated analysis of the
transcriptome and proteome highlighted Ngp, Chil3, S100a9, and Ltf as potential
targets of IGF2R. Given the selective expression of these genes in bone marrow

granulocytes but not in normal cardiac cells, we reasoned that IGF2R could be a key

player in regulating the interplay between cardiomyocytes and granulocytes after

myocardial damage. Immune cell infiltration is a common response to stress signals in

the myocardium and is observed across various types of myocardial injuries.[35]

Neutrophils and other granulocytes can rapidly and massively infiltrate the site of

injury and release reactive oxygen species, proteolytic enzymes, and other cytotoxic

molecules that can cause further damage to the myocardium and contribute to the

development of cardiac fibrosis.[36, 37] Ngp, Chil3, S100a9, and Ltf are among several

proteins secreted by granulocytes that have been implicated in a variety of

pathological processes underlying heart disease, including inflammation, immune

response, and fibrosis.[38-42] Therefore, evidence suggests that IGF2R-mediated protein

capture may lead to activation of inflammation pathways in cardiomyocytes and

subsequent impairment of heart function following I/R injury. However, further

investigation is needed to elucidate the precise mechanisms involved.

Furthermore, our findings indicated that myoblasts subjected to hypoxia

experienced an increase in IGF2R expression, which was reversed upon

reoxygenation. This pattern was consistent with the expression trend of IGF2R

observed in heart tissue during the I/R processes. It is possible that the regulation of

IGF2R expression during hypoxia and reoxygenation is mediated by different

transcription factors or signaling pathways. However, previous studies investigating

the impact of hypoxia on the expression of the IGF2R gene have yielded conflicting

results. Some studies have shown increased expression of IGF2R during hypoxia in

myoblast cells,[17, 18] while others have found that hypoxia did not affect IGF2R

expression in ectoplacental cones.[43] Therefore, the effects of hypoxia on IGF2R

expression appear to be cell type-specific and involve multiple signaling pathways.

Aside from its overall gene expression level, the functionality of IGF2R is also

dependent on its intracellular localization.[44] The results of our study demonstrated that

hypoxia enhanced the intracellular trafficking of IGF2R from the perinuclear region

to the plasma membrane, and this effect was primarily attributed to membrane

attachment rather than cytoplasmic solubility.

To assess the therapeutic implications of specifically targeting IGF2R

expression, we utilized CRISPRi complex through the Dox-controlled transactivator

system to block the IGF2R promoter region in myoblasts. By enabling targeted gene

regulation, the CRISPRi technique offers remarkable specificity, versatility, and

reversibility, and presents a promising approach for treating a diverse range of diseases.[45, 46] IGF2R expression was significantly eliminated by CRISPRi which was found to attenuate cell death under hypoxic conditions, yet the molecular pathways involved in this protective effect remain to be determined. Future studies using I/R animal models will be also needed to determine the potential therapeutic effect of this approach.

In summary, this study investigates the expression and function of IGF2R in various models of myocardial I/R injury. We observed dynamic changes in IGF2R expression and intracellular distribution in response to ischemic or hypoxic conditions. Deletion of IGF2R resulted in improved heart function and reduced cell death, indicating its involvement in myocardial I/R injury. RNA sequencing analysis revealed that IGF2R deletion altered transcriptional profiles in response to I/R, attenuating innate immune responses while promoting cell cycle progression and self-healing processes. Proteomic analysis identified granulocyte-specific factors as potential targets of IGF2R binding. Taken together, these findings suggest that myocardial IGF2R plays an essential role in I/R injury and that its deletion may offer a cardioprotective effect against inflammation or fibrosis. This study contributes to our understanding of the pathogenesis of myocardial I/R injury and identifies IGF2R as a potential therapeutic target for ischemic heart disease.

ACKNOWLEDGMENTS

This work was supported by the American Heart Association award 22TPA968781 (J.L.) and the National Institutes of Health (NIH) grants including R01HL143490 and R01HL157456 (Y.W.). We would like to thank Dr. Randy Jirtle and Dr. David Skaar (North Carolina State University) for providing the *Igf2r*$^{fl/fl}$ mice.

DECLARATION OF INTERESTS

The authors declare no competing interests.

SUPPLEMENTAL METHODS

All research protocols conformed to the Guidelines for the Care and Use of Laboratory Animals published by the National Institutes of Health (National Academies Press, eighth edition, 2011). All animal use protocols and experiments conducted in this study were approved and oversight by the University of Cincinnati Animal Care and Use Committee. Detailed methods are available in the supplemental information.

REFERENCES

1. Eltzschig HK, Eckle T. Ischemia and reperfusion--from mechanism to translation. *Nat Med*. 2011;17:1391-1401

2. Ibanez B, Heusch G, Ovize M, Van de Werf F. Evolving therapies for myocardial ischemia/reperfusion injury. *J Am Coll Cardiol*. 2015;65:1454-1471

3. Kleinbongard P, Heusch G. Extracellular signalling molecules in the ischaemic/reperfused heart - druggable and translatable for cardioprotection? *Br J Pharmacol*. 2015;172:2010-2025

4. Heusch G. Myocardial ischaemia-reperfusion injury and cardioprotection in perspective. *Nat Rev Cardiol*. 2020;17:773-789

5. Laustsen PG, Russell SJ, Cui L, Entingh-Pearsall A, Holzenberger M, Liao R, Kahn CR. Essential role of insulin and insulin-like growth factor 1 receptor signaling in cardiac development and function. *Mol Cell Biol*. 2007;27:1649-1664

6. Guo CA, Guo S. Insulin receptor substrate signaling controls cardiac energy metabolism and heart failure. *The Journal of endocrinology*. 2017;233:R131-r143

7. O'Sullivan JF, Leblond AL, Kelly G, Kumar AH, Metharom P, Buneker CK, Alizadeh-Vikali N, Hristova I, Hynes BG, O'Connor R, Caplice NM. Potent long-term cardioprotective effects of single low-dose insulin-like growth factor-1 treatment postmyocardial infarction. *Circulation. Cardiovascular interventions*. 2011;4:327-335

8. Ludwig T, Eggenschwiler J, Fisher P, D'Ercole AJ, Davenport ML, Efstratiadis A. Mouse mutants lacking the type 2 igf receptor (igf2r) are rescued from perinatal lethality in igf2 and igf1r null backgrounds. *Dev Biol*. 1996;177:517-535

9. El-Shewy HM, Luttrell LM. Insulin-like growth factor-2/mannose-6 phosphate receptors. *Vitamins and hormones*. 2009;80:667-697

10. De Meyts P, Whittaker J. Structural biology of insulin and igf1 receptors: Implications for drug design. *Nat Rev Drug Discov*. 2002;1:769-783

11. Salminen PR, Dahle GO, Moen CA, Wergeland A, Jonassen AK, Haaverstad R, Matre K, Grong K. Reperfusion therapy with low-dose insulin or insulin-like growth factor 2; myocardial function and infarct size in a porcine model

of ischaemia and reperfusion. *Basic & clinical pharmacology & toxicology*.

2014;115:438-447

12.   Zhang HX, Zang YM, Huo JH, Liang SJ, Zhang HF, Wang YM, Fan Q, Guo

WY, Wang HC, Gao F. Physiologically tolerable insulin reduces myocardial

injury and improves cardiac functional recovery in myocardial

ischemic/reperfused dogs. *J Cardiovasc Pharmacol*. 2006;48:306-313

13.   White SJ, Chong JJH. Growth factor therapy for cardiac repair: An overview

of recent advances and future directions. *Biophys Rev*. 2020;12:805-815

14.   Song HP, Zhang L, Dang YM, Yan H, Chu ZG, Huang YS. The

phosphatidylinositol 3-kinase-akt pathway protects cardiomyocytes from

ischaemic and hypoxic apoptosis via mitochondrial function. *Clinical and*

experimental pharmacology & physiology. 2010;37:598-604

15.   Gomez L, Paillard M, Thibault H, Derumeaux G, Ovize M. Inhibition of

gsk3beta by postconditioning is required to prevent opening of the

mitochondrial permeability transition pore during reperfusion. *Circulation*.

2008;117:2761-2768

16.   Yamashita K, Kajstura J, Discher DJ, Wasserlauf BJ, Bishopric NH, Anversa

P, Webster KA. Reperfusion-activated akt kinase prevents apoptosis in

transgenic mouse hearts overexpressing insulin-like growth factor-1. *Circ Res*.

2001;88:609-614

17.   Chen WK, Kuo WW, Hsieh DJ, Chang HN, Pai PY, Lin KH, Pan LF, Ho TJ,

Viswanadha VP, Huang CY. Creb negatively regulates igf2r gene expression

and downstream pathways to inhibit hypoxia-induced h9c2 cardiomyoblast

cell death. *International journal of molecular sciences*. 2015;16:27921-27930

18.   Lin KH, Kuo CH, Kuo WW, Ho TJ, Pai P, Chen WK, Pan LF, Wang CC,

Padma VV, Huang CY. Nfil3 suppresses hypoxia-induced apoptotic cell death

by targeting the insulin-like growth factor 2 receptor. *J Cell Biochem*.

2015;116:1113-1120

19.   Chen Z, Ge Y, Kang JX. Down-regulation of the m6p/igf-ii receptor increases

cell proliferation and reduces apoptosis in neonatal rat cardiac myocytes. *BMC*

cell biology. 2004;5:15

20.   Hille-Rehfeld A. Mannose 6-phosphate receptors in sorting and transport of

lysosomal enzymes. *Biochim Biophys Acta*. 1995;1241:177-194

21. Ghosh P, Dahms NM, Kornfeld S. Mannose 6-phosphate receptors: New
twists in the tale. *Nat Rev Mol Cell Biol*. 2003;4:202-212

22. Takeda T, Komatsu M, Chiwaki F, Komatsuzaki R, Nakamura K, Tsuji K,
Kobayashi Y, Tominaga E, Ono M, Banno K, Aoki D, Sasaki H. Upregulation
of igf2r evades lysosomal dysfunction-induced apoptosis of cervical cancer
cells via transport of cathepsins. *Cell Death Dis*. 2019;10:876

23. Pluvinage JV, Sun J, Claes C, Flynn RA, Haney MS, Iram T, Meng X,
Lindemann R, Riley NM, Danhash E, Chadarevian JP, Tapp E, Gate D,
Kondapavulur S, Cobos I, Chetty S, Pasca AM, Pasca SP, Berry-Kravis E,
Bertozzi CR, Blurton-Jones M, Wyss-Coray T. The cd22-igf2r interaction is a
therapeutic target for microglial lysosome dysfunction in niemann-pick type c.
Sci Transl Med. 2021;13:eabg2919

24. Tabula Muris C, Overall c, Logistical c, Organ c, processing, Library p,
sequencing, Computational data a, Cell type a, Writing g, Supplemental text
writing g, Principal i. Single-cell transcriptomics of 20 mouse organs creates a
tabula muris. *Nature*. 2018;562:367-372

25. Brown J, Jones EY, Forbes BE. Interactions of igf-ii with the igf2r/cation-
independent mannose-6-phosphate receptor mechanism and biological
outcomes. *Vitamins and hormones*. 2009;80:699-719

26. Timmers L, Pasterkamp G, de Hoog VC, Arslan F, Appelman Y, de Kleijn DP.
The innate immune response in reperfused myocardium. *Cardiovasc Res*.
2012;94:276-283

27. Arslan F, de Kleijn DP, Pasterkamp G. Innate immune signaling in cardiac
ischemia. *Nat Rev Cardiol*. 2011;8:292-300

28. Kearns NA, Genga RM, Enuameh MS, Garber M, Wolfe SA, Maehr R. Cas9
effector-mediated regulation of transcription and differentiation in human
pluripotent stem cells. *Development*. 2014;141:219-223

29. Xie X, Yang X, Wu J, Ma J, Wei W, Fei X, Wang M. Trib1 contributes to
recovery from ischemia/reperfusion-induced acute kidney injury by regulating
the polarization of renal macrophages. *Frontiers in immunology*. 2020;11:473

30. Yang Y, Li T, Vu TH, Ulaner GA, Hu JF, Hoffman AR. The histone code
regulating expression of the imprinted mouse igf2r gene. *Endocrinology*.
2003;144:5658-5670

31.   Latos PA, Pauler FM, Koerner MV, Senergin HB, Hudson QJ, Stocsits RR,
Allhoff W, Stricker SH, Klement RM, Warczok KE, Aumayr K, Pasierbek P,
Barlow DP. Airn transcriptional overlap, but not its lncrna products, induces
imprinted igf2r silencing. *Science*. 2012;338:1469-1472

32.   Szentirmay MN, Yang HX, Pawar SA, Vinson C, Sawadogo M. The igf2
receptor is a usf2-specific target in nontumorigenic mammary epithelial cells
but not in breast cancer cells. *J Biol Chem*. 2003;278:37231-37240

33.   Zhang Y, Zhang Y, Xiao L, Yu TX, Li JZ, Rao JN, Turner DJ, Gorospe M,
Wang JY. Cooperative repression of insulin-like growth factor type 2 receptor
translation by microrna 195 and rna-binding protein cugbp1. *Mol Cell Biol*.
2017;37

34.   Young LE, Fernandes K, McEvoy TG, Butterwith SC, Gutierrez CG, Carolan
C, Broadbent PJ, Robinson JJ, Wilmut I, Sinclair KD. Epigenetic change in
igf2r is associated with fetal overgrowth after sheep embryo culture. *Nat
Genet*. 2001;27:153-154

35.   Swirski FK, Nahrendorf M. Cardioimmunology: The immune system in
cardiac homeostasis and disease. *Nat Rev Immunol*. 2018;18:733-744

36.   Vafadarnejad E, Rizzo G, Krampert L, Arampatzi P, Arias-Loza AP, Nazzal Y,
Rizakou A, Knochenhauer T, Bandi SR, Nugroho VA, Schulz DJJ, Roesch M,
Alayrac P, Vilar J, Silvestre JS, Zernecke A, Saliba AE, Cochain C. Dynamics
of cardiac neutrophil diversity in murine myocardial infarction. *Circ Res*.
2020;127:e232-e249

37.   Puhl SL, Steffens S. Neutrophils in post-myocardial infarction inflammation:
Damage vs. Resolution? *Frontiers in cardiovascular medicine*. 2019;6:25

38.   Marinkovic G, Koenis DS, de Camp L, Jablonowski R, Graber N, de Waard V,
de Vries CJ, Goncalves I, Nilsson J, Jovinge S, Schiopu A. S100a9 links
inflammation and repair in myocardial infarction. *Circ Res*. 2020;127:664-676

39.   Lee SY, Lee CM, Ma B, Kamle S, Elias JA, Zhou Y, Lee CG. Targeting
chitinase 1 and chitinase 3-like 1 as novel therapeutic strategy of pulmonary
fibrosis. *Frontiers in pharmacology*. 2022;13:826471

40.   Kruzel ML, Zimecki M, Actor JK. Lactoferrin in a context of inflammation-
induced pathology. *Frontiers in immunology*. 2017;8

41.   Frodermann V, Nahrendorf M. Neutrophil–macrophage cross-talk in acute
myocardial infarction. *European Heart Journal*. 2016;38:198-200

42.   Othman A, Sekheri M, Filep JG. Roles of neutrophil granule proteins in

orchestrating inflammation and immunity. *FEBS J*. 2022;289:3932-3953

43.   Pringle KG, Kind KL, Thompson JG, Roberts CT. Complex interactions

between hypoxia inducible factors, insulin-like growth factor-ii and oxygen in

early murine trophoblasts. *Placenta*. 2007;28:1147-1157

44.   Kang JX, Bell J, Leaf A, Beard RL, Chandraratna RA. Retinoic acid alters the

intracellular trafficking of the mannose-6-phosphate/insulin-like growth factor

ii receptor and lysosomal enzymes. *Proc Natl Acad Sci U S A*. 1998;95:13687-

13691

45.   Chavez M, Chen X, Finn PB, Qi LS. Advances in crispr therapeutics. *Nature*

reviews. Nephrology. 2023;19:9-22

46.   Musunuru K. Crispr and cardiovascular diseases. *Cardiovasc Res*.

2023;119:79-93

FIGURE LEGENDS

Fig.1 IGF2R expression in ischemic or reperfused hearts. (**A**) Immunoblotting of

IGF2R and GAPDH (loading control) in mice after 45min myocardial ischemia (MI)

induced by left anterior descending artery ligation or releasing ligature (I/R, ischemia-

reperfusion) for 1 to 6 hours. (**B**) Immunostaining of IGF2R and cTnT in the ischemic

heart reperfused for 1 hour. Scale bar, 50μm. Arrow indicates noncardiac cells. (**C**)

Single-cell profiling of IGF2R expression in normal mouse heart (reanalysis of

GSM2967050). Cell phenotype annotation: 1-cardiac muscle cell; 2-fibroblast; 3-

endothelial cell; 4-Endocardial cell; 5-cardiac neuron; 6-leukocyte; 7-myofibroblast;

8-smooth muscle cell. (**D**) Schematic of transgenic mouse models for conditional

knockout (cKO) of myocardial IGF2R. (**E**) Expression of IGF2R mRNA in the Ctrl

(Igf2r$^{fl/fl}$) or cKO (*Myh6^{MerCreMer};Igf2r$^{fl/fl}$*) hearts after the tamoxifen (TAM) diet for 4

weeks. (**F**) Expression of IGF2R protein in the Ctrl or cKO hearts after the TAM diet

for 4 weeks.

Fig.2 IGF2R-cKO enhances cardiac contractile functions. (**A**) Schematic of the

warm I/R model. Myocardial ischemia (MI) induced by left anterior descending artery

(LAD) ligation for 45min and reperfusion by releasing the ligature. (**B**) Heart function

of control and cKO mice assessed by echocardiography in 1 week after the warm I/R.

(**C-D**) Heart function of control and cKO mice assessed by echocardiography in 4

weeks after the warm I/R. EF-ejection fraction; FS-fractional shortening; IVSd-

interventricular septal end-diastole; IVSs-interventricular septal end-systole. (**E**)

Schematic of the cold I/R model. Collecting and perfusing donor hearts with 4℃ HTK

solution and keeping on ice until transplant into the abdomen of syngeneic recipients.

(**F**) The rebeating time (after anastomosis) for ventricular contraction restoration of

control or cKO heart grafts (preserved for 6, 12, or 24 hours in cold HTK). (**G**)

Survival curves of control or cKO heart grafts after transplant. The complete cessation

of a heartbeat is the endpoint. (**H-I**) Heart rate, pressure difference (PD), left ventricle

systolic pressure (LVSP), left ventricle end-diastolic pressure (LVEDP), maximum

dp/dt in the systolic period, and maximum dp/dt in the diastolic period of control or

cKO heart grafts (preserved for 12 hours in HTK) on day 1 post-transplant. (**J**) The

serum creatine kinase (CK) or lactate dehydrogenase (LDH) levels of control or cKO

heart grafts (preserved for 12 hours in HTK) on day 1 post-transplant.

Fig.3 Transcriptional signatures changed by IGF2R-cKO. (**A**) Principal

component (PC) analysis of the global gene profiling in the control or cKO heart at

normal or I/R (12-hour cold HTK and transplant reperfusion for 12 hours) conditions.

(**B-C**) Heatmap and hierarchical clustering analysis of differentially expressed genes

in cKO heart as compared to the control. The examples of different gene clusters are

shown with their functional annotations (GO). (**D**) Top enriched GO terms of

upregulated or downregulated genes in the cKO heart compared with the control. (**E**)

GSEA analysis of various GO terms related to the upregulated or downregulated

genes in the cKO heart compared with the control.

Fig.4 IGF2R binds to granulocyte-derived inflammatory factors. (A)

Representative silver staining image of immunoblotting of IgG or anti-IGF2R

antibody pulled-down proteins from the lysed control or cKO hearts. Red rectangles

labeling the bands for mass spectrometry analysis. (**B**) Venn's diagram comparing the

lists of genes upregulated or downregulated in cKO hearts and identified peptides or

proteins from the selected bands from (A). Five factors identified from the integrated

analysis are listed. (**C**) Single-cell profiling of Ngp, Chil3 (Chi3l3), S100a9, or Ltf

expression in normal mouse hearts (GSM2967050) or bone marrow (GSM2967055).

Cell phenotype annotation: 1-granulocytes; 2-granulocytopoietic cell; 3-monocyte.

(**D-E**) Representative hematoxylin and eosin staining (D) or Masson's trichrome

staining (E) of the control or cKO heart grafts after I/R (12-hour cold HTK and

transplant reperfusion for 4 weeks).

Fig.5 Inhibition of IGF2R expression reduces cell death under hypoxia (**A**)
Immunostaining of IGF2R in H9C2 cardiomyoblasts before or after hypoxia-
reoxygenation. Scale bar, 50μm. (**B**) Immunoblotting of IGF2R in total cell proteins
from H9C2 myoblasts treated with or without hypoxia-reoxygenation. (**C**)
Immunostaining of IGF2R with co-staining of wheat germ agglutinin (WGA) in
H9C2 cells before or after hypoxia. Arrowhead indicates the cell plasma membrane.
Scale bar, 50μm. (**D**) Immunoblotting of IGF2R after subcellular (membrane or
cytosolic) fractions from H9C2 cells before or after hypoxia and reoxygenation. (**E**):
H9C2 myoblasts were transduced with the doxycycline (Dox) inducible CRISPRi
targeting IGF2R's promoter region. P-Tet-ON: Pomoter with tetracycline-inducible
activation element. (**F**): Immunoblotting of IGF2R and caspase3 in H9C2 cells
exposure to hypoxia/reoxygenation.
500

Figure 1

A

Sham | MI | I/R
1h | 3h | 6h

Igf2r

Gapdh

B

Igf2r cTnT DAPI

Igf2r DAPI

C

Igf2r

tSNE_2

tSNE_1

6
4
2
0

D

Transgene: Myh6 CreER

Igf2r locus: 9 — LoxP — 10 — LoxP — 11

+TAM

cKO: 9 — LoxP — 11

E

Relative Igf2r level

p<0.01

Ctrl cKO

1.5
1.0
0.5
0.0

F

Ctrl | cKO

Igf2r

Gapdh

Figure 2

Figure 3

A

Legend: ● Ctrl, ▲ cKO, I/R, Normal

PC2: 15% variance vs PC1: 60% variance

C

GO of cluster-a:
DNA repair (Rpain, Fen1, Prkdc, Sirt6, Lig4, etc.)
Cell cycle (Rb1, Rabgap1, Cdkn1, Cdk5, etc.)
Negative regulation of cell death (Ucn, Fgf2, etc.)
Autophagy (Tfeb, Atg13, Ulk3, Atp13a2, etc.)

GO of cluster-b:
Cell division (Dca3, Aurka, Septin10, Nek1, etc.)
Cilium assembly (Ift140, Fbf1, Ift80, Wdr35, etc.)
Autophagosome assembly (Ift20, Nprl3, etc.)
Fatty acid metabolic process (Lypla1, Cpt1a, etc.)

GO of cluster-c:
Inflammatory response (Csf1r, Ptgir, Ly86, etc.)
Apoptotic mitochondrial changes (Ppif, Hk2, etc.)
Cellular response to hypoxia (Egln3, Fndc1, etc.)
Angiogenesis (Dll4, Nrp2, Esm1, Vegfa, Apln , etc.)

GO of cluster-d:
Innate immune response (Cd84, Il23a, Tlr1, etc.)
Neutrophil chemotaxis (Ccl11, Cxcl1, S100a8, etc.)
Apoptotic process (Bcl2l1, Casp8, Nfkb1, etc.)
IL1β production (Nlrp3, Casp1, Casp4, Aim2, etc.)

B

Heatmap with clusters a, b, c, d. Legend: Ctrl, cKO, I/R, Normal. Normalized counts scale from -3 to 3.

D GO: cKO *vs* Ctrl

Up-regulated

mitotic nuclear division
nuclear division
DNA repair
chromosome segregation
organelle fission
cilium organization
intraciliary transport
chromatin organization
proteasome-mediated ubiquitin-dependent protein catabolic process
cilium assembly
peptidyl-lysine modification
nuclear chromosome segregation
glycerolipid metabolic process
sister chromatid segregation
chromosome separation
microtubule cytoskeleton organization involved in mitosis
cell cycle phase transition
autophagy
process utilizing autophagic mechanism
regulation of chromosome organization

Count: 0, 20, 40, 60

p.adjust: 4e-07, 3e-07, 2e-07, 1e-07

Down-regulated

positive regulation of cytokine production
leukocyte migration
positive regulation of response to external stimulus
leukocyte cell-cell adhesion
myeloid leukocyte activation
regulation of cell-cell adhesion
positive regulation of cell adhesion
myeloid leukocyte migration
cytokine-mediated signaling pathway
negative regulation of immune system process
regulation of immune effector process
granulocyte migration
neutrophil migration
regulation of leukocyte cell-cell adhesion
regulation of inflammatory response
mononuclear cell differentiation
positive regulation of cell-cell adhesion
leukocyte chemotaxis
positive regulation of defense response
response to molecule of bacterial origin

Count: 0, 50, 100, 150

p.adjust: 1.505758e-23, 1.129318e-23, 7.528788e-24, 3.764394e-24, 1.538923e-49

E GSEA: cKO *vs* Ctrl

autophagy ($p>0.05$); mitotic cell cycle process ($p<0.001$); innate immune response ($p<0.001$); inflammatory response ($p<0.001$); chemotaxis ($p<0.001$)

Figure 4

A

Ctrl — IP: IgG | Igf2r
cKO — IP: IgG | Igf2r

(kDa)
250 ▶
130 ▶
100 ▶
70 ▶
55 ▶
35 ▶
25 ▶
15 ▶

B

Down-regulated genes

Identified proteins from (Ctrl)

Up-regulated genes

Identified proteins from (cKO)

Actb
Ngp
Chil3
(Chi3l3)
S100a9
Ltf

C

Heart

Ngp   Ltf   Chi3l3   S100a9

Marrow

Ngp   Ltf   Chi3l3   S100a9

D

Ctrl   cKO
2X   2X
40X   40X

E

Ctrl   cKO
2X   2X
40X   40X

Figure 5