Long-term Hematopoietic Transfer of the Anti-Cancer and Lifespan-Extending Capabilities of A Genetically Engineered Blood System by Transplantation of Bone Marrow Mononuclear Cells

Jing-Ping Wang[#1,2], Chun-Hao Hung[#1,2*], Yao-Huei Liou[2], Ching-Chen Liu[2], Kun-Hai Yeh[2], Keh-Yang Wang[1,2], Zheng-Sheng Lai[2], Biswanath Chatterjee[1,2], Tzu-Chi Hsu[1,2], Tung-Liang Lee[2,6], Yu-Chiau Shyu[2,3,8*], Pei-Wen Hsiao[4,5], Liuh-Yow Chen[2], Trees-Juen Chuang[#,7], Chen-Hsin Albert Yu[2], Nah-Shih Liao[2], and Che-Kun James Shen[1,2*]

1. The Ph.D. Program in Medicine Neuroscience, Taipei Medical University, Taipei 115, Taiwan
2. Institute of Molecular Biology, Academia Sinica, Nankang, Taipei 115, Taiwan
3. Department of Nursing, Chang Gung University of Science and Technology, Taoyuan City 333, Taiwan
4. Agricultural Biotechnology Research Center, Academia Sinica, Nankang, Taipei 115, Taiwan
5. Graduate Institute of Life Sciences, National Defense Medical Center, Taipei 115, Taiwan
6. Chang Gung Memorial Hospital, Pro-Clintech Co. Ltd., Keelung 204, Taiwan
7. Genomics Research Center, Academia Sinica, Nankang, Taipei 115, Taiwan
8. Community Medicine Research Center, Chang Gung Memorial Hospital, Keelung Branch, Keelung 204, Taiwan

# Contributed equally

* Corresponding Authors

Please send all correspondence to:

Che-Kun James Shen

The PhD Program for Neural Regenerative Medicine

Taipei Medical University

Taipei 115, Taiwan, ROC

E-mail: ckshen@gate.sinica.edu.tw

Yu-Chiau Shyu

Community Medicine Research Center,

Chang Gung Memorial Hospital, Keelung Branch,

Keelung 204, Taiwan (R.O.C.)

E-mail: yuchiaushyu@gmail.com

39
40   Chun-Hao Hung

41   The PhD Program for Neural Regenerative Medicine

42   Taipei Medical University

43   Taipei 115, Taiwan, ROC

44   E-mail: libur777@hotmail.com

Abstract

A causal relationship exists among the aging process, organ decay and dis-function, and the

occurrence of various diseases including cancer. A genetically engineered mouse model,

termed $Eklf^{K74R/K74R}$ or $Eklf$(K74R), carrying mutation on the well-conserved sumoylation site

of the hematopoietic transcription factor KLF1/ EKLF has been generated that possesses

extended lifespan and healthy characteristics including cancer resistance. We show that the

healthy longevity characteristics of the $Eklf$(K74R) mice, as exemplified by their higher

anti-cancer capability, are likely gender-, age- and genetic background-independent.

Significantly, the anti-cancer capability, in particular that against melanoma as well as

hepatocellular carcinoma, and lifespan-extending property of $Eklf$(K74R) mice could be

transferred to wild-type mice via transplantation of their bone marrow mononuclear cells at

young age of the latter. Furthermore, NK(K74R) cells carry higher *in vitro* cancer cell-killing

ability than wild type NK cells. Targeted/global gene expression profiling analysis has

identified changes of the expression of specific proteins, including the immune checkpoint

factors PD-1 and PD-L1, and cellular pathways in the leukocytes of the $Eklf$(K74R) that are in

the directions of anti-cancer and/or anti-aging. This study demonstrates the feasibility of

developing a transferable hematopoietic/ blood system for long-term anti-cancer and,

potentially, for anti-aging.

Introduction

Aging of animals, including humans, is accompanied by lifespan-dependent organ deterioration and the occurrence of chronic diseases such as cancer, diabetes, cardiovascular failure and neurodegeneration[1,2]. To extend healthspan and lifespan, various biomedical- and biotechnology-related strategies have been intensively developed and applied, including the therapy of different diseases such as cancer[3-5]. The hematopoietic/ blood system is an important biomedical target for anti-aging and anti-cancer research development. Multiple blood cell lineages arise from hematopoietic stem cells (HSCs), with the lymphoid lineage giving rise to T, B, and natural killer (NK) cell populations, whereas the myeloid lineage differentiates into megakaryocytes, erythrocytes, granulocytes, monocytes and macrophages[6-8]. The genetic constituents and homeostasis of the hematopoietic system are regulated epigenetically and via environmental factors to maintain animal health[6,9].

EKLF, also named KLF1, is a Krüppel-like factor that is expressed in a range of blood cells including erythrocytes, megakaryocytes, T cells, NK cells, as well as in various hematopoietic progenitors including common-myeloid-progenitor (CMP), megakaryocyte-erythroid-progenitor (MEP), and granulocyte-macrophage-progenitor (GMP)[10-12]. The factor regulates erythropoiesis[13] and the differentiation of MEP to megakaryocytes and erythrocytes[10,14] as well as of monocytes to macrophages[15]. EKLF is also expressed in HSC and regulates their differentiation[16]. The factor can positively or negatively regulate transcription, including the adult β globin genes, through binding of its canonical zinc finger domain to the CACCC motif of the regulatory regions of a diverse array of genes[16-19]. In fact, the zinc finger of EKLF could recognize/bind to genomic sites with the consensus sequence CCNCNCCC, and recruit different co-activator or co-repressor chromatin remodeling complexes[14,17].

EKLF could be sumoylated *in vitro* and *in vivo*, and sumoylation of the lysine at codon 74 of mouse EKLF altered the transcriptional regulatory function[10] as well as nuclear import[20] of the factor. Surprisingly, homozygosity of a single amino acid substitution, lysine(K) to arginine(R), at the sumoylation site of EKLF results in the generation of a novel mouse model with healthy longevity. These mice, termed *Eklf*[K74R/K74R] or *Eklf*(K74R), exhibited extended healthspan and lifespan. In particular, the *Eklf*(K74R) mice showed delay of the

age-dependent decline of physical performance, such as the motor function and spatial
learning/memory capability, and deterioration of the structure/function of tissues including
the heart, liver, and kidney. Furthermore, the *Eklf*(K74R) mice appeared to have significantly
higher anti-cancer capability than the WT mice[21].

As described in the following, we have since characterized the high anti-cancer
capability of the *Eklf*(K74R) mice with respect to its dependence on the age, gender and
genetic background. More importantly, we have demonstrated that the high anti-cancer ability
of these genetically engineered mice could be transferred to wild type mice (WT) through
hematopoietic transplantation of the bone marrow mononuclear cells (BMMNC).
Furthermore, we show that the higher anti-cancer capability and extended life span of
Eklf(K74R) mice are associated with changes of the global protein expression profile and
specific aging-/cancer-associated cellular signaling pathways in their white blood cells (WBC)
or leukocytes.
118
119

Result

Characterization of the cancer resistance of *Eklf*(K74R) mice in relation to age, gender,
and genetic background

The *Eklf*(K74R) mice appeared to be cancer resistant to carcinogenesis as manifested by
their lower spontaneous cancer incidence (12.5%) in life than WT mice (75%). The
Eklf(K74R) mutation also protected the mice from metastasis of melanoma cells and it
reduced melanoma growth in the subcutaneous cancer cell inoculation assay[21]. Here we used
the pulmonary melanoma foci assay, which has been commonly used for quick analysis of
early metastasis[22], to further characterize the higher cancer resistance of the *Eklf*(K74R) mice
with respect to the effects of gender/age/genetic background of the mice and the requirement
for the homozygosity of the K74R mutation. The cultured melanoma B16-F10 cells used in
this assay are poorly immunogenic and highly aggressive thus being useful for monitoring
early metastasis in various immunotherapy studies[22,23].

It appeared that male as well as female *Eklf*(K74R) mice in the B6 genetic background
had significantly fewer pulmonary melanoma foci than the corresponding WT mice (Figure 1).
Because of this result, we used male mice for all of the studies describe below. First, both
young (2 month-old) and aged (24 month-old) *Eklf*(K74R) mice had higher anti-metastasis

ability against the injected melanoma cells than WT mice of age-dependent groups (Figure

1A and 1B). Secondly, homozygosity of the K74R substitution was required for the higher

cancer resistance of the *Eklf*(K74R) mice (Figure S1). Importantly, the *Eklf*(K74R) mice in

the FVB background also exhibited high cancer resistance than FVB WT mice by this assay

(Figure 1C), suggesting that cancer resistance of *Eklf*(K74R) mice conferred by the

homozygous K74R substitution was likely genetic background-independent. Finally, the

higher anti-cancer capability of the *Eklf*(K74R) mice did not appear to depend on the arginine

at codon 74, since *Eklf*(K74A) mice carrying K→A amino acid substitution at the K74

sumoylation site also exhibited higher anti-metastasis capability than WT mice in the

pulmonary foci assay (Figure 1D).

The anti-cancer capability of *Eklf*(K74R) mice was not restricted to melanoma. As

shown by the subcutaneous inoculation assay of Hepa1-6 cells (Figure 1E), *Eklf*(K74R) mice

also carry a higher anti-cancer capability against hepatocellular carcinoma than the WT mice.

It thus appears that the *Eklf*(K74R) mice are resistant to the carcinogenesis of a range of

different cancers.

Transfer of the anti-cancer capability and extended lifespan of *Eklf*(K74R) mice to WT

mice via BMT

Since EKLF is a hematopoietic transcription factor expressed not only in mature blood

cells but also in HSCs and hematopoietic stem progenitor cells, this cancer resistance may be

transferable by means of BMT. This possibility was tested with uses of male mice and a

standard BMT protocol[24]. BMMNC were purified from the bone marrow of 2 month-old

CD45.2 *Eklf*(K74R) or WT mice and injected into the tail vein of CD45.1 WT recipient mice.

Blood replacement of recipient mice with 10Gy γ-irradiation by that of the donor mice

reached 90% at 7th-week (Figure 2A and 2B). After 2 weeks, the recipient mice were injected

with B16-F10 cells and then sacrificed a further 2 weeks later to quantify pulmonary tumor

foci. We found that WT mice transplanted with WT BMMNC had similarly high numbers of

tumor foci relative to WT mice without BMT (Figure 2C and 1A). However, similar to

Eklf(K74R) mice challenged with B16-F10 cells (Figure 1B), WT mice that received

BMMNC from *Eklf*(K74R) mice presented significantly fewer tumor foci on their lungs

(Figure 2C). Notably, BMT using 24 month-old donor mice gave similar result although the

effect is relatively small (Figure S2A).

In order to determine if WT mice having more restricted blood replacement upon BMT

from *Eklf*(K74R) mice still exhibited a higher anti-cancer capability, we also carried out BMT

experiments with lower doses of γ-irradiation. BMT using two lower doses of γ-irradiation

(2.5Gy/5Gy) still resulted in transfer of cancer resistance from *Eklf*(K74R) to WT mice.

Approximately 40% of recipient blood cells were substituted by donor cells upon BMT with

5Gy γ-irradiation. Consequently, at that irradiation dosage, BMT from *Eklf*(K74R) mouse

donors reduced the average number of pulmonary tumor foci in recipient WT mice to 5. On

the other hand, only 20% blood replacement was achieved in the recipient mice with 2.5Gy

γ-irradiation (Figure 2D). However, the WT mice receiving BMT from *Eklf*(K74R) mice

again developed less number (~10/mouse) of pulmonary tumor foci than those WT mice

receiving BMT from the WT mice (Figure 2D). Thus, even at a low level of 20% blood

replacement, BMT still enabled effective transfer of cancer resistance from *Eklf*(K74R) mice

to WT mice.

In addition, we also attempted to transfer the extended lifespan characteristics of the

Eklf(K74R) mice to WT mice by BMT. Significantly, the median lifespan of WT mice

receiving BMT from *Eklf*(K74R) mice was 5 months longer than that of WT mice receiving

BMT from WT mice (Figure S2B). Thus, the longer lifespan characteristics of the *Eklf*(K74R)

mice was also transferable via BMT.

Inhibition of tumor growth by transplanted BMMNC from *Eklf* K74R) mice

Our experiments indicated that *Eklf*(K74R) bone marrow carried the anti-metastasis

capability that prevented melanoma cell colonization on the lungs of recipient mice (Figures.

1 and 2). To determine if *Eklf*(K74R) BMT could inhibit tumor growth, we examined the

effect of BMT on growth of tumors with B16-F10-luc cells. As outlined in Figure 3A, ten

days after injection of cancerous cells, the formation of bioluminescent signals in the recipient

mice were confirmed by the observation of *in vivo* bioluminescence. The following day, we

transplanted the recipient mice with BMMNC from WT or *Eklf*(K74R) mice and then

measured the intensities of bioluminescence signals from tumors 7 and 14 days later. As

shown, tumor growth in mice subjected to BMT from *Eklf*(K74R) mice was significantly

slower relative to those receiving BMMNC from WT mice (Figure 3B and 3C). Thus,

Eklf(K74R) BMMNC indeed can inhibit the growth of tumor more effectively than WT

BMMNC.

Differential expression of specific immune-, ageing- and/or cancer- associated

biomolecules in the blood of *Eklf*(K74R) mice

In general, the cell populations in PB were not affected much by the K74R substitution,

as shown by CBC analysis[21], although flow cytometry analysis of PB from WT and

Eklf(K74R) mice of different ages showed that the population of NK(K74R) cells in 24

month-old *Eklf*(K74R) mice was higher than WT NK cells in 24 month-old WT mice (Figure

S3). In parallel, the K74R substitution did not affect much the expression of *Eklf* gene in the

different types of hematopoietic cells. As seen in Figure S4, the K74R substitution did not

alter the expression levels of EKLF protein in leukocytes from peripheral blood (PB) (left

panels, Figure S4A), although the *Eklf* mRNA level of *Eklf*(K74R) leukocytes was 10-20%

lower than that of WT leukocytes (right panel, Figure S4A). Previous studies and database

have shown that *Eklf* mRNA and EKLF protein are expressed in the erythroid lineage[10] and

megakaryocytes[10,14], and the *Eklf* mRNA is expressed in HSC[16](bio-GPS database), NK cells

(bio-GPS database), naïve T cells[12] and Treg cells[12]. By Western Blotting (WB) analysis

(Figure S4B) and RT-qPCR analysis (Figure S4C), we found that the *Eklf* gene is expressed

in B220$^+$ B cells, CD3$^+$ T cells, and NK cells of both WT and *Eklf*(K74R) mice, but the levels

are much lower than that of the erythroid MEL cells.

We first used RT-qPCR to analyze the levels in PB cells of mRNAs encoding the

immune checkpoint genes (ICGs) PD-1/PD-L1[25] in view of the cancer resistance of

Eklf(K74R) mice (Figure 4) as well as increased levels of PD-1 and PD-L1 in aged or

tumorigenic mice[26]. As shown in Figure 4, the mRNA levels of *Pd-1* and *Pd-l1* in the PB, B

cells and T cells of WT mice were both increased during aging. In great contrast, the mRNA

levels and protein levels of these two genes were lower in 3 month-old *Eklf*(K74R) mice than

the age-matched WT mice, and they remained low during ageing of the *Eklf*(K74R) mice.

Significantly, RNAi knockdown of *Eklf* expression reduced *pd-1* and *pd-l1* mRNA levels in

splenic CD3$^+$ T cells (Figure 4C). In interesting correlation, overexpression of EKLF would

increase the expression of *pd-l1* in mouse CD4$^+$ T cells[12]. There data together indicate that

EKLF positively regulates the transcription of the *Pd-1* and *Pd-l1* genes, directly or indirectly,

and likely the transcriptomes of a diverse range of hematopoietic cells. Since the levels of

EKLF protein in CD3$^+$ T (K74R) cells and WT CD3$^+$ T cells were similar (Figure S4B),

change of the expression of *Pd-1* and *Pd-l1* genes (Figure 4) was most likely caused by the

loss of the sumoylation of EKLF(K74R), which would alter its transactivation capability[17].

Interestingly, *pd-1* appeared to be a direct regulatory target of EKLF. As shown by

ChIP-qPCR analysis, EKLF was bound at the CACCC box (Box3) at -103 of the *Pd-1*

promoter in CD3$^+$ T cells from both WT and *Eklf*(K74R) mice (Figure S5). The detail of the

role of sumoylation in the transcriptional activation of *Pd-1* by EKLF awaits future

investigation. As expected, lower levels of *Pd-1* and *Pd-11* expression were also observed in

the PB of mice receiving BMT from *Eklf*(K74R) mice (Figure S4D). These findings together

suggest that the low tumorigenesis rate of *Eklf*(K74R) mice arises in part from low expression

of the two ICGs, *Pd-1* and *Pd-l1*, as the result of K74R substitution in the EKLF protein.

We have also examined, by bead-based multiplex assay[27,28], the expression patterns of

several ageing- and/or cancer- associated cytokines. As shown in Figure S6, there was no

significant difference in the serum levels of IL-1β, IL-2, IL-10, IL-12p70, INF-γ or TNF-α

between WT and *Eklf*(K74R) mice at 24 month-old. In contrast, the level of IL-4, an

anti-inflammatory cytokine[29] beneficial to the hippocampus of aging mice[30], in 24 month-old

Eklf(K74R) mice was 3-4 fold higher than the 24 month-old WT mice. On the other hand, the

level of IL-6, a key factor in chronic inflammatory diseases, autoimmunity, cytokine storm

and cancer[28,31], increased only moderately during aging of the *Eklf*(K74R) mice (Figure S6).

Thus, similar to PD-1 and PD-L1, the altered expression of some of the cytokines in the blood

likely also contributes to the anti-aging and/or anti-cancer characteristics of the *Eklf*(K74R)

mice.

Comparative proteomics analysis of the leukocytes of *Eklf*(K74R) mice and WT mice

We proceeded to examine age-dependent cell-intrinsic changes in the proteomes of the

leukocytes from the WT and *Eklf*(K74R) mice in two different age groups. 259 and 306

differentially expressed proteins (DEPs) were identified between the two age groups for the

WT and *Eklf*(K74R) mice, respectively (Figure 5A). To understand the correlations of these

proteins with aging and cancer, we performed the GSEA and found that the age-dependent

DEPs changed in the concordant direction in the WT and *Eklf*(K74R) mice were enriched for

several known aging-related pathways, e.g. IL-6-JAK-STAT3 signaling, DNA repair, etc[32]

(Figure 5B). Meanwhile, the age-dependent DEPs changed in the reverse directions in the

WT and *Eklf*(K74R) mice were enriched for nine other aging-related pathways (Figure 5C).

We further performed DEP analyses between WT and *Eklf*(K74R) mice and identified

strain-dependent DEPs in the two age groups. As shown in Figure 5D, only 7 DEPs were

identified in the 3 month-old mice but 40 DEPs in the 24 month-old ones. Of the 40 DEPs in

the elder mice, 3 and 37 were upregulated in *Eklf*(K74R) and WT mice, respectively (Figure

5D). Significantly, GSEA analysis of these DEPs showed that elder *Eklf*(K74R) leukocytes

were enriched for the anti-aging pathways related to hypoxia, and p53 signaling, etc.[33,34],

while the elder WT leukocytes were enriched for the aging-associated pathways related to

apoptosis, and mTORC1 signaling, etc[33,35]. On the other hand, the DEPs in elder *Eklf*(K74R)

leukocytes were also enriched for anti-cancer pathways related to the interferon-α response,
and TGF-β signaling, etc.[36-38] (Figure 5E), while DEPs in the elder WT leukocytes were
enriched for the pro-cancer pathways related to IL-6-JAK-STAT3 signaling and
angiogenesis[39]. These data together have demonstrated that *Eklf*(K74R) leukocytes contribute
to their anti-cancer capability and long lifespan through several specific cellular signaling
pathways.
279
Higher *in vitro* cancer cell cytotoxicity of the NK(K74R) cells than WT NK cells
The higher anti-cancer capability of the *Eklf*(K74R) mice could result from combined
contributions by different types of the hematopoietic cells, e.g. the NK(K74R) cells. NK cells
are effector lymphocytes of the innate immune system that can rapidly destroy tumor cells
and against infectious pathogen[40]. Also, NK cells do not require pre-stimulation for the
treatment of cancer cells[41]. Further, the NK lineage could be fully reconstituted 2 weeks after
BMT[42]. We tested the *in vitro* cancer cell cytotoxicity ability of NK(K74R) cells from 3
month-old *Eklf*(K74R) mice in comparison to the WT NK cells. As shown in Figure 6,
NK(K74R) cells exhibited significantly higher cytotoxicity against the B16-F10 melanoma
cells as well as Hepa1-6 hepatocellular carcinoma cells than WT NK cells. This data together
with the higher level of NK cells in aged *Eklf*(K74R) mice (Figure S3) indicated that in
parallel to the leukocytes[21], NK cells also play an important role in the higher anti-cancer
capability of the hematopoietic blood system of *Eklf*(K74R) mice, as exemplified in Figures 2
and 3.
294
Discussion
Because of the complexity and intercrosses of the different pathways regulating the
health and the aging process, genetic manipulation of non-human animals[43,44] and non-genetic
approaches on animals including human[4,45,46] targeting these pathways inevitably lead to
moderate-to-severe side effects such as body weight loss, adiposity, etc.. With respect to the
above, the *Eklf*(K74R) mice[21] is ideal as an animal model for further insightful understanding
of the ageing process as well as for biomedical development of new anti-ageing tools and
approaches. In the current study, we demonstrate that the healthy longevity of the *Eklf*(K74R)
mice, as exemplified by their higher tumor resistance, is likely to be gender/genetic
background independent and irrespective of the amino acid change at K74 of EKLF. Further,
we show that the *Eklf*(K74R) mice are resistant to carcinogenesis of not only melanoma, but

also other cancers such as hepatocellular carcinoma. This correlates well with the higher

cytotoxicity *in vitro* of NK(K74R) cells than WT NK cells against the same two types of

cancer cells. Significantly, we also show that one-time transplantation of the bone marrow

mononuclear cells or BMMNC (BMT) could confer the recipient WT mice extended lifespan

and higher tumor-resistance capability, the latter of which could be achieved by 20% of

substitution of the blood of the recipient WT mice with that from the *Eklf*(K74R) mice. These

findings together have interesting implications in the development of this novel hematopoietic

blood system for future medical application in anti-disease and anti-aging.

The anti-cancer capability of the *Eklf*(K74R) mice have rendered them relatively free

from spontaneous cancer occurrence[21], which is also reflected by their resistance to

tumorigenesis of the B16-F10 cells and Hepa1-6 in the anti-cancer assays (Figures 1-3).

Furthermore, the cancer resistance of *Eklf*(K74R) mice appears to be independent of the

gender, age, and genetic background (Figure 1). The anti-metastasis property of the

Eklf(K74A) mice in the pulmonary foci assay (Figure 1D) also indicates that the anti-cancer

capability of the *Eklf*(K74R) mice is not due to the structural and/or post-translational

properties of the arginine introduced at codon 74 of EKLF. Importantly, we have shown that

the anti-cancer capability and the extended lifespan characteristics of *Eklf*(K74R) mice are

transferrable through BMT (Figure 2, 3, and S2). In particular, we show that BMMNC from

Eklf(K74R) mice (Figure 2 and S2A) could confer 2 month-old WT recipient mice with the

anti-cancer capability against melanoma. Furthermore, ~20% of blood substitution would

allow the recipient mice to become cancer-resistant in the pulmonary foci assay (Figure 2D).

It is likely that homozygosity, but not heterozygosity, of K74R substitution causes one or

more types of the hematopoietic cells to gain new functions, such as the higher cancer

cell-killing capability of NK(K74R) cells (Figure 6). The data of Figure 2D further suggest

that the amount of specific types of cells such as NK(K74R) in the 20% of blood cells

carrying homozygous *Eklf*(K74R) alleles in the recipient mice upon BMT is sufficient to

confer the mice a higher anti-cancer ability.

Hematopoietic stem cell therapy for different diseases[47-50] including cancer has been

intensively explored and practiced such as leukemia, and neuroblastoma, etc[51,52]. Also, certain

characteristics of the young mice could be transferred to old mice via heterochronic

parabiosis or heterochronic transplantation[53-56]. Similarly, plasma proteins from human

umbilical cord blood can revitalize hippocampal function and neuroplasticity in aged mice[57,58].

Previously, we have shown that multiple injections of LSK-HSC(K74R) from 3 month-old

Eklf(K74R) mice to 25 month-old WT mice would extend the lifespan of the latter by about 4

months[21]. In the current study, we further show that 2 month-old WT mice receiving one-time

transplantation of BMMNCs from 2 month-old *Eklf*(K74R) mice would live 5.5 months

longer than the controls (Figure S2B). Thus, transplantation/transfer of the blood MNC

carrying homozygous mutation at the sumoylation site of EKLF could be developed as a new

approach for long-term anti-aging, anti-cancer and likely rejuvenation.

The tumorigenesis resistance and long lifespan exhibited by the *Eklf*(K74R) mice are

most likely due to changes in the transcription regulatory properties of the mutant

EKLF(K74R) protein relative to WT[59]. As exemplified in Figure 4A and 4B, expression

levels of the ICGs *Pd-1* and *Pd-l1* in the PB, B, and T cells of *Eklf*(K74R) mice are reduced

in comparison to the WT mice. Notably, cancer incidence increases with aging[60], which is

accompanied by increased expression of PD-1 and PD-L1[61]. The lower expression of ICGs

would contribute to the anti-cancer capabilities of the *Eklf*(K74R) blood to fight against

cancer (Figures. 1, 2, and 3) and to extension of the lifespan of cancer-bearing mice[21]. Indeed,

similar to ICGs, the expression levels of several cytokines in the *Eklf*(K74R) blood/serum are

also different from the WT blood and some of the changes during ageing or carcinogenesis in

the *Eklf*(K74R) blood are opposite to the blood/serum of WT mice [28] (Figure S6).

The transcriptome data have been used to dissect the regulation of leukocyte aging [32,62,63].

In addition, proteomic analysis has revealed the signaling pathways that regulate aging of

specific types of leukocyte such as the lymphocyte and neutrophils cells[64,65]. In this study, we

have performed proteomics analysis of leukocytes from WT and *Eklf*(K74R) mice in two age

groups, and found that for the elder mice, the strain-dependent DEPs in the leukocytes are

enriched for a number of signaling pathways. Among these signaling pathways, at least 12 of

them are closely associated with the aging process, which include hypoxia, DNA repair, etc.

(Figure 5E). As summarized by the model in Figure 5F, it appears that changes of these

pathways in the elder *Eklf*(K74R) leukocytes relative to the elder WT leukocytes are mostly

in the direction of anti-aging. The data of Figure 5 together strongly suggest that the

EKLF(K74R) amino acid substitution causes a change in the global gene expression profile of

the leukocytes, which contributes to the high anti-cancer capability and long lifespan of the

Eklf(K74R) mice.

In sum, we have characterized the cancer resistance of the *Eklf*(K74R) mice, among their

other healthy characteristics, in relation to gender, age, and genetic background. We also have

identified cell populations, gene expression profiles and cellular signaling pathways of the

white blood cells of young and old mutant mice, in comparison to the WT ones, that are

changed in the anti-cancer and/or anti-ageing directions. Finally, the transferability of the

cancer resistance and extended life-span of the mutant mice via transplantation of BMMNC

suggests the possibility of future development of hematopoietic blood cells genome-edited at

the conserved sumoylation site of EKLF for anti-cancer and the extension of healthspan

and/or lifespan in animals including human.

Acknowledgments

We thank Ya-Min Lin and her colleagues at the FACS Core for their efforts on the cell

sorting. We also thank Taiwan Mouse Clinic facility for expertise help on the IVIS

experiments. The effort by Dr. Ching-Yen Tsai at the Transgenic Core Facility, IMB, is

greatly appreciated. The Immune Monitoring Core of Taipei Medical University provided

great help in the analysis of the blood cell populations. We also thank the Bioinformatics

Core Facility, IMB, for its efforts on the proteomics analysis, and Dr. Da-Liang Ou at

National Taiwan University for providing us the Hepa1-6 cell line.

Competing interests

The authors declare there is no conflict of interest.

Animal

All the animal procedures were approved by the Institute of Animal Care and Use

Committees, IACUC, at Academia Sinica. The IACUC numbers are 17-02-1052, 17-12-1142,

20-10-1528, BioTReC-110-M-005, and BioTReC-111-M-004, respectively.

Author Contributions

C.-H.H., designed and performed experiments, analyzed data and co-wrote the helped paper.;

K.-Y.W., J.-P.W. and T.-L.L. performed experiments and analyzed data.; Y.-H.L. provided

technology expertise; Z.-S.L. and Y.-H.L. provided essential mouse strains and experimental

knowledge.; P.-W.H. provided essential cell-line strains and experimental knowledge.;

Y.-H.L., N.-S.L., and Y.-C.S., provided essential mouse strains.; C.-C.L., K.-H.Y., C.-H.A.Y.

and T.-J.C. analyses of the bioinformatics data and co-wrote the paper. Y.-H.L. and N.-S.L.

provided technology expertise, ideas and essential materials.; Y.-C.S. and C.-K.J.S. provided

ideas, supervised the research and co-wrote the paper.

Funding

This work was supported by Taipei Medical University (to C.-K. J. S.), Academia Sinica (to

C.-K. J. S.), Chang-Gung Memorial Hospital Research Project (CMRPG2L0081 and

CMRPG2G0311 to Y.-C. S.), and grants from the National Science and Technology Council,

Taiwan, ROC (NSC 102-2320-B-001-010 to N.-S. L., MOST 103-2311-B-010-003 Y.-C. S.,

and MOST 110-2628-B-001-024 to L.-Y. C.).

Availability of data and materials

The data that support the findings of this study are available from the corresponding authors

upon request.

Material and Methods

Mice

C57BL/6, B6, and FVB mice were purchased from Jackson Laboratories (Bar Harbor,

Maine). The B6 *Eklf*(K74R), B6 *Eklf*(K74A) and FVB *Eklf*(K74R) mice were established

with the assistance of the Transgenic Core Facility (TCF), IMB, Academia Sinica, Taiwan.

As described previously[21], the K74R mutation was introduced by homologous recombination

into exon 2 (E2) of the *Eklf* gene of B6 mice by means of a recombinant retrovirus containing

the construct loxP-PGK-gb2-neo-loxP-E2 (K74R), before excising the neomycin (neo)

selection marker by crossing with Ella-Cre mice. The heterozygous *Eklf*(K74R/+) mice were

then crossed to obtain homozygous mutant *Eklf*(K74R/K74R) mice, hereafter termed

Eklf(K74R) mice.

On the other hand, *Eklf*(K74A) mice were generated by using the CRISPR/Cas9 system.

Female B6 mice (7 to 8 week-old) were mated with B6 males and the fertilized embryos were

collected from the oviducts. For oligos injection, Cas mRNA (100 ng/μl), sgRNA (50 ng/μl),

and donor oligos (100 ng/μl) were mixed and injected into the zygotes at the pronuclei stage.

The F0 mice were genotyped by PCR and DNA sequencing. The heterozygous *Eklf*(K74A/+)

mice were crossed to establish the germ-line stable homozygous *Eklf*(K74A) F1 strain.

Eklf(K74R) mice in the FVB background were generated using an *in vitro* fertilization

strategy. Briefly, sperm from male B6 *Eklf*(K74R) mice was used to fertilize FVB mouse

oocytes. *In vitro* fertilizations of FVB oocytes were carried out consecutively for five

generations. The resulting chimeric mice with >90% FVB background were then crossed with

FVB mice for another five generations or more.

Cell lines

Murine B16-F10 melanoma cell line was purchased from ATCC (CRL-6475). B16-F10 cells expressing luciferase (B16-F10-luc) were generated by infection of recombinant lentivirus stably expressing luciferase luc2 under the control of hEF1A-eIF4G promoter (InvivoGen)[61]. All cell lines were derived from cryopreserved stocks split fewer than three times and they were confirmed as mycoplasma-free prior to use. B16-F10 cells were cultured at 37 °C and 5 % CO_2 in DMEM medium supplemented with 10 % FBS, 1 % penicillin/streptomycin, and 2 mM L-glutamine. B16-F10-luc cells were selected at 37 °C and 5 % CO2 in a DMEM medium supplemented with 0.2 mg/mL zeocin (Invitrogen), 10 % FBS, 1 % penicillin/streptomycin, and 2 mM L-glutamine. The hepatocellular carcinoma cell line, Hepa1-6, was from Dr. Da-Liang Ou 's lab at College of Medicine, National Taiwan University. The line was maintained at 37°C and 5% CO_2 in Gibico[TM] DMEM, a high glucose commercial medium containing 10% FBS (Gibico[TM]) and 1% penicillin-streptomycin.

Pulmonary melanoma foci assay

Cultured B16-F10 melanoma cells ($1x10^5$ cells/mouse) were injected into the tail vein of 8 week- to 9 week-old or 24 month-old *Eklf*(K74R) and WT mice with/ without bone marrow transplantation. Two weeks after injection, the mice were sacrificed and the number of tumor foci on their lungs was quantified[66].

Subcutaneous cancer cell inoculation assay of Hepa1-6

The protocol follows that used by Han *et al*[67]. After anaesthetization of the mice with 3% isoflurane, $2x10^6$/ 50uL of Hepa1-6 cells in serum-free DMEM medium were mixed with 50ul of Corning® Matrigel® Matrix and injected between skin and muscle of the hind limb. The size of the tumors was measured and recorded once a week with Mitutoyo® Vernier Caliper. The volume was calculated using the formula: length × weight × height × π/6.

Flow cytometric analysis and cell sorting

Single cell suspensions of the peripheral blood cells and spleen tissue of B6 mice were prepared by lysing red blood cells and then passing them through a 40-μm cell strainer (Falcon®). Bone marrow mononuclear cells (BMMNCs) were prepared as described below in the section **Bone marrow transplantation (BMT)**. The peripheral blood cells and splenocytes were stained extracellularly for 30 min at room temperature using different combinations of the following antibodies: anti-CD45.1 (eBioscience); anti-CD45.2

(eBioscience); anti-CD3e (eBioscience); anti-CD45R (eBioscience); anti-NK1.1

(eBioscience); anti-NKp46 (eBioscience); anti-Gr-1 (eBioscience); CD11b (eBioscience);

anti-PD-1 (eBioscience) and anti-PD-L1 (eBioscience). The various hematopoietic progenitor

cell compartments of bone marrow were also stained extracellularly for 30 min at room

temperature by using different combination of the following antibodies: anti-Lineage

(eBioscience), anti-c-Kit (eBioscience), anti-Sca-1 (eBioscience), anti-CD34 (eBioscience),

anti-Flt-3 (eBioscience). All the immuno-stained cells were subsequently washed with 1%

PBS three times and resuspended for FACS analysis and sorting. Small amounts of the cell

samples were run on a FACS Analyzer LSRII-12P (BD Bioscience) to determine the

proportions of different cell preparations. FACS AriaII SORP (BD Bioscience) was then used

to sort the indicated cell populations. The detail gating subsets for all the cell as described

above are shown in Table S1. Data analysis was performed using FlowJo software.

RNAi knockdown of *Eklf* mRNA from T cells

CD3$^+$ T cells isolated by sorting were cultured in RPMI 1640 medium for one day for

recovery and then transfected with EGFP-plasmid (control), scrambled oligonucleotides (SC

control), *Eklf* knockdown oligonucleotide 1 (oligo 1), or *Eklf* knockdown oligonucleotide 2

(oligo 2) in a 96-well plate for 48 h using a LONZA electroporation kit (P3 Primary Cell

4D-NucleofectorTM X Kit) and machine (4D-NucleofectorTM Core Unit). Then the cells

were lysed using a PureLink® RNA Mini kit (Life Technologies) and analyzed by RT-qPCR.

RT-qPCR analysis

Total RNAs of B cells, T cells, and leukocytes/ WBC from the peripheral blood or

spleen of *Eklf*(K74R) mice and WT mice, respectively, were extracted using a PureLink®

RNA Mini kit (Life Technologies). The RNAs were reverse-transcribed by means of oligo-dT

primers, Maxima H Minus Reverse Transcriptase (Thermo Scientific™) and SYBR Green

reagents (Applied Biosystems). RT-qPCR was performed using a LightCycler® Nano

machine (Roche). Gene-specific primers for *Eklf*, *Pd-1*, *Pd-l1*, *Pd-l2,* and *Gapdh* were

designed using Vector NTI Advance 9 software according to respective mRNA sequences in

the NCBI database (primer sequences are available upon request). The PCR primers for

analysis of *Eklf* mRNA had been used and verified in a previous study[16]. Expression levels of

mRNAs were normalized to that of endogenous *Gapdh* mRNA.

ChIP-qPCR

The ChIP-qPCR analysis followed the procedures by Daftari et al[68]. Extracts were

prepared from formaldehyde cross-linked mouse CD3[+] T cells and erythroleukemia cell line

(MEL), sonicated, and then immune-precipitated (IP) with anti-EKLF (Thermo) or purified

rabbit IgG. The chromatin DNAs in the precipitates were purified and analyzed by qPCR in

the Roche LightCycle real-time system. Sequences of the DNA primers used for qPCR are

shown in Supplementary Table 2.

Western blotting (WB)

We adopted a previously described WB procedure[69]. Leukocytes/ WBC of Eklf(K74R)

mice and WT B6 mice, respectively, were collected from RBC lysis buffer-treated peripheral

blood. CD3[+] T cells, B220[+] B cells and NK cells from the spleen were sorted by FACS AriaII

SORP (BD Bioscience). The cell pellets were lysed in sample buffer and run on SDS-PAGE

gels. WB with anti-EKLF (Abcam) and anti-actin (Sigma) antibodies was then used to

analyze and compare the levels of EKLF and actin proteins.

In vivo **bioluminescence imaging**

Eklf(K74R) and WT B6 mice were physically restrained and 1×10^5 B16-F10-luc

cells/mouse were intravenously injected into their tail vein. Ten days after melanoma cell

inoculation, mice were anesthetized for 5 min and injected intraperitoneally with D-luciferin

(300 mg/Kg of body weight). Fifteen minutes after maximum luciferin uptake, the mice were

subjected to imaging of the lung and liver regions in an IVIS 50 Bioluminescence imager

(Caliper Life Sciences) to determine metastatic burden. The same mice were used the next

day as recipients of bone marrow transplantation (BMT) from donor WT or Eklf(K74R) mice.

Following BMT, bioluminescence imaging was performed on days 0, 10, 17 and 24.

Bone marrow transplantation (BMT)

BMT followed the standard protocol described in Imado et al. (2004)[24]. B6, CD45.1 or

CD45.2 donor mice were sacrificed and their femurs were removed. Bone marrow cells were

harvested by flashing the femurs with RPM I1640 medium (GIBCO) using a 27-gauge needle

and syringe. The cells were then incubated at 37 °C for 30 min in murine complement buffer

containing antibodies against B cells, T cells and NK cells, washed twice with PBS, and then

subjected to Ficoll-Paque PLUS gradient centrifugation to collect bone marrow mononuclear

cells (BMMNCs). BMMNCs (1×10^6 cells/mouse) from donor mice were injected into the tail

veins of recipient B6, CD45.2 or CD45.1 mice that had been exposed to total body

γ-irradiation of 10, 5 or 2.5 Gy.

Calcein-AM cytotoxicity assay of NK cells

1×10^6 target cells (B16-F10 and Hepa1-6) were suspended in 1ml DPBS containing 5

µM Calcein-AM (BioLegend) and incubated at 37℃ for 30 minutes in the dark. The target

cells were then washed by RPMI for three times and seeded at 0.3×10^5 cells per well on

V-bottom 96-well plate. NK cells from the spleen were sorted by the surface markers, NK1.1$^+$,

CD3$^-$, B220$^-$ and NKp46$^+$ (Table S1), centrifuged at 500 g for 5 minutes, and cultured with

IL-15 in RPMI. They were co-cultured with the target cells through different E (Effector): T

(Target) ratios (0.5:1, 1:1, 3:1 and 9:1). The plates were centrifuged at 250 g for 1 minute to

accelerate the contact of NK cells and target cells and then co-cultured for 4hrs at 37 ℃.

Spontaneous release of Calcein-AM was measured in the target cells without NK cells and the

maximal release was determined by complete lysis of the target cells in RPMI containing 2%

Triton-X 100. After co-culturing, the plates were centrifuged at 120 g for 1 minute and 100

µl/ well of the supernatant was transferred to 96-well assay plate (Corning). The 488/520 nm

values were determined by EnSpire (PerkinElmer).

Bead-based multiplex assay of serum cytokines

Serum samples were obtained via submandibular blood collection and allowed to clot in

uncoated tubes for two hours at room temperature. The tubes were centrifuged at 6,000 rpm

and the supernatants were collected for cytokine analysis by bead-based multiplex assay

(MILLIPLEX MAP Mouse High Sensitivity T Cell Panel, Millipore) following the

manufacturer protocol[27].

Protein extraction

The cell pellets were resuspended in protein extraction buffer (20 mM HEPES, 0.2%

SDS, 1 mM EDTA, 1 mM glycerophosphate, 1 mM Na_3VO_4, and 2.5 mM $Na_4P_2O_7$) with

protease inhibitor cocktail (Sigma-Aldrich) and 1 mM phenylmethylsulfonyl fluoride (PMSF).

The lysates were further homogenized using a Bioruptor (Diagenode) at 4 °C for 15 min, and

then centrifuged at 14,000 × g at 4 °C for 20 min. The supernatant was transferred to a new

tube before determining protein concentration by means of BCA protein assay (Pierce,

Thermo Fisher). Protein aliquots were stored at -30 °C until use.

 In-solution digestion

Protein solutions were first diluted with 50 mM ammonium bicarbonate (ABC) and

reduced with 5 mM dithiothreitol (DTT, Merck) at 60 °C for 45 min, followed by

cysteine-blocking with 10 mM iodoacetamide (IAM, Sigma) at 25°C for 30 min. The samples

were then diluted with 25 mM ABC and digested with sequencing-grade modified porcine

trypsin (Promega) at 37 °C for 16 h. The peptides were desalted using a homemade C18

microcolumn (SOURCE 15RPC, GE Healthcare) and stored at -30 °C until use.

LC-MS/MS analysis

The desalted peptides were diluted in HPLC buffer A (0.1% formic acid in 30%

acetonitrile) and loaded onto a homemade SCX column (0.6 × 5 mm, Luna 5 μm SCX 100 Å,

Phenomenex). The eluted peptides were then trapped in a reverse-phase column (Zorbax

300SB-C18, 0.3 × 5 mm; Agilent Technologies), and separated on a homemade column

(HydroRP 2.5 μm, 75 μm I.D. × 15 cm with a 15 μm tip) using a multi-step gradient of HPLC

buffer B (99.9% acetonitrile/0.1% formic acid) for 90 min with a flow rate of 0.3 μl/min. The

LC apparatus was coupled to a 2D linear ion trap mass spectrometer (Orbitrap Elite ETD;

Thermo Fisher) operated using Xcalibur 2.2 software (Thermo Fisher). Full-scan MS was

performed in the Orbitrap over a range of 400 to 2,000 Da and a resolution of 120,000 at m/z

400. Internal calibration was performed using the ion signal of $[Si(CH_3)_2O]6H^+$ at m/z

536.165365 as lock mass. The 20 data-dependent MS/MS scan events were followed by one

MS scan for the 20 most abundant precursor ions in the preview MS scan. The m/z values

selected for MS/MS were dynamically excluded for 40 sec with a relative mass window of 10

ppm. The electrospray voltage was set to 2.0 kV, and the temperature of the capillary was set

to 200 °C. MS and MS/MS automatic gain control was set to 1,000 ms (full scan) and 200 ms

(MS/MS), or to 3 × 106 ions (full scan) and 3,000 ions (MS/MS), for maximum accumulated

time or ions, respectively.

Protein identification

Data analysis was carried out using Proteome Discoverer software (version 1.4, Thermo

Fisher Scientific). The MS/MS spectra were searched against the SwissProt database using

the Mascot search engine (Matrix Science, version 2.5). For peptide identification, 10 ppm

mass tolerance was permitted for intact peptide masses, and 0.5 Da for CID fragment ions

with an allowance for two missed cleavages arising from trypsin digestion, oxidized

methionine and acetyl (protein N-terminal) as variable modifications, and carbamidomethyl

(cysteine) as a static modification. Peptide spectrum matches (PSM) were then filtered based
on high confidence and a Mascot search engine ranking of 1 for peptide identification to
ensure an overall false discovery rate <0.01. Proteins with single peptide hits were removed
from further analysis.

Gene Set Enrichment Analysis

The absolute abundance of each peptide was calculated from respective peak intensity
based on the PSM abundance. The protein abundance of each sample was calculated from the
sum of the peptide abundance. The abundance data were then background-corrected and
normalized according to variance stabilizing transformation by using the function
"normalize_vsn" in the R package *DEP*[70]. Differential expression across groups was
determined using the function "test_diff" based on protein-wise linear models combined with
empirical Bayes statistics. Significantly differentially-expressed proteins were determined
according to a P-value threshold of 0.01 and a fold-change (FC) >1.5. To establish functional
pathways enriched across groups, normalized data for each pair of compared groups were
used to perform Gene Set Enrichment Analysis (GSEA v4.2.0)[71] on selected MSigDB gene
sets, including Hallmark (H), curated (C2), and immunologic signature (C7) gene sets, by
using the default parameters. Normalized enrichment scores (NES) were used to plot a
heatmap in the R package *pheatmap* (v1.0.12).

Statistical analysis

Data are shown as mean ± standard deviation (SD) or standard error of the mean (SEM).
Comparisons of data under different experimental conditions were carried out using
GraphPad Prism 6.0 software (GraphPad). Each error bar represents SEM unless otherwise
indicated. Significant differences in tumor growth on mouse lungs were assessed by Student's
t test. A difference between groups was considered statistically significant when the p value
was lower than 0.05.

References

1.   Aman Y, Schmauck-Medina T, Hansen M, et al. Autophagy in healthy aging and
disease. *Nat Aging*. Aug 2021;1(8):634-650. doi:10.1038/s43587-021-00098-4
2.   López-Otín C, Blasco MA, Partridge L, Serrano M, Kroemer G. The hallmarks
of aging. *Cell*. Jun 6 2013;153(6):1194-217. doi:10.1016/j.cell.2013.05.039

3.   Bashor CJ, Hilton IB, Bandukwala H, Smith DM, Veiseh O. Engineering the
next generation of cell-based therapeutics. *Nat Rev Drug Discov*. Sep
2022;21(9):655-675. doi:10.1038/s41573-022-00476-6

4.   Fontana L, Partridge L. Promoting health and longevity through diet: from model
organisms to humans. *Cell*. Mar 26 2015;161(1):106-118.
doi:10.1016/j.cell.2015.02.020

5.   Longo VD, Anderson RM. Nutrition, longevity and disease: From molecular
mechanisms to interventions. *Cell*. Apr 28 2022;185(9):1455-1470.
doi:10.1016/j.cell.2022.04.002

6.   Mitchell E, Spencer Chapman M, Williams N, et al. Clonal dynamics of
haematopoiesis across the human lifespan. *Nature*. Jun 2022;606(7913):343-350.
doi:10.1038/s41586-022-04786-y

7.   Pálovics R, Keller A, Schaum N, et al. Molecular hallmarks of heterochronic
parabiosis at single-cell resolution. *Nature*. Mar 2022;603(7900):309-314.
doi:10.1038/s41586-022-04461-2

8.   Seita J, Weissman IL. Hematopoietic stem cell: self-renewal versus
differentiation. *Wiley Interdiscip Rev Syst Biol Med*. Nov-Dec 2010;2(6):640-53.
doi:10.1002/wsbm.86

9.   Montazersaheb S, Ehsani A, Fathi E, Farahzadi R. Cellular and Molecular
Mechanisms Involved in Hematopoietic Stem Cell Aging as a Clinical Prospect. *Oxid*
Med Cell Longev. 2022;2022:2713483. doi:10.1155/2022/2713483

10. Frontelo P, Manwani D, Galdass M, et al. Novel role for EKLF in
megakaryocyte lineage commitment. *Blood*. Dec 1 2007;110(12):3871-80.
doi:10.1182/blood-2007-03-082065

11. Nishizawa M, Chonabayashi K, Nomura M, et al. Epigenetic Variation between
Human Induced Pluripotent Stem Cell Lines Is an Indicator of Differentiation
Capacity. *Cell Stem Cell*. Sep 1 2016;19(3):341-54. doi:10.1016/j.stem.2016.06.019

12. Teruya S, Okamura T, Komai T, et al. Egr2-independent, Klf1-mediated
induction of PD-L1 in CD4(+) T cells. *Sci Rep*. May 4 2018;8(1):7021.
doi:10.1038/s41598-018-25302-1

13. Perkins A, Xu X, Higgs DR, et al. Krüppeling erythropoiesis: an unexpected
broad spectrum of human red blood cell disorders due to KLF1 variants. *Blood*. Apr
14 2016;127(15):1856-62. doi:10.1182/blood-2016-01-694331

14. Neuwirtova R, Fuchs O, Holicka M, et al. Transcription factors Fli1 and EKLF

in the differentiation of megakaryocytic and erythroid progenitor in 5q- syndrome and

in Diamond-Blackfan anemia. *Ann Hematol*. Jan 2013;92(1):11-8.

doi:10.1007/s00277-012-1568-1

15. Luo Q, Ma X, Wahl SM, Bieker JJ, Crossley M, Montaner LJ. Activation and

repression of interleukin-12 p40 transcription by erythroid Kruppel-like factor in

macrophages. *J Biol Chem*. Apr 30 2004;279(18):18451-6.

doi:10.1074/jbc.M400320200

16. Hung CH, Wang KY, Liou YH, et al. Negative Regulation of the Differentiation

of Flk2(-) CD34(-) LSK Hematopoietic Stem Cells by EKLF/KLF1. *Int J Mol Sci*.

Nov 10 2020;21(22)doi:10.3390/ijms21228448

17. Siatecka M, Bieker JJ. The multifunctional role of EKLF/KLF1 during

erythropoiesis. *Blood*. Aug 25 2011;118(8):2044-54.

doi:10.1182/blood-2011-03-331371

18. Pilon AM, Ajay SS, Kumar SA, et al. Genome-wide ChIP-Seq reveals a

dramatic shift in the binding of the transcription factor erythroid Kruppel-like factor

during erythrocyte differentiation. *Blood*. Oct 27 2011;118(17):e139-48.

doi:10.1182/blood-2011-05-355107

19. Tallack MR, Magor GW, Dartigues B, et al. Novel roles for KLF1 in

erythropoiesis revealed by mRNA-seq. *Genome Res*. Dec 2012;22(12):2385-98.

doi:10.1101/gr.135707.111

20. Shyu YC, Lee TL, Chen X, et al. Tight regulation of a timed nuclear import

wave of EKLF by PKCθ and FOE during Pro-E to Baso-E transition. *Dev Cell*. Feb

24 2014;28(4):409-22. doi:10.1016/j.devcel.2014.01.007

21. Shyu YC, Liao PC, Huang TS, et al. Genetic Disruption of KLF1 K74

SUMOylation in Hematopoietic System Promotes Healthy Longevity in Mice. *Adv*

Sci (Weinh). Sep 2022;9(25):e2201409. doi:10.1002/advs.202201409

22. Vaseghi G, Dana N, Ghasemi A, Abediny R, Laher I, Javanmard SH. Morphine

promotes migration and lung metastasis of mouse melanoma cells. *Braz J Anesthesiol*.

Jul-Aug 2023;73(4):441-445. doi:10.1016/j.bjane.2021.10.019

23. Díaz-García VM, Guerrero S, Díaz-Valdivia N, et al. Biomimetic quantum

dot-labeled B16F10 murine melanoma cells as a tool to monitor early steps of lung

metastasis by in vivo imaging. *Int J Nanomedicine*. 2018;13:6391-6412.

doi:10.2147/ijn.S165565

24.   Imado T, Iwasaki T, Kataoka Y, et al. Hepatocyte growth factor preserves

graft-versus-leukemia effect and T-cell reconstitution after marrow transplantation.

Blood. Sep 1 2004;104(5):1542-9. doi:10.1182/blood-2003-12-4309

25.   Iwai Y, Hamanishi J, Chamoto K, Honjo T. Cancer immunotherapies targeting

the PD-1 signaling pathway. *J Biomed Sci*. Apr 4 2017;24(1):26.

doi:10.1186/s12929-017-0329-9

26.   Jiang X, Wang J, Deng X, et al. Role of the tumor microenvironment in

PD-L1/PD-1-mediated tumor immune escape. *Mol Cancer*. Jan 15 2019;18(1):10.

doi:10.1186/s12943-018-0928-4

27.   Aira C, Ruiz T, Dixon L, Blome S, Rueda P, Sastre P. Bead-Based Multiplex

Assay for the Simultaneous Detection of Antibodies to African Swine Fever Virus

and Classical Swine Fever Virus. *Front Vet Sci*. 2019;6:306.

doi:10.3389/fvets.2019.00306

28.   Menees KB, Earls RH, Chung J, et al. Sex- and age-dependent alterations of

splenic immune cell profile and NK cell phenotypes and function in C57BL/6J mice.

Immun Ageing. Jan 8 2021;18(1):3. doi:10.1186/s12979-021-00214-3

29.   Ul-Haq Z, Naz S, Mesaik MA. Interleukin-4 receptor signaling and its binding

mechanism: A therapeutic insight from inhibitors tool box. *Cytokine Growth Factor*

Rev. Dec 2016;32:3-15. doi:10.1016/j.cytogfr.2016.04.002

30.   Boccardi V, Westman E, Pelini L, et al. Differential Associations of IL-4 With

Hippocampal Subfields in Mild Cognitive Impairment and Alzheimer's Disease.

Front Aging Neurosci. 2018;10:439. doi:10.3389/fnagi.2018.00439

31.   Rea IM, Gibson DS, McGilligan V, McNerlan SE, Alexander HD, Ross OA.

Age and Age-Related Diseases: Role of Inflammation Triggers and Cytokines. *Front*

Immunol. 2018;9:586. doi:10.3389/fimmu.2018.00586

32.   Fabian DK, Fuentealba M, Dönertaş HM, Partridge L, Thornton JM. Functional

conservation in genes and pathways linking ageing and immunity. *Immun Ageing*.

May 14 2021;18(1):23. doi:10.1186/s12979-021-00232-1

33.   Yeo EJ. Hypoxia and aging. *Exp Mol Med*. Jun 20 2019;51(6):1-15.

doi:10.1038/s12276-019-0233-3

34.   Zhu X, Chen Z, Shen W, et al. Inflammation, epigenetics, and metabolism

converge to cell senescence and ageing: the regulation and intervention. *Signal*

Transduct Target Ther. Jun 28 2021;6(1):245. doi:10.1038/s41392-021-00646-9

35.   Hamarsheh S, Groß O, Brummer T, Zeiser R. Immune modulatory effects of

oncogenic KRAS in cancer. *Nat Commun*. Oct 28 2020;11(1):5439.

doi:10.1038/s41467-020-19288-6

36.   Ni L, Lu J. Interferon gamma in cancer immunotherapy. *Cancer Med*. Sep

2018;7(9):4509-4516. doi:10.1002/cam4.1700

37.   Principe DR, Doll JA, Bauer J, et al. TGF-β: duality of function between tumor

prevention and carcinogenesis. *J Natl Cancer Inst*. Feb 2014;106(2):djt369.

doi:10.1093/jnci/djt369

38.   Zitvogel L, Galluzzi L, Kepp O, Smyth MJ, Kroemer G. Type I interferons in

anticancer immunity. *Nat Rev Immunol*. Jul 2015;15(7):405-14. doi:10.1038/nri3845

39.   Zhao G, Zhu G, Huang Y, et al. IL-6 mediates the signal pathway of

JAK-STAT3-VEGF-C promoting growth, invasion and lymphangiogenesis in gastric

cancer. *Oncol Rep*. Mar 2016;35(3):1787-95. doi:10.3892/or.2016.4544

40.   Roncagalli R, Taylor JE, Zhang S, et al. Negative regulation of natural killer cell

function by EAT-2, a SAP-related adaptor. *Nat Immunol*. Oct 2005;6(10):1002-10.

doi:10.1038/ni1242

41.   Yu Y. The Function of NK Cells in Tumor Metastasis and NK Cell-Based

Immunotherapy. *Cancers (Basel)*. Apr 16 2023;15(8)doi:10.3390/cancers15082323

42.   Ferreira FM, Palle P, Vom Berg J, Prajwal P, Laman JD, Buch T. Bone marrow

chimeras-a vital tool in basic and translational research. *J Mol Med (Berl)*. Jul

2019;97(7):889-896. doi:10.1007/s00109-019-01783-z

43.   Bin-Jumah MN, Nadeem MS, Gilani SJ, et al. Genes and Longevity of Lifespan.

Int J Mol Sci. Jan 28 2022;23(3)doi:10.3390/ijms23031499

44.   Hofmann JW, Zhao X, De Cecco M, et al. Reduced expression of MYC

increases longevity and enhances healthspan. *Cell*. Jan 29 2015;160(3):477-88.

doi:10.1016/j.cell.2014.12.016

45.   Ocampo A, Reddy P, Martinez-Redondo P, et al. In Vivo Amelioration of

Age-Associated Hallmarks by Partial Reprogramming. *Cell*. Dec 15

2016;167(7):1719-1733.e12. doi:10.1016/j.cell.2016.11.052

46.   Varady KA, Cienfuegos S, Ezpeleta M, Gabel K. Clinical application of

intermittent fasting for weight loss: progress and future directions. *Nat Rev*

Endocrinol. May 2022;18(5):309-321. doi:10.1038/s41574-022-00638-x

47. Hifumi T, Yamamoto A, Ato M, et al. Clinical Serum Therapy: Benefits,

Cautions, and Potential Applications. *Keio J Med*. Dec 25 2017;66(4):57-64.

doi:10.2302/kjm.2016-0017-IR

48. Morgan RA, Gray D, Lomova A, Kohn DB. Hematopoietic Stem Cell Gene

Therapy: Progress and Lessons Learned. *Cell Stem Cell*. Nov 2 2017;21(5):574-590.

doi:10.1016/j.stem.2017.10.010

49. Giannaccare G, Carnevali A, Senni C, Logozzo L, Scorcia V. Umbilical Cord

Blood and Serum for the Treatment of Ocular Diseases: A Comprehensive Review.

Ophthalmol Ther. Jun 2020;9(2):235-248. doi:10.1007/s40123-020-00239-9

50. Wu SY, Fu T, Jiang YZ, Shao ZM. Natural killer cells in cancer biology and

therapy. *Mol Cancer*. Aug 6 2020;19(1):120. doi:10.1186/s12943-020-01238-x

51. Daver N, Wei AH, Pollyea DA, Fathi AT, Vyas P, DiNardo CD. New directions

for emerging therapies in acute myeloid leukemia: the next chapter. *Blood Cancer J*.

Oct 30 2020;10(10):107. doi:10.1038/s41408-020-00376-1

52. Mora J. Autologous Stem-Cell Transplantation for High-Risk Neuroblastoma:

Historical and Critical Review. *Cancers (Basel)*. May 24

2022;14(11)doi:10.3390/cancers14112572

53. Conboy IM, Conboy MJ, Wagers AJ, Girma ER, Weissman IL, Rando TA.

Rejuvenation of aged progenitor cells by exposure to a young systemic environment.

Nature. Feb 17 2005;433(7027):760-4. doi:10.1038/nature03260

54. Das MM, Godoy M, Chen S, et al. Young bone marrow transplantation preserves

learning and memory in old mice. *Commun Biol*. 2019;2:73.

doi:10.1038/s42003-019-0298-5

55. Goodell MA, Rando TA. Stem cells and healthy aging. *Science*. Dec 4

2015;350(6265):1199-204. doi:10.1126/science.aab3388

56. Kang S, Moser VA, Svendsen CN, Goodridge HS. Rejuvenating the blood and

bone marrow to slow aging-associated cognitive decline and Alzheimer's disease.

Commun Biol. Feb 13 2020;3(1):69. doi:10.1038/s42003-020-0797-4

57. Castellano JM, Mosher KI, Abbey RJ, et al. Human umbilical cord plasma

proteins revitalize hippocampal function in aged mice. *Nature*. Apr 27

2017;544(7651):488-492. doi:10.1038/nature22067

58. Mehdipour M, Skinner C, Wong N, et al. Rejuvenation of three germ layers

tissues by exchanging old blood plasma with saline-albumin. *Aging (Albany NY)*. May

30 2020;12(10):8790-8819. doi:10.18632/aging.103418

59. Siatecka M, Xue L, Bieker JJ. Sumoylation of EKLF promotes transcriptional

repression and is involved in inhibition of megakaryopoiesis. *Mol Cell Biol*. Dec

2007;27(24):8547-60. doi:10.1128/mcb.00589-07

60. Aunan JR, Cho WC, Søreide K. The Biology of Aging and Cancer: A Brief

Overview of Shared and Divergent Molecular Hallmarks. *Aging Dis*. Oct

2017;8(5):628-642. doi:10.14336/ad.2017.0103

61. Lages CS, Lewkowich I, Sproles A, Wills-Karp M, Chougnet C. Partial

restoration of T-cell function in aged mice by in vitro blockade of the PD-1/PD-L1

pathway. *Aging Cell*. Oct 2010;9(5):785-98. doi:10.1111/j.1474-9726.2010.00611.x

62. Schaum N, Lehallier B, Hahn O, et al. Ageing hallmarks exhibit organ-specific

temporal signatures. *Nature*. Jul 2020;583(7817):596-602.

doi:10.1038/s41586-020-2499-y

63. Aira C, Ruiz T, Dixon L, Blome S, Rueda P, Sastre P. Bead-Based Multiplex

Assay for the Simultaneous Detection of Antibodies to African Swine Fever Virus

and Classical Swine Fever Virus. Original Research. *Frontiers in Veterinary Science*.

2019-September-13 2019;6doi:10.3389/fvets.2019.00306

64. Zhou J, Zhu Z, Bai C, Sun H, Wang X. Proteomic profiling of lymphocytes in

autoimmunity, inflammation and cancer. *J Transl Med*. Jan 7 2014;12:6.

doi:10.1186/1479-5876-12-6

65. Zhang D, Chen G, Manwani D, et al. Neutrophil ageing is regulated by the

microbiome. *Nature*. Sep 24 2015;525(7570):528-32. doi:10.1038/nature15367

66. Narasimhan PB, Eggert T, Zhu YP, et al. Patrolling Monocytes Control NK Cell

Expression of Activating and Stimulatory Receptors to Curtail Lung Metastases. *J

Immunol*. Jan 1 2020;204(1):192-198. doi:10.4049/jimmunol.1900998

67. Han Z, Yang D, Trivett A, Oppenheim JJ. Therapeutic vaccine to cure large

mouse hepatocellular carcinomas. *Oncotarget*. Aug 8 2017;8(32):52061-52071.

doi:10.18632/oncotarget.19367

68. Daftari P, Gavva NR, Shen CK. Distinction between AP1 and NF-E2

factor-binding at specific chromatin regions in mammalian cells. *Oncogene*. Sep 23

1999;18(39):5482-6. doi:10.1038/sj.onc.1202916

69. Green MR, Sambrook J. *Molecular Cloning: A Laboratory Manual*. Cold Spring

Harbor Laboratory Press; 2012.

70.   Zhang X, Smits AH, van Tilburg GB, Ovaa H, Huber W, Vermeulen M.

Proteome-wide identification of ubiquitin interactions using UbIA-MS. *Nat Protoc*.

Mar 2018;13(3):530-550. doi:10.1038/nprot.2017.147

71.   Subramanian A, Tamayo P, Mootha VK, et al. Gene set enrichment analysis: a

knowledge-based approach for interpreting genome-wide expression profiles. *Proc*

Natl Acad Sci U S A. Oct 25 2005;102(43):15545-50. doi:10.1073/pnas.0506580102

<u>Figure Legends</u>

Figure 1. <u>Anti-cancer capability of *Eklf*(K74R) mice as analyzed by the experimental</u>

<u>melanoma metastasis assay.</u>

(A) Flow chart illustrating the strategy of the pulmonary tumor foci assay. Left

panels, representative photographs of pulmonary metastatic foci on the lungs of

WT and *Eklf*(K74R) male mice in the B6 background two weeks after intravenous

injection of B16-F10 cells (10^5 cells/ mouse). Statistical comparison of the

numbers of pulmonary foci is shown in the two histograms on the right. N=10

(male) and N=7 (female), **, $p<0.01$. Note that only the numbers of large

pulmonary foci (>1mm diameter) were scored. N>6, **, $p<0.01$. (B) Pulmonary

tumor foci assay of 24 month-old WT and *Eklf*(K74R) male mice. Statistical

comparison is shown in the two histograms. N=10 (male), *, $p<0.05$. (C)

Pulmonary tumor foci assay of male mice in the FVB background. Statistical

comparison is shown in the histograph on the right. N=10, **, $p<0.01$. (D)

Pulmonary tumor foci assay of *Eklf*(K74A) male mice. Statistical comparison of

the 3 month-old WT and *Eklf*(K74A) mice numbers of pulmonary foci is shown in

the two histograms. N=10 (male), **, $p<0.01$. (E) The *Eklf*(K74R) mice and WT

mice were subcutaneously injected with Hepa1-6 cells to form tumors. The tumor

volumes were measured once a week using the formula: length x weight x height x

$\pi/6$. The curves of tumor growth started to show difference between the

Eklf(K74R) and WT mice at 49 days after Hepa1-6 cell injection. N>6, *$p < 0.05$,

**$p < 0.01$ and ***$p < 0.001$.

Figure 2. <u>Transfer of cancer resistance of *Eklf*(K74R) mice to WT mice by bone marrow transplantation (BMT)</u>

(A) Flow chart illustrating the experimental strategy. (B) FACS analysis of the efficiency of BMT with use of 10Gy γ-irradiation. The percentages of CD45.1/CD45.2 cells in the PB of the recipient male mice were analyzed by flow cytometry, with the representative FACS charts shown on the left and the statistical histobar diagram on the right. (C) Transfer of the anti-metastasis capability of 8 week-old *Eklf*(K74R) male mice to age-equivalent WT male mice by BMT with use of 10Gy γ-irradiation. Left panels, representative photographs of lungs with pulmonary metastatic foci in the recipient WT (CD45.1) mice after BMT from WT (CD45.2) or *Eklf*(K74R) (CD45.2) donor mice and challenged with B16-F10 cells. Statistical analysis of the numbers of pulmonary B16-F10 metastatic foci on the lungs is shown in the right histogram. n=10, *, p<0.05. (D) Transplantation of 8 week-old male WT (CD45.1) mice with BMMNC from age-equivalent WT (CD45.2) male mice or from *Eklf*(K74R) (CD45.2) male mice with use of the γ-irradiation dosage 2.5Gy or 5Gy. The histobar diagram comparing the percentages of CD45.1 and CD45.2 PB cells of the recipient WT mice after BMT is shown on the left. The statistical analysis of the average numbers of pulmonary foci on the lungs of recipient WT mice after BMT and injected with the B16-F10 cells is shown in the right histogram, N=6. **, p<0.01, ***, p<0.001.

Figure 3. <u>Inhibition of tumor growth in WT mice by BMT from *Eklf*(K74R) mice</u>

(A) A flow chart of the experiments. Luciferase-positive B16-F10 cells were injected into the tail vein of 8 week-old WT male mice (day 0). The mice were then transplanted with BMMNC from WT or *Eklf*(K74R) male mice on day 11 after the luciferase-positive B16-F10 cell injection. *In vivo* imaging system (IVIS) was used to follow the tumor growth in mice on day 0, 10, 17 and 24, respectively. (B) Representative images of bioluminescence reflecting the luciferase activity from melanoma cancer cells in mice. The color bar indicates the scale of the bioluminescence intensity. (C) Statistical analysis of the intensities of bioluminescence in the cancer-bearing mice (WT→WT, purple, N=7; *Eklf*(K74R)→WT, blue, N=8; Control (no BMT), red, N=3).

Figure 4. <u>Decrease of *Pd-1* and *Pd-l1* expression in blood cells of *Eklf*(K74R) mice</u>

(A) Levels of *Pd-1* and *Pd-l1* mRNAs in the PB of WT and *Eklf*(K74R) male mice at the ages of 3 months and 24 months, respectively, as analyzed by RT-qPCR. Note the relatively low levels of *Pd-1* and *Pd-l1* mRNAs in the *Eklf*(K74R) mice at both ages in comparison to the WT mice. (B) Upper panels, comparison of the mRNA levels of *Pd-1* and *Pd-l1* of CD3$^+$ T cells and B220$^+$ B cells isolated from the PB of 8 week-old WT and *Eklf*(K74R) male mice. N=5. *, p<0.05; **, p<0.01. Lower panels, comparison of the protein levels of PD-1 and PD-L1, as analyzed by flow cytometry, of CD3$^+$ T cells and B220$^+$ B cells from 8 week-old WT and *Eklf*(K74R) male mice. N=3. *, p<0.05; **, p<0.01. (C) Comparison of the levels of *Pd-1, Pd-l1* and *Eklf* mRNAs, as analyzed by RT-qPCR, in CD3$^+$ T cells, which were isolated from splenocytes, without or with RNAi knockdown of *Eklf* mRNA. Two oligos (oligo-1 and oligo-2) were used to knockdown *Eklf* mRNA by ~60-70%, which resulted in the reduction of *Pd-1* mRNA level by 30-60% and nearly complete depletion of *Pd-l1* mRNA. Control, T cells transfected with GFP-plasmid. SC, T cells transfected with scrambled oligos. N>3. *, p<0.05; **, p<0.01; ***, p<0.001.

Figure 5. <u>Proteomics analysis of mouse leukocytes</u>

(A) Heatmap plots representing age-dependent differentially expressed proteins (DEPs) in the leukocytes from the WT male mice (left) and *Eklf*(K74R) male mice (right). Differential Test Threshold: expression fold change > 1.5 and *p*-value < 0.01. (B) and (C), pathway analysis of the age-dependent DEPs changed in the concordant (B) and reverse (C) directions, respectively, in the WT and *Eklf*(K74R) mice. NES, normalized enrichment score. (D) Heatmap plots representing strain-dependent DEPs in leukocytes of 3 month-old (left) and 24 month-old (right) mice. (E) Pathway analyses of the strain-dependent DEPs in elder WT leukocytes (enrichment indicated by blue color) and elder *Eklf*(K74R) leukocytes (enrichment indicated by red color). The aging- and cancer-associated pathways are presented in the left and right diagrams, respectively. NES, normalized enrichment score. (F) A model of the regulation of the mouse lifespan by different cellular pathways in the leukocytes. The inter-relationship of 11 aging-associated cellular pathways differentially expressed in the leukocytes of *Eklf*(K74R) male mice in comparison

to the WT male mice, among themselves and with respect to the regulation of

lifespan (for references, see text), are depicted here. The directions and lengths of

the arrows indicate changes of the individual pathways, up-regulation (upward

arrows) or repression (downward arrows), in the leukocytes during aging of the WT

male mice (blue arrows) and *Eklf*(K74R) male mice (red arrows), respectively.

Figure 6. <u>Higher *in vitro* cancer cell cytotoxicity of NK(K74R) cells than WT NK cells</u>

B16-F10 and Hepa1-6 cells were labeled with Calcein-AM (BioLegend) and

co-cultured with NK cells in different E:T ratio for 4hrs. Killing rates of B16-F10

and Hepa 1-6 cancer cells by NK cells were determined by the intensities of

fluorescence in the medium in comparison to the controls of spontaneous release

and maximal release.

Fig. 1

(A)

(B) B6 (24-month male)

(C) FVB (Male)

(D) *Eklf* (K74A) (Male)

(E) Solid-tumor formation (Hepa1-6)

(A)

(B)

(C)

(A)

(B)

(C)

Fig. 5