**Luteinizing hormone stimulates ingression of mural granulosa cells**

**within the mouse preovulatory follicle**

Corie M. Owen* and Laurinda A. Jaffe*

Department of Cell Biology, University of Connecticut Health Center, Farmington, CT 06030

USA

***Correspondence**: Department of Cell Biology, University of Connecticut Health Center, 263

Farmington Ave., Farmington, CT 06030 USA. Email: coowen@uchc.edu, ljaffe@uchc.edu.

**Running title**: Ingression of granulosa cells in ovarian follicles

**Summary sentence**: LH-induced ingression of LH receptor-expressing cells within the mural

granulosa layer of the ovarian follicle is a new component in the complex sequence of structural

changes that lead to ovulation.

**Keywords:** Ovarian follicle, luteinizing hormone receptor, granulosa cell, ingression, ovulation,

migration

**Abbreviations**: LH (luteinizing hormone), LHR (luteinizing hormone receptor), HA-LHR

(hemagglutinin-tagged luteinizing hormone receptor), eCG (equine chorionic gonadotropin),

hCG (human chorionic gonadotropin)

**Grant Support**: This work was supported by the *Eunice Kennedy Shriver* National Institute of

Child Health and Human Development (R37 HD014939 to L.A.J., and F31 HD107918 to

C.M.O.).

## Abstract

Luteinizing hormone (LH) induces ovulation by acting on its receptors in the mural granulosa cells that surround a mammalian oocyte in an ovarian follicle. However, much remains unknown about how activation of the LH receptor modifies the structure of the follicle such that the oocyte is released and the follicle remnants are transformed into the corpus luteum. The present study shows that the preovulatory surge of LH stimulates LH receptor-expressing granulosa cells, initially located almost entirely in the outer layers of the mural granulosa, to rapidly extend inwards, intercalating between other cells. The cellular ingression begins within 30 minutes of the peak of the LH surge, and the proportion of LH receptor-expressing cell bodies in the inner half of the mural granulosa layer increases until the time of ovulation, which occurs at about 10 hours after the LH peak. During this time, many of the initially flask-shaped cells appear to detach from the basal lamina, acquiring a rounder shape with multiple filipodia. Starting at about 4 hours after the LH peak, the mural granulosa layer at the apical surface of the follicle where ovulation will occur begins to thin, and the basolateral surface develops invaginations and constrictions. Our findings raise the question of whether LH stimulation of granulosa cell ingression may contribute to these changes in the follicular structure that enable ovulation.

## Introduction

Preovulatory mammalian follicles are comprised of many concentric layers of cells. Directly around the oocyte are cumulus cells, and outside of these are a fluid-filled antrum, mural granulosa cells, a basal lamina, and a theca layer including steroidogenic cells, fibroblasts, smooth muscle cells, and vasculature [1] (Figure 1A). Ovulation is triggered when luteinizing hormone (LH), which is released from the pituitary and delivered to the ovary through blood vessels in the theca layer, acts on its receptors in a subset of the outer mural granulosa cells [2–8]. LH binding to its receptors initiates a G-protein-coupled signaling cascade that induces meiotic resumption in the oocyte and changes in gene expression in the granulosa cells that lead to ovulation [9–14].

Activation of the LH receptor also causes complex structural changes in the follicle. These include chromosomal and cytoskeletal rearrangements as meiosis progresses in the oocyte [15], secretion of an extracellular matrix from the cumulus cells that causes the cumulus mass to expand [16], and loss of connections between these cells and the oocyte as transzonal projections retract [17]. As ovulation approaches, the mural granulosa and theca layers in the apical region of the follicle, directly adjacent to the surface epithelium where the oocyte will be released, become thinner [18–21], correlated with vasoconstriction and reduced blood flow in the apical region of the theca [21]. The mural granulosa and theca layers in the basolateral region (that outside of the apical region) develop constrictions, and these are thought to contribute to expelling the oocyte and surrounding cumulus cells [22–27]. The preovulatory constrictions may be mediated in part by contraction of smooth muscle cells in the overlying theca [23], in response to endothelin-2 produced by the granulosa cells [26,28,29]. However, granulosa-specific deletion of endothelin-2 in mice only partially inhibits ovulation [30], suggesting that additional factors may also contribute to the follicular constriction. At the time of

ovulation, the basal lamina around the granulosa cells breaks down, and blood vessels from the

theca grow inwards as the remnants of the follicle transform into the corpus luteum, which

produces progesterone to support pregnancy [31].

One unexplored factor that could contribute to the preovulatory changes in follicle structure

is suggested by evidence that LH receptor activation induces migration [32,33] and cytoskeletal

shape changes [34,35] in isolated mural granulosa cells. These cytoskeletal shape changes are

detected within 30 minutes after LH receptor activation [35].

However, LH-stimulated granulosa cell motility has not been investigated in intact ovarian

follicles. By using mice with an HA epitope tag on the endogenous LH receptor such that its

localization could be visualized [8], and by collecting ovaries for imaging at defined times after

eliciting a physiological LH surge by injection of kisspeptin [36], we obtained a precise

timecourse of changes in localization of mural granulosa cells expressing the LH receptor over

the period leading to ovulation. We discovered that LH stimulates granulosa cells that express

its receptor to extend inwards in the follicle, intercalating between other granulosa cells. This

LH-stimulated granulosa cell motility may influence the structure of the follicle as it prepares for

ovulation and transformation into a corpus luteum.

## Materials and Methods

### Mice

Protocols covering the maintenance and experimental use of mice were approved by the Institutional Animal Care Committee at the University of Connecticut Health Center. The mice were housed in a room in which lights were turned on at 6 AM and off at 4 PM. Generation, genotyping, and characterization of mice with an HA tag on the endogenous LH receptor (HA-LHR) have been previously described [8]; homozygotes were used for breeding pairs and for all experiments. The HA-LHR and wildtype mouse background was C57BL/6J. These mice have been deposited at the Mutant Mouse Resource and Research Centers repository at The Jackson Laboratory (Bar Harbor, ME); MMRRC #71301, C57BL/6J-Lhcgr[em1Laj]/Mmjax, JR#038420.

### Hormone injections of prepubertal mice for collection of ovaries and assessment of ovulation

Ovaries containing preovulatory follicles were obtained by intraperitoneal injection of 22–24-day-old mice with 5 I.U. equine chorionic gonadotropin (eCG) (ProSpec #HOR-272). 44 hours later, the mice were injected intraperitoneally with 1 nmol kisspeptin-54 (Cayman Chemical, #24477) to stimulate an endogenous LH surge [36]. Ovaries were dissected at indicated time points after kisspeptin injection.

To assess ovulation, cumulus-oocyte complexes were collected from oviducts dissected after hormone injections as described above. For Figure 1B, some mice were injected with human chorionic gonadotropin (hCG, 5 I.U., ProSpec #HOR-250) instead of kisspeptin. Cumulus-oocyte complexes were dissociated by pipetting and then counted.

**Collection of ovaries from proestrus adult mice before and after the LH surge**

Vaginal cytology of 6-8 week old females was examined daily at approximately 10 AM to

determine the stage of the estrous cycle [37]. Ovaries from proestrus mice were collected at

either 12 noon (pre-LH) or 10 PM (post-LH).

**Preparation of ovary cryosections for immunofluorescence microscopy**

Ovaries were frozen, fixed, cryosectioned, and labelled for immunofluorescence microscopy

as previously described [8]. Sections were 10 μm thick. Antibody sources and concentrations

are listed in Table S1. Equatorial sections of follicles, defined as sections including the oocyte,

were used for imaging.

**Confocal and Airyscan imaging**

Ovarian cryosections were imaged with a confocal microscope (LSM800 or LSM980, Carl

Zeiss Microscopy). To image entire follicle cross-sections, a 20x/0.8 NA Plan-Apochromat

objective was used. Small regions of follicles were imaged using a 63x/1.4 NA Plan-Apochromat

objective with an Airyscan detector to enhance resolution. Images of optical sections were

reconstructed using the 2-dimensional Airyscan processing at standard strength. Except as

indicated, all images were captured using an interval space of 1 μm. Brightness and contrast

were adjusted after analysis in Fiji software [38].

**Image analysis**

Preovulatory follicles were identified in equatorial cryosections by the presence of LH

receptor-expressing cells in their mural granulosa cells. To quantify the percentage of LH

receptor-expressing cell bodies that were located in the inner half of the mural granulosa layer

(Figure 3A,B), inner and outer halves were defined as previously described [8]. In brief, the

width of the mural granulosa layer from antrum to basal lamina was measured at 8 radial

positions for each follicle, and halfway points were marked. These points were then connected

to define the boundary of the inner and outer mural. The basal lamina position was identified by

locating the outer edge of the outermost layer of granulosa cells, or by labeling with an antibody

against laminin gamma 1. LH receptor-expressing cell bodies were identified as DAPI-stained

nuclei that  were surrounded by HA labeling, and were counted in inner and outer mural

granulosa cell regions using the Cell Counter Tool in Fiji [38]. Counts were performed on the

series of 10 optical sections with 1 µm interval space. The percentage of LH receptor-

expressing cells in the inner mural was calculated by dividing the number of LH receptor-

expressing cells in the inner mural by the total number of LH receptor-expressing cells in the

inner and outer mural regions. The percentage of mural granulosa cells that expressed the LH

receptor (Figure 3C) was calculated by dividing the total number LH receptor-expressing cells

by the total number of DAPI-stained nuclei.

Basal lamina invaginations were analyzed using images of equatorial sections in which the

basal lamina was labelled with an antibody against laminin gamma 1 (Table S1). Invaginations

were defined as regions where the basal lamina curved inwards to a depth of 5-100 µm, with a

width of ≤150 µm. The depth of each invagination (Figure 5C) was determined by drawing a line

connecting the 2 points on the basal lamina at which it curved inwards, and then measuring the

distance between that line and the deepest point of the invagination.  For counts of the number

of invaginations per cross-section (Figure 5D), the apical region of the follicle was defined as the

mural granulosa region adjacent to the surface epithelium. The basolateral region was defined

as the rest of the mural granulosa. Both measurements and counts were performed on the

series of 10 optical sections with 1 µm interval space.

To measure the width of the mural granulosa layer in apical and basal regions, the center of

the follicle was located by measuring the height and width of each follicle. A line was then drawn

from the base through the center of the follicle to the apex, and beginning and end points were

marked as the middle of the base and apex. Two points were marked at 50 μm along the basal

lamina to the left and right of each point, and the width of the mural layer was measured at each

point (see right hand panels of Figure 6A). The averages of the three measurements on the

apical and basal sides were used for comparisons (Figure 6B).  Measurements were performed

on maximum projections of a series of 10 optical sections spaced 1 μm apart.

**Western blotting**

For measurement of HA-LHR protein content (Figure S9), ovaries were collected at

designated time points and sonicated in 1% SDS with protease inhibitors [39]. Protein

concentrations were determined with a BCA assay (Thermo Scientific, #23227) and 40 μg of

total protein was loaded per lane. Western blots were probed with an antibody against the HA

epitope, developed using a fluorescent secondary antibody (Table S1), and detected with an

Odyssey imager (LICOR, Lincoln, NE). Blots were co-imaged with the Revert stain for total

protein (LICOR), and HA-LHR fluorescence intensity was normalized to the Revert fluorescence

intensity for each lane. Values were then normalized to that for the ovary without kisspeptin

injection.

**Statistics and graphics**

Analyses were conducted as indicated in the figure legends using Prism 9 (GraphPad

Software, Inc, La Jolla, CA). Values in graphs are presented as mean ± standard error of the

mean (SEM), and values indicated by different letters are significantly different ($P < 0.05$).

Diagrams for the graphical abstract and for Figures 1A,C and 3A were generated using

BioRender.com.

## Results and Discussion

Except as indicated, LH release was stimulated by intraperitoneal injection of kisspeptin into ~25 day old mice that had been injected 44 hours previously with eCG to stimulate follicle growth and LH receptor expression [36]. Kisspeptin is a neuropeptide that causes gonadotropin releasing hormone to be secreted from the hypothalamus, which in turn causes release of LH into the bloodstream [40]. The rise in serum LH peaks ~1.5 hours after kisspeptin injection and is comparable in amplitude and duration to the endogenous LH surge [36]. As previously described, kisspeptin injection causes ovulation [36]. We chose to use kisspeptin rather than the more standard induction of ovulation by injection of human chorionic gonadotropin (hCG), because kisspeptin causes release of mouse LH, closely mimicking the natural ovulatory stimulus. To determine the kinetics of kisspeptin-induced ovulation, cumulus-oocyte complexes in the oviduct were counted at defined time points. With the experimental conditions used here, ovulation occurred between 11 and 12 hours after kisspeptin or hCG injection, and similar numbers of oocytes were released by each stimulus (Figure 1B).

The localization of cells expressing the LH receptor was investigated using a recently developed mouse line with a hemagglutinin (HA) tag on the endogenous LH receptor (HA-LHR), to allow specific immunolocalization of the LH receptor protein [8]. The heterogenous expression of the LH receptor within the mural epithelium allows individual cells to be well visualized [8]. The timing and number of oocytes released in response to kisspeptin were similar comparing HA-LHR and wild-type mice (Figure 1B). Ovaries from HA-LHR mice were collected, fixed, and frozen, either before kisspeptin injection, or at 2 hour intervals afterwards (Figure 1C). Ovary cryosections were then labelled with an HA antibody, and the localization of HA-LHR expressing cells was analyzed.

**LH induces ingression of LH receptor-expressing granulosa cells within the preovulatory follicle**

The mural granulosa region of a mouse preovulatory follicle is comprised of ~5-15 layers of cells, located between the basal lamina and the fluid-filled antrum [8,41,42] (Figures 1A, 2A, S1). Previously, we showed that before LH receptor stimulation, almost all of the LH receptor-expressing cell bodies in the mural granulosa are located in its outer half, and within this region, expression is heterogeneous [8]. Many of the LH receptor-expressing granulosa cells have an elongated flask-like shape, with the cell bodies containing their nuclei located a few cell layers away from the basal lamina and long processes extending back to the basal lamina, forming a pseudostratified epithelium [8,18] (Figures 2A, S1, S10). In the present study, only ~7% of the total LH receptor-expressing granulosa cell bodies were found in the inner half of the mural granulosa region before LH receptor stimulation (Figure 3A,B).

At 2 hours after kisspeptin injection (~30 minutes after the peak of the induced LH surge), the percentage of LH receptor-expressing cell bodies in the inner mural layer had increased to ~21% (Figures 2B, 3B, S2). At later time points after kisspeptin injection, this percentage continued to increase, reaching ~35% at 10 hours (Figures 2C-F, 3B, S3-S6). Ovulation occurred between 11 and 12 hours after kisspeptin (Figures 1B, 2G, S7). Injection with PBS instead of kisspeptin did not change the percentage of LH receptor-expressing cells in the inner mural granulosa layer at 6 hours, indicating that the localization change is dependent on LH (Figures 3B, S8).

The increase in the fraction of LH receptor-expressing cells in the inner vs. outer mural granulosa region in response to the LH surge is most easily explained by migration into the inner region of cells originally in the outer region. This conclusion is consistent with previous studies showing that LH receptor stimulation causes granulosa cells in culture to become migratory [32,33]. Future live imaging of granulosa cell migration in intact follicles, using mice in

which cells expressing the LH receptor are fluorescently tagged, could further test this

conclusion.

Alternatively, the observed increase in the ratio of LH receptors in the inner vs. outer mural

granulosa regions could potentially be a consequence of synthesis of LH receptor mRNA and

protein in the inner mural granulosa region. Our previously published in situ hybridization data

showed no detectable LHR mRNA in the inner mural granulosa cells in follicles prior to the LH

surge [8], so such an explanation would require new transcription of LHR mRNA. In rat ovaries,

LHR mRNA synthesis has not been detected until 24-72 hours after hCG injection [6,43],

making this an unlikely explanation of the relocalization of LHR protein that we observe in

mouse preovulatory follicles beginning within ~30 minutes after the peak of the LH surge (2

hours after kisspeptin injection).

Nevertheless, to examine this alternative hypothesis, we compared the percentage of the

total population of mural granulosa cells that expressed LH receptor protein over the 10-hour

period after kisspeptin injection. No significant increase was seen over this time period, and

mean values at 6 hours after injection of kisspeptin or a control solution of PBS were identical

(Figure 3C). These results are consistent with the conclusion that the LH surge did not stimulate

synthesis of its receptor over this time period. We also measured the LH receptor protein

content of ovaries after kisspeptin injection, using quantitative western blotting, and found that

LH receptor protein levels were unchanged over the 12-hour period after injection (Figure S9). If

new LHR protein synthesis occurred in the inner mural region, we would expect to have seen an

increase in the total LH receptor content of the ovary, unless an equal amount of LH receptor

protein in the outer mural region was proteolyzed at the same time. Thus, new synthesis of LH

receptor protein in the inner mural granulosa layer is an unlikely explanation of the observed

redistribution of LH receptor-expressing cells in the follicle.

**LH induces an epithelial-to-mesenchymal-like transition in LH receptor-expressing granulosa cells within the follicle**

High resolution Airyscan images of ovaries from mice before and after kisspeptin injection showed that by 6 hours after kisspeptin injection, many LH receptor-expressing cells in the follicle interior were rounder compared to the predominantly flask-shaped cells seen near the basal lamina prior to LH receptor stimulation (Figures 3D, S10, S11). At 6 hours, many of the LH receptor-expressing cells had lost a visible attachment to the basal lamina, suggesting that these cells had detached, undergoing an epithelial-to-mesenchymal-like transition (Figures 3D,E, S11). Alternatively, the cellular processes connecting these cell bodies to the basal lamina might have been too thin to see with light microscopy or might not have been contained within the 10 μm thick section. These cells often had numerous filopodia extending in many directions, as well as membrane blebs (Figure 3D,E), consistent with the presence of filopodia and blebs on other migratory cells [44–46].

**The LH surge also induces ingression of LH receptor-expressing cells in preovulatory follicles of naturally cycling adult mice**

To confirm that the ingression of granulosa cells that was seen in hormonally stimulated prepubertal mice also occurred in naturally cycling mice, we collected ovaries from adult mice in proestrus. Based on measurements of serum LH, the peak of the proestrous LH surge occurs close to the time that the lights are turned off in the room where the mice are housed, although the exact time is variable [47]. Mice were determined to be in proestrus based on vaginal cytology, and we collected ovaries at either 4 hours before lights off (pre-LH) or 6 hours after lights off (post-LH). In the pre-LH ovaries, ~11% of LH receptor expressing cells were located in the inner mural, whereas in the post-LH ovaries, the LH receptor-expressing cell bodies had moved inwards, with ~35% of cells expressing the LH receptor localized in the inner mural

(Figures 4A,B, S12). This was comparable to the change in localization that occurred in the

prepubertal mice (Figures 2, 3B), confirming that the ingression occurs during the natural

ovulation process.

**LH induces invaginations of the basolateral surface of the follicle, starting ~6 hours**

**before ovulation.**

The LH-induced ingression of the LH receptor-expressing granulosa cells into the inner half

of the mural layer, and the associated changes in cell shape, raised the question of how these

cellular events might correlate with LH-induced changes in the shape of the follicle as a whole.

Constrictions in the basolateral surface of follicles have been previously observed during or

within 2 hours of ovulation, but reports at earlier time points are lacking [24–29]. By fixing

ovaries before or 6-10 hours after injection of mice with kisspeptin, labelling the basal lamina in

ovary cryosections, and imaging equatorial cross-sections of preovulatory follicles, we observed

that invaginations of the follicle surface began much earlier than previously reported.

In follicles in ovaries from mice that had not been injected with kisspeptin, the basal lamina

was smooth, with only occasional inward deflections (Figure 5A,B). However, at 6-10 hours after

kisspeptin injection, the basal lamina showed numerous invaginations, defined as regions where

the basal lamina curved inwards to a depth of 5-100 μm with a width of ≤150 μm (Figure 5A-C).

The invaginations occurred only in basolateral regions of the follicle, and were seen as early as

6 hours after kisspeptin injection, corresponding to ~4.5 hours after the peak of the LH surge,

and ~6 hours before ovulation (Figures 5A-D, S4B, S5B, S6B). Their number increased

between 6 and 10 hours after kisspeptin injection (Figure 5D). In addition to the invaginations,

the basolateral region of follicles at 8 and 10 hours after kisspeptin injection usually also showed

larger scale constrictions (Figures 5B, S5, S6). Similar invaginations and constrictions of the

basolateral surface were seen in follicles of ovaries after the LH surge in naturally cycling mice

(Figures 4A, S12).

**LH induces thinning of the apical surface of the follicle, starting ~6 hours before**

**ovulation.**

In addition to the basolateral constrictions of the follicle surface, LH stimulation caused

thinning of the mural granulosa layer at the apex of the follicle where the oocyte will be released

(Figures 2, 5A, 6, S4-S7). As previously reported for hamster follicles [20], the apical thinning

begins early, showing a statistically significant decrease in mean thickness by 6 hours after

injection of kisspeptin, and not occurring in the basal region (Figure 6B). At 8 hours after

kisspeptin, many follicles also had a concavity on the apical side, which was wider and

smoother than the invaginations seen in basolateral regions (Figures 5A,B, S5). Apical thinning

was evident in some of the follicles of naturally cycling mice after the LH surge; variability in the

timing of the LH surge with respect to the time of lights off may explain why it was not seen in all

examples (Figures 4A, S12).

**Does LH-induced ingression of LH receptor-expressing mural granulosa cells contribute**

**to LH-induced basolateral invaginations or apical thinning?**

Our findings that basolateral invaginations and apical thinning are both evident by 6 hours

after kisspeptin injection (Figures 5 and 6) raise the question of whether the ingression of LH

receptor-expressing cells might contribute to causing these events. Inwardly extending

granulosa cells could be speculated to pull the attached basal lamina inwards, contributing to

LH-induced invaginations and constrictions of the basolateral follicle surface. Likewise,

migration of granulosa cells out of the apical region of the mural granulosa could be imagined to

contribute to thinning of the follicle apex. While previous studies have provided evidence for the

combined action of extracellular proteases [13,21,27] and endothelin-induced contractions of

smooth muscle cells in the theca [29] and theca blood vessels [21] in causing both basolateral

invaginations and apical thinning, definitive identification of the cells that are responsible

remains uncertain.

We hypothesize that LH-induced ingression of LH receptor-expressing mural granulosa cells

may be an additional factor that contributes to LH-induced changes in follicle shape. In support

of this concept, cell migration has been found to regulate tissue architecture in other

developmental systems [48,49], including *Drosophila* ovarian follicles [50] and the mammalian

placenta [51]. LH-stimulated granulosa cell migration could provide a complementary

mechanism that acts in parallel with proteases [13,21,27], and with contraction of vascular and

non-vascular smooth muscle cells in the theca layer [21,23,24,28–30] to contribute to rupture of

the follicle surface at ovulation.

Importantly, the early changes in follicular shape reported here start around 6 hours after

kisspeptin injection, apparently preceding the synthesis of endothelin mRNA by the granulosa

cells, which has been reported to begin at ~11 hours after injection of mice with the LH receptor

agonist hCG [28]. Based on these kinetics, endothelin-induced responses do not appear to fully

account for the LH-induced changes constrictions in the basolateral region of the follicle and the

thinning of the mural granulosa layer in the apical region, although additional spatial-temporal

analysis of the expression of endothelins and their receptors would be useful.

Understanding of the possible contribution of LH-induced ingression of mural granulosa cells

to causing the invaginations in the basolateral region of the follicle and the thinning of the apical

mural granulosa cell layer could be furthered by knowledge of the cytoskeletal changes that

cause the cells to migrate, such that the migration could be inhibited. Notably, activation of the

actin severing and depolymerizing protein cofilin is required for LH receptor-stimulated

granulosa cell shape changes in vitro [35], suggesting a possible function of cofilin in granulosa

cell ingression within the follicle in vivo. Phosphoproteomic analysis of rat follicles has shown

that LH signaling decreases the phosphorylation of cofilin on serine 3 to ~10% of baseline within

30 minutes [52], and dephosphorylation at this site increases cofilin activity [53]. Cofilin is also

essential for other developmental processes involving cellular extension, such as axon growth

[54]. Mice with genetically modified cofilin or other cytoskeletal proteins in their granulosa cells

could be used to investigate the mechanisms that mediate LH-induced granulosa cell ingression

in preovulatory follicles, and the consequences for follicular shape changes and ovulation.

Happening over a similar time frame as LH-induced changes in follicle shape, LH-induced

ingression of LH receptor-expressing cells within the mural granulosa layer is a new component

in the complex sequence of structural changes in the follicle that occur in the hours preceding

ovulation.

379
380

**Supplementary material**

**Data availability**

All relevant data can be found within the article and its supplementary information.

**Author contributions**

CMO and LAJ designed and carried out the studies, analyzed the data, and wrote the manuscript.

**Conflict of interest**

The authors declare that no conflict of interest exists.

**Acknowledgements**

We thank Deb Kaback and Tracy Uliasz for technical assistance, Lydia Sorokin and Siegmund Budny for generously providing the laminin-gamma 1 antibody, Mark Terasaki, Jeremy Egbert, Rachael Norris, Siu-Pok Yee, Lisa Mehlmann, Valentina Baena, Allan Herbison, Tabea Marx,

## References

[1]  Richards JS, Ren YA, Candelaria N, Adams JE, Rajkovic A. Ovarian follicular theca cell recruitment, differentiation, and impact on fertility: 2017 update. Endocr Rev 2018; 39:1–20.

[2]  Richards JS, Ireland JJ, Rao MC, Bernath GA, Midgley AR, Reichert LE. Ovarian follicular development in the rat: hormone receptor regulation by estradiol, follicle stimulating hormone and luteinizing hormone. Endocrinology 1976; 99:1562–1570.

[3]  Bortolussi M, Marini G, Dal Lago A. Autoradiographic study of the distribution of LH(HCG) receptors in the ovary of untreated and gonadotrophin-primed immature rats. Cell Tissue Res 1977; 183:329–342.

[4]  Bortolussi M, Marini G, Reolon ML. A histochemical study of the binding of 125I-HCG to the rat ovary throughout the estrous cycle. Cell Tissue Res 1979; 197:213–226.

[5]  Uilenbroek JT, Richards JS. Ovarian follicular development during the rat estrous cycle: gonadotropin receptors and follicular responsiveness. Biol Reprod 1979; 20:1159–1165.

[6]  Peng X-R, Hsueh AJW, Lapolt PS, Bjersing L, Ny T. Localization of luteinizing hormone receptor messenger ribonucleic acid expression in ovarian cell types during follicle development and ovulation. Endocrinology 1991; 129:3200–3207.

[7]  Eppig JJ, Wigglesworth K, Pendola FL. The mammalian oocyte orchestrates the rate of ovarian follicular development. PNAS 2002; 99:2890–2894.

[8]  Baena V, Owen CM, Uliasz TF, Lowther KM, Yee S-P, Terasaki M, Egbert JR, Jaffe LA. Cellular heterogeneity of the luteinizing hormone receptor and its

significance for cyclic GMP signaling in mouse preovulatory follicles. Endocrinology

2020; 161:bqaa074.

[9]  Liu X, Xie F, Zamah AM, Cao B, Conti M. Multiple pathways mediate luteinizing

hormone regulation of cGMP signaling in the mouse ovarian follicle. Biol Reprod

2014; 91:9.

[10] Jaffe LA, Egbert JR. Regulation of mammalian oocyte meiosis by intercellular

communication within the ovarian follicle. Annu Rev Physiol 2017; 79:237–260.

[11] Richards JS, Ascoli M. Endocrine, paracrine, and autocrine signaling pathways that

regulate ovulation. Trends Endocrinol Metab 2018; 29:313–325.

[12] Sposini S, Hanyaloglu AC. Driving gonadotrophin hormone receptor signalling: the

role of membrane trafficking. Reproduction 2018; 156:R195–R208.

[13] Duffy DM, Ko C, Jo M, Brannstrom M, Curry TE. Ovulation: parallels with

inflammatory processes. Endocrine Reviews 2019; 40:369–416.

[14] Hughes CHK, Murphy BD. Nuclear receptors: Key regulators of somatic cell

functions in the ovulatory process. Molecular Aspects of Medicine 2021;

78:100937.

[15] Harasimov K, Uraji J, Mönnich EU, Holubcová Z, Elder K, Blayney M, Schuh M.

Actin-driven chromosome clustering facilitates fast and complete chromosome

capture in mammalian oocytes. Nat Cell Biol 2023; 25:439–452.

[16] Eppig JJ. The relationship between cumulus cell-oocyte coupling, oocyte meiotic

maturation, and cumulus expansion. Dev Biol 1982; 89:268–272.

[17] Clarke HJ. Transzonal projections: Essential structures mediating intercellular

communication in the mammalian ovarian follicle. Mol Reprod Dev 2022; 89:509–

525.

[18] Lipner H, Cross NL. Morphology of the membrana granulosa of the ovarian follicle.

Endocrinology 1968; 82:638–641.

[19] Downs SM, Longo FJ. An ultrastructural study of preovulatory apical development

in mouse ovarian follicles: effects of indomethacin. Anat Rec 1983; 205:159–168.

[20] Martin GG, Talbot P. Formation of the rupture site in preovulatory hamster follicles:

morphological and morphometric analysis of thinning of the granulosa and thecal

layers. Gamete Res 1987; 17:303–320.

[21] Migone FF, Cowan RG, Williams RM, Gorse KJ, Zipfel WR, Quirk SM. In vivo

imaging reveals an essential role of vasoconstriction in rupture of the ovarian

follicle at ovulation. Proc Natl Acad Sci U S A 2016; 113:2294–2299.

[22] Gothié S. Contribution à l'étude cinétique de l'ovulation chez la souris prépubère. C

R Seances Soc Biol Fil 1967; 161:554–556.

[23] Martin GG, Talbot P. The role of follicular smooth muscle cells in hamster

ovulation. J Exp Zool 1981; 216:469–482.

[24] Talbot P, Chacon RS. In vitro ovulation of hamster oocytes depends on contraction

of follicular smooth muscle cells. J Exp Zool 1982; 224:409–415.

[25] Szołtys M, Galas J, Jabłonka A, Tabarowski Z. Some morphological and hormonal

aspects of ovulation and superovulation in the rat. J Endocrinol 1994; 141:91–100.

[26] Ko C, Gieske MC, Al-Alem L, Hahn Y, Su W, Gong MC, Iglarz M, Koo Y.

Endothelin-2 in ovarian follicle rupture. Endocrinology 2006; 147:1770–1779.

[27] Brown HM, Dunning KR, Robker RL, Boerboom D, Pritchard M, Lane M, Russell

DL. ADAMTS1 cleavage of versican mediates essential structural remodeling of

the ovarian follicle and cumulus-oocyte matrix during ovulation in mice. Biol Reprod

2010; 83:549–557.

[28] Palanisamy GS, Cheon Y-P, Kim J, Kannan A, Li Q, Sato M, Mantena SR, Sitruk-

Ware RL, Bagchi MK, Bagchi IC. A novel pathway involving progesterone receptor,

endothelin-2, and endothelin receptor B controls ovulation in mice. Mol Endocrinol

2006; 20:2784–2795.

[29] Ko CJ, Cho YM, Ham E, Cacioppo JA, Park CJ. Endothelin 2: a key player in

ovulation and fertility. Reproduction 2022; 163:R71–R80.

[30] Cacioppo JA, Lin P-CP, Hannon PR, McDougle DR, Gal A, Ko C. Granulosa cell

endothelin-2 expression is fundamental for ovulatory follicle rupture. Sci Rep 2017;

7:817.

[31] Stocco C, Telleria C, Gibori G. The molecular control of corpus luteum formation,

function, and regression. Endocrine Reviews 2007; 28:117–149.

[32] Grossman H, Chuderland D, Ninio☐Many L, Hasky N, Kaplan☐Kraicer R, Shalgi R.

A novel regulatory pathway in granulosa cells, the LH/human chorionic

gonadotropin-microRNA-125a-3p-Fyn pathway, is required for ovulation. FASEB

2015; 29:3206–3216.

[33] Bianco S, Bellefleur A-M, Beaulieu É, Beauparlant CJ, Bertolin K, Droit A,

Schoonjans K, Murphy BD, Gévry N. The ovulatory signal precipitates LRH-1

transcriptional switching mediated by differential chromatin accessibility. Cell

Reports 2019; 28:2443-2454.e4.

[34] Ben-Ze'ev A, Amsterdam A. Regulation of cytoskeletal protein organization and

expression in human granulosa cells in response to gonadotropin treatment.

Endocrinology 1989; 124:1033–1041.

[35] Karlsson AB, Maizels ET, Flynn MP, Jones JC, Shelden EA, Bamburg JR,

Hunzicker-Dunn M. Luteinizing hormone receptor-stimulated progesterone

production by preovulatory granulosa cells requires protein kinase A-dependent

activation/dephosphorylation of the actin dynamizing protein cofilin. Mol Endocrinol

2010; 24:1765–1781.

[36] Owen CM, Zhou X, Bernard DJ, Jaffe LA. Kisspeptin-54 injection induces a

physiological luteinizing hormone surge and ovulation in mice. Biol Reprod 2021;

104:1181–1183.

[37] Wall EG, Desai R, Khant Aung Z, Yeo SH, Grattan DR, Handelsman DJ, Herbison

AE. Unexpected plasma gonadal steroid and prolactin levels across the mouse

estrous cycle. Endocrinology 2023; 164:bqad070.

[38] Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T,

Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, et al. Fiji - an

Open Source platform for biological image analysis. Nat Methods 2012;

9:10.1038/nmeth.2019.

[39] Norris RP, Freudzon M, Mehlmann LM, Cowan AE, Simon AM, Paul DL, Lampe

PD, Jaffe LA. Luteinizing hormone causes MAP kinase-dependent phosphorylation

and closure of connexin 43 gap junctions in mouse ovarian follicles: one of two

paths to meiotic resumption. Development 2008; 135:3229–3238.

[40] Stevenson H, Bartram S, Charalambides MM, Murthy S, Petitt T, Pradeep A,

Vineall O, Abaraonye I, Lancaster A, Koysombat K, Patel B, Abbara A. Kisspeptin-

neuron control of LH pulsatility and ovulation. Front Endocrinol (Lausanne) 2022;

13:951938.

[41] Norris RP, Freudzon L, Freudzon M, Hand AR, Mehlmann LM, Jaffe LA. A $G_s$-

linked receptor maintains meiotic arrest in mouse oocytes, but luteinizing hormone

does not cause meiotic resumption by terminating receptor-$G_s$ signaling. Dev Biol

2007; 310:240–249.

[42] Baena V, Terasaki M. Three-dimensional organization of transzonal projections

and other cytoplasmic extensions in the mouse ovarian follicle. Sci Rep 2019; 9:1–

13.

[43] LaPolt PS, Oikawa M, Jia XC, Dargan C, Hsueh AJ. Gonadotropin-induced up- and

down-regulation of rat ovarian LH receptor message levels during follicular growth,

ovulation and luteinization. Endocrinology 1990; 126:3277–3279.

[44] Xue F, Janzen DM, Knecht DA. Contribution of filopodia to cell migration: a

mechanical link between protrusion and contraction. Int J Cell Biol 2010;

2010:507821.

[45] Paluch EK, Raz E. The role and regulation of blebs in cell migration. Curr Opin Cell

Biol 2013; 25:582–590.

[46] Truszkowski L, Batur D, Long H, Tarbashevich K, Vos BE, Trappmann B, Raz E.

Primordial germ cells adjust their protrusion type while migrating in different tissue

contexts in vivo. Development 2023; 150:dev200603.

[47] Czieselsky K, Prescott M, Porteous R, Campos P, Clarkson J, Steyn FJ, Campbell

RE, Herbison AE. Pulse and surge profiles of luteinizing hormone secretion in the

mouse. Endocrinology 2016; 157:4794–4802.

[48] Lecuit T, Lenne P-F. Cell surface mechanics and the control of cell shape, tissue

patterns and morphogenesis. Nat Rev Mol Cell Biol 2007; 8:633–644.

[49] Walck-Shannon E, Hardin J. Cell intercalation from top to bottom. Nat Rev Mol Cell

Biol 2014; 15:34–48.

[50] Parsons TT, Mosallaei S, Raftery LA. Two phases for centripetal migration of

Drosophila melanogaster follicle cells: initial ingression followed by epithelial

migration. Development 2023; 150:dev200492.

[51] Kannan A, Beal JR, Neff AM, Bagchi MK, Bagchi IC. Runx1 regulates critical

factors that control uterine angiogenesis and trophoblast differentiation during

placental development. PNAS Nexus 2023; 2:pgad215.

[52] Egbert JR, Silbern I, Uliasz TF, Lowther KM, Yee S-P, Urlaub H, Jaffe LA.

Phosphatases modified by LH signaling in ovarian follicles: testing their role in

regulating the NPR2 guanylyl cyclase. bioRxiv 2023:2023.06.12.544636.

[53] Huang TY, DerMardirossian C, Bokoch GM. Cofilin phosphatases and regulation of

actin dynamics. Curr Opin Cell Biol 2006; 18:26–31.

[54] Tedeschi A, Dupraz S, Curcio M, Laskowski CJ, Schaffran B, Flynn KC, Santos TE,

Stern S, Hilton BJ, Larson MJE, Gurniak CB, Witke W, et al. ADF/Cofilin-Mediated

Actin Turnover Promotes Axon Regeneration in the Adult CNS. Neuron 2019;

103:1073-1085.e6.

[55] Sixt M, Hallmann R, Wendler O, Scharffetter-Kochanek K, Sorokin LM. Cell

adhesion and migration properties of beta 2-integrin negative polymorphonuclear

granulocytes on defined extracellular matrix molecules. Relevance for leukocyte

extravasation. J Biol Chem 2001; 276:18878–18887.

[56] Irving-Rodgers HF, Hummitzsch K, Murdiyarso LS, Bonner WM, Sado Y, Ninomiya

Y, Couchman JR, Sorokin LM, Rodgers RJ. Dynamics of extracellular matrix in

ovarian follicles and corpora lutea of mice. Cell Tissue Res 2010; 339:613–624.

[57] Segaloff DL, Wang HY, Richards JS. Hormonal regulation of luteinizing

hormone/chorionic gonadotropin receptor mRNA in rat ovarian cells during follicular

development and luteinization. Mol Endocrinol 1990; 4:1856–1865.

[58] Camp TA, Rahal JO, Mayo KE. Cellular localization and hormonal regulation of

follicle-stimulating hormone and luteinizing hormone receptor messenger RNAs in

the rat ovary. Mol Endocrinol 1991; 5:1405–1417.

[59] Zheng C, Gulappa T, Menon B, Menon KMJ. Association between LH receptor

regulation and ovarian hyperstimulation syndrome in a rodent model. Reproduction

2020; 160:239–245.

586
587

## Figure Legends

**Figure 1.** Follicle organization, time course of kisspeptin-induced ovulation, and experimental design. A) Tissue layers of a preovulatory mouse follicle. B) Time course of kisspeptin-induced ovulation in wild-type and HA-LHR mice. The time course of hCG-induced ovulation in wild-type mice is shown for comparison. Oviducts were collected at the indicated time points, and the numbers of ovulated oocytes were counted (n= 3 mice per each condition). C) Collection of ovaries after kisspeptin injection, for analysis of the localization of LH receptor-expressing cells by confocal microscopy.

**Figure 2.** LH receptor-expressing cells extend inwards within the preovulatory follicle in response to LH. A-G) Confocal images of representative 10 μm thick equatorial cryosections of follicles in ovaries before and 2-12 hours after injection of HA-LHR mice with kisspeptin. The sections were labelled for HA immunofluorescence (green), and nuclei were labelled with DAPI (blue). Each panel shows a maximum projection of a stack of 10 optical sections imaged at 1 μm intervals with a 20x/0.8 NA objective. Scale bars = 100 μm. F and G were captured using a lower zoom than the other images to account for increased follicle size.

**Figure 3.** LH induces ingression of LH receptor-expressing cells and an epithelial-mesenchymal-like transition. A) Schematic of ingression of LH receptor-expressing cells within the follicle. B) Time course of the increase in the percentage of LH receptor (LHR)-expressing cell bodies in the inner half of the mural granulosa region after kisspeptin injection. No change in the distribution of LH receptor-expressing cells was seen at 6 hours after a control injection of PBS. C) No effect of kisspeptin on the percentage of mural granulosa cells that express the LH receptor. Measurements in B and C were made from images in Figures S1-S6 and S8. For each

time point after kisspeptin, 1100-3000 cells from each of 7-14 follicles from 3-7 mice were

analyzed. Each point on the graphs represents an individual follicle. Data were analyzed via

one-way ANOVA with the Holm-Sidak correction for multiple comparisons. There were no

significant differences among time points for data in C. D) Images of representative 10 µm thick

equatorial cryosections of preovulatory follicles in ovaries from LH receptor-expressing mice

before or 6 hours after kisspeptin injection. Each panel shows a maximum projection of a stack

of 10 x 1 µm optical sections taken with a 63x/1.4 NA objective using the Airyscan detector.

Scale bars = 10 µm. E) Filopodia and blebbing observed at the 6-hour time point. Images are

maximum projections of a stack of 30 x 0.69 µm optical sections taken with 63x/1.4 NA objective

using the Airyscan detector. Scale bar = 5 µm.

**Figure 4.** LH induces ingression of LH receptor-expressing cells in follicles of naturally cycling

mice. A) Representative images of follicles from mice collected either 4 hours before lights off

(pre-LH) or 6 hours after lights off (post-LH) on the day of proestrus. The sections were labelled

for HA immunofluorescence (green), and nuclei were labelled with DAPI (blue). Each panel

shows a maximum projection of a stack of 10 optical sections imaged at 1 µm intervals with a

20x/0.8 NA objective. Scale bars = 100 µm. Note that the section through the edge of the follicle

at the far left of panel A was not equatorial, accounting for the difference in appearance. B) The

percentage of LH receptor-expressing cell bodies in the inner half of the mural granulosa region

in mice before or after the endogenous LH surge. Measurements were made from follicles in

Figure S12. Each point on the graphs represents an individual follicle; data were collected from

2 mice at each time point. Data were analyzed via an unpaired t-test (p < 0.0005).

**Figure 5.** LH induces basal lamina invaginations and constrictions in the basolateral region of

the follicle. A) Representative images of follicles before kisspeptin injection and 8 hours

afterwards, labeled for laminin gamma 1 (white). Scale bars = 100 µm. B) Tracings of the basal

lamina of follicles in equatorial sections at 0, 6, 8, or 10 hours after kisspeptin injection (from

images in Figures S1B, S4B, S5B, S6B). The tracings are aligned such that the apical region is

on the right, as indicated by the double line. Apical was defined as the region that is directly

adjacent to the surface epithelium, while basolateral was defined as any portion of the follicle

that is not in contact with the surface epithelium. Scale bar = 100 µm. C) Depth of invaginations

in follicles at 0, 6, 8, and 10 hr post kisspeptin. Each symbol represents one invagination. D)

Total number of invaginations in apical and basolateral regions of follicles at 0, 6, 8, and 10 hr

post kisspeptin. Each symbol represents one follicle. Data for C and D were generated using

datasets in Figures S1B, S4B, S5B, S6B). Each point on the graphs represents an individual

follicle; data were collected from 6 - 7 mice at each time point.  Data were analyzed via two-way

ANOVA with the Holm-Sidak correction for multiple comparisons.

**Figure 6.** LH induces thinning of the mural granulosa layer at the apex but not the base. A)

Representative images of follicles before kisspeptin injection and 10 hours after, with nuclei

labelled with DAPI (white). Apical and basal regions (blue and yellow boxes) are enlarged at the

right. Pink lines represent measurements of the mural granulosa width, taken 50 µm apart and

then averaged to generate the points in (B). Scale bars = 100 µm for full follicle images on the

left, 50 µm for insets on the right. B) Width of mural granulosa layer in apical and basal regions

of follicles before and 2-10 hours after kisspeptin injection. Each point on the graph represents

the average measurements from one follicle from Figures S1-S6. Data were analyzed via one-

way ANOVA with the Holm-Sidak correction for multiple comparisons.

**A**

Outer mural granulosa
Inner mural granulosa
Basal lamina
Theca
Vasculature
Interstitium

Antrum
Cumulus
Oocyte

**B**

# of oocytes retrieved per mouse

- HA-LHR + kisspeptin
- WT + kisspeptin
- WT + hCG

Hours after kisspeptin or hCG injection

**C**

5 I.U. eCG
1 nmol kisspeptin

HA-LHR mouse

Peak of LH release (1.5 hr post kisspeptin)

Ovulation starts

Collect ovaries at designated times after kisspeptin injection

44 hr

0 hr    2 hr    4 hr    6 hr    8 hr    10 hr    12 hr

Fix
Freeze
Cryosection

Immunostain for HA-LHR
Image with confocal microscopy

Analyze location of HA-LHR expressing cells in follicles

**A**

LH

**B**

% of total LHR-expressing cell bodies that are in the inner mural granulosa

a   b   b   b   b   c          a

0   2   4   6   8   10         6
Kisspeptin              PBS

Hours after injection

**C**

% of mural granulosa cells that express LHR

0   2   4   6   8   10         6
Kisspeptin              PBS

Hours after injection

**D**

HA-LHR/DAPI

0 hr post kisspeptin          6 hr post kisspeptin

**E**

HA-LHR

6 hr post kisspeptin

**A**

| 4 hr before lights off | 6 hr after lights off |

HA-LHR/DAPI

**B**

% of total LHR-expressing cell bodies that are in the inner mural granulosa

**A**

0 hr post kisspeptin

10 hr post kisspeptin

| | 0 hr post kisspeptin | 10 hr post kisspeptin |
| Apical | | |
| Basal | | |

**B**

Mural granulosa width (µm)

Apical — a, ab, abc, bc, c, c

Basal

Hours after kisspeptin injection