TITLE

Immediate effects of light on circadian eclosion and locomotor activity depend on distinct sensory input pathways

AUTHORS

Daniel Bidell[1], Natalie-Danielle Feige[1], Tilman Triphan[2], Dennis Pauls[1], Charlotte Helfrich-Förster[3] and Mareike Selcho[1,#]

AFFILIATIONS

[1] Department of Animal Physiology, Institute of Biology, Leipzig University, Talstraße 33, 04103 Leipzig, Germany

[2] Department of Genetics, Institute of Biology, Leipzig University, Talstraße 33, 04103 Leipzig, Germany

[3] Neurobiology and Genetics, Biocenter, University of Würzburg, 97074 Würzburg, Germany

[#] Corresponding author: MS (mareike.selcho@uni-leipzig.de)

ABSTRACT

Animals need to be able to sharpen circadian behavioural output in the adaptation to the variable environment. Light is the main entraining signal of the circadian clock, but can also directly increase alertness, locomotor activity, body temperature and heart rate in diurnal animals including humans. Thus, immediate effects of light can enhance or even overwrite circadian output and thereby mask circadian behaviour.

In *Drosophila melanogaster*, immediate light effects are most evident as a lights-on response in two well described behavioural rhythms of the fly – the emergence rhythm of the adult insect from the pupa, called eclosion, and the diurnal rhythm of locomotor activity. Here, we show that the immediate effect of light on rhythmic eclosion depends on the R8 photoreceptor cells of the compound eyes, while the light response of locomotor activity is triggered by different light detecting cells and organs, that seem to compensate for the loss of each other.

INTRODUCTION

An appropriate daily timing of behaviour is of critical importance for animal fitness (Daan and Tinbergen 1979; DeCoursey et al. 2000). Light shapes the daily timing in two ways: (1) the cyclic change in light intensity acts as an entraining signal, synchronizing the endogenous (circadian) timing system to appropriately adjust an organism to the 24h environmental period; (2) light directly modulates behaviour i.e. increases alertness, locomotor activity, body temperature and heart rate in diurnal animals including humans (Erkert and Gröber 1986; Aschoff and von Goetz 1989; Dijk et al. 1991; Badia et al. 1991; Cajochen et al. 2005) and supresses activity and promotes sleep in nocturnal animals (Mrosovsky 1994; Altimus et al. 2008). These immediate light effects are independent of the circadian clock and since they often obscure ("mask") circadian behaviours, they are also referred to as "masking" (Mrosovsky 1999). The immediate light effects are essential for appropriate responses of the animal to changes in the environment and for sharpening the behavioural output. Interestingly, light responses are not only shown to be independent of the circadian clock, they can even provoke quasi-wildtype activity rhythms in clock-less and thus arrhythmic fruit flies and mice under natural-like conditions (Daan et al. 2011; Vanin et al. 2012; Schlichting et al. 2015). This clearly underlines their importance.

In *Drosophila melanogaster*, the immediate light effects are most evident as a lights-on response in the two well described behavioural rhythms of the fly – the emergence rhythm of the adult insect from the pupa, called eclosion, and the adult activity rhythm (Engelmann and Honegger 1966; Rieger et al. 2003; McNabb and Truman 2008). Eclosion is gated by the circadian clock to the early day, most probably to prevent desiccation and enhance survival rate. A light stimulus induces a rapid increase in eclosion rate (lights-on effect) that is eliminated in flies without eyes and potentially in mutants lacking ocelli (Engelmann and Honegger 1966; McNabb and Truman 2008). Even though the lights-on effect is gone, eclosion remains synchronized to the light-dark cycle in eyeless flies by the circadian blue-light photopigment Cryptochrome (CRY; Engelmann and Honegger 1966; reviewed by Helfrich-Förster and Engelmann 2002). Similarly, the locomotor activity rhythm of adult eyeless flies remains synchronized to the light-dark cycle due to entrainment by CRY (reviewed by Helfrich-Förster 2002), but the immediate increase of activity after lights-on, also known as "startle response" disappears after elimination of the eyes (Helfrich and Engelmann 1983; Rieger et al., 2003). Subsequent studies showed that several

rhodopsins contribute to this immediate light effect in adult flies (Schlichting et al. 2014,
2015).

Flies receive light information via two external organs, the compound eyes and ocelli as
well as the internal extraretinal Hofbauer-Buchner (H-B) eyelets. In addition, a small
subset of central brain neurons express the blue-light sensitive photopigment CRY (Emery
et al. 1998; Stanewsky et al. 1998). The fruit fly's compound eye consists of around 800
ommatidia, each of them equipped with six outer (R1-6) and two inner (R7, R8)
photoreceptor cells. While the outer photoreceptor cells express the photoreceptor protein
Rhodopsin 1 (Rh1), the inner ones express specific combinations of Rh3, Rh4, Rh5 and
Rh6 (reviewed by Rister and Desplan 2011). Photoreceptor proteins are specifically
expressed in two main ommatidia types: pale ommatidia express Rh3 in R7 and Rh5 in R8
photoreceptors and yellow ommatidia Rh4 in R7 and Rh6 in R8 cells. A third type, the
DRA ommatidia, positioned in the dorsal rim area (DRA) expresses Rh3 in R7 and R8
cells. The photoreceptors R1-6 were shown to be involved in dim light vision and the
perception of motion (Heisenberg and Buchner 1977), while R7/R8 are involved in colour
vision (Heisenberg and Buchner 1977; Schnaitmann et al. 2013; Melnattur et al. 2014) and
DRA photoreceptors are involved in polarization vision (Wernet et al. 2012). Besides the
large compound eyes, fruit flies contain three simple dorsal eyes called ocelli. Ocelli
consist of only one photoreceptor type, containing Rh2 (Feiler et al. 1988a; Pollock and
Benzer 1988). The ocelli do not form an image or perceive objects in the environment,
instead they are sensitive to changes in light intensities and seem to response to polarized
light. As Rh2 is highly sensitive to UV light, the ocelli provide information to distinguishing
between sky and the ground (reviewed by Sabat et al. 2016) . This enables the fly to
maintain its orientation in space. The HB eyelets evolve from the larval Bolwig's organ and
express Rh6 (Hofbauer and Buchner 1989; Yasuyama and Meinertzhagen 1999; Helfrich-
Förster et al. 2002; Sprecher and Desplan 2008). These extraretinal light detecting cells
are involved in entrainment at high light intensities (Schlichting et al. 2019). In addition to
the light detecting organs, Drosophila contains CRY expressing neurons in the brain and
compound eyes, sensitive to blue light (Yoshii et al. 2008; Hoang et al. 2008; Benito et al.
2008; Fogle et al. 2011). Beside the compound eyes, CRY is the main photoreceptor
involved in light entrainment of the circadian clock (Emery et al. 2000; Rieger et al. 2003).
In our study, we aim to identify the neuronal correlates of the immediate light effects that
trigger specific behavioural responses counteracting or reinforcing circadian rhythms. For

this, we use eclosion and locomotor activity as behavioural readouts to examine
immediate light effects on circadian behaviours. We provide evidence that the light effect
on eclosion is mediated by Rh5-positive R8 photoreceptor neurons of the compound eyes.
In contrast, the ocelli, the H-B eyelets, as well as light sensitive CRY-positive cells are not
required to elicit this lights-on response. Interestingly, the light response on locomotor
activity in the night remains in flies without eyes, ocelli or CRY-positive cells. We
hypothesize redundant signalling pathways and propose that light perceiving cells and
organs compensate the loss of each other, enabling flies to react to changes in their
environment. Thus, immediate effects of light on circadian behaviours seem to depend on
different underlying neuronal networks.
111
RESULTS
Light elicits an immediate increase in eclosion at lights-on in fly populations monitored
under 14:10 light:dark cycles (Fig.1A). This immediate effect is also visible in flies kept
under constant darkness when a light pulse is given at times around expected lights-on
(Fig.1B-D). In detail, in line with previous studies (McNabb and Truman 2008), we could
show that a twenty minutes light pulse one hour before or after expected lights-on elicits
an immediate eclosion response in control flies (Fig.1B-D). The light response thus
enables the flies to eclose promptly in response to their environment, reinforcing the
behaviour controlled by the clock. Next, we tested whether the pigmentation of the eyes
plays a role in the detection of light necessary to elicit immediate eclosion. For this, we
monitored eclosion in red-eyed CantonS (CS) and white-eyed (w^{1118}) flies. Both groups
showed a clear eclosion peak in response to light, even though the distribution within the
twenty minutes is different (Fig.1C,D). Nevertheless, eye-colour does not influence the
immediate response to light. Further, we tested different mutants to discover the light
perceiving cells and organs perceiving light triggering immediate eclosion. The lights-on
response is gone in flies lacking the compound eyes (cli^{eya}, Fig.2B; Engelmann and
Honegger 1966; McNabb and Truman 2008) and in *norpA* (*no receptor potential*) mutants
with disturbed phospholipase C function (*norpA^{p41}*, Fig.2C). In contrast, flies with disabled
light perception in photoreceptors of the ocelli (*rh2^{1}*, Fig.2D), *cry*-positive cells (*cry^{01}*,
Fig.2E) or the Rh6-positive eyelet (*rh6^{1}*, Fig.S1), respectively, showed an immediate
increase in eclosion in response to light indicating the exclusive requirement of the
compound eyes in the immediate effect of light on eclosion.
134

eclosion in light phase; dashed bars: % eclosion in controls. n_{exp}, n_{ctrl} = 524, 544 (A); 502, 508 (B); 520, 539
(C); 604, 584 (D); 510, 556 (E).

Our data so far suggest that the compound eyes are required for the immediate light effect

on eclosion. Light information is received by photoreceptor cells of the compound eyes

and converted into neuronal activity. The outer photoreceptor cells R1-R6 contain Rh1, the

inner R7 cells express Rh3 (R7pale) or Rh4 (R7yellow) and proximal R8 cells express

either Rh5 (R8pale) or Rh6 (R8yellow), while DRA ommatidia express Rh3 in R7 and R8.

Thus, the different photoreceptor cells respond to light of a specific spectrum as the five

different rhodopsin's of the compound eyes have different absorption spectra (Feiler et al.

1988, 1992; Salcedo et al. 1999). Therefore, illumination by monochromatic light of

different wavelength will allow to address specific sets of Rh cells only and thereby give

insights into the sufficiency of the different photoreceptor cells in this context. Flies showed

an immediate behavioural response to blue (455– 475 nm), green (510– 545 nm) and red

(625– 642 nm) light (Fig.3A-C). In contrast to mammals, where immediate light-effects

depend exclusively on blue light-sensitive melanopsin-expressing retinal ganglion cells

(Redlin et al. 1999; Redlin and Mrosovsky 1999; Mrosovsky and Hattar 2003; Panda et al.

2003; Gall et al. 2013), flies are able to additionally respond to longer wavelengths. As red

light of 633nm could only be absorbed by Rh1 or Rh6, we monitored the immediate light

responses in flies without Rh1 (*ninaE^{17}*), flies that lack Rh6 (*rh6^1*) and double mutants

without both of these rhodopsins (*ninaE^{17}; rh6^1*). As expected, the lights-on response is

missing in the *rh1,rh6* double mutant (Fig.3F). Lacking just one of the red light detecting

rhodopsins leads to a reduced lights-on response (Fig.3D,E). Interestingly, however, flies

lacking Rh1 and Rh6 show an immediate behavioural response to a white light pulse

(Fig.3G), indicating that Rh1 and Rh6 are both involved in the immediate effect of red light,

but are dispensable for the immediate effect of white light on eclosion. In addition, as Rh1

is the only rhodopsin expressed in the outer photoreceptor cells, R1-R6 might not be

required for the lights-on response to white light. To further disentangle the role of the R7,

R8 cells, we screened rhodopsin and photoreceptor mutants for their behavioural

response to white light (Fig.3G-J). The quadruple mutant (*rh5^2;rh3^1,rh4^1,rh6^1*) lacks all

rhodopsins of the inner photoreceptors, so that light perception is only possible via R1-R6.

The lack of the lights-on response (Fig.3H) indicates that (1) R7 and/ or R8 transmit light

information for the lights-on response and (2) functional outer photoreceptors are not

sufficient to trigger immediate eclosion. To address if R7 is involved in the immediate

effect of light, we monitored eclosion in *sevenless* mutants (*sev^{LY3}*, Fig.3I). Flies without

R7 photoreceptor cells show an immediate response, so that R8 cells are considered for
the transmission of light information. R8 cells express Rh5 or Rh6. Since the absence of
Rh6 has no effect on the immediate response to light (Fig.3G and Fig.S1), we tuned our
attention to the role of *rh5* mutants on eclosion (*rh5^2*, Fig.3J). As expected, flies lacking
Rh5 show no lights-on response. Thus, pale R8 ommatidia expressing functional Rh5
turned out to be essential for the immediate response to light. In line with this hypothesis,
we optogenetically activated *rh5*-expressing R8 neurons (Fig.3K,L). Photostimulation of
rh5-positive neurons via Channelrhodopsin-2^{XXL} (*rh5^G>chop2^{XXL}*, Fig.3K; Nagel et al.
2003; Dawydow et al. 2014) using a blue light pulse triggered immediate eclosion, while
the response is absent in control flies (Fig.3L). This demonstrates that activation of *rh5*-
positive R8 cells triggers immediate eclosion. Our data demonstrate that Rh5-expressing
R8 cells are necessary and sufficient for the immediate light effect on eclosion.
198

Fig.3: The immediate light effect on eclosion behaviour depends on R8 cells. (A-C) Twenty minutes blue (A, 455– 475 nm), green (B, 510– 545 nm) and red (C, 625– 642 nm) light pulses elicit immediate eclosion. **(D-F)** The eclosion response to red light is visible in *rh1* (D, *ninaE[17]*) and *rh6* mutants (E, *rh6[1]*), but gone in *rh1,rh6* double mutants (F, *ninaE[1]; rh6[1]*). **(G)** In contrast, *rh1,rh6* double mutants (*ninaE[17]; rh6[1]*) respond with an increase of eclosion to white light. **(H)** The quadruple mutant (*rh5[2]; rh3[1], rh4[1], rh6[1]*), lacking all rhodopsins of the inner photoreceptors, shows no reaction to light. **(I)** The lights-on response in flies without R7 cells (*sev[LY3]*). **(J)** Flies lacking Rh5 (*rh5[2]*) do not respond to light with increased eclosion. **(K)** Optogenetic activation of *rh5*-positive neurons with a 2 min blue light pulse (blue line) one hour after expected lights-on elicits eclosion. **(L)** Control flies without Gal4 expression (*w[1118]; chop2[XXL]*) do not respond to the 2 min light pulse. n_{exp}, n_{ctrl} = 534, 1543 (A); 521, 1543 (B); 547, 1543 (C); 511, 512 (D); 579, 731 (E); 604, 567 (F); 561, 567 (G); 595, 518 (H), 525, 531 (I); 585, 506 (J); n= 516 (K), 515 (L).

Eclosion is not the only circadian behaviour that can be modulated by light. To explore whether immediate light effects are generally mediated by R8 cells of the compound eyes

as demonstrated above, we turned our attention to locomotor activity. Locomotor activity in
Drosophila is regulated by the circadian clock so that flies display low levels of nocturnal
activity and higher levels of diurnal activity. Importantly, flies display a lights-on response
in the transition from night to day (startle response, Fig.4A). To investigate immediate light
effects, we applied a twenty minutes light pulse two hours before the anticipated lights-on
response in the transition from night to day. The light pulse elicited an immediate increase
in locomotor activity, similar to the light effect on eclosion (Fig.4A, Fig.1). Interestingly, in
contrast to eclosion, the immediate light response in locomotor activity is also present in
eyeless flies that lack additionally functional CRY (*cli^{eya};cry^b*; Fig.4B). Thus, neither
photoreceptor cells of the compound eyes nor CRY are required for the increase in activity
in response to light (Fig.4B and Fig.S2 for *rh* and *norpA* mutants). In addition, we analysed
locomotor activity in two different *cry* mutants to clarify the role of photosensitive neurons
outside the visual system (*cry^b* and *cry^{01}*). Flies without functional CRY responded to light
with an immediate increased locomotor activity (Fig.4C and Fig.S2A). Further, immediate
light effects can be seen in flies with disabled light perception in photoreceptors of the
ocelli (*rh2^1*, Fig.4D) and flies without Rh5 and Rh6 and therefore without functional eyelets
(*rh5^2, rh6^1*; Fig.4E). Thus, all the different cells and organs that perceive light appear to
mediate its immediate effects and compensate for the failure of individual photoreceptors,
so that flies can respond immediately to a light stimulus. In contrast, flies without a
functional histidine decarboxylase (*hdc^{JK910}*), the enzyme necessary for histamine
synthesis, show no increase in locomotor activity in response to light (Fig.4F). Histamine is
the main transmitter of the photoreceptor cells of the eyes, ocelli and eyelets (Pollack and
Hofbauer 1991), leaving the hypothesis that *cry*-positive cells alone may not be sufficient
to elicit the lights-on response.
In summary, our data provides evidence that immediate light effects are present on
eclosion and locomotor activity. However, the underlying signalling pathways appear to be
different.

Fig. 4: The immediate light effect on locomotor activity is visible in flies without functional eyes, photosensation in *cry*-positive cells or ocelli. (A-F) Activity pattern in ten minutes intervals at the time around expected lights-on (0). First and second column show bar plots of mean $\pm$ SEM activity at the day the light pulse was applied (-2, second column) and the activity of the same flies a day before (first column). The third column visualizes the comparison between the mean activity of flies in dark and in light at night at the corresponding timepoints. **(A)** Activity data of control (w^{1118}) and **(B)** eyeless (cli^{eya}; cry^b) flies, **(C)** flies lacking Cryptochrome (cry^{01}), **(D)** flies without functional ocelli photoreceptors ($rh2^1$), **(E)** flies without Rh5 and Rh6 ($rh5^2$; $rh6^1$) **(F)** without Rh5 and Rh6 ($rh5^2$; $rh6^1$) that do respond to light. **(F)** The light response is absent in flies that lack the histidine decarboxylase (hdc^{JK910}) and therefore histamine, the transmitter of photoreceptor cells. n = 27- 32; asterisks denote level of significance: **$p \leq 0.01$, ***$p \leq 0.001$, ****$p \leq 0.0001$.

DISCUSSION

Even though light triggers immediate eclosion and activity, modifying circadian behaviour, the necessary underlying networks seem to differ already at sensory input level. The lights-on response on eclosion depends on Rh5-positive R8 cells of the compound eyes, while the light response on locomotor activity works without eyes, ocelli, eyelets or photosensitive *cry*-positive brain cells and is only absent when histamine is missing. For activity increase at night the different light detecting organs and cells can compensate the loss of each other, probably to ensure the ability to react to the unexpected external stimulus.

Interestingly, former studies showed, that the immediate light effects on locomotor activity at lights-on (also called startle responses) depend on the compound eyes and were completely absent in eyeless flies (Rieger et al., 2003; Helfrich-Förster, 2014). Here, we observed the same: eyeless flies (cli^{eya};cry^b mutants, which lack functional CRY in addition to the eyes) as well as flies without histamine (hdc^{JK910} mutants) lack the startle response at lights-on (time point 0 first column, Fig.4B,F). Nevertheless, a light pulse during the night two hours before lights-on, provokes an immediate increase in activity in cli^{eya};cry^b mutants (Fig.4B). Only the disruption of neuronal communication of photoreceptor cells of the eyes, ocelli and eyelets by the loss of histamine (hdc^{JK910} mutants) prevents an immediate response to light (Fig. 4F). The immediate increase in activity in response to light at night is thus mediated by functionally redundant photoreceptor cells, whereas the startle response depends purely on the receptors in the eyes.

The functional redundancy of photoreceptors was also described in nocturnal mice. Here, bright light at night leads to an inhibition of locomotor activity, whereas dim light increases activity (Mrosovsky et al., 1999; Mrosovsky, 2003; Redlin et al., 2003). Mice with

degenerated retina (*rd/rd* mice), devoid of rods and cones, lack the increase of activity by

dim light, but still show activity inhibition by bright light. In addition, mice lacking the

photopigment melanopsin (*Opn4$^{-/-}$*) in the retinal ganglion cells also show inhibition of

activity by bright light, while double mutants (*Opn4$^{-/-}$; rd/rd*) loose the ability to respond to

light (Panda et al. 2003; Mrosovsky and Hattar 2003; Hattar et al. 2003). Thus, also in

mice, several photopigments contribute to the immediate light responses and these are

able to compensate to some degree the loss of each other to ensure immediate responses

to unexpected external stimuli.

Interestingly, recent data suggests that CRY suppresses activity during the night in flies,

as the absence of CRY enhances activity during moonlit nights (Zurl et al. 2022). Flies

were as active during moonlit nights as they were during the day, suggesting that CRY is

important for distinguishing nocturnal low light of moonlight intensity from day light, and

most interestingly this was also true for marine bristle worms (Zurl et al. 2022). If CRY is

present, it seems to suppress activity during the night in the diurnal *D. melanogaster* and

to suppress swarming activity during the day in the nocturnal marine bristle worms. This

finding for *Drosophila* is corroborated by the present study: *cry^b* mutants show strong

immediate light effects upon a light pulse 2h before lights-on, but flies that lack the function

of all photoreceptors except of CRY (*hdc^{JK910}* mutants) don't show this response. In

summary, we conclude that the immediate light effects of adult flies in response to

nocturnal light are mediated by all rhodopsins (those in the compound eyes, the H-B

eyelets and putatively also the ocelli), while they are inhibited by CRY. This is different for

the lights-on effect at the beginning of the day. Here we could not see any inhibiting effect

of CRY.

For the immediate effect of light on eclosion, only Rh5 in R8 cells of the compound eyes is

needed. Possibly, not all photoreceptors are yet mature at the time of eclosion. This is

certainly true for the H-B eyelets that are fully functional only several days after eclosion

(Helfrich-Förster et al. 2002). For the other photoreceptors such a delayed maturation is

not known, but it is possible that not all the connections to the central brain, in particular

those that connect to activity-promoting centres, are fully established.

METHODS

Fly stocks

Flies were raised on standard cornmeal and molasses medium at 25°C and 65% relative

humidity at a 14:10 light-dark (L:D) cycle unless otherwise stated. For optogenetic

experiments, vials have been covered with a light filter foil (Nr. 026; LEE Filters Worldwide,

UK). The following flies were used in this study: cli^{eya} (Bonini et al. 1993), cli^{eya}; cry^b, cry^{01}

(Dolezelova et al. 2007), cry^b (Stanewsky et al. 1998, kind gift of R. Stanewsky), hdc^{JK910}

(Burg et al. 1993),

$nina^{E17}$ (O´Tousa et al. 1985; Kumar and Ready 1995), $nina^{E17}$; $rh6^1$ (Hanai et al. 2008),

$norpA^{p41}$ (Zhu et al. 1993, kind gift of R. Stanewsky), $rh2^1$ (Sokabe et al. 2016; kind gift of

C. Montell), $rh5^2$ (Yamaguchi et al. 2008; kind gift of R. Stanewsky), $rh5^2$; $rh3^1$, $rh4^1$, $rh6^1$

(Schlichting et al. 2014), $rh5^2$; $rh6^1$ (Yamaguchi et al. 2008), $rh5^G$ (Sokabe et al. 2016; kind

gift of C. Montell; BDSC #66671), $rh6^1$ (Cook et al. 2003; Hanai et al. 2008; kind gift of C.

Montell), sev^{LY3} (Harris et al. 1976; Chou et al. 1999), $UAS\text{-}chop2^{XXL}$ (Dawydow et al.

2014), $10xUAS\text{-}IVS\text{-}myr::GFP$ (Pfeiffer et al. 2010; BDSC #32197)

Eclosion behaviour

To investigate the immediate light effect on eclosion behaviour, an eclosion monitor based

on the WEclMon-System (Ruf et al. 2017; Selcho et al. 2017) was used. 7 to 9 days old

pupae were placed onto a transparent acrylic plate. This plate was placed on an area light

with an RGB or White-LED illumination (LED-color: λ(blue) = 455 – 475 nm, λ(green) =

510 – 545 nm, λ(red) = 625 – 642 nm; white = 500K (± 10%); Hansen GmbH, Germany)

and an additional IR LED strip was installed at the bottom of the lighting unit (SOLAROX®

LED, λ(infrared) = 850nm, IR1-60-850, Winger Electronics GmbH & Co. KG, Germany)

and covered with an aluminum box. Flies were kept in darkness and received a twenty

minutes light pulse one hour after expected lights-on on the second day, or remained in

darkness (darkness control). The eclosion monitor is equipped with a camera (DMK

27AUC02 with a TPL 0420 6MP objective; DMK 37BUX287 with a TPL 0620 6MP

objective, The Imaging Source Europe GmbH) that took images every two minutes (IC

Capture 64bit, V2.5.1547.4007, The Imaging Source Europe GmbH, Germany). We

developed a Python-script for Fiji (V2.9.0, (Schindelin et al. 2012)) that was able to

independently scan a selected image sequence for eclosion events (detailed description

below).

Eclosion data analysis

The detection of hatching events is based on the difference in brightness between a pupa
and an empty pupal case. The latter is almost transparent while a late pupa is
considerably darker (Fig.S3A, empty pupal case marked with an asterisk). In the first step
of the analysis all pupae are detected. The pupae are darker than the background and can
be easily extracted in a single image. After setting a threshold (Fig.S3B), objects are
detected, and their outlines stored. Objects that are outside of the parameters defined for
a pupa (area too big or too small) are excluded. Note the empty pupal case that is
excluded from further analysis. As the pupae are stationary, we can follow them through
time and calculate the median grey value of the area contained in their respective outlines
(Fig.S3C) for each frame. Alternatively, the average or mode (i.e. most frequent) grey
values can be used. A large jump in brightness indicates an eclosion event (Fig.S3D,E).
The median grey values are more than doubled from around 40 to almost 100. For each
eclosion event a few frames before and after the event are extracted to help with manual
confirmation of real hatching events or to exclude inaccurate detections (Fig.S3E). The
time for each eclosion event is stored in a csv file for further analysis. Additional checks
are implemented to reduce the number of incorrect detections and handle changes in
lighting. See the commented source code for details. The complete workflow is
implemented as a Python script for Fiji. Specific parameters that depend on camera
resolution, optics and lighting can be set manually (e.g. area of the pupae in pixel,
difference in brightness for full and empty pupae, etc.). All code and further information
can be found at https://github.com/triphant/EclosionDetector.
For the evaluation of eclosion events, a time window of five hours was chosen, from two
hours before expected lights-on to three hours after expected lights-on. To calculate the
eclosion percentage (% eclosion), the number of flies eclosed in a 10 min time interval was
normalized to the number of flies eclosed during the 5 h time window. The bar plots in
Fig.1D'', Fig.2 and Fig.3 visualize the eclosion of flies perceiving a 20 min light pulse and
of appropriate control flies kept in darkness (Fig.1D-D''). n refers to the number of
individuals tested.
374
Optogenetics
For optogenetic activation, Channelrhodopsin-2^{XXL} (UAS-*chop2*^{XXL}; Nagel et al. 2003;
Dawydow et al. 2014) has been used to depolarize *rh5*-positive neurons by blue light.
Pupae have been collected under red light during their subjective day to monitor eclosion.
One hour after expected lights-on pupae received a 2 min blue light stimulus (455 – 475

nm, I = 3,41 μW/mm^2) and eclosion was monitored and calculated as described above. n

refers to the number of individuals tested.

Locomotor Activity

Flies were entrained to a 12:12 LD rhythm. To investigate the lights-on effect on locomotor

activity, we placed two to four days old flies in small glass tubes with 2% agarose and 4%

sugar on one side into the *Drosophila* Activity Monitoring System (DAM, V2, TriKinetics

Inc., USA) and recorded locomotor activity for the next four night-day cycles. In the first

three night-day cycles the light regime did not differ from the entrained rhythm (12:12 LD).

On the fourth night, a 20 min light-pulse (100 lux) was given two hours before normal

lights-on. The data received from the DAM system was analyzed by taking the number of

counts of a 10 min interval of each tube and calculating the average activity. This was

done for both the 6 h time window (-3 h to +3 h from lights-on) in which the light pulse was

given and the same time window on the day before, which was used as control.

Immunohistochemistry

Whole heads have been fixed for 2 hours in 4% PFA. After washing in PBS (Phosphate

Buffered Saline) specimens have been embedded in 7% agarose and cut into 70-100 μm

sections using a vibratome (Leica VT1000S; Pauls et al. 2018). Sections have been

washed three times for 10 min in 0.3% PBT (PBS with 0.3% TritonX-100), blocked for 1,5

h in 5% normal goat serum in PBT. Afterwards the first antibody solution was incubated

overnight at 4°C. Specimens were washed six times and the second antibody solution was

added and incubated overnight at 4°C. After another washing step, specimens were

mounted in Vectashield (Vector Laboratories) and stored at 4°C until scanning. Probes

have been imaged using a LSM 800 (Zeiss). Afterwards the images have been edited for

brightness and contrast using FIJI and Adobe Photoshop (V23.5.3).

The following antibodies were used: rabbit-α-GFP (1:1000; life technologies, A11122),

3C11 α-Synapsin (1:50, Klagges et al. 1996), goat-α-rabbit AlexaFluor 488 (1:250, life

technologies, A11034) and goat-α-mouse STAR RED (1:250, Abberior, STRED-1001).

Statistical Analysis

Data was analysed with Excel (Microsoft) and Prism 8.2 (GraphPad). The Shapiro-Wilk

test was used to analyse normal distribution. Group means were compared by an ordinary

one-way ANOVA with Tukey correction and for not normally distributed data by a Kruskal-

Wallis test. Prism was used to plot data. Significance levels refer to the raw p values

obtained in the statistical tests.

REFERENCES

Altimus, C. M., A. D. Güler, K. L. Villa, D. S. McNeill, T. A. Legates *et al.*, 2008 Rods-cones and melanopsin
detect light and dark to modulate sleep independent of image formation. Proc. Natl. Acad. Sci.
U.S.A. 105: 19998–20003.
Aschoff, J., and C. von Goetz, 1989 Masking of circadian activity rhythms in canaries by light and dark. J.
Biol. Rhythms 4: 29–38.
Badia, P., B. Myers, M. Boecker, J. Culpepper, and J. R. Harsh, 1991 Bright light effects on body
temperature, alertness, EEG and behavior. Physiol. Behav. 50: 583–588.
Benito, J., J. H. Houl, G. W. Roman, and P. E. Hardin, 2008 The Blue-Light Photoreceptor
CRYPTOCHROME Is Expressed in a Subset of Circadian Oscillator Neurons in the *Drosophila* CNS.
J Biol Rhythms 23: 296–307.
Bonini, N. M., W. M. Leiserson, and S. Benzer, 1993 The eyes absent gene: Genetic control of cell survival
and differentiation in the developing Drosophila eye. Cell 72: 379–395.
Burg, M. G., P. V. Sarthy, G. Koliantz, and W. L. Pak, 1993 Genetic and molecular identification of a
Drosophila histidine decarboxylase gene required in photoreceptor transmitter synthesis. The EMBO
Journal 12: 911–919.
Cajochen, C., M. Münch, S. Kobialka, K. Kräuchi, R. Steiner *et al.*, 2005 High sensitivity of human melatonin,
alertness, thermoregulation, and heart rate to short wavelength light. J. Clin. Endocrinol. Metab. 90:
1311–1316.
Daan, S., K. Spoelstra, U. Albrecht, I. Schmutz, M. Daan *et al.*, 2011 Lab mice in the field: unorthodox daily
activity and effects of a dysfunctional circadian clock allele. J. Biol. Rhythms 26: 118–129.
Daan, S., and J. Tinbergen, 1979 Young Guillemots (Uria lomvia) Leaving Their Arctic Breeding Cliffs: A
Daily Rhythm in Numbers and Risk. Ardea 67: 96–100.
Dawydow, A., R. Gueta, D. Ljaschenko, S. Ullrich, M. Hermann *et al.*, 2014 Channelrhodopsin-2–XXL, a
powerful optogenetic tool for low-light applications. Proc. Natl. Acad. Sci. U.S.A. 111: 13972–13977.
DeCoursey, P. J., J. K. Walker, and S. A. Smith, 2000 A circadian pacemaker in free-living chipmunks:
essential for survival? J. Comp. Physiol. A 186: 169–180.
Dijk, D. J., C. Cajochen, and A. A. Borbély, 1991 Effect of a single 3-hour exposure to bright light on core
body temperature and sleep in humans. Neurosci. Lett. 121: 59–62.
Dolezelova, E., D. Dolezel, and J. C. Hall, 2007 Rhythm defects caused by newly engineered null mutations
in Drosophila's cryptochrome gene. Genetics 177: 329–345.
Emery, P., W. V. So, M. Kaneko, J. C. Hall, and M. Rosbash, 1998 CRY, a Drosophila Clock and Light-
Regulated Cryptochrome, Is a Major Contributor to Circadian Rhythm Resetting and
Photosensitivity. Cell 95: 669–679.
Emery, P., R. Stanewsky, C. Helfrich-Förster, M. Emery-Le, J. C. Hall *et al.*, 2000 Drosophila CRY Is a Deep
Brain Circadian Photoreceptor. Neuron 26: 493–504.
Engelmann, W., and H. Honegger, 1966 Tagesperiodische Schlupfrhythmik einer augenlosen Drosophila
melanogaster- Mutante. Naturwissenschaften 588.
Erkert, H. G., and J. Gröber, 1986 Direct modulation of activity and body temperature of owl monkeys (Aotus
lemurinus griseimembra) by low light intensities. Folia Primatol. 47: 171–188.
Feiler, R., R. Bjornson, K. Kirschfeld, D. Mismer, G. M. Rubin *et al.*, 1992 Ectopic expression of ultraviolet-
rhodopsins in the blue photoreceptor cells of Drosophila: visual physiology and photochemistry of
transgenic animals. J. Neurosci. 12: 3862–3868.
Feiler, R., W. A. Harris, K. Kirschfeld, C. Wehrhahn, and C. S. Zuker, 1988 Targeted misexpression of a
Drosophila opsin gene leads to altered visual function. Nature 333: 737–741.
Fogle, K. J., K. G. Parson, N. A. Dahm, and T. C. Holmes, 2011 CRYPTOCHROME Is a Blue-Light Sensor
That Regulates Neuronal Firing Rate. Science 331: 1409–1413.
Gall, A. J., L. Smale, L. Yan, and A. A. Nunez, 2013 Lesions of the Intergeniculate Leaflet Lead to a

Reorganization in Circadian Regulation and a Reversal in Masking Responses to Photic Stimuli in the Nile Grass Rat. PLoS ONE 8: e67387.

Heisenberg, M., and E. Buchner, 1977 The rôle of retinula cell types in visual behavior ofDrosophila melanogaster. J. Comp. Physiol. 117: 127–162.

Helfrich, C., and W. Engelmann, 1983a Circadian rhythm of the locomotor activity in Drosophila melanogaster and its mutants 'sine oculis' and 'small optic lobes.' Physiological Entomology 8: 257–272.

Helfrich-Förster, C., 2002 The circadian system of Drosophila melanogaster and its light input pathways1. Zoology 105: 297–312.

Helfrich-Förster, C., T. Edwards, K. Yasuyama, B. Wisotzki, S. Schneuwly *et al.*, 2002 The extraretinal eyelet of Drosophila: development, ultrastructure, and putative circadian function. J. Neurosci. 22: 9255–9266.

Helfrich-Förster, C., and W. Engelmann, 2002 Photoreceptors for the Circadian Clock of the Fruitfly, pp. 94–106 in *Biological Rhythms*, edited by V. Kumar. Springer Berlin Heidelberg, Berlin, Heidelberg.

Hoang, N., E. Schleicher, S. Kacprzak, J.-P. Bouly, M. Picot *et al.*, 2008 Human and Drosophila Cryptochromes Are Light Activated by Flavin Photoreduction in Living Cells (U. Schibler, Ed.). PLoS Biol 6: e160.

Hofbauer, A., and E. Buchner, 1989 Does Drosophila have seven eyes? Naturwissenschaften 76: 335–336.

Klagges, B. R. E., G. Heimbeck, T. A. Godenschwege, A. Hofbauer, G. O. Pflugfelder *et al.*, 1996 Invertebrate Synapsins: A Single Gene Codes for Several Isoforms in *Drosophila* . J. Neurosci. 16: 3154–3165.

McNabb, S. L., and J. W. Truman, 2008 Light and peptidergic eclosion hormone neurons stimulate a rapid eclosion response that masks circadian emergence in Drosophila. J. Exp. Biol. 211: 2263–2274.

Melnattur, K. V., R. Pursley, T.-Y. Lin, C.-Y. Ting, P. D. Smith *et al.*, 2014 Multiple redundant medulla projection neurons mediate color vision in Drosophila. J. Neurogenet. 28: 374–388.

Mrosovsky, N., 1994 In praise of masking: behavioural responses of retinally degenerate mice to dim light. Chronobiol. Int. 11: 343–348.

Mrosovsky, N., 1999 Masking: history, definitions, and measurement. Chronobiol. Int. 16: 415–429.

Mrosovsky, N., and S. Hattar, 2003 Impaired masking responses to light in melanopsin-knockout mice. Chronobiol. Int. 20: 989–999.

Nagel, G., T. Szellas, W. Huhn, S. Kateriya, N. Adeishvili *et al.*, 2003 Channelrhodopsin-2, a directly light-gated cation-selective membrane channel. Proc. Natl. Acad. Sci. U.S.A. 100: 13940–13945.

Panda, S., I. Provencio, D. C. Tu, S. S. Pires, M. D. Rollag *et al.*, 2003 Melanopsin is required for non-image-forming photic responses in blind mice. Science 301: 525–527.

Pauls, D., C., Blechschmidt, F., Frantzmann, B., elJundi, and S. Selcho, 2018 Comprehensive anatomical map of the peripheral octopaminergic/tyraminergic system of Drosophila melanogaster. Sci Rep. 8:15314.

Pollack, I., and A. Hofbauer, 1991 Histamine-like immunoreactivity in the visual system and brain of Drosophila melanogaster. Cell Tissue Res 266: 391–398.

Pollock, J. A., and S. Benzer, 1988 Transcript localization of four opsin genes in the three visual organs of Drosophila; RH2 is ocellus specific. Nature 333: 779–782.

Redlin, U., and N. Mrosovsky, 1999 Masking by light in hamsters with SCN lesions. J. Comp. Physiol. A 184: 439–448.

Redlin, U., N. Vrang, and N. Mrosovsky, 1999 Enhanced masking response to light in hamsters with IGL lesions. J. Comp. Physiol. A 184: 449–456.

Rister, J., and C. Desplan, 2011 The retinal mosaics of opsin expression in invertebrates and vertebrates. Dev Neurobiol 71: 1212–1226.

Ruf, F., M. Fraunholz, K. Öchsner, J. Kaderschabek, and C. Wegener, 2017 WEclMon - A simple and robust camera-based system to monitor Drosophila eclosion under optogenetic manipulation and natural conditions. PLoS ONE 12: e0180238.

Sabat, D., S. Priyadarsini, and M. Mishra, 2016a Understanding the Structural and Developmental Aspect of Simple Eye of Drosophila: The Ocelli. Journal of Cell Signaling 1: 1–10.

Salcedo, E., A. Huber, S. Henrich, L. V. Chadwell, W. H. Chou *et al.*, 1999 Blue- and green-absorbing visual

pigments of Drosophila: ectopic expression and physiological characterization of the R8
photoreceptor cell-specific Rh5 and Rh6 rhodopsins. J. Neurosci. 19: 10716–10726.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair *et al.*, 2012 Fiji: an open-source platform
for biological-image analysis. Nat Methods 9: 676–682.

Schlichting, M., R. Grebler, P. Menegazzi, and C. Helfrich-Förster, 2015 Twilight Dominates Over Moonlight
in Adjusting *Drosophila*'s Activity Pattern. J Biol Rhythms 30: 117–128.

Schlichting, M., R. Grebler, N. Peschel, T. Yoshii, and C. Helfrich-Förster, 2014 Moonlight Detection by
Drosophila's Endogenous Clock Depends on Multiple Photopigments in the Compound Eyes. J Biol
Rhythms 29: 75–86.

Schlichting, M., P. Menegazzi, M. Rosbash, and C. Helfrich-Förster, 2019 A distinct visual pathway mediates
high light intensity adaptation of the circadian clock in *Drosophila*. J. Neurosci. 1497–18.

Schnaitmann, C., C. Garbers, T. Wachtler, and H. Tanimoto, 2013 Color discrimination with broadband
photoreceptors. Curr. Biol. 23: 2375–2382.

Selcho, M., C. Millán, A. Palacios-Muñoz, F. Ruf, L. Ubillo *et al.*, 2017 Central and peripheral clocks are
coupled by a neuropeptide pathway in Drosophila. Nat Commun 8: 15563.

Sprecher, S. G., and C. Desplan, 2008 Switch of rhodopsin expression in terminally differentiated Drosophila
sensory neurons. Nature 454: 533–537.

Stanewsky, R., M. Kaneko, P. Emery, B. Beretta, K. Wager-Smith *et al.*, 1998 The cryb Mutation Identifies
Cryptochrome as a Circadian Photoreceptor in Drosophila. Cell 95: 681–692.

Vanin, S., S. Bhutani, S. Montelli, P. Menegazzi, E. W. Green *et al.*, 2012 Unexpected features of Drosophila
circadian behavioural rhythms under natural conditions. Nature 484: 371–375.

Wernet, M. F., M. M. Velez, D. A. Clark, F. Baumann-Klausener, J. R. Brown *et al.*, 2012 Genetic dissection
reveals two separate retinal substrates for polarization vision in Drosophila. Curr. Biol. 22: 12–20.

Yasuyama, K., and I. A. Meinertzhagen, 1999 Extraretinal Photoreceptors at the Compound Eye's Posterior
Margin inDrosophila melanogaster. J. Comp. Neurol. 412: 193–202.

Yoshii, T., T. Todo, C. Wülbeck, R. Stanewsky, and C. Helfrich-Förster, 2008 Cryptochrome is present in the
compound eyes and a subset ofDrosophila's clock neurons. J. Comp. Neurol. 508: 952–966.

Zhu, L., R. R. McKay, and R. D. Shortridge, 1993 Tissue-specific expression of phospholipase C encoded by
the norpA gene of Drosophila melanogaster. Journal of Biological Chemistry 268: 15994–16001.

547
548
549

Acknowledgments

551 We thank Claude Desplan, Craig Montell, Christopher Schnaitmann and Ralf Stanewsky

552 for providing flies and Simon Sprecher for sharing antibodies. This work was supported by

553 grants from the German Research Foundation (DFG) to MS (PA3241/2-1), CHF

554 (FO207/18-1), DP (PA1979/2-1). Stocks obtained from the Bloomington Drosophila Stock

555 Center (NIH P40OD018537) were used in this study.

Author contributions

558 MS, DB, CHF and DP conceived and designed the experiments. DB, NDF performed the

559 experiments. TT wrote code for analysis. MS wrote the main manuscript text with

560 significant input from DP and CHF. MS and DB prepared the figures. All authors reviewed

561 the manuscript.

Data availability statement

The datasets generated during and analysed during the current study are available from the corresponding author on reasonable request.

All data generated and analysed during this study are included in this published article (and its Supplementary Information files).

Additional Information

571
572

Fig.S1: The immediate light effect on eclosion behaviour. (A) The light response is gone in flies lacking the compound eyes and CRY function (cli^{eya}; cry^b). **(B)** Flies without Rh6 ($rh6^1$) respond to light. **(C)** Projection of a head section visualizing Gal4-expression in $rh5$-Gal4-positive cells. The expression can be seen in a subset of R8 cells. Orange = α-Synapsin; white = α-GFP, scale bar: 100µm. n_{exp}, n_{ctrl} = 506, 549 (A), 504,731 (B)

578
579
580

Fig.S2: The immediate light effect on locomotor activity is visible in flies lacking photosensation in
***cry*-positive or photoreceptor cells. (A-F)** All tested flies respond to the light pulse by a significant
increase in locomotor activity compared to the appropriate controls. **(A)** Activity data of flies lacking
Cryptochrome (*cry^b*), **(B)** flies with impaired phospholipase C activity (*norpA^{p41}*), **(C)** flies lacking Rh1
(*ninaE^{17}*), **(C)** Rh3, Rh4, Rh5 and Rh6 (*rh5^2; rh3^1, rh4^1, rh6^1*), **(D)** R7 (*sev^{LY3}*) and **(E)** Rh5 (*rh5^2*). n = 14- 32;
asterisks denote level of significance: **p≤0.01, ***p≤0.001, ****p≤0.0001.
588
589

590
Fig.S3: Analysis of the eclosion monitor data. (A) Several pupae before hatching. Notice the empty pupal
case marked with an asterisk. **(B)** Pupae are darker than the background. Pixels with a value below the
threshold are coloured red. Yellow outlines mark the different objects (i.e. pupae) that fall within the given
size constraints. Notice that the empty pupal chase (marked with an asterisk) is excluded. **(C)** Outlines of the
different pupae. Median grey values are calculated of each of these areas over time (i.e. for every frame).
The pupae marked 1-4 are shown as examples in D-F. **(D)** Median grey values for pupae 1 to 4 over time.
Notice the big jump in brightness from around 40 to around 100 at different points in time. **(E)** Montage of
each 2 frames before and after hatching for 4 different pupae. In row 4 the eclosion process can be seen. **(F)**
Median grey values for the pupae shown in E.
600
601
602

603 **Competing interests**

604 The authors declare no competing interests.