Development of genomic resources for cattails (*Typha*), a globally important macrophyte

genus

Running title: Development of genomic resources for *Typha*

Alberto Aleman[1*], Marcel E. Dorken[1,2], Aaron B. A. Shafer[1,3], Tulsi Patel[2], Polina A.

Volkova[4], Joanna R. Freeland[1,2*]

[1]Environmental and Life Sciences Graduate Program, Trent University, Peterborough,

Ontario, Canada

[2]Department of Biology, Trent University, Peterborough, Ontario, Canada

[3]Department of Forensic Sciences, Trent University, Peterborough, Ontario, Canada

[4]Papanin Institute for Biology of Inland Waters, Russian Academy of Sciences, Borok,

Nekouz District, Yaroslavl Region, Russia

*Corresponding authors:

Joanna Freeland: joannafreeland@trentu.ca

Alberto Aleman: albertolopezaleman@trentu.ca

Abstract

A critical knowledge gap in freshwater plants research is the lack of genetic resources

necessary to answer fundamental questions about their demographic histories, adaptation, and

taxonomy. One example of this is *Typha*, a macrophyte genus essential to wetlands that is

also becoming an increasingly problematic biological invader in numerous regions

worldwide; while important insights have been discovered for this genus, currently available

genetic markers are insufficient to resolve its phylogenetic relationships, population structure,

and hybridization dynamics. We performed a cost and time–accessible library preparation for

high–throughput sequencing to develop a suite of genomic resources for *Typha*. Genome–

wide nuclear SNPs of 140 *Typha* samples from North America, temperate Eurasia and Africa

revealed three independent genetic clusters corresponding to *T. angustifolia*, *T. domingensis*

and *T. latifolia*. Data from ~40% of the nuclear genome was obtained, permitting the

characterization of 119,324 nuclear diagnostic markers (SNPs that differentiate the three

species). A reference–guided workflow to reconstruct whole–chloroplast–genome sequences

was implemented, recovering ~60% of the genome per sample. Three genetic lineages were

identified from the cpDNA phylogenetic analysis, agreeing with those determined from

nrDNA. With a cost below 15 USD per sample and a processing time of two hours for the

library preparation, this is a rapid and cost–effective protocol for population genomic studies.

The genomic locations of diagnostic markers and the chloroplast sequences produced in this

study will be permanent resources that can be incorporated into future studies of *Typha,* a

globally important and evolutionarily dynamic genus.

Keywords: Hybridization, diagnostic markers, chloroplast–genome assembly, high–

throughput sequencing, *Typha* phylogenomics, introgression.

Introduction

Freshwater plants are essential to aquatic ecosystems, shaping their habitats' structure and ecological functions (Chambers et al., 2008; Christie et al., 2009; Rejmankova, 2011). Although freshwater plants have been increasingly incorporated into applications that include habitat restoration and invasive species management, they remain highly understudied compared to terrestrial plants (Evangelista et al., 2014; Iversen et al., 2022). One key knowledge gap in freshwater plant research is the lack of genetic tools necessary to answer fundamental questions about their demographic histories, adaptation, and phylogenetic relationships (Fay et al., 2019; Maréchal, 2019; O'Hare et al., 2018; Yannelli et al., 2022).

Genomic characterization of freshwater plants has been hampered by biological and technical challenges, plus biases in scientific research (Matheson & McGaughran, 2022; Troudet et al., 2017). Hundreds to thousands of molecular markers are often required for genomic–based research on topics such as gene flow and adaptation of non–model organisms (da Fonseca et al., 2016; Stapley et al., 2010), but the *de novo* development of genomic resources can be both time–consuming and expensive (Hu et al., 2020; Ortega et al., 2020; Prieto et al., 2021). Overcoming these challenges is feasible with novel, rapid and cost–effective methods that capture genome–wide genetic variation, allowing researchers to address questions related to taxonomy, evolution, and conservation (Andrews et al., 2016; Goodwin et al., 2016).

Typha L. (cattails) is a global genus of freshwater plants foundational to wetlands (reviewed in Bansal et al., 2019). Cattails are a valuable ecosystemic and human resource that play a fundamental ecological role by cycling nutrients, preventing erosion, maintaining stable water levels and providing food and shelter for wildlife (Andrews & Pratt, 1978; Bonanno & Cirelli, 2017; Dieye et al., 2017; Kimmerer, 2013; Svedarsky et al., 2019). While

their importance is undisputed, one major challenge in *Typha* research has been taxonomic

identification, which cannot be fully accomplished using morphological characters due to

their high intraspecific variability and interspecific hybridization. Consequently, the richness

of cattail species, their taxonomy, their provenance (e.g., alien versus native lineages), and

their phylogenetic relationships remain incompletely resolved (Ciotir & Freeland, 2016;

Volkova & Bobrov, 2022; Zhou et al., 2018). A refined *Typha* taxonomy accompanied by

diagnostic markers, i.e., those loci that have alternative fixed alleles between two species, is

necessary to identify lineages of both introduced and hybrid taxa, which are increasingly

documented as invasive, e.g., *T. domingensis* Pers. in Western Europe and Central America,

T. latifolia L. in Oceania, and *T. × glauca* Godr. (*T. angustifolia* L. × *T. latifolia* L.) in North

America (Bansal et al., 2019; *GISD*; Govaerts, 2004; Hall, 2009; Maldonado, 2019).

High–throughput sequencing technologies, novel, cost and time–accessible genome

library preparations, and the recent assembly of the *T. latifolia* genome (Goodwin et al., 2016;

Rowan et al., 2019; Widanagama et al., 2022) collectively present an opportunity to develop a

suite of genomic resources for *Typha*. In addition to taxonomic resolution, these resources can

facilitate investigations of the genetic variation, local adaptation, population structure,

evolutionary history, and hybridization dynamics in this genus. We applied a high–throughput

sequencing protocol for enzymatic library preparation and data processing to produce

genome–wide resources in *Typha* spp. By optimizing the method from Rowan et al. (2019),

we generated a catalogue of nuclear SNPs that differentiate *T. angustifolia*, *T. domingensis*,

and *T. latifolia*, plus extensive chloroplast genome sequences in a fast, straightforward, and

cost–efficient manner. These permanent resources will be incorporated into future

evolutionary and population genetics research in *Typha*.

Materials and methods

Reference genome

We used the reference–based scaffolder Chromosomer 0.1.4a (Tamazian et al., 2016) to align the 1158 *T. latifolia* scaffolds from Widanagama et al. (2022) (Genbank accession: JAIOKV000000000.1) with the *T. latifolia* isolate L0001 (15 chromosomes, GenBank accession: JAAWWQ000000000.1) to produce a local chromosome–level *T. latifolia* genome. Widanagama et al. (2022) had a significantly higher mapping success of unrelated re–sequenced *Typha* spp. compared to the isolate L0001, suggesting it was a more representative *Typha* genome assembly. The scaffolds were aligned as chromosomes using BLAST+ 2.12.0 (Camacho et al., 2009) with the software default settings, and the alignments were anchored in Chromosomer, establishing a gap length = 0 and a ratio threshold = 1.

Sampling, DNA extraction, and sequencing

DNA extractions were obtained either from previous studies (Bhargav et al., 2022; Ciotir et al., 2017; Pieper et al., 2020, 2017; Tangen et al., 2022; Tisshaw et al., 2020) or from samples collected in Eurasia and extracted at Trent University following the published protocols. We obtained DNA from 140 *T. angustifolia*, *T. domingensis*, and *T. latifolia* samples (Figure 1), previously identified using a combination of morphological characteristics (Grace & Harrison, 1986; Smith, 1967) and genetic analyses of microsatellite loci (Bansal et al., 2019; Kirk et al., 2011; Pieper et al., 2020; Snow et al., 2010). Extracted DNA was quantified using a Qubit fluorometer (Thermofisher Scientific) and calculated as the mean of three independent readings from each sample. All samples were either standardized to 2 ng/μL by dilution with nuclease–free water or, if at concentrations less than 2 ng/μL (0.4 – 1.9 ng/μL), left undiluted.

Each DNA sample was prepared for Nextera XT sequencing libraries by enzymatic

fragmentation followed by ligation of short adapter sequences with the Illumina Tagment

DNA enzyme (TD) and buffer kit (small kit, #20034210). As the ratio of TD enzyme to DNA

is crucial for the reaction, we initially followed the recommendations of Rowan et al. (2019)

and subsequently optimized the reagent volumes for our library preparation as 5.5 μL of 5×

TD buffer, 0.5 μL of 1× TD enzyme, and 4 μL of DNA (standardized or undiluted), keeping

all reagents on ice during the preparation. Samples were incubated at 55°C for 10 minutes and

left at room temperature for five minutes. 5 μL of each sample were run on an agarose gel to

confirm the efficacy of the tagmentation reaction, evidenced by visible smears. The

fragmented DNA was then amplified using unique dual indexing based on combinations from

a total of 24 N7 (47 bases) and 8 S5 (51 bases) adapters (Alpha DNA, Canada). The PCR

cocktail included 0.2 μM of each index, 0.5U of KAPA HiFi HotStart DNA polymerase

(Roche), 12.5 μL of 5× KAPA reagent, 5 μL of tagmented DNA, and 6.5 μL of nuclease–free

water to a final volume of 25 μL. The PCR cycle comprised 72°C (3 minutes); 95°C (30

seconds); and 14 cycles of 95°C (10 seconds), 55°C (30 seconds), and 72°C (30 seconds).

Visible smears confirmed amplification success after running 5 μL of the PCR product on an

agarose gel; then, 10 μL from each sample were pooled, and the remaining PCR products

were stored at -20°C. The pooled library was purified with a QIAquick PCR purification kit

(QIAGEN) following the manufacturer's protocol, with a final elution in 50 μL of buffer. The

library was quantified using a D1000 Tapestation assay (Agilent Technologies, USA) and a

Qubit fluorometer (Thermo Fisher Scientific). A quality–control paired–end sequencing was

executed using a Miseq (151 bp) to ensure the library was compiled successfully. Finally,

paired–end sequencing was performed on a Novaseq 6000 (126 bp) at The Centre for Applied

Genomics (Toronto, Ontario).

Raw data processing, filtering, and SNP–calling

The quality of the demultiplexed raw sequences was evaluated using FastQC 0.11.9

(Andrews, 2017) and MultiQC 1.14 (Ewels et al., 2016). Read pairing and adapter removal

were carried out with trimmomatic 0.39 (Bolger et al., 2014), removing any cleaned reads

shorter than 100 bp. Paired and remaining unpaired reads were mapped to our chromosome–

level *T. latifolia* nuclear genome and the *T. latifolia* chloroplast reference (Genbank

accession: NC_013823.1) using the mem module of BWA 0.7.17 (Li & Durbin, 2009).

Mapped reads from Miseq and Novaseq 6000 sequencers were merged, and mapping statistics

were evaluated with the flagstat and coverage modules of SAMtools 1.15.1 (Li et al., 2009).

SNP–calling was performed with ANGSD 0.93 (Korneliussen et al., 2014) following

the SAMtools model, exclusively retrieving SNPs with a minimum *P*–value of $1e^{-6}$, minimum

mapping and sequencing qualities of 20, omitting any triallelic sites, and outputting a variant

call format (VCF) file (*-doGeno 4 -gl 1 -skipTriallelic 1 -SNP_pval 1e-6 -minMapQ 20 -minQ*

20 -doMajorMinor 1 -domaf 1 -doPost 1 -doCounts 1 -dovcf 1). For the nuclear analyses,

SNPs with > 50% missing data across all samples and sites that mapped to the chloroplast

genome were removed with vcftools 0.1.16 (Danecek et al., 2011). We did not apply

additional filters to SNP identification: our samples represented broad geographical sampling

and thus were not expected to be in Hardy–Weinberg equilibrium; additionally, as allele

frequencies were unlikely to be representative of regional allele frequencies we did not apply

a minor allele frequency filter. Neither did we filter for linkage equilibrium, as eliminating

alleles that are in linkage disequilibrium is likely to decrease the power of analyses to detect

hybridization and introgression (Alexander, 2020; Pearman et al., 2022).

Genetic structure and diagnostic markers

We first used nuclear SNPs to assess the most likely number of genetic clusters across all samples and the membership of each plant to these clusters using three complementary approaches. ADMIXTURE 1.3.0 (Alexander & Lange, 2011) was run with K = 1 – 10, and the optimal number of clusters was chosen via the cross–validation procedure. A neighbour–joining tree from the samples' pairwise genetic distance matrix (transformed on the R 4.2.2 package ape (Paradis et al., 2004; R Core Team, 2022)), and a Principal Component Analysis (PCA) without an *a priori* assumption were produced in Plink 1.90 (Purcell et al., 2007). We verified that the assignment of samples to genetic clusters (corresponding to three species, see *Results*) was consistent for each approach.

Potential introgression was tested by running ADMIXTURE (K = 1 – 5) on three datasets, each comprising a unique combination of two genetic clusters, using only those SNPs that remained variable based on the two clusters being compared. We confirmed that the cross–validation procedure for the runs of each cluster pair chose the optimal number of clusters as two and used the admixture proportion (Q score in $K = 2$) as an index of potential introgression of each sample. Applying Senn & Pemberton (2009) and Smith et al. (2018) thresholds, individuals whose Q score was $0.05 \le Q \le 0.95$, were considered as introgressed.

To compare the levels of differentiation between clusters, values of Weir and Cockerham's genetic differentiation (F_{ST}; Weir & Cockerham, 1984) for every variable site and genetic divergence (d_{xy}; Nei & Miller, 1990) in 10 Kbp windows between species pairs were computed using pixy 1.2.7 (Korunes & Samuk, 2021), and the means were calculated. Species–specific SNPs for each pair of genetic clusters were identified with DiagnoSNPs 1.0 (Arce-Valdés, 2022). We removed introgressed individuals before estimating genetic differentiation levels and identifying species–specific SNPs.

Chloroplast genome reconstruction and phylogenetic analysis

We implemented a reference–guided workflow to reconstruct whole–chloroplast–

genome sequences. Nucleotide calling was performed individually for each of the 140

samples in ANGSD using the reads that mapped to the chloroplast genome reference,

requiring minimum mapping and base qualities of 20, and using Ns as missing data (*-dofasta*

2 -minMapQ 20 -minQ 20 -doCounts 1). Sequences were aligned to the chloroplast genomes

of *T. przewalskii* Skvortsov, *T. lugdunensis* P. Chabert, *T. orientalis* C. Presl, and

Sparganium natans (GenBank accessions: NC_061354.1, NC_061353.1, NC_050678.1, and

NC_058577.1), following Smith et al. (2021) by applying MAFFT 7.0 default settings (plus

the flag *--nwildcard*) (Katoh et al., 2019). Snp–sites 2.5.1 (Page et al., 2016) was used to

remove sections of the genome with ambiguous positions, gaps, and missing data, such that if

any of those was found in a sequence, that position was removed for all sequences.

Nucleotide diversity (π) was calculated in the R package pegas (Paradis, 2010).

The phylogenetic relationships of the chloroplast genome sequences were

reconstructed through RAxML-NG 1.1 (Kozlov et al., 2019). Model selection was based on

jmodeltest 2.1.10 results (Darriba et al., 2012) using the AIC. A transversion model was

established, assuming gamma–distributed rates with four substitution categories and

estimating the proportion of invariable sites (TVM + G4 + I). RAxML was run with a

thorough bootstrap starting from 100 random trees, with *Sparganium natans* as the outgroup,

and letting automatic bootstrapping (--autoMRE). The best–scoring ML tree was plotted.

Results

Genome scaffolding, mapping statistics, and genotyping

Approximately 99.81% of the scaffold sequences were aligned to the template genome. The scaffolds were anchored to 15 chromosomes producing a genome of 285.11 Mb (Pending GenBank accession: JAIOKV000000000.2). The total assembled size was comparable to the *T. latifolia* genome sizes of Widanagama et al. (2022) (287.19 Mb) and the isolate L0001 (214.13 Mb). Updating the chromosome–level assembly genome simplified our downstream analyses while keeping the highest mapping success of unrelated re–sequenced *Typha* spp. and facilitating an accurate recombination map for future studies of speciation, hybridization, and the genomic landscape of introgression in *Typha*.

After quality control, 982 M clean paired–end reads were retained, and ~98% mapped to the reference genome. With minimum mapping and sequencing qualities = 20, the average depth and breadth of coverage of the nuclear sequences were 4× and 42%, respectively. Over 60% of the chloroplast genome breadth was covered across all samples (mean depth = 711×), enabling us to use 96,591 bp for the phylogenetic reconstruction. We assembled 12,177,703 bi–allelic nuclear SNPs across the 140 *Typha* samples (7,122,151 with a MAF > 0.05). The total genotyping rate, i.e., the mean proportion of samples with data for each SNP, was 0.68.

Genetic structure and diagnostic markers

The admixture analysis, the PCA, and the neighbour–joining tree each established the most likely number of genetic clusters as three ($K = 3$) (Figure 1): in line with previous taxonomic identifications, 38 samples were identified within the *T. angustifolia* cluster (15 of which had *T. latifolia* introgression and 5 of which had both *T. domingensis* and *T. latifolia* introgression); 25 samples were in the *T. domingensis* cluster (one with *T. angustifolia* introgression, 12 with *T. latifolia* introgression, and 3 with both *T. angustifolia* and *T. latifolia*

introgression); and 77 samples were in the *T. latifolia* cluster (one with both *T. angustifolia*

and *T. domingensis* introgression, and one with *T. angustifolia* introgression). Using only the

18 *T. angustifolia*, 9 *T. domingensis*, and 75 *T. latifolia* not introgressed individuals, the mean

pairwise interspecific F_{ST} and d_{xy} values ranged from 0.25 to 0.49 and 0.28 to 0.35,

respectively, with *T. latifolia* showing the highest differentiation with both *T. angustifolia* and

T. domingensis. We identified 119,324 nuclear species-specific SNPs (Table 1) based on the

comparisons between species pairs.

Chloroplast genome reconstruction and phylogenetic relationships

After removing all ambiguities and missing data from the chloroplast genomes, we

were left with an alignment of 96,591 bp across all 140 sequences and the four reference

genomes, with 4,916 segregating sites and $\pi = 0.003$. The phylogenetic reconstruction was

consistent with the nuclear genetic structure results, grouping the 143 *Typha* samples into

three lineages, with *T. angustifolia* in one clade, *T. domingensis* and *T. orientalis* sharing

another, and *T. latifolia* and *T. przewalskii* in a third clade (Figure 2). All interspecific nodes

were strongly supported (100%, Felsenstein bootstrap proportion).

Discussion

We aimed to produce a suite of genome–wide resources to facilitate investigations into

the taxonomy and population genetics of *Typha* and to advance the general understanding of

wetland plants. Following a fast, straightforward, and cost–efficient genomic library

preparation protocol (Rowan et al., 2019), we sequenced 140 *Typha* samples, obtaining an

average breadth of 42% of the nuclear genome, characterizing 119,324 nuclear SNPs that

collectively differentiate three *Typha* spp., and producing chloroplast sequences with a

breadth of > 60% per sample. With a cost below 15 USD per sample and a processing time of

two hours for the library preparation, our workflow is a rapid and cost–effective protocol that

can be applied in population genomic studies for investigating levels of genetic diversity and

differentiation, identifying conservation units and alien taxa, characterizing hybrid classes,

and examining hybridization dynamics, among other purposes.

249   Three genetic clusters were identified from both nuclear and chloroplast genomes,

corresponding to *T. angustifolia*, *T. domingensis*, and *T. latifolia*; and potential introgression

was detected among the three species. Hybridization of *T. domingensis* with *T. angustifolia*

and *T. latifolia* has been historically documented (Govaerts, 2004; Smith, 1967); however,

those inferences were based on experimental and taxonomical observations and recently

exposed with microsatellite loci by Ciotir et al. (2017). Future investigations should address

the extent to which hybridization is shaping the genetic variation among these three species.

Furthermore, *T. domingensis* is increasingly invading Costa Rica (Trama et al., 2017), Nigeria

(Ringim et al., 2016), and North America, potentially expanding its range across the latter

(Spencer & Vincent, 2013; Zhang et al., 2008). However, the taxonomic identity of these

plants is unclear –are they hybrids, non–native lineages, or native lineages responding to

environmental change? By identifying SNPs that characterize *T. domingensis*, we provide a

valuable resource to answer this and other questions across the evolutionary and hybridization

history of *Typha*.

263   The markers that differentiate *T. angustifolia* from *T. latifolia* will have important

applications in North America, where the two species interbreed across a large area and

produce an invasive interspecific hybrid (*T.* × *glauca*) that dominates wetlands, alters nutrient

cycling, and reduces biodiversity across the Great Lakes Region (Bansal et al., 2019);

additionally, this hybrid is expanding throughout the Prairie Pothole Region, causing native

plant diversity to decrease in invaded potholes (Jones et al., 2023), and may impact essential

habitat for millions of breeding and migratory waterfowl species (Tangen et al., 2022). Until

now, molecular resources to differentiate *T. angustifolia*, *T. latifolia*, and *T. × glauca* were

limited to sets of relatively few individual markers that have produced important insights:

RAPDs, chloroplast DNA sequences, and codominant SSR loci have contributed to exposing

the sexual fertility of first–generation hybrids (F1) (Snow et al., 2010), asymmetric

hybridization (with *T. angustifolia* being mainly the maternal parent) (Ball & Freeland, 2013;

Kuehn et al., 1999; Pieper et al., 2017), overall comparable levels of sexual and clonal

reproduction in parents and F1s (Pieper et al., 2020; Travis et al., 2011), heterosis in F1s

(Bunbury-Blanchette et al., 2015; Travis et al., 2010; Zapfe & Freeland, 2015), a high

frequency of F1s in natural populations (Kirk et al., 2011; Travis et al., 2010), and partial

sterility of F1s coupled with hybrid breakdown of F2s and advanced–generation hybrids

(Bhargav et al., 2022; Pieper et al., 2017). However, critical questions remain unresolved

because existent markers are insufficient to expose the prevalence of advanced–generation

hybrids and backcrosses in wild populations. The expansive suite of SNPs identified in this

study will facilitate investigations into the extent of hybridization, hybrid breakdown

dynamics, and adaptive introgression across the *T. × glauca* hybrid zone, allowing

researchers to understand the evolutionary processes shaping this genus speciation and

species boundaries, and to inform conservation and management strategies.

Fundamental genetic tools are essential for understanding macrophytes biology,

management, and conservation (O'Hare et al., 2018). The protocol described in this paper, the

updated chromosome–level assembly genome of *T. latifolia*, the SNPs catalogue, and the

chloroplast sequences produced for each sample comprise permanent resources that can be

applied to study the genetic composition of *Typha* populations and hybrid zones. Genome–

wide sequencing techniques and reference–based chloroplast genome assemblies are

promising tools to clarify the demographic histories, dispersal, adaptation, and taxonomy of

multiple congeneric macrophyte species (Russello et al., 2015; Straub et al., 2012), and

substantial genome–wide research will allow us to tackle these and other knowledge gaps in

Typha and other taxa.

Acknowledgements

We acknowledge that the laboratory procedures and data analyses were conducted at

Trent University, which is on the traditional territory of the Mississauga Anishinaabeg. The

Natural Sciences and Engineering Research Council of Canada (NSERC) financially

supported this work, and Alberto Aleman is funded by the Environmental & Life Sciences

Graduate Program at Trent University. The work of Polina A. Volkova was supported by the

Russian Science Foundation grant no. 23-14-00115. We thank V. Bhargav, N. Tikhomirov,

and M. Ivanova for providing plant tissue samples; T. Pimenov, M. Aksyonova and the staff

of the Dagestansky Nature Reserve, in particular, G. S. Dzhamirzoyev, for their help in the

field, AO "IEPI" for organizing fieldwork in Krasnodar Region, and SHARCNET and

Compute Canada for providing computational resources. Finally, we thank Camille Kessler

for her comments on the manuscript.

References

Alexander, A. (2020). *GBS_SNP_filter v1.17* [R].
https://github.com/laninsky/GBS_SNP_filter

Alexander, D. H., & Lange, K. (2011). Enhancements to the ADMIXTURE algorithm for
individual ancestry estimation. *BMC Bioinformatics*, *12*(1), 246.
https://doi.org/10.1186/1471-2105-12-246

Andrews, K. R., Good, J. M., Miller, M. R., Luikart, G., & Hohenlohe, P. A. (2016).
Harnessing the power of RADseq for ecological and evolutionary genomics. *Nature*
Reviews Genetics, *17*(2), Article 2. https://doi.org/10.1038/nrg.2015.28

Andrews, N., & Pratt, D. (1978). Energy Potential of Cattails (Typha spp.) and Productivity in
Managed Stands. *Journal of the Minnesota Academy of Science*, *44*(2), 5–8.

Andrews, S. (2017). *FastQC: A quality control tool for high throughput sequence data. 2010.*

Arce-Valdés, L. R. (2022). *LuisRodrigoArce-Valdes/DiagnoSNPs* [Shell].
https://github.com/LuisRodrigoArce-Valdes/DiagnoSNPs (Original work published
2021)

Ball, D., & Freeland, J. R. (2013). Synchronous flowering times and asymmetrical
hybridization in Typha latifolia and T. angustifolia in northeastern North America.
Aquatic Botany, *104*, 224–227. https://doi.org/10.1016/j.aquabot.2012.08.006

Bansal, S., Lishawa, S. C., Newman, S., Tangen, B. A., Wilcox, D., Albert, D., Anteau, M. J.,
Chimney, M. J., Cressey, R. L., DeKeyser, E., Elgersma, K. J., Finkelstein, S. A.,
Freeland, J., Grosshans, R., Klug, P. E., Larkin, D. J., Lawrence, B. A., Linz, G.,
Marburger, J., … Windham-Myers, L. (2019). Typha (Cattail) Invasion in North
American Wetlands: Biology, Regional Problems, Impacts, Ecosystem Services, and
Management. *Wetlands*, *39*(4), 645–684. https://doi.org/10.1007/s13157-019-01174-7

Bhargav, V. V., Freeland, J. R., & Dorken, M. E. (2022). Evidence of hybrid breakdown
among invasive hybrid cattails (Typha × glauca). *Heredity*, *129*(3), Article 3.
https://doi.org/10.1038/s41437-022-00557-7

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for
Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170

Bonanno, G., & Cirelli, G. L. (2017). Comparative analysis of element concentrations and
translocation in three wetland congener plants: Typha domingensis, Typha latifolia
and Typha angustifolia. *Ecotoxicology and Environmental Safety*, *143*, 92–101.
https://doi.org/10.1016/j.ecoenv.2017.05.021

Bunbury-Blanchette, A. L., Freeland, J. R., & Dorken, M. E. (2015). Hybrid Typha×glauca
outperforms native T. latifolia under contrasting water depths in a common garden.
Basic and Applied Ecology, *16*(5), 394–402.
https://doi.org/10.1016/j.baae.2015.04.006

Camacho, C., Coulouris, G., Avagyan, V., Ma, N., Papadopoulos, J., Bealer, K., & Madden,
T. L. (2009). BLAST+: Architecture and applications. *BMC Bioinformatics*, *10*(1),
421. https://doi.org/10.1186/1471-2105-10-421

Chambers, P. A., Lacoul, P., Murphy, K. J., & Thomaz, S. M. (2008). Global diversity of
aquatic macrophytes in freshwater. In E. V. Balian, C. Lévêque, H. Segers, & K.
Martens (Eds.), *Freshwater Animal Diversity Assessment* (pp. 9–26). Springer
Netherlands. https://doi.org/10.1007/978-1-4020-8259-7_2

Christie, H., Norderhaug, K. M., & Fredriksen, S. (2009). Macrophytes as habitat for fauna.
Marine Ecology Progress Series, *396*, 221–233. https://doi.org/10.3354/meps08351

Ciotir, C., & Freeland, J. (2016). Cryptic intercontinental dispersal, commercial retailers, and
the genetic diversity of native and non-native cattails (Typha spp.) in North America.
Hydrobiologia, *768*(1), 137–150. https://doi.org/10.1007/s10750-015-2538-0
Ciotir, C., Szabo, J., & Freeland, J. (2017). Genetic characterization of cattail species and
hybrids (Typha spp.) in Europe. *Aquatic Botany*, *141*, 51–59.
https://doi.org/10.1016/j.aquabot.2017.03.005
da Fonseca, R. R., Albrechtsen, A., Themudo, G. E., Ramos-Madrigal, J., Sibbesen, J. A.,
Maretty, L., Zepeda-Mendoza, M. L., Campos, P. F., Heller, R., & Pereira, R. J.
(2016). Next-generation biology: Sequencing and data analysis approaches for non-
model organisms. *Marine Genomics*, *30*, 3–13.
https://doi.org/10.1016/j.margen.2016.04.012
Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker,
R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000
Genomes Project Analysis Group. (2011). The variant call format and VCFtools.
Bioinformatics, *27*(15), 2156–2158. https://doi.org/10.1093/bioinformatics/btr330
Darriba, D., Taboada, G. L., Doallo, R., & Posada, D. (2012). jModelTest 2: More models,
new heuristics and high-performance computing. *Nature Methods*, *9*(8), 772.
https://doi.org/10.1038/nmeth.2109
Dieye, Y., Sambou, V., Faye, M., Thiam, A., Adj, M., & Azilinon, D. (2017). Thermo-
mechanical characterization of a building material based on Typha Australis. *Journal
of Building Engineering*, *9*, 142–146. https://doi.org/10.1016/j.jobe.2016.12.007
Evangelista, H. B. A., Thomaz, S. M., & Umetsu, C. A. [UNESP. (2014). An analysis of
publications on invasive macrophytes in aquatic ecosystems. *Aquatic Invasions*, 521.
https://doi.org/10.3391/ai.2014.9.4.10
Ewels, P., Magnusson, M., Lundin, S., & Käller, M. (2016). MultiQC: Summarize analysis
results for multiple tools and samples in a single report. *Bioinformatics*, *32*(19), 3047–
3048. https://doi.org/10.1093/bioinformatics/btw354
Fay, M. F., Gargiulo, R., & Viruel, J. (2019). The present and future for population genetics,
species boundaries, biogeography and conservation. *Botanical Journal of the Linnean
Society*, *191*(3), 299–304. https://doi.org/10.1093/botlinnean/boz076
GISD. (n.d.). Retrieved March 13, 2023, from
http://www.iucngisd.org/gisd/speciesname/Typha+latifolia
Goodwin, S., McPherson, J. D., & McCombie, W. R. (2016). Coming of age: Ten years of
next-generation sequencing technologies. *Nature Reviews Genetics*, *17*(6), Article 6.
https://doi.org/10.1038/nrg.2016.49
Govaerts, R. (2004). The Monocot Checklist Project. *TAXON*, *53*(1), 144–146.
https://doi.org/10.2307/4135499
Grace, J., & Harrison, J. (1986). The biology of Canadian weeds. 73. Typha latifolia L.,
Typha angustifolia L. and Typha xglauca Godr. *Canadian Journal of Plant Science -
CAN J PLANT SCI*, *66*, 361–379. https://doi.org/10.4141/cjps86-051
Hall, S. J. (2009). Cultural Disturbances and Local Ecological Knowledge Mediate Cattail
(Typha domingensis) Invasion in Lake Pátzcuaro, México. *Human Ecology*, *37*(2),
241–249. https://doi.org/10.1007/s10745-009-9228-3
Hu, Z.-M., Zhong, K.-L., Weinberger, F., Duan, D.-L., Draisma, S. G. A., & Serrão, E. A.
(2020). Linking Ecology to Genetics to Better Understand Adaptation and Evolution:
A Review in Marine Macrophytes. *Frontiers in Marine Science*, *7*.
https://www.frontiersin.org/articles/10.3389/fmars.2020.545102

Iversen, L. L., Girón, J. G., & Pan, Y. (2022). Towards linking freshwater plants and ecosystems via functional biogeography. *Aquatic Botany*, *176*, 103454. https://doi.org/10.1016/j.aquabot.2021.103454

Jones, S. A., DeKeyser, E. S., Dixon, C., & Kobiela, B. (2023). Invasive Species Change Plant Community Composition of Preserved Prairie Pothole Wetlands. *Plants*, *12*(6), Article 6. https://doi.org/10.3390/plants12061281

Katoh, K., Rozewicki, J., & Yamada, K. D. (2019). MAFFT online service: Multiple sequence alignment, interactive sequence choice and visualization. *Briefings in Bioinformatics*, *20*(4), 1160–1166. https://doi.org/10.1093/bib/bbx108

Kimmerer, R. (2013). *Braiding Sweetgrass: Indigenous Wisdom, Scientific Knowledge and the Teachings of Plants*. Milkweed Editions.

Kirk, H., Connolly, C., & Freeland, J. R. (2011). Molecular genetic data reveal hybridization between Typha angustifolia and Typha latifolia across a broad spatial scale in eastern North America. *Aquatic Botany*, *95*(3), 189–193. https://doi.org/10.1016/j.aquabot.2011.05.007

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation Sequencing Data. *BMC Bioinformatics*, *15*(1), 356. https://doi.org/10.1186/s12859-014-0356-4

Korunes, K. L., & Samuk, K. (2021). pixy: Unbiased estimation of nucleotide diversity and divergence in the presence of missing data. *Molecular Ecology Resources*, *21*(4), 1359–1368. https://doi.org/10.1111/1755-0998.13326

Kozlov, A. M., Darriba, D., Flouri, T., Morel, B., & Stamatakis, A. (2019). RAxML-NG: A fast, scalable and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics*, *35*(21), 4453–4455. https://doi.org/10.1093/bioinformatics/btz305

Kuehn, M. M., Minor, J. E., & White, B. N. (1999). An examination of hybridization between the cattail species Typha latifolia and Typha angustifolia using random amplified polymorphic DNA and chloroplast DNA markers. *Molecular Ecology*, *8*(12), 1981–1990. https://doi.org/10.1046/j.1365-294x.1999.00792.x

Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: An online tool for phylogenetic tree display and annotation. *Nucleic Acids Research*, *49*(W1), W293–W296. https://doi.org/10.1093/nar/gkab301

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows–Wheeler transform. *Bioinformatics*, *25*(14), 1754–1760. https://doi.org/10.1093/bioinformatics/btp324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., Durbin, R., & 1000 Genome Project Data Processing Subgroup. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics*, *25*(16), 2078–2079. https://doi.org/10.1093/bioinformatics/btp352

Maldonado, G. (2019). *The Paradox of Culturally Useful Invasive Species: Chuspatel (Typha Domingensis) Crafts of Lake Patzcuaro, Mexico*.

Maréchal, E. (2019). Marine and Freshwater Plants: Challenges and Expectations. *Frontiers in Plant Science*, *10*, 1545. https://doi.org/10.3389/fpls.2019.01545

Matheson, P., & McGaughran, A. (2022). Genomic data is missing for many highly invasive species, restricting our preparedness for escalating incursion rates. *Scientific Reports*, *12*(1), Article 1. https://doi.org/10.1038/s41598-022-17937-y

Nei, M., & Miller, J. C. (1990). A simple method for estimating average number of nucleotide substitutions within and between populations from restriction data. *Genetics*, *125*(4), 873–879. https://doi.org/10.1093/genetics/125.4.873

O'Hare, M. T., Aguiar, F. C., Asaeda, T., Bakker, E. S., Chambers, P. A., Clayton, J. S.,
Elger, A., Ferreira, T. M., Gross, E. M., Gunn, I. D. M., Gurnell, A. M., Hellsten, S.,
Hofstra, D. E., Li, W., Mohr, S., Puijalon, S., Szoszkiewicz, K., Willby, N. J., &
Wood, K. A. (2018). Plants in aquatic ecosystems: Current trends and future
directions. *Hydrobiologia*, *812*(1), 1–11. https://doi.org/10.1007/s10750-017-3190-7
Ortega, A., Geraldi, N. R., Díaz-Rúa, R., Ørberg, S. B., Wesselmann, M., Krause-Jensen, D.,
& Duarte, C. M. (2020). A DNA mini-barcode for marine macrophytes. *Molecular*
Ecology Resources, *20*(4), 920–935. https://doi.org/10.1111/1755-0998.13164
Page, A. J., Taylor, B., Delaney, A. J., Soares, J., Seemann, T., Keane, J. A., & Harris, S. R.
(2016). SNP-sites: Rapid efficient extraction of SNPs from multi-FASTA alignments.
Microbial Genomics, *2*(4), e000056. https://doi.org/10.1099/mgen.0.000056
Paradis, E. (2010). pegas: An R package for population genetics with an integrated–modular
approach. *Bioinformatics*, *26*(3), 419–420.
https://doi.org/10.1093/bioinformatics/btp696
Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and
Evolution in R language. *Bioinformatics*, *20*(2), 289–290.
https://doi.org/10.1093/bioinformatics/btg412
Pearman, W. S., Urban, L., & Alexander, A. (2022). Commonly used Hardy–Weinberg
equilibrium filtering schemes impact population structure inferences using RADseq
data. *Molecular Ecology Resources*, *22*(7), 2599–2613. https://doi.org/10.1111/1755-
0998.13646
Pieper, S., Dorken, M., & Freeland, J. (2020). Genetic structure in hybrids and progenitors
provides insight into processes underlying an invasive cattail (Typha × glauca) hybrid
zone. *Heredity*, *124*(6), Article 6. https://doi.org/10.1038/s41437-020-0307-y
Pieper, S. J., Nicholls, A. A., Freeland, J. R., & Dorken, M. E. (2017). Asymmetric
Hybridization in Cattails (Typha spp.) and Its Implications for the Evolutionary
Maintenance of Native Typha latifolia. *Journal of Heredity*, *108*(5), 479–487.
https://doi.org/10.1093/jhered/esx036
Prieto, A. E., Hardion, L., & Beisel, J.-N. (2021). Designing robust DNA barcode libraries for
metabarcoding of freshwater plants by integrating herbarium collections and
contemporary floristic inventories. *ARPHA Conference Abstracts*, *4*, e64713.
https://doi.org/10.3897/aca.4.e64713
Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J.,
Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A Tool Set
for Whole-Genome Association and Population-Based Linkage Analyses. *The*
American Journal of Human Genetics, *81*(3), 559–575.
https://doi.org/10.1086/519795
R Core Team. (2022). *R: The R Project for Statistical Computing*. https://www.r-project.org/
Rejmankova, E. (2011). The role of macrophytes in wetland ecosystems. *Journal of Ecology*
and Environment, *34*(4), 333–345. https://doi.org/10.5141/JEFB.2011.044
Ringim, A., Babura, B., & Harry. (2016). Implication of Invasive Plant Typha domingensis
on Biodiversity: An Ecological Study of the Hadejia-Nguru Wetlands, Nigeria.
Scholarly Journal of Biological Science, *4*, 40–46.
Rowan, B. A., Heavens, D., Feuerborn, T. R., Tock, A. J., Henderson, I. R., & Weigel, D.
(2019). An Ultra High-Density Arabidopsis thaliana Crossover Map That Refines the
Influences of Structural Variation and Epigenetic Features. *Genetics*, *213*(3), 771–787.
https://doi.org/10.1534/genetics.119.302406

Russello, M. A., Waterhouse, M. D., Etter, P. D., & Johnson, E. A. (2015). From promise to
practice: Pairing non-invasive sampling with genomics in conservation. *PeerJ*, *3*,
e1106. https://doi.org/10.7717/peerj.1106
Senn, H. V., & Pemberton, J. M. (2009). Variable extent of hybridization between invasive
sika (Cervus nippon) and native red deer (C. elaphus) in a small geographical area.
Molecular Ecology, *18*(5), 862–876. https://doi.org/10.1111/j.1365-
294X.2008.04051.x
Smith, C. I., McKain, M. R., Guimond, A., & Flatz, R. (2021). Genome-scale data resolves
the timing of divergence in Joshua trees. *American Journal of Botany*, *108*(4), 647–
663. https://doi.org/10.1002/ajb2.1633
Smith, S. G. (1967). Experimental and Natural Hybrids in North American Typha
(Typhaceae). *The American Midland Naturalist*, *78*(2), 257–287.
https://doi.org/10.2307/2485231
Smith, S. L., Senn, H. V., Pérez-Espona, S., Wyman, M. T., Heap, E., & Pemberton, J. M.
(2018). Introgression of exotic Cervus (nippon and canadensis) into red deer (Cervus
elaphus) populations in Scotland and the English Lake District. *Ecology and*
Evolution, *8*(4), 2122–2134. https://doi.org/10.1002/ece3.3767
Snow, A. A., Travis, S. E., Wildová, R., Fér, T., Sweeney, P. M., Marburger, J. E., Windels,
S., Kubátová, B., Goldberg, D. E., & Mutegi, E. (2010). Species-specific SSR alleles
for studies of hybrid cattails (Typha latifolia × T. angustifolia; Typhaceae) in North
America. *American Journal of Botany*, *97*(12), 2061–2067.
Spencer, J. M., & Vincent, M. A. (2013). Southern cat-tail (Typha domingensis, Typhaceae)
discovered in Ohio. *Phytoneuron*, *2013–22*, 1–5.
Stapley, J., Reger, J., Feulner, P. G. D., Smadja, C., Galindo, J., Ekblom, R., Bennison, C.,
Ball, A. D., Beckerman, A. P., & Slate, J. (2010). Adaptation genomics: The next
generation. *Trends in Ecology & Evolution*, *25*(12), 705–712.
https://doi.org/10.1016/j.tree.2010.09.002
Straub, S. C. K., Parks, M., Weitemier, K., Fishbein, M., Cronn, R. C., & Liston, A. (2012).
Navigating the tip of the genomic iceberg: Next-generation sequencing for plant
systematics. *American Journal of Botany*, *99*(2), 349–364.
https://doi.org/10.3732/ajb.1100335
Svedarsky, D., Grosshans, R., Venema, H., Ellis-Felege, S., Bruggman, J., Ostlund, A., &
Lewis, J. (2019). Integrated management of invasive cattails (Typha spp.) for wetland
habitat and biofuel in the Northern Great Plains of the United States and Canada: A
review. *Mires and Peat*, *25*, 1–14. https://doi.org/10.19189/MaP.2018.APG.367
Tamazian, G., Dobrynin, P., Krasheninnikova, K., Komissarov, A., Koepfli, K.-P., &
O'Brien, S. J. (2016). Chromosomer: A reference-based genome arrangement tool for
producing draft chromosome sequences. *GigaScience*, *5*(1), 38.
https://doi.org/10.1186/s13742-016-0141-6
Tangen, B. A., Bansal, S., Freeland, J. R., Travis, S. E., Wasko, J. D., McGonigle, T. P.,
Goldsborough, L. G., Gow, K., Marburger, J. E., & Meier, J. A. (2022). Distributions
of native and invasive Typha (cattail) throughout the Prairie Pothole Region of North
America. *Wetlands Ecology and Management*, *30*(1), 1–17.
https://doi.org/10.1007/s11273-021-09823-7
Tisshaw, K., Freeland, J., & Dorken, M. (2020). Salinity, not genetic incompatibilities, limits
the establishment of the invasive hybrid cattail Typha × glauca in coastal wetlands.
Ecology and Evolution, *10*(21), 12091–12103. https://doi.org/10.1002/ece3.6831

Trama, F. A., Viale, F. L. S. R. P., Kumar, A. S., Stynoski, J. L., Colton, M. B. M., &
Springer, M. C. (2017). The Management of Typha domingensis (Typhaceae) affects
Macroinvertebrate Assemblages in the Palo Verde Wetland, Guanacaste, Costa Rica.
Ecological Restoration, *35*(2), 175–189. https://doi.org/10.3368/er.35.2.175

Travis, S. E., Marburger, J. E., Windels, S. K., & Kubátová, B. (2011). Clonal Structure of
Invasive Cattail (Typhaceae) Stands in the Upper Midwest Region of the US.
Wetlands, *31*(2), 221–228. https://doi.org/10.1007/s13157-010-0142-7

Travis, S. E., Marburger, J. E., Windels, S., & Kubátová, B. (2010). Hybridization dynamics
of invasive cattail (Typhaceae) stands in the Western Great Lakes Region of North
America: A molecular analysis. *Journal of Ecology*, *98*(1), 7–16.
https://doi.org/10.1111/j.1365-2745.2009.01596.x

Troudet, J., Grandcolas, P., Blin, A., Vignes-Lebbe, R., & Legendre, F. (2017). Taxonomic
bias in biodiversity data and societal preferences. *Scientific Reports*, *7*(1), Article 1.
https://doi.org/10.1038/s41598-017-09084-6

Volkova, P. A., & Bobrov, A. A. (2022). Easier than it looks: Notes on the taxonomy of
Typha L. (Typhaceae) in East Europe. *Aquatic Botany*, *176*, 103453.
https://doi.org/10.1016/j.aquabot.2021.103453

Weir, B. S., & Cockerham, C. C. (1984). Estimating F-Statistics for the Analysis of
Population Structure. *Evolution*, *38*(6), 1358–1370. https://doi.org/10.2307/2408641

Wickham, H. (2016). *Ggplot2*. Springer International Publishing. https://doi.org/10.1007/978-
3-319-24277-4

Widanagama, S. D., Freeland, J. R., Xu, X., & Shafer, A. B. A. (2022). Genome assembly,
annotation, and comparative analysis of the cattail Typha latifolia. *G3*
Genes|Genomes|Genetics, *12*(2), jkab401. https://doi.org/10.1093/g3journal/jkab401

Yannelli, F. A., Bazzichetto, M., Conradi, T., Pattison, Z., Andrade, B. O., Anibaba, Q. A.,
Bonari, G., Chelli, S., Ćuk, M., Damasceno, G., Fantinato, E., Geange, S. R., Guuroh,
R. T., Holle, M. J. M., Küzmič, F., Lembrechts, J. J., Mosyaftiani, A., Šikuljak, T.,
Teixeira, J., … Sperandii, M. G. (2022). Fifteen emerging challenges and
opportunities for vegetation science: A horizon scan by early career researchers.
Journal of Vegetation Science, *33*(1), e13119. https://doi.org/10.1111/jvs.13119

Zapfe, L., & Freeland, J. R. (2015). Heterosis in invasive F1 cattail hybrids (Typha × glauca).
Aquatic Botany, *125*, 44–47. https://doi.org/10.1016/j.aquabot.2015.05.004

Zhang, X.-H., Tapia, M., Webb, J. B., Huang, Y.-H., & Miao, S. (2008). Molecular signatures
of two cattail species, Typha domingensis and Typha latifolia (Typhaceae), in South
Florida. *Molecular Phylogenetics and Evolution*, *49*(1), 368–376.
https://doi.org/10.1016/j.ympev.2008.03.032

Zhou, B., Tu, T., Kong, F., Wen, J., & Xu, X. (2018). Revised phylogeny and historical
biogeography of the cosmopolitan aquatic plant genus Typha (Typhaceae). *Scientific*
Reports, *8*(1), Article 1. https://doi.org/10.1038/s41598-018-27279-3

Data Accessibility Statement

The updated genome assembly, raw reads supporting this study's findings, SNP information and chloroplast sequences produced in the present study will be deposited into online databases upon publication. Upon publication, related metadata and all the scripts used will be available (https://gitlab.com/WiDGeT_TrentU). We will provide unique sample identifier tags that can be matched to the deposited genetic data, metadata, and scripts.

Benefit–Sharing Statement

Benefits from this research accrue from sharing our data and results on public databases, as described above.

Author Contributions

Joanna R. Freeland, Aaron B. A. Shafer and Marcel E. Dorken conceived this study. Joanna R. Freeland and Polina A. Volkova collected and characterized the plant tissue samples. Tulsi Patel performed the lab work. Alberto Aleman executed the bioinformatics and wrote the manuscript. All authors contributed to the manuscript and approved this version.

Conflict of Interest

The authors declare no conflict of interest.

601   **Supporting Information**

602   **Tables and Figures**

603   **Table 1.** Mean Weir and Cockerham's pairwise genetic differentiation (F_{ST}) for every variable

604   site, mean genetic divergence (d_{xy}) measured in 10 Kbp windows, number of SNPs, and

605   diagnostic markers (SNPs with fixed opposite alleles) between three *Typha* spp.

Pairwise comparison	F_{ST}	d_{xy}	SNPs	Diagnostic SNPs
T. angustifolia – T. domingensis	0.25	0.28	10,358,977	33,436
T. angustifolia – T. latifolia	0.44	0.35	8,786,870	30,113
T. domingensis – T. latifolia	0.49	0.34	9,380,607	55,775

608
Figure 1. Genetic structure results of 12,177,703 nuclear SNPs obtained for three *Typha* spp.
(a) Sampling locations in this study. Coloured areas of the pie charts reflect the proportion of
samples from each taxon, and the number of samples is written top–right of each pie chart. (b)
Principal Component Analysis for PC1 and 2. Shapes represent individuals, colours represent
taxa, (c) Neighbour–joining tree. Tips represent individuals; colours represent clusters as a
and b, (d) ADMIXTURE analysis for *K = 3*. Vertical bars represent individuals, and the
probability of belonging to different taxa is coded with different colours. Arrows in b and d
indicate the location of potentially introgressed individuals. Colour and shape code as the box
in b: *T. angustifolia* (orange, triangles), *T. domingensis* (pink, circles), *T. latifolia* (green,
squares), and potentially introgressed individuals (cyan, diamonds). a-b plotted in ggplot2
(Wickham, 2016), c plotted with iTOL v6 (Letunic & Bork, 2021), and d plotted in R.

620
Figure 2. Chloroplast phylogeny of 140 *Typha* samples and four NCBI references, based on
96,591 bp. The tree was produced with *Sparganium natans* as the outgroup and drawn
without it. Coloured lines recreate the genetic structure results in Figure 1, i.e., *T. latifolia*
(green), *T. angustifolia* (orange), and *T. domingensis* (pink). Black circles indicate branch
support = 100% (Felsenstein bootstrap proportion). Arrows indicate the location of potentially
introgressed individuals (genetic structure results).