ESCRT disruption provides evidence against transsynaptic signaling functions for extracellular vesicles

Erica C. Dresselhaus[1], Kathryn P. Harris[2], Cassandra R. Blanchette[1], Kate Koles[1], Steven J. Del Signore[1], Matthew F. Pescosolido[1], Biljana Ermanoska[1], Mark Rozencwaig[1], Rebecca C. Soslowsky[1], Michael J. Parisi[4], Bryan A. Stewart[3], Timothy J. Mosca[4], Avital A. Rodal[1,*]

[1] Department of Biology, Brandeis University, Waltham, MA

[2] Office of the Vice-Principal, Research and Innovation, University of Toronto, Mississauga, Mississauga, Canada

[3] Department of Biology, University of Toronto Mississauga, Mississauga, Canada; Department of Cell and Systems Biology University of Toronto, Toronto, Canada

[4] Department of Neuroscience, Vickie and Jack Farber Institute of Neuroscience, Thomas Jefferson University, Bluemle Life Sciences Building, Philadelphia, PA

*Corresponding author:

Email: arodal@brandeis.edu

Key words: ESCRT, Tsg101, Hrs, Shrub, Vps4, Amyloid Precursor Protein, Synaptotagmin-4, Neuroglian, Evi, Wingless, Draper, *Drosophila*, extracellular vesicle, endosome, exosome, synapse

Abstract

Extracellular vesicles (EVs) are released by many cell types including neurons, carrying cargoes involved in signaling and disease. It is unclear whether EVs promote intercellular signaling or serve primarily to dispose of unwanted materials. We show that loss of multivesicular endosome-generating ESCRT (endosomal sorting complex required for transport) machinery disrupts release of EV cargoes from *Drosophila* motor neurons. Surprisingly, ESCRT depletion does not affect the signaling activities of the EV cargo Synaptotagmin-4 (Syt4) and disrupts only some signaling activities of the EV cargo Evenness Interrupted (Evi). Thus, these cargoes may not require intercellular transfer via EVs, and instead may be conventionally secreted or function cell autonomously in the neuron. We find that EVs are phagocytosed by glia and muscles, and that ESCRT disruption causes compensatory autophagy in presynaptic neurons, suggesting that EVs are one of several redundant mechanisms to remove cargoes from synapses. Our results suggest that synaptic EV release serves primarily as a proteostatic mechanism for certain cargoes.

Introduction

Neurons release extracellular vesicles (EVs) that can mediate intercellular communication, dispose of unwanted neuronal components, and propagate pathological factors in neurodegenerative disease (Budnik et al., 2016; Holm et al., 2018; Song et al., 2020). Many elegant functional studies of neuronal EVs involve their purification from donor cells and subsequent application to target cells for tests of biological activity e.g. (Gong et al., 2016; Vilcaes et al., 2021). These experiments demonstrate that EVs containing specific cargoes are sufficient to cause functional changes in the recipient cell, but do not rigorously show that traffic into EVs is necessary for the functions of cargoes *in vivo*. In the donor cell, EV cargoes are typically trafficked through the secretory system, plasma membrane, and endosomal network, where they might execute intracellular activities in the donor cell before

being released (van Niel et al., 2018). Therefore, to test the physiological functions of EVs *in vivo*, it will be essential to uncouple potential donor cell-autonomous from transcellular functions of these cargoes, using tools that specifically block EV release. Developing such tools will require a deeper understanding of how cargoes are packaged into EVs, and released in a spatially and temporally controlled fashion, especially within the complex morphology of neurons (Blanchette and Rodal, 2020).

Exosomes are a type of EV that arise when multivesicular endosomes (MVEs) fuse with the plasma membrane, releasing their intralumenal vesicles (ILVs) into the extracellular space. Spatial and temporal regulation of the machinery that controls formation of MVEs is therefore likely to be critical for exosome cargo selection, packaging, and release. MVEs can form via multiple nonexclusive mechanisms for budding of vesicles into the endosomal lumen (van Niel et al., 2018). One such pathway relies on Endosomal Sorting Complex Required for Traffic (ESCRT) proteins. In this pathway, ESCRT-0, -I, and -II components cluster cargoes, deform membranes, and then recruit ESCRT-III components, which form a helical polymer that drives fission of the ILV. The VPS4 ATPase remodels and finally disassembles the ESCRT-III filaments (Gruenberg, 2020; Vietri et al., 2020). The ESCRT-I component Tsg101 (Tumor susceptibility gene 101) is incorporated into and serves as a common marker for EVs, highlighting the link between ESCRT and EVs (van Niel et al., 2018). A neutral sphingomyelinase (nSMase)-mediated pathway may operate together with or in parallel to ESCRT to generate EVs by directly modifying lipids and altering their curvature, and indeed EV release of many neuronal cargoes is sensitive to nSMase depletion or inhibition (Asai et al., 2015; Dinkins et al., 2016; Goncalves et al., 2015; Guo et al., 2015; Men et al., 2019; Sackmann et al., 2019). The ESCRT machinery also has functions beyond MVE formation, including autophagosome closure and organelle repair, which are in turn involved in alternative modes of EV biogenesis (Arbo et al., 2020; Lefebvre et al., 2018; Leidal and Debnath, 2021). Further, ESCRT is involved in budding of EVs directly from the plasma membrane (van Niel et al., 2018). However, as there is evidence both for and against a role for ESCRT in EV biogenesis in different neuronal cell types (Cone et al., 2020; Coulter et al., 2018; Gong et al., 2016), it remains unclear whether organism-level physiological defects arising from ESCRT disruption (including ESCRT-linked human neurological disease) could arise from defects in EV traffic (Brugger et al., 2024; Rodger et al., 2020; Sadoul et al., 2018; Ugbode and West, 2021).

At the *Drosophila* larval neuromuscular junction (NMJ), EVs are released from presynaptic motor neurons into extrasynaptic space within the muscle membrane subsynaptic reticulum, and can also be taken up by muscles and glia (Fuentes-Medel et al., 2009; Koles et al., 2012). These EVs are likely to be exosomes, as cargoes are found in presynaptic MVEs, and depend on endosomal sorting machinery for their release and regulation (Blanchette et al., 2022; Koles et al., 2012; Korkut et al., 2009; Lauwers et al., 2018; Walsh et al., 2021). This system provides the powerful advantage of investigating endogenous or exogenous EV cargoes with known physiological functions in their normal tissue and developmental context. EV cargoes characterized to date at the *Drosophila* NMJ include Synaptotagmin-4 (Syt4, which mediates functional and structural plasticity), Amyloid Precursor Protein (APP, a signaling protein involved in Alzheimer's Disease), Evenness Interrupted/Wntless/Sprinter (Evi, which carries Wnt/Wingless (Wg) to regulate synaptic development and plasticity), and Neuroglian (Nrg, a cell adhesion molecule) (Koles et al., 2012; Korkut et al., 2013; Walsh et al., 2021). Mutants in membrane trafficking machinery (e.g. *evi*, *rab11 (*a recycling endosome GTPase))*, and *nwk* (a component of the endocytic machinery) cause reduced levels of cargo in EVs and show defects in the physiological activities of EV cargoes (Blanchette et al., 2022; Korkut et al., 2009; Korkut et al., 2013), leading to the hypothesis that trans-synaptic transfer of these cargoes into the postsynaptic muscle is required for their

signaling functions (Budnik et al., 2016). However, we and others have shown that these
mutants also have a dramatic local presynaptic decrease in cargo levels, making it difficult to
rule the donor neuron out as their site of action (Ashley et al., 2018; Blanchette et al., 2022;
Koles et al., 2012; Korkut et al., 2009; Korkut et al., 2013; Walsh et al., 2021). Here we show
that disruption of the ESCRT machinery can cause a specific loss of EV release without
strongly depleting presynaptic cargo levels, and use this system to test whether EV release
is required for cargo signaling functions.

Results

ESCRT machinery promotes EV release from synapses

To determine if the ESCRT pathway is involved in EV release at the *Drosophila* NMJ
synapse, we first used GAL4/UAS-driven RNAi to knock down the ESCRT-I component
Tsg101 (Tumor susceptibility gene 101) specifically in neurons (Tsg101^{KD}). We then used our
previously established methods to measure the levels of the endogenously tagged EV cargo
Syt4-GFP, both in the donor presynaptic compartment and in neuron-derived EVs in the
adjacent postsynaptic cleft and muscle (Walsh et al., 2021). Neuronal knockdown of Tsg101
(Tsg101^{KD}) led to accumulation of Syt4-GFP in the presynaptic compartment, together with a
striking loss of postsynaptic Syt4-GFP EVs (**Fig 1A, E**). We next tested the effects of
Tsg101^{KD} on three other known EV cargoes: neuronal UAS-driven Evi-GFP **(1B, F)** or human
APP-GFP **(1C, G)**, and endogenous Nrg **(1D, H)** (Korkut et al., 2009; Korkut et al., 2013;
Walsh et al., 2021). For all three cargoes, we observed a similar phenotype to Syt4-GFP:
presynaptic redistribution in large structures (accumulating to particularly high levels for
Syt4-GFP and Evi-GFP), together with loss of postsynaptic EV signal. Similarly, neuronal
UAS-driven Tsp42Ej/Sunglasses (a *Drosophila* tetraspanin EV marker (Walsh et al., 2021))
was strongly depleted from the postsynaptic region upon Tsg101^{KD} (**Fig. S1A**).

We then used super-resolution STED microscopy to measure the size and number of
presynaptically-derived EV cargoes in the 3 µm region surrounding the presynaptic terminal.
The density of postsynaptic puncta was strongly reduced at Tsg101^{KD} synapses (**Fig. S1B-**
E), and was not significantly different from background signal. Further, the diameters of Nrg
and APP-GFP extraneuronal puncta were ~125 nm, consistent with the size of MVE-derived
exosomes (Welsh et al., 2024)). Given their size, the observation that cargoes accumulate in
internal structures upon Tsg101^{KD}, and previous observations of MVE fusion with the plasma
membrane at this synapse (Koles et al., 2012; Korkut et al., 2009; Lauwers et al., 2018), it is
most likely that NMJ EVs are exosomes derived from intracellular MVEs, rather than
budding directly from the plasma membrane. Thus, multiple EV cargoes, either
endogenously or exogenously expressed, require the ESCRT-I component Tsg101 for
release in neuronally-derived EVs.

In addition to its functions in MVE biogenesis, Tsg101 also plays roles in numerous cellular
processes including membrane repair, lipid transfer, neurite pruning, and autophagy, each
depending on a specific subset of other ESCRT machinery (Vietri et al., 2020). We therefore
tested if EV release depends on other canonical ESCRT components. Hrs (Hepatocyte
growth factor receptor substrate) is a component of the ESCRT-0 complex and is required to
cluster EV cargo on the delimiting membrane of the endosome (Vietri et al., 2020). Similar to
Tsg101^{KD}, *Hrs* loss-of-function mutants caused a strong decrease in postsynaptic Syt4-GFP,
Evi-GFP, and Nrg, though interestingly their presynaptic levels were also partially depleted,
unlike the Tsg101^{KD} condition (**Fig 2A-F**). Notably, direct comparison showed that *Hrs*
mutants exhibited a postsynaptic decrease in Syt4-GFP nearly as severe as *nwk* mutants,
but with a comparatively modest decrease in presynaptic Syt4-GFP (**Fig 2A,D**).

Next, we tested ESCRT-III, which forms the polymer involved in constriction and scission of the ILV neck. The *Drosophila* genome encodes several ESCRT-III proteins, of which *shrub* is homologous to mammalian CHMP4B. Shrub is likely to play an important role at synapses, since its loss leads to defects in NMJ morphogenesis and ILV formation (Sweeney et al., 2006). Pan-neuronal RNAi of *shrub* (Shrub[KD]) caused a dramatic loss of postsynaptic Syt4-GFP and Nrg signals (**Fig. 2G-I**). Finally, we examined the role of Vps4, which catalyzes remodeling and disassembly of the ESCRT-III polymer, finalizing the formation of the ILV. Pan-neuronal expression of a dominant negative Vps4 fragment (Vps4[DN], (Rodahl et al., 2009)) strongly reduced postsynaptic levels of both Syt4-GFP and Nrg, and increased their presynaptic levels (**Fig. 2J-L**). Together, these results demonstrate that multiple components of the ESCRT pathway are required for release of EV cargoes at neuronal synapses, with variable effects on presynaptic accumulation of these cargoes.

Loss of Tsg101 or Hrs uncouples autophagic and EV functions of ESCRT machinery

To explore the nature of the presynaptic accumulations of EV cargoes at Tsg101[KD] NMJs, we examined their co-localization with early (Rab5) and recycling (Rab11) endosomes, which drive an endosome-to-plasma membrane recycling flux that supplies the EV biogenesis pathway at this synapse (Korkut et al., 2009; Walsh et al., 2021). We also examined cargo co-localization with late endosomes (Rab7), which play less important roles in NMJ EV traffic (Walsh et al., 2021). We found that EV cargoes exhibited increased co-localization with all these endosomal markers at Tsg101[KD] synapses, in what appeared to be multi-endosome clusters (**Fig 3A, S2A-C**). These results argue against formation of a single type of arrested MVE such as the canonical Class E compartment in ESCRT-deficient yeast and mammalian cells (Doyotte et al., 2005; Raymond et al., 1992) and instead suggest a more global defect in endosome maturation or turnover. To test this hypothesis, we measured the overall mean intensity as well as puncta number and size for Rab5, Rab7, and Rab11 upon Tsg101 knockdown. We saw no changes in the total intensity or puncta parameters for Rab7 (**Fig S2D-F**). However, for both Rab5 and Rab11, we observed a significant increase in Rab puncta intensity, mean intensity over the whole NMJ, and an increase in puncta size, with no change (Rab11) or a slight decrease (Rab5) in the number of puncta (**Fig S2D-F)**. These results suggest defects in early and recycling endosome maturation and/or turnover upon loss of ESCRT function.

We next examined Tsg101[KD] NMJs using transmission electron microscopy (TEM). Tsg101[KD] boutons retained typical mitochondria, synaptic vesicles and active zone "T-bars", and were surrounded by an apparently normal subsynaptic reticulum, representing the infolded postsynaptic muscle membrane. However, within Tsg101[KD] boutons we observed striking clusters of double membrane-surrounded or electron-dense structures, typical of autophagic vacuoles at various stages of maturation (Klionsky et al., 2021), including those with unclosed phagophores (**Fig. 3B**; three or more autophagic vacuoles were observed in 58.9% of mutant boutons (n=56) compared to 2.5% of control boutons (n=40), p<0.001). Given that secretion of autophagosomal contents has been described as an EV-generating mechanism (Buratta et al., 2020), we next tested whether core autophagy machinery might play a role in EV release at the *Drosophila* NMJ, and could therefore be linked to the Tsg101[KD] EV phenotype. Atg1 is a kinase that is required for the initiation of phagophore assembly, and acts as a scaffold for recruitment of subsequent proteins, while Atg2 is required for phospholipid transfer to the phagophore (Nakatogawa, 2020). We observed a modest but significant decrease in both pre- and post-synaptic levels of the EV marker Nrg upon disruption of autophagy by RNAi-mediated knockdown of Atg1 (knockdown validated in **Fig. S2G**, Nrg results in **Fig. 3C-D**), as well as by loss-of-function *Atg2* mutations (**Fig. S2H-I**). Notably, these mutants did not recapitulate the ESCRT mutant phenotype of dramatic

depletion of postsynaptic EVs and presynaptically trapped cargoes. These results indicate that autophagic machinery does not play a major role in EV biogenesis or release at the NMJ, and that the autophagic defects at Tsg101^{KD} NMJs are likely separable from its roles in EV release.

To further explore these autophagic defects, we next used the reporter GFP-mCherry-Atg8/LC3 to assess autophagic flux in Tsg101^{KD} neurons. Under normal circumstances, the GFP moiety in this reporter is quenched when autophagosomes fuse with acidic endosomes and lysosomes, while mCherry retains its fluorescence. By contrast, defects in autophagic flux lead to accumulation of structures with both GFP and mCherry fluorescence (Klionsky et al., 2021). We examined this flux in motor neuron cell bodies, where mature autolysosomes are predicted to accumulate (Sidibe et al., 2022). In wild-type animals we observed mCherry-positive/GFP-negative puncta reflecting mature autolysosomes. By contrast we observed an increased volume of intense puncta in the cell bodies of Tsg101^{KD} motor neurons, most of which were labeled by both mCherry and GFP (**Fig. 3E-F, S2J**). These data suggest that Tsg101^{KD} causes a defect in autophagic flux.

Since autophagy is normally rare at wild-type *Drosophila* NMJ synapses (Soukup et al., 2016), we hypothesized that ESCRT mutants might activate a compensatory "endosomophagy" or "simaphagy" pathway to degrade ESCRT-deficient endosomes (Migliano et al., 2023; Millarte et al., 2022). However, since Tsg101 is also required for phagophore closure (Takahashi et al., 2018), this process is likely unable to dispose of defective endosomes upon Tsg101 knockdown. By contrast, ESCRT-0/Hrs is not required for autophagy in some cell types, such as *Drosophila* fat body (Rusten et al., 2007). To test if this also applies to motor neurons, we examined GFP-mCherry-Atg8 in *Hrs* mutant motor neuron cell bodies, and found mCherry-positive, GFP-negative structures, similar to controls (**Fig. 3E, G, S3J**), suggesting that Hrs is not required for autophagic flux in motor neurons. Interestingly, we did observe an increase in the area covered by puncta, indicating that autophagy is induced in *Hrs* mutants (**Fig. 3G**). Overall, *Tsg101* and *Hrs* have different autophagy phenotypes but similar EV release defects, and canonical autophagy mutants do not phenocopy ESCRT mutants in trapping EV cargoes presynaptically. Together, these results suggest that autophagy and EV traffic are separable functions of ESCRT at the synapse, and that a compensatory (and Tsg101-dependent) autophagy mechanism might be activated to remove defective endosomes in *Hrs* mutants.

Finally, we further explored whether accumulation of EV cargoes in arrested structures was local to the synapse or occurring throughout the neuron. First, we examined Syt4-GFP levels in motor neuron cell bodies and axons after Tsg101 knockdown, and found that Syt4-GFP accumulated significantly at both locations (**Fig. 4A-B**). To ask whether the presynaptic accumulations could be due to faster anterograde or slowed retrograde transport of EV cargo-containing compartments, we next conducted live imaging and kymograph analysis of motor neuron-driven APP-GFP, as well as a mitochondrial marker. We found that Tsg101 knockdown led to a large increase in the number of stationary APP-GFP puncta in axons without affecting the number of compartments undergoing retrograde or anterograde transport (**Fig 4C-D**), though we observed a small decrease in the retrograde transport rate (**Fig 4E**). By contrast, we did not observe an increase in the steady state intensity of the mitochondrial marker or see any changes in its transport behavior (**Fig S3A-D**), suggesting that axonal accumulations are specific to EV cargo. Thus, loss of *tsg101* leads to accumulation of stationary EV cargo-containing compartments throughout the neuron, without affecting the transport rates of moving cargoes, suggesting that altered axonal transport kinetics do not underlie synaptic accumulation.

evi and *wg* signaling are not correlated with EV release

Specific depletion of cargo in postsynaptic EVs (but not the donor presynaptic terminal) upon ESCRT disruption provided us with a valuable tool to determine if these cargoes require trans-synaptic transfer for their known synaptic functions. Neuron-derived Wg provides anterograde (to the muscle) and autocrine (to the neuron) signals, promoting NMJ growth, active zone development, and assembly of the postsynaptic apparatus (Miech et al., 2008; Packard et al., 2002). Evi is a multipass transmembrane protein that serves as a carrier for Wg through the secretory system, ultimately leading to Wg release from the cell, either by conventional exocytosis or via EVs (Das et al., 2012). At the NMJ, Evi co-transports with Wg into EVs, and *evi* mutants phenocopy *wg* signaling defects, providing support for the hypothesis that Evi/Wg EVs are required for Wg signaling (Koles et al., 2012; Korkut et al., 2009). However, since Evi is broadly required for many steps of Wg traffic, *evi* mutants trap Wg in the somatodendritic compartment and prevent its transport into presynaptic terminals (Korkut et al., 2009). Therefore, Wg signaling defects in *evi* mutants may be due to generalized loss of Wg secretion rather than specific loss of its trans-synaptic transfer. *Wg* or *evi* mutants exhibit dramatic reductions in bouton number, together with the appearance of immature boutons with abnormal or missing active zones, fewer mitochondria, aberrant swellings or pockets in the postsynaptic region opposing active zones, and missing areas of PSD95/Discs-Large (DLG)-positive postsynaptic subsynaptic reticulum (Korkut et al., 2009; Packard et al., 2002). We found that in *evi* mutants, the number of synaptic boutons and the number of active zones (marked by ELKS/CAST/ Bruchpilot (BRP)) were both significantly reduced compared to controls, and the postsynaptic scaffolding molecule DLG frequently exhibited a "feathery" appearance, suggesting defects in postsynaptic assembly (**Fig. 5A-E**). By contrast, we found that bouton and active zone numbers at Tsg101[KD] NMJs (which have presynaptic Evi but strongly diminished Evi EVs (**Fig. 1B**)) were not significantly different from controls. Further, active zones in Tsg101[KD] appeared morphologically normal by TEM (**Fig. 3B**). We did not observe a significant frequency of "feathery" DLG distribution in control or Tsg101[KD] larvae (2.3% of control NMJs (n=87) and 5.5% of Tsg101[KD] NMJs (n=87), compared to 51.7% of *evi*[2] mutant NMJs (n=91), p=0.27 for control versus Tsg101[KD]). We also did not observe significant differences between control and Tsg101[KD] NMJs in the appearance of subsynaptic reticulum by EM (**Fig. 3B**) These results indicate that some neuronal Evi and Wg functions are unexpectedly maintained despite loss of detectable postsynaptic Evi EVs upon Tsg101[KD].

In addition to an overall decrease in the number of synaptic boutons, both *Wg* and *evi*[2] mutants show increased numbers of developmentally arrested or "ghost" boutons that feature presynaptic markers such as α-HRP antigens, but lack a postsynaptic apparatus defined by DLG (Korkut et al., 2009). We found that in *evi*[2] mutants, these ghost boutons are more prevalent in anterior segments of the larvae, where overall synaptic growth is more exuberant. Similarly, Tsg101[KD] animals exhibited a significant increase in ghost boutons in abdominal segment A2 (but not in A3), partially phenocopying the *evi*[2] mutant (**Fig, 5 D,E**). In sharp contrast, *Hrs* mutants did not exhibit a significant change in ghost bouton number despite having a similar decrease in postsynaptic Evi-GFP to Tsg101[KD] (**Fig, 5 D,E**). These results suggest that Evi release in EVs and *wg*-related phenotypes can be uncoupled.

To further explore Wg signaling in ESCRT mutants, we directly measured the output of this pathway. In *Drosophila* muscles, Wg does not signal via the conventional β-catenin pathway. Instead, neuronally-derived Wg activates cleavage of its receptor Fz2, resulting in translocation of a Fz2 C-terminal fragment into muscle nuclei (Mathew et al., 2005; Mosca and Schwarz, 2010). Using an antibody specific to the Fz2 C-terminus, we measured the number of nuclear Fz2-C puncta (**Fig. 5F-G**). *Hrs* mutants showed a similar number of

puncta compared to controls. By contrast neuronal knockdown of Tsg101 caused a dramatic loss of Fz2-C puncta, consistent with our findings for ghost boutons. Given the similarly strong loss of EVs in *Hrs* mutants and Tsg101[KD], these results indicate that EVs are not required for transynaptic signaling by Wg, and suggest that a separable membrane trafficking pathway for Wg secretion is defective only in the Tsg101[KD] condition.

ESCRT loss does not recapitulate *syt4* phenotypes in activity-dependent structural or functional plasticity

We next explored the functions of the EV cargo Syt4, which is required for activity-dependent structural and functional plasticity at the NMJ (Barber et al., 2009; Korkut et al., 2013; Piccioli and Littleton, 2014; Yoshihara et al., 2005). Endogenous Syt4 is thought to be generated only by the presynaptic motor neuron, based on the absence of Syt4 transcript in muscle preparations, and the finding that presynaptic RNAi of Syt4 depletes both presynaptic and postsynaptic signals (Korkut et al., 2013). We independently verified that all the Syt4 at the NMJ was derived from the neuron, using a strain in which the endogenous Syt4 locus is tagged at its 3' end with a switchable TagRFP-T tag, which could be converted to GFP in the genome via tissue-specific GAL4/UAS expression of the Rippase recombinase (Koles et al., 2015; Walsh et al., 2021) (**Fig. S4A**). Conversion of the tag only in neurons resulted in a bright Syt4-GFP signal both presynaptically and postsynaptically, together with disappearance of the TagRFP-T signal. However, conversion of the tag in muscles did not result in any GFP signal, and the TagRFP-T signal remained intact (**Fig. S4B**). These results indicate that Syt4 is only expressed in neurons, and support the previous conclusion that the postsynaptic signal is derived from a presynaptically-expressed product (Korkut et al., 2013).

Membrane trafficking mutants such as *rab11* and *nwk* deplete Syt4 from presynaptic terminals, secondarily reducing its traffic into EVs, and phenocopy *syt4* null mutant plasticity phenotypes (Blanchette et al., 2022; Korkut et al., 2013; Walsh et al., 2021). This does not provide conclusive evidence that signaling by Syt4 explicitly requires its transfer via EVs, since Syt4 is missing from both the donor and recipient compartment and could therefore be signaling in the presynaptic cell. Given that loss of *Hrs* and *tsg101* leads to a similar postsynaptic decrease in Syt4 to *nwk* mutants but without a strong presynaptic decrease, these mutants present an opportunity to challenge the hypothesis that Syt4 must transfer via EVs to exert its functions. We first tested the effect of Tsg101[KD], which depletes the majority of EVs without diminishing presynaptic Syt4 (**Fig. 1A, E**), on Syt4- dependent structural plasticity. In this paradigm, spaced high potassium stimulation promotes acute formation of nascent ghost boutons (Ataman et al., 2006; Korkut et al., 2013; Piccioli and Littleton, 2014). These are likely transient structures, and thus are not directly comparable to developmentally arrested ghost boutons such as those found in *evi* mutants (Fernandes et al., 2023). However, to avoid the confounding presence of these immature boutons, we explored the activity-dependent synaptic growth paradigm on muscle 4, where the Tsg101[KD] animals do not have significantly more ghost boutons than controls under baseline conditions. Unexpectedly, Tsg101[KD] animals behaved similarly to controls, and exhibited a significant increase in ghost boutons following high K+ spaced stimulation compared to mock stimulation (**Fig. 6A-B**), suggesting that Syt4 function is preserved in these synapses despite depletion of EV Syt4. We were surprised by these results and contacted another laboratory (KPH, BAS) to replicate this experiment independently at muscle 6/7 in segments A3 and A4, and again readily observed activity-dependent ghost bouton formation (**Fig. S4C-D**). KPH next tested the effect of Tsg101[KD] on Syt4-dependent functional plasticity. In this paradigm, stimulation with 4x100 Hz pulses causes a Syt4-dependent increase in the frequency of miniature excitatory junction potentials (mEJPs), in a phenomenon termed High Frequency-Induced Miniature Release (HFMR) (Korkut et al., 2013; Yoshihara et al., 2005).

Tsg101^{KD} animals exhibited similar HFMR to wild type controls, indicating that Syt4 function
was not disrupted (**Fig. 6C-D**). *Hrs* mutants, despite being very sickly, also exhibited similar
HFMR to wild type controls, in sharp contrast to *syt4* null animals which did not exhibit any
HFMR (**Fig. 6E-F**). Taken together, our results show that Syt4-dependent structural and
functional plasticity at the larval NMJ can occur despite dramatic depletion of EVs containing
Syt4.

Syt4 is not detectable in the muscle cytoplasm and is taken up by phagocytosis

If trans-synaptic transfer of Syt4 in EVs serves a calcium-responsive signaling function in the
muscle, one would expect to find neuronally-derived Syt4 in the muscle cytoplasm.
Therefore, we tested whether neuronally-derived Syt4-GFP (for which the GFP moiety is
topologically maintained on the cytoplasmic side of membranes in both donor and recipient
cells) could be found in the muscle cytoplasm. Using the GAL4/UAS system, we expressed
a proteasome-targeted anti-GFP nanobody (deGradFP (Caussinus et al., 2011), **Fig. 7A**)
only in neurons or only in muscles. We observed strong depletion of Syt4-GFP fluorescence
upon presynaptic deGradFP expression, including a reduction in Syt4 postsynaptic puncta
intensity and number, consistent with the presynaptic source of EV Syt4-GFP protein(**Fig.
7B, D, F**). This result also demonstrates the effectiveness of deGradFP in depleting Syt4-
GFP. However, we did not observe any difference in either presynaptic or postsynaptic Syt4-
GFP levels or puncta number upon deGradFP expression in the muscle (**Fig. 7C, E, F**),
though deGradFP could efficiently deplete DLG as a control postsynaptic protein (**Fig. S5A**).
These results suggest that the majority of postsynaptic Syt4 is not exposed to the muscle
cytoplasm.

Conversely, if EVs serve primarily as a proteostatic mechanism to shed neuronal Syt4 for
subsequent uptake and degradation in recipient cells, then Syt4 would not need to be
exposed to the muscle cytoplasm, as it could be taken up by phagocytosis in double
membrane compartments for degradation via fusion with recipient cell lysosomes. Indeed, α-
HRP positive neuronal "debris" is taken up via the phagocytic receptor Draper (Fuentes-
Medel et al., 2009). This debris was not previously directly linked to EVs, though it co-
localizes strongly with EV cargo (Walsh et al., 2021). We used cell type-specific Draper
RNAi to test directly if known EV cargoes are cleared by Draper-dependent phagocytosis.
First, we established that Draper is expressed in neurons, glia, and muscles at the NMJ.
RNAi of Draper in each of these tissues depleted a subset of Draper immunostaining at the
NMJ and axon, indicating that Draper is expressed in each of these cell types (**Fig. S5B-C**).
We then quantified Syt4 levels in Draper RNAi larvae. Depletion of Draper in either muscles,
glia, or neurons led to an increase in postsynaptic Syt4-GFP, indicating that the normal
destination of Syt4 following release from the neuron is phagocytosis by multiple adjacent
cell types (**Fig. 7G-H**). Interestingly, we found that EV cargoes also accumulated
presynaptically upon Draper knockdown in glia, neurons, or muscles – this could be due to
presynaptic reuptake of EVs by bulk endocytosis when they cannot be cleared by
phagocytosis. Overall, these results show that Syt4 EVs are cleared by phagocytosis, similar
to previously characterized α-HRP-labeled debris, but are not transported at detectable
levels into the cytoplasm of muscle cells.

Discussion

Here we show that the ESCRT pathway is required for EV cargo packaging at the
Drosophila larval NMJ, and that these EVs are likely MVE-derived exosomes. We found that
ESCRT depletion caused presynaptic accumulation of cargoes, defects in their axonal
transport, and a dramatic loss of trans-synaptic transfer in EVs. Surprisingly, we found that
this trans-synaptic transfer is not required for several physiological functions of EV cargoes

Evi and Syt4. Further, neuronally-derived Syt4 is taken up by phagocytosis and could not be detected in the muscle cytoplasm, consistent with findings from HeLa cells that the majority of EV cargoes remain in the endosomal system of the recipient cell (O'Brien et al., 2022). Our results suggest that neuronal EV release for these cargoes at this developmental stage serves primarily proteostatic and not signaling functions.

Functions of ESCRT in MVE biogenesis and EV release at synapses

We found that ESCRT is required for EV generation and release at the *Drosophila* larval NMJ. ESCRT components are also required for EV/exosome cargo release from primary neurons in culture (Gong et al., 2016) and Purkinje neurons *in vivo* (Coulter et al., 2018), but not for EV release of pathogenic APP variants or Evi from cell lines (Beckett et al., 2013; Cone et al., 2020), underscoring the importance of studying membrane traffic in *bona fide* neurons. Further, we found that upon ESCRT depletion, cargoes accumulate in intracellular compartments, suggesting that this population of NMJ EVs are MVE-derived exosomes rather than plasma membrane-derived microvesicles. This is consistent with the requirement for endosomal sorting machinery, such as retromer, in their regulation (Walsh et al., 2021).

One major open question is whether EV-precursor MVEs are generated on-demand in response to local cues at presynaptic terminals, or if they arise in response to global cues and are transported to synapses from other regions of the neuron. Answering this question will require tools to visualize the timecourse of MVE biogenesis in neurons, as have been developed in cultured non-neuronal cells (Wenzel et al., 2018). In addition, future methods (e.g. optogenetic) for acute and localized inhibition of ESCRT will reveal whether arrested structures first appear locally at the synapse and are only later transported into axons and cell bodies, and/or if they are generated far from the site of release at the synapse. These experiments will be critical for understanding when and where local or global signaling events impinge on EV biogenesis. Interestingly, activity-dependent delivery of Hrs to presynaptic terminals is critical for proteostasis of synaptic vesicle proteins (Birdsall et al., 2022; Sheehan et al., 2016). If MVEs are generated on-demand at synapses, Hrs transport could similarly underlie the activity-dependence of EV release, which has been reported in many (but not all) neuronal experimental systems, and remains poorly understood (Ataman et al., 2008; Faure et al., 2006; Lachenal et al., 2011; Lee et al., 2018; Vilcaes et al., 2021).

Other synaptic functions of ESCRT

ESCRT is best-known for its functions in MVE biogenesis, but has many other potential synaptic roles including in autophagy, lipid transfer and membrane repair (Vietri et al., 2020). Tsg101 is involved in lipid transfer to mitochondria (Wang et al., 2021), but we did not detect obvious defects in mitochondria in motor neuron axons, as were seen in *Tsg101*-mutant *Drosophila* adult wing sensory neurons (Lin et al., 2021). Our results also show that the function of ESCRT in EV release is likely separate from its roles in autophagy, since several canonical autophagy mutants do not phenocopy presynaptic trapping of EV cargoes as seen upon ESCRT depletion, and *Hrs* mutants exhibit EV but not autophagic flux defects. Interestingly, we found that *atg* mutants led to a moderate presynaptic and postsynaptic reduction in levels of the EV cargo Nrg. This raises the possibility that other degradative pathways are upregulated at synapses when autophagy is blocked. Thus, while ESCRT has many cellular activities, our experiments separate these functions, and specifically narrow down its role in neuronal EV release.

Many organelles are selectively targeted for macroautophagy via compartment-specific receptors (Lamark and Johansen, 2021), but such a process has not been specifically described for neuronal endosomes/MVEs. Our data suggest that synapses use a

proteostatic mechanism called endosomophagy or simaphagy that has been previously
observed in cell culture (Migliano et al., 2023; Millarte et al., 2022; Wang et al., 2022; Zellner
et al., 2021), adding to the numerous intersections between endolysosomal traffic and
autophagy in neurons (Boecker and Holzbaur, 2019). We found that autophagy is induced in
ESCRT mutant synapses, presumably to dispose of aberrant endosomes, with different
outcomes in Tsg101^{KD} versus *Hrs* mutants: Tsg101^{KD} led to aberrant autophagic vacuoles
and reduced autophagic flux, perhaps due to a secondary role for ESCRT-1/Tsg101 in
phagophore closure or another step of autophagy (Takahashi et al., 2018). By contrast we
found that *Hrs* mutants do not show these structures, either by light microscopy of the
autophagic flux reporter GFP-mCherry-ATG8, or in previously published TEM of the NMJ
(Lloyd et al., 2002). Instead, *Hrs* mutants exhibit induction of autophagy but normal
autophagic flux in motor neurons, together with a moderate reduction in EV cargo levels.
Together, these results suggest that aberrant EV-cargo-containing MVEs may be removed
from *Hrs* mutant synapses by a compensatory, Tsg101-dependent autophagy pathway.

Implications for the signaling roles of EVs

The majority of functional studies of EVs involve isolating EV subpopulations (at various
degrees of homogeneity) from cell culture supernatants, applying them to target cells or
tissues, and assessing their biological effects (Welsh et al., 2024). Additional mechanistic
insight has been obtained by eliminating specific cargo molecules from the donor cells
before EV isolation, to determine if these molecules are required for EV bioactivity. While
these approaches are very useful for determining therapeutic uses for EVs, they have
several major limitations for understanding their normal functions *in vivo*. First, it is difficult to
determine the concentration of EVs that a target cell would normally encounter, in order to
design a physiologically relevant experiment. Second, while these types of experiments
inform what EVs can do, they do not show that EV transfer is necessary for that signaling
function *in vivo*. Removing the signaling cargo from the donor cell also does not show the
necessity of EV transfer for biological functions, since the cargo could be acting cell
autonomously in the donor cell, or could signal to a neighboring cell by another trafficking
route. Indeed, previous studies at the *Drosophila* larval NMJ, which has been an important
model system for the *in vivo* functions of EV traffic, have conducted tests for EV cargo
activity in *evi* or *rab11* mutants, though we and others have shown that this results in
depletion of cargo from the presynaptic donor cell in addition to loss of EVs (Ashley et al.,
2018; Blanchette et al., 2022; Koles et al., 2012; Korkut et al., 2009; Korkut et al., 2013;
Walsh et al., 2021). Ultimately, determining if transfer of a cargo in EVs is necessary for its
signaling function requires blocking EV transfer specifically, which we were able to achieve
at ESCRT-depleted synapses.

Neuronally derived Wg is required and sufficient for synaptic growth, and functions together
with glia-derived Wg to organize postsynaptic glutamate receptor fields (Kerr et al., 2014;
Korkut et al., 2009; Miech et al., 2008; Packard et al., 2002). Therefore, if transsynaptic
transfer of Evi and Wg in EVs was required for Wg signaling, we would expect to see a
reduction in synaptic growth at ESCRT-depleted synapses, as well as disruptions in
postsynaptic development and organization. However, we observed no significant change in
bouton or active zone number relative to controls upon either Tsg101 or Hrs depletion,
indicating that they do not phenocopy either *evi* or *wg* mutants, and that at least some Wg
activity is maintained even when Evi-GFP transfer is strongly inhibited. *wg*-phenocopying
defects in subsynaptic reticulum were also not observed by electron microscopy in
CHMPIIB^{intron5} (West et al., 2015) or *Hrs*-mutant synapses (Lloyd et al., 2002), or in our data
from ESCRT mutants. Similarly, Hsp90 mutants attenuate Evi EV release by disrupting
MVE-plasma membrane fusion, but do not result in disruption of active zone or subsynaptic

reticulum structure (Lauwers et al., 2018). Therefore, loss of EVs does not phenocopy many *wg* or *evi* defects, suggesting that the primary function of Evi is likely to traffic Wg to the presynaptic terminal and maintain its levels there, rather than specifically to mediate its release via EVs.

Importantly, it is likely that Hsp90 and ESCRT mutant synapses do secrete Wg, albeit by a non-EV mechanism (Beckett et al., 2013; Won and Cho, 2021). Interestingly, we found that Tsg101[KD] leads to loss of Fz2-C nuclear import and an increase in baseline ghost boutons, consistent with some defects in Wg signaling. Our finding that Tsg101[KD] causes additional membrane trafficking defects (e.g. autophagy) compared to *Hrs* suggests that non-EV modes of Wg release may be disrupted in this mutant. Likely possibilities include conventional secretion or secretory autophagy (Beckett et al., 2013; Chang et al., 2024; Won and Cho, 2021). However, since *Hrs* mutants do not show any deficit in Fz2-C nuclear import or ghost boutons despite exhibiting a similar loss of postsynaptically transferred Evi-GFP to Tsg101[KD], we conclude that Wg signaling and EV release are separable functions.

Syt4 protein is thought to act in the postsynaptic muscle (Adolfsen et al., 2004; Barber et al., 2009; Harris et al., 2016), but its endogenous transcript is not expressed in this tissue, leading to the prevailing model of transynaptic transfer from the presynaptic neuron in EVs (Korkut et al., 2013). Our results show that transynaptic transfer in EV can be blocked without affecting the signaling activities of Syt4, and that the majority of postsynaptic Syt4 is not exposed to the muscle cytoplasm. The main evidence for a muscle requirement for Syt4 is that re-expression of Syt4 using muscle-specific GAL4 drivers is sufficient to rescue structural and functional plasticity defects of the *syt4* null mutant (Korkut et al., 2013; Piccioli and Littleton, 2014; Yoshihara et al., 2005). This is difficult to reconcile with our findings that Tsg101[KD] and *Hrs* animals lack detectable postsynaptic Syt4, but do not phenocopy *syt4* mutants. There are several possible explanations for this conundrum. First, we cannot completely rule out the possibility that small amounts of residual Syt4 EVs are sufficient to drive a transynaptic signal. This is unlikely, since *nwk* and *rab11* mutants also have trace amounts of Syt4 postsynaptically, and do strongly phenocopy the *syt4* null mutant (presumably since they also deplete Syt4 from the presynaptic compartment) (Blanchette et al., 2022; Korkut et al., 2013; Walsh et al., 2021). Therefore, trace Syt4 is insufficient for signaling. Second, Syt4 could be transferred by a non-EV pathway such as conventional secretion, tunneling nanotubes, or cytonemes, and be distributed diffusely in the muscle such that we cannot detect its presence or degradation by cytoplasmic deGradFP (Dagar and Subramaniam, 2023; Daly et al., 2022)). Third, it is possible that the muscle GAL4 drivers and UAS lines used in these previous rescue studies have some leaky expression in the neuron. Fourth, ectopically muscle-expressed Syt4 might have a neomorphic function in the muscle that bypasses the loss of neuronal Syt4, or else it could be retrogradely transported to the neuron. Indeed, muscle-expressed Syt4 is localized in close apposition to the presynaptic membrane (Harris et al., 2016).

Conclusions

Why are Evi and Syt4 trafficked into EVs, if not for a signaling function? Local EV release could serve as a proteostatic mechanism for synapse-specific control of signaling cargo levels, in cooperation with other degradative mechanisms. Our data show that EVs are taken up by glial, muscle, and neuronal phagocytosis, and that cargoes are protected from the muscle cytoplasm. Indeed, the amount of cargo loaded into EVs could be tuned by regulating endosomal sorting via retromer (Walsh et al., 2021), or by controlling the rate of endocytic flux into the Rab11-dependent recycling pathway (Blanchette et al., 2022). Our results also show that EVs are one of several intersecting and complementary mechanisms

for synaptic proteostasis of membrane-bound cargoes; when EV release is reduced, we found that compensatory autophagy pathways are upregulated to degrade unwanted endosomal components. Further, endosomes that are not competent for EV biogenesis can be targeted for dynein-mediated retrograde transport (Blanchette et al., 2022), perhaps to bring cargoes to the cell body where lysosomal degradation is more active (Ferguson, 2018). Through these mechanisms, neurons might achieve local control of synaptogenic or plasticity-inducing signaling pathways, in a much more rapid and spatially controlled fashion than transcriptional or translational regulation.

Importantly, our results do not rule out signaling functions for Syt4 or Evi/Wg EVs in other contexts or neuronal cell types, or for other EV cargoes. For example, ESCRT disruption suppresses the pathological functions of APP in *Drosophila*, perhaps due to its reduced propagation in EVs (Zhuang et al., 2023). Indeed, extensive evidence supports signaling and pathological functions for neuronal EVs in multiple contexts (Gassama and Favereaux, 2021; Lizarraga-Valderrama and Sheridan, 2021; Schnatz et al., 2021). However, our data warrant future hypothesis-challenging experiments for EV functions using membrane trafficking mutants that disrupt EV release specifically.

Materials and Methods

Drosophila culture

Flies were cultured using standard media and techniques, except larvae for **Fig. 5F-G** (FzC-2 nuclear import), which were cultured on Gerber Natural for Baby, Peach, 2nd Foods (Sitter). Flies used for experiments were maintained at 25°C, except for experiments using Shrub-RNAi, which were maintained at 20°C. Suitable reagents were not available to assess the extent of Tsg101 or Shrub knockdown. However, given that we observe very similar phenotypes for ESCRT RNAi, genomic mutants, and dominant negative mutants (**Figs. 1-2**), we conclude that these RNAi tools phenocopy strong loss-of-function, and that the phenotypes we observe are specific. For detailed information on fly stocks used, see **Table S1**, and for detailed genotype information for each figure panel, see **Table S3**.

Immunohistochemistry

Wandering 3rd instar larvae were dissected in HL3.1 and fixed in HL3.1 with 4% paraformaldehyde for 45 minutes (except **Figs. 2A,B,D,E, Fig. 7** and **Fig. S5** which were fixed for 10 minutes). For α-Fz2 staining, wandering 3rd instar larvae were dissected in 0 mM Ca^{2+} modified *Drosophila* saline (Restrepo et al., 2022) and fixed in 4% paraformaldehyde for 20 minutes. Washes and antibody dilutions were conducted using PBS containing 0.2% Triton X-100 (0.2% PBX), except Fz2-C stain washes and antibody dilutions which were conducted using PBS containing 0.3% Triton X-100. Primary antibody incubations were conducted overnight at 4°C, and secondary antibody incubations for 1-2 hours at room temperature. α-HRP incubations were conducted either overnight at 4°C or for 1-2 hours at room temperature. Prior to imaging, fillets were mounted on slides with Vectashield (Vector Labs) or Abberior Liquid Antifade (Abberior). For detailed information on antibodies used in this study, see **Table S2**.

Electron microscopy

Wandering 3rd instar larvae were dissected and fixed in 1% glutaraldehyde and 4% paraformaldehyde in 1% (0.1M) sodium cacodylate buffer overnight at 4°C. Samples were postfixed in 1% osmium tetroxide and 1.5% potassium ferrocyanide for 1 h, then 1% aqueous uranyl acetate for 0.5 h. Stepwise dehydration was conducted for 10 min each in 30%, 50%, 70%, 85%, and 95% ethanol, followed by 2× 10 min in 100% ethanol. Samples

were transferred to 100% propylene oxide for 1 h, then 3:1 propylene oxide and 812 TAAB
Epon Resin (epon, TAAB Laboratories Equipment Ltd.) for 1 h, then 1:1 propylene
oxide:epon for 1 h and then left overnight in a 1:3 mixture of propylene oxide:epon. Samples
were then transferred to fresh epon for 2 h. Samples were then flat-embedded and
polymerized at 60°C for 48 h, and remounted for sectioning. 70-μm-thin sections were cut on
a Leica UC6 Ultramicrotome (Leica Microsystems), collected onto 2X1 mm slot grids coated
with formvar and carbon, and then poststained with lead citrate (Venable and Coggeshall,
1965). Grids were imaged using a FEI Morgagni transmission electron microscope (FEI)
operating at 80 kV and equipped with an AMT Nanosprint5 camera.

Activity-induced synaptic growth

High K^+ spaced stimulation was performed as described (Piccioli and Littleton, 2014). Briefly,
3^{rd} instar larvae were dissected in HL3 saline (Stewart et al., 1994) at room temperature (in
mM, 70 $NaCl_2$, 5 KCl, 0.2 $CaCl_2$, 20 $MgCl_2$, 10 $NaHCO_3$, 5 trehalose, 115 sucrose, and 5
HEPES (pH=7.2)). Dissecting pins were then moved inward to relax the fillet to 60% of its
original size, and then stimulated 3 times in high K^+ solution (in mM, 40 $NaCl_2$, 90 KCl, 1.5
$CaCl_2$, 20 $MgCl_2$, 10 $NaHCO_3$, 5 trehalose, 5 sucrose, and 5 HEPES (pH=7.2)) for 2 minutes
each, with 10-minute HL3 incubations in between stimulation while on a shaker at room
temperature. Following the 3^{rd} and final stimulation, larvae were incubated in HL3
(approximately 2 minutes) and stretched to their initial length. Mock stimulations were
performed identically to the high K^+ stimulation assay, except HL3 solution was used in place
of high K^+ solution. Larvae were then fixed in 4% PFA in HL3 solution for 15 minutes and
then stained and mounted as above.

Electrophysiology

Wandering 3^{rd} instar larvae were dissected in HL3 saline. Recordings were taken using an
AxoClamp 2B amplifier (Axon Instruments, Burlingame, CA). A recording electrode was filled
with 3M KCl and inserted into muscle 6 at abdominal segments A3 or A4. A stimulating
electrode filled with saline was used to stimulate the severed segmental nerve using an
isolated pulse stimulator (2100; A-M Systems). HFMR was induced by four trains of 100 Hz
stimuli spaced 2 s apart in 0.3 mM extracellular Ca^{2+}. Miniature excitatory junctional
potentials (minis) were recorded 2 min before and 10 min after HFMR induction for Tsg101.
Many *hrs* mutant larvae did not maintain quality mini recordings over 10 minutes, so we
recorded for only 5 minutes. Mini frequency at indicated time points was calculated in 10-s
bins. Fold enhancement was calculated by normalizing to the baseline mini frequency
recorded prior to HFMR induction. Analyses were performed using Clampfit 10.0 software
(Molecular Devices, Sunnyvale, CA). Each n value represents a single muscle recording,
with data generated from at least six individual larvae of each genotype arising from at least
two independent crosses. Resting membrane potentials were between -50 mV and -75 mV
and were not different between genotypes. Input resistances were between 5 MΩ and 10
MΩ and were not different between genotypes.

Imaging and quantification

Acquisition: Analysis of EV cargoes and bouton morphology were conducted from larval
abdominal muscles and segments as indicated in **Table S3**. Z-stacks were acquired using a
Nikon Ni-E upright microscope equipped with a Yokogawa CSU-W1 spinning disk head, an
Andor iXon 897U EMCCD camera, and Nikon Elements AR software. A 60X (n.a. 1.4) oil
immersion objective was used to image NMJs, cell bodies, and fixed axons. Data in **Fig.**
S4C-D were acquired with a Zeiss LSM 800 confocal microscope using a 40x (n.a. 1.4) oil
immersion objective and Zen Black 2.3 software. For colocalization and puncta analysis

branches from muscle 6/7 NMJ from segments A2 and A3 were imaged using Zen Blue software on a Zeiss LSM880 Fast Airyscan microscope in super resolution acquisition mode, using a 63X (n.a. 1.4) oil immersion objective.

Data in **Fig. S1B-E** were acquired on an Abberior FACILITY line STED microscope with 60x (NA1.3) silicone immersion objective, pulsed excitation lasers (561 nm and 640 nm), and a pulsed depletion laser (775 nm) to deplete all signals. Nrg was labeled with STAR ORANGE (Abberior, Inc.) and APP-GFP was labeled with anti-GFP antibodies and STAR RED-labeled secondaries (Abberior, Inc.). Pixel size was set to 40 nm, and single 2D slices were acquired of terminal branches of NMJs that lay in a single focal plane. α-HRP signal was detected using conventional confocal imaging.

For axonal transport, wandering 3^{rd} instar larvae were dissected one at a time in HL3.1. For APP-GFP, larvae were mounted between a slide and coverslip in HL3.1. For Mito-GFP axons the larvae were pinned in a sylgard coated dish covered with HL3.1. Dissection and imaging for each larva was completed within 30 minutes. Timelapse images were taken on the same Nikon Ni-E microscope described above. Images were taken of axon bundles proximal to the ventral ganglion (roughly within 100-300 µm). For APP transport, timelapse images were acquired for 3 minutes using 60X (n.a. 1.4) oil immersion objective. For mitochondria timelapse, images were acquired for 7 minutes using 60X (n.a. 1.0) water immersion objective. 9 Z slices were collected per frame (Step size 0.3 µm, with no acquisition delay between timepoints, resulting in a frame rate of 2.34-2.37 sec/frame). To visualize moving particles for mitochondria, a third of the axon in the field of view was photobleached using an Andor Mosaic digital micromirror device operated by Andor IQ software, to eliminate fluorescence from stationary particles that would interfere with visualization of particles moving into the bleached region. Image acquisition settings were identical for all images in each independent experiment.

EV cargo quantification and colocalization: Volumetric analysis was performed using Volocity 6.0 software. For each image, both type 1s and 1b boutons were retained for analysis while axons were cropped out, (except in limited cases where 1s boutons were very faint and therefore were cropped as their inclusion would cause the HRP threshold to be excessively dilated for other branches). The presynaptic volume was defined by an HRP threshold, excluding objects smaller than 7 µm³ and closing holes. The postsynaptic region was defined by a 3.3 µm dilation of the HRP mask. However, for Evi-GFP, where the presynaptic signal vastly exceeded postsynaptic signal, we analyzed only the distal 2.9 µm of this postsynaptic dilation region to eliminate the bleed-over haze from the presynaptic signal. EV cargo and Rab signals were manually thresholded to select particles brighter than the muscle background. EV cargo integrated density in these thresholded puncta was normalized to the overall presynaptic volume. These values were further normalized to the mean of the control to produce a "normalized puncta intensity" value for each NMJ. For colocalization, the overlap of the two channels was measured in Volocity 6.0 and used for calculation of Mander's coefficients.

To perform particle-based analyses of EV puncta density and width, 2D-STED micrographs were denoised using Noise2Void (Krull et al., 2019). Briefly, a model was trained using the Nrg channel, using 10 control and 10 Tsg101^{KD} images as a training set. This model was used to denoise both Nrg and APP channels. Presynaptic regions were segmented using a conventional confocal image of HRP, as described above. This mask was dilated by 3 µm to generate a postsynaptic mask containing the vast majority of EV signal, from which a 10% dilation of the presynaptic mask was subtracted to remove any presynaptic signal. Finally, the postsynaptic mask was further dilated by 10% to generate a 300 nm buffer, and the

remainder of the image was defined as background (e.g. nonspecific antibody signal) (BG, see **Fig. S1C** for schematic). To detect particles, each channel was independently rescaled from 0-1 (min and max pixel values), processed by a Mexican Hat filter (radius=4), and local intensity maxima were detected using a prominence value of 1.25. Local maxima were used as seeds to fit a 2D gaussian on the original (unscaled) denoised images, using the plugin GaussFit on Spot (https://imagej.net/ij/plugins/gauss-fit-spot/index.html). Parameters for fitting were as follows: shape=Circle fitmode=NelderMead rectangle=2.5 pixel=40 max=500 cpcf=1 base=0.

Quantification of electron micrographs: A single medial section of each bouton was selected for analysis. Two experimenters, blinded to genotype, together recorded the presence of autophagic vacuoles, including phagopores (double or dense membrane but not closed; note that depending on the plane of section, some of these may appear as autophagosomes), autophagosomes (contents with similar properties to the cytoplasm, fully enclosed in the section by a double membrane), and autolyososomes (contents are electron dense) (Lucocq and Hacker, 2013; Nagy et al., 2015). We also evaluated whether boutons lacked subsynaptic reticulum, or featured postsynaptic pockets (electron-lucent areas extending at least 300 nm from the presynaptic membrane (Packard et al., 2002).

Quantification of GFP-mCherry-Atg8 distribution: A single field-of-view confocal stack (62x62 µm) from the larval ventral ganglion, containing 10-15 Vglut-expressing cell bodies, was manually thresholded in Volocity 6.0 software to segment and measure the volume and integrated fluorescence density of soma, GFP puncta, and mCherry puncta. The overlap between the GFP and mCherry channels was used for the calculation of the Mander's coefficient (fraction of total mCherry-puncta integrated density found in the GFP-puncta positive volume).

Axon and cell body measurements: To measure intensity of EV cargoes, axons proximal to the ventral ganglion (within 100-300 µm) were imaged as described above. Images were analyzed in Fiji by making sum projections, cropping out unwanted debris or other tissue and generating a mask from the α-HRP signal. The total intensity of the EV cargo was then measured within the masked HRP area. For cell bodies, EV cargo intensity was measured from a middle slice through the motor neuron cell body layer of the ventral ganglion using Fiji.

Quantification of live axonal trafficking of APP-GFP and Mito-GFP puncta: To quantify APP-GFP and mitochondria dynamics in live axons, maximum intensity projections of time course images were processed in Fiji to subtract background and adjust for XY drift using the StackReg plugin. Kymographs were generated from 1-4 axons per animal using the Fiji plugin KymographBuilder. Kymographs were blinded and number of tracks were manually counted. The minimum track length measured was 3 µm with most tracks above 5 µm. Velocity was measured by calculating the slope of the identified tracks.

Bouton quantification: The experimenter was blinded to genotypes and then manually counted the total number of type 1 synaptic boutons on the NMJ on muscle 6 and 7 in the abdominal segments A2 and A3 of third instar wandering larvae. A synaptic bouton was considered a spherical varicosity, defined by the presence of the synaptic vesicle marker Synaptotagmin 1, the active zone marker Bruchpilot (Brp) or the neuronal membrane marker Hrp. For quantifying ghost boutons (basal and activity-induced), the experimenter was blinded to genotype and condition and ghost boutons were quantified as α-HRP-positive structures with a visible connection to the main NMJ arbor, and without α-DLG staining. For quantifying DLG "featheriness", the experimenter was blinded to genotype and scored the

number of NMJs with at least one region of fenestrated Dlg that extended far from the
bouton periphery.

Active zone quantification: To count the active zones in fluorescence micrographs, Brp-
stained punctae were assessed on maximum intensity projection images. The Trainable
Weka Segmentation (TWS) machine-learning tool
(https://doi.org/10.1093/bioinformatics/btx180) in Fiji was used to manually annotate Brp-
positive punctae with different fluorescence intensities, and to train a classifier that
automatically segmented the Brp-positive active zones. The objects segmented via the
applied classifier were subjected to Huang auto thresholding to obtain binary masks. Next,
we applied a Watershed processing on the binary image, to improve the isolation of
individual neighboring active zones from the diffraction limited images. We performed
particle analysis on the segmented active zones and obtained their number, area, and
integrated density. To determine the NMJ area using TWS, we trained the classifier by
annotating the HRP positive NMJ on maximum intensity projections of the HRP channel.
Axons were manually cropped from the image before TWS. The segmented HRP area was
subjected to Huang auto thresholding, the binary masks were selected and the NMJ area
was obtained via the "Analyze particle" function in FIJI of particles larger than 5 µm (to
eliminate from the analysis residual HRP EV debris segmented in a very few images).

Quantification of Frizzled2 C-terminus nuclear localization: To quantify Fz2-C puncta,
muscles 6 and 7 were imaged from segments A2 and A3 in larvae where the experimenter
was blinded to genotype. Muscle nuclei were identified by the boundaries of LamC staining
(which recognizes the nuclear envelope). Nuclear puncta were quantified as aggregates of
staining that exceeded the size and fluorescence intensity of non-specific background
staining from the rabbit α-Fz2-C antibody. No nuclei were excluded from quantification and
all nuclei pooled for final statistical analysis. In all genotypes, >250 individual nuclei were
scored.

Quantification of Draper knockdown: To analyze Draper levels in axonal bundles (which
include neurons and glia), a region of axon proximal to muscle 4 was cropped to 100x100
pixels in a 3D spinning disk confocal Z-stack, using Fiji. This image was masked on the HRP
channel and the mean α-Draper intensity was calculated in the 3D volume. To analyze
Draper levels at the NMJ (which includes the presynaptic neuron and adjacent or optically
overlapping postsynaptic muscle membrane), images were cropped to include only type 1b
bouton branches (excluding axon bundles, axon, type 1s bouton branches, or any non-
bouton material). The image was masked on the HRP channel, dilated by 0.22 µm, and the
mean α-Draper intensity was calculated in the 3D volume.

Statistics

All statistical measurements were performed in GraphPad Prism (see **Table S3**).
Comparisons were made separately for presynaptic and postsynaptic datasets, due to
differences between these compartments for intensity, signal-to-noise ratio, and variance.
Datasets were tested for normality, and statistical significance was tested as noted for each
experiment in **Table S3**. Statistical significance is indicated as *P < 0.05; **P < 0.01; ***P <
0.001.

Acknowledgements

44 We thank the Developmental Studies Hybridoma Bank created by the NICHD of the NIH,
45 and the Bloomington *Drosophila* Stock Center (Indiana University, Bloomington, IN, NIH
46 P40OD018537). This work was supported by grant 1S10OD034223-01 for the Abberior

Facility Line STED microscope, housed in the Brandeis Light Microscopy Core Facility. We thank Berith Isaac and the Brandeis Electron Microscopy Facility for assistance with EM, Michael Boutros, Josie Clowney, and Tor Erik Ruston for fly lines, and Harald Stenmark for discussion of simaphagy. This work was supported by NINDS grants R01 NS103967 to A.A.R., F32 NS120909 to E.C.D., and R01 NS110907 to T.J.M.

Author contributions

ECD, KPH, KK, MFP, CRB, BAS, TJM, and AAR designed the study and experiments. ECD, KPH, KK, MFP, BE, CRB, MR, RCS, TJM conducted the experiments. ECD, KPH, KK, MFP, BE, CRB, SJDS, MR, RCS, TJM and AAR performed the analyses. ECD and AAR wrote the manuscript, and all authors edited the manuscript. This article contains supporting information (5 Supplemental Figures and 3 tables).

Competing interest statement

The authors declare no competing financial interests.

References

Adolfsen, B., S. Saraswati, M. Yoshihara, and J.T. Littleton. 2004. Synaptotagmins are trafficked to distinct subcellular domains including the postsynaptic compartment. *J Cell Biol*. 166:249-260.

Arbo, B.D., L.R. Cechinel, R.P. Palazzo, and I.R. Siqueira. 2020. Endosomal dysfunction impacts extracellular vesicle release: Central role in Abeta pathology. *Ageing Res Rev*. 58:101006.

Asai, H., S. Ikezu, S. Tsunoda, M. Medalla, J. Luebke, T. Haydar, B. Wolozin, O. Butovsky, S. Kugler, and T. Ikezu. 2015. Depletion of microglia and inhibition of exosome synthesis halt tau propagation. *Nat Neurosci*. 18:1584-1593.

Ashley, J., B. Cordy, D. Lucia, L.G. Fradkin, V. Budnik, and T. Thomson. 2018. Retrovirus-like Gag Protein Arc1 Binds RNA and Traffics across Synaptic Boutons. *Cell*. 172:262-274 e211.

Ataman, B., J. Ashley, D. Gorczyca, M. Gorczyca, D. Mathew, C. Wichmann, S.J. Sigrist, and V. Budnik. 2006. Nuclear trafficking of Drosophila Frizzled-2 during synapse development requires the PDZ protein dGRIP. *Proc Natl Acad Sci U S A*. 103:7841-7846.

Ataman, B., J. Ashley, M. Gorczyca, P. Ramachandran, W. Fouquet, S.J. Sigrist, and V. Budnik. 2008. Rapid activity-dependent modifications in synaptic structure and function require bidirectional Wnt signaling. *Neuron*. 57:705-718.

Barber, C.F., R.A. Jorquera, J.E. Melom, and J.T. Littleton. 2009. Postsynaptic regulation of synaptic plasticity by synaptotagmin 4 requires both C2 domains. *J Cell Biol*. 187:295-310.

Beckett, K., S. Monier, L. Palmer, C. Alexandre, H. Green, E. Bonneil, G. Raposo, P. Thibault, R. Le Borgne, and J.P. Vincent. 2013. Drosophila S2 cells secrete wingless on exosome-like vesicles but the wingless gradient forms independently of exosomes. *Traffic*. 14:82-96.

Birdsall, V., K. Kirwan, M. Zhu, Y. Imoto, S.M. Wilson, S. Watanabe, and C.L. Waites. 2022. Axonal transport of Hrs is activity dependent and facilitates synaptic vesicle protein degradation. *Life Sci Alliance*. 5.

Blanchette, C.R., and A.A. Rodal. 2020. Mechanisms for biogenesis and release of neuronal extracellular vesicles. *Curr Opin Neurobiol*. 63:104-110.

Blanchette, C.R., A.L. Scalera, K.P. Harris, Z. Zhao, E.C. Dresselhaus, K. Koles, A. Yeh, J.K. Apiki, B.A. Stewart, and A.A. Rodal. 2022. Local regulation of extracellular vesicle traffic by the synaptic endocytic machinery. *J. Cell Biol*. 10.1083/jcb.202112094.

Boecker, C.A., and E.L. Holzbaur. 2019. Vesicular degradation pathways in neurons: at the crossroads of autophagy and endo-lysosomal degradation. *Curr Opin Neurobiol*. 57:94-101.

Brugger, M., A. Lauri, Y. Zhen, L.L. Gramegna, B. Zott, N. Sekulic, G. Fasano, R. Kopajtich, V. Cordeddu, F.C. Radio, C. Mancini, S. Pizzi, G. Paradisi, G. Zanni, G. Vasco, R. Carrozzo, F. Palombo, C. Tonon, R. Lodi, C. La Morgia, M. Arelin, C. Blechschmidt, T. Finck, V. Sorensen, K. Kreiser, G. Strobl-Wildemann, H. Daum, R. Michaelson-Cohen, L. Ziccardi, G. Zampino, H. Prokisch, R. Abou Jamra, C. Fiorini, T. Arzberger, J. Winkelmann, L. Caporali, V. Carelli, H. Stenmark, M. Tartaglia, and M. Wagner. 2024. Bi-allelic variants in

SNF8 cause a disease spectrum ranging from severe developmental and epileptic encephalopathy to syndromic optic atrophy. *Am J Hum Genet*. 111:594-613.

Budnik, V., C. Ruiz-Canada, and F. Wendler. 2016. Extracellular vesicles round off communication in the nervous system. *Nat Rev Neurosci*. 17:160-172.

Buratta, S., B. Tancini, K. Sagini, F. Delo, E. Chiaradia, L. Urbanelli, and C. Emiliani. 2020. Lysosomal Exocytosis, Exosome Release and Secretory Autophagy: The Autophagic- and Endo-Lysosomal Systems Go Extracellular. *Int J Mol Sci*. 21.

Caussinus, E., O. Kanca, and M. Affolter. 2011. Fluorescent fusion protein knockout mediated by anti-GFP nanobody. *Nat Struct Mol Biol*. 19:117-121.

Chang, Y.C., Y. Gao, J.Y. Lee, Y.J. Peng, J. Langen, and K.T. Chang. 2024. Identification of secretory autophagy as a mechanism modulating activity-induced synaptic remodeling. *Proc Natl Acad Sci U S A*. 121:e2315958121.

Cone, A.S., S.N. Hurwitz, G.S. Lee, X. Yuan, Y. Zhou, Y. Li, and D.G. Meckes, Jr. 2020. Alix and Syntenin-1 direct amyloid precursor protein trafficking into extracellular vesicles. *BMC Mol Cell Biol*. 21:58.

Coulter, M.E., C.M. Dorobantu, G.A. Lodewijk, F. Delalande, S. Cianferani, V.S. Ganesh, R.S. Smith, E.T. Lim, C.S. Xu, S. Pang, E.T. Wong, H.G.W. Lidov, M.L. Calicchio, E. Yang, D.M. Gonzalez, T.M. Schlaeger, G.H. Mochida, H. Hess, W.A. Lee, M.K. Lehtinen, T. Kirchhausen, D. Haussler, F.M.J. Jacobs, R. Gaudin, and C.A. Walsh. 2018. The ESCRT-III Protein CHMP1A Mediates Secretion of Sonic Hedgehog on a Distinctive Subtype of Extracellular Vesicles. *Cell Rep*. 24:973-986 e978.

Dagar, S., and S. Subramaniam. 2023. Tunneling Nanotube: An Enticing Cell-Cell Communication in the Nervous System. *Biology (Basel)*. 12.

Daly, C.A., E.T. Hall, and S.K. Ogden. 2022. Regulatory mechanisms of cytoneme-based morphogen transport. *Cell Mol Life Sci*. 79:119.

Das, S., S. Yu, R. Sakamori, E. Stypulkowski, and N. Gao. 2012. Wntless in Wnt secretion: molecular, cellular and genetic aspects. *Front Biol (Beijing)*. 7:587-593.

Dinkins, M.B., J. Enasko, C. Hernandez, G. Wang, J. Kong, I. Helwa, Y. Liu, A.V. Terry, Jr., and E. Bieberich. 2016. Neutral Sphingomyelinase-2 Deficiency Ameliorates Alzheimer's Disease Pathology and Improves Cognition in the 5XFAD Mouse. *J Neurosci*. 36:8653-8667.

Doyotte, A., M.R. Russell, C.R. Hopkins, and P.G. Woodman. 2005. Depletion of TSG101 forms a mammalian "Class E" compartment: a multicisternal early endosome with multiple sorting defects. *J Cell Sci*. 118:3003-3017.

Faure, J., G. Lachenal, M. Court, J. Hirrlinger, C. Chatellard-Causse, B. Blot, J. Grange, G. Schoehn, Y. Goldberg, V. Boyer, F. Kirchhoff, G. Raposo, J. Garin, and R. Sadoul. 2006. Exosomes are released by cultured cortical neurones. *Mol Cell Neurosci*. 31:642-648.

Ferguson, S.M. 2018. Axonal transport and maturation of lysosomes. *Curr Opin Neurobiol*. 51:45-51.

Fernandes, A.R., J.P. Martins, E.R. Gomes, C.S. Mendes, and R.O. Teodoro. 2023. Drosophila motor neuron boutons remodel through membrane blebbing coupled with muscle contraction. *Nat Commun*. 14:3352.

Fuentes-Medel, Y., M.A. Logan, J. Ashley, B. Ataman, V. Budnik, and M.R. Freeman. 2009. Glia and muscle sculpt neuromuscular arbors by engulfing destabilized synaptic boutons and shed presynaptic debris. *PLoS Biol*. 7:e1000184.

Gassama, Y., and A. Favereaux. 2021. Emerging Roles of Extracellular Vesicles in the Central Nervous System: Physiology, Pathology, and Therapeutic Perspectives. *Front Cell Neurosci*. 15:626043.

Goncalves, M.B., T. Malmqvist, E. Clarke, C.J. Hubens, J. Grist, C. Hobbs, D. Trigo, M. Risling, M. Angeria, P. Damberg, T.P. Carlstedt, and J.P. Corcoran. 2015. Neuronal RARbeta Signaling Modulates PTEN Activity Directly in Neurons and via Exosome Transfer in Astrocytes to Prevent Glial Scar Formation and Induce Spinal Cord Regeneration. *J Neurosci*. 35:15731-15745.

Gong, J., R. Korner, L. Gaitanos, and R. Klein. 2016. Exosomes mediate cell contact-independent ephrin-Eph signaling during axon guidance. *J Cell Biol*. 214:35-44.

Gruenberg, J. 2020. Life in the lumen: The multivesicular endosome. *Traffic*. 21:76-93.

Guo, B.B., S.A. Bellingham, and A.F. Hill. 2015. The neutral sphingomyelinase pathway regulates packaging of the prion protein into exosomes. *J Biol Chem*. 290:3455-3467.

Harris, K.P., Y.V. Zhang, Z.D. Piccioli, N. Perrimon, and J.T. Littleton. 2016. The postsynaptic t-SNARE Syntaxin 4 controls traffic of Neuroligin 1 and Synaptotagmin 4 to regulate retrograde signaling. *Elife*. 5.

Holm, M.M., J. Kaiser, and M.E. Schwab. 2018. Extracellular Vesicles: Multimodal Envoys in Neural Maintenance and Repair. *Trends Neurosci*. 41:360-372.

Kerr, K.S., Y. Fuentes-Medel, C. Brewer, R. Barria, J. Ashley, K.C. Abruzzi, A. Sheehan, O.E. Tasdemir-Yilmaz, M.R. Freeman, and V. Budnik. 2014. Glial wingless/Wnt regulates glutamate receptor clustering and synaptic physiology at the Drosophila neuromuscular junction. *J Neurosci*. 34:2910-2920.

Klionsky, D.J., A.K. Abdel-Aziz, S. Abdelfatah, M. Abdellatif, A. Abdoli, S. Abel, H. Abeliovich, M.H. Abildgaard, Y.P. Abudu, A. Acevedo-Arozena, I.E. Adamopoulos, K. Adeli, T.E. Adolph, A. Adornetto, E. Aflaki, G. Agam, A. Agarwal, B.B. Aggarwal, M. Agnello, P. Agostinis, J.N. Agrewala, A. Agrotis, P.V. Aguilar, S.T. Ahmad, Z.M. Ahmed, U. Ahumada-Castro, S. Aits, S. Aizawa, Y. Akkoc, T. Akoumianaki, H.A. Akpinar, A.M. Al-Abd, L. Al-Akra, A. Al-Gharaibeh, M.A. Alaoui-Jamali, S. Alberti, E. Alcocer-Gomez, C. Alessandri, M. Ali, M.A. Alim Al-Bari, S. Aliwaini, J. Alizadeh, E. Almacellas, A. Almasan, A. Alonso, G.D. Alonso, N. Altan-Bonnet, D.C. Altieri, E.M.C. Alvarez, S. Alves, C. Alves da Costa, M.M. Alzaharna, M. Amadio, C. Amantini, C. Amaral, S. Ambrosio, A.O. Amer, V. Ammanathan, Z. An, S.U. Andersen, S.A. Andrabi, M. Andrade-Silva, A.M. Andres, S. Angelini, D. Ann, U.C. Anozie, M.Y. Ansari, P. Antas, A. Antebi, Z. Anton, T. Anwar, L. Apetoh, N. Apostolova, T. Araki, Y. Araki, K. Arasaki, W.L. Araujo, J. Araya, C. Arden, M.A. Arevalo, S. Arguelles, E. Arias, J. Arikkath, H. Arimoto, A.R. Ariosa, D. Armstrong-James, L. Arnaune-Pelloquin, A. Aroca, D.S. Arroyo, I. Arsov, R. Artero, D.M.L. Asaro, M. Aschner, M. Ashrafizadeh, O. Ashur-Fabian, A.G. Atanasov, A.K. Au, P. Auberger, H.W. Auner, L. Aurelian, et al. 2021. Guidelines for the use and interpretation of assays for monitoring autophagy (4th edition)(1). *Autophagy*. 17:1-382.

Koles, K., J. Nunnari, C. Korkut, R. Barria, C. Brewer, Y. Li, J. Leszyk, B. Zhang, and V. Budnik. 2012. Mechanism of evenness interrupted (Evi)-exosome release at synaptic boutons. *J Biol Chem*. 287:16820-16834.

Koles, K., A.R. Yeh, and A.A. Rodal. 2015. Tissue-specific tagging of endogenous loci in Drosophila melanogaster. *Biol Open*. 5:83-89.

Korkut, C., B. Ataman, P. Ramachandran, J. Ashley, R. Barria, N. Gherbesi, and V. Budnik. 2009. Trans-synaptic transmission of vesicular Wnt signals through Evi/Wntless. *Cell*. 139:393-404.

Korkut, C., Y. Li, K. Koles, C. Brewer, J. Ashley, M. Yoshihara, and V. Budnik. 2013. Regulation of postsynaptic retrograde signaling by presynaptic exosome release. *Neuron*. 77:1039-1046.

Krull, A., T. Buchholz, and F. Jug. 2019. Noise2Void - Learning Denoising from Single Noisy Images.10.48550/arXiv.41811.10980.

Lachenal, G., K. Pernet-Gallay, M. Chivet, F.J. Hemming, A. Belly, G. Bodon, B. Blot, G. Haase, Y. Goldberg, and R. Sadoul. 2011. Release of exosomes from differentiated neurons and its regulation by synaptic glutamatergic activity. *Mol Cell Neurosci*. 46:409-418.

Lamark, T., and T. Johansen. 2021. Mechanisms of Selective Autophagy. *Annu Rev Cell Dev Biol*. 37:143-169.

Lauwers, E., Y.C. Wang, R. Gallardo, R. Van der Kant, E. Michiels, J. Swerts, P. Baatsen, S.S. Zaiter, S.R. McAlpine, N.V. Gounko, F. Rousseau, J. Schymkowitz, and P. Verstreken. 2018. Hsp90 Mediates Membrane Deformation and Exosome Release. *Mol Cell*. 71:689-702 e689.

Lee, S.H., S.M. Shin, P. Zhong, H.T. Kim, D.I. Kim, J.M. Kim, W.D. Heo, D.W. Kim, C.Y. Yeo, C.H. Kim, and Q.S. Liu. 2018. Reciprocal control of excitatory synapse numbers by Wnt and Wnt inhibitor PRR7 secreted on exosomes. *Nat Commun*. 9:3434.

Lefebvre, C., R. Legouis, and E. Culetto. 2018. ESCRT and autophagies: Endosomal functions and beyond. *Semin Cell Dev Biol*. 74:21-28.

Leidal, A.M., and J. Debnath. 2021. Emerging roles for the autophagy machinery in extracellular vesicle biogenesis and secretion. *FASEB Bioadv*. 3:377-386.

Lin, T.H., D.M. Bis-Brewer, A.E. Sheehan, L.N. Townsend, D.C. Maddison, S. Zuchner, G.A. Smith, and M.R. Freeman. 2021. TSG101 negatively regulates mitochondrial biogenesis in axons. *Proc Natl Acad Sci U S A*. 118.

Lizarraga-Valderrama, L.R., and G.K. Sheridan. 2021. Extracellular vesicles and intercellular communication in the central nervous system. *FEBS Lett*. 595:1391-1410.

Lloyd, T.E., R. Atkinson, M.N. Wu, Y. Zhou, G. Pennetta, and H.J. Bellen. 2002. Hrs regulates endosome membrane invagination and tyrosine kinase receptor signaling in Drosophila. *Cell*. 108:261-269.

Lucocq, J.M., and C. Hacker. 2013. Cutting a fine figure: On the use of thin sections in electron microscopy to quantify autophagy. *Autophagy*. 9:1443-1448.

Mathew, D., B. Ataman, J. Chen, Y. Zhang, S. Cumberledge, and V. Budnik. 2005. Wingless signaling at synapses is through cleavage and nuclear import of receptor DFrizzled2. *Science*. 310:1344-1347.

Men, Y., J. Yelick, S. Jin, Y. Tian, M.S.R. Chiang, H. Higashimori, E. Brown, R. Jarvis, and Y. Yang. 2019. Exosome reporter mice reveal the involvement of exosomes in mediating neuron to astroglia communication in the CNS. *Nat Commun*. 10:4136.

Miech, C., H.U. Pauer, X. He, and T.L. Schwarz. 2008. Presynaptic local signaling by a canonical wingless pathway regulates development of the *Drosophila* neuromuscular junction. *J Neurosci*. 28:10875-10884.

Migliano, S.M., S.W. Schultz, E.M. Wenzel, S. Takats, D. Liu, S. Mork, K.W. Tan, T.E. Rusten, C. Raiborg, and H. Stenmark. 2023. Removal of hypersignaling endosomes by simaphagy. *Autophagy*:1-23.

Millarte, V., S. Schlienger, S. Kalin, and M. Spiess. 2022. Rabaptin5 targets autophagy to damaged endosomes and Salmonella vacuoles via FIP200 and ATG16L1. *EMBO Rep*. 23:e53429.

Mosca, T.J., and T.L. Schwarz. 2010. The nuclear import of Frizzled2-C by Importins-beta11 and alpha2 promotes postsynaptic development. *Nat Neurosci*. 13:935-943.

Nagy, P., A. Varga, A.L. Kovacs, S. Takats, and G. Juhasz. 2015. How and why to study autophagy in Drosophila: it's more than just a garbage chute. *Methods*. 75:151-161.

Nakatogawa, H. 2020. Mechanisms governing autophagosome biogenesis. *Nat Rev Mol Cell Biol*. 21:439-458.

Packard, M., E.S. Koo, M. Gorczyca, J. Sharpe, S. Cumberledge, and V. Budnik. 2002. The Drosophila Wnt, wingless, provides an essential signal for pre- and postsynaptic differentiation. *Cell*. 111:319-330.

Piccioli, Z.D., and J.T. Littleton. 2014. Retrograde BMP signaling modulates rapid activity-dependent synaptic growth via presynaptic LIM kinase regulation of cofilin. *J Neurosci*. 34:4371-4381.

Raymond, C.K., I. Howald-Stevenson, C.A. Vater, and T.H. Stevens. 1992. Morphological classification of the yeast vacuolar protein sorting mutants: evidence for a prevacuolar compartment in class E vps mutants. *Mol Biol Cell*. 3:1389-1402.

Restrepo, L.J., A.T. DePew, E.R. Moese, S.R. Tymanskyj, M.J. Parisi, M.A. Aimino, J.C. Duhart, H. Fei, and T.J. Mosca. 2022. gamma-secretase promotes Drosophila postsynaptic development through the cleavage of a Wnt receptor. *Dev Cell*. 57:1643-1660 e1647.

Rodahl, L.M., K. Haglund, C. Sem-Jacobsen, F. Wendler, J.P. Vincent, K. Lindmo, T.E. Rusten, and H. Stenmark. 2009. Disruption of Vps4 and JNK function in Drosophila causes tumour growth. *PLoS One*. 4:e4354.

Rodger, C., E. Flex, R.J. Allison, A. Sanchis-Juan, M.A. Hasenahuer, S. Cecchetti, C.E. French, J.R. Edgar, G. Carpentieri, A. Ciolfi, F. Pantaleoni, A. Bruselles, R. Onesimo, G. Zampino, F. Marcon, E. Siniscalchi, M. Lees, D. Krishnakumar, E. McCann, D. Yosifova, J. Jarvis, M.C. Kruer, W. Marks, J. Campbell, L.E. Allen, S. Gustincich, F.L. Raymond, M. Tartaglia, and E. Reid. 2020. De Novo VPS4A Mutations Cause Multisystem Disease with Abnormal Neurodevelopment. *Am J Hum Genet*. 107:1129-1148.

Rusten, T.E., T. Vaccari, K. Lindmo, L.M. Rodahl, I.P. Nezis, C. Sem-Jacobsen, F. Wendler, J.P. Vincent, A. Brech, D. Bilder, and H. Stenmark. 2007. ESCRTs and Fab1 regulate distinct steps of autophagy. *Curr Biol*. 17:1817-1825.

Sackmann, V., M.S. Sinha, C. Sackmann, L. Civitelli, J. Bergstrom, A. Ansell-Schultz, and M. Hallbeck. 2019. Inhibition of nSMase2 Reduces the Transfer of Oligomeric alpha-Synuclein Irrespective of Hypoxia. *Front Mol Neurosci*. 12:200.

Sadoul, R., M.H. Laporte, R. Chassefeyre, K.I. Chi, Y. Goldberg, C. Chatellard, F.J. Hemming, and S. Fraboulet. 2018. The role of ESCRT during development and functioning of the nervous system. *Semin Cell Dev Biol*. 74:40-49.

Schnatz, A., C. Muller, A. Brahmer, and E.M. Kramer-Albers. 2021. Extracellular Vesicles in neural cell interaction and CNS homeostasis. *FASEB Bioadv*. 3:577-592.

Sheehan, P., M. Zhu, A. Beskow, C. Vollmer, and C.L. Waites. 2016. Activity-Dependent Degradation of Synaptic Vesicle Proteins Requires Rab35 and the ESCRT Pathway. *J Neurosci*. 36:8668-8686.

Sidibe, D.K., M.C. Vogel, and S. Maday. 2022. Organization of the autophagy pathway in neurons. *Curr Opin Neurobiol*. 75:102554.

Song, Z., Y. Xu, W. Deng, L. Zhang, H. Zhu, P. Yu, Y. Qu, W. Zhao, Y. Han, and C. Qin. 2020. Brain Derived Exosomes Are a Double-Edged Sword in Alzheimer's Disease. *Front Mol Neurosci*. 13:79.

Soukup, S.F., S. Kuenen, R. Vanhauwaert, J. Manetsberger, S. Hernandez-Diaz, J. Swerts, N. Schoovaerts, S. Vilain, N.V. Gounko, K. Vints, A. Geens, B. De Strooper, and P. Verstreken. 2016. A LRRK2-Dependent EndophilinA Phosphoswitch Is Critical for Macroautophagy at Presynaptic Terminals. *Neuron*. 92:829-844.

Stewart, B.A., H.L. Atwood, J.J. Renger, J. Wang, and C.F. Wu. 1994. Improved stability of Drosophila larval neuromuscular preparations in haemolymph-like physiological solutions. *J Comp Physiol [A]*. 175:179-191.

Sweeney, N.T., J.E. Brenman, Y.N. Jan, and F.B. Gao. 2006. The coiled-coil protein shrub controls neuronal morphogenesis in Drosophila. *Curr Biol*. 16:1006-1011.

Takahashi, Y., H. He, Z. Tang, T. Hattori, Y. Liu, M.M. Young, J.M. Serfass, L. Chen, M. Gebru, C. Chen, C.A. Wills, J.M. Atkinson, H. Chen, T. Abraham, and H.G. Wang. 2018. An autophagy assay reveals the ESCRT-III component CHMP2A as a regulator of phagophore closure. *Nat Commun*. 9:2855.

Ugbode, C., and R.J.H. West. 2021. Lessons learned from CHMP2B, implications for frontotemporal dementia and amyotrophic lateral sclerosis. *Neurobiol Dis*. 147:105144.

van Niel, G., G. D'Angelo, and G. Raposo. 2018. Shedding light on the cell biology of extracellular vesicles. *Nat Rev Mol Cell Biol*. 19:213-228.

Vietri, M., M. Radulovic, and H. Stenmark. 2020. The many functions of ESCRTs. *Nat Rev Mol Cell Biol*. 21:25-42.

Vilcaes, A.A., N.L. Chanaday, and E.T. Kavalali. 2021. Interneuronal exchange and functional integration of synaptobrevin via extracellular vesicles. *Neuron*. 109:971-983 e975.

Walsh, R.B., E.C. Dresselhaus, A.N. Becalska, M.J. Zunitch, C.R. Blanchette, A.L. Scalera, T. Lemos, S.M. Lee, J. Apiki, S. Wang, B. Isaac, A. Yeh, K. Koles, and A.A. Rodal. 2021. Opposing functions for retromer and Rab11 in extracellular vesicle traffic at presynaptic terminals. *J Cell Biol*. 220:e202012034.

Wang, B.B., H. Xu, S. Isenmann, C. Huang, X. Elorza-Vidal, G.Y. Rychkov, R. Estevez, R.B. Schittenhelm, G.L. Lukacs, and P.M. Apaja. 2022. Ubr1-induced selective endophagy/autophagy protects against the endosomal and Ca(2+)-induced proteostasis disease stress. *Cell Mol Life Sci*. 79:167.

Wang, J., N. Fang, J. Xiong, Y. Du, Y. Cao, and W.K. Ji. 2021. An ESCRT-dependent step in fatty acid transfer from lipid droplets to mitochondria through VPS13D-TSG101 interactions. *Nat Commun*. 12:1252.

Welsh, J.A., D.C.I. Goberdhan, L. O'Driscoll, E.I. Buzas, C. Blenkiron, B. Bussolati, H. Cai, D. Di Vizio, T.A.P. Driedonks, U. Erdbrugger, J.M. Falcon-Perez, Q.L. Fu, A.F. Hill, M. Lenassi, S.K. Lim, M.G. Mahoney, S. Mohanty, A. Moller, R. Nieuwland, T. Ochiya, S. Sahoo, A.C. Torrecilhas, L. Zheng, A. Zijlstra, S. Abuelreich, R. Bagabas, P. Bergese, E.M. Bridges, M. Brucale, D. Burger, R.P. Carney, E. Cocucci, R. Crescitelli, E. Hanser, A.L. Harris, N.J. Haughey, A. Hendrix, A.R. Ivanov, T. Jovanovic-Talisman, N.A. Kruh-Garcia, V. Ku'ulei-Lyn Faustino, D. Kyburz, C. Lasser, K.M. Lennon, J. Lotvall, A.L. Maddox, E.S. Martens-Uzunova, R.R. Mizenko, L.A. Newman, A. Ridolfi, E. Rohde, T. Rojalin, A. Rowland, A. Saftics, U.S. Sandau, J.A. Saugstad, F. Shekari, S. Swift, D. Ter-Ovanesyan, J.P. Tosar, Z. Useckaite, F. Valle, Z. Varga, E. van der Pol, M.J.C. van Herwijnen, M.H.M. Wauben, A.M. Wehman, S. Williams, A. Zendrini, A.J. Zimmerman, C. Thery, and K.W. Witwer. 2024. Minimal information for studies of extracellular vesicles (MISEV2023): From basic to advanced approaches. *J Extracell Vesicles*. 13:e12404.

Wenzel, E.M., S.W. Schultz, K.O. Schink, N.M. Pedersen, V. Nahse, A. Carlson, A. Brech, H. Stenmark, and C. Raiborg. 2018. Concerted ESCRT and clathrin recruitment waves define the timing and morphology of intraluminal vesicle formation. *Nat Commun*. 9:2932.

West, R.J., Y. Lu, B. Marie, F.B. Gao, and S.T. Sweeney. 2015. Rab8, POSH, and TAK1 regulate synaptic growth in a Drosophila model of frontotemporal dementia. *J Cell Biol*. 208:931-947.

Won, J.H., and K.O. Cho. 2021. Wg secreted by conventional Golgi transport diffuses and forms Wg gradient whereas Wg tethered to extracellular vesicles do not diffuse. *Cell Death Differ*. 28:1013-1025.

Yoshihara, M., B. Adolfsen, K.T. Galle, and J.T. Littleton. 2005. Retrograde signaling by Syt 4 induces presynaptic release and synapse-specific growth. *Science*. 310:858-863.

Zellner, S., M. Schifferer, and C. Behrends. 2021. Systematically defining selective autophagy receptor-specific cargo using autophagosome content profiling. *Mol Cell*. 81:1337-1354 e1338.

Zhuang, L., C. Li, F. Peng, E. Xue, W. Li, X. Sun, P. Chen, Q. Zhou, and L. Xue. 2023. Depletion of ESCRT ameliorates APP-induced AD-like symptoms in Drosophila. *J Cell Physiol*. 238:1567-1579.

Figure Legends

Fig 1. Tsg101 is required for release of EV cargoes from presynaptic terminals. (A-D) Representative confocal images from larvae expressing UAS-Tsg101-RNAi (Tsg101^{KD}) or a control RNAi either pan-neuronally (C380-GAL4) or in motor neurons (Vglut-GAL4) together with the following EV cargoes: **(A)** Syt4-GFP expressed from its endogenous locus, **(B)** UAS-driven Evi-GFP, **(C)** UAS-driven APP-GFP, **(D)** endogenous Neuroglian (Nrg, neuronal isoform Nrg180) detected by antibody. **(E-H)** Quantification of EV cargo puncta intensity.

All images show MaxIPs of muscle 6/7 segments A2 or A3. Scale bars are 5 μm. **(A-D)** Blue outlines represent the neuronal membrane as marked from an HRP mask; yellow line in **(A)** shows a 3.3 μm dilation of the HRP mask, representing the postsynaptic region. Arrows show examples of postsynaptic EVs. Data are represented as mean +/- s.e.m.; n represents NMJs. All intensity measurements are normalized to their respective controls. *p<0.05, ***p<0.001. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure S1: Characterization of EV structures upon Tsg101 depletion (associated with Figure 1). (A) Representative Airyscan images of larvae expressing UAS-Tsp42Ej-HA, and labeled with α-HA and α-Nrg. Scale bar is 5 μm. Yellow outline represents the neuronal membrane as marked from an HRP mask. **(B)** Representative 2D-STED images of muscle 6/7 labeled with α-GFP and α-Nrg antibodies. Scale bar is 2.5 μm. **(C)** Noise2Void denoised images and depiction of image regions used for quantification (left panel, scale bar is 2.5 μm). Pre: presynaptic; Post: Postsynaptic; BG: Background. Buffers (between double lines in the top left panel) generated by a 10% dilation of the presynaptic or postsynaptic area were used to eliminate signal that overlapped between regions. Boxes indicate zoomed areas in middle and right panels showing automated spot detections (green dots) and the presynaptic boundary (dotted line). **(D**) Quantification of APP-GFP and Nrg puncta number. Data are represented as mean +/- s.e.m.; n represents NMJs; ***p<0.001. **(E)** Cumulative distribution of Nrg and APP puncta diameter. Graph shows fraction of particles under the indicated size; numbers indicate mean and standard deviation of all detected puncta. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Fig 2. Multiple ESCRT components are required for release of EV cargoes from presynaptic terminals. (A) Representative confocal images of control, *Hrs* and *nwk* mutant larvae expressing Syt4-GFP from its endogenous locus. **(B)** Representative confocal images of control and *Hrs* mutant larvae expressing motor neuron (Vglut-GAL4)-driven UAS-Evi-GFP. **(C)** Representative confocal images of control and *Hrs* mutant larvae labeled with antibodies against endogenous Nrg. **(D-F)** Quantification of EV cargo puncta intensity. **(G)** Representative confocal images of larvae pan-neuronally expressing UAS-Shrub-RNAi (Shrub^{KD}) or a control RNAi. **(H-I)** Quantification of Syt4-GFP and Nrg puncta intensity. **(J)** Representative confocal images of larvae pan-neuronally expressing UAS-Vps4^{DN}. **(K-L)** Quantification of Syt4-GFP and Nrg puncta intensity.

All images show MaxIPs of muscle 6/7 segments A2 or A3. Scale bars are 5 μm. Outline represents the neuronal membrane as marked from an HRP mask. Data are represented as mean +/- s.e.m.; n represents NMJs. All fluorescence intensity values are normalized to their respective controls. *p<0.05, **p<0.01, ***p<0.001. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure 3: Loss of ESCRT causes compensatory autophagy of presynaptic cargoes. (A) Representative Airyscan images showing co-localization of EV cargoes Syt4-GFP or α-Nrg with endosomal markers α-Rab11, GFP-Rab5 (endogenous tag), or YFP-Rab7 (endogenous tag). Scale bars are 5 μm and outline represents the neuronal membrane as marked from an HRP mask. **(B)** Representative TEM images of boutons from muscle 6/7 from wild-type and neuronal Tsg101^{KD} larvae. Examples of autophagic vacuoles are marked with arrowheads, blue = autophagosome, magenta = autolysosome, and green = unclosed phagophore. Other notable features include Az = active zone, S = synaptic vesicles, M = mitochondria, SSR = subsynaptic reticulum. Scale bar is 400 nm. **(C)** Representative images of the EV cargo Nrg following motor neuron knockdown of *Atg1*. Scale bar is 5 μm. **(D)** Quantification of Nrg intensity from **(C)**, normalized to control. **(E)** Representative images from neuronal cell bodies in the ventral ganglion expressing motor neuron-driven GFP-mCherry-Atg8. Scale bar is 10 μm. Brightness/contrast are matched for each mutant with its paired

control (**see Table S3**). **(F-G)** Quantification of GFP-mCherry-Atg8 levels in **(F)** Tsg101[KD] and **(G)** *Hrs[D28]* mutant larvae**.**

Data are represented as mean +/- s.e.m.; n represents NMJs in **(C)** and animals in **(F-G)**. *p<0.05, **p<0.01, ***p<0.001. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure S2: Quantification of endosomal accumulation and autophagy controls (associated with Figure 3). (A-C) Quantification of co-localization of Nrg or Syt4 and Rab GTPases upon neuronal Tsg101[KD] (representative images in **Fig. 3A**). Mander's coefficient for the colocalization of Nrg and Rab5 **(A)**, Nrg and Rab7 **(B)**, Syt4 and Rab11 **(C)**, where M1 indicates the fraction of EV cargo in the Rab-positive thresholded area and M2 is the fraction of the Rab marker in the EV cargo-positive thresholded area. **(D-F)** Quantification of Rab compartment properties: **(D)** normalized Rab puncta intensity, **(E)** density of Rab puncta in the presynaptic compartment, and **(F)** average size of Rab puncta. **(G)** Representative confocal images of motor neuron cell bodies to validate that pan-neuronal Atg1 RNAi effectively blocks autophagic flux, assessed by GFP-mCherry-Atg8. **(H)** Representative confocal images of Nrg in muscle 6/7 NMJs **(I)** Quantification of Nrg intensity from **(H)**. **(J)** Colocalization of GFP and mCherry in cell bodies from motor neurons expressing GFP-mCherry-Atg8 (representative images in **Fig 3E**).

All scale bars = 5 μm. Data are represented as mean +/- s.e.m.; n represents NMJs in **(A-F, I)** and animals in **(J)**. Intensity measurements **(D, I)** are normalized to their respective controls. *p<0.05, **p<0.01, ***p<0.001. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure 4. Tsg101[KD] causes neuronal accumulation of EV cargoes (A) (Left) Representative confocal images of Syt4-GFP in a single slice through motor neuron cell bodies of the ventral ganglion. Scale bar is 10 μm. **(Right)** Quantification of total Syt4-GFP intensity in the brain. **(B) (Left)** Maximum intensity projection of axon segment proximal to the ventral ganglion. Scale bar is 10 μm. **(Right)** Quantification of total Syt4-GFP intensity in the axon. **(C)** Representative kymographs showing tracks of APP-GFP in the axon proximal to the ventral ganglion. Bottom panels show color coded traces. **(D)** Quantification of directionality of APP-GFP tracks. **(E)** Quantification of the velocity of retrograde and anterograde APP-GFP tracks upon neuronal Tsg101[KD]**.**

Data are represented as mean +/- s.e.m.; n represents animals. Intensity measurements **(A, B)** are normalized to their respective controls. *p<0.05, ***p<0.001. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure S3. Controls for axonal transport in Tsg101[KD] larvae (associated with Figure 4). (A) Representative kymographs showing tracks of Mito-GFP in axonal region proximal to the ventral ganglion, following photobleaching. Lower panels show color coded traces. **(B)** Percent of mitochondria tracks moving retrograde and anterograde. **(C)** Velocities of mitochondria tracks. **(D) (Top)** Representative images of the first frame of Mito-GFP videos. Scale bar = 10 μm. **(Bottom)** Quantification of Mito-GFP intensity.

Data are represented as mean +/- s.e.m.; n represents axons. Intensity measurements are normalized to their respective controls. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure 5. Tsg101[KD] phenocopies a subset of *evi* and *wg* synaptic morphology and signaling defects, while loss of Hrs has no effect. (A) Representative confocal images of muscle 6/7 NMJs labeled with α-HRP and α-BRP antibodies (left). Magnification of the yellow boxed area (right). HRP brightness was adjusted independently. Large image scale bar is 20 μm, small image scale bar is 5 μm. **(B)** Quantification of total bouton number (top) and active zone number (bottom) on muscle 6/7. **(C)** Representative confocal images of muscle 6/7 NMJ highlighting α-DLG pattern. Arrows indicate location of "feathery" DLG. Scale bar is 5 μm. **(D)** Representative confocal images of muscle 6/7 NMJ (abdominal segment A2) labeled with α-HRP and α-DLG antibodies. α-DLG and α-HRP signals were acquired in the linear range but adjusted independently and displayed near saturation to highlight DLG-negative ghost boutons, which are indicated with yellow arrows. **(E)** Quantification of baseline (i.e. unstimulated) ghost boutons. Top and bottom graphs represent independent experiments. **(F)** Single slices of muscle 6/7 nuclei labeled with α-LamC (nuclear periphery) and α-Fz2-C antibodies.

Dotted line represents LamC-defined nuclear boundary. Scale bars are 10µm. **(G)** Quantification of Fz2-C puncta per nucleus. Number of nuclei quantified are indicated in the bar graph.
A2 and A3 indicate the larval abdominal segment. Data are represented as mean +/- s.e.m.; n represents nuclei in **(G)** and NMJs in **(B-E)**. *p<0.05, **p<0.01, ***p<0.001. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure 6. Loss of ESCRT does not phenocopy *syt4* functional defects,. A) Representative confocal images from muscle 4 in mock and spaced K⁺-stimulated larvae. Arrows indicate examples of activity-dependent ghost boutons. Scale bar = 10 µm. **(B)** Quantification of ghost bouton numbers per NMJ. **(C)** Representative traces of mEJPs before (top trace) and after (bottom trace) high frequency stimulation (4 x 100 Hz) from control and Tsg101^{KD}. **(D)** Timecourse of mEJP frequency after stimulation. **(E)** Representative traces of mEJPs before (top trace) and after (bottom trace) high frequency stimulation (4 x 100 Hz) from control, *Hrs*^{D28}, and *syt4*^{BA1}. **(F)** Timecourse of mEJP frequency after stimulation.
Data are represented as mean +/- s.e.m.; n represents NMJs. *p<0.05, **p<0.01. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure S4. Additional controls showing presynaptic source of Syt4 and structural plasticity upon Tsg101^{KD} (associated with Figure 6). (A-B) Syt4 protein is derived from the presynaptic neuron. **(A)** Schematic for Tissue-Specific Tagging of Endogenous Proteins (T-STEP). Scissors indicate a Prescission protease cleavage site and * indicates stop codons. **(B)** Representative confocal images from muscle 6/7, showing Syt4^{TSTEP} expressed from its endogenous promoter, and switched from TagRFPt to GFP using either presynaptically (neuronal, C380-GAL4) or postsynaptically (muscle, C57-Gal4)-expressed recombinase (Rippase). Scale bar = 10 µm. **(C)** Representative confocal images from muscle 6/7 in spaced K⁺-stimulated larvae. Arrows indicate ghost boutons. Scale bar = 20 µm. **(D)** Quantification of ghost bouton numbers per NMJ.
Scale bars = 10 µm. Data is represented as mean +/- s.e.m.; n represents NMJs. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure 7. Neuronally derived EV cargoes are targeted for phagocytosis and are not detectable in the cytoplasm of recipient cells. (A) Schematic for DeGradFP system. **(B-C)** Representative images of Syt4-GFP with neuronal (C380-GAL4, **B**) or muscle (C57-GAL4, **C**) expressed DeGradFP. **(D)** Quantification of Syt4-GFP intensity from **(B)**. **(E)** Quantification of Syt4-GFP intensity from **(C)**. **(F)** Quantification of normalized presynaptic puncta number from **(B)** and **(C)**. **(G)** Representative confocal images of Syt4-GFP at muscle 4 NMJs following knockdown of Draper in different cell types. Outlines represent the neuronal membrane as marked from an HRP mask **(H)** Quantification of Syt4 puncta intensity.
All scale bars = 10 µm. Intensity measurements are normalized to their respective controls. Data are represented as mean +/- s.e.m.; n represents NMJs. **p<0.01, ***p<0.001. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Figure S5. Controls for DeGradFP and validation of Draper RNAi (associated with Figure 7). (A) Representative images of Dlg^{MiMIC} (a postsynaptically-localized GFP knock-in) with muscle-expressed DeGradFP. **(B)** Representative confocal images of muscle 4 NMJs labeled with α-Draper antibodies**. (C)** Quantification of α-Draper intensity at NMJs and axon bundles proximal to the NMJ upon Draper RNAi under the control of the indicated drivers. Axon bundles represent a combination of glial and neuronal signal; NMJs represent a combination of neuronal and muscle signal.
Scale bars are 20 µm. Intensity measurements are normalized to their respective controls. Data are represented as mean +/- s.e.m.; n represents NMJs. **p<0.01, ***p<0.001. **See Tables S1 and S3 for detailed genotypes and statistical analyses.**

Table S1: *Drosophila* mutants and genetic tools

Chromosome for transgene insertion indicated in roman numerals.

GAL4^{C380} (X)	(Budnik et al., 1996)	Flybase ID:FBti0016294
GAL4^{C155} (X)	(Lin and Goodman, 1994)	Flybase ID: FBti0002575
GAL4^{Vglut} (X)	{Daniels, 2008 #424}	Flybase ID: FBti0129146
GAL4^{C57} (III)	(Budnik et al., 1996)	Flybase ID: FBti0016293
GAL4^{repo} (III)	(Sepp et al., 2001)	Flybase ID: FBti0018692
GAL4^{Tub} (III)	(Lee and Luo, 1999)	Flybase ID: FBal0097158
w^{1118}	(Hazelrigg et al., 1984)	Flybase ID: FBal0018186
Syt4$^{EGFP-KI}$	(Walsh et al., 2021)	Flybase ID: FBal0368282
UAS-Evi-EGFP (II)	(Bartscherer et al., 2006)	Flybase ID: FBal0194740
UAS-APP-EGFP (II)	(Walsh et al., 2021)	Flybase ID: FBal0368284
pUASt-Tsp42Ej-3xHA (AttP1)	(Walsh et al., 2021)	Flybase ID: FBal0368283
GFP-Rab5-KI (II)	(Fabrowski et al., 2013)	Flybase ID: FBAl0288693
YFP-myc-Rab7-KI (III)	(Dunst et al., 2015)	Flybase ID: FBal0314198
UAS-Mito-HA-GFP (II)	(Rizzuto et al., 1995)	Flybase ID: FBti0040803
UAS-GFP-mCherry-Atg8 (II)	(Nezis et al., 2010)	Flybase ID: FBti0147141
UAS-mCherry-RNAi (III)	(Perkins et al., 2015)	Flybase ID: FBal0260847
UAS-Luciferase-RNAi (III)	(Perkins et al., 2015)	Flybase ID: FBti0143388
UAS-Tsg101-RNAi^{GLV21075} (III) (Tsg101^{KD})	(Perkins et al., 2015)	Flybase ID: FBti0144671
UAS-Shrub-RNAi^{HMS01767} (II) (Shrub^{KD})	(Perkins et al., 2015)	Flybase ID: FBti0149514
UAS-Vps4^{DN} (II)	(Rodahl et al., 2009)	Flybase ID: FBal0221338
Hrs^{D28}	(Lloyd et al., 2002)	Flybase ID: FBal0039519
evi^2	(Bartscherer et al., 2006)	Flybase ID: FBal0194741
Syt4^{BA1}	(Adolfsen et al., 2004)	Flybase ID: FBal0191284
Df(2L)Exel6277 (removes Hrs)		Flybase ID: FBab0037889
UAS-Atg1-RNAi^{GL00047} (III) (Atg1^{KD})		Flybase ID: FBti0144152
Atg2^{EP3697}	(Shen and Ganetzky, 2009)	Flybase ID: FBti0011778
Df(3L)Exel6091 (removes *Atg2*)		Flybase ID: FBab0038111
UAS-DegradGFP (II) (nSLimb-vhhGFP4)	(Caussinus et al., 2011)	Flybase ID: FBal0299649
Dlg1-MiMIC (X)	(Nagarkar-Jaiswal et al., 2015)	Flybase ID: FBti0168863
UAS-Rippase (III)	(Nern et al., 2011)	Flybase ID: FBst0055795
Syt4^{TSTEP}	(Walsh et al., 2021)	Flybase ID: FBti0216043
UAS-Draper-RNAi (II)	(MacDonald et al., 2006)	Flybase ID: FBal0258016

Table S2: Antibodies

REAGENT	SOURCE	IDENTIFIER	Concentration
Antibodies			
α-HRP-647	Jackson ImmunoResearch	123-605-021	1:250-1:500
α-Rab11	BD Biosciences	BD 610657	1:100
α-Dlg (mouse)	(Parnas et al., 2001), DSHB	4F3	1:500
α-Dlg (rabbit)	(Koh et al., 1999)		1:10000
α-Nrg	(Hortsch et al., 1990), DSHB	BP104	1:100
α-BRP	(Wagh et al., 2006), DSHB	nc82	1:100
α-Draper	DSHB	5D14	1:200
α-HA		HA.11	1:500
α-GFP	Abcam	Ab6556	1:500
α-Frizzled-C	(Mathew et al., 2005)		1:300
α-rabbit STAR RED	Aberrior	1002	1:250
α-mouse STAR ORANGE	Aberrior	1001	1:250

Supplemental Table References

Adolfsen, B., S. Saraswati, M. Yoshihara, and J.T. Littleton. 2004. Synaptotagmins are trafficked to distinct subcellular domains including the postsynaptic compartment. *J Cell Biol*. 166:249-260.

Bartscherer, K., N. Pelte, D. Ingelfinger, and M. Boutros. 2006. Secretion of Wnt ligands requires Evi, a conserved transmembrane protein. *Cell*. 125:523-533.

Budnik, V., Y.H. Koh, B. Guan, B. Hartmann, C. Hough, D. Woods, and M. Gorczyca. 1996. Regulation of synapse structure and function by the Drosophila tumor suppressor gene dlg. *Neuron*. 17:627-640.

Caussinus, E., O. Kanca, and M. Affolter. 2011. Fluorescent fusion protein knockout mediated by anti-GFP nanobody. *Nat Struct Mol Biol*. 19:117-121.

Dunst, S., T. Kazimiers, F. von Zadow, H. Jambor, A. Sagner, B. Brankatschk, A. Mahmoud, S. Spannl, P. Tomancak, S. Eaton, and M. Brankatschk. 2015. Endogenously tagged rab proteins: a resource to study membrane trafficking in Drosophila. *Dev Cell*. 33:351-365.

Fabrowski, P., A.S. Necakov, S. Mumbauer, E. Loeser, A. Reversi, S. Streichan, J.A. Briggs, and S. De Renzis. 2013. Tubular endocytosis drives remodelling of the apical surface during epithelial morphogenesis in Drosophila. *Nat Commun*. 4:2244.

Hazelrigg, T., R. Levis, and G.M. Rubin. 1984. Transformation of white locus DNA in drosophila: dosage compensation, zeste interaction, and position effects. *Cell*. 36:469-481.

Hortsch, M., A.J. Bieber, N.H. Patel, and C.S. Goodman. 1990. Differential splicing generates a nervous system-specific form of Drosophila neuroglian. *Neuron*. 4:697-709.

Koh, Y.H., E. Popova, U. Thomas, L.C. Griffith, and V. Budnik. 1999. Regulation of DLG localization at synapses by CaMKII-dependent phosphorylation. *Cell*. 98:353-363.

Lee, T., and L. Luo. 1999. Mosaic analysis with a repressible cell marker for studies of gene function in neuronal morphogenesis. *Neuron*. 22:451-461.

Lin, D.M., and C.S. Goodman. 1994. Ectopic and increased expression of Fasciclin II alters motoneuron growth cone guidance. *Neuron*. 13:507-523.

Lloyd, T.E., R. Atkinson, M.N. Wu, Y. Zhou, G. Pennetta, and H.J. Bellen. 2002. Hrs regulates endosome membrane invagination and tyrosine kinase receptor signaling in Drosophila. *Cell*. 108:261-269.

MacDonald, J.M., M.G. Beach, E. Porpiglia, A.E. Sheehan, R.J. Watts, and M.R. Freeman. 2006. The Drosophila cell corpse engulfment receptor Draper mediates glial clearance of severed axons. *Neuron*. 50:869-881.

Mathew, D., B. Ataman, J. Chen, Y. Zhang, S. Cumberledge, and V. Budnik. 2005. Wingless signaling at synapses is through cleavage and nuclear import of receptor DFrizzled2. *Science*. 310:1344-1347.

Nagarkar-Jaiswal, S., P.T. Lee, M.E. Campbell, K. Chen, S. Anguiano-Zarate, M.C. Gutierrez, T. Busby, W.W. Lin, Y. He, K.L. Schulze, B.W. Booth, M. Evans-Holm, K.J. Venken, R.W. Levis, A.C. Spradling, R.A. Hoskins, and H.J. Bellen. 2015. A library of MiMICs allows tagging of genes and reversible, spatial and temporal knockdown of proteins in Drosophila. *Elife*. 4.

Nern, A., B.D. Pfeiffer, K. Svoboda, and G.M. Rubin. 2011. Multiple new site-specific recombinases for use in manipulating animal genomes. *Proc Natl Acad Sci U S A*. 108:14198-14203.

Nezis, I.P., B.V. Shravage, A.P. Sagona, T. Lamark, G. Bjorkoy, T. Johansen, T.E. Rusten, A. Brech, E.H. Baehrecke, and H. Stenmark. 2010. Autophagic degradation of dBruce controls DNA fragmentation in nurse cells during late Drosophila melanogaster oogenesis. *J Cell Biol*. 190:523-531.

Parnas, D., A.P. Haghighi, R.D. Fetter, S.W. Kim, and C.S. Goodman. 2001. Regulation of postsynaptic structure and protein localization by the Rho-type guanine nucleotide exchange factor dPix. *Neuron*. 32:415-424.

Perkins, L.A., L. Holderbaum, R. Tao, Y. Hu, R. Sopko, K. McCall, D. Yang-Zhou, I. Flockhart, R. Binari, H.S. Shim, A. Miller, A. Housden, M. Foos, S. Randkelv, C. Kelley, P. Namgyal, C. Villalta, L.P. Liu, X. Jiang, Q. Huan-Huan, X. Wang, A. Fujiyama, A. Toyoda, K. Ayers, A. Blum, B. Czech, R. Neumuller, D. Yan, A. Cavallaro, K. Hibbard, D. Hall, L. Cooley, G.J. Hannon, R. Lehmann, A. Parks, S.E. Mohr, R. Ueda, S. Kondo, J.Q. Ni, and N. Perrimon. 2015. The Transgenic RNAi Project at Harvard Medical School: Resources and Validation. *Genetics*. 201:843-852.

Rizzuto, R., M. Brini, P. Pizzo, M. Murgia, and T. Pozzan. 1995. Chimeric green fluorescent protein as a tool for visualizing subcellular organelles in living cells. *Curr Biol*. 5:635-642.

Rodahl, L.M., K. Haglund, C. Sem-Jacobsen, F. Wendler, J.P. Vincent, K. Lindmo, T.E. Rusten, and H. Stenmark. 2009. Disruption of Vps4 and JNK function in Drosophila causes tumour growth. *PLoS One*. 4:e4354.

Sepp, K.J., J. Schulte, and V.J. Auld. 2001. Peripheral glia direct axon guidance across the CNS/PNS transition zone. *Dev Biol*. 238:47-63.

Shen, W., and B. Ganetzky. 2009. Autophagy promotes synapse development in Drosophila. *J Cell Biol*. 187:71-79.

Wagh, D.A., T.M. Rasse, E. Asan, A. Hofbauer, I. Schwenkert, H. Durrbeck, S. Buchner, M.C. Dabauvalle, M. Schmidt, G. Qin, C. Wichmann, R. Kittel, S.J. Sigrist, and E. Buchner. 2006. Bruchpilot, a protein with homology to ELKS/CAST, is required for structural integrity and function of synaptic active zones in Drosophila. *Neuron*. 49:833-844.

Walsh, R.B., E.C. Dresselhaus, A.N. Becalska, M.J. Zunitch, C.R. Blanchette, A.L. Scalera, T. Lemos, S.M. Lee, J. Apiki, S. Wang, B. Isaac, A. Yeh, K. Koles, and A.A. Rodal. 2021. Opposing functions for retromer and Rab11 in extracellular vesicle traffic at presynaptic terminals. *J Cell Biol*. 220:e202012034.

Table S3: Genotypes and Statistics by Dataset

Experiments were done at muscle 6/7 unless otherwise noted

All larvae are male unless otherwise noted

Analysis performed in Volocity unless otherwise noted

Presynaptic volume: α-HRP objects > 7µm^3

Postsynaptic volume: 3µm dilated from presynaptic volume

A: Sum intensity of signal in thresholded objects in presynaptic volume, normalized to presynaptic volume

B: Sum intensity of signal in thresholded objects in postsynaptic volume, normalized to presynaptic volume

n.s.: not significantly different; SDCM: spinning disk confocal microscopy; LSCM: laser scanning confocal microscopy; TEM: transmission electron microscopy; STED: Stimulated Emission Depletion Microscopy

Figure	Genotype/Conditions	N	Measurement	Statistical Test(s)
1A, E **SDCM**	Gal4^{C380}/Y;; Syt4-GFP/UAS-mcherry-RNAi -VALIUM20 Gal4^{C380}/Y;; Syt4-GFP/UAS-Tsg101-RNAi	8 larvae/ 21 NMJs 8 larvae/ 17 NMJs	Syt4-GFP intensity levels A, B	Pre: Unpaired t-test, p<0.05 Post: Mann-Whitney, p<0.0001
1B, F **SDCM**	Gal4^{Vglutx}/Y; UAS-Evi-GFP/+; UAS-mcherry-RNAi-VALIUM20/+ Gal4^{Vglutx}/Y; UAS-Evi-GFP/+; UAS-Tsg101-RNAi/+	7 larvae/ 21 NMJs 8 larvae/ 25 NMJs	Evi-GFP intensity levels A, B	Pre: Mann-Whitney, p<0.001 Post: Mann-Whitney, p<0.001
1C, D, G, H **SDCM**	Gal4^{Vglutx}/Y; UAS-APP-GFP/+; UAS-mcherry-RNAi-VALIUM20/+ Gal4^{Vglutx}/Y; UAS-APP-GFP/+; UAS-Tsg101-RNAi/+	6 larvae/ 15 NMJs 6 larvae/ 19 NMJs	APP-GFP intensity levels A, B	Pre: Mann-Whitney, p=0.56 n.s. Post: Unpaired t-test, p<0.05
			Nrg intensity levels A, B	Pre: Mann-Whitney, p=0.31 n.s. Post: Mann-Whitney, p<0.0001
S1A **Airyscan**	Gal4^{Vglutx}/w; UAS- Tsp42Ej-3xHA/+; UAS-mcherry-RNAi-VALIUM20/+ Gal4^{Vglutx}/w; UAS- Tsp42Ej-3xHA/+; UAS-Tsg101-RNAi/+ (Female)	6 larvae/ 17 NMJs 6 larvae/ 19 NMJs	No measurements	
S1B-E **STED**	Gal4^{Vglutx}/Y; UAS-APP-GFP/+; UAS-mcherry-RNAi-VALIUM20/+ Gal4^{Vglutx}/Y; UAS-APP-GFP/+; UAS-Tsg101-RNAi/+	6 larvae/ 19 NMJs 6 larvae/ 17 NMJs	Quantification of density of APP-GFP and Nrg puncta	Nrg puncta density: 2-way ANOVA with Sidak's multiple comparison's test, p<0.001 APP puncta density: 2-way ANOVA with Sidak's multiple comparison's test, p<0.001
			APP-GFP puncta diameter	No statistical comparison made

2A,D **SDCM**	Syt4-GFP	6 larvae /22 NMJs	Syt4-GFP intensity levels A, B	Pre: One-way ANOVA with Tukey's multiple comparisons test $p<0.001$ Post: Kruskal-Wallis with Dunn's multiple comparisons test $p<0.001$
	Hrs^{D28}/Df(2L)Exel6277; Syt4-GFP	7 larvae/ 24 NMJs		
	nwk^2; Syt4GFP	6 larvae/ 20 NMJs		
2B,E **SDCM**	Gal4^{Vglutx}/Y; UAS-Evi-GFP/+	7 larvae /24 NMJs	Evi-GFP intensity levels A, B	Pre: Mann-Whitney $p<0.001$ Post: Mann-Whitney $p<0.001$
	Gal4^{Vglutx}/Y; UAS-Evi-GFP, Hrs^{D28}/Df(2L)Exel6277	7 larvae/ 26 NMJs		
2C,F **SDCM**	w^{1118}	8 larvae/ 24 NMJs	Nrg intensity levels A, B	Pre: Mann-Whitney, $p<0.001$ Post**:** Mann-Whitney, $p<0.0001$
	Hrs^{D28}/Df(2L)Exel6277	8 larvae/ 26 NMJs		
2G-I **SDCM**	Gal4^{C380}/Y; Syt4-GFP/UAS-mcherry-RNAi-VALIUM20	6 larvae/ 22 NMJs	Syt4-GFP intensity levels A, B	Pre: Mann-Whitney, $p=0.68$ n.s. Post: Mann-Whitney, $p<0.001$
	Gal4^{C380}/Y; UAS-Shrub-RNAi/+ ; Syt4-GFP/+	6 larvae/ 18 NMJs		
			Nrg intensity levels A, B	Pre: Mann-Whitney, $p<0.01$ Post: Mann-Whitney, $p<0.001$
2J-L **SDCM**	Gal4^{C380}/Y;; Syt4-GFP/UAS-mcherry-RNAi- VALIUM20	8 larvae/ 25 NMJs	Syt4-GFP intensity levels A, B	Pre: Unpaired t-test, $p<0.001$ Post: Mann-Whitney, $p<0.0001$
	Gal4^{C380}/Y;; UAS-Vps4^{DN}/+ ; Syt4-GFP/+	8 larvae/ 27 NMJs (Female)		
			Nrg intensity levels A, B	Pre: Mann-Whitney, $p<0.0001$ Post: Mann-Whitney, $p<0.0001$
3A, S2A **Airyscan**	Gal4^{Vglutx}/Y; GFP-Rab5-KI/+	6 larvae/ 20 NMJs	Mander's coefficients (fraction of Nrg in GFP-Rab5 positive puncta and fraction of GFP-Rab5 in Nrg positive puncta)	M1: Mann-Whitney, $p<0.0001$ M2: Unpaired t-test, $p<0.0001$
	Gal4^{Vglutx}/Y; GFP-Rab5-KI/+; UAS-Tsg101-RNAi/+	6 larvae/ 17 NMJs		
3A, S2B **Airyscan**	Gal4^{Vglutx}/Y;; YFP-myc-Rab7KI/+	7 larvae/ 20 NMJs	Mander's coefficients (fraction of Nrg in YFP-Rab7 positive puncta and fraction of YFP-Rab7 in Nrg positive puncta)	M1: Mann-Whitney, $p<0.01$ M2: Unpaired t-test, $p<0.01$
	Gal4^{Vglutx}/Y;; YFP-myc-Rab7KI, UAS-Tsg101-RNAi/+	7 larvae/ 17 NMJs		

Figure	Genotype	Sample size	Measurement	Statistical test
3A, S2C Airyscan	Gal4^{C380}/Y;; Syt4-GFP/UAS-mcherry-RNAi- VALIUM20 Gal4^{C380}/Y;; Syt4GFP/UAS-Tsg101-RNAi	7 larvae/ 25 NMJs 9 larvae/ 25 NMJs	Mander's coefficients (fraction of Syt4-GFP in anti-Rab11 positive puncta and fraction of anti-Rab11 in Syt4-GFP positive puncta)	M1: Unpaired t-test, p<0.0001 M2: Mann-Whitney: p<0.0001
3A, S2D-F Rab11 Airyscan	Gal4^{C380}/Y;; Syt4-GFP/UAS-mcherry-RNAi- VALIUM20 Gal4^{C380}/Y;; Syt4GFP/UAS-Tsg101-RNAi	7 larvae/ 23 NMJs 9 larvae/ 25 NMJs	Rab11 intensity levels A	Unpaired t-test, p<0.001
			Number of anti-Rab11 positive puncta in HRP volume (puncta count/um^3)	Mann-Whitney, P>0.05 ns
			Average anti-Rab11 puncta size (total Rab11 volume um^3/puncta number)	Mann-Whitney, p<0.05
3A, S2D-F Rab5 Airyscan	Gal4^{Vglutx}/Y; GFP-Rab5-KI/+ Gal4^{Vglutx}/Y; GFP-Rab5-KI/+; UAS-Tsg101-RNAi/+	6 larvae/ 20 NMJs 6 larvae/ 17 NMJs	GFP-Rab5 intensity levels A	Mann-Whitney, p<0.05
			Number of GFP-Rab5 positive puncta in HRP volume (puncta count/um^3)	Unpaired t-test, p<0.05
			Average GFP-Rab5 puncta size (total Rab5 volume um^3/puncta number)	Unpaired t-test, p<0.001
3A, S2D-F Rab7 Airyscan	Gal4^{Vglutx}/Y;; YFP-myc-Rab7KI/+ Gal4^{Vglutx}/Y;; YFP-myc-Rab7KI, UAS-Tsg101-RNAi/+	7 larva/ 20 NMJs 7 larva/ 17 NMJs	YFP-Rab7 intensity levels A	Mann-Whitney, p=0.14 n.s.
			Number of YFP-Rab7 positive puncta in HRP volume (puncta count/um^3)	Mann-Whitney, p=0.73 n.s.
			Average YFP-Rab7 puncta size (total Rab7 volume um^3/puncta number)	Mann-Whitney, p=0.25 n.s.
3B TEM	Gal4^{Vglutx}/Y;; UAS-mcherry-RNAi- VALIUM20/+ Gal4^{Vglutx}/Y;; UAS-Tsg101RNAi/+	40 boutons from 3 larvae 56 boutons from 3 larvae	Fraction of images with three or more autophagic vacuoles	Fisher's exact test, p<0.0001
3C-D SDCM	Gal4^{Vglutx}/Y;; UAS-mcherry-RNAi- VALIUM20/+ Gal4^{Vglutx}/Y;; UAS-Atg1-RNAi/+	8 larvae/ 30 NMJs 8 larvae/ 22 NMJs	Nrg intensity levels A, B	Pre: Unpaired t-test, p<0.01 Post: Mann-Whitney, p<0.05

3E-G, S2J Airyscan	Gal4^{Vglutx}/Y; UAS-GFP-mCherry-Atg8/+; UAS-Luciferase-RNAi-VALIUM10/+ Gal4^{Vglutx}/Y; UAS-GFP-mCherry-Atg8/+; UAS-Tsg101-RNAi/+ Gal4^{Vglutx}/Y; UAS-GFP-mCherry-Atg8/+ Gal4^{Vglutx}/Y; UAS-GFP-mCherry-Atg8, Hrs^{D28}/Df(2L)Exel6277	8 larvae each 4 male, 4 female per genotype	(2E-G) Fraction of grouped cell bodies volume occupied by GFP signal or mCherry signal. (S2D) Mander's coefficient for fraction total mCherry signal in GFP-positive volume	GFP in Tsg101: Mann-Whitney, $p<0.001$ mCherry in Tsg101: Mann-Whitney, $p<0.001$ GFP in Hrs: Mann-Whitney, $p=0.35$ n.s. mCherry in Hrs: Unpaired t-test, $p<0.001$ Mander's Hrs: Mann-Whitney, $p=0.3$ n.s. Mander's Tsg101: Mann-Whitney, $p<0.001$
S2G SDCM	Gal4^{Vglutx}/Y; UAS-GFP-mCherry-Atg8/+; UAS-Luciferase-RNAi-VALIUM10/+ Gal4^{Vglutx}/Y; UAS-GFP-mCherry-Atg8/+; UAS-Atg1-RNAi/+	2 larvae 3 larvae	No measurements	NA
S2H-I SDCM	w¹¹¹⁸ Atg2^{EP3697}/Df(3L)Exel6091	8 larvae/ 18 NMJs 7 larvae/ 17 NMJs Mixed males and females	Nrg intensity levels A, B	Pre: Mann-Whitney, $p<0.01$ Post: Unpaired t-test, $p<0.001$
4A SDCM	Gal4^{C380}/Y;; Syt4-GFP/UAS-mcherryRNAi-VALIUM20 Gal4^{C380}/Y;; Syt4-GFP/UAS-Tsg101RNAi	10 larvae 8 larvae	Intensity measurement from an ROI containing 6-8 cell bodies in ImageJ	Unpaired t-test, $p<0.0001$
4B SDCM	Gal4^{C380}/Y;; Syt4-GFP/UAS-mcherryRNAi- VALIUM20 Gal4^{C380}/Y;; Syt4-GFP/UAS-Tsg101RNAi	10 larvae 9 larvae	Intensity measurement from HRP thresholded axon bundles proximal (within $\sim$100-300μm) to the ventral ganglion in ImageJ	Unpaired t-test, $p<0.05$
4C-E SDCM	Gal4^{Vglutx}/Y; UAS-APPGFP/+; UAS-mcherryRNAi-VALIUM20/+ Gal4^{Vglutx}/Y; UAS-APPGFP/+; UAS-Tsg101RNAi/+	12 larvae 12 larvae	Kymograph analysis, manual count of anterograde, retrograde, stationary, and complex moving APP-GFP particles in ImageJ	Unpaired t-test, Stationary: $p<0.001$ Retrograde: p0.48 n.s. Anterograde: $p=0.50$ Complex: $p=0.12$
			Kymograph analysis, calculation of the slope of particle tracks, including segments from complex tracks in ImageJ	Retrograde: Mann-Whitney, $p<0.05$ Antergrade: Unpaired t-test $p=0.68$ n.s.
S3A-D SDCM	Gal4^{Vglutx}/Y; UAS-Mito-GFP/+; UAS-mcherry-RNAi-VALIUM20/+ Gal4^{Vglutx}/Y; UAS-Mito-GFP/+; UAS-Tsg101-RNAi/+	11 larvae 11 larvae	Kymograph analysis, manual count of percentage of anterograde and retrograde mito-GFP particles in ImageJ	Retrograde: Unpaired t-test, $p=0.70$ n.s. Anterograde: Unpaired t-test, $p=0.70$ n.s.

			Kymograph analysis, calculation of the slope of particle tracks in ImageJ	Retrograde: Unpaired t-test, p=0.65 n.s. Anterograde: Unpaired t-test, p=0.36 n.s.
			Intensity measurement from first frame of videos of axon bundles proximal (within ~100-300μm) to the ventral ganglion in ImageJ	Unpaired t-test, p=0.79 n.s.
5A-B SDCM	Gal4^{Vglutx}/Y;; UAS-mcherryRNAi- VALIUM20/+ Gal4^{Vglutx}/Y;; UAS-Tsg101RNAi/+ evi^2/evi^2	7 larvae/14 A2 NMJs, 12 A3 NMJs 7 larvae/11 A2 NMJs, 12 A3 NMJs 7 larvae/8 A2 NMJs, 12 A3 NMJs	Active zone number on muscle 6/7	A2: One-way ANOVA with Tukey's multiple comparison's test, p<0.001 A3: One-way ANOVA with Tukey's multiple comparison's test, p<0.001
			Number of active zones per NMJ area on muscle 6/7	A2: Kruskal-Wallis with Dunn's multiple comparison's test, p<0.001 A3: One-way ANOVA with Tukey's multiple comparison's test, p<0.001
			Number of boutons on muscle 6/7	A2: One-way ANOVA with Tukey's multiple comparisons test, p<0.001 A3: One-way ANOVA with Tukey's multiple comparisons test, p<0.001
5C-E SDCM	Gal4^{Vglutx}/Y;; UAS-mcherryRNAi- VALIUM20/+ Gal4^{Vglutx}/Y;; UAS-Tsg101RNAi/+ evi^2/evi^2	26 larvae/ 87 NMJs 26 larvae/ 87 NMJs 29 larvae/ 91 NMJs (Pooled data from 3 independent experiments, mix of males and females)	Fraction of images displaying expanded "feathery" DLG pattern	Fishers exact test WT vs evi^2 p<0.0001 WT vs Tsg101^{KD} p=0.27 n.s.
			Ghost bouton number on muscle 6/7 (HRP-positive bouton with no postsynaptic Dlg)	A2: Kruskal-Wallis test with Dunn's multiple comparison's test, p<0.001 A3: Kruskal-Wallis test with Dunn's multiple comparison's test, p<0.05

	Hrs^{D28}/Df(2L)Exel6277 Gal4^{Vglut}/Y;; UAS-Tsg101RNAi/+	12 larvae/40 NMJs 10 larvae/32 NMJs	Ghost bouton number on muscle 6/7 (HRP-positive bouton with no postsynaptic Dlg)	A2: Mann-Whitney test, p=0.005 A3: Mann-Whitney test, p=0.68 n.s.
5F-G **LSCM**	y^1 , w^{1118} Gal4^{Vglut}/Y;; UAS-Tsg101RNAi/+ Hrs^{D28}/Df(2L)Exel6277	4 larvae, 16 muscles, 264 nuclei 4 larvae, 16 muscles, 303 nuclei 4 larvae, 16 muscles, 274 nuclei	Fz2-C puncta number per nucleus (muscles 6 and 7)	One-way ANOVA with Tukey's multiple comparisons test, p<0.001
6A, B **SDCM**	Gal4^{C380}/Y;; Syt4GFP/UAS-mcherry-RNAi-VALIUM20 Gal4^{C380}/Y;; Syt4-GFP/UAS-Tsg101RNAi	WT Mock: 6 larvae/ 22 NMJs WT Stim: 5 larvae/ 23 NMJs Tsg101 Mock: 6 larvae/ 25 NMJs Tsg101 Stim: 5 larvae/ 23 NMJs	Ghost bouton number on muscle 4 (HRP-positive bouton with no postsynaptic Dlg)	WT stim vs mock: Mann-Whitney, p<0.001 Tsg101 stim vs mock: Mann-Whitney, p<0.001 WT stim vs Tsg101 stim: Mann-Whitney, p=0.81 n.s.
6C, D **electro-physiology**	Gal4^{Vglut}/+;; UAS-mcherry-RNAi-VALIUM20/+ Gal4^{Vglut}/+;; UAS-Tsg101RNAi/+	9 larvae / 12 NMJs 8 larvae / 12 NMJs	Fold change in mEJP frequency	No statistical measurement
6E, F **electro-physiology**	w^{1118} Hrs^{D28}/Df(2L)Exel6277 $syt4^{BA1}$	9 larvae / 11 NMJs 12 larvae/12 NMJs 10 larvae/10 NMJs	Fold change in mEJP frequency	No statistical measurement
S4B **SDCM**	Gal4^{C380}/Y or +;; Syt4^{TSTEP} Gal4^{C380}/Y or +;; Syt4 ^{TSTEP}/UAS-Rippase, Syt4^{TSTEP} w;; Gal4^{C57}, Syt4 ^{TSTEP}/UAS-Rippase, Syt4^{TSTEP}	No quantification	No measurements	NA
S4C, D **LSCM**	Gal4^{Vglut}/+;; UAS-mcherry-RNAi-VALIUM20/+ Gal4^{Vglut}/+;; UAS-Tsg101-RNAi/+	7 larvae / 13 NMJs 7 larvae / 13 NMJs	Ghost bouton number on muscle 6/7 (HRP-positive bouton with no postsynaptic Dlg)	Unpaired t-test, p=0.65 n.s.
7B, D **SDCM**	Gal4^{C380}/Y; Syt4-GFP Gal4^{C380}/Y; UAS-DegradGFP, Syt4-GFP	29 NMJ branches 25 NMJ branches	Syt4-GFP intensity levels A, B	Pre: Unpaired t-test, p<0.001 Post: Unpaired t-test, p<0.001
7C, E **SDCM**	Syt4-GFP UAS-DegradGFP; Gal4^{C57}, Syt4-GFP	8 larvae/ 33 NMJ branches 8 larvae/ 30 NMJ branches	Syt4-GFP intensity levels A, B	Pre: Mann-Whitney, p=0.59 n.s. Post: Mann-Whitney, p=0.56 n.s.

7F **SDCM**	Genotypes from 7B and 7C	N from 4F and 4G	Number of Syt4-GFP positive puncta in HRP volume (puncta count/um^3)	C380: Unpaired t-test, p<0.001 C57: Mann-Whitney, p=0.84 n.s.
7G-H **SDCM**	UAS-DraperRNAi/+; Syt4-GFP/+ GAL4^{C155}; UAS-DraperRNAi/+ Syt4-GFP UAS-DraperRNAi/+ Syt4-GFP/GAL4^{C57} UAS-DraperRNAi/+ Syt4-GFP/Gal4^{repo} UAS-DraperRNAi/+ Syt4-GFP/GAL4^{tub}	6 larvae/ 26 NMJs 6 larvae/ 18 NMJs 6 larvae/ 26 NMJs 6 larvae/ 25 NMJs 6 larvae/ 28 NMJs	Syt4-GFP intensity levels A, B	Pre: Kruskal-Wallis P<0.001 Post: Kruskal-Wallis P<0.001
S5A **SDCM**	Dlg1-MiMIC/+; CyOGFP/+; Gal4^{C57}, Syt4-GFP/TM6 Dlg1-MiMIC/+; UAS-DegradGFP/+; Gal4^{C57}, Syt4-GFP/ Gal4^{C57}	4 larvae/ 14 NMJs 4 larvae/ 13 NMJs	No measurements	NA
S5B-C **SDCM**	w^{1118} GAL4^{C155}; UAS-DraperRNAi/+ UAS-DraperRNAi/+ GAL4^{C57}/+ UAS-DraperRNAi/+ GAL4^{repo}/+	6 larvae/ 27 NMJs 6 larvae/ 25 NMJs 6 larvae/ 29 NMJs 6 larvae/ 31 NMJs	Draper mean intensity	NMJ: One-way ANOVA p<0.001 Axon: Kruskal-Wallis p<0.001

A Syt4-GFP **B** Evi-GFP **C** APP-GFP **D** α-Nrg

Control

Tsg101^KD

E Pre Post
Syt4-GFP intensity/HRP Vol
* ***
Control Tsg101^KD Control Tsg101^KD

F Pre Post
Evi-GFP intensity/HRP Vol
*** ***
Control Tsg101^KD Control Tsg101^KD

G Pre Post
APP-GFP intensity/HRP Vol
n.s. *
Control Tsg101^KD Control Tsg101^KD

H Pre Post
α–Nrg intensity/HRP Vol
n.s. ***
Control Tsg101^KD Control Tsg101^KD

A DeGradFP-mediated degradation

B C380-GAL4 | + UAS-DeGradFP
Syt4-GFP
α-HRP
Merge

C C57-GAL4 | + UAS-DeGradFP
Syt4-GFP
α-HRP
Merge

D Neuron (C380-GAL4)
Normalized Syt4-GFP intensity/HRP vol
Pre *** Post ***
UAS-DeGradFP

E Muscle (C57-GAL4)
Normalized Syt4-GFP intensity/HRP vol
Pre n.s. Post n.s.
UAS-DeGradFP

F Neuron (C380-GAL4) Muscle (C57-GAL4)
Postsynaptic Syt4-GFP puncta #/HRP vol
***
UAS-DeGradFP

G Draper RNAi α-HRP Syt4-GFP
Control (No Driver)
Glia (repo-Gal4)
Muscle (C57-Gal4)
Neuron (C155-Gal4)
Ubiquitous (Tubulin-Gal4)

H Syt4-GFP intensity/HRP Vol
Pre *** *** *** *
Post *** * ***
No driver Glia Muscle Neuron Ubiquitous
Draper RNAi

A

SYT4 | Tag-RFPt | 3'UTR | selection marker | | GFP | 3'UTR
Rippase recombinae recognition site

Locus after tissue-specific Rippase expression

SYT4 | | GFP | 3'UTR

B

| | Syt4^TSTEP | +C380-GAL4, UAS-R (neuronal Rippase) | +BG57-GAL4, UAS-R (muscle Rippase) |

Syt4-GFP

Syt4-TagRFPt

merge

C After stimulation

Control

Tsg101^KD

α-HRP α-DLG

D

Number of ghost boutons (muscle 6/7)

n.s.

Control   Tsg101^KD

A

DLG^{MIMIC}

α-HRP α-DLG

+ C57-GAL4, UAS-DeGradFP

B

α-HRP α-Draper

α-Draper

Control

Draper RNAi

Glia (repo-Gal4)

Muscle (C57-Gal4)

Neuron (C155-Gal4)

C

NMJ

Axon bundle