Blood and cerebellar abundance of *ATXN3* splice variants in spinocerebellar ataxia type 3/Machado-Joseph disease

Mafalda Raposo, PhD[1,2*], Jeannette Hübener-Schmid, PhD[3,4*], Rebecca Tagett, MSc[5], Ana F. Ferreira, MSc[2], Ana Rosa Vieira Melo, MSc[2], João Vasconcelos, MD[6], Paula Pires, MD[7], Teresa Kay, MD, PhD[8], Hector Garcia-Moreno, MD[9,10], Paola Giunti, MD, PhD[9,10], Magda M. Santana, PhD[11,12,13], Luis Pereira de Almeida, PhD[11,12,14], Jon Infante, MD, PhD[15], Bart P. van de Warrenburg, MD, PhD[16], Jeroen J. de Vries, MD[17], Jennifer Faber, MD[18,19], Thomas Klockgether, MD[18,19], Nicolas Casadei, PhD[3,20], Jakob Admard, MSc[3,20], Ludger Schöls, MD[21,22], Olaf Riess, MD[3,4,20], European Spinocerebellar ataxia type 3/Machado-Joseph disease Initiative (ESMI) study group, Maria do Carmo Costa, PhD[23*], Manuela Lima, PhD[2*]

1. Instituto de Biologia Molecular e Celular (IBMC), Instituto de Investigação e Inovação em Saúde (i3S), Universidade do Porto, Porto, Portugal.
2. Faculdade de Ciências e Tecnologia, Universidade dos Açores, Ponta Delgada, Portugal.
3. Institute of Medical Genetics and Applied Genomics, University of Tübingen, Tübingen, Germany.
4. Centre for Rare Diseases, University of Tübingen, Tübingen, Germany.
5. Bioinformatics Core, Michigan Medicine, University of Michigan, Ann Arbor, MI, USA
6. Serviço de Neurologia, Hospital do Divino Espírito Santo, Ponta Delgada, Portugal.
7. Serviço de Neurologia, Hospital do Santo Espírito da Ilha Terceira, Angra do Heroísmo, Portugal.
8. Serviço de Genética Clínica, Hospital D. Estefânia, Lisboa, Portugal.
9. Ataxia Centre, Department of Clinical and Movement Neurosciences, UCL Queen Square Institute of Neurology, University College London, London, UK.
10. Department of Neurogenetics, National Hospital for Neurology and Neurosurgery, University College London Hospitals NHS Foundation Trust, London, UK.
11. Center for Neuroscience and Cell Biology (CNC), University of Coimbra, Coimbra, 3000-075, Portugal.
12. Center for Innovative Biomedicine and Biotechnology (CIBB), University of Coimbra, Coimbra, Portugal.
13. Institute for Interdisciplinary Research, University of Coimbra (IIIUC), Coimbra, Portugal.
14. Faculty of Pharmacy, University of Coimbra (FFUC), Coimbra, Portugal
15. Neurology Service, University Hospital Marqués de Valdecilla-IDIVAL, Universidad de Cantabria, Centro de Investigación en Red de Enfermedades Neurodegenerativas (CIBERNED), Santander, Spain.
16. Radboud University Medical Centre, Donders Institute for Brain, Cognition and Behaviour, Department of Neurology, Nijmegen, The Netherlands.
17. Department of Neurology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands.
18. Department of Neurology, University Hospital Bonn, Bonn, Germany.
19. German Center for Neurodegenerative Diseases (DZNE), Bonn, Germany.
20. NGS Competence Center Tübingen, Tübingen, Germany.
21. Department for Neurodegenerative Diseases, Hertie-Institute for Clinical Brain Research and Center for Neurology, University of Tübingen, Germany.
22. German Center for Neurodegenerative Diseases (DZNE), Tübingen, Germany.
23. Department of Neurology, Michigan Medicine, University of Michigan, Ann Arbor, MI, USA.

*Corresponding authors:

Mafalda Raposo

Instituto de Investigação e Inovação em Saúde (i3S)

Rua Alfredo Allen, 208

4200-135 Porto, Portugal

Phone: +351 226 074 900

E-mail address: msraposo@i3s.up.pt

Jeannette Hübener-Schmid

Institute of Medical Genetics and Applied Genomics

Nägelestraße 5

72074 Tübingen, Germany

Phone: +49 7071 29 72276

E-mail address: jeannette.huebener@med.uni-tuebingen.de

Maria do Carmo Costa, Ph.D.

Department of Neurology, University of Michigan

A. Alfred Taubman Biomedical Sciences Research Building, Room 4027

109 Zina Pitcher Place, Ann Arbor, MI 48109-2200, USA

Phone: +1 734-615-5632

E-mail address: mariadoc@med.umich.edu

Manuela Lima

Faculdade de Ciências e Tecnologia, Universidade dos Açores

Rua Mãe de Deus, N.13

9500-321 Ponta Delgada, Portugal

Phone: +351 296 650 118

E-mail address: maria.mm.lima@uac.pt

Word count: 3981 words

Running title: *ATXN3* splice variants in SCA3/MJD

Financial Disclosures/Conflict of Interest

TK is receiving research support from the Bundesministerium für Bildung und Forschung (BMBF),

the National Institutes of Health (NIH) and Servier. Within the last 24 months, he has received

consulting fees from Biogen, UCB and Vico Therapeutics. BvdW is supported by grants from

ZonMw, NOW, Hersenstichting, Gossweiler Foundation, and Radboud university medical center;

he has served on the scientific advisory boards or steering committees of uniQure, VICO

Therapeutics, and Servier. LS is receiving research support from the European Commission, the

Bundesministerium für Bildung und Forschung (BMBF) and the Bundesministerium für

Gesundheit (BMG) as well as the Deutsche Forschungsgemeinschaft (DFG), Servier and Vigil

Neuroscience. Within the last 24 months, he has received consulting fees from Vico

Therapeutics.

The remaining authors declare that they have nothing to report.

Funding Sources

This work is an outcome of ESMI, an EU Joint Programme - Neurodegenerative Disease Research

(JPND) project. The ESMI project was supported through the following funding organisations

under the aegis of JPND: Germany, Federal Ministry of Education and Research (BMBF; funding

codes 01ED1602A/B); Netherlands, The Netherlands Organisation for Health Research and

Development; Portugal, Fundação para a Ciência e a Tecnologia (FCT; funding codes

JPCOFUND/0002/2015); United Kingdom, Medical Research Council. This project has received

funding from the European Union's Horizon 2020 research and innovation program under grant

agreement No 643417. MR is supported by FCT (CEECIND/03018/2018). AFF and ARVM received

PhD fellowships from the FCT (SFRH/BD/121101/2016 and SFRH/BD/129547/2017,

respectively). Fundo Regional para a Ciência e Tecnologia (FRCT, Governo Regional dos Açores)

is currently supporting ESMI in the Azores, under the PRO-SCIENTIA program. NGS sequencing

methods were performed with the support of the DFG-funded NGS Competence Center

Tübingen (INST 37/1049-1). Bioinformatic analysis was funded by Fundo Regional para a Ciência

e Tecnologia (FRCT, Governo Regional dos Açores), under the PRO-SCIENTIA program (ML) and

by discretionary funds from the University of Michigan (MCC). MCC is supported by the Heeringa

Ataxia Research Fund (University of Michigan). Several authors of this publication are members

of the European Reference Network for Rare Neurological Diseases - Project ID No 739510.

ABSTRACT

Background: Spinocerebellar ataxia type 3 (SCA3)/Machado-Joseph disease (MJD) is an

autosomal dominant polyglutamine disease. SCA3/MJD causative gene, *ATXN3*, is known to

undergo alternative splicing (AS) and 54 transcripts are currently annotated. Differences in the

toxicity of ataxin-3 protein isoforms, harbouring on its C-terminus two or three ubiquitin

interacting motifs (UIMs), were previously uncovered, raising the hypothesis that specific

ATXN3 splice variants play key roles in promoting the selective toxicity displayed in SCA3/MJD.

Methods: Using RNA-seq datasets we identified and determined the abundance of annotated

ATXN3 transcripts in blood (n=60) and cerebellum (n=12) of SCA3/MJD subjects and controls.

Results: Globally, the number and the abundance of individual *ATXN3* transcripts were higher

in the cerebellum than in the blood. While the most abundant transcript in the cerebellum was

a protein with a coding sequence not defined of unknown function (*ATXN3*-208), the transcript

with the highest abundance in blood was the reference transcript (*ATXN3*-251) which

translates into an ataxin-3 isoform harboring three UIMs. Noteworthy, the abundance of

ATXN3-251 and *ATXN3*-214, two out of the four transcripts that encode full-length ataxin-3

protein isoforms but differ in the C-terminus were strongly related with tissue expression

specificity: *ATXN3-251* (3UIM) was expressed in blood 50-fold more than in cerebellum,

whereas *ATXN3*-214 (2UIM) was expressed in the cerebellum 20-fold more than in blood.

Conclusions: These findings provide new insights into the elucidation of *ATXN3* AS in different

tissues, contributing for a better understanding of SCA3/MJD pathogenesis and providing

information for the development of future effective *ATXN3* mRNA-lowering therapies.

KEYWORDS: ataxin-3, polyQ diseases, neurodegenerative disease, RNA-seq, mRNA, alternative

splicing

INTRODUCTION

Alternative splicing (AS) is a fundamental step in the regulation of eukaryotic gene expression through which introns are excised and exons are ligated together to form the messenger RNA (mRNA) (Vuong *et al*, 2016). Multiple evidence indicates that genetic variation in *cis*-acting splicing sequence elements or in *trans*-acting splicing factors plays a relevant role in patients with neurodegenerative conditions (reviewed in (Daguenet *et al*, 2015; Nik & Bowman, 2019)). Importantly, *post-mortem* brains of Huntington disease (HD) patients and of mouse models show aberrant splicing of the HD causative gene (huntingtin, *HTT*) itself, leading to the production of high levels of a splice variant only containing exon 1 that encodes a highly aggregation-prone protein and thus contributes to disease pathogenesis (Mangiarini *et al*, 1996; Sathasivam *et al*, 2013; Barbaro *et al*, 2015; Neueder *et al*, 2017). Currently, several pathologies triggered by splicing defects, including neurodegenerative diseases, can be treated using different types of RNA molecules, such as antisense oligonucleotide (ASO) or small interference RNA (siRNA), either by reducing the levels of the defective protein or by producing the corrected protein (reviewed in (Dhuri *et al*, 2020; Zhu *et al*, 2022)). In fact, ASO-mediated therapies are currently available for some common and rare diseases, including neurological diseases such as spinal muscular atrophy and Duchenne muscular dystrophy (reviewed in (Zhu *et al*, 2022)).

SCA3/MJD is an autosomal dominant neurodegenerative polyglutamine (polyQ) disorder and the most common dominant ataxia worldwide (Sequeiros *et al*, 2012). SCA3/MJD is caused by an abnormal polyQ-encoding CAG repeat motif, consensually containing more than 60 trinucleotides within exon 10 of the *ATXN3* gene (Takiyama *et al*, 1993; Kawaguchi *et al*, 1994; Maciel *et al*, 2001). The *ATXN3* gene (ENSG00000066427) originates 54 transcripts (Martin *et al*, 2023) in the GRCh38.p13 human genome assembly including the canonical transcript (ENST00000644486/*ATXN3*-251) containing 11 exons and spanning a genomic region about 48 kb (Ichikawa *et al*, 2001), encoding the ataxin-3 canonical protein isoform (P54252-2, UniProt).

Ataxin-3 is a ubiquitously expressed deubiquitinating enzyme (*e.g.*, (Burnett *et al*, 2003))

harbouring on its N-terminus the globular Josephin domain (JD) that includes the catalytic

residues, and on its C-terminus a protein segment of undetermined structure containing the

polyQ tract and two or three ubiquitin interacting motifs (UIM) (Goto *et al*, 1997; Masino *et al*,

2003). Aggregation of the ataxin-3 protein is a hallmark of SCA3/MJD (reviewed in (Costa &

Paulson, 2012)) and occurs not only in brain regions with neuronal loss, such as the cerebellar

dentate nucleus and the motor and pontine nuclei of the brainstem, but also in brain areas

spared of neuronal destruction, like the cerebellar cortex (Koeppen, 2018).

Despite nearly 30 years of research following the identification of *ATXN3* as the causative gene

of SCA3/MJD (Takiyama *et al*, 1993; Kawaguchi *et al*, 1994), the characterization of the

transcriptional species of *ATXN3* remains very incomplete. While *ATXN3* is known to undergo

AS (Kawaguchi *et al*, 1994; Goto *et al*, 1997; Ichikawa *et al*, 2001; Harris *et al*, 2010;

Bettencourt *et al*, 2010), the extent and regulation of AS, the relative abundance of alternative

transcripts in health and disease conditions, the existence of transcript tissue/cell-specificity,

and the impact of AS in SCA3/MJD neurotoxicity remain poorly understood. To the best of our

knowledge, at least 24 *ATXN3* transcripts, likely to be translated into different isoforms of

ataxin-3, have been detected in the human brain and/or peripheral tissues including the blood

(Kawaguchi *et al*, 1994; Goto *et al*, 1997; Ichikawa *et al*, 2001; Harris *et al*, 2010; Bettencourt *et*

al, 2010), although validation of a large number of such transcripts has not been conducted. *In*

silico analysis performed to infer the functional impact of the several putative ataxin-3 protein

isoforms showed that some variants were predicted to be "protective" (lacking the CAG tract),

whereas others were foreseen to show increased toxicity (Bettencourt *et al*, 2010). Among the

ATXN3 transcripts which are predicted to be translated, only two transcripts encoding ataxin-3

isoforms that differ on the C-terminal containing either two (2UIM) or three UIMs (3UIM) have

been mostly studied so far (Harris *et al*, 2010; Johnson *et al*, 2019; Weishäupl *et al*, 2019).

These studies using cellular, mouse or *Drosophila* models and *post-mortem* human brain

tissues from healthy individuals, though controversial, uncovered differences in the toxicity of

these two ataxin-3 protein isoforms by mainly revealing a faster degradation rate but higher

aggregation propensity of the 2UIM isoform compared to the 3UIM (Harris *et al*, 2010;

Johnson *et al*, 2019; Weishäupl *et al*, 2019). The fact that these two ataxin-3 isoforms seem to

differentially impact SCA3/MJD pathogenesis, raises the possibility that other species

generated by *ATXN3* AS play key roles in promoting the selective neuronal toxicity displayed by

SCA3/MJD patients.

Several phase I and II clinical trials are ongoing for polyQ disorders aiming to reduce the levels

of disease-related transcripts/proteins using ASO-based drugs, but none was yet been

approved for clinical use (reviewed in (Helm *et al*, 2022)). Non-allele-specific as well as allele-

specific ASO-mediated reduction of *ATXN3* have been successfully tested *in vivo* and *in vitro*

models of MJD (McLoughlin *et al*, 2018; Moore *et al*, 2019, 2017; Hauser *et al*, 2022).

Noteworthy, the pharmacokinetics and safety study of a non-selective ASO for SCA3/MJD is

currently ongoing (NCT05160558). The identification of specific *ATXN3* transcripts, whose

expression needs to be blocked is, therefore, paramount to unveil better strategies in future

ASO-based therapies of SCA3/MJD.

Hence, the goal of this study was to profile annotated *ATXN3* AS transcripts in blood and

cerebellum of mutation carriers of SCA3/MJD and healthy individuals using RNA sequencing

(RNA-seq) data.

MATERIAL AND METHODS

<u>Samples and RNA-seq datasets</u>

Two RNA-seq datasets, generated from a total of 72 samples, namely 60 samples from whole

blood and 12 samples from *post-mortem* cerebellum (cerebellar lobules) were used in this

study: i) a RNA-seq dataset from whole blood samples of 40 SCA3/MJD subjects (30 patients

and 10 pre-ataxic) and 20 control individuals (Raposo *et al*, 2023), obtained in the scope of the

European spinocerebellar ataxia type 3/Machado-Joseph Initiative (ESMI); and ii) RNA-seq

dataset from *post-mortem* cerebellar samples of six SCA3/MJD patients and six controls

publicly available from the work of Haas and colleagues (Haas *et al*, 2022). Detailed

information about the blood and cerebellum samples from SCA3/MJD subjects (and

corresponding healthy controls), is provided in supplementary Table 1. Briefly, RNA extraction

and cDNA library preparation was performed according to manufacturer's instructions, as

described elsewhere (Haas *et al*, 2022; Raposo *et al*, 2023). Libraries from blood samples were

sequenced as paired end 100bp reads on an Illumina NovaSeq6000 (Illumina), whereas

libraries from cerebellum samples were sequenced as paired-end 68 bp reads on a HiSeq2000

(Illumina), both next-generation sequencing (NGS) platforms with limited read length (short-

reads).

Ataxin-3 protein levels in plasma (n=28) and in cerebrospinal fluid (CSF, n=8) from a subset of

the total 72 SCA3/MJD subjects were available from previous studies(Gonsior *et al*, 2021;

Hübener-Schmid *et al*, 2021). Briefly, total full-length ataxin-3 and/or mutant ataxin-3 were

measured using a time-resolved fluorescence energy transfer (TR-FRET)-based immunoassay

(Gonsior *et al*, 2021) and a single molecule counting (SMC) immunoassay (Hübener-Schmid *et*

al, 2021).

The study was approved by local ethics committees of all participating centers; all subjects

provided written informed consent.

<u>Bioinformatic analyses of *ATXN3* splicing</u>

The reads were trimmed using Cutadapt v2.3. Next, reads were mapped to the reference

genome GRCh38 (Ensembl), using STAR v2.7.8a (Dobin *et al*, 2013) and assigned count

estimates to genes with RSEM v1.3.3(Li & Dewey, 2011). Alignment options followed ENCODE

standards for RNA-seq (Dobin *et al*, 2013). FastQC (v0.11.8) (Andrews, 2010) was run on .bam

files in a post-alignment step, including both aligned and un-aligned reads, to ensure data

quality. From the raw reads, TPM was calculated according to Zhao and colleagues (Zhao *et al*,

2020), considering the denominator as the sum over all samples (brain and blood).

<u>Ensembl classification biotype</u>

ATXN3 transcripts were clustered according to the Ensembl classification biotype (Ensembl

release 108 – Jan 2023 (Martin *et al*, 2023)) as: (i) protein coding (a transcript that contains an

open reading frame), (ii) nonsense mediated decay (NMD; a transcript with a premature stop

codon, predicted as most likely to be subjected to targeted degradation because the in-frame

termination codon is found more than 50 bp upstream of the final splice junction), (iii) protein

coding sequence (CDS) not defined (a transcript of a protein coding gene for which a CDS has

not yet been defined), and (iv) retained intron (a transcript that lacks an open reading frame

believed to contain intronic sequence relative to other, coding, transcripts of the same gene)

which is included in the major category of long non-coding RNA transcripts. Additionally, using

the Ensembl annotation for the *ATXN3* gene (Martin *et al*, 2023), *ATXN3* transcripts were

clustered according to the presence or absence of the CAG tract, inferred from the presence or

absence of exon 10 where the CAG repeat is located.

<u>Statistical analyses</u>

To increase the sample size and the power of statistical tests, *ATXN3* transcripts were labelled

according to its frequency by sample. Transcripts not detected in at least half of our set of

samples (i.e., 5 or less samples from cerebellum and 29 or less samples for blood) were not

included in statistical analyses (Suppl. Table 2, Fig. S1). Transcripts levels were compared

between biological groups (pre-ataxic *versus* controls and patients *versus* controls) and

between tissues (cerebellum *versus* blood), and statistical differences were determined by the

Mann-Whitney U test. Among the blood samples from controls (n=20), a subset (n=10) was

paired-matched by age and sex to pre-ataxic (CTRL-PA) and used for group comparisons (pre-

ataxic *versus* CTRL-PA). The significance of the association between absence/presence of

transcripts within each tissue by biological group (contingency table) was tested using a

Fisher's exact test. In SCA3/MJD subjects' group, the relationship between transcript and

protein levels was determined using a Spearman rank correlation test. The ROUT method

(Q=1%) was used to exclude outliers previously to correlation test. Statistical analyses were

performed in GraphPad Prism version 8.0.1 for Windows (GraphPad Software, San Diego,

California USA). The significance level of all tests was set to 5%. Graphic bars are shown as

median ± 95% CI (confidence interval).

<u>Availability of data and materials</u>

Data supporting the findings of this study are available in this manuscript and in

supplementary material.

RESULTS

The abundance and the frequency of annotated *ATXN3* transcripts are higher in cerebellum

than in blood regardless of the disease status

Data from RNA-seq of whole blood (n=60) and *post-mortem* cerebellum (n=12) from

SCA3/MJD subjects and age- and sex-matched controls was used to identify the currently

annotated 54 *ATXN3* transcripts and quantify their abundance (Fig. 1). All annotated

transcripts were identified in our study. According to Ensembl annotation (Martin *et al*, 2023),

these 54 transcripts can be classified into four main biotypes: protein-coding (n=16), nonsense

mediated decay (NMD; n=19), protein CDS not defined (n=16) and retained intron (long

noncoding) (n=3) transcripts (Fig. 1). In the 16 protein-coding transcripts group, some

structural-related particularities are highlighted (Fig. 1): four transcripts encode full-length

ataxin-3 protein isoforms (*ATXN3*-214, *ATXN3*-215, *ATXN3*-227, *ATXN3*-251), including an

intact Josephin domain (JD), whereas the remaining 12 transcripts might encode proteins with

variations of the N-terminus (*ATXN3*-201, *ATXN3*-203, *ATXN3*-204, *ATXN3*-206, *ATXN3*-207,

ATXN3-209, *ATXN3*-213, *ATXN3*-219, *ATXN3*-228, *ATXN3*-231, *ATXN3*-235, *ATXN3*-246). Nine

transcripts lack exon 11 and, therefore, translate ataxin-3 isoforms only with 2UIM (*ATXN3*-

209, *ATXN3*-213, *ATXN3*-214, *ATXN3*-219, *ATXN3*-227, *ATXN3*-228, *ATXN3*-231, *ATXN3*-235,

ATXN3-246), while seven transcripts retain exon 11 (*ATXN3*-201, *ATXN3*-203, *ATXN3*-204,

ATXN3-206, *ATXN3*-207, *ATXN3*-215, *ATXN3*-251), and thus encode 3UIMs isoforms. Finally,

and although the annotation of the 3' untranslated (UTR) is incomplete, two *ATXN3* transcripts

seem to lack exon 10, which contains the polyQ-encoding tract (*ATXN3*-209, *ATXN3*-213) (Fig.

1).

To explore putative tissue-specific patterns of *ATXN3* AS regardless of the disease status, the

frequency of each annotated *ATXN3* transcript detected in samples from SCA3/MJD subjects

and controls was determined within each tissue (blood or cerebellum). Expression levels of

ATXN3 transcripts ranged from 0.01 (blood) to 50.49 (cerebellum) transcripts per million

(TPM). Globally, cerebellum samples displayed higher expression levels than blood samples

(Fig. 2; Suppl. Table 2). The biotype with more abundant transcripts in blood and cerebellum is

the retained intron and the noncoding, respectively (Fig. 2). Thirty out of the 46 (74%)

transcripts identified in the cerebellum and 44 out of the 52 (85%) transcripts expressed in the

blood could only be detected in less than half of samples (Fig. S1; Suppl. Table 2); for example,

the protein-coding transcript *ATXN3*-204 was only expressed in one out of 12 cerebellum

samples (8%) and in six out of 60 blood samples (10%) (Suppl. Table 2).

The protein coding *ATXN3*-251 reference transcript and the protein CDS not defined *ATXN3*-

208 transcript showed, respectively, the highest expression in blood and in cerebellum.

Additionally, the transcripts more abundant in blood by biotype were the protein coding

ATXN3-251, the NMD *ATXN3*-202, the noncoding *ATXN3*-208 and the retained intron *ATXN3*-

210, while in cerebellum the more abundant were the protein coding *ATXN3*-214, the NMD

ATXN3-234, the noncoding *ATXN3*-208 and the retained intron *ATXN3*-205 (Fig. 2).

Eight of the overall 54 annotated transcripts were exclusively expressed in blood: (i) two

corresponded to protein-coding transcripts (*ATXN3*-215, which is one of the four transcripts

encoding the full-length ataxin-3 protein isoforms, and *ATXN3*-228); (ii) four were transcripts

targeted for NMD (*ATXN3*-212, *ATXN3*-218, *ATXN3*-221, *ATXN3*-236) and (iii) two were protein

CDS not defined (*ATXN3*-220, *ATXN3*-233). *ATXN3*-212 transcript showed the highest

expression among all transcripts detected in the blood; (Fig. 2, Suppl. Table 2, and Fig. S1). Two

NMD transcripts were exclusively expressed in the cerebellum (*ATXN3*-242, *ATXN3*-244)

(Suppl. Table 2, Fig. S1).

Cerebellar abundance of *ATXN3* transcripts is similar in SCA3 patients and controls

Analysis of the distribution of *ATXN3* transcripts by biological group (SCA3/MJD patients

and/or controls) in cerebellum samples revealed that several transcripts were detected only in

individual or in a limited number of samples (Suppl. Table 2).

To compare distributions of cerebellar *ATXN3* transcripts between SCA3/MJD patients and

controls, transcripts detected in more than 50% of all cerebellum samples (six or more of 12

samples) were selected. Following this criterium, 16 transcripts were identified in cerebellum

samples corresponding to seven protein-coding (the reference variant - *ATXN3*-251, plus

ATXN3-201, *ATXN3*-206, *ATXN3*-207, *ATXN3*-214, *ATXN3*-219, *ATXN3*-227), four targeted for

NMD (*ATXN3*-234, *ATXN3*-243, *ATXN3*-253, *ATXN3*-254), three protein CDS not defined

(*ATXN3*-208, *ATXN3*-217, *ATXN3*-226) and two retained intron (*ATXN3*-205, *ATXN3*-210; Fig. 2).

While all 16 transcripts showed similar abundance in the cerebellum of SCA3/MJD patients and

controls (p>0.05; Fig. 3), the protein CDS not defined transcript *ATXN3*-226 was expressed in all

samples from controls but was not expressed in four out of the six patients (Fisher's exact test,

p=0.06; Suppl. Table 2).

Blood abundance of *ATXN3* transcripts is similar in SCA3/MJD subjects and controls.

Analysis of the distribution of *ATXN3* transcripts in blood samples from SCA3/MJD subjects

(pre-ataxic and patients) and controls revealed that, similarly to the cerebellum, several

transcripts were only detected in individual or in a limited number of samples (Suppl. Table 2).

Following the same criterium used for the cerebellum, eight *ATXN3* transcripts detected in

more than 50% of all blood samples (SCA3/MJD subjects and controls) were selected for

comparative analyses of patients versus controls and pre-ataxic carriers versus controls. These

eight transcripts corresponded to two protein coding (*ATXN3*-214 and the reference transcript

- *ATXN3*-251), two transcripts targeted for NMD (*ATXN3*-202, *ATXN3*-254), one protein CDS not

defined (*ATXN3*-249), and three retained introns (*ATXN3*-210, *ATXN3*-205, *ATXN3*-248) (Fig. 2).

All eight transcripts showed similar levels between SCA3/MJD patients or pre-ataxic carriers

compared to their respective controls (Fig. 4).

The amount of *ATXN3*-205 transcript in blood is directly associated with CSF levels of soluble

mutant ataxin-3.

The relationship between *ATXN3* transcript levels from blood cells and soluble levels of ataxin-

3 protein (mutant and/or total) quantified in plasma, peripheral mononuclear blood cells

(PBMCs) and CSF samples from the same SCA3/MJD subjects was explored in this study (Fig.

S2a). Interestingly, levels of the retained intron transcript *ATXN3*-205 in blood cells directly

correlated with the abundance of soluble mutant ataxin-3 in the CSF of SCA3/MJD subjects

(p=0.031, Fig. S2c). Although not reaching statistical significance, high levels of *ATXN3*-248 and

of the reference transcript *ATXN3*-251 displayed a trend to subtly associate with low levels of

total ataxin-3 measured in PBMCs of SCA3/MJD individuals (Fig. S2b, d).

Expression of the four transcripts encoding full-length ataxin-3 2UIM and 3UIM protein

isoforms are tissue-specific, irrespective of disease status.

From the 16 protein-coding transcripts, only four encode a full-length ataxin-3 protein with an

intact JD and a preserved C-terminus (*ATXN3*-214, *ATXN3*-215, *ATXN3*-227 and *ATXN3*-251; Fig.

5a). *ATXN3*-214 and *ATXN3*-227 transcripts encode ataxin-3 2UIM isoforms, whereas *ATXN3*-

215 and *ATXN3*-251 transcripts encode for ataxin-3 3UIM isoforms (Fig. 5a). To elucidate

whether the individual expression levels of the four transcripts encoding full-length ataxin-3

2UIM and 3UIM isoforms contribute to the tissue-specific vulnerability observed in SCA3/MJD,

the abundance of these transcripts in blood and cerebellum was compared between

SCA3/MJD patients and controls (Fig. 5b).

Interestingly, two of these four transcripts were only expressed in some samples. The *ATXN3*-

215 transcript (3UIM) was only identified in one blood sample (Supp. Table 2); while the

ATXN3-227 transcript (2UIM) was detected at distinct frequencies in both tissues, it was only

measured in three out of 60 (5%) blood samples and in seven out of 12 (58%) in cerebellum

samples (Supp. Table 2; Fig. 3). Although being overall commonly detected in the cerebellum,

ATXN3-227 transcript showed similar levels and frequency in SCA3/MJD patients and controls

(Fig. 3).

The reference transcript, *ATXN3*-251 (3UIM), showed a ~50-fold increase of abundance in

blood samples over cerebellum samples, whereas the *ATXN3*-214 (2UIM) transcript was more

expressed in the cerebellum about 20-fold more than in blood (Fig. 5b). The magnitude of the

fold change values, however, could be partially reflecting technical differences due to different

RNA-seq experimental setups. Remarkably, in cerebellum samples, the *ATXN3*-214 (2UIM) was

expressed 265-fold more than the *ATXN3*-251 (3UIM) transcript, whereas in blood samples the

ATXN3-251 (3UIM) transcript was expressed 4-fold more than *ATXN3*-214 (2UIM). These

observed differences in transcript abundances between tissues were similar in SCA3/MJD

subjects and controls.

DISCUSSION

Here, we describe the diversity and abundance of annotated *ATXN3* transcripts detected in

blood and cerebellum samples from SCA3/MJD subjects and controls. Globally, the number of

different transcripts and the abundance of transcripts were higher in the cerebellum than in

blood, both in SCA3/MJD subjects and controls. Interestingly, in SCA3/MJD subjects and

controls pooled samples altogether, while the most abundant transcript in the cerebellum

(*ATXN3*-208) is categorized as a "protein coding with a CDS not defined of unknown function

transcript" by Ensembl, the transcript showing the highest abundance in the blood is the

protein-coding *ATXN3* reference transcript (*ATXN3*-251) that is translated into an ataxin-3

3UIM protein isoform. No differences in the abundance of the most frequent transcripts were

found in blood or cerebellum samples of SCA3/MJD subjects and controls. Noteworthy, the

abundance of *ATXN3*-251 and *ATXN3*-214 transcripts, two out of the four transcripts that

encode full-length ataxin-3 protein isoforms but differ in the C-terminus (2UIM *versus* 3UIM)

were strongly associated with tissue expression specificity: *ATXN3-251* (3UIM) transcript was

expressed in blood 50-fold more than in cerebellum, whereas *ATXN3*-214 (2UIM) transcript

was expressed in the cerebellum 20-fold more than in blood.

Studies using a *Drosophila* model of SCA3/MJD indicate that in addition to a gain-of-function of

the mutant ataxin-3 protein, *ATXN3* transcripts harbouring expanded CAG repeats show

independent and progressive cellular toxicity (Li *et al*, 2008). However, whether such toxic

transcript action occurs in human cells remains to be elucidated. Hence, the characterization

of *ATXN3* AS in mutation carriers of SCA3/MJD might provide fundamental knowledge to

unveil the contribution of such mechanism in SCA3/MJD pathogenesis.

Globally, AS shows its highest complexity (high diversity of transcripts), in the brain amongst

other tissues (Porter *et al*, 2018). As expected, *ATXN3* transcripts diversity and abundance (a

greater number of transcripts, which are expressed in a larger number of samples with the

highest levels of expression) was higher in the cerebellum than in the blood and might indicate

an important role of *ATXN3* AS and its regulation in the neurodegenerative process of

SCA3/MJD.

Twelve out of 16 protein-coding transcripts are predicted to encode proteins with truncations

in the JD. Such modifications imply that several ataxin-3 protein forms with a variable N-

terminus might exist, displaying for example different forms of the globular JD or other N-

terminal domains that show cysteine protease activity or even another yet unknown activity.

The fact that ataxin-3 interacts with so many proteins in cells (reviewed in (Costa & Paulson,

2012)) and is ubiquitously expressed (*e.g.,* (Paulson *et al*, 1997)) together with our data

indicates that several ataxin-3 isoforms might exist with distinct functions other than the

currently known deubiquitinating enzyme role. Interestingly, four out of these 12 transcripts

(*ATXN3*-201, *ATXN3*-206, *ATXN3*-207, *ATXN3*-219) are highly expressed in the cerebellum

compared to blood suggesting that these potential encoded proteins may play specific roles in

the cerebellum and that the correspondent mutant forms could potentially contribute for the

observed selective cerebellar toxicity observed in SCA3/MJD patients.

Four out of 54 transcripts, *ATXN3*-214, *ATXN3*-215, *ATXN3*-227 and *ATXN3*-251, encode full-

length ataxin-3 protein isoforms, which differ only in the C-terminus. In the present study, the

ATXN3-214 and the *ATXN3*-227 transcripts, both encoding 2UIM isoforms, are preferentially

abundant in cerebellum samples, whereas the *ATXN3*-251 transcript (3UIM) is more expressed

in blood. Altogether, these findings suggest that ataxin-3 isoforms with 2UIM might have a

unique role in the cerebellum and consequently in neurodegeneration. The correlation

between transcript and protein abundance is still not known and future studies exploring the

role of the *ATXN3*-214 and *ATXN3*-227 transcripts and respective protein isoforms in brain

tissues could contribute for the elucidation of mechanisms involved in cellular vulnerability to

mutant *ATXN3* gene products in SCA3/MJD.

Of note, a significant direct relationship between the abundance of the retained intron *ATXN3*-

205 transcript in blood and soluble mutant ataxin-3 in CSF samples was observed in a small

sub-set of SCA3/MJD subjects. Because *ATXN3*-205 is classified by Ensembl as a long non-

coding RNA, such finding suggests that this transcript might have a regulatory function in the

transcription and/or translation of protein coding transcripts in the nervous system and,

therefore, contribute to the observed correlation with the soluble mutant ataxin-3 protein in

the CSF. More comprehensive studies about the function of these type of transcripts are

needed as well as larger number of CSF samples should be further analyzed to better explore

this preliminary but important finding.

In this study, transcripts more frequently found in blood (n=8) and in cerebellum (cerebellar

lobules, n=16) showed similar levels in SCA3/MJD patients and controls. Mort and colleagues

also described similar amount of two *HTT* transcripts in cerebellum samples of HD patients and

controls (Mort *et al*, 2015). In SCA3/MJD, neuronal loss in the cerebellar lobules is not as

severe as, for example, in the dentate nucleus or pons (Koeppen, 2018). Whether our findings

are specific of this brain area is unknown and, thus, characterization of *ATXN3* transcripts in

other SCA3/MJD patients brain tissues showing more severe lesions would be important to

conduct in the future.

In this work, the diversity and abundance of annotated *ATXN3* transcripts in SCA3/MJD

patients and controls is provided. The methodology used in this study did not allow to

specifically analyze the *ATXN3* transcripts with CAG repeat expansions. Future studies using

other NGS approaches, including novel bioinformatic pipelines to correctly detect CAG

expansions are needed to elucidate the contribution of the expanded CAG repeat itself in

ATXN3 AS. Moreover, functional studies to define the role of the most promising transcripts

found in the cerebellum, for example for its most abundant known non-coding *ATXN3*-208

transcript, would be important to conduct in the future. In fact, the majority of annotated

ATXN3 transcripts might have distinct functions or regulatory properties and its

comprehension will reveal novel mechanisms of SCA3/MJD pathogenesis.

We show that *ATXN3* AS is complex and, thus, its elucidation in the SCA3/MJD-related tissues

should become an integral part of future design of *ATXN3* transcripts-targeted therapeutics to

overcome potential limitations of efficacy of current RNA-based therapies targeting a limited

number of *ATXN3* transcripts. Our findings further aid to the clarification of the specific tissue

vulnerability observed in SCA3/MJD and, therefore, may help unravelling novel mechanisms

involved in SCA3/MJD pathogenesis.

ACKNOWLEDGEMENTS

The ESMI consortium would like to thank Nina Roy for coordination and managing of the

project. We gratefully thank Dr. Aires Raposo for the collaboration on blood collection in

Azores islands.

Additional members of the European Spinocerebellar Ataxia type 3/Machado-Joseph Initiative

(ESMI) Study Group are listed in Supplementary Material (Table S3) and participated in subject

recruitment and acquisition of participants data as well as read and approved the final

manuscript.

AUTHORS' ROLES

Design and conceptualization of the study: MR, JH-S, RT, MCC, ML; Subject

recruitment/Acquisition of participants biomaterials/data: MR, AFF, ARVM, JV, PP, TK, HG-M,

PG, MMS, LPA, JI, BPW, JV, JF, TK, NC, JA, LS, OR, ML; Bioinformatic/Statistical analysis of data:

MR, RT; Drafting of the manuscript: MR, JH-S, MCC, ML; Revision of the Manuscript: MR, JH-S,

RT, AFF, ARVM, JV, PP, TK, HG-M, PG, MMS, LPA, JI, BPW, JV, JF, TK, NC, JA, LS, OR, MCC, ML.

All authors read and approved the final manuscript.

REFERENCES

Andrews S (2010) FastQC: A Quality Control Tool for High Throughput Sequence Data [Online].
(http://www.bioinformatics.babraham.ac.uk/projects/fastqc/) [PREPRINT]

Barbaro BA, Lukacsovich T, Agrawal N, Burke J, Bornemann DJ, Purcell JM, Worthge SA,
Caricasole A, Weiss A, Song W, *et al* (2015) Comparative study of naturally occurring
huntingtin fragments in Drosophila points to exon 1 as the most pathogenic species in
Huntington's disease. *Hum Mol Genet* 24: 913–925

Bettencourt C, Santos C, Montiel R, Costa M do C, Cruz-Morales P, Santos LR, Simões N, Kay T,
Vasconcelos J, Maciel P, *et al* (2010) Increased transcript diversity: novel splicing variants
of Machado–Joseph Disease gene (ATXN3). *Neurogenetics* 11: 193–202

Burnett B, Li F & Pittman RN (2003) The polyglutamine neurodegenerative protein ataxin-3
binds polyubiquitylated proteins and has ubiquitin protease activity. *Hum Mol Genet* 12:
3195–205

Costa M do C & Paulson HL (2012) Toward understanding Machado-Joseph disease. *Prog*
Neurobiol 97: 239–257 doi:10.1016/j.pneurobio.2011.11.006 [PREPRINT]

Daguenet E, Dujardin G & Valcárcel J (2015) The pathogenicity of splicing defects: mechanistic
insights into pre- <scp>mRNA</scp> processing inform novel therapeutic approaches.
EMBO Rep 16: 1640–1655

Dhuri K, Bechtold C, Quijano E, Pham H, Gupta A, Vikram A & Bahal R (2020) Antisense
Oligonucleotides: An Emerging Area in Drug Discovery and Development. *J Clin Med* 9:
2004

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M & Gingeras
TR (2013) STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* 29: 15–21

Gonsior K, Kaucher GA, Pelz P, Schumann D, Gansel M, Kuhs S, Klockgether T, Forlani S, Durr A,
Hauser S, *et al* (2021) PolyQ-expanded ataxin-3 protein levels in peripheral blood
mononuclear cells correlate with clinical parameters in SCA3: a pilot study. *J Neurol* 268:
1304–1315

Goto J, Watanabe M, Ichikawa Y, Yee S-B, Ihara N, Endo K, Igarashi S, Takiyama Y, Gaspar C,

Maciel P, *et al* (1997) Machado–Joseph disease gene products carrying different carboxyl
termini. *Neurosci Res* 28: 373–377

Haas E, Incebacak RD, Hentrich T, Huridou C, Schmidt T, Casadei N, Maringer Y, Bahl C,
Zimmermann F, Mills JD, *et al* (2022) A Novel SCA3 Knock-in Mouse Model Mimics the
Human SCA3 Disease Phenotype Including Neuropathological, Behavioral, and
Transcriptional Abnormalities Especially in Oligodendrocytes. *Mol Neurobiol* 59: 495–522

Harris GM, Dodelzon K, Gong L, Gonzalez-Alegre P & Paulson HL (2010) Splice isoforms of the
polyglutamine disease protein ataxin-3 exhibit similar enzymatic yet different aggregation
properties. *PLoS One* 5

Hauser S, Helm J, Kraft M, Korneck M, Hübener-Schmid J & Schöls L (2022) Allele-specific
targeting of mutant ataxin-3 by antisense oligonucleotides in SCA3-iPSC-derived neurons.
Mol Ther - Nucleic Acids 27: 99–108

Helm J, Schöls L & Hauser S (2022) Towards Personalized Allele-Specific Antisense
Oligonucleotide Therapies for Toxic Gain-of-Function Neurodegenerative Diseases.
Pharmaceutics 14: 1708

Hübener-Schmid J, Kuhlbrodt K, Peladan J, Faber J, Santana MM, Hengel H, Jacobi H, Reetz K,
Garcia-Moreno H, Raposo M, *et al* (2021) Polyglutamine-Expanded Ataxin-3: A Target
Engagement Marker for Spinocerebellar Ataxia Type 3 in Peripheral Blood. *Mov Disord*
36: 2675–2681

Ichikawa Y, Goto J, Hattori M, Toyoda A, Ishii K, Jeong SY, Hashida H, Masuda N, Ogata K, Kasai
F, *et al* (2001) The genomic structure and expression of MJD, the Machado-Joseph
disease gene. *J Hum Genet* 46: 413–22

Johnson SL, Blount JR, Libohova K, Ranxhi B, Paulson HL, Tsou W-L & Todi S V. (2019)
Differential toxicity of ataxin-3 isoforms in Drosophila models of Spinocerebellar Ataxia
Type 3. *Neurobiol Dis* 132: 104535

Kawaguchi Y, Okamoto T, Taniwaki M, Aizawa M, Inoue M, Katayama S, Kawakami H,
Nakamura S, Nishimura M & Akiguchi I (1994) CAG expansions in a novel gene for
Machado-Joseph disease at chromosome 14q32.1. *Nat Genet* 8: 221–8

Koeppen AH (2018) The Neuropathology of Spinocerebellar Ataxia Type 3/Machado-Joseph
Disease. In pp 233–241.

Li B & Dewey CN (2011) RSEM: accurate transcript quantification from RNA-Seq data with or
without a reference genome. *BMC Bioinformatics* 12: 323

Li L-B, Yu Z, Teng X & Bonini NM (2008) RNA toxicity is a component of ataxin-3 degeneration
in Drosophila. *Nature* 453: 1107–1111

Maciel P, Costa MC, Ferro A, Rousseau M, Santos CS, Gaspar C, Barros J, Rouleau GA, Coutinho
P & Sequeiros J (2001) Improvement in the molecular diagnosis of Machado-Joseph
disease. *Arch Neurol* 58: 1821–1827

Mangiarini L, Sathasivam K, Seller M, Cozens B, Harper A, Hetherington C, Lawton M, Trottier
Y, Lehrach H, Davies SW, *et al* (1996) Exon 1 of the HD Gene with an Expanded CAG
Repeat Is Sufficient to Cause a Progressive Neurological Phenotype in Transgenic Mice.
Cell 87: 493–506

Martin FJ, Amode MR, Aneja A, Austine-Orimoloye O, Azov AG, Barnes I, Becker A, Bennett R,
Berry A, Bhai J, *et al* (2023) Ensembl 2023. *Nucleic Acids Res* 51: D933–D941

Masino L, Musi V, Menon RP, Fusi P, Kelly G, Frenkiel TA, Trottier Y & Pastore A (2003) Domain
architecture of the polyglutamine protein ataxin-3: a globular domain followed by a
flexible tail. *FEBS Lett* 549: 21–5

McLoughlin HS, Moore LR, Chopra R, Komlo R, McKenzie M, Blumenstein KG, Zhao H,
Kordasiewicz HB, Shakkottai VG & Paulson HL (2018) Oligonucleotide therapy mitigates
disease in spinocerebellar ataxia type 3 mice. *Ann Neurol* 84: 64–77

Moore LR, Keller L, Bushart DD, Delatorre RG, Li D, McLoughlin HS, do Carmo Costa M,
Shakkottai VG, Smith GD & Paulson HL (2019) Antisense oligonucleotide therapy rescues
aggresome formation in a novel spinocerebellar ataxia type 3 human embryonic stem cell
line. *Stem Cell Res* 39: 101504

Moore LR, Rajpal G, Dillingham IT, Qutob M, Blumenstein KG, Gattis D, Hung G, Kordasiewicz
HB, Paulson HL & McLoughlin HS (2017) Evaluation of Antisense Oligonucleotides
Targeting ATXN3 in SCA3 Mouse Models. *Mol Ther - Nucleic Acids* 7: 200–210

Mort M, Carlisle FA, Waite AJ, Elliston L, Allen ND, Jones L & Hughes AC (2015) Huntingtin
Exists as Multiple Splice Forms in Human Brain. *J Huntingtons Dis* 4: 161–171

Neueder A, Landles C, Ghosh R, Howland D, Myers RH, Faull RLM, Tabrizi SJ & Bates GP (2017)
The pathogenic exon 1 HTT protein is produced by incomplete splicing in Huntington's
disease patients. *Sci Rep* 7: 1307

Nik S & Bowman T V. (2019) Splicing and neurodegeneration: Insights and mechanisms. *WIREs*
RNA 10

Paulson HL, Das SS, Crino PB, Perez MK, Patel SC, Gotsdiner D, Fischbeck KH & Pittman RN
(1997) Machado-Joseph disease gene product is a cytoplasmic protein widely expressed
in brain. *Ann Neurol* 41: 453–62

Porter RS, Jaamour F & Iwase S (2018) Neuron-specific alternative splicing of transcriptional
machineries: Implications for neurodevelopmental disorders. *Mol Cell Neurosci* 87: 35–45

Raposo M, Hübener-Schmid J, Ferreira AF, Vieira Melo AR, Vasconcelos J, Pires P, Kay T, Garcia-
Moreno H, Giunti P, Santana MM, *et al* (2023) Blood transcriptome sequencing identifies
biomarkers able to track disease stages in spinocerebellar ataxia type 3. *Brain* accepted

Sathasivam K, Neueder A, Gipson TA, Landles C, Benjamin AC, Bondulich MK, Smith DL, Faull
RLM, Roos RAC, Howland D, *et al* (2013) Aberrant splicing of HTT generates the
pathogenic exon 1 protein in Huntington disease. *Proc Natl Acad Sci* 110: 2366–2370

Sequeiros J, Martins S & Silveira I (2012) Epidemiology and population genetics of degenerative
ataxias. In pp 227–251.

Takiyama Y, Nishizawa M, Tanaka H, Kawashima S, Sakamoto H, Karube Y, Shimazaki H,
Soutome M, Endo K & Ohta S (1993) The gene for Machado-Joseph disease maps to
human chromosome 14q. *Nat Genet* 4: 300–4

Vuong CK, Black DL & Zheng S (2016) The neurogenetics of alternative splicing. *Nat Rev*
Neurosci 17: 265–281

Weishäupl D, Schneider J, Peixoto Pinheiro B, Ruess C, Dold SM, von Zweydorf F, Gloeckner CJ,
Schmidt J, Riess O & Schmidt T (2019) Physiological and pathophysiological characteristics
of ataxin-3 isoforms. *J Biol Chem* 294: 644–661

Zhao S, Ye Z & Stanton R (2020) Misuse of RPKM or TPM normalization when comparing across
samples and sequencing protocols. *RNA* 26: 903–909

Zhu Y, Zhu L, Wang X & Jin H (2022) RNA-based therapeutics: an overview and prospectus. *Cell*
Death Dis 13: 644

Figure 1. Schematic representation of the annotated 54 *ATXN3* transcripts (adapted from

Ensembl release 108 – Jan 2023) detected in blood and cerebellum of pooled SCA3/MJD patients

and controls. *ATXN3* transcripts in both SCA3/MJD subjects and controls are clustered according

to Ensembl-related biotype - protein coding, nonsense mediated decay, protein coding

sequence (CDS) not defined, and retained intron. Moreover, transcripts are clustered by

descending order of frequency within tissue calculated in the present study as well as transcripts

with exclusive expression in blood and transcripts with exclusive expression in cerebellum are

highlighted in orange and green, respectively. The Ensembl's *ATXN3* reference transcript

ENST00000644486.2|*ATXN3*-251 (MANE Select v0.95) is highlighted in blue. Protein-coding

transcripts not containing exon 10 (which contains the CAG tract) are highlighted in bold. *The

annotation of the transcript is incomplete at 5' untranslated (UTR), 3'UTR or both. The catalytic

amino acids cysteine (C) 14, histidine (H) 119 and asparagine (N) 134 are shown in the respective

exons; red diamonds represent a premature termination codon (PTC). Grey box = coding exon;

light grey box = UTR; white box = non-coding exon; UIM = ubiquitin interacting motif; polyQ =

polyglutamine tract; NES = nuclear export signal; NLS = nuclear localization signal. The size of

exons/introns/UTR are not drawn to scale.

Figure 2. Heatmap of the 54 annotated *ATXN3* transcripts detected in cerebellum and blood

samples of SCA3/MJD subjects and healthy individuals. Transcripts are clustered by biotype

(protein coding, nonsense mediated decay, protein sequence (CDS) not defined, and retained

intron) and disease/health status. Expression values, shown in transcripts per million (TPM),

were obtained for 72 samples (12 from cerebellum - lines one to 12) and 60 from whole blood

(lines 13 to 72). The magnitude of the expression values was set independently for each

transcript type cluster. Protein-coding transcripts not containing exon 10 (which contains the

CAG tract) are highlighted in bold. The brain (🧠) and/or blood (🩸) symbols indicate transcripts

detected in more than 50% of all samples in each tissue. *ATXN3* transcripts with exclusive

expression in blood and transcripts with exclusive expression in cerebellum are highlighted in

orange and green, respectively. The Ensembl's *ATXN3* reference transcript

ENST00000644486.2|*ATXN3*-251 (MANE Select v0.95) is highlighted in blue.

Figure 3. Levels of the 16 *ATXN3* transcripts expressed in more than half of all cerebellum

samples from SCA3/MJD patients (n=6) and controls (n=6). The transcripts are clustered

according to the type of transcript, as defined by Ensembl classification biotype (Ensembl release

108 - Jan 2023, REF). Transcript expression levels are shown in transcript per million (TPM). Bars

in the graphs represent the median ± 95% CI (confidence interval, error bars).

Figure 4. Expression levels of eight *ATXN3* transcripts in blood samples from pre-ataxic (PA)

carriers (n=10), SCA3/MJD patients (n=30) and respective controls (CTRL-P, n=20 and CTRL-PA,

n=10). CTRL-PA was defined as a subset of age-matched controls with pre-ataxic carriers within

the CTRL-P group. The transcripts are clustered according to biotype, as defined by Ensembl

classification biotype (Ensembl release 108 - Jan 2023(Martin *et al*, 2023)). Expression levels are

shown in transcript per millions (TPM). Bars in the graphs represent the median ± 95% CI

(confidence interval, error bars).

Figure 5. (a) Sequence alignment of the four full-length ataxin-3 protein isoforms harbouring an

intact Josephin domain (JD) and a preserved C-terminus. Ataxin-3 protein isoforms, namely the

P54252-2, F5H211, P54252-1 and G3V3R7 are encoded by the *ATXN3*-251, *ATXN3*-215, *ATXN3*-

214 and *ATXN3*-227 transcripts, respectively. Multiple sequence alignment was performed using

Clustal Omega program (Sievers F. et al., 2011). JD (green line), the three catalytic sites (orange

dot) and the three UIMs (blue boxes) are highlighted. (b) Levels of *ATXN3*-251 (3UIM) and of the

ATXN3-214 as well as *ATXN3*-227 (both 2UIM) transcripts in the blood and cerebellum samples

from SCA3/MJD subjects and controls. Transcript levels are shown in transcripts per million

(TPM). Bars in the graphs represent the median ± 95% CI (confidence interval, error bars).

reverse strand

92.11Mb   92.09Mb   92.07Mb   92.05Mb

Protein coding

ATXN3-251
ATXN3-214
ATXN3-219
ATXN3-201
ATXN3-207
ATXN3-227
ATXN3-206
ATXN3-235
ATXN3-203
ATXN3-204
ATXN3-231
ATXN3-246
ATXN3-215
ATXN3-228
ATXN3-209
ATXN3-213

Ataxin-3 protein (P54252-2, UniProt canonical sequence)

N ^{14}C ^{119}H ^{134}N   UIM UIM   UIM

Josephin

NES   NES   NLS

polyQ

C

Nonsense mediated decay

ATXN3-254
ATXN3-243
ATXN3-234
ATXN3-253
ATXN3-202
ATXN3-242
ATXN3-232
ATXN3-230
ATXN3-244
ATXN3-218
ATXN3-221
ATXN3-225
ATXN3-236
ATXN3-237
ATXN3-238
ATXN3-245
ATXN3-240
ATXN3-229
ATXN3-212

Protein CDS not defined

ATXN3-217
ATXN3-226
ATXN3-208
ATXN3-249
ATXN3-216
ATXN3-220
ATXN3-223
ATXN3-224
ATXN3-233
ATXN3-241
ATXN3-239
ATXN3-222
ATXN3-211
ATXN3-247
ATXN3-252
ATXN3-250

Retained intron

ATXN3-210
ATXN3-205
ATXN3-248

protein coding

nonsense mediated decay

protein CDS not defined

retained intron

a.

```
P54252-2|ATXN3-251    MESIFHEKQEGSLCAQHCLNNLLQGEYFSPVELSSIAHQLDEEERMRMAEGGVTSEDYRT    60
F5H211|ATXN3-215      MESIFHEKQEGSLCAQHCLNNLLQGEYFSPVELSSIAHQLDEEERMRMAEGGVTSEDYRT    60
P54252-1|ATXN3-214    MESIFHEKQEGSLCAQHCLNNLLQGEYFSPVELSSIAHQLDEEERMRMAEGGVTSEDYRT    60
G3V3R7|ATXN3-227      MESIFHEKQEGSLCAQHCLNNLLQGEYFSPVELSSIAHQLDEEERMRMAEGGVTSEDYRT    60
                      ************************************************************

P54252-2|ATXN3-251    FLQQPSGNMDDSGFFSIQVISNALKVWGLELILFNSPEYQRLRIDPINERSFICNYKEHW   120
F5H211|ATXN3-215      FLQQPSGNMDDSGFFSIQVISNALKVWGLELILFNSPEYQRLRIDPINERSFICNYKEHW   120
P54252-1|ATXN3-214    FLQQPSGNMDDSGFFSIQVISNALKVWGLELILFNSPEYQRLRIDPINERSFICNYKEHW   120
G3V3R7|ATXN3-227      FL-QPSGNMDDSGFFSIQVISNALKVWGLELILFNSPEYQRLRIDPINERSFICNYKEHW   119
                      **  *******************************************************

P54252-2|ATXN3-251    FTVRKLGKQWFNLNSLLTGPELISDTYLALFLAQLQQEGYSIFVVKGDLPDCEADQLLQM   180
F5H211|ATXN3-215      FTVRKLGKQWFNLNSLLTGPELISDTYLALFLAQLQQEGYSIFVVKGDLPDCEADQLLQM   180
P54252-1|ATXN3-214    FTVRKLGKQWFNLNSLLTGPELISDTYLALFLAQLQQEGYSIFVVKGDLPDCEADQLLQM   180
G3V3R7|ATXN3-227      FTVRKLGKQWFNLNSLLTGPELISDTYLALFLAQLQQEGYSIFVVKGDLPDCEADQLLQM   179
                      ************************************************************

P54252-2|ATXN3-251    IRVQQMHRPKLIGEELAQLKEQRVHKTDLERVLEANDGSGMLDEDEEDLQRALALSRQEI   240
F5H211|ATXN3-215      IRVQQMHRPKLIGEELAQLKEQRVHKTDLERVLEANDGSGMLDEDEEDLQRALALSRQEI   240
P54252-1|ATXN3-214    IRVQQMHRPKLIGEELAQLKEQRVHKTDLERVLEANDGSGMLDEDEEDLQRALALSRQEI   240
G3V3R7|ATXN3-227      IRVQQMHRPKLIGEELAQLKEQRVHKTDLERVLEANDGSGMLDEDEEDLQRALALSRQEI   239
                      ************************************************************

P54252-2|ATXN3-251    DMEDEEADLRRAIQLSMQGSSRNISQDMTQTSGTNLTSEELRKRREAYFEKQ--------   292
F5H211|ATXN3-215      DMEDEEADLRRAIQLSMQGSSRNISQDMTQTSGTNLTSEELRKRREAYFEKVRLRLQKYI   300
P54252-1|ATXN3-214    DMEDEEADLRRAIQLSMQGSSRNISQDMTQTSGTNLTSEELRKRREAYFEKQ-------   292
G3V3R7|ATXN3-227      DMEDEEADLRRAIQLSMQGSSRNISQDMTQTSGTNLTSEELRKRREAYFEKQ--------   291
                      ***************************************************

P54252-2|ATXN3-251    -QQKQQQQQQQQQQGDLSGQSSHPCERPATSSGALGSDLGDAMSEEDMLQAAVTMSLETV   351
F5H211|ATXN3-215      QQQKQQQQQQQQQQGDLSGQSSHPCERPATSSGALGSDLGDAMSEEDMLQAAVTMSLETV   360
P54252-1|ATXN3-214    -QQKQQQQQQQQQQGDLSGQSSHPCERPATSSGALGSDLGKACSPFIMFATFTLYLTYEL   351
G3V3R7|ATXN3-227      -QQKQQQQQQQQQQGDLSGQSSHPCERPATSSGALGSDL--------------------   329
                      **************************************

P54252-2|ATXN3-251    RND--LKTEGKK-   361
F5H211|ATXN3-215      RND--LKTEGKK-   370
P54252-1|ATXN3-214    HVIFALHYSSFPL   364
G3V3R7|ATXN3-227      -------------   329
```

b.

ATXN3-251 (3UIM)   <0.0001

ATXN3-214 (2UIM)   <0.0001

ATXN3-227

▲ Controls   ● Patients