The cortical microtubules of *Toxoplasma* contribute to the helicity, but not the speed or persistence of parasite movement

Isadonna F. Tengganu[1#], Luisa F. Arias Padilla[1#], Jonathan Munera Lopez[1#], Jun Liu[2,3], Peter T. Brown[4], Ke Hu[1*]

[1]Biodesign Center for Mechanisms of Evolution/School of Life Sciences, Arizona State University, USA
[2]Department of Biology, Indiana University, USA; [3]Current address: Berkgen Biopharmaceuticals, Shenzhen, China
[4]Department of Physics and Center for Biological Physics, Arizona State University, USA

#: equal contribution

*: Address correspondence to:
Email: kehu4@asu.edu

Running title: *cortical microtubules facilitate helical parasite movement*

ABSTRACT

Apicomplexans parasitize a wide range of hosts. To infect, the parasite needs to travel through different types of tissues and invade into various types of cells. In tissues and three-dimensional (3-D) matrix, the apicomplexan parasite *Toxoplasma gondii* moves along a helical path. While many genes, including actin and myosins, have been shown to be important for parasite motility, it remains unknown what allows the parasite to travel over a long distance along a helical path. The cortical microtubules, which are ultra-stable, sprially arranged, and form extensive lateral interaction with the parasite cortex, have been considered to be a candidate structure for guiding the long-distance movement of the parasite.

In wild-type parasites, the cortical microtubules in mature parasites are impervious to destabilization by cold-treatment or depolymerizing drugs, which makes it difficult to assess their function. Previously, we generated a mutant (dubbed "TKO" for succinctness) that lacks three microtubule-associated proteins. The loss of these three proteins destabilizes the cortical microtubules in mature parasites. Here we quantify the proportion of parasites with various levels of defects in the microtubule array under different conditions. We found that ~ 80% of the non-dividing TKO parasites have severely curtailed cortical microtubules. The extent of depolymerization is further exacerbated upon the initiation of daughter construction or cold treatment. The cold-induced depolymerization is reversible, with noticeable restoration of the cortical microtubules within 20 min of temperature shift to 37°C. While microtubule polymerization is essential for generating viable daughter cells, the destabilization of the cortical microtubules in the mature parasite does not affect parasite replication. In a 3-D Matrigel matrix, the TKO mutant parasites can travel directionally over long distances. However, their trajectories are significantly more linear than those of wild-type parasites. In tissue culture, the TKO parasite displays a defect in infection and cytolytic efficiency. Interestingly, the speed and behavior of the parasite's entry into and egress from the host cell are similar to that of the wild-type parasite. These results indicate that the cortical microtubules contribute to the helicity but not the persistence of parasite long-distance movement. Furthermore, host cell entry is less sensitive to structural changes in the parasite than overall infection efficiency, which also include extracellular migration and orient the parasite for proper contact with the host cell.

INTRODUCTION

Apicomplexans are obligate intracellular parasites, many of which are important pathogens. For example, the malaria parasites, *Plasmodium* spp., kill more than half a million people each year [1]. *Toxoplasma gondii* permanently infects ~ 20% of the global human population and can infect any warm-blooded animal [2]. As obligate intracellular organisms, a successful infection requires the parasite to travel over a long distance through tissues with varying biochemical and biophysical properties as well as to enter into host cells, which requires shorter travel distance, but perhaps more finesse, as the host cell plasma membrane needs to remain intact to allow the parasite to use host cell resources for proliferation.

Toxoplasma gondii moves along an irregular helical path in tissues and a three-dimensional (3-D) matrix over tens of microns [3]. Chemical inhibition as well as genetic manipulations showed that the basic components driving the motility of *Toxoplasma* include actin and a series of apical and cortical myosin motors for providing the driving force, secreted adhesins on the parasite surface for engaging with the environment, as well as associated proteins for coupling the intracellular domains of the adhesins to the actomyosin complex [4-13]. The characterization of these players has significantly advanced the understanding of how energy is injected into the system to fuel motility. However, while actin polymerization is critical for parasite movement, no native actin filaments have been detected, suggesting that the polymerization is likely to occur dynamically. Thus, it is still not known how the parasite is capable of traveling over long distance along a helical path. Intuitively, a stable, inherently helical cortical structure would be required for guiding continuous parasite movement over a long distance.

In *Toxoplasma,* one of the main cytoskeletal structures with a roughly helical arrangement is the set of cortex-associating (denoted as "subpellicular" or "cortical") microtubules that are anchored at one end in a ring-like structure (the apical polar ring) at the parasite apex [14-16]. Among the apicomplexans, the microtubule cytoskeleton displays intriguing species and life-stage specific variations that coincide with substantial differences in physiology. The mature *Plasmodium* ookinete has an intricate cage of ~ 60 cortical microtubules [17-19], but the asexual blood-stage parasites only have a narrow band of 2-4 cortical microtubules that extends down one side of the parasite [14, 20]. The *Toxoplasma* tachyzoite has consistently 22 cortical microtubules that are arranged in a helical, or more accurately, a spiral pattern. The parasite invests considerable amount of resources in building and maintaining these polymers. Aside from the core building blocks, the tubulins, there is a suite of proteins that associate with the cortical microtubules in a region-specific manner [15, 21-23]. In mature parasites, these microtubules form an ultra-stable "rib cage" that form extensive lateral interaction with the inner membrane complex (IMC). They remain intact under treatments that depolymerize microtubules in mammalian cells within minutes [24, 25]. This extraordinary stability makes it difficult to assess the function of these polymers in the mature parasite.

Previously, we generated a mutant (dubbed "TKO" for succinctness) that lacks three microtubule-associated proteins, TLAP2, SPM1 and TLAP3 [22]. The removal of these three proteins dramatically reduces the stability of the cortical microtubules in mature parasites. In the present work, we quantified the proportion of parasites with various levels of defects in the microtubule array under different conditions and found that ~ 80% of the non-dividing TKO parasites have severely curtailed cortical microtubules. Depolymerization in mature parasites becomes even more pronounced upon the initiation of daughter construction or cold treatment,

the latter effect being reversible. While microtubule polymerization is essential for generating
viable daughter cells, parasite replication is not affected by the destabilization of the cortical
microtubules in the mature parasite. In a 3-D Matrigel matrix, the TKO mutant parasite moves
as persistently as the wild-type parasite with similar velocity and net displacement. However, it
travels along a path significantly more linear than the wild-type parasite. Interestingly, even
though the cytolytic efficiency of the TKO parasite is significanty lower, live cell imaging
revealed that the parasite's behaviors during entry into and egress from the host cell is
indistinguishable from those of the wild-type. These results indicate that the cortical
microtubules contribute to the helicity but not the persistence of parasite long-distance
movement. On the other hand, host cell entry and egress, which involve short-distance travel,
is less sensitive to structural changes in the parasite than overall infection efficiency.
121
122

123 **RESULTS**
124

125 *The TKO parasite replicates normally, likely due to the cell-stage dependent nature of the*
126 *destabilization of cortical microtubules*
127

Previously, we showed qualitatively that the cortical microtubules in mature TKO parasites are
destabilized compared with those in the wild-type parasites [22]. Here to quantify the extent of
destabilization, we determined the proportions of TKO and wild-type parasites that display
various characteristics in the microtubule array. We found that the cortical microtubules are
severely curtailed in 79% (±8.5% SEM) of non-dividing TKO parasites and sparse in the rest,
whereas 100% of the wild-type parasites have a full, regularly arranged array (Fig 1A) (Three
independent experiments, total 490 parasites analyzed for the TKO and 246 parasites
analyzed for the wild-type). The extent of depolymerization in mature TKO parasites is more
pronounced during cell division (Fig 1B-C). The proportion of parasites with nearly fully
depolymerized cortical microtubules ("absent" category) was ~ 42% at the initiation stage of
daughter construction and increased to ~ 98% prior to daughter emergence from the mother
parasite (Fig 1C). For wild-type parasites, the cortical microtubules of the mother parasite in
the initiation and elongation stages of daughter construction are similar to that of non-dividing
stage (Fig 1B-C). However, ~ 24% of mature parasites at the pre-emergence stage and 100%
during daughter emergence have short microtubules.
143

We examined the average number of rounds of replication at 12, 24, and 36 hours after
infection and found that the rate of replication of the TKO parasite is very similar to that of the
wild-type parasite (Fig 1D). At first glance, it is surprising that a mutant that has destabilized
cortical microtubules does not have a replication defect, because microtubule polymerization is
coupled with cortex formation, and thus is required for the daughter formation [26-29].
However, as we observed previously [22], in the same cell in which the cortical microtubules of
the mother TKO parasite are depolymerized, the microtubules do polymerize in growing
daughters, which develop normally. These results reveal a cell-stage dependent microtubule
depolymerizing activity that acts on the cortical microtubules in mature parasites but not the
polymerizing microtubules in growing daughters. The depolymerization of cortical microtubules
in mature TKO parasites becomes apparent at a much earlier stage and is much more
complete than the wild-type parasites likely because they are already destabilized due to the
lack of TLAP2, TLAP3, and SPM1.
157

In our previous work, we showed that incubation at ~ 8°C can further destabilize the
cortical microtubules of the TKO parasite [22]. Here we characterize the time course of cold-

induced disassembly of cortical MTs in mature TKO parasites and found that, as expected, the
extent of depolymerization increased with time of cold-treatment (Fig 1E). After ~ 2 hr of cold-
treatment (C120), cortical MTs in ~ 72% of the parasites are nearly completely depolymerized
with no visible MTs aside from a small tubulin-positive "cap" at the apical end ("absent"
category). On the other hand, the array of cortical microtubules in wild-type parasite remains
similar before and after the cold treatment. The cold-induced depolymerization in the TKO
parasite is reversible (Fig 1E). When TKO cultures cold-treated for 2 hr were returned to 37°C,
the cortical MTs were restored significantly after 20 min of the temperature shift. At 1hr of 37°C
incubation (R60), ~ 80% of the parasites showed patterns ("short or sparse") similar to those of
untreated parasites, with the microtubules concentrated at the apical end of the parasite (Fig
3E), suggesting strong preference for elongation from pre-existing nucleation sites.
171
172
173 *The trajectory of movement of TKO parasite is more linear than that of the wild-type parasite in*
174 *3-D matrigel*
175
In wild-type parasites, the stable, spirally arranged cortical microtubules have been proposed
to guide directionality of parasite movement [30] or to function as a spring that transmits
energy to drive gliding motility [31]. We tested this hypothesis by comparing the 3-D motile
behaviors of the wild-type parental with those of the TKO parasite, taking advantage of the
destabilized and curtailed microtubule array in the TKO. By modifying an established Matrigel-
based 3-D motility assay [3], we obtained optical sections of the moving parasites under a
label-free condition through Differential Interference Contrast (DIC) imaging. DIC imaging not
only minimizes phototoxicity, but also enables the observation of parasite morphological
changes as they move. However, while fluorescence images exhibit high-contrast that
facilitates cell-tracking, computerization of cell positions in DIC image stacks are not as
straightforward. We applied a series of edge detection filters, followed by erosion and dilation
filters on the DIC images to derive images with contrast similar to fluorescence images (Fig
2A-B) that are amenable to 3-D cell tracking. Detection and tracking were carried out using the
TrackMate plugin in Fiji which generates the 3-D coordinates of the cell trajectories for further
analysis and 3-D visualization (Fig 2C and Fig 3, Video S1-S4).
191
In these experiments, 40.2% of the wild-type parental (RHΔhx) parasites were motile
(413 parasites from 6 independent experiments analyzed) (Table 1). The average net
displacement is 31.4 μm. The average run length (i.e., distance between pauses of >5.4 sec)
is 22.7 μm and the average run speed (average speed during a run) is 1.28 μm/sec (Fig 3A).
Similarly, ~42.3% of the TKO parasites were motile (395 parasites from 6 independent
experiments analyzed) with an average net displacement of 31.8 μm, average run length of
23.3 μm and average run speed of 1.3 μm/sec. As previously reported [3], the wild-type
parasite moves along an irregular helical path. The average dispersal ratio (ratio of net
displacement to total distance traveled of motile segments) of the trajectories is 0.71(a value of
1 indicates a completely linear track). The average turn angle (relative angle of parasite
between two consecutive steps) is ~50°. The movement of the TKO parasite was visibly more
linear than that of the wild-type parasite (Fig 3B, Video S1-S2). Confirming this visual
impression, we found that the dispersal ratio of the TKO parasites is 0.78, significantly higher
than that of the wild-type parasite (p << 0.0001, Fig 3C). The turn angle of the TKO is ~32.6°,
significantly lower than the wild-type parasite (p << 0.0001). Both are consistent with the
trajectories being more linear. For segments of trajectories during which the TKO parasite
displays near helical movement, the average pitch of the helical turns is 9.9 μm, significantly
higher than that of the wild-type parasite, which is 7.5 μm (p << 0.0001).

As shown in Figure 1A, there are remnant cortical microtubules in untreated TKO parasites. To determine whether these remaining microtubules influence parasite movement, we cold-treated the parasites for 2 hr, under which condition the cortical microtubules are nearly fully depolymerized in >70% of the TKO parasites but those in the wild-type parasites are not affected (see Fig 1E). The parasites were then harvested and mixed with Matrigel for 3-D motility assay. We found that the trajectories of the cold-treated TKO parasites are qualitatively similar to those of untreated TKO parasites (Fig 3B).

In addition to the cortical microtubules, another helically arranged cytoskeletal structure in the parasite is the conoid (Fig 1B), which is a substructure in the apical complex formed of helically arranged tubulin fibers [25, 28, 32, 33]. When a structural component (TgDCX) was removed, the conoid structure becomes stunted and disorganized and the parasite displays severe invasion defects [33]. To determine whether the helicity of the trajectory is sensitive to changes in the conoid structure, we examined the motile behavior of the DCX knockout parasite (*Δdcx*). We found that on average, the motile percentage of the *Δdcx* is ~31%, slightly lower than that of its "wild-type" parental (RH*ΔhxΔku80*, abbreviated as "RH*Δku80*") (Table 1). The average net displacement, run-length and run speed of the *Δdcx* parasite are similar to those of the wild-type parasites (30.2 vs 30.7 μm, ~26 vs 25 μm and 1.21 vs 1.28 μm/sec, respectively, Fig 3A). The trajectories of the *Δdcx* are helical in nature (Fig 3B-C, Video S3-S4), with an average dispersal ratio of 0.68 and turn angle of 52°, indicating slightly tighter turns compared with its parental parasite (average dispersal ratio of 0.72 and turn angle of 48.4°, $p < 0.05$, Fig 3C). This indicates that the conoid is not a major contributor to the helicity of parasite movement.

The infection efficiency of the TKO is lower than that of the wild-type parasite, while entry into and calcium-induced egress from the host cell are not affected

Parasite motility is critical for invasion into and egress from the host cell [4-9, 12, 13, 34]. In addition, as the cortical microtubules are stable polymers that associate extensively with the parasite cortex, it is conceivable that they might provide mechanical support to the cortex when the parasites penetrate the membrane of the host cell during entry or exit. We therefore used video microscopy to determine whether the destabilization of the cortical microtubules in the TKO parasite might affect any of these processes. To examine parasite egress, the cultures infected by TKO or RH*Δhx* (wild-type parental) parasites were subjected to cold-treatment for 1 hr and then placed at 37°C for egress induced by the calcium ionophore A23187. Both TKO and wild-type parasites egressed actively and efficiently (Fig 4A, Video S5). Within ~ 1 min, egress was completed in >93% of vacuoles for the wild-type parasites (n=15) and in 100% of vacuoles for the TKO parasites (n=15) (Fig 4B).

To assess the parasite behavior during entry into the host cell, freshly harvested extracellular parasites were settled onto the host cell monolayer by a ~ 10 min pre-incubation on ice and then imaged by DIC at 37°C (Fig 4C, Video S6). The number of entry events was recorded for four parasite lines: wild-type RH*Δhx* parental, TKO, wild-type RH*ΔhxΔku80* parental, and *Δdcx* parasite. We observed 58 events for RH*Δhx* in 10 videos, 40 for TKO in 13 videos, 56 for RH*Δku80* in 9 videos, and 35 for *Δdcx* in 22 videos. The median speed of host cell entry was 25.5 second for RH*Δhx*, 22.5 second for the TKO, 25 second for the RH*ΔhxΔku80* and 33 second for the *Δdcx* parasites (Fig 4D). We also did not observe any notable differences in morphological changes (*e.g.* the formation and resolution of constrictions) during host cell entry among these parasite lines.

The lower incidence of entry attempts observed in the time-lapse experiments for the
TKO and Δ*dcx* indicates lower overall infection efficiency. Indeed, Δ*dcx* parasite was shown in
our previous work to invade at ~ 25% of the efficiency of the wild-type parasites when
assessed by an immunofluorescence based assay [33]. Consistent with lower efficiency in
infection, the TKO parasite generates ~ 60% of the number of plaques as the wild-type
parasite (p ≈ 0.005) when examined by the plaque assay, where the parasites were incubated
with the host cell monolayer unperturbed for 8 days (Fig 4E). The total area of the host cell
monolayer lysed by the TKO is 40% of the wild-type parasite (p < 0.0001), indicating significant
lower cytolytic efficiency. Thus, regarding the parasite's ability to move helically or conoid
structure, the short-distance host cell entry is less sensitive than infection efficiency.

DISCUSSION
274

The array of cortical microtubules is one of the major features in the cytoskeleton of
Toxoplasma and other apicomplexan parasites. In *Toxoplasma,* the roots of the cortical
microtubules are precisely located at evenly spaced intervals separated by cogwheel-like
projections at the circumference of the apical polar ring [14-16]. The cortical microtubules are
co-constructed and associated extensively with the inner membrane complex (IMC), guiding its
formation during parasite replication [24, 25]. The inhibition of microtubule polymerization in
the daughters results in distorted daughter cortex and failed cell division [26-29]. In mature
Toxoplasma, the cortical microtubules are extraordinarily stable and resistant to drug or cold-
temperature treatments that readily depolymerize mammalian microtubules [26, 28, 29]. It was
proposed that the cortical microtubules might have various mechanical roles in facilitating
parasite motility and invasion [30, 31, 35]. However, the extraordinary stability of these
polymers poses a major barrier against directly testing these hypotheses.

Previously, we generated a mutant (Δ*tlap2*Δ*spm1*Δ*tlap3,* a.k.a." TKO") in which the
cortical microtubules are destabilized in the mature parasites while continuing to grow in
developing daughters [22]. Here, by quantifying the proportion of TKO parasites with different
levels of defects in the array of cortical microtubules, we now have a more complete view of
the extent of depolymerization in the non-dividing and dividing parasites. We found that 1) ~
80% of untreated, non-dividing parasites have severely shortened microtubules and 2) the
extent of depolymerization in the mature parasite becomes increasingly marked when the
daughters start to form, but the development of daughters is not affected. Examining cortical
microtubules in the wild-type also revealed a small proportion of mature parasites with
shortened cortical microtubules at the end of daughter construction. This suggests that
microtubule depolymerization in the mother parasite might be a part of the normal preparation
for disassembling maternal structures into components parts for possible recycle in the new
generation. The cortical microtubules in mature TKO parasites are likely further sensitized to
the depolymerizing activity due to the lack of TLAP2, SPM1, and TLAP3. Remarkably, in the
same cell, while the maternal microtubules are destabilized, the daughter microtubules grow
normally. This indicates the local nature of this activity, possibly due to either depolymerizing
factors specific for the mother parasite and/or stabilizing factors specific for daughter cells.

When treated with low temperature, the cortical microtubules in the TKO parasite can
be fully depolymerized [22]. Here we show that this depolymerization is reversible, when the
parasite was placed at 37°C after cold-treatment, short cortical microtubules are found
concentrated at the apical end of the parasite. This is consistent with the current hypothesis

that the apical polar ring is the nucleating center for the cortical microtubules [14-16]. It is
tempting to suggest that the nucleation complexes in the apical polar ring in the mature
parasite remain active and are capable of re-nucleating new cortical microtubules after
complete depolymerization. However, we cannot exclude the possibility that the new growth
occurred from short microtubules that remain associated with the apical polar ring, as a small
cap of tubulin antibody labeling remains at the apical end even after prolonged cold treatment.
If the latter were true, it suggests differential regulation of stability along the length of the
cortical microtubules.
318
In the 3-D Matrigel, the TKO mutant moves much more linearly than the wild-type
parasite, which travels along an irregular helical path. The stable and spirally arranged cortical
microtubules, with their extensive interactions with the cortex, could contribute to the helicity of
the parasite movement by maintaining non-isotropic cortical properties (e.g. stiffness), and
thus biasing the orientation at which the parasite interacts with its environment. In plant cells,
the transmembrane cellulose synthesis complex is dynamically coupled to the cortical
microtubules via cellulose synthases interacting (CSI1) protein [36]. This results in the co-
alignment of the extracellular cellulose fibers with the cortical microtubules on the other side of
the plasma membrane. One could imagine that a similar mechanism might be involved in the
confinement of the force-generating actomyosin apparatus associated with the inner
membrane complex in *Toxoplasma*, although currently a CSI1-like coupler is unknown. The
trajectory of the parasite movement might also be affected by the parasite shape. For example,
in *Plasmodium berghei,* the knockout of IMC1h, which changes the shape of the ookinete,
results in more linear movement [37]. We do not think this is the likely explanation for the
change in movement trajectory of the TKO parasite, as the TKO parasite still maintains the
crescent shape characteristic of *Toxoplasma* (e.g. Video S1)*.* In addition, a number of
observations complicate the interpretation of the exact contribution of parasite shape in the
trajectory of movement. First, in the IMC1h *Plasmodium* mutant, while the cortical microtubules
are still present and have a recognizable twist, they are also disorganized, which makes it
difficult to disentangle the individual contributions from changes in parasite shape vs in the
organization of the microtubules. Second, while it has been shown that the TKO mutant
becomes shorter than the wild-type parasite when treated with low temperature [38], the
linearity of the movement of the TKO parasite does not differ significantly with or without
prolonged cold-treatment.
343
It was proposed that the cortical microtubules act as a spring to store and release
energy for helical gliding [31]. However, our data suggest that these microtubules do not
appear to act as a link in the energy relay vital for parasite movement. While the TKO mutant
does not move helically, it is clearly capable of sustaining long-distance movement even under
conditions where the TKO parasites were pre-treated with cold to nearly fully depolymerize the
microtubules. Therefore the cortical microtubules do not play a role in maintaining directional
long-distance movement. This means other higher-order structures are needed to limit the
direction of force generation by the actomyosin complex. Whatever these structures might be,
they need to be regularly arranged and stably associated with the parasite cortex. One
candidate is the array of inner membrane particles (IMPs), which form periodic rows of IMPs
that run the length along the longitudinal axis of the parasite [24]. Supporting this hypothesis,
the depletion of a potential IMP component (GAPM1a) in mature parasite results in decreased
parasite displacement in the 3-D Matrigel [38]. A subset of the IMP rows co-align with the
cortical microtubules. In the GAPM1a mutant, the cortical microtubules are disordered and
destabilized, also indicating intimate interaction between the two sets of structures. It is worth

noting that, in contrast to the GAPM1a mutant, the net displacement of the TKO parasite is
similar to that of the wild-type (Fig 3A). This separation of helicity from sustained parasite
movement might be due to the cell-stage dependent nature of the stability of the cortical
microtubules in the TKO parasite. The IMC cortex of the parasite is built concomitantly with the
cortical micotubules during daughter formation. Because the daughter cortical microtubules
grow normally in the TKO parasite, the structures in the IMC also likely form largely normally.
Thus, we think that, in the mature TKO parasites, we are seeing mainly the impact of the
microtubule destabilization itself, which manifests in the change from helical to linear
movement.
368
Given the precision by which the cortical microtubules are constructed, the
sophisticated suite of associated proteins [15, 21-23], and their extraordinary stability, we had
expected that they would provide important mechanical support for the cortex in the mature
parasite. To our surprise, the shape of the TKO parasite is not grossly different from the wild-
type parasite (e.g. Video S1), and there is also no detectable difference in the behavior of the
parasite during host cell entry and egress when the parasite experiences considerable
mechanical stress pushing through host cell cortex. This indicates that other cortical structures,
such as the inner membrane complex and associated intermediate filament-like network, are
more critical as structural support in mature parasites. Notably, the conoid structure mutant,
Δ*dcx,* which invades with a much lower efficiency [33], also showed wild-type like behavior
during entry. This suggests that host cell entry is considerably less sensitive to the change in
the structure of the cortical microtubules and conoid compared with extracellular migration
during infection.
382
Lastly, TLAP2, SPM1, and TLAP3, the three proteins collectively responsible for the
stability of cortical microtubules in mature *Toxoplasma*, have well-conserved orthologs in
Plasmodium spp [22]. It would be intriguing to determine whether the microtubule stabilizing
function of these proteins, and the influence of cortical microtubule stability to parasite
movement are also conserved among the apicomplexans. As the cortical microtubules in the
developing daughter and the mature TKO parasites display distinct polymerization and
depolymerization states, identifying the activities responsible for this maternal-filial distinction
will generate insights into how the cortical microtubules of the mother and developing
daughters (composed of the same polymer and less than half a micron apart) coexist in distinct
biochemical states and how the switch from one state to the other occurs as the daughters
become mature.
394
395
396 **MATERIALS AND METHODS**
397
398 *T. gondii mutant strains and host cell cultures*
399
400 Host cell and *T. gondii* tachyzoite parasite cultures were maintained as described previously
401 [15, 22, 34, 39]. The TKO (Δ*tlap2*Δ*spm1*Δ*tlap3*) mutant was generated in [22] and the Δ*dcx*
402 parasite was generated in [33].
403
404 *Temperature treatment of T. gondii for testing microtubule stability*
405
406 Cultures of intracellular parasites growing in HFF cells plated in 3.5-cm glass-bottom dishes
407 (MatTek Corporation, CAT# P35G-1.5-14-C or CAT# P35G-1.5-21-C) were incubated at 7°C
408 for 20, 60, 90 or 120 min in the L15 imaging media, which is Leibovit's L-15 (21083-027,
409 Gibco- Life Technologies, Grand Island, NY) supplemented with 1% (vol/vol) heat-inactivated

cosmic calf serum (CCS; SH30087.3; Hyclone, Logan, UT). For recovery, the intracellular parasites incubated at 7°C for 2 hours were subsequently placed at 37°C for 5, 10, 20, 30, 60 or 120 min in the L15 imaging media. The samples were then processed for immunofluorescence as described below. Total number of parasites analyzed: 444, 422, 468, 423 of TKO and 246, 274, 300, 390 of wild-type cold-treated at 7°C for 20, 60, 90 or 120 min, respectively; 440, 428, 456, 496, 396, 360 of TKO, and 295, 168, 172, 166, 205, 230 of wild-type recovered at 37°C for 5, 10, 20, 30, 60, or 120 min, respectively.

Immunofluorescence

All immunofluorescence labeling steps were performed at room temperature. For intracellular parasites, *T. gondii*–infected HFF monolayers growing in glass-bottom dishes were fixed in 3% (vol/vol) formaldehyde and 50% (vol/vol) methanol in 0.5X PBS for 15 min, permeabilized in 0.5% (vol/vol) Triton X-100 (TX-100) in PBS for 15 min, washed with PBS twice, followed by incubating with the primary antibody (mouse monoclonal anti-acetylated tubulin, 6-11B-1, 1:1000, Sigma T6793) and then with the secondary antibody (goat anti-mouse IgG- Alexa Fluor 488, A11029, Molecular Probes) at 1:1000, for 1 hr each. For labeling of the Inner Membrane Complex Protein 1 (IMC1), the samples were subsequently blocked with 1.5% BSA in PBS for 1 hour, then incubated for 1 hour with rabbit anti-IMC1 (a kind gift from Dr. Con Beckers, William Carey University) at 1:1000, and finally with goat anti-rabbit IgG -Alexa Fluor 568 (A11036, Molecular Probes) at 1:1000 for 1 hour. The antibodies were diluted in the blocking solution (i.e. 1.5% BSA in PBS).

Three-dimensional structured-illumination microscopy

Three-dimensional structured-illumination microscopy (3D-SIM) image stacks were collected at z-increments of 0.125 µm using a DeltaVision OMX Flex imaging station (GE Healthcare-Applied Precision). A 60x oil immersion lens (numerical aperture [NA] 1.42) and immersion oil at refractive index 1.520 were used. Structured illumination image reconstructions and 3D projections were processed using the functions and software supplied by the manufacturer.

Replication assay

The intracellular replication assay was performed as described in [5]. Four independent experiments were performed. In each replicate, the number of parasites in ~ 100 vacuoles was counted for each strain at each time point.

Egress assay

For induced egress assay, parasites were added to MatTek dishes containing a confluent HFF or BS-C-1 monolayer and grown for ~ 35 hours. Cells were washed once and incubated in 2 ml of L15 imaging media and placed at ~7°C for 1 hr, before being moved to the environmental chamber kept at 37ºC which was then replaced with ~ 2 ml of 5 µM A23187 in L15 imaging media. Images were collected at 37ºC with 1 sec interval for 10 minutes on an Applied Precision Delta Vision imaging station equipped with an environmental chamber.

Live invasion assays

To analyze the behavior of live parasites during invasion, extracellular parasites were used to infect a MatTek dish with a nearly confluent monolayer of HFF cells. The dishes were incubated on ice for ~10 min to facilitate parasite settling on the host cells. The culture was then imaged at 37°C with 1 sec intervals for 15 min to capture invasion events.

Plaque assays

A total of 50, 100 and 150 freshly harvested parasites were added to each well of 6-well plates with confluent HFF monolayers and grown undisturbed for 8 days at 37°C, then fixed with 70% ethanol for 15 min and stained with 1% (wt/vol) crystal violet in 25% (vol/vol) methanol, rinsed with PBS, air-dried, and imaged.

3-D motility assay and image acquisition

The 3-D motility assay for *Toxoplasma,* originally developed in [3], was carried out as described in [34]. For cold-treated parasites, cultures with intracellular parasites were placed at ~ 7°C for 2 hr, before harvesting as described in [34]. Parasites were harvested and imaged immediately at 37°C. For each imaging experiment, 151-181 image stacks were acquired at 1.36 sec intervals. Each stack consisted of 41 optical sections, spaced 1 μm apart in the Z-plane (40 μm stacks), with field of view of 1500x1500 pixels, binning 2x2, with a final dimension of 750 x 750 pixels. Image stacks were acquired from at least ~10 μm above the coverslip. To represent the 3-D DIC images in 2-D format (Fig 2 and Video S1&3), we performed focus-stacking using Fiji by applying the Extended Depth of Field Sobel Projection plugin from [40, 41]

DIC image stacks processing and cell tracking

DIC images were processed using our custom scripts extended from [34]. High contrast images were generated by subjecting the DIC images to 1) a 3-D median filter (footprint nx = ny = nz = 2 px) to reduce noise, 2) a 2-D Laplacian filter to attenuate out-of-focus portions of the cells, and 3) a 3-D gradient filter of Sobel type for edge detection. A dilation and erosion process using a maximum filter followed by a minimum filter (footprint nz = 3 px, nx = ny =13 px) was then applied. 2 pixels from all x and y edges were cropped before tracking to avoid edge artifacts. We implemented the imaging filters using Python 3 to enable GPU-acceleration using the CuPy plugin [42-44]. The TrackMate plugin in Fiji was then used to detect and track the parasites in the processed image stacks [40, 45, 46]. The Difference-of-Gaussian (DoG) detector was employed for spot detection with a diameter of 7 μm. To avoid artifacts associated with edge-effects and out-of-focus signal, we excluded spots detected within the first and last 1 μm of the z-dimension and 3 μm of each of the x and y dimension. The Kalman filter tracker was used to reconstruct the cell tracks, and false reconstructions (i.e., "tracks" between debris or random noise) were filtered out based on quality, track length, and duration. The values for these parameters were determined empirically. The resulting tracks were manually verified and errors such as misdetections and gaps corrected by adding or deleting individual spots. To refine spot positions, we used the TrackMate spot fitting feature to fit 3-D Gaussian with fixed radius on each spot. After conducting a second round of manual verification, we then separated the motile parasite tracks from the non-motile tracks by applying another set of filters based on displacement and speed. The 3-D coordinates of both the non-motile and motile tracks were copied to different comma-separated value text files (.csv) for analysis in R version 4.1.2 in RStudio [47, 48].

Trajectory analysis

We used the R package celltrackR v 1.1.0 and our own custom functions to process and analyze the parasite trajectories [47-49]. The complete list of the rest of the R packages we used for our analysis is at the end of this section. First, the global drifts in the data were corrected according to the workflow developed by [49] and we calculated the drift velocity from

the corresponding set of non-motile tracks. Then, non-motile ends (the very start or very end) of the parasite trajectories were trimmed to eliminate biases introduced by the asynchronous nature of the parasite motility. After trimming, only tracks with more than four steps (i.e., five tracking points) were included in the subsequent analysis.

In Fig 3A, the net displacement is a simple Euclidean distance between the first point and the last point of the trajectories. The run length of the parasite is the total distance traveled between two "pauses", which we defined as an event when the parasite stops moving for more than 5.4 sec (>= 4 frames). "stop moving" means that when the speed of a step (sub-track of length 1) is lower than a threshold. For each data set, the threshold value was calculated by taking the average step-speed of the non-motile tracks. We removed the track points that correspond to pause events, introducing gaps. The motile segments were then obtained by splitting the tracks around those gaps. The run speed was obtained by dividing the motile segment's total distance (run length) by its total duration. For Fig 3C, the dispersal ratio was calculated by taking the net displacement of the motile segments and dividing it by the run length, where a value of 1 indicates a completely linear track.

For analyzing the turning angle (Fig 3C), any stops that were not filtered out before (i.e., stops that lasted less than 4 frames) and steps with displacement of less than 10% of the parasite body length (~ 0.75 µm) were cut out from the motile segments before splitting into sub-segments. Sub-segments that are less than 3-frames long were excluded from the turning angle analysis, which was done based on [49].

The pitch length is the height of one complete helical turn, which was quantified as the straight-line distance between the starting and ending points of the turn. These points were identified through visual inspection of the trajectories and parasite orientation in 3-D space for the pitch calculation in Fig 3C.

The 3-D reconstruction of the processed DIC image stacks, along with the tracks exported from TrackMate (Fig 2C, Fig 3B, Video S1-S4), was done using the 3-D viewer and Track Processor in Icy [50]. Gaussian blur with a sigma of 2 was applied to the processed stacks before Icy rendering.

The Kruskal-Wallis test was done to compare the groups for all of the track measurements in Fig 3A&C [47]. We used R version 4.1.2 [48] and the following R packages: data.table v. 1.14.6 [51], gridExtra v. 2.3 [52], onewaytests v. 2.7 [53], pracma v. 2.4.2 [54], rmarkdown v. 2.20 [55-57], rstatix v. 0.7.2 [58], scales v. 1.2.1 [59], tidyverse v. 2.0.0 [60].

Acknowledgments
We thank Dr. John M. Murray for critical discussions and readings of the manuscript, and Melissa Molina for tissue culture support. This study was supported by funding from the National Institutes of Health/National Institute of Allergy and Infectious Diseases (R01-AI132463) awarded to K.H.

Conflict of Interest Statement
The authors declare that they have no conflict of interest.

Figure Legends

Figure 1. The destabilization of cortical microtubules in mature TKO parasites
A. Projections of image stacks acquired by three-dimensional structured illumination microscopy (3D-SIM) of the non-dividing wild-type and TKO parasites. While 100% of the wild-type parasites have the typical spirally arranged full microtubule array, ~79% (±8.5% SEM) of the TKO parasites have significantly curtailed/short microtubules. The cortical microtubules are sparse in the rest 21%. Green: tubulin labeling by a mouse anti-acetylated tubulin antibody, which also labels the microtubules in the host cell (arrowhead). Red: cortex labeling by a rabbit anti-IMC1 antibody.
B. Diagrams of wild-type parasites at the initiation (INT), elongation (EL), pre-emergence (pre-EM), and emergence (EM) stages of daughter construction. Cortical microtubules and the conoid are shown for the mother and daughter parasites.
C. *Left*: Projections of 3D-SIM images of the TKO and wild-type parasites at the initiation, elongation, pre-emergence, and emergence stage of daughter construction. Hollow arrow: cortical microtubules in the mother parasite. Arrow: daughter parasites. Arrowhead: host cell microtubules. Insets (0.7X) are projections that include a subset of the image stack (sub pj) to better display the structure of the daughter parasites, which are obscured in the full projections (full pj) for the wild-type parasites due to the cortical microtubules in the mother. Note that for the TKO parasite, due to the shortness or complete absence of the maternal microtubules, the daughters are as fully visible in the full projection as in the subset projection.
Right: Quantification of the proportions of wild-type (top) and TKO (bottom) parasites that display various characteristics in the array of cortical microtubules at different stages, including the non-dividing (non-DIV) stage or the initiation (INT), elongation (EL), pre-emergence (pre-EM), and emergence (EM) stages of daughter construction. For the TKO parasites, the two patterns observed are "short or sparse" (*i.e.* similar to non-dividing TKO parasite) and "absent". For the wild-type parasites, the two patterns observed are "non-dividing wild-type-like" or "short or sparse". Error bars: standard error from three independent experiments. Total number of parasites analyzed: 88, 68, 50, 36 of TKO and 52, 62, 54, 49 of wild-type at the initiation, elongation, pre-emergence, or emergence stage, respectively. The data for the non-dividing stage is the same as shown in Fig 1A.
D. Average number of cell divisions at 12, 24 or 36 hrs after infection in four independent experiments for RH*Δhx* (*WT*) and TKO parasites. Error bars: standard deviation
E. *Left*: Quantification of the percentage of non-dividing wild-type and TKO parasites that display various characteristics in the array of the cortical microtubules after being placed at ~ 7°C for 0, 20, 60, 90 or 120 min (C0-C120) as well as after recovery at 37°C for 5, 10, 20, 30, 60, or 120 min after 120 min of ~ 7°C incubation (R5-R120). For the TKO parasites, the two patterns observed are "short or sparse" or "absent". For wild-type parasites, the cortical microtubules remain similar to those in untreated parasites under all conditions tested here (after cold-treatment and cold-treatment plus 37°C recovery). Error bars: standard error from three independent experiments. *Middle*: projection of 3D-SIM images of TKO parasites in which cortical microtubules are absent. *Right*: projection of 3D-SIM images of TKO parasites after 1hr of 37°C (R60) recovery, in which short cortical microtubules at the apical end are seen. Arrowhead: host cell microtubules.

Figure 2. Workflow for image processing and automatic tracking of 3-D motility data acquired by DIC imaging
A. Schematic diagram of the image processing (red outline and box), detection and tracking (blue outline), and tracks analysis (black outline) workflow for 3-D motility data acquired by DIC

imaging. (Red) A representative area of a slice from a DIC image stack before processing and the resulting images corresponding to the filters applied on each processing step using Python3, showing the resulting high contrast image. For details, see Materials and Methods section.

B. Focused composite image of a DIC image stack (41 slices; 40 *μ*m) of extracellular wild-type parasites in Matrigel and corresponding processed image (represented by its maximum projection) (see also Video S1).

C. Examples of 3-D tracks for motile RH*Δhx* parental, TKO, RH*Δku80* parental and *Δdcx* parasites reconstructed in the Icy software (see also Video S1- S4). Length of arrow = 16 μm.

Figure 3 The 3-D trajectories of TKO parasites are significantly more linear than those of parental parasites.

A. Frequency distribution and statistical analysis of motion-related parameters including net displacement, run length (distance between pauses >5.4 sec), and run speed of RH*Δhx* parental, TKO, RH*ΔhxΔku80* parental and *Δdcx* parasites. The mean values for individual histograms are shown (in red) and also marked by the red dashed line. ns: p-value not significant. P-values were calculated by the Kruskal-Wallis test.

B. Representative tracks projected on the (x, y) plane for RH*Δhx* parental, TKO and cold-treated TKO, RH*Δku80* parental and *Δdcx* parasites with superimposed origins (red circles). For cold-treated TKO, intracellular parasite cultures were placed at 7 degree for two hours before harvesting and mixed with Matrigel. For the other samples/conditions, the parasites were harvested as described previously in [34]. All motility assays were carried out at 37°C. Bottom right: enlarged insets (~ 6X) that include the segment indicated by the dashed boxes in the trajectories for RH*Δhx* parental, TKO and cold-treated (CT) TKO parasites.

C. Frequency distribution and statistical analysis of helical curve-related parameters including turning angle, dispersal ratio, and the pitch of helical turn of RH*Δhx* parental, and TKO, with an illustration defining pitch. Data for all parameters are also displayed for RH*Δku80* parental and *Δdcx* parasites except for the pitch. The mean values for individual histograms are shown (in red) and also marked by the red dashed line. *p<0.05; ****p<0.0001. p values were calculated by the Kruskal-Wallis test.

Figure 4. The speeds of entry and egress of TKO parasites are similar to those of parental parasites despite lower plaquing efficiency

A. Images selected from time-lapse experiments of intracellular RH*Δhx* parental and TKO parasite treated with 5 µM A23187 (also see Video S5). Parasitophorous vacuoles are indicated by the orange arrows.

B. Dot plots of time taken to disperse after treatment with 5 µM A23187. The intracellular wild-type and TKO parasite cultures were placed at ~ 7°C for 1 hr prior to the egress experiment, which was conducted at 37°C. Error bars: standard error.

C. Images selected from time-lapse recording of RH*Δhx* parental, TKO, RH*Δku80* parental and *Δdcx* parasites in the process of invasion (also see Video S6). Frames marked in yellow: entry completed. Green arrows: constrictions formed during invasion.

D. Dot plots of time taken for entry (i.e. time from parasite contact with the host cell at its apical end to completion of entry) for RH*Δhx* parental, TKO, RH*ΔhxΔku80* parental and *Δdcx* parasites. Error bars: standard error.

E. *Top*: Plaques formed by the RH*Δhx* parental and TKO parasites. "plaques" are areas of the host cell culture lysed by repeated cycles of parasite invasion, replication, and egress. The cultures were fixed with 70% ethanol and stained with crystal violet 8 days after inoculation. *Bottom*: Bar graphs that compare the number of plaques (left) and total plaque area (right) of

659 the RH*Δhx* parental and TKO parasites. Five independent experiments were conducted. Error
660 bars: standard error. **p<0.01; ****p<0.0001. p values were calculated by unpaired Student's t-
661 tests.
662
663
664

Table 1. Fraction of motile parasites in 3-D motility assay

Parasite strain	Mean percent motile ± SEM (# of independent replicates)	Total number of parasites analyzed
RH*Δhx* (TKO parental)	40.2 ± 6.36 (*N=6*)	413
TKO	42.3 ± 6.77 (*N=6*)	395
RH*ΔhxΔku80* (*Δdcx* parental)	36.5 ± 4.00 (*N=8*)	404
Δdcx	30.9 ± 4.97 (*N=9*)	486

665
666
667 **Supplemental Videos:**
668
669
670 **Video S1:** Time-lapse microscopy of 3-D motility in 50% Matrigel for the RH*Δhx* (RH parental
671 1) and TKO parasites. *Left:* Videos of projected DIC images (projections of 3-D stack
672 generated by Stack focuser in ImageJ/Fiji). *Right:* Corresponding processed images with a
673 subset of 3-D trajectories. Time interval: ~ 1.36 sec. Video speed: 10 frames/s.
674
675 **Video S2:** Rotational view of 3-D trajectories for the RH*Δhx* (RH parental 1) and TKO
676 parasites included in Video S1. The traveling sphere associated with each track represents the
677 centroid of the parasite traveling along that path. Length of arrow= 16 μm
678
679 **Video S3:** Time-lapse microscopy of 3-D motility in 50% Matrigel for the RH*Δku80* (RH
680 parental 2) and *Δdcx* (*dcx KO*) parasites. *Left:* Videos of projected DIC images (projections of
681 3-D stack generated by Stack focuser in ImageJ/Fiji). *Right:* Corresponding processed images
682 with a subset of 3-D trajectories. Time interval: ~ 1.36 sec. Video speed: 10 frames/s.
683
684 **Video S4:** Rotational view of 3-D trajectories for the RH*Δku80* (RH parental 2) and *Δdcx* (*dcx
685 KO*) parasites included in Video S3. The traveling sphere associated with each track
686 represents the centroid of the parasite traveling along that path. Length of arrow= 16 μm
687
688 **Video S5:** Time-lapse microscopy of A23187 induced-egress for RH*Δhx* (RH parental) and
689 TKO parasites. A23187 was added at the beginning of the movies to a final concentration of 5
690 μM. The intracellular wild-type and TKO parasite cultures were placed at 7°C for 1 hr prior to
691 the egress experiment, which was conducted at 37°C. Time interval: 1 sec. Video speed: 60
692 frames/s. Scale bar: 5 μm.
693
694 **Video S6:** Time-lapse microscopy of host cell entries for RH*Δhx* (RH parental 1), TKO and
695 RH*Δku80* (RH parental 2) and *Δdcx* (*dcx KO*) parasites. Fig 4C includes images selected from
696 the timelapses in the leftmost column. Time interval: 1 sec. Video speed: 6 frames/s. Scale
697 bar: 5 μm.
698

REFERENCES

1. World-Health-Organization, *World Malaria Report 2022.* 2022: p. 293.
2. Dubey, J.P., *The history of Toxoplasma gondii--the first 100 years.* The Journal of eukaryotic microbiology, 2008. **55**(6): p. 467-75.
3. Leung, J.M., M.A. Rould, C. Konradt, C.A. Hunter, and G.E. Ward, *Disruption of TgPHIL1 alters specific parameters of Toxoplasma gondii motility measured in a quantitative, three-dimensional live motility assay.* PloS one, 2014. **9**(1): p. e85763.
4. Dobrowolski, J.M., V.B. Carruthers, and L.D. Sibley, *Participation of myosin in gliding motility and host cell invasion by Toxoplasma gondii.* Mol Microbiol, 1997. **26**(1): p. 163-73.
5. Heaslip, A.T., M. Nishi, B. Stein, and K. Hu, *The motility of a human parasite, Toxoplasma gondii, is regulated by a novel lysine methyltransferase.* PLoS Pathog, 2011. **7**(9): p. e1002201.
6. Meissner, M., D. Schluter, and D. Soldati, *Role of Toxoplasma gondii myosin A in powering parasite gliding and host cell invasion.* Science, 2002. **298**(5594): p. 837-40.
7. Gaskins, E., S. Gilk, N. DeVore, T. Mann, G. Ward, and C. Beckers, *Identification of the membrane receptor of a class XIV myosin in Toxoplasma gondii.* J Cell Biol, 2004. **165**(3): p. 383-93.
8. Graindorge, A., K. Frenal, D. Jacot, J. Salamun, J.B. Marq, and D. Soldati-Favre, *The Conoid Associated Motor MyoH Is Indispensable for Toxoplasma gondii Entry and Exit from Host Cells.* PLoS Pathogens, 2016. **12**(1): p. e1005388.
9. Jacot, D., N. Tosetti, I. Pires, J. Stock, A. Graindorge, Y.F. Hung, et al., *An Apicomplexan Actin-Binding Protein Serves as a Connector and Lipid Sensor to Coordinate Motility and Invasion.* Cell Host & Microbe, 2016. **20**(6): p. 731-743.
10. Huynh, M.-H. and V.B. Carruthers, *Toxoplasma MIC2 is a major determinant of invasion and virulence.* PLoS Pathogens, 2006. **2**(8): p. e84.
11. Huynh, M.H., K.E. Rabenau, J.M. Harper, W.L. Beatty, L.D. Sibley, and V.B. Carruthers, *Rapid invasion of host cells by Toxoplasma requires secretion of the MIC2-M2AP adhesive protein complex.* Embo J, 2003. **22**(9): p. 2082-90.
12. Frenal, K., V. Polonais, J.B. Marq, R. Stratmann, J. Limenitakis, and D. Soldati-Favre, *Functional dissection of the apicomplexan glideosome molecular architecture.* Cell Host Microbe, 2010. **8**(4): p. 343-57.
13. Tosetti, N., N. Dos Santos Pacheco, D. Soldati-Favre, and D. Jacot, *Three F-actin assembly centers regulate organelle inheritance, cell-cell communication and motility in Toxoplasma gondii.* eLife, 2019. **8**: p. e42669.
14. Morrissette, N.S. and L.D. Sibley, *Cytoskeleton of apicomplexan parasites.* Microbiol Mol Biol Rev, 2002. **66**(1): p. 21-38.
15. Leung, J.M., Y. He, F. Zhang, Y.C. Hwang, E. Nagayasu, J. Liu, et al., *Stability and function of a putative microtubule organizing center in the human parasite Toxoplasma gondii.* Molecular Biology of the Cell (Featured as Journal Cover), 2017. **28**(10): p. 1361-1378.
16. Nichols, B.A. and M.L. Chiappino, *Cytoskeleton of Toxoplasma gondii.* J. Protozool., 1987. **34**(2): p. 217-26.
17. Wang, X., P. Qian, H. Cui, L. Yao, and J. Yuan, *A protein palmitoylation cascade regulates microtubule cytoskeleton integrity in Plasmodium.* The EMBO journal, 2020. **39**(13): p. e104168-e104168.
18. Bounkeua, V., F. Li, and J.M. Vinetz, *In vitro generation of Plasmodium falciparum ookinetes.* The American journal of tropical medicine and hygiene, 2010. **83**(6): p. 1187-1194.

19. Garnham, P.C., R.G. Bird, and J.R. Baker, *Electron microscope studies of motile stages*
of malaria parasites. III. The ookinetes of Haemamoeba and Plasmodium. Trans R Soc
Trop Med Hyg, 1962. **56**: p. 116-20.
20. Bannister, L.H. and G.H. Mitchell, *The role of the cytoskeleton in Plasmodium*
falciparum merozoite biology: an electron-microscopic view. Annals of tropical medicine
and parasitology, 1995. **89**(2): p. 105-11.
21. Liu, J., L. Wetzel, Y. Zhang, E. Nagayasu, S. Ems-McClung, L. Florens, et al., *Novel*
Thioredoxin-Like Proteins Are Components of a Protein Complex Coating the Cortical
Microtubules of Toxoplasma gondii. Eukaryotic cell, 2013. **12**(12): p. 1588-99.
22. Liu, J., Y. He, I. Benmerzouga, W.J. Sullivan, Jr., N.S. Morrissette, J.M. Murray, et al.,
An ensemble of specifically targeted proteins stabilizes cortical microtubules in the
human parasite Toxoplasma gondii. Molecular Biology of the Cell (Featured as Journal
Cover), 2015. **27**(3): p. 549-71.
23. Tran, J.Q., C. Li, A. Chyan, L. Chung, and N.S. Morrissette, *SPM1 stabilizes*
subpellicular microtubules in Toxoplasma gondii. Eukaryot Cell, 2012. **11**(2): p. 206-16.
24. Morrissette, N.S., J.M. Murray, and D.S. Roos, *Subpellicular microtubules associate*
with an intramembranous particle lattice in the protozoan parasite Toxoplasma gondii. J.
Cell Sci., 1997. **110**(Pt 1): p. 35-42.
25. Hu, K., D.S. Roos, and J.M. Murray, *A novel polymer of tubulin forms the conoid of*
Toxoplasma gondii. J Cell Biol, 2002. **156**(6): p. 1039-50.
26. Morrissette, N.S. and L.D. Sibley, *Disruption of microtubules uncouples budding and*
nuclear division in Toxoplasma gondii. J Cell Sci, 2002. **115**(Pt 5): p. 1017-25.
27. Stokkermans, T.J., J.D. Schwartzman, K. Keenan, N.S. Morrissette, L.G. Tilney, and
D.S. Roos, *Inhibition of Toxoplasma gondii replication by dinitroaniline herbicides.* Exp.
Parasitol., 1996. **84**(3): p. 355-70.
28. Hu, K., J. Johnson, L. Florens, M. Fraunholz, S. Suravajjala, C. DiLullo, et al.,
Cytoskeletal components of an invasion machine – the apical complex of Toxoplasma
gondii. PloS pathogens, 2006. **2**(2): p. 0121-0139.
29. Hu, K., *Organizational Changes of the Daughter Basal Complex during the Parasite*
Replication of Toxoplasma gondii. PLoS Pathogens, 2008. **4**((1)): p. 108-121.
30. Stadler, R.V., L.A. White, K. Hu, B.P. Helmke, and W.H. Guilford, *Direct measurement*
of cortical force generation and polarization in a living parasite. Mol Biol Cell, 2017. **E16-**
07-0518.
31. Pavlou, G., B. Touquet, L. Vigetti, P. Renesto, A. Bougdour, D. Debarre, et al., *Coupling*
Polar Adhesion with Traction, Spring, and Torque Forces Allows High-Speed Helical
Migration of the Protozoan Parasite Toxoplasma. ACS Nano, 2020. **14**(6): p. 7121-
7139.
32. Swedlow, J.R., K. Hu, P.D. Andrews, D.S. Roos, and J.M. Murray, *Measuring tubulin*
content in Toxoplasma gondii: a comparison of laser-scanning confocal and wide-field
fluorescence microscopy. Proc Natl Acad Sci U S A, 2002. **99**(4): p. 2014-9.
33. Nagayasu, E., Y.-c. Hwang, J. Liu, J. Murray, and K. Hu, *Loss of a doublecortin (DCX)*
domain containing protein causes structural defects in a tubulin-based organelle of
Toxoplasma gondii and impairs host cell invasion. Molecular Biology of the Cell, 2016.
E16-08-0587
34. Munera Lopez, J., I.F. Tengganu, J. Liu, J.M. Murray, L.F. Arias Padilla, Y. Zhang, et
al., *An apical protein, Pcr2, is required for persistent movement by the human parasite*
Toxoplasma gondii. PLoS Pathog, 2022. **18**(8): p. e1010776.

35. Del Rosario, M., J. Periz, G. Pavlou, O. Lyth, F. Latorre-Barragan, S. Das, et al.,
Apicomplexan F-actin is required for efficient nuclear entry during host cell invasion.
EMBO reports, 2019. **20**(12): p. e48896.

36. Bringmann, M., B. Landrein, C. Schudoma, O. Hamant, M.T. Hauser, and S. Persson,
Cracking the elusive alignment hypothesis: the microtubule-cellulose synthase nexus
unraveled. Trends Plant Sci, 2012. **17**(11): p. 666-74.

37. Kan, A., Y.-H. Tan, F. Angrisano, E. Hanssen, K.L. Rogers, L. Whitehead, et al.,
Quantitative analysis of Plasmodium ookinete motion in three dimensions suggests a
critical role for cell shape in the biomechanics of malaria parasite gliding motility.
Cellular Microbiology, 2014. **16**(5): p. 734-750.

38. Harding, C.R., M. Gow, J.H. Kang, E. Shortt, S.R. Manalis, M. Meissner, et al., *Alveolar*
proteins stabilize cortical microtubules in Toxoplasma gondii. Nature Communications,
2019. **10**(1): p. 401.

39. Roos, D.S., R.G. Donald, N.S. Morrissette, and A.L. Moulton, *Molecular tools for genetic*
dissection of the protozoan parasite Toxoplasma gondii. Methods Cell Biol., 1994. **45**: p.
27-78.

40. Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, et al.,
Fiji: an open-source platform for biological-image analysis. Nature methods, 2012. **9**(7):
p. 676-82.

41. Forster, B., D. Van De Ville, J. Berent, D. Sage, and M. Unser, *Complex wavelets for*
extended depth‐of‐field: A new method for the fusion of multichannel microscopy
images. Microscopy research and technique, 2004. **65**(1‐2): p. 33-42.

42. Harris, C.R., K.J. Millman, S.J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau,
et al., *Array programming with NumPy.* Nature, 2020. **585**(7825): p. 357-362.

43. Okuta, R., Y. Unno, D. Nishino, S. Hido, and Crissman. *CuPy : A NumPy-Compatible*
Library for NVIDIA GPU Calculations. 2017.

44. Van Rossum, G.a.D., Fred L., *Python 3 Reference Manual.* 2009, Scotts Valley, CA,
USA: CreateSpace.

45. Tinevez, J.Y., N. Perry, J. Schindelin, G.M. Hoopes, G.D. Reynolds, E. Laplantine, et
al., *TrackMate: An open and extensible platform for single-particle tracking.* Methods,
2017. **115**: p. 80-90.

46. Ershov, D., M.-S. Phan, J.W. Pylvänäinen, S.U. Rigaud, L. Le Blanc, A. Charles-
Orszag, et al., *TrackMate 7: integrating state-of-the-art segmentation algorithms into*
tracking pipelines. Nature Methods, 2022. **19**(7): p. 829-832.

47. RStudioTeam, *RStudio: Integrated Development for R.* RStudio, PBC, Boston, MA
2020. **URL http://www.rstudio.com/**.

48. Team, R.C., *{R}: A Language and Environment for Statistical Computing*, 2021, R
Foundation for Statistical Computing: Vienna, Austria.

49. Wortel, I.M.N., A.Y. Liu, K. Dannenberg, J.C. Berry, M.J. Miller, and J. Textor,
CelltrackR: An R package for fast and flexible analysis of immune cell migration data.
ImmunoInformatics, 2021. **1-2**: p. 100003.

50. de Chaumont, F., S. Dallongeville, N. Chenouard, N. Hervé, S. Pop, T. Provoost, et al.,
Icy: an open bioimage informatics platform for extended reproducible research. Nat
Methods, 2012. **9**(7): p. 690-6.

51. Dowle, M. and A. Srinivasan, *{data.table}: Extension of '{data.frame}'*, 2022.

52. Auguie, B., *{gridExtra}: Miscellaneous Functions for "{Grid}" Graphics*, 2017.

53. Dag, O., A. Dolgun, and N.M. Konar, *{onewaytests: An R Package for One-Way Tests*
in Independent Groups Designs}. {The R Journal}, 2018. **10**(1): p. 175-199.

54. Borchers, H.W., *{pracma}: Practical Numerical Math Functions*, 2022.

55.   Allaire, J.J., Y. Xie, J. McPherson, J. Luraschi, K. Ushey, A. Atkins, et al., *{rmarkdown}:*
Dynamic Documents for R, 2023.
56.   Xie, Y., J.J. Allaire, and G. Grolemund, *R Markdown: The Definitive Guide*. 2018, Boca
Raton, Florida: Chapman and Hall/CRC.
57.   Xie, Y., C. Dervieux, and E. Riederer, *R Markdown Cookbook*. 2020, Boca Raton,
Florida: Chapman and Hall/CRC.
58.   Kassambara, A., *{rstatix}: Pipe-Friendly Framework for Basic Statistical Tests*, 2023.
59.   Wickham, H. and D. Seidel, *{scales}: Scale Functions for Visualization*, 2022.
60.   Wickham, H., M. Averick, J. Bryan, W. Chang, L.D.A. McGowan, R. François, et al.,
Welcome to the {tidyverse}. Journal of Open Source Software, 2019. **4**(43): p. 1686.
854
855

Figure 1

Figure 2

A — **Python 3** — DIC image stacks processing

Raw DIC image (representative slice)

Background subtraction
1. Median
2. Laplacian

Edge detection
3. Sobel-Feldman

Dilation and erosion
4. Maximum
5. Minimum

Fiji; TrackMate

3-D Spots detection and tracking

R

Drift correction and further analysis of reconstructed tracks

B — DIC — processed — 20 µm

C — RHΔ*hx* — TKO — RHΔ*ku80* — Δ*dcx*

Figure 3

Figure 4