1 **Females exhibit smaller and less variable patterns of brain activation across a range of motor**

2 **tasks**

3 Justin W. Andrushko[1,2], Shie Rinat [1], Eric D. Kirby[3], Julia Dahlby [1], Chelsea Ekstrand[4,*], Lara A.
4 Boyd[1,2,*]
5

1. Department of Physical Therapy, Faculty of Medicine, University of British Columbia,
Vancouver, Canada
2. Djavad Mowafaghian Centre for Brain Health, University of British Columbia, Vancouver,
Canada
3. Faculty of Individualized Interdisciplinary Studies, Simon Fraser University, Burnaby, BC,
Canada
4. Department of Neuroscience, University of Lethbridge, Lethbridge, Alberta, Canada
13
14 * Co-senior authors
15
16
17
18 **Correspondence**
19
20 Lara A. Boyd, Ph.D.
21 Email: lara.boyd@ubc.ca
22 ORCID: https://orcid.org/0000-0002-2828-4549
23 Department of Physical Therapy
24 The University of British Columbia
25
26 and
27
28 Chelsea Ekstrand, Ph.D.
29 Email: Chelsea.ekstrand@uleth.ca
30 ORCID: https://orcid.org/0000-0002-1330-1924
31 Department of Neuroscience
32 The University of Lethbridge

**Keywords:**
Sex differences, functional magnetic resonance imaging, somatotopic motor mapping

**Abstract**

Past work has shown that there are both structural and functional differences in the brains of

females versus males (Ritchie *et al.*, 2018). Specifically, males have larger cortical and sub-cortical

volume and surface area (both total and subregional), while females have greater cortical thickness in

most brain regions.  Given these differences it is possible that males may show larger volumes of brain

activation and greater variability in patterns of brain activity. Yet to date little work has systematically

considered whether patterns of brain activity indexed with functional MRI differ between females and

males. The current study sought to remediate this issue by employing brain somatotopic motor

mapping from a previously published (Ma *et al.*, 2022) and openly available dataset to test differences

in patterns of functional brain activity associated with 12 voluntary movement patterns in females

versus males. We discovered that females exhibited smaller volumes of brain activation across all 12

movement tasks, and lower patterns of variability between female participants in 10 of the 12

movements. We also observed that in these data females had greater cortical thickness which is in

alignment with previous structural sex difference analyses (Ritchie *et al.*, 2018; Wierenga *et al.*, 2022).

Overall, these findings provide a basis for considering sex differences in future fMRI research and

provide a foundation of understanding for which various neurological pathologies may present different

between females and males.

## Introduction

Females and males have anatomical and functional differences in their nervous systems (Aboitiz *et al.*, 1992; Kulynych *et al.*, 1994; Swaab *et al.*, 2001). These differences manifest in lateralization, motor planning, motor skill, language, memory, and spatial ability (Diamond, 1991; Harasty *et al.*, 1997; Jones *et al.*, 2003; Andreano & Cahill, 2009; Cohen *et al.*, 2010). However, our understanding of the neuronal or somatotopic representations of the body within the brain with respect to biological sex is limited. Although there has been a push to modernize our understanding of the classic sensorimotor "homunculus", including the proposal of an updated female "hermunculus", the focus has been largely on the sexual and reproductive dimorphisms between the sexes and their respective neuronal representations (Di Noto *et al.*, 2013).

Sex differences in brain structure have been observed (Lotze *et al.*, 2019; Liu *et al.*, 2020; Wierenga *et al.*, 2022). Males show differences in grey matter volume, cortical surface area, and increased variability of these measures compared to females. Previous research suggests that males have significantly greater grey matter volume in areas such as the amygdala, temporal pole, fusiform gyrus, putamen, and premotor cortex than females (Lotze *et al.*, 2019; Liu *et al.*, 2020). Further, a large meta-analysis of 16,683 healthy participants found that males exhibit greater subcortical volume than females, with the largest differences found in the thalamus and pallidum, bilaterally (Wierenga *et al.*, 2022). Males also exhibit greater cortical surface area than females. Importantly, males not only exhibit greater mean differences in subcortical volume and cortical surface area as compared to females, but also significantly greater *variability* in these measures (Wierenga *et al.*, 2022). This variability is most pronounced in motor related regions such as the pallidum, right inferior parietal cortex, and paracentral region (Wierenga *et al.*, 2022). This suggests that, compared to females, males have less consistent brain structure, particularly in motor related regions.

Females also show important differences in grey matter volume and cortical thickness compared to males. Specifically, females have greater grey matter volume than males primarily in

prefrontal and superior parietal cortices, as well as the superior temporal sulcus, orbitofrontal cortex,

and posterior insula (Lotze *et al.*, 2019; Liu *et al.*, 2020). Females also exhibit greater cortical thickness

compared to males (Ritchie *et al.*, 2018; Wierenga *et al.*, 2022). Further, Wierenga et al. (2022) noted

regional differences in cortical thickness between females and males; females had greater cortex

thickness for 38 of 68 regions, which were primarily in the frontal and parietal cortices. These

structural differences suggest that brain function may also differ by sex, which in turn may have

important implications for understanding sex differences in brain health and disease.

Sex related differences may affect the efficiency of neural activation patterns, with females

potentially having more localized and precise, and males having more distributed and varied, neuronal

processing. Several sex differences in brain activation have been observed with functional magnetic

resonance imaging (fMRI) across a range of tasks. Sex differences in mental rotation tasks (Thomsen *et*

*al.*, 2000), auditory verbal working memory (Goldstein *et al.*, 2005), task switching (Kuptsova *et al.*,

2015), and resting-state spontaneous brain activity  (Xu *et al.*, 2015) have been observed. However, to

date, few have considered whether sex differences affect somatotopic sensorimotor mapping across the

entire brain. Researching sex specific patterns of brain activation with respect to area and variability

across a range of motor tasks may shed light on key differences in brain function that informs future

research. For instance, in fMRI research, a common approach to investigate brain activation patterns is

through region of interest (ROI) analyses. However, using ROIs that are derived from an atlas that is

not sex-specific could potentially lead to data selection bias that does not fully capture differences in

functional organization between females and males.

Using fMRI whole brain somatotopic motor mapping from a previously published and openly

available dataset [(Ma *et al.*, 2022), https://openneuro.org/datasets/ds004044/versions/2.0.3], the

current study investigated the differences in cortical activation patterns between females and males

during a variety of motor tasks. Based on previous observations of structural differences in the brain

between females and males, we hypothesized that across movement tasks, females would exhibit lower

variability and smaller volumes of brain activation than males, reflecting more focal neural processing.

This concerted effort to further quantify the anatomical localization and functional differences will aid

our understanding of the inherent differences in somatotopic maps between the sexes, and inform ROI

selection strategies for future fMRI research by illustrating whether sex specific somatotopic brain

parcellations are needed.

**Methods**

*Participants*

A total of 62 neurologically intact right-handed young adults ($22.76 \pm 2.22$ years) were used

from the Ma *et al.* (2022) database. Of the 62 participants, 34 were female ($22.56 \pm 2.16$ years), and 28

were male ($23.00 \pm 2.34$ years). The original study from which these data were collected was approved

by the Institutional Review Board at Beijing Normal University (Ma *et al.*, 2022).

*Motor tasks*

Twelve voluntary movement patterns were each performed for 16 seconds twice during every

fMRI scan. A total of six fMRI scans were acquired for each participant, resulting in a total of 12 event

blocks for each of the 12 movement types (i.e.,192 seconds of task evoked data for each movement).

These movement conditions included bilateral toes, ankles, fingers, wrists, forearms, upper arms, and

eyes, in addition to jaw, lip, and tongue movements, and separate unilateral movements for the left and

right legs. Detailed movement instructions can be found in table 1. All movements were performed at a

self-selected pace between 0.5 and 5 Hz. For further details about the nature of the experiment and

motor tasks please review the published dataset manuscript (Ma *et al.*, 2022).

**Table 1**. Movement conditions and instructions

| Body movements | Movement patterns and instructions |
| --- | --- |
| Eyes | Blink or saccade eyes. |
| Jaw | Bite or twist jaws. |
| Lips | Expand and contract the lips with the teeth being bitten and tongue still. |
| Tongue | Circular tongue with the teeth being bitten and lips closed. |
| Upper arms | Lift and lower the upper arms (maximum 20°) with the upper arms, forearms, wrist, and fingers straight. |
| Forearms | Flex and release both forearms at the elbow (maximum 20°) with the wrists and fingers straight. |

| | |
|---|---|
| **Wrists** | Pitch and roll both wrists with clenched fists. |
| **Fingers** | Clench and loose both fists. |
| **Left leg** | Lift and lower the left leg (maximum 10°) with the leg, ankle, and toes straight. |
| **Right leg** | Lift and lower the right leg (maximum 10°) with the leg, ankle, and toes straight. |
| **Ankles** | Dorsiflex and release both ankles. |
| **Toes** | Flex and extend the toes from both feet. |
| **Rest** | Fixate on the dot presented in the center of the screen. |

*Preprocessing structural MRI data*

The T1-weighted (T1w) anatomical scans were preprocessed using FSL's (Jenkinson *et al.*,

2012) anatomical preprocessing script *fsl_anat*. Briefly, this script performs image reorientation to

align with standard space template coordinates, crops the T1w image in the z-direction to remove the

neck and other non-brain aspects of the image, implements bias-field correction using FSL's FAST

(Zhang *et al.*, 2001), and then extracts the brain using a non-linear reverse transformation of a standard-

space brain mask.

*First level functional MRI processing*

The previously preprocessed and manually denoised fMRI data [for more information regarding

the original preprocessing and denoising please review Ma et al. (2022)] were linearly registered to the

bias-field corrected and brain extracted T1w image using FSL's FLIRT (Jenkinson & Smith, 2001;

Jenkinson *et al.*, 2002), and then non-linearly registered to the MNI152_2mm brain template using

FNIRT (Andersson *et al.*, 2007). Each of the 12 motor tasks were modelled using the event timing files

provided in the downloaded data with a double-gamma function and with a temporal derivative applied

to the model in FSL's FEAT (Woolrich *et al.*, 2001). White matter BOLD signal was not treated as a

nuisance regressor in the present work even though this remains a controversial technique (Rostrup *et*

*al.*, 2000; Tettamanti *et al.*, 2002; Helenius *et al.*, 2003; Mazerolle *et al.*, 2010, 2013; Newman *et al.*,

2010; Gawryluk *et al.*, 2011*a*, 2011*b*, 2014; Weis *et al.*, 2011; Ding *et al.*, 2013; Fabri *et al.*, 2014; Peer

*et al.*, 2017; Grajauskas *et al.*, 2019; Gore *et al.*, 2019; Frizzell *et al.*, 2020; Jiang *et al.*, 2021).

Therefore, to prevent a bias towards grey matter BOLD signal and to truly investigate whole-brain

fMRI signal we retained white matter BOLD in our analyses to avoid the potential historical blind spot

in fMRI research (Grajauskas *et al.*, 2019).

*Second level fixed effects functional MRI processing*

Since each participant performed the same tasks over six separate runs, a fixed effects analysis

was carried out in FEAT (Woolrich *et al.*, 2001) prior to performing group-level analyses. The fixed

effects analysis was used to average the parameter estimates of the task-based activations across the six

runs to create a single activation map and effect size for each motor task for every participant. These

averaged parameter estimates were then carried forward for group level analyses in FEAT (Woolrich *et*

*al.*, 2004).

*Structural MRI analysis*

T1w images for each participant were segmented using Fastsurfer (Henschel *et al.*,

2020). Overall brain volume without ventricles was first analyzed to determine if there were differences

between females and males in brain size. Next, cortical surface area, cortical thickness, and grey matter

volume for each hemisphere were extracted for analysis and normalized as a ratio to overall brain

volume on a participant-by-participant basis to account for differences in brain size. We then compared

each of these normalized metrics for each hemisphere between females and males separately using two-

tailed independent samples *t*-tests implemented using the *permutation_test* module from SciPy stats in

Python 3.10 (Virtanen *et al.*, 2020) with 20,000 permutations. A manual Bonferroni correction to adjust

for multiple comparisons was performed and a result was deemed to be significant if the *p*-value was $\leq$

0.007 ($\alpha = 0.05/7$).

*Group level functional MRI analysis*

For group-level analyses, two one-sample *t*-tests were performed to compute spatial maps for

males and females separately. Additionally, two one-tailed independent samples *t*-tests were carried out

to determine if there were differences in the somatotopic representations of each of the 12 motor

movements between male and females (*t*-test #1: females > males; *t*-test #2: males > females contrasts).

Non-parametric permutation testing was performed on each contrast using *randomise* with 5,000

permutations, and threshold free cluster enhancement was used for statistical inference with a family

wise error rate set to $\alpha \leq 0.05$ for each contrast.

*Sex difference analyses*

To address the question of sex differences for somatotopic motor mapping, two approaches

were taken. First, the two *t*-tests previously described (*t*-test #1: females > males; *t*-test #2: males >

females contrasts) were assessed and reported when a significant contrast was detected (figure 1).

Second, to further investigate sex differences, the mean activation map for each movement was

computed separately for males and females and the spatial maps were correlated to assess their degree

of similarity. To do this, for males and females the group-level *p*-value threshold statistical maps ($\alpha \leq$

0.05) for each movement condition were temporally merged together using *fslmerge*. Then correlation

analyses between the spatial maps was carried out with *fslcc*. The correlation coefficients for same

movement conditions between females and males were then reported and the inter-task correlations

were disgarded.

*Sex differences in brain activation variability*

To assess within group variability in motor responses, we performed pairwise correlations for

both females and males. To do so, we used the raw *t*-statistic image for each participant for each

contrast and calculated the voxel-to-voxel correlation of these images for each pair of participants

separately for each group using in-house python code. We then compared the pairwise correlations

between females and males for each contrast using two-tailed independent samples *t*-tests implemented

using the *permutation_test* module from SciPy stats in Python 3.10 (Virtanen *et al.*, 2020) with 20,000

permutations. A manual Bonferroni correction to adjust for multiple comparisons was performed and a

result was deemed to be significant if the *p*-value was $\leq 0.004$ ($\alpha = 0.05/12$).

**Results**

*Sex differences in brain structure*

Permutation *t*-testing with 20,000 permutations revealed sex differences in total brain volume

[Females: 1,099,719.242 ± 55040.247 mm$^3$; Males: 1,230,966.679 ± 110273.860 mm$^3$; $t(59) = -5.912$,

$p = 1.00E-04$, $d = -1.480$]. There was also a significant difference in normalized cortical thickness in

the left [Females: 2.18E-06 ± 1.17E-07; Males: 1.99E-06 ± 1.75E-07; $t(59) = 4.950$, $p = 1.00E-04$, $d =$

1.251] and right [Females: 2.19E-06 ± 1.17E-07; Males: 1.99E-06 ± 1.70E-07; $t(59) = 5.268$, $p =$

1.00E-04, d = 1.333] hemispheres with females having thicker cortex. However, no differences were

observed for normalized grey matter volume in the left [Females: 0.211 ± 0.008; Males: 0.214 ± 0.007;

$t(59) = 1.451$, $p = 0.155$, $d = -0.376$], or right [Females: 0.212 ± 0.008; Males: 0.215 ± 0.007; $t(59) = -$

1.740, $p = 0.089$, $d = -0.451$] hemisphere, or normalized surface area in the left [Females: 0.075 ±

0.002; Males: 0.075 ± 0.002; $t(59) = 0.653$, $p = 0.509$, $d = 0.165$], or right [Females: 0.075 ± 0.002;

Males: 0.075 ± 0.002; $t(59) = 0.653$, $p = 0.509$, $d = 0.165$] hemisphere. These findings suggest that

although males have larger overall brains, females exhibited thicker cortex in both hemispheres.

*Sex differences for group-level contrasts*

Significant differences were found in male > female contrasts in seven out of the 12 movements

including ankles, fingers, jaw, lips, tongue, upper arms, and wrists (figure 1; supplementary data tables

S1-S8). The only contrast that showed a significant female > male difference was right leg movements.

It is important to note that the figure only shows six of the eight contrasts. The two contrasts not

represented only contained cluster sizes of 1 and 2 voxels respectively and were not visible in the

figures (see supplementary data tables S1-S8).

**Figure 1.** *Sex difference contrasts for 6 of the 8 significant conditions. The other two contrasts were not visible and are therefore found in the supplemental data. Significantly greater BOLD signal for males (Red/Yellow) and females (White/Blue) are plotted. Images are family-wise error corrected p-value maps with a p value threshold of 0.05. Figure was made in MRIcoGL (Rorden & Brett, 2000).*

 *Sex differences for group-level spatial maps*

Across all 12 movement tasks, males (mean across 12 tasks: 32,974 ± 13,181.47 voxels)

exhibited larger volumes of activation compared to females (mean across 12 tasks: 19,528.67 ±

10,000.85 voxels), which equates to males having a mean 87.18 ± 69.68% larger volumes of activation

across the whole brain (figure 2; table 8). The spatial maps between females and males were

moderately correlated (Akoglu, 2018) with a mean Pearsons correlation strength $r = 0.62 ± 0.06$, and a

range across the 12 tasks from $r = 0.48$ for ankle movements to $r = 0.72$ for left leg movements (table

2).

**Table 2.** Group-level whole brain spatial maps and correlations between sexes

| Condition | Total number of significant voxels* | | Spatial map similarity | |
|---|---|---|---|---|
| | Females | Males | Difference (%)** | *r*-values*** |
| Eyes | 33,429 | 47,035 | 40.70 | 0.65 |
| Jaw | 17,674 | 30,752 | 74.00 | 0.62 |
| Lips | 25,939 | 29,277 | 12.87 | 0.65 |
| Tongue | 31,624 | 43,223 | 36.68 | 0.62 |
| Upper Arms | 9,339 | 23,022 | 146.51 | 0.53 |
| Forearms | 19,865 | 39,633 | 99.51 | 0.58 |
| Wrists | 24,690 | 45,753 | 85.31 | 0.63 |
| Fingers | 26,591 | 43,500 | 63.59 | 0.66 |
| Left Leg | 6,050 | 8,637 | 42.76 | 0.72 |
| Right Leg | 4,838 | 10,067 | 108.08 | 0.64 |
| Ankles | 9,187 | 34,538 | 275.94 | 0.48 |
| Toes | 25,118 | 40,251 | 60.25 | 0.60 |

*Voxel size = 2 × 2 × 2 mm
**Percent difference of voxels activated in each condition for males compared to females
*** Pearson's *r* correlation coefficients from a cross-correlation analyses between females and males

235
236
237

*Figure 2.* P-value spatial maps for all 12 movement tasks. Female spatial maps are in blue on the left, male spatial maps are in red/orange on the right. Cerebral and cerebellar flatmaps showing unique and overlap are in the centre two columns respectively. Cerebral inflated brains and flatmaps were

*made in pycortex* (Gao *et al.*, 2015)*, and cerebellar flatmaps were made in SUIT* (Diedrichsen & Zotow, 2015).

239    *Sex differences in brain activation variability*

240    Using permutation testing to assess sex differences in correlations derived from whole-brain

241    voxelwise *t*-statistics for each contrast, we found that females had significantly higher between subject

242    consistency in 10 of the 12 motor tasks (table 3; $p < 0.004$, two-tailed, 20,000 permutations).

Table 3. Permutation testing of within group, between subject whole-brain voxelwise brain activation correlations

| Condition | Mean Correlation ± Standard Deviation | | |
|---|---|---|---|
| | Females | Males | Independent samples t-tests |
| Eyes | 0.205 ± 0.073 | 0.177 ± 0.058 | $t(904) = 6.210$, $p = 1.00E{-}04$, $d = 0.426$ * |
| Jaw | 0.200 ± 0.052 | 0.155 ± 0.051 | $t(904) = 12.779$, $p = 1.00E{-}04$, $d = 0.863$ * |
| Lips | 0.197 ± 0.056 | 0.164 ± 0.063 | $t(904) = 8.347$, $p = 1.00E{-}04$, $d = 0.557$ * |
| Tongue | 0.234 ± 0.067 | 0.218 ± 0.054 | $t(904) = 3.994$, $p = 3.00E{-}04$, $d = 0.274$ * |
| Upper Arms | 0.162 ± 0.066 | 0.153 ± 0.050 | $t(904) = 2.155$, $p = 0.029$, $d = 0.148$ |
| Forearms | 0.182 ± 0.055 | 0.158 ± 0.049 | $t(904) = 6.975$, $p = 1.00E{-}04$, $d = 0.474$ * |
| Wrists | 0.226 ± 0.054 | 0.198 ± 0.051 | $t(904) = 7.948$, $p = 1.00E{-}04$, $d = 0.538$ * |
| Fingers | 0.220 ± 0.060 | 0.203 ± 0.053 | $t(904) = 4.203$, $p = 1.00E{-}04$, $d = 0.286$ * |
| Left Leg | 0.171 ± 0.077 | 0.147 ± 0.071 | $t(904) = 4.772$, $p = 1.00E{-}04$, $d = 0.324$ * |
| Right Leg | 0.155 ± 0.080 | 0.156 ± 0.070 | $t(904) = -0.352$, $p = 0.724$, $d = -0.024$ |
| Ankles | 0.158 ± 0.064 | 0.142 ± 0.064 | $t(904) = 3.550$, $p = 9.00E{-}04$, $d = 0.239$ * |
| Toes | 0.171 ± 0.056 | 0.143 ± 0.050 | $t(904) = 7.806$, $p = 1.00E{-}04$, $d = 0.531$ * |

Note: * significant at $p < 0.05$ Bonferroni corrected [$\alpha = 0.004$; (0.05/12)], two-tailed, 20,000 permutations.

**Discussion**

In this study, we sought to determine if there were sex differences in somatotopic motor maps

with a range of movements across the human body. Based on previous research, there are distinct

structural differences in cortical organization between females and males, with females having greater

cortical thickness (Ritchie *et al.*, 2018; Wierenga *et al.*, 2022), and males having larger cortical surface

area; this is particularly the case for the motor and somatosensory regions (Ritchie *et al.*, 2018;

Wierenga *et al.*, 2022) which are known to have somatotopic organization (Penfield & Rasmussen,

1950). Given these previously observed differences, we hypothesized that females would exhibit more

focal (i.e., smaller volumes of activation) and consistent (i.e., lower variability between females for

each task) patterns of brain activation, compared to males across the 12 different body movements. To

answer these questions, we used a freely available fMRI dataset that included a total of 62 participants,

of which 34 were females and 28 were males.

*Males generally exhibit larger magnitudes of brain activation*

In seven of the movements under investigation (jaw, tongue, upper arms, wrists, fingers, ankles,

and toes) we observed greater BOLD signal in males than females. In contrast, females only exhibited

greater activation than males in one small cluster during right leg movements, and no differences were

seen with the remaining four movements (eyes, lips, forearms, and left leg).

*Females exhibit smaller volumes of brain activation across all movement conditions*

Smaller activation maps were observed in females across all 12 motor tasks (on average males

had $87.18 \pm 68.68\%$ larger activation maps than females, ranging from 12.87% larger for lip

movements to 275.94% larger for ankle movements; table 8, figure 2). These data support our

hypothesis that females would have more focal volumes of brain activation given the previous findings

that males have larger overall cortical and subcortical volume and surface area (Ritchie *et al.*, 2018;

Wierenga *et al.*, 2022), and provide support for females having better cortical efficiency (Luders *et al.*,

2006) during simple movements across the body.

*Lower whole-brain activation variability in females during all movement conditions*

Based on our novel whole-brain voxelwise correlation analysis, females consistently had higher

correlations between subjects compared to males (table 2), and these differences were significantly

different between sex. These findings should be viewed as complementary to the volume of activation

data reported in this work, given the smaller volumes of brain activation paired with the lower between

subject variability seen in females may be indicative of greater neural efficiency compared to males.

Our findings contribute to the growing body of evidence indicating that extensive sex

differences in brain organization and function exist (Cahill, 2017; Ritchie *et al.*, 2018; Murray *et al.*,

2018; Liu *et al.*, 2020). Importantly, Ritchie *et al.*, (2018) noted that the observed sex differences in

cortical measures in their work may be related compensatory interactions between sex-specific

hormones and the neural underpinnings that govern human voluntary movement (De Vries, 2004;

McCarthy & Arnold, 2011; Hyde, 2014). It is plausible that these hormonal differences may impact

both behaviour and the structure and function of the human brain, which could explain at least in part

the differences observed in the present study.

It is well known that the primary motor cortex is organized somatotopically, with adjacent body

parts represented by neighbouring regions of the cortex (Penfield & Rasmussen, 1950). Recently, the

accuracy of the motor homunculus has been brought into question, with a proposed update to the

cortical representation maps (Gordon *et al.*, 2023). However, little is known about the extent to which

this organization may vary between females and males. There is accumulating evidence from recent

years suggesting sex differences in brain structure, and specifically in motor areas, including the

premotor cortex, putamen, and cerebellum [GM volume higher in males > females (Ruigrok *et al.*,

2014; Ritchie *et al.*, 2018; Lotze *et al.*, 2019)]. Yet, the majority of these findings refer to structural

differences in volume, surface area or cortical thickness, with limited data regarding functional imaging

differences or their associations with motor behaviour. Greater spatial distributions and/or magnitudes

of BOLD signal have been observed, without differences in task performance, for males during verbal

and spatial tasks (Gur *et al.*, 2000), in addition to cognitive and motor tasks (Bell *et al.*, 2006). These

findings paired with the present work suggest that females typically show lower magnitudes, and

smaller volumes of brain activation across cognitive-motor tasks.

*Implications of sex differences in functional brain imaging research analyses*

Importantly, there are many methods to analyze functional brain imaging data. These analysis

and interpretation approaches often rely on previous findings, and as a result neglect to acknowledge

sex differences. For example, a ROI based analysis that is derived from an atlas, could prevent true

ROI identification if it does not consider sex differences in functional organization of the brain, and is

instead a product of group averaging. Some brain atlases that are commonly used in fMRI research for

ROI selection include the Harvard-Oxford cortical and subcortical atlases (16 females, 21 males,

between the ages of 18-50), the Brainnectome atlas (10 females, 10 males, all right handed, between

the ages of 19-25), and the Shaefer atlas (665 males, 905 females, between the ages of 18-35). Each of

these atlases were created by mixing female and male participants, which may lead to suboptimal

parcellations given our findings that males tend to exhibit larger volumes of BOLD signal across a

range of motor tasks. Our data suggests that future studies should consider selection of sex specific

ROIs based on functional activation (Nieto-Castanon *et al.*, 2003; Poldrack, 2007; Nieto-Castañón &

Fedorenko, 2012), when possible.

*Implications of sex differences in neurological impairments*

As discussed, a knowledge gap exists regarding the dimorphisms in somatotopic maps between

females in males. While filling this gap is generally useful for research purposes, it cannot be

understated how important these findings may be for the understanding of the etiology, prognosis, and

recovery from injury or disease in females versus males. The incidence and presentation of many

neurological conditions, and movement disorders differ between sexes, suggesting underlying

differences in brain function or structure (Meoni *et al.*, 2020). For example, females experience

increased severity, more impairment, and longer recovery from stroke (Kim *et al.*, 2010; Persky *et al.*,

2010; Phan *et al.*, 2019). Although many explanatory factors have been identified, incorporating

somatotopic differences into this phenomenon should be discussed. As identified in this research,

females use less of their cortex to generate voluntary movements, and do so with significantly lower

brain activation variability. Although this may be beneficial in healthy, neurologically intact

populations, this approach may be detrimental in the presence of a neurological injury. For example,

with stroke, injury to the central nervous system interrupts the normal motor output pathways and leads

to motor impairment. Acute stroke severity is often classified using clinical tests of neurological

function, such as the National Institutes of Health Stroke Scale (NIHSS; (Kwah & Diong, 2014). This

scale includes assessments of motor function in the arm and leg, with a higher score indicating more

impairment. Females present with higher NIHSS scores on average at stroke onset (Tomita *et al.*,

2015), indicating a greater degree of motor impairment than males, however one study has suggested

that males have, on average, larger infarcts than females (Appelros *et al.*, 2009). Our data showing that

females have smaller volumes of cortical activation than males, and lower variability, shows that

comparisons against standard ROI atlases may lead to misinterpretation of data and fundamentally limit

our understanding of brain function.

**Limitations and future directions**

335           As this study used a publicly available data, our analysis was restricted to the previously

collected data. Therefore, we were unable to include gender in our analysis. Recent studies indicate that

gender differences in both brain structure and brain function exist (Zhang *et al.*, 2020), and future

studies should collect the appropriate data to allow for both sex- and gender-based analyses.

Additionally, force output can influence brain activation (Noble *et al.*, 2011, 2013; Andrushko *et al.*,

2021), and in the present data there are no behavioural metrics to quantify the movement force,

velocity, or movement quality. This is a noted limitation that future work should consider measuring to

better link movement kinematics to cortical measures of activation.

**Conclusions**

344           In this study we found significant sex differences in the magnitude of brain activation for eight

out of the 12 movement conditions, with males having significantly greater brain activation in seven

out of the 12 conditions and females showing greater brain activation in one condition. Interestingly,

we observed smaller volumes of brain activation in all 12 movement conditions and significantly lower

whole-brain activation variability in 10 of the 12 movement conditions for females. With a cross-

correlation analysis to compare the spatial map similarity between sexes, we found the spatial maps

were moderately correlated for each movement conditions between sexes. The difference in volume is

an important consideration when creating ROIs based on functional localizers, as clear and visible

differences can be observed between sexes, and has potentially significant implications for

understanding the neurobiology of various neurological impairments in females and males (figure 2).

We propose these data provide important insight into the need for sex specific ROIs for functional

localization to accurately study brain-behaviour relationships using fMRI, and likely have important

clinical implications for understanding impairment severity and rehabilitation strategies for each sex

independently.

**Acknowledgements**

The authors would like to acknowledge the original authors of the open dataset Sai Ma,

Taicheng Huang, Yukun Qu, Xiayu Chen, Yajie Zhang and Zonglei Zhen. Without their support of open

science practices, this work would not have been made possible.

**Funding sources**

This work was funded by the Canadian Institutes of Health Research (PI L.A.B., PJT-148535)

to L.B. and an NSERC Discovery Grant and NSERC Discovery Launch Supplement to C.E. (RGPIN-

2021-03568 and DGECR-2021-00297). J.W.A., is funded by a Canadian Institutes of Health Research

(CIHR) fellowship and a Michael Smith Foundation for Health Research postdoctoral award, S.R. is

supported by awards from the University of British Columbia, E. K. is funded by an NSERC Canada

Graduate Master's award, J.D. is funded by a CIHR Canada Graduate Master's award.

**Author contributions**

J.W.A. processed and analyzed the data, and assisted in writing the manuscript. S.R. processed

and analyzed the data and assisted in writing the manuscript. E.K. created the figures and assisted in

writing the manuscript. J.D. assisted with writing the manuscript and approving the final version. C.E.

assisted in data analysis and writing the manuscript, and C.E. and L.B. were involved with the overall

conception and direction of this study, and edited and approved the manuscript.

**References**

Aboitiz F, Scheibel AB & Zaidel E (1992). Morphometry of the sylvian fissure and the corpus

callosum, with emphasis on sex differences. *Brain* **115,** 1521–1541.

Akoglu H (2018). User's guide to correlation coefficients. *Turkish J Emerg Med* **18,** 91–93.

Andersson JLR, Jenkinson M & Smith S (2007). *Non-linear registration aka spatial normalisation*

*FMRIB Technial Report TR07JA2*.

Andreano JM & Cahill L (2009). Sex influences on the neurobiology of learning and memory. *Learn*

*Mem* **16,** 248–266.

Andrushko JW, Gould LA, Renshaw DW, Ekstrand C, Hortobágyi T, Borowsky R & Farthing JP

(2021). High force unimanual handgrip contractions increase ipsilateral sensorimotor activation

and functional connectivity. *Neuroscience* **452,** 111–125.

Appelros P, Stegmayr B & Terent A (2009). Sex differences in stroke epidemiology. *Stroke* **40,** 1082–

1090.

Bell EC, Willson MC, Wilman AH, Dave S & Silverstone PH (2006). Males and females differ in brain

activation during cognitive tasks. *Neuroimage* **30,** 529–538.

Cahill L (2017). An issue whose time has come. *J Neurosci Res* **95,** 12–13.

Cohen NR, Pomplun M, Gold BJ & Sekuler R (2010). Sex differences in the acquisition of complex

skilled movements. *Exp Brain Res* **205,** 183–193.

Diamond MC (1991). Hormonal effects on the development of cerebral lateralization.

*Psychoneuroendocrinology* **16,** 121–129.

Diedrichsen J & Zotow E (2015). Surface-based display of volume-averaged cerebellar imaging data.

*PLoS One*; DOI: 10.1371/JOURNAL.PONE.0133402.

Ding Z, Newton AT, Xu R, Anderson AW, Morgan VL & Gore JC (2013). Spatio-temporal correlation

tensors reveal functional structure in human brain. *PLoS One* **8,** e82107.

Fabri M, Pierpaoli C, Barbaresi P & Polonara G (2014). Functional topography of the corpus callosum

investigated by DTI and fMRI. *World J Radiol* **6,** 895–906.

Frizzell TO, Grajauskas LA, Liu CC, Ghosh Hajra S, Song X & D'Arcy RCN (2020). White matter neuroplasticity: Motor learning activates the internal capsule and reduces hemodynamic response variability. *Front Hum Neurosci* **14,** 456.

Gao JS, Huth AG, Lescroart MD & Gallant JL (2015). Pycortex: An interactive surface visualizer for fMRI. *Front Neuroinform*; DOI: 10.3389/FNINF.2015.00023.

Gawryluk JR, D'Arcy RCN, Mazerolle EL, Brewer KD & Beyea SD (2011*a*). Functional mapping in the corpus callosum: A 4 T fMRI study of white matter. *Neuroimage* **54,** 10–15.

Gawryluk JR, Mazerolle EL, Brewer KD, Beyea SD & D'Arcy RCN (2011*b*). Investigation of fMRI activation in the internal capsule. *BMC Neurosci* **12,** 1–7.

Gawryluk JR, Mazerolle EL & D'Arcy RCN (2014). Does functional MRI detect activation in white matter? A review of emerging evidence, issues, and future directions. *Front Neurosci* **8,** 239.

Goldstein JM, Poldrack R, Breiter HC, Makris N, Goodman JM, Jerram M, Anagnoson R, Tsuang MT & Seidman LJ (2005). Sex differences in prefrontal cortical brain activity during fMRI of auditory verbal working memory. *Neuropsychology* **19,** 509–519.

Gordon EM et al. (2023). A somato-cognitive action network alternates with effector regions in motor cortex. *Nat 2023*1–9.

Gore JC, Li M, Gao Y, Wu TL, Schilling KG, Huang Y, Mishra A, Newton AT, Rogers BP, Chen LM, Anderson AW & Ding Z (2019). Functional MRI and resting state connectivity in white matter - a mini-review. *Magn Reson Imaging* **63,** 1–11.

Grajauskas LA, Frizzell T, Song X & D'Arcy RCN (2019). White matter fMRI activation cannot be treated as a nuisance regressor: Overcoming a historical blind spot. *Front Neurosci* **13,** 1024.

Gur RC, Alsop D, Glahn D, Petty R, Swanson CL, Maldjian JA, Turetsky BI, Detre JA, Gee J & Gur RE (2000). An fMRI study of sex differences in regional activation to a verbal and a spatial task. *Brain Lang* **74,** 157–170.

Harasty J, Double KL, Halliday GM, Kril JJ & McRitchie DA (1997). Language-associated cortical

regions are proportionally larger in the female brain. *Arch Neurol* **54,** 171–176.

Helenius J, Perkiö J, Soinne L, Østergaard L, Carano RAD, Salonen O, Savolainen S, Kaste M, Aronen

HJ & Tatlisumak T (2003). Cerebral hemodynamics in a healthy population measured by dynamic

susceptibility contrast MR imaging. *Acta radiol* **44,** 538–546.

Henschel L, Conjeti S, Estrada S, Diers K, Fischl B & Reuter M (2020). FastSurfer - A fast and

accurate deep learning based neuroimaging pipeline. *Neuroimage* **219,** 117012.

Hyde JS (2014). Gender similarities and differences. *Annu Rev Psychol* **65,** 373–398.

Jenkinson M, Bannister P, Brady M & Smith S (2002). Improved optimization for the robust and

accurate linear registration and motion correction of brain images. *Neuroimage* **17,** 825–841.

Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW & Smith SM (2012). FSL. *Neuroimage* **62,**

782–790.

Jenkinson M & Smith S (2001). A global optimisation method for robust affine registration of brain

images. *Med Image Anal* **5,** 143–156.

Jiang Y, Duan M, Li X, Huang H, Zhao G, Li X, Li S, Song X, He H, Yao D & Luo C (2021).

Function–structure coupling: White matter functional magnetic resonance imaging hyper-

activation associates with structural integrity reductions in schizophrenia. *Hum Brain Mapp* **42,**

4022–4034.

Jones CM, Braithwaite VA & Healy SD (2003). The evolution of sex differences in spatial ability.

*Behav Neurosci* **117,** 403–411.

Kim JS, Lee KB, Roh H, Ahn MY & Hwang HW (2010). Gender differences in the functional recovery

after acute stroke. *J Clin Neurol* **6,** 183.

Kulynych JJ, Vladar K, Jones DW & Weinberger DR (1994). Gender differences in the normal

lateralization of the supratemporal cortex: MRI surface-rendering morphometry of heschl's gyrus

and the planum temporale. *Cereb Cortex* **4,** 107–118.

Kuptsova S V., Ivanova M V., Petrushevsky AG, Fedina ON & Zhavoronkova LA (2015). Sex-related

differences in task switching: An fMRI study. *Hum Physiol* **41,** 611–624.

Kwah LK & Diong J (2014). National institutes of health stroke scale (NIHSS). *J Physiother* **60,** 61.

Liu S, Seidlitz J, Blumenthal JD, Clasen LS & Raznahan A (2020). Integrative structural, functional,

and transcriptomic analyses of sex-biased brain organization in humans. *Proc Natl Acad Sci U S A*

**117,** 18788–18798.

Lotze M, Domin M, Gerlach FH, Gaser C, Lueders E, Schmidt CO & Neumann N (2019). Novel

findings from 2,838 adult brains on sex differences in gray matter brain volume. *Sci Rep*; DOI:

10.1038/S41598-018-38239-2.

Luders E, Narr KL, Thompson PM, Rex DE, Woods RP, DeLuca H, Jancke L & Toga AW (2006).

Gender effects on cortical thickness and the influence of scaling. *Hum Brain Mapp* **27,** 314–324.

Ma S, Huang T, Qu Y, Chen X, Zhang Y & Zhen Z (2022). An fMRI dataset for whole-body

somatotopic mapping in humans. *Sci data*; DOI: 10.1038/S41597-022-01644-4.

Mazerolle EL, Beyea SD, Gawryluk JR, Brewer KD, Bowen C V. & D'Arcy RCN (2010). Confirming

white matter fMRI activation in the corpus callosum: Co-localization with DTI tractography.

*Neuroimage* **50,** 616–621.

Mazerolle EL, Gawryluk JR, Dillen KNH, Patterson SA, Feindel KW, Beyea SD, Stevens MTR,

Newman AJ, Schmidt MH & D'Arcy RCN (2013). Sensitivity to white matter fMRI activation

increases with field strength. *PLoS One* **8,** e58130.

McCarthy MM & Arnold AP (2011). Reframing sexual differentiation of the brain. *Nat Neurosci* **14,**

677–683.

Meoni S, Macerollo A & Moro E (2020). Sex differences in movement disorders. *Nat Rev Neurol 2020*

*162* **16,** 84–96.

Murray SO, Schallmo MP, Kolodny T, Millin R, Kale A, Thomas P, Rammsayer TH, Troche SJ,

Bernier RA & Tadin D (2018). Sex differences in visual motion processing. *Curr Biol* **28,** 2794-

2799.e3.

Newman AJ, Supalla T, Hauser P, Newport EL & Bavelier D (2010). Dissociating neural subsystems

for grammar by contrasting word order and inflection. *Proc Natl Acad Sci U S A* **107,** 7539–7544.

Nieto-Castañón A & Fedorenko E (2012). Subject-specific functional localizers increase sensitivity and

functional resolution of multi-subject analyses. *Neuroimage* **63,** 1646–1669.

Nieto-Castanon A, Ghosh SS, Tourville JA & Guenther FH (2003). Region of interest based analysis of

functional imaging data. *Neuroimage* **19,** 1303–1316.

Noble JW, Eng JJ & Boyd LA (2013). Effect of visual feedback on brain activation during motor tasks:

an FMRI study. *Motor Control* **17,** 298–312.

Noble JW, Eng JJ, Kokotilo KJ & Boyd LA (2011). Aging effects on the control of grip force

magnitude: An fMRI study. *Exp Gerontol* **46,** 453–461.

Di Noto PM, Newman L, Wall S & Einstein G (2013). The hermunculus: What is known about the

representation of the female body in the brain? *Cereb Cortex* **23,** 1005–1013.

Peer M, Nitzan M, Bick AS, Levin N & Arzy S (2017). Evidence for functional networks within the

human brain's white matter. *J Neurosci* **37,** 6394–6407.

Penfield W & Rasmussen T (1950). The cerebral cortex of man: A clinical study of localization of

function. *Macmillan*; DOI: 10.1001/jama.1950.02920160086033.

Persky RW, Turtzo LC & McCullough LD (2010). Stroke in women: Disparities and outcomes. *Curr*

*Cardiol Rep* **12,** 6.

Phan HT, Reeves MJ, Blizzard CL, Thrift AG, Cadilhac DA, Sturm J, Otahal P, Rothwell P, Bejot Y,

Cabral NL, Appelros P, Kõrv J, Vibo R, Minelli C & Gall SL (2019). Sex differences in severity of

stroke in the INSTRUCT study: A meta-analysis of individual participant data. *J Am Heart Assoc*;

DOI: 10.1161/JAHA.118.010235.

Poldrack RA (2007). Region of interest analysis for fMRI. *Soc Cogn Affect Neurosci* **2,** 67–70.

Ritchie SJ, Cox SR, Shen X, Lombardo M V., Reus LM, Alloza C, Harris MA, Alderson HL, Hunter S,

Neilson E, Liewald DCM, Auyeung B, Whalley HC, Lawrie SM, Gale CR, Bastin ME, McIntosh

AM & Deary IJ (2018). Sex differences in the adult human brain: Evidence from 5216 UK

biobank participants. *Cereb Cortex* **28,** 2959–2975.

Rorden C & Brett M (2000). Stereotaxic display of brain lesions. *Behav Neurol* **12,** 191–200.

Rostrup E, Law I, Blinkenberg M, Larsson HBW, Born AP, Holm S & Paulson OB (2000). Regional

differences in the CBF and BOLD responses to hypercapnia: A combined PET and fMRI study.

*Neuroimage* **11,** 87–97.

Ruigrok ANV, Salimi-Khorshidi G, Lai MC, Baron-Cohen S, Lombardo M V., Tait RJ & Suckling J

(2014). A meta-analysis of sex differences in human brain structure. *Neurosci Biobehav Rev* **39,**

34–50.

Swaab DF, Chung WCJ, Kruijver FPM, Hofman MA & Ishunina TA (2001). Structural and functional

sex differences in the human hypothalamus. *Horm Behav* **40,** 93–98.

Tettamanti M, Paulesu E, Scifo P, Maravita A, Fazio F, Perani D & Marzi CA (2002). Interhemispheric

transmission of visuomotor information in humans: fMRI evidence. *J Neurophysiol* **88,** 1051–

1058.

Thomsen T, Hugdahl K, Ersland L, Barndon R, Lundervold A, Smievoll AI, Roscher BE & Sundberg H

(2000). Functional magnetic resonance imaging (fMRI) study of sex differences in a mental

rotation task. *Med Sci Monit* **6,** 1186–1196.

Tomita H, Hagii J, Metoki N, Saito S, Shiroto H, Hitomi H, Kamada T, Seino S, Takahashi K, Baba Y,

Sasaki S, Uchizawa T, Iwata M, Matsumoto S, Shoji Y, Tanno T, Osanai T, Yasujima M &

Okumura K (2015). Impact of sex difference on severity and functional outcome in patients with

cardioembolic stroke. *J Stroke Cerebrovasc Dis* **24,** 2613–2618.

Virtanen P et al. (2020). SciPy 1.0: Fundamental algorithms for scientific computing in Python. *Nat*

*Methods* **17,** 261–272.

De Vries GJ (2004). Minireview: Sex differences in adult and developing brains: compensation,

compensation, compensation. *Endocrinology* **145,** 1063–1068.

Weis S, Leube D, Erb M, Heun R, Grodd W & Kircher T (2011). Functional neuroanatomy of sustained

memory encoding performance in healthy aging and in alzheimer's disease. *Int J Neurosci* **121,**

384–392.

Wierenga LM et al. (2022). Greater male than female variability in regional brain structure across the

lifespan. *Hum Brain Mapp* **43,** 470–499.

Woolrich MW, Behrens TEJ, Beckmann CF, Jenkinson M & Smith SM (2004). Multilevel linear

modelling for FMRI group analysis using Bayesian inference. *Neuroimage* **21,** 1732–1747.

Woolrich MW, Ripley BD, Brady M & Smith SM (2001). Temporal autocorrelation in univariate linear

modeling of FMRI data. *Neuroimage* **14,** 1370–1386.

Xu C, Li C, Wu H, Wu Y, Hu S, Zhu Y, Zhang W, Wang L, Zhu S, Liu J, Zhang Q, Yang J & Zhang X

(2015). Gender differences in cerebral regional homogeneity of adult healthy volunteers: A

resting-state fMRI study. *Biomed Res Int*; DOI: 10.1155/2015/183074.

Zhang X, Liang M, Qin W, Wan B, Yu C & Ming D (2020). Gender differences are encoded differently

in the structure and function of the human brain revealed by multimodal MRI. *Front Hum*

*Neurosci* **14,** 244.

Zhang Y, Brady M & Smith S (2001). Segmentation of brain MR images through a hidden Markov

random field model and the expectation-maximization algorithm. *IEEE Trans Med Imaging* **20,**

45–57.