Females exhibit smaller volumes of brain activation and lower inter-subject variability during

motor tasks

Justin W. Andrushko[1,2], Shie Rinat [1], Eric D. Kirby[3], Julia Dahlby [1], Chelsea Ekstrand[4,*], Lara A. Boyd[1,2,*]

1. Department of Physical Therapy, Faculty of Medicine, University of British Columbia, Vancouver, Canada
2. Djavad Mowafaghian Centre for Brain Health, University of British Columbia, Vancouver, Canada
3. Faculty of Individualized Interdisciplinary Studies, Simon Fraser University, Burnaby, BC, Canada
4. Department of Neuroscience, University of Lethbridge, Lethbridge, Alberta, Canada

* Co-senior authors

Correspondence

Lara A. Boyd, Ph.D.
Email: lara.boyd@ubc.ca
ORCID: https://orcid.org/0000-0002-2828-4549
Department of Physical Therapy
The University of British Columbia

and

Chelsea Ekstrand, Ph.D.
Email: Chelsea.ekstrand@uleth.ca
ORCID: https://orcid.org/0000-0002-1330-1924
Department of Neuroscience
The University of Lethbridge

Keywords:
Sex differences, functional magnetic resonance imaging, somatotopic motor mapping

Abstract

Past work has shown that brain structure and function differ between females and males. Males have

larger cortical and sub-cortical volume and surface area (both total and subregional), while females

have greater cortical thickness in most brain regions. Functional differences are also reported in the

literature, yet to date little work has systematically considered whether patterns of brain activity

indexed with functional magnetic resonance imaging (fMRI) differ between females and males.

Recently, an attempt to modernize and improve our understanding of the homunculus, a somatotopic

representation of the human body within the brain, was made. Still, in the decades since it was

introduced, no research has considered sex differences in the somatotopic representation. The current

study sought to remediate this issue by employing brain somatotopic motor mapping from a previously

published and openly available dataset. We tested differences in patterns of functional brain activity

associated with 12 voluntary movement patterns in females versus males. Results suggest that females

exhibited smaller volumes of brain activation across all 12 movement tasks, and lower patterns of

variability in 10 of the 12 movements. We also observed that females had greater cortical thickness,

which is in alignment with previous analyses of structural differences. Overall, these findings provide a

basis for considering biological sex in future fMRI research and provide a foundation of understanding

differences in how neurological pathologies present in females vs males.

Introduction

Females and males have anatomical and functional differences in their nervous systems [1–5]. These differences, attributed to differential sex chromosome expression and sex hormones as early as in utero [6–8], manifest in lateralization, motor planning, motor skill, language, memory, and spatial ability [9–14]. However, our understanding of how biological sex affects the somatotopic representations of the body within the brain is limited.

The homunculus, a somatotopic representation of the human body within the sensory and motor cortex, was first introduced in the late 1930s [15]. It maps out the cortical areas responsible for processing information from different body parts, where adjunct body parts are represented by adjacent brain regions. Although there has been a push to modernize our understanding of the classic sensorimotor "homunculus" [16–20], most studies have not considered sex differences. When sex has been included as a factor, the focus was largely on the sexual and reproductive dimorphisms and their respective neuronal representations [20–22].

A comparison of female and male brain structure reveals differences in volume, cortical surface area, cortical thickness, and structural variability [23]. Previous research suggests that males have significantly greater grey matter volume in areas such as the amygdala, temporal pole, fusiform gyrus, putamen, and premotor cortex [3,4]. Further, a large meta-analysis of 16,683 healthy participants found that males exhibit greater subcortical volume as compared to females, with the largest differences found in the thalamus and pallidum, bilaterally [5]. Males also exhibit greater cortical surface area. Importantly, males not only exhibit greater mean differences in subcortical volume and cortical surface area as compared to females, but also significantly greater variability in these measures [5,23]. This variability in surface area is most pronounced in motor related regions such as the pallidum, right inferior parietal cortex, and paracentral region [5]. This suggests that, compared to females, males have less consistent brain structure, particularly in motor related regions.

While most studies suggest greater total and subregional volume in males, some research found

regions with greater relative volume in females. When controlling for total brain volume, these studies

suggest females show greater grey matter volume than males in prefrontal and superior parietal corti-

ces, as well as the superior temporal sulcus, orbitofrontal cortex, and posterior insula [3,4]. Females

also exhibit greater cortical thickness compared to males [5,23]. Further, Wierenga et al. [5] noted re-

gional differences in cortical thickness between females and males; females had greater cortex thick-

ness in 38 of 68 regions, which were primarily in the frontal and parietal cortices. These structural dif-

ferences suggest that brain function may also differ by sex.

Sex related differences may affect the efficiency of neural activation patterns, with females po-

tentially having more localized and precise, and males having more distributed and varied, neuronal

processing. Several sex differences in brain activation have been observed with functional magnetic

resonance imaging (fMRI) across a range of tasks and resting-state brain activity [12,24,25]. However,

to date, little work has considered whether sex differences affect somatotopic sensorimotor mapping

across the entire brain. Researching sex specific patterns of brain activation with respect to area and

variability across a range of motor tasks may shed light on key differences in brain function. For in-

stance, in fMRI research, a common approach to investigate brain activation patterns is through region

of interest (ROI) analyses. However, using ROIs that are derived from an atlas that is not sex-specific

could potentially lead to data selection bias that does not fully capture differences in functional organi-

zation in females or males.

Using fMRI whole brain somatotopic motor mapping from a previously published and openly

available dataset [[26], https://openneuro.org/datasets/ds004044/versions/2.0.3], the current study in-

vestigated the differences in cortical activation patterns between females and males during a variety of

motor tasks. Based on previous observations of structural differences, we hypothesized that across

movement tasks, females would exhibit lower variability and smaller volumes of brain activation than

males, reflecting more focal neural processing. This concerted effort to further quantify the anatomical

localization and functional differences will aid our understanding of the inherent differences in somato-

topic maps between the sexes, and inform ROI selection strategies for future fMRI research by illustrat-

ing whether sex specific somatotopic brain parcellations are needed.

Methods

Participants

A total of 68 neurologically intact right-handed young adults (age range 19-29 years) were

recruited and screened for the Ma *et al.* [26] study. The original study from which these data were

collected was approved by the Institutional Review Board at Beijing Normal University [26], and

writen informed consent was provided by the participants.

Motor tasks

Twelve voluntary movement patterns were each performed for 16 seconds twice during every

fMRI scan. A total of six fMRI scans were acquired for each participant, resulting in 12 event blocks

for each of the 12 movement types (i.e.,192 seconds of task evoked data for each movement). These

movement conditions included bilateral toes, ankles, fingers, wrists, forearms, upper arms, and eyes, in

addition to jaw, lip, and tongue movements, and separate unilateral movements for the left and right

legs. Detailed movement instructions can be found in table 1. All movements were performed at a self-

selected pace between 0.5 and 5 Hz. Importantly, each participant underwent two behavioural training

sessions before undergoing the MRI portion of the study. These sessions involved watching a video

tutorial on how to perform each of movements, followed by lying supine and practicing each of the

movements under experimenter supervision. During these training sessions feedback was given to

ensure mastery of movement patterns, magnitude, and speed, while avoiding head or unrelated body

movements. To proceed to the MRI portion of the study, movement performance was evaluated and

verified by two experimenters. For further details about the nature of the experiment and motor tasks

please review the published dataset manuscript [26].

Table 1. Movement conditions and instructions

Body movements	Movement patterns and instructions
Eyes	Blink or saccade eyes.
Jaw	Bite or twist jaws.
Lips	Expand and contract the lips with the teeth being bitten and tongue still.
Tongue	Circular tongue with the teeth being bitten and lips closed.
Upper arms	Lift and lower the upper arms (maximum 20°) with the upper arms, forearms, wrist, and fingers straight.
Forearms	Flex and release both forearms at the elbow (maximum 20°) with the wrists and fingers straight.
Wrists	Pitch and roll both wrists with clenched fists.
Fingers	Clench and loose both fists.
Left leg	Lift and lower the left leg (maximum 10°) with the leg, ankle, and toes straight.
Right leg	Lift and lower the right leg (maximum 10°) with the leg, ankle, and toes straight.
Ankles	Dorsiflex and release both ankles.
Toes	Flex and extend the toes from both feet.
Rest	Fixate on the dot presented in the center of the screen.

Preprocessing structural MRI data

The T1-weighted (T1w) anatomical scans were preprocessed using FSL's [27] anatomical

preprocessing script *fsl_anat*. Briefly, this script performs image reorientation to align with standard

space template coordinates, crops the T1w image in the z-direction to remove the neck and other non-

brain aspects of the image, implements bias-field correction using FSL's FAST [28], and then extracts

the brain using a non-linear reverse transformation of a standard-space brain mask.

First level functional MRI processing

The previously preprocessed and manually denoised fMRI data (for more information regarding

the original preprocessing and denoising please review Ma et al. [26]) were linearly (ridged body +

affine) registered to the bias-field corrected and brain extracted T1w image and then non-linearly (rigid

+ affine + deformable syn) registered to the MNI152_2mm brain template using antsRegistrationSyN

and antsApplyTransforms from Advanced Normalization Tools (ANTs) [29,30]. Each of the 12 motor

tasks were modelled using the event timing files provided in the downloaded data with a double-

gamma function and with a temporal derivative applied to the model in FSL's FEAT [31]. White matter

BOLD signal was not treated as a nuisance regressor in the present work as this remains a controversial

technique [32,33,42–48,34–41]. Therefore, to prevent a bias towards grey matter BOLD signal and to

truly investigate whole-brain fMRI signal we retained white matter BOLD in our analyses [46].

Second level fixed effects functional MRI processing

Since each participant performed the same tasks over six separate runs, a fixed effects analysis was carried out in FEAT [31] prior to performing group-level analyses. The fixed effects analysis was used to average the parameter estimates of the task-based activations across the six runs to create a single activation map and effect size for each motor task for every participant. These averaged parameter estimates were then carried forward for group level analyses in FEAT [49].

Structural MRI analysis

T1w images for each participant were segmented using Fastsurfer [50]. Overall brain volume without ventricles was first analyzed to determine if there were differences between females and males in brain size. Next, cortical surface area, cortical thickness, and grey matter volume for each hemisphere were extracted for analysis and normalized as a ratio to overall brain volume on a participant-by-participant basis to account for differences in brain size. We then compared each of these normalized metrics for each hemisphere between females and males separately using two-tailed independent samples *t*-tests implemented using the *permutation_test* module from SciPy stats in Python 3.10 [51] with 20,000 permutations. A manual Bonferroni correction to adjust for multiple comparisons was performed and a result was deemed to be significant if the *p*-value was ≤ 0.007 ($\alpha = 0.05/7$).

Group level functional MRI analysis

For group-level analyses, two one-sample *t*-tests were performed to compute spatial maps for males and females separately. Additionally, two one-tailed independent samples *t*-tests were carried out to determine if there were differences in the somatotopic representations of each of the 12 motor movements between male and females (*t*-test #1: females > males; *t*-test #2: males > females contrasts). Non-parametric permutation testing was performed on each contrast using *randomise* with 5,000 permutations, and threshold free cluster enhancement was used for statistical inference with a family wise error rate set to $\alpha \leq 0.05$ for each contrast.

Sex difference analyses

To address the question of sex differences for somatotopic motor mapping, two approaches

were taken. First, the two *t*-tests previously described (*t*-test #1: females > males; *t*-test #2: males >

females contrasts) were assessed and reported when a significant contrast was detected (figure 1).

Second, to further investigate sex differences, the mean activation map for each movement was

computed separately for males and females and the spatial maps were correlated to assess their degree

of similarity. To do this, for males and females the group-level *p*-value threshold statistical maps ($\alpha \leq$

0.05) for each movement condition were temporally merged together using *fslmerge*. Then correlation

analyses between the spatial maps was carried out with *fslcc*. The correlation coefficients for same

movement conditions between females and males were then reported and the inter-task correlations

were disgarded.

Sex differences in brain activation variability

Variabililty in brain activation was assessed in two ways. First, within group between-subject

variability in brain activation was assessed for each motor task. To do this we performed pairwise

correlations for both females and males using the raw *t*-statistic image for each participant and

calculated the voxel-to-voxel correlation of these images for each pair of participants separately for

each group using in-house python code. This process was repeated for each motor task. We then

compared the pairwise correlations between females and males for each contrast using two-tailed

independent samples *t*-tests implemented using the *permutation_test* module from SciPy stats in

Python 3.10 [51] with 20,000 permutations. A manual Bonferroni correction to adjust for multiple

comparisons was performed and a result was deemed to be significant if the *p*-value was ≤ 0.004 ($\alpha =$

0.05/12).

Second, we analyzed brain activation variability within-subject between-runs (6 runs). To do

this, we once again perfomed parwise correlations for each participant using the raw t-statistic image

for each of the runs. We then calculated the mean correlation coefficient from these pairwise

correlations (mean of 15 comparisons). This process was repeated for each of the 12 motor tasks and

the rest condition. We then compared the pairwise correlations between females and males for each

contrast using two-tailed independent samples *t*-tests implemented using the *permutation_test* module

from SciPy stats in Python 3.10 [51] with 20,000 permutations. A manual Bonferroni correction to

adjust for multiple comparisons was performed and a result was deemed to be significant if the *p*-value

was ≤ 0.004 ($\alpha = 0.05/12$).

Results

Participants

214   Of the 68 participants from the Ma et al. [26] study, 61 participants were included in the present

studies analyses, of which 33 were female (22.58 ± 2.19 years), and 28 were male (23.00 ± 2.34 years).

Six participants were excluded from the study because they failed to master the movement patterns as

determined by two experimenters prior to the MRI session, and one particpant was exluded from the

analyses in this work because they were missing demographics information and their sex could not be

determined. Detailed demographic information is provided in original paper [26].

Sex differences in brain structure

221   Permutation *t*-testing with 20,000 permutations revealed sex differences in total brain volume

[Females: 1,099,719.242 ± 55040.247 mm^3; Males: 1,230,966.679 ± 110273.860 mm^3; $t(59) = -5.912$,

$p = 1.00E-04$, $d = -1.480$]. There was also a significant difference in normalized cortical thickness in

the left [Females: 2.18E-06 ± 1.17E-07; Males: 1.99E-06 ± 1.75E-07; $t(59) = 4.950$, $p = 1.00E-04$, $d =$

1.251] and right [Females: 2.19E-06 ± 1.17E-07; Males: 1.99E-06 ± 1.70E-07; $t(59) = 5.268$, $p =$

1.00E-04, d = 1.333] hemispheres with females having thicker cortex. However, no differences were

observed for normalized grey matter volume in the left [Females: 0.211 ± 0.008; Males: 0.214 ± 0.007;

$t(59) = 1.451$, $p = 0.155$, $d = -0.376$], or right [Females: 0.212 ± 0.008; Males: 0.215 ± 0.007; $t(59) = -$

1.740, $p = 0.089$, $d = -0.451$] hemisphere, or normalized surface area in the left [Females: 0.075 ±

0.002; Males: 0.075 ± 0.002; $t(59) = 0.653$, $p = 0.509$, $d = 0.165$], or right [Females: 0.075 ± 0.002;

Males: 0.075 ± 0.002; $t(59) = 0.653$, $p = 0.509$, $d = 0.165$] hemisphere. These findings suggest that

although males have larger overall brains, females exhibited thicker cortex in both hemispheres.

Sex differences for group-level contrasts

Significant differences were found in males > females contrasts in seven out of the 12

movements including tongue, jaw, upper arms, forearms, wrists, fingers, and ankles (figure 1;

supplementary data tables S1-S7). The only contrast that showed a significant females > males

difference was right leg movements (data table S8).

Figure 1. Sex difference contrasts for 8 significant conditions. Significantly greater BOLD signal for males (Red/Yellow) and females (White/Blue) are plotted. Images are family-wise error corrected p-value maps with a p value threshold of 0.05. Figure was made in MRIcoGL [52].

Sex differences for group-level spatial maps

Across all 12 movement tasks, males (mean across 12 tasks: 36,632.58 ± 14,469.06 voxels)

exhibited larger volumes of activation compared to females (mean across 12 tasks: 21,315.33 ±

11,738.90 voxels), which equates to males having a mean 92.68 ± 66.81% larger volumes of activation

across the whole brain (figure 2; table 8). The spatial maps between females and males were

moderately correlated [53] with a mean Pearsons correlation strength $r = 0.61 ± 0.06$, and a range

across the 12 tasks from $r = 0.48$ for upper arms movements to $r = 0.70$ for left leg movements (table

2).

Table 2. Group-level whole brain spatial maps and correlations between sexes

Condition	Total number of significant voxels*		Spatial map similarity	
	Females	Males	Difference (%)**	*r*-values***
Eyes	38,843	53,649	38.12	0.66
Jaw	14,336	34,774	142.56	0.56
Lips	27,593	36,828	33.47	0.64
Tongue	37,443	52,374	39.88	0.64
Upper Arms	9,434	30,045	218.48	0.48
Forearms	22,018	38,586	75.25	0.60
Wrists	26,317	47,832	81.75	0.64
Fingers	27,359	47,557	73.83	0.65
Left Leg	6,267	8,892	41.89	0.70
Right Leg	5,325	10,933	105.31	0.64
Ankles	11,112	35,301	217.68	0.51
Toes	29,737	42,820	44.00	0.64

*Voxel size = 2 × 2 × 2 mm
**Percent difference of voxels activated in each condition for males compared to females
*** Pearson's *r* correlation coefficients from a cross-correlation analyses between females and males

247
248
249

Figure 2. *P-value spatial maps for all 12 movement tasks. Female spatial maps are in blue on the left, male spatial maps are in red/orange on the right. Cerebral and cerebellar flatmaps showing unique*

and overlap are in the centre two columns respectively. Cerebral inflated brains and flatmaps were made in pycortex [54], and cerebellar flatmaps were made in SUIT [55].

Sex differences in brain activation variability

Using permutation testing to assess sex differences in correlations derived from whole-brain

voxelwise *t*-statistics for each contrast, we found that females had significantly higher between subject

consistency in 10 of the 12 motor tasks (table 3; $p < 0.004$, two-tailed, 20,000 permutations). However

an analysis of within subject between run variability resulted in no differences between sex for any of

the motor tasks (table 4; two-tailed, 20,000 permutations), suggesting that there were no differences

between females and males in how consistent their patterns of brain activation were across the six runs

for each of the motor tasks.

Table 3. Permutation testing of within group, between subject whole-brain voxelwise brain activation correlations

	Mean Correlation ± Standard Deviation		
Condition	Females	Males	Independent samples t-tests
Eyes	0.259 ± 0.085	0.226 ± 0.064	$t(904) = 6.247, p = 1.00\text{E-}04, d = 0.430$
Jaw	0.228 ± 0.056	0.187 ± 0.055	$t(904) = 11.021, p = 1.00\text{E-}04, d = 0.744$
Lips	0.231 ± 0.066	0.201 ± 0.063	$t(904) = 6.883, p = 1.00\text{E-}04, d = 0.465$
Tongue	0.278 ± 0.069	0.264 ± 0.056	$t(904) = 3.274, p = 0.001, d = 0.224$
Upper Arms	0.216 ± 0.079	0.200 ± 0.061	$t(904) = 3.483, p = 5.00\text{E-}04, d = 0.240$
Forearms	0.239 ± 0.072	0.209 ± 0.047	$t(904) = 7.253, p = 1.00\text{E-}04, d = 0.505$
Wrists	0.302 ± 0.063	0.266 ± 0.062	$t(904) = 8.530, p = 1.00\text{E-}04, d = 0.576$
Fingers	0.303 ± 0.068	0.280 ± 0.066	$t(904) = 5.179, p = 1.00\text{E-}04, d = 0.350$
Left Leg	0.217 ± 0.085	0.183 ± 0.083	$t(904) = 5.886, p = 1.00\text{E-}04, d = 0.398$
Right Leg	0.199 ± 0.096	0.209 ± 0.083	$t(904) = -1.598, p = 0.108, d = -0.109$
Ankles	0.216 ± 0.072	0.188 ± 0.066	$t(904) = 5.906, p = 1.00\text{E-}04, d = 0.401$
Toes	0.244 ± 0.070	0.184 ± 0.055	$t(904) = 9.023, p = 1.00\text{E-}04, d = 0.620$

Note: * significant at $p < 0.05$ Bonferroni corrected [$\alpha = 0.004$; (0.05/12)], two-tailed, 20,000 permutations.

259
260

Table 4. Permutation testing of within subject, between runs whole-brain voxelwise brain activation correlations

	Mean Correlation ± Standard Deviation		
Condition	Females	Males	Independent samples t-tests
Eyes	0.190 ± 0.101	0.181 ± 0.100	$t(59) = 0.369, p = 0.700, d = 0.095$
Jaw	0.174 ± 0.063	0.172 ± 0.092	$t(59) = 0.096, p = 0.929, d = 0.024$
Lips	0.189 ± 0.086	0.180 ± 0.082	$t(59) = 0.452, p = 0.638, d = 0.116$
Tongue	0.269 ± 0.085	0.226 ± 0.103	$t(59) = 1.774, p = 0.081, d = 0.452$
Upper Arms	0.241 ± 0.079	0.225 ± 0.089	$t(59) = 0.700, p = 0.489, d = 0.179$
Forearms	0.195 ± 0.080	0.186 ± 0.073	$t(59) = 0.452, p = 0.654, d = 0.116$
Wrists	0.223 ± 0.092	0.219 ± 0.087	$t(59) = 0.161, p = 0.870, d = 0.041$

Fingers	0.206 ± 0.092	0.199 ± 0.071	$t(59) = 0.335, p = 0.731, d = 0.087$
Left Leg	0.195 ± 0.067	0.219 ± 0.099	$t(59) = -1.129, p = 0.264, d = -0.285$
Right Leg	0.181 ± 0.054	0.242 ± 0.120	$t(59) = -2.588, p = 0.012, d = -0.646$
Ankles	0.148 ± 0.073	0.161 ± 0.071	$t(59) = -0.705, p = 0.479, d = -0.181$
Toes	0.154 ± 0.073	0.141 ± 0.079	$t(59) = 0.685, p = 0.494, d = 0.175$
Rest	0.059 ± 0.033	0.048 ± 0.022	$t(59) = 1.511, p = 0.138, d = 0.394$

Note: Significance set to Bonferroni corrected $\alpha = 0.004$ [$\alpha = 0.004$; (0.05/12)], two-tailed, 20,000 permutations.

Discussion

In this study, we sought to determine if there were sex differences in somatotopic motor maps

with a range of movements across the human body. Based on previous research, there are distinct

structural differences in cortical organization between females and males, with females having greater

cortical thickness [5,23], and males having larger cortical surface area and cortical volume; this is

particularly the case for the motor and somatosensory regions [5,23] which are known to have

somatotopic organization [56]. Given these previously observed differences, we hypothesized that

females would exhibit more focal (i.e., smaller volumes of activation) and consistent (i.e., lower

variability between females for each task) patterns of brain activation, compared to males across the 12

different body movements. To answer these questions, we used a freely available fMRI dataset that

included a total of 61 participants, of which 33 were females and 28 were males.

Males generally exhibit larger magnitudes of brain activation

In seven of the movements under investigation (jaw, tongue, upper arms, forearms, wrists,

fingers, and ankles) we observed greater BOLD signal in males than females. In contrast, females only

exhibited greater activation than males during right leg movements, and no differences were seen with

the remaining four movements (eyes, lips, left leg, and toes).

Females exhibit smaller volumes of brain activation across all movement conditions

Smaller activation maps were observed in females across all 12 motor tasks (on average males

had 92.68 ± 66.81% larger activation maps than females, ranging from 33.47% larger for lip

movements to 218.48% larger for upper arms movements; table 2, figure 2). These data support our

hypothesis that females would have more focal volumes of brain activation given the previous findings

that males have larger overall cortical and subcortical volume and surface area [5,23], and provide

support for females having better cortical efficiency [57] during simple movements across the body.

Lower whole-brain activation variability in females during all movement conditions

Based on our novel whole-brain voxelwise correlation analysis, females consistently showed

higher correlations between subjects compared to males (table 2); these differences were statistically

significant. These findings should be viewed as complementary to the volume of activation data

reported in this work, given the smaller volumes of brain activation paired with the lower between

subject variability seen in females may be indicative of greater neural efficiency compared to males.

Additionally, to determine if there were sex differences for how consistent patterns of brain activation

were during the 12 motor tasks, we assessed intra-subject variability for each motor task across the six

separate fMRI runs. Based on these intra-subject variability analyses, there were no sex differences and

both females and males exhibited similar levels of variability. These analysis were important to ensure

any sex differences observed throughout the study were not a product of intra-subject variability

differences between sexes.

Our findings contribute to the growing body of evidence indicating that extensive sex

differences in brain organization and function exist [3,23,58,59]. Importantly, Ritchie *et al.*, [23] noted

that the observed sex differences in cortical measures in their work may be related compensatory

interactions between sex-specific hormones and the neural underpinnings that govern human voluntary

movement [60–62]. It is plausible that these hormonal differences may impact both behaviour and the

structure and function of the human brain, which could explain at least in part the differences observed

in the present study.

It is well known that the primary motor cortex is organized somatotopically, with adjacent body

parts represented by neighbouring regions of the cortex [56]. Recently, the accuracy of the motor

homunculus has been brought into question, with a proposed update to the cortical representation maps

[16]. However, little is known about the extent to which this organization may vary between females

and males. There is accumulating evidence from recent years suggesting sex differences in brain

structure, and specifically in motor areas, including the premotor cortex, putamen, and cerebellum [GM

volume higher in males > females [4,23,63]]. Yet, the majority of these findings refer to structural

differences in volume, surface area or cortical thickness, with limited data regarding functional imaging

differences or their associations with motor behaviour. Greater spatial distributions and/or magnitudes

of BOLD signal have been observed, without differences in task performance, for males during verbal

and spatial tasks [64], in addition to cognitive and motor tasks [65]. These findings paired with the

present work suggest that females often show lower magnitudes, and smaller volumes of brain

activation across cognitive-motor tasks.

Implications of sex differences in functional brain imaging research analyses

Importantly, there are many methods to analyze functional brain imaging data. These analysis

and interpretation approaches often rely on previous findings, and as a result neglect to acknowledge

sex differences. For example, a ROI based analysis that is derived from an atlas, could prevent true

ROI identification if it does not consider sex differences in functional organization of the brain, and is

instead a product of group averaging. Some brain atlases that are commonly used in fMRI research for

ROI selection include the Harvard-Oxford cortical and subcortical atlases (16 females, 21 males,

between the ages of 18-50), the Brainnectome atlas (10 females, 10 males, all right handed, between

the ages of 19-25), and the Shaefer atlas (665 males, 905 females, between the ages of 18-35). Each of

these atlases were created by mixing female and male participants, which may lead to suboptimal

parcellations given our findings that males tend to exhibit larger volumes of BOLD signal across a

range of motor tasks. Our data suggests that future studies should consider selection of sex specific

ROIs based on functional activation [66–68], when possible.

Implications of sex differences in neurological impairments

As discussed, a knowledge gap exists regarding the dimorphisms in somatotopic maps between

females in males. While filling this gap is generally useful for research purposes, it cannot be

understated how important these findings may be for the understanding of the etiology, prognosis, and

recovery from injury or disease in females versus males. The incidence and presentation of many

neurological conditions and movement disorders differ between sexes, suggesting underlying

differences in brain function or structure [69]. For example, females experience increased severity,

more impairment, and longer recovery from stroke [70–73]. Although many explanatory factors have

been identified, incorporating somatotopic differences into this phenomenon should be discussed. As

identified in this research, females use less of their cortex to generate voluntary movements, and do so

with significantly lower brain activation variability. Although this may be beneficial in healthy,

neurologically intact populations, this approach may be detrimental in the presence of a neurological

injury. For example, with stroke, injury to the central nervous system interrupts the normal motor

output pathways and leads to motor impairment. Our data showing that females have smaller volumes

of cortical activation than males, and lower variability, shows that comparisons against standard ROI

atlases may lead to misinterpretation of data and fundamentally limit our understanding of brain

function.

Limitations and future directions

As this study used a publicly available data, our analysis was restricted to the previously

collected data. Therefore, we were unable to include gender in our analysis. Recent studies indicate that

gender differences in both brain structure and brain function exist [74], and future studies should

collect the appropriate data to allow for both sex- and gender-based analyses. Additionally, force output

can influence brain activation [75–77], and in the present data there are no behavioural metrics to

quantify the movement force, velocity, or quality. This is a noted limitation that future work should

consider measuring to better link movement kinematics to cortical measures of activation. However,

past work [75] suggests that small differences in force (e.g., between 25% and 50%, or 50% and 75%

MVC) are not associated with significant differences in brain activation. Only large differences in

movement characteristics (e.g., 25% and 75% MVC) result in activation changes. Such differences are

unlikely to occur in this study, as participants were instructed to perform the movements at a

comfortable self-selected pace, without any additional resistance. Further, even if individual differences

exist, group differences in movement characteristics are not assumed, especially in light of the lack of

sex differences in intra-subject variability. In light of our sample size, we presume that such

differences, if they exist, will be balanced between the two groups.

Conclusions

In this study we found significant sex differences in the magnitude of brain activation for eight

out of the 12 movement conditions, with males having significantly greater brain activation in seven

out of the 12 conditions and females showing greater brain activation in one condition. Interestingly,

we observed smaller volumes of brain activation in all 12 movement conditions and significantly lower

whole-brain activation variability in 10 of the 12 movement conditions for females. With a cross-

correlation analysis to compare the spatial map similarity between sexes, we found that spatial maps

were moderately correlated for each movement conditions between sexes. The difference in volume is

an important consideration when creating ROIs based on functional localizers, as clear and visible

differences can be observed between sexes, and has potentially significant implications for

understanding the neurobiology of various neurological impairments in females and males (figure 2).

We propose these data provide important insight into the need for sex specific ROIs for functional

localization to accurately study brain-behaviour relationships using fMRI, and likely have important

clinical implications for understanding impairment severity and rehabilitation strategies for each sex

independently.

Acknowledgements

The authors would like to acknowledge the original authors of the open dataset Sai Ma,

Taicheng Huang, Yukun Qu, Xiayu Chen, Yajie Zhang and Zonglei Zhen. Without their support of open

science practices, this work would not have been made possible.

Funding sources

This work was funded by the Canadian Institutes of Health Research (PI L.A.B., PJT-148535)

to L.B. and an NSERC Discovery Grant and NSERC Discovery Launch Supplement to C.E. (RGPIN-

2021-03568 and DGECR-2021-00297). J.W.A., is funded by a Canadian Institutes of Health Research

(CIHR) fellowship and a Michael Smith Foundation for Health Research postdoctoral award, S.R. is

supported by awards from the University of British Columbia, E. K. is funded by an NSERC Canada

Graduate Master's award, J.D. is funded by a CIHR Canada Graduate Master's award.

Author contributions

J.W.A. processed and analyzed the data, and assisted in writing the manuscript. S.R. processed

and analyzed the data and assisted in writing the manuscript. E.K. created the figures and assisted in

writing the manuscript. J.D. assisted with writing the manuscript and approving the final version. C.E.

assisted in data analysis and writing the manuscript, and C.E. and L.B. were involved with the overall

conception and direction of this study, and edited and approved the manuscript.

References

1. Williams CM, Peyre H, Toro R, Ramus F. Sex differences in the brain are not reduced to differences in body size. Neurosci Biobehav Rev. 2021;130: 509–511. doi:10.1016/J.NEUBIOREV.2021.09.015

2. DeCasien AR, Guma E, Liu S, Raznahan A. Sex differences in the human brain: A roadmap for more careful analysis and interpretation of a biological reality. Biol Sex Differ. 2022;13. doi:10.1186/S13293-022-00448-W

3. Liu S, Seidlitz J, Blumenthal JD, Clasen LS, Raznahan A. Integrative structural, functional, and transcriptomic analyses of sex-biased brain organization in humans. Proc Natl Acad Sci U S A. 2020;117: 18788–18798. doi:10.1073/PNAS.1919091117/SUPPL_FILE/PNAS.1919091117.SM01.MP4

4. Lotze M, Domin M, Gerlach FH, Gaser C, Lueders E, Schmidt CO, et al. Novel findings from 2,838 adult brains on sex differences in gray matter brain volume. Sci Rep. 2019;9. doi:10.1038/S41598-018-38239-2

5. Wierenga LM, Doucet GE, Dima D, Agartz I, Aghajani M, Akudjedu TN, et al. Greater male than female variability in regional brain structure across the lifespan. Hum Brain Mapp. 2022;43: 470–499. doi:10.1002/HBM.25204

6. Cook KM, De Asis-Cruz J, Lopez C, Quistorff J, Kapse K, Andersen N, et al. Robust sex differences in functional brain connectivity are present in utero. Cereb Cortex. 2023;33: 2441–2454. doi:10.1093/CERCOR/BHAC218

7. Mallard TT, Liu S, Seidlitz J, Ma Z, Moraczewski D, Thomas A, et al. X-chromosome influences

on neuroanatomical variation in humans. Nat Neurosci. 2021;24: 1216–1224.

doi:10.1038/S41593-021-00890-W

8.   McCarthy MM, Pickett LA, VanRyzin JW, Kight KE. Surprising origins of sex differences in the

brain. Horm Behav. 2015;76: 3–10. doi:10.1016/J.YHBEH.2015.04.013

9.   Hegarty M. Ability and sex differences in spatial thinking: What does the mental rotation test

really measure? Psychon Bull Rev. 2018;25: 1212–1219. doi:10.3758/S13423-017-1347-

Z/TABLES/6

10. Yuan L, Kong F, Luo Y, Zeng S, Lan J, You X. Gender differences in large-scale and small-scale

spatial ability: A systematic review based on behavioral and neuroimaging research. Front Behav

Neurosci. 2019;13. doi:10.3389/FNBEH.2019.00128

11. Rogojin A, Gorbet DJ, Sergio LE. Sex differences in the neural underpinnings of unimanual and

bimanual control in adults. Exp Brain Res. 2023;241: 793–806. doi:10.1007/S00221-023-06561-

5

12. Spets DS, Slotnick SD. Are there sex differences in brain activity during long-term memory? A

systematic review and fMRI activation likelihood estimation meta-analysis. Cogn Neurosci.

2021;12: 163–173. doi:10.1080/17588928.2020.1806810

13. Xu M, Liang X, Ou J, Li H, Luo YJ, Tan LH. Sex differences in functional brain networks for

language. Cereb Cortex. 2020;30: 1528–1537. doi:10.1093/CERCOR/BHZ184

14. Gur RC, Gur RE. Complementarity of sex differences in brain and behavior: From laterality to

multimodal neuroimaging. J Neurosci Res. 2017;95: 189–199. doi:10.1002/JNR.23830

15. Penfield W, Boldrey E. Somatic motor and sensory representation in the cerebral cortex of man

as studied by electrical stimulation. Brain. 1937;60: 389–443. doi:10.1093/BRAIN/60.4.389

16. Gordon EM, Chauvin RJ, Van AN, Rajesh A, Nielsen A, Newbold DJ, et al. A somato-cognitive

action network alternates with effector regions in motor cortex. Nature. 2023; 1–9.

doi:10.1038/s41586-023-05964-2

17.   Roux FE, Niare M, Charni S, Giussani C, Durand JB. Functional architecture of the motor

homunculus detected by electrostimulation. J Physiol. 2020;598: 5487–5504.

doi:10.1113/JP280156

18.   Zeharia N, Hertz U, Flash T, Amedi A. Negative blood oxygenation level dependent homunculus

and somatotopic information in primary motor cortex and supplementary motor area. Proc Natl

Acad Sci U S A. 2012;109: 18565–18570. doi:10.1073/PNAS.1119125109

19.   Kell CA, Von Kriegstein K, Rösler A, Kleinschmidt A, Laufs H. The sensory cortical

representation of the human penis: revisiting somatotopy in the male homunculus. J Neurosci.

2005;25: 5984–5987. doi:10.1523/JNEUROSCI.0712-05.2005

20.   Lima SQ. Genital cortex: Development of the genital homunculus. Curr Biol. 2019;29: R1122–

R1124. doi:10.1016/J.CUB.2019.09.051

21.   Di Noto PM, Newman L, Wall S, Einstein G. The hermunculus: What is known about the

representation of the female body in the brain? Cereb Cortex. 2013;23: 1005–1013.

doi:10.1093/CERCOR/BHS005

22.   Komisaruk BR, Wise N, Frangos E, Liu WC, Allen K, Brody S. Women's clitoris, vagina, and

cervix mapped on the sensory cortex: fMRI evidence. J Sex Med. 2011;8: 2822–2830.

doi:10.1111/J.1743-6109.2011.02388.X

23.   Ritchie SJ, Cox SR, Shen X, Lombardo M V., Reus LM, Alloza C, et al. Sex differences in the

adult human brain: Evidence from 5216 UK biobank participants. Cereb Cortex. 2018;28: 2959–

2975. doi:10.1093/CERCOR/BHY109

24.   Xu C, Li C, Wu H, Wu Y, Hu S, Zhu Y, et al. Gender differences in cerebral regional

homogeneity of adult healthy volunteers: A resting-state fMRI study. Biomed Res Int.

2015;2015. doi:10.1155/2015/183074

25.   Kuptsova S V., Ivanova M V., Petrushevsky AG, Fedina ON, Zhavoronkova LA. Sex-related

differences in task switching: An fMRI study. Hum Physiol. 2015;41: 611–624.

478   doi:10.1134/S0362119715050084/METRICS

26.   Ma S, Huang T, Qu Y, Chen X, Zhang Y, Zhen Z. An fMRI dataset for whole-body somatotopic

mapping in humans. Sci Data. 2022;9: 515. doi:10.1038/s41597-022-01644-4

27.   Jenkinson M, Beckmann CF, Behrens TEJ, Woolrich MW, Smith SM. FSL. Neuroimage.

2012;62: 782–790. doi:10.1016/j.neuroimage.2011.09.015

28.   Zhang Y, Brady M, Smith S. Segmentation of brain MR images through a hidden Markov

random field model and the expectation-maximization algorithm. IEEE Trans Med Imaging.

2001;20: 45–57. doi:10.1109/42.906424

29.   Avants B, Tustison N, Johnson H. Advanced Normalization Tools (ANTS). Insight J. 2009.

30.   Avants BB, Epstein CL, Grossman M, Gee JC. Symmetric diffeomorphic image registration with

cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. Med

Image Anal. 2008;12: 26–41. doi:10.1016/J.MEDIA.2007.06.004

31.   Woolrich MW, Ripley BD, Brady M, Smith SM. Temporal autocorrelation in univariate linear

modeling of FMRI data. Neuroimage. 2001;14: 1370–1386. doi:10.1006/nimg.2001.0931

32.   Gore JC, Li M, Gao Y, Wu TL, Schilling KG, Huang Y, et al. Functional MRI and resting state

connectivity in white matter - a mini-review. Magn Reson Imaging. 2019;63: 1–11.

doi:10.1016/J.MRI.2019.07.017

33.   Gawryluk JR, Mazerolle EL, D'Arcy RCN. Does functional MRI detect activation in white

matter? A review of emerging evidence, issues, and future directions. Front Neurosci. 2014;8:

239. doi:10.3389/FNINS.2014.00239/ABSTRACT

34.   Rostrup E, Law I, Blinkenberg M, Larsson HBW, Born AP, Holm S, et al. Regional differences

in the CBF and BOLD responses to hypercapnia: A combined PET and fMRI study.

Neuroimage. 2000;11: 87–97. doi:10.1006/NIMG.1999.0526

35.   Helenius J, Perkiö J, Soinne L, Østergaard L, Carano RAD, Salonen O, et al. Cerebral

hemodynamics in a healthy population measured by dynamic susceptibility contrast MR

503  imaging. Acta radiol. 2003;44: 538–546. doi:10.1080/J.1600-0455.2003.00104.X

504 36. Tettamanti M, Paulesu E, Scifo P, Maravita A, Fazio F, Perani D, et al. Interhemispheric

505  transmission of visuomotor information in humans: fMRI evidence. J Neurophysiol. 2002;88:

506  1051–1058. doi:10.1152/JN.2002.88.2.1051/ASSET/IMAGES/LARGE/9K0822479104.JPEG

507 37. Newman AJ, Supalla T, Hauser P, Newport EL, Bavelier D. Dissociating neural subsystems for

508  grammar by contrasting word order and inflection. Proc Natl Acad Sci U S A. 2010;107: 7539–

509  7544. doi:10.1073/PNAS.1003174107/SUPPL_FILE/SM01.MP4

510 38. Gawryluk JR, D'Arcy RCN, Mazerolle EL, Brewer KD, Beyea SD. Functional mapping in the

511  corpus callosum: A 4 T fMRI study of white matter. Neuroimage. 2011;54: 10–15.

512  doi:10.1016/J.NEUROIMAGE.2010.07.028

513 39. Gawryluk JR, Mazerolle EL, Brewer KD, Beyea SD, D'Arcy RCN. Investigation of fMRI

514  activation in the internal capsule. BMC Neurosci. 2011;12: 1–7. doi:10.1186/1471-2202-12-

515  56/FIGURES/3

516 40. Weis S, Leube D, Erb M, Heun R, Grodd W, Kircher T. Functional neuroanatomy of sustained

517  memory encoding performance in healthy aging and in alzheimer's disease. Int J Neurosci.

518  2011;121: 384–392. doi:10.3109/00207454.2011.565892

519 41. Mazerolle EL, Gawryluk JR, Dillen KNH, Patterson SA, Feindel KW, Beyea SD, et al.

520  Sensitivity to white matter fMRI activation increases with field strength. PLoS One. 2013;8:

521  e58130. doi:10.1371/JOURNAL.PONE.0058130

522 42. Fabri M, Pierpaoli C, Barbaresi P, Polonara G. Functional topography of the corpus callosum

523  investigated by DTI and fMRI. World J Radiol. 2014;6: 895–906. doi:10.4329/WJR.V6.I12.895

524 43. Peer M, Nitzan M, Bick AS, Levin N, Arzy S. Evidence for functional networks within the

525  human brain's white matter. J Neurosci. 2017;37: 6394–6407. doi:10.1523/JNEUROSCI.3872-

526  16.2017

527 44. Frizzell TO, Grajauskas LA, Liu CC, Ghosh Hajra S, Song X, D'Arcy RCN. White matter

neuroplasticity: Motor learning activates the internal capsule and reduces hemodynamic

response variability. Front Hum Neurosci. 2020;14: 456.

doi:10.3389/FNHUM.2020.509258/BIBTEX

45.   Jiang Y, Duan M, Li X, Huang H, Zhao G, Li X, et al. Function–structure coupling: White matter

functional magnetic resonance imaging hyper-activation associates with structural integrity

reductions in schizophrenia. Hum Brain Mapp. 2021;42: 4022–4034. doi:10.1002/HBM.25536

46.   Grajauskas LA, Frizzell T, Song X, D'Arcy RCN. White matter fMRI activation cannot be

treated as a nuisance regressor: Overcoming a historical blind spot. Front Neurosci. 2019;13:

1024. doi:10.3389/FNINS.2019.01024/BIBTEX

47.   Ding Z, Newton AT, Xu R, Anderson AW, Morgan VL, Gore JC. Spatio-temporal correlation

tensors reveal functional structure in human brain. PLoS One. 2013;8: e82107.

doi:10.1371/JOURNAL.PONE.0082107

48.   Mazerolle EL, Beyea SD, Gawryluk JR, Brewer KD, Bowen C V., D'Arcy RCN. Confirming

white matter fMRI activation in the corpus callosum: Co-localization with DTI tractography.

Neuroimage. 2010;50: 616–621. doi:10.1016/J.NEUROIMAGE.2009.12.102

49.   Woolrich MW, Behrens TEJ, Beckmann CF, Jenkinson M, Smith SM. Multilevel linear

modelling for FMRI group analysis using Bayesian inference. Neuroimage. 2004;21: 1732–

1747. doi:10.1016/j.neuroimage.2003.12.023

50.   Henschel L, Conjeti S, Estrada S, Diers K, Fischl B, Reuter M. FastSurfer - A fast and accurate

deep learning based neuroimaging pipeline. Neuroimage. 2020;219: 117012.

doi:10.1016/J.NEUROIMAGE.2020.117012

51.   Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0:

Fundamental algorithms for scientific computing in Python. Nat Methods. 2020;17: 261–272.

doi:10.1038/s41592-019-0686-2

52.   Rorden C, Brett M. Stereotaxic display of brain lesions. Behav Neurol. 2000;12: 191–200.

doi:10.1155/2000/421719

53.   Akoglu H. User's guide to correlation coefficients. Turkish J Emerg Med. 2018;18: 91–93.

doi:10.1016/J.TJEM.2018.08.001

54.   Gao JS, Huth AG, Lescroart MD, Gallant JL. Pycortex: An interactive surface visualizer for

fMRI. Front Neuroinform. 2015;9. doi:10.3389/FNINF.2015.00023

55.   Diedrichsen J, Zotow E. Surface-based display of volume-averaged cerebellar imaging data.

PLoS One. 2015;10. doi:10.1371/JOURNAL.PONE.0133402

56.   Penfield W, Rasmussen T. The cerebral cortex of man: A clinical study of localization of

function. Macmillan. 1950. doi:10.1001/jama.1950.02920160086033

57.   Luders E, Narr KL, Thompson PM, Rex DE, Woods RP, DeLuca H, et al. Gender effects on

cortical thickness and the influence of scaling. Hum Brain Mapp. 2006;27: 314–324.

doi:10.1002/HBM.20187

58.   Cahill L. An issue whose time has come. J Neurosci Res. 2017;95: 12–13.

doi:10.1002/JNR.23972

59.   Murray SO, Schallmo MP, Kolodny T, Millin R, Kale A, Thomas P, et al. Sex differences in

visual motion processing. Curr Biol. 2018;28: 2794-2799.e3. doi:10.1016/j.cub.2018.06.014

60.   De Vries GJ. Minireview: Sex differences in adult and developing brains: compensation,

compensation, compensation. Endocrinology. 2004;145: 1063–1068. doi:10.1210/EN.2003-1504

61.   McCarthy MM, Arnold AP. Reframing sexual differentiation of the brain. Nat Neurosci.

2011;14: 677–683. doi:10.1038/NN.2834

62.   Hyde JS. Gender similarities and differences. Annu Rev Psychol. 2014;65: 373–398.

doi:10.1146/ANNUREV-PSYCH-010213-115057

63.   Ruigrok ANV, Salimi-Khorshidi G, Lai MC, Baron-Cohen S, Lombardo M V., Tait RJ, et al. A

meta-analysis of sex differences in human brain structure. Neurosci Biobehav Rev. 2014;39: 34–

50. doi:10.1016/J.NEUBIOREV.2013.12.004

64.   Gur RC, Alsop D, Glahn D, Petty R, Swanson CL, Maldjian JA, et al. An fMRI study of sex

579   differences in regional activation to a verbal and a spatial task. Brain Lang. 2000;74: 157–170.

580   doi:10.1006/BRLN.2000.2325

581   65.   Bell EC, Willson MC, Wilman AH, Dave S, Silverstone PH. Males and females differ in brain

582   activation during cognitive tasks. Neuroimage. 2006;30: 529–538.

583   doi:10.1016/J.NEUROIMAGE.2005.09.049

584   66.   Nieto-Castañón A, Fedorenko E. Subject-specific functional localizers increase sensitivity and

585   functional resolution of multi-subject analyses. Neuroimage. 2012;63: 1646–1669.

586   doi:10.1016/J.NEUROIMAGE.2012.06.065

587   67.   Nieto-Castanon A, Ghosh SS, Tourville JA, Guenther FH. Region of interest based analysis of

588   functional imaging data. Neuroimage. 2003;19: 1303–1316. doi:10.1016/S1053-8119(03)00188-

589   5

590   68.   Poldrack RA. Region of interest analysis for fMRI. Soc Cogn Affect Neurosci. 2007;2: 67–70.

591   doi:10.1093/SCAN/NSM006

592   69.   Meoni S, Macerollo A, Moro E. Sex differences in movement disorders. Nat Rev Neurol 2020

593   162. 2020;16: 84–96. doi:10.1038/s41582-019-0294-x

594   70.   Phan HT, Reeves MJ, Blizzard CL, Thrift AG, Cadilhac DA, Sturm J, et al. Sex differences in

595   severity of stroke in the INSTRUCT study: A meta-analysis of individual participant data. J Am

596   Heart Assoc. 2019;8. doi:10.1161/JAHA.118.010235

597   71.   Persky RW, Turtzo LC, McCullough LD. Stroke in women: Disparities and outcomes. Curr

598   Cardiol Rep. 2010;12: 6. doi:10.1007/S11886-009-0080-2

599   72.   Kim JS, Lee KB, Roh H, Ahn MY, Hwang HW. Gender differences in the functional recovery

600   after acute stroke. J Clin Neurol. 2010;6: 183. doi:10.3988/JCN.2010.6.4.183

601   73.   Tomita H, Hagii J, Metoki N, Saito S, Shiroto H, Hitomi H, et al. Impact of sex difference on

602   severity and functional outcome in patients with cardioembolic stroke. J Stroke Cerebrovasc Dis.

603 2015;24: 2613–2618. doi:10.1016/j.jstrokecerebrovasdis.2015.07.016

604 74. Zhang X, Liang M, Qin W, Wan B, Yu C, Ming D. Gender differences are encoded differently in

605 the structure and function of the human brain revealed by multimodal MRI. Front Hum

606 Neurosci. 2020;14: 244. doi:10.3389/FNHUM.2020.00244/BIBTEX

607 75. Andrushko JW, Gould LA, Renshaw DW, Ekstrand C, Hortobágyi T, Borowsky R, et al. High

608 force unimanual handgrip contractions increase ipsilateral sensorimotor activation and functional

609 connectivity. Neuroscience. 2021;452: 111–125. doi:10.1016/j.neuroscience.2020.10.031

610 76. Noble JW, Eng JJ, Kokotilo KJ, Boyd LA. Aging effects on the control of grip force magnitude:

611 An fMRI study. Exp Gerontol. 2011;46: 453–461.

612 77. Noble JW, Eng JJ, Boyd LA. Effect of visual feedback on brain activation during motor tasks: an

613 FMRI study. Motor Control. 2013;17: 298–312. doi:10.1123/MCJ.17.3.298