Dynamics of the epigenetic landscape during development and in response to drought

stress in sorghum

Yongfeng Hu[1, †, *], Chao He[2, †], Yuning Shen[2], Gongjian Zeng[1], Siteng Bi[2], Quanjun Huang[1],

Xiner Qin[1], Zhuying Deng[1], Zhengquan He[1], Xiangling Shen[1,*]

[1]Key Laboratory of Three Gorges Regional Plant Genetics & Germplasm Enhancement

(CTGU)/ Biotechnology Research Center, College of Biological and Pharmaceutical Sciences,

China Three Gorges University, Yichang, Hubei 443002, China.

[2]National Key Laboratory of Crop Genetic Improvement, Hubei Hongshan Laboratory,

Huazhong Agricultural University, Wuhan, China.

[†]These authors contributed equally to this work.

[*]Corresponding author: huyongfeng@ctgu.edu.cn; shenxl1982@hotmail.com

Abstract

Sorghum bicolor is a C4 plant with the characteristics of high stress tolerance, which may be

conferred partly by the underlying epigenetic mechanism unique to sorghum. In this study, we

revealed some epigenomic features in sorghum that have never been reported before. The

long H3K27me3 regions clustered in four areas, which we defined as H3K27me3 islands,

were identified in sorghum. H3K36me3 plays some role in inhibiting the deposition of both

H3K27me3 and H2A.Z, which may serve as partial motivation for the removal of H3K27me3

and H2A.Z in leaf and root. All the 7 histone marks are involved in the regulation of

tissue-specific genes, especially the specific expression of C4 genes in leaf and peroxidase

(POD) encoding genes in root, which are involved in the photosynthesis in leaf and lignin

synthesis in root, respectively. These marks except H3K36me3 and H3K27me3 also engage in

the regulation of stress genes in response to PEG treatment. However, we found that

differential enrichment of histone marks on many tissue-specific genes was observed only

between leaf and root but hardly in response to PEG treatment, although expression of these

genes changed after PEG treatment.

Key words: sorghum, histone modification, H2A.Z, root and leaf, drought stress

Introduction

Histone marks play important roles in gene regulation, and they exhibit different distribution patterns in intragenic regions. Histone acetylation and H3K4me3 tend to mark transcription start site (TSS) regions (a short region immediately downstream from TSS), while H3K36me3 are more likely to be enriched in gene body regions (a longer region from TSS to transcription end site (TES)) (Barski et al. 2007; Guenther et al. 2007; Hu et al. 2020), suggesting that they are involved in governing different processes of gene transcription. Histone acetylation and H3K4me3 have been reported to be involved in transcription initiation by affecting the assembly of pre-initiation complex (PIC), or stimulating the transition from initiation to elongation (Ding et al. 2012; Lauberth et al. 2013; Gates et al. 2017; Wang et al. 2023). H3K36me3, generally considered as a hallmark of transcription elongation, is required for co-transcriptional splicing (Guenther et al. 2007; Iwamori et al. 2016; Sorenson et al. 2016). The distribution patterns of some histone marks are similar in different species while the others are not. For example, in humans and yeast, H3K36me3 is excluded from TSS of genes while there is a clear enrichment peak of the mark around TSS in plants (Barski et al. 2007; Li et al. 2015; Liu et al. 2016; Suzuki et al. 2016). In animals, H3K27me3 can spread to hundreds of kilobases and cover several genes per region (Schwartz et al. 2006). However, most of H3K27me3 regions in *Arabidopsis* are limited to single genes (Zhang et al. 2007). The different distribution patterns of these histone marks may reflect diverse deposition mechanism of them in different species.

The complex crosstalk between different histone modifications is established, since the activity of histone modifiers depositing one histone mark could be either promoted or inhibited by other histone marks. For instance, H2B ubiquitylation is required for simulating the activity of the MLL complex to catalyze H3K4me3 (Hsu et al. 2019). H3K4me3 provides the binding platform for the HAT complexes, facilitating histone acetylations (Taverna et al. 2006; Ringel et al. 2015). The interplay between repressive marks such as H3K27me3 and H2AK119ub1, which are deposited by Polycomb repressive complex 2 (PRC2) and Polycomb repressive complex 1 (PRC1) respectively, have also been unraveled (Blackledge and Klose 2021). Interestingly, both PRC2 and PRC1 can read the two marks to establish or reinforce these modifications (Blackledge and Klose 2021). By contrast, H3K4me3 and H3K36me3

inhibit the activity of PRC2 (Schmitges et al. 2011; Finogenova et al. 2020). Furthermore, it

has been demonstrated that H3K36me3 and H3K27me3 are rarely present on the same histone

tail in various species (Yuan et al. 2011; Yang et al. 2014), suggesting that the inhibitory effect

of H3K36me3 on deposition of H3K27me3 is conserved. Indeed, genome-wide analysis of

chromatin states indicates that the distributed regions of some histone marks highly co-exist

while the others are seldom overlapped, which is well consistent with the crosstalk between

them (Ernst et al. 2011; Sequeira-Mendes et al. 2014).

Sorghum is an ideal model species for analyzing genetic features of C4 plants due to its

small genome and natural diploid. Besides, it also exhibits high tolerance to various abiotic

stresses, making it a good candidate for studying mechanism of plant stress tolerance. In this

paper, genome-wide analysis of various histone modification enrichment was conducted in

sorghum root and leave before and after PEG treatment to explore the role of epigenetic

marks in the regulation of sorghum development and stress response. We found that

distribution pattern of these marks in sorghum was similar to that reported in other plant

species. However, sorghum-specific epigenomic features were also revealed.

Results

Distribution pattern analysis of histone marks reveals four H3K27me3 islands in

sorghum

Three replicates for RNA-seq and two biological replicates for ChIP-seq in leaf and root

of sorghum were performed with antibodies against various histone modifications (H3K9ac,

H3K27ac, H3K4me3, H3K36me3, H3K4me2 and H3K27me3) and H2A.Z. The three

replicates of RNA-seq analysis were highly correlated (Fig. S1a). For most of the ChIP-seq

samples, more than 20 million reads were obtained, the overall mapping rate was more than

85% (Table S1), the FRiP value and the SPOT value were greater than 0.5 (Fig. S1b). The two

biological replicates of ChIP-seq were also highly correlated (Fig. S1c). Genome-wide view

of histone mark enrichment demonstrate that the distribution of all the 7 histone marks was

highly correlated with gene density, which is relatively higher at both ends of each

chromosome and lower in the middle of the chromosome (Fig. S2). These results suggest that

all the data are reliable for subsequent analysis.

To attempt to investigate the distribution pattern of histone marks, intragenic peaks

(located from 1.5 kb upstream from gene TSS to 1.5 kb downstream from gene TES) were

called for creating metaplots and heat maps (Fig. 1a and 1b). The results showed that H3K9ac,

H3K27ac, H3K4me3 were predominantly enriched in TSS regions. H3K4me2 and H2A.Z

were distributed either TSS regions or gene body regions. H3K36me3 enrichment extended

from TSS in gene body but not to TES. The distribution patterns of these histone marks are

similar to those in *Arabidopsis* and rice (Zilberman et al. 2008; Roudier et al. 2009; Li et al.

2015; Hu et al. 2020). We also found strong H3K27me3 enrichment in gene body in both root

and leaf. Interestingly, low H3K27me3 enrichment in TSS regions was observed in leaf but

not in root (Fig. 1b), which has never been reported in plants before. Besides, genome-wide

enrichment of H3K4me2, H3K36me3, H3K27me3 and H2A.Z was decreased in root,

suggesting their possible regulatory roles in the fine-tuning of developmental genes (Fig. 1a).

In animals, H3K27me3 enrichment can spread to more than one hundred kilobases,

which usually cover several genes (Schwartz et al. 2006). It seems that in *Arabidopsis*

H3K27me3 coverage is confined to single gene (Zhang et al. 2007). To determine whether

H3K27me3 in sorghum is distributed similarly, we analyzed length of H3K27me3 regions and

number of genes per region. The results showed that most of H3K27me3 regions were short

and covered single genes in sorghum (Fig. 1c and 1d), which is similar as in *Arabidopsis*.

However, we found considerable long H3K27me3 regions in sorghum (Fig. 1c and 1d). The

maximum length can reach more than two hundred kilobases. Three H3K27me3 regions

covered more than 20 genes (Fig. 1d). In addition, there were more short regions and less long

regions in leaf than those in root, suggesting that H3K27me3 enrichment in the two organs are

reprogrammed. When we visualized genome-wide H3K27me3 regions, we found four areas

on Chromosomes 2, 5 and 8 where long H3K27me3 regions clustered, which we termed the

H3K27me3 islands (Fig. 1e). The other long H3K27me3 regions scattered in the other part of

the genome. The similar analysis was performed in *Arabidopsis* and rice by using the recently

deposited high quality data (see Material and methods). The long H3K27me3 regions were

also identified in *Arabidopsis* and rice despite that the number was less than that in sorghum

(Fig. S3a), especially the regions longer than 50kb. In addition, the H3K27me3 island was not

observed in *Arabidopsis* and rice (Fig. S3c). However, the genes covered by long H3K27me3

regions were greater in *Arabidopsis* (Fig. S3b), which may be due to the higher gene density.

This indicates that plant epigenomes contain both gene-specific and long region of

H3K27me3 enrichment.

Antagonistic relationship between H3K36me3 enrichment and H3K27me3 enrichment

as well as H2A.Z deposition

To analyze the relationship between different histone marks in sorghum, we clustered

distribution patterns of each histone mark in intragenic region (from 0.5kb upstream TSS to

3kb downstream TSS) into three categories (C1, C2 and C3), and observe how the other

histone marks are distributed in the same category. We found that H3K9ac, H3K27ac,

H3K4me3 enriched in the TSS region and H3K36me3 enriched in the gene body region are

highly overlapped and associated with actively transcribed genes (Fig. 2a-2d, S4a-S4d). The

gene-body- enriched H3K27me3 is exclusive with the above histone modifications especially

H3K36me3, and is associated with silent genes (Fig. 2g, Fig. S4g). The partial overlapping of

H3K4me3 and H3K27me3 suggests the existence of the bivalent mark in sorghum. The

distributions of H3K4me2 and H2A.Z in different regions displayed distinct relationships

with the other marks (Fig. 2e-2f, Fig. S4e-S4f). The strong enrichment of H2A.Z in the gene

body region (C1) overlapped with the repressive mark (H3K27me3), whereas the weak

enrichment of H2A.Z in the TSS region (C3) overlapped with the active marks (H3K9ac,

H3K27ac, H3K36me3) (Fig. 2e, Fig. S4e). By contrast, H3K4me2 deposited in the TSS

region (C1) co-existed with the repressive mark, whereas that deposited in the gene body

region (C2) co-existed with the active marks (Fig. 2f, Fig. S4f). Furthermore, we found that

H3K4me2 and H3K4me3 co-existed in many genes although distribution patterns of the two

marks in these genes were slightly different. This suggests that the nucleosome

simultaneously carrying H3K4me2 and H3K4me3 is common in sorghum, but the deposition

of the two marks may be independent.

Considering the strictly antagonistic distribution of H3K27me3 and H3K36me3, we

wonder how H3K36me3 is distributed in H3K27me3 islands. We selected one area in each

H3K27me3 island to visualize the distribution of H3K27me3 and H3K36me3. The diagram

showed that H3K27me3 and H3K36me3 seldom co-existed at the same gene (Fig. 2h, Fig.

S4h), suggesting that they may exclude each other. Indeed, the long H3K27me3 regions are

disrupted due to the deposition of H3K36me3. Thus, we speculate that there should be much

more and longer H3K27me3 regions than expected in sorghum. As H2A.Z deposition can also

extend to generate long H2A.Z regions (data not shown) and H2A.Z is absent in the gene

body regions marked by H3K36me3 (Fig. 2e), we randomly selected three long H2A.Z

regions and we found that enrichment of H3K36me3 excluded H2A.Z in these regions (Fig.

2i, Fig. S4i). However, the antagonistic relationship between H3K36me3 and H2A.Z is not as

rigorous as H3K36me3 and H3K27me3, as at some loci H3K36me3 and H2A.Z co-existed.

Taken together, these data suggest that H3K36me3 inhibits the deposition of both H3K27me3

and H2A.Z. To confirm whether the scenario is conserved in plants, we performed the similar

analysis in *Arabidopsis* and rice. The results also revealed the antagonistic distributions

between H3K27me3 and H3K36me3 as well as H2A.Z and H3K36me3 (Fig. S5a-d).

Dynamic change of histone marks during development in sorghum

In order to identify which histone marks are more likely to be related with specific gene

expression during developmental processes, correlation analysis of differential enrichment of

histone marks and gene expression in leaf and root was performed. The metaplots indicated

that the overall levels of H3K9ac, H3K27ac, H3K4me3 and H3K36me3 were increased both

at root-up-DEGs (Gene expression is up-regulated in roots compared to leaves) in root and

leaf-up-DEGs (Gene expression is up-regulated in leaves compared to roots) in leaf (Fig. 3a).

The number of genes whose expression changes were positively correlated with changes in these

marks was much greater than the number of genes whose expression was negatively correlated

with them (Fig. 3b). These supports the positive role of these marks in the regulation of gene

expression. In contrast, H2A.Z, H3K4me2 and H3K37me3 levels were significantly

decreased at root-up-DEGs in root but slightly decreased at leaf-up-DEGs in leaf (Fig. 3a),

and genes negatively regulated by these marks were more than those positively regulated (Fig.

3b), suggesting negative correlation of these marks with gene expression.

To investigate how different histone marks cooperate to regulate root-leaf-DEGs

(differentially expressed genes between root and leaf), we calculated the proportions of genes

that were co-regulated by two histone marks. Considering the opposite roles of different

histone marks in gene regulation as described above, the number of genes with increased

H3K9ac, H3K27ac, H3K4me3, H3K36me3 or decreased H2A.Z, H3K4me2, H3K27me3

between leaf and root were used for the calculation. The higher proportions of overlapping

(more than 30%) were observed between the increase of H3K9ac and the other active marks

(H3K27ac, H3K4me3 and H3K36me3) as well as the increase of H3K4me3 and H3K36me3

in both leaf (vs root) and root (vs leaf) (Fig. 3c), suggesting that these histone marks tend to

activate gene expression collaboratively. The decrease of H3K4me2 and H3K27me3 also

showed a high percentage (30.63%) of overlapping only in leaf (vs root) (Fig. 3c), which

demonstrates that they cooperate to repress more gene expression in root. We found that the

proportions of overlapping between the increase of H3K36me3 and the decrease of

H3K27me3 as well as the decrease of H2A.Z deposition were not as high as expected (8.57%

and 15.69% in leaf (vs root), 8.15% and 10.15% root (vs leaf)) (Fig. 3c). This indicates that

H3K36me3 enrichment contribute to small part of decline of H3K27me3 enrichment or

H2A.Z deposition. In addition, the heat map showed that nearly half of genes with increased

H3K36me3 were not marked with H3K27me3 (Fig. 3d). However, most of strong H3K27me3

enrichment were inhibited by the increase of H3K36me3. Similarly, the relatively higher

levels of H2A.Z deposition were decreased to moderate levels after H3K36me3 enrichment

was increased (Fig. 3e). These results further support the notion that H3K36me3 play part of

the roles in the inhibition of H3K27me3 enrichment or H2A.Z deposition.

As shown in Fig. 3a, the differential enrichment of H3K27me3 in root-up-DEGs and

leaf-up-DEGs in leaf and root was also obvious outside the gene body region (upstream of

TSS and downstream of TES), suggesting that the dynamic change of H3K27me3 could

spread outside the intragenic region. Alternatively, H3K27me3 deposition changes in units of

long segments instead of single genes. To confirm the suspicion, we examined the number of

H3K27m3 peaks with different length differentially enriched in root and leaf and found that

although most of differentially enriched H3K27me3 peaks were short (less than 10 kb), more

than 300 long H3K27me3 regions (more than 10 kb), consisting of 278 (10-20 kb), 87 (20-40

kb), 12 (40-60 kb), 5 (60-80 kb) and 4 (80-100 kb), were also differentially deposited (Fig.

S6a). Genome-wide screenshots indicated that the changes of long H3K27me3 regions

scattered in 10 chromosomes and negatively co-related with gene expression in general (Fig.

S6b). But close observation on individual regions showed that a number of genes covered by

long H3K27me3 regions were not activated despite removal of H3K27me3 (Fig. S6b). For

example, two neighboring long H3K27me3 regions decreased in root and localized in

chr5:70,880 kb-71,000 kb cover 11 genes, among which only 3 genes were up-regulated (Fig.

S6b). This suggests that the alteration of H3K27me3 enrichment in long regions is not always

associated with transcriptional activity.

Leaf-specific expression of photosynthesis genes and root-specific expression lignin

biosynthesis genes involves epigenetic regulation

The Kyoto Encyclopedia of Genes and Genomes (KEGG) and Gene Ontology (GO)

analysis of root-leaf-DEGs with changed histone marks were performed to discover which

pathways were more likely to be subjected to epigenetic regulation. The result showed that

leaf-up-DEGs with various changed histone marks were enriched in photosynthesis pathway

and localized to chloroplast or plastid. The root-up-DEGs were enriched in Phenylpropanoid

biosynthesis pathway and localized to cell wall (Fig. 4a and Fig. S7). For a more detailed

analysis of which genes in these pathways were epigenetically regulated, we investigated

changes of expression and histone marks of all the photosynthesis genes encoding

Photosystem I, Photosystem II, Cytochrome b6/f, ATP synthase, plastocyanin, cytc6 as well

as chlorophyll biosynthesis genes. We found that expression of most photosynthesis genes

and chlorophyll biosynthesis genes encoding the enzymes that catalyze the reactions from

protoporphyrin IX to chlorophyll was dramatically increased in leaf (Fig. 4b). The increased

expression of these genes was strongly associated with increased enrichment of H3K9ac,

H3K27ac, H3K4me3 and H3K36me3, and partially with decreased enrichment H2A.Z,

H3K4me2 and H3K27me3 (Fig. 4b). In particular, dynamic change of H3K27me3 occurred

only on a very small proportion of these leaf-specific genes. Besides, a few of these genes

were marked with high level of H3K27me3 in root (data not shown), suggesting that

H3K27me3 is not required for repression of most of photosynthesis genes in root compared to

the other histone marks.

As sorghum is a C4 plant, we would like to learn how C4 genes are regulated by

different histone marks. In sorghum there were nine C4 genes: *carbonic anhydrase 2* (*CA2*),

CA3, *NADP-malate dehydrogenase 2* (*NADP-MDH2*), *nicotinamide adenine dinucleotide*

phosphate (NADP)-malic enzyme 2 (*NADP-ME2*), *phosphoenolpyruvate carboxylase 6*

(*PEPC6*), *PEPC kinase 2* (*PPCK2*), *pyruvate orthophosphate dikinase 2* (*PPDK2*), *PPDK*

regulatory protein 1 (*PPDK-RP1*) and *RbcS* (Tao et al. 2020). We observed that seven C4

genes (except *CA3* and *PPCK2*) exhibited strong leaf expression specificity, and were mostly

associated with differential enrichment of H3K9ac, H3K4me3 and H3K36me3 (Fig. 4c and

4d). Among these histone marks, H3K4me3 varied more significantly than the other marks,

implying that H3K4me3 acts as the major epigenetic mark for the leaf-specific regulation of

C4 gene expression. In addition, similar to the other photosynthesis genes, H3K27me3 was

negatively associated with the expression of *CA2* and *PEPC6* only (Fig. 4c and 4d),

suggesting a limited regulatory role of H3K27me3 on C4 genes. By contrast, the non-C4

genes homologous to C4 genes showed weaker leaf expression specificity and fewer of them

were regulated by dynamic change of histone marks (Fig. 4c). These results suggest that the

active histone marks may confer the leaf expression specificity of photosynthesis genes in

sorghum.

When we looked into the root-up-DEGs with differentially enriched histone marks in the

Phenylpropanoid biosynthesis pathway, we found many of these genes encode lignin

biosynthesis enzymes, especially peroxidases (PODs) (data not shown). Therefore, we

examined changes of expression and histone marks of all the lignin biosynthesis genes (Fig.

5a). It was shown that for 20 of 52 lignin biosynthesis genes, up-regulation of them in root

involves dynamic change of at least one histone mark (Fig. 5b). And the epigenetically

regulated genes encode all the enzymes for lignin biosynthesis except C3H, echoing the

importance of epigenetic regulation in the lignin biosynthesis in root. More interestingly, we

found that vast majority of *POD* genes showed root expression specificity that were

associated with most of histone marks (Fig. 5c), although it is not clear how many of them are

involved in lignin biosynthesis, especially in root. Particularly, all the 7 histone marks were

differentially enriched on four *POD* genes (Fig. 5d). However, It seems that H3K36me3 is not

involved in the regulation of most *POD* genes, as they were not marked with H3K36me3

even when they were significantly up-regulated in root (Fig. 5c). Unlike photosynthesis genes,

we found decreased enrichment of H3K27me3 on considerable *POD* genes (Fig. 5c),

suggesting that repression of these *POD* genes concerns H3K27me3.

Gene regulation in response to drought stress concerns histone marks except H3K36me3

and H3K27me3

To explore how histone marks engage in the induction of drought-responsive genes, we

investigated the levels of histone marks in PEG-up-DEGs and PEG-down-DEGs in response

to PEG treatment in root and leaf. The metaplots showed that PEG treatment only slightly

affects the overall levels of some histone marks (Fig. 6a and 6b). For example, H3K9ac,

H3K27ac, H3K4me3, H3K36me3 levels in PEG-up-DEGs in leaf and H3K36me3 level in

PEG-up-DEGs in root showed a little increase. H3K9ac and H3K27ac levels in

PEG-down-DEGs in leaf and root was slightly decreased. H2A.Z and H3K4me3 levels in

PEG-down-DEGs in leaf and root rose and H2A.Z level in PEG-up-DEGs in leaf and root

declined. These results are consistent with the regulatory roles of these marks. Then we

performed association analysis of changes of histone marks and gene expression. In both leaf

and root, only a few genes had altered H3K36me3 and H3K27me3 responding to PEG

treatment (Fig. 6c and 6d), which suggests that these two marks are probably not involved in

the PEG-responsive gene expression regulation. H3K4me3 was more likely to be associated

with gene activation in response to PEG treatment, as the number of genes with increased

H3K4me3 levels and gene expression in leaf (n = 138) and root (n = 363) was far more than

the number of genes with the other relationships between H3K4me3 and gene expression (Fig.

6c and 6d). H3K9ac and H3K27ac were associated with either gene activation or repression

when the level of these marks was increased or decreased responding to PEG treatment in leaf

and root (Fig. 6c and 6d). Surprisingly, for certain number of genes in root but few in leaf

changes of H3K9ac and H3K27ac were negatively correlated with changes of gene

expression, which seems to contradict to the positive role of these marks. Finally, H2A.Z and

H3K4me2 were negatively associated with some gene activation or repression in leaf and root

(Fig. 6c and 6d). These results demonstrate that only a small proportion of gene induction in

response to PEG treatment require dynamic change of histone marks.

In order to find out whether histone marks affect the degree of gene induction by PEG

treatment, we classified PEG-up-DEGs into three groups: genes without enrichment of

histone marks, genes with unchanged histone marks after PEG treatment, genes with changed

histone marks after PEG treatment, and investigated fold change of each group gene

expression. As PEG-responding gene up-regulation are more associated with increase of

H3K9ac, H3K27ac, H3K4me3 and decrease of H2A.Z, the effect of these marks on gene

induction was analyzed. We found that the degree of induction of the genes with changed

histone marks after PEG treatment was significantly higher than that of the other groups (Fig.

S8), suggesting that the dynamic change of histone marks can elevate gene induction to some

extent. However, the genes with unchanged H3K9ac, H3K27ac and H3K4me3 were induced

to a relatively lower level than the genes without histone marks (Fig. S8). Conversely, the

genes with unchanged H2A.Z enrichment were induced to a higher level (Fig. S8). These

results demonstrate that the basic level of histone marks is not related to the gene

responsiveness.

The direct regulation of many stress genes by transcription factors has been

characterized in different plant species such as the genes involved in abscisic acid (ABA)

metabolism, signaling and reactive oxygen species (ROS)-related and aquaporins, *late*

embryogenesis abundant (LEA), wax biosynthese, polyamine biosynthesis genes (Hu et al.

2022). We found that induction of some of these stress genes also involves epigenetic

regulation. For example, *9-cis-epoxycarotenoid dioxygenase 3* (*NCED3*), the key gene in

ABA biosynthesis pathway, and *delta1-PYRROLINE-5-CARBOXYLATE SYNTHASE 1*

(*P5CS1*), the key gene in proline biosynthesis pathway, were up-regulated by PEG treatment

in both leaf and root (Table S2). The up-regulation was associated with the increase of

H3K9ac, H3K27ac and H3K4me3 (Table S2). Surprisingly, expression of 9 in 10 clade A type

2C protein phosphatases (*PP2C*s) genes, the repressors of ABA signaling, was also increased

(Table S2), which usually entails increased enrichment of H3K9ac, H3K27me3, H3K4me3

and decreased enrichment H2A.Z in leaf and root. In addition, epigenetic regulation of many

LEA genes in response to PEG was also observed (Table S2), although the scenario was

different between leaf and root. The expression of these *LEA* genes was induced to a higher

degree in leaf where change of epigenetic marks was more significant (Table S2). It is of great

interest to ascertain if the higher degree of *LEA* expression is determined by changed

epigenetic marks, and whether the differential induction of *LEA* genes in leaf and root has

some physiological significance.

325 Dynamic change of histone marks is not necessary for the response of tissue-specific

326 genes to drought stress

To learn that for the genes both differentially expressed between leaf and root and

responsive to PEG treatment whether the histone marks also change in both conditions

(developmentally regulated and PEG responsive), we identified 780 and 2825 Root-up-DEGs

that were also PEG-up-DEGs in leaf and PEG-down-DEGs in root respectively, 1359 and 950

Leaf-up-DEGs that were also PEG-down-DEGs in leaf and PEG-up-DEGs in root

respectively (Fig. 7a), which were defined as four groups of common DEGs. Then we

investigated the number of genes where histone marks changed uniquely in each condition or

in both conditions among each of four groups of common DEGs. For the up-regulated genes

increase of H3K9ac, H3K27me3, H3K4me3, H3K36me3 and decrease of H2A.Z, H3K4me2,

H3K27me3 were analyzed, and for the down-regulated genes decrease of H3K9ac,

H3K27me3, H3K4me3, H3K36me3 and increase of H2A.Z, H3K4me2, H3K27me3 were

analyzed. The results revealed that for all four groups of common DEGs, the number of genes

with change of histone marks uniquely regulated developmentally was much greater than

those uniquely responsive to PEG and those in both conditions (Fig. 7b). Moreover, we found

some common DEGs where 7 histone marks changed developmentally but none of them

changed in response to PEG (Fig. S9). In particular, as we revealed that leaf-specific

expression of photosynthesis genes and root-specific expression of lignin biosynthesis genes

were associated with dynamic change of histone marks, we attempt to understand how these

gene expression was affected by stress. We found that many photosynthesis genes including

four C4 genes and lignin biosynthesis genes were slightly down-regulated (Fig. 7c and 7d, Fig.

S10), and more than half of *POD* gene expression was decreased to a varying degree after

PEG treatment (Fig. 7e). However, histone marks on only a small proportion of these genes

were decreased. This indicates that dynamic change of histone marks important for

tissue-specific regulation of gene expression is not necessary for the gene induction in

response to environmental cues, and some tissue-specific regulatory factors are possibly

required for the alteration of histone marks.

Cis-elements enriched in the promoter of genes with changed histone marks

Through above observations that the responses of chromatin dynamics to developmental

or environmental cues were distinct in leaf and root, we considered that tissue-specific factors

may direct the change of chromatin structure. Transcription factors (TFs) are the best

candidates for the directing factors. *Cis*-elements analysis of promoter of genes (1.5kb

upstream from TSS) with changed histone marks was performed to predict which kinds of TFs are responsible for directing the writing or erasing of histone marks in different tissues. The five most enriched *cis*-elements for each variation of histone marks were shown in Fig. 8. For developmental genes, the *cis*-elements of Dof, AT-hook, MADS families TFs were associated with most of histone mark variations (Fig. 8a), suggesting that they were generally involved in the deposition or removal of histone marks in different tissues. The C2H2 Zinc finger TFs binding *cis*-element was inclined to be enriched in the genes with down-regulated histone marks in leaf (Fig. 8a), making it a good candidate as the tissue-specific guider for changing histone marks. In addition, ARID/BRIGHT TFs were responsible for some of histone mark variation (Fig. 8a). More specifically, YABBY TFs were associated with deposition of H3K36me3 in leaf (Fig. 8a). For PEG-induced genes, Dof, AT-hook, C2H2 Zinc finger and AP2 TFs were identified to be the most enriched *cis*-elements binding TFs (Fig. 8b). The bZIP TFs were more likely to be associated with up-regulation of H3K4me3 in leaf (Fig. 8b). MADS box *cis*-element was mostly enriched in the genes with decreased H3K9ac in leaf (Fig. 8b).

Discussion

The unique feature of H3K27me3 enrichment and function in sorghum

In this study we investigate the genome-wide enrichment of histone marks including H3K9ac, H3K27ac, H3K4me3, H3K36me3, H3K4me2, H3K27me3 and H2A.Z in sorghum. As reported in other plants, the distribution patterns of these marks in intragenic region are conserved in sorghum. For instance, H3K9ac, H3K27ac and H3K4me3 tend to be enriched in the TSS region, and H3K36me3, H3K4me2, H3K27me3 and H2A.Z were distributed in the whole transcribed region (Fig. 1a and 1b). However, early report revealed that H3K27me3 enrichment peaks in *Arabidopsis* was much shorter than that in animals (usually 20-50kb) and limited to single genes mostly (Zhang et al. 2007). But our analysis in *Arabidopsis*, rice and sorghum indicates that there are considerable long H3K27me3 regions in these species despite much less than the short regions (Fig. 2a and Fig. S3a). The maximum length of H3K37me3 region can reach more than 200 kb in sorghum. These results suggest that H3K27me3 enrichment can also spread to a long region covering multiple genes in plants. In Drosophila,

spreading of H3K27me3 requires specific *cis*-regulatory sequences, called Polycomb

response elements (PREs) and pre-existing H3K27me3 (Margueron et al. 2009). In

mammalian, spatially interacted nucleation sites are targeted by PRC2 complex to create

H3K27me3-forming Polycomb foci for initiation of H3K27me3 spreading (Oksuz et al. 2018).

Indeed, we found four H3K27me3 islands where the long H3K27me3 regions were clustered

on three chromosomes of sorghum but not in *Arabidopsis* and rice (Fig. 2c and Fig. S3c). It is

intriguing to unveil whether the H3K27me3 islands in sorghum are generated by spreading

from PREs as in Drosophila or nucleation sites as in mammalian in the future. It has been

disclosed that H3K27me3-mediated chromatin interaction takes place in *Arabidopsis* (Huang

et al. 2021; Yang et al. 2022), but whether the interaction of chromatin is the prerequisite for

H3K27me3 spreading remains to be elucidated. In our opinion, analysis of 3D genome

structure of sorghum to reveal whether the four H3K27me3 islands are interacted is necessary

for the next step. Besides, LIKE HETEROCHROMATIN PROTEIN 1 (LHP1), the reader of

H3K27me3 in plant, has been reported to be responsible for the spreading of H3K27me3

towards the 3' end of the gene body in *Arabidopsis* (Veluchamy et al. 2016). It is of great

importance to investigate whether the homolog of LHP1 in sorghum is involved in the

spreading of H3K27me3 enrichment to a longer region.

H3K36me3 deposition inhibit both H3K27me3 and H2A.Z enrichment

Inhibition of PRC2 activity by H3K36m3 has been reported in animal and plant by

biochemical analysis (Yuan et al. 2011; Yang et al. 2014), but whether H3K36me3 can serve

as an active mark to restrict H3K27me3 spreading is not clear. Comparative analysis of

genome-wide distribution of the two marks is missing. Here, by comparing genome-wide

distribution of 7 histone marks in intragenic regions we found that gene-body-enriched

H3K27me3 is depleted in the genes marked with H3K9ac, H3K27ac, H3K36me3 (Fig. 3a, 3b

and 3d). But among these active marks only H3K36me3 is distributed in the entire gene body

region which is similar to the distribution pattern of H3K27me3 (Fig. 3d and 3g), making it a

suitable marker for inhibiting H3K27me3 deposition. When we visualized H3K36me3

enrichment in H3K27me3 island, we obtained mutually exclusive distribution patterns of

H3K27me3 and H3K36me3 (Fig. 3h). In the long H3K27me3 regions, a gap would be present

where H3K36me3 is deposited, indicating the inhibitory effect of H3K36me3 deposition on

H3K37me3 modification. But on the other hand, this also demonstrates that H3K36me3

cannot prevent the spreading of H3K27me3 since H3K27me3 is abundant at both sides of

H3K36me3 region. It is of great interest to explore how H3K27me3 spreads over H3K36me3

regions. We also observed that strong enrichments of H3K36me3 were most associated with

weak H2A.Z enrichments in the TSS region but hardly associated with gene-body-enriched

H2A.Z (Fig. 3d and 3e). This reminds us of the antagonistic relationship between H2A.Z and

DNA methylation in *Arabidopsis* (Zilberman et al. 2008). We found that similar to

H3K27me3, strong enrichment of H3K36me3 disrupted the long H2A.Z regions (Fig. 3i),

documenting that the enrichment of H3K36me3 also inhibit H2A.Z deposition. In mammals,

H3K36me2/3 serves as platform to recruit DNMT3A for DNA methylation in the gene body

in order for preventing spurious transcription initiation (Yano et al. 2022), but the scenario has

not been revealed in plants. It is possible that one of the main functions of H3K36me3 and

DNA methylation is to promote gene body H2A.Z eviction in plants. In *Arabidopsis*

enrichment of H3K27me3 and deposition of H2A.Z are mutually reinforced (Carter et al.

2018), raising the possibility that the active marks H3K36me3 and DNA methylation can

counteract both H2A.Z and H3K27me3. However, our results showed that H3K36me3

variation overlapped with part of changes of H2A.Z and H3K27me3 between leaf and root

(Fig. 4c), suggesting that H3K36me3 deposition is not the main cause of removal of H2A.Z

and H3K27me3.

The role of histone marks in the regulation of tissue-specific and stress-responsive genes

in sorghum

We compared genome-wide enrichment of histone marks in leaf and root to unravel to

what extent do histone marks affect transcriptional reprogramming of tissue-specific genes.

The number of differential intragenic enrichments of each histone marks is much less than

that of DEGs, suggesting that specific genes were epigenetically regulated in response to

developmental cues. A large number of photosynthesis genes including C4 genes and lignin

biosynthesis genes were among these genes (Fig. 4b-4d, Fig. 5b). The photosynthesis genes

exhibit leaf-specific expression which is consistent with their function. Epigenetic regulation

of some C4 genes have been reported in several species including sorghum (Offermann et al.

2006; Heimann et al. 2013; Li et al. 2017). Here we reported differential enrichment of

histone marks, particularly H3K4me3, on all C4 genes except PPCK2 in leaf and root of sorghum. But the homologous non-C4 genes were not expressed in a leaf-specific pattern and fewer of them were associated with change in histone marks. This suggest that the introduction of epigenetic marks onto C4 genes during evolution of C4 plants may be critical for their leaf-specific expression pattern. The single cell ChIP analysis is further required to determine whether cell-type-specific expression of C4 genes is conferred by cell-type-specific H3K4me3 enrichment, as C4 genes show mesophyll or bundle sheath cell-type-specificity in leaf. Many lignin biosynthesis genes showed root-specific expression. In sorghum, it has been reported that mutation of *brown midrib 12* (*bmr12*), which encodes caffeic acid O-methyltransferase (COMT) and is one of the key enzymes in lignin biosynthesis, leads to reduced lateral root density and altered root anatomy (Saluja et al. 2021). In this study, we revealed that activation of *bmr12* (SORBI_3007G047300) in root mainly implicated the increase of H3K4me3, H3K36me3 and the decrease of H3K27me3. We also identified many *POD* genes that showed strong root-specific expression pattern and were regulated by various histone marks, although whether they are involved in lignin biosynthesis remains to be elucidated. These results indicate that lignin biosynthesis under epigenetic regulation is important for sorghum root development. However, considering the multiple function of PODs (Cosio and Dunand 2009), it is very likely that they control root development through other pathways instead of lignin biosynthesis. Indeed, the role of POD in root development has been characterized in *Arabidopsis* (Marzol et al. 2022).

The dynamic change of histone marks in response to PEG in leaf and root was also compared in this study. H3K9ac, H3K27ac, H3K4me3, H3K4me2, and H2A.Z are associated with gene induction in varying degree in response to PEG (Fig. 5a-5d). The fold changes of the genes with changed histone marks were greater than those of the genes with unchanged histone marks or without enrichment of histone marks (Fig. S8), suggesting that the variation of histone marks can improve the degree of gene induction. However, H3K36me3 and H3K27me3 enrichments in few DEGs changed both in leaf and root after PEG treatment (Fig. 5c and 5d), suggesting that these two marks are probably not involved in PEG-responsive gene regulation in the process of response in sorghum. The study in moss Physcomitrella patens also showed that H3K27me3 change was not significant in response to drought stress

(Widiez et al. 2014). However, in *Arabidopsis* salt priming results in shortening and

fractionation of H3K27me3 islands (Sani et al. 2013). The H3K27me3 dynamics responding

to salt and cold stress has also been revealed in rice (Zheng et al. 2019; Dasgupta et al. 2022).

These demonstrate that different stress signals in different species induce distinct responses of

H3K27me3 landscape.

Materials and methods

Plant growth conditions and PEG treatment

Sorghum (BTx623 variety) seeds were surface-sterilized and then soaked for

germination. The germinated seeds were transferred to the nursery box (1/2 MS) to continue

growing in a growth room kept at 28□ with a 12h light/12 dark cycle. The 14-day-old

seedlings were transferred to the nursery box containing 20% PEG6000 that was prepared

with 1/2 MS. The seedlings transferred to fresh 1/2MS solution was used as the control. After

6 hours treatment, the third leaves and roots were harvested respectively for subsequent

experiments.

RNA-seq

The leaves and roots were harvested and frozen immediately in liquid nitrogen. Total

RNA was extracted using TRNzol Universal Reagent (TIANGEN) following the

manufacturer's instructions. The quality of RNA was measured by gel electrophoresis,

Nanodrop analyzer, Labchip and Qubit analyzer respectively. mRNA was purified by using

oligo(dT) and then fragmented by incubating in fragmentation buffer. The fragmented mRNA

was primed with random hexamer primers and reverse-transcribed with Reverse Transcriptase.

After end repair, adenylation, adaptor ligation, purification,PCR amplification, and quality

control, sequencing was performed on an Illumina HiSeq system.

ChIP-seq

Chromatin immunoprecipitation (ChIP) was performed as described in Hu et al. (Hu et al.

2020), In brief, the samples were crosslinked with 1% (v/v) formaldehyde. The nuclei were

extracted and the chromatin DNA was broken into 200-500bp fragments by sonication

(Soniprep 150, energy 5, 15s on and 1min off for 5 times). The antibodies H3K9ac (07-352,

Millipore), H3K27ac (07-360, Millipore), H3K4me2 (07-030, Millipore), H3K4me3 (07-473,

Millipore), H3K27me3 (A2363, ABclonal), H3K36me3 (ab9050, abcam), H2A.Z (preparation using antigen "GKGLLAAKTTAAKSAEKDKGKKAPV") were incubated with Protein A agarose (16-157, Millipore) before mixed with sonicated chromatin. The immunoprecipitated DNA was purified for constructing libraries and sequenced on an Illumina HiSeq system.

Raw sequencing reads filtering.

To obtain high quality clean reads of RNA-seq, ChIP-seq and MNase-seq, the raw sequencing reads were trimmed by Trimmomatic v0.32 (https://github.com/usadellab/Trimmomatic) with parameters of "TruSeq3-PE.fa:2:30:10 LEADING:20 TRAILING:20 SLIDINGWINDOW:4:15 MINLEN:36".

Analyses of RNA-seq data.

Transcript abundance was quantified using pseudoalignment of high quality clean RNA-seq reads to the reference cDNA sequences and gene models from Sorghum_bicolor_NCBIv3 assembly of the variety BTx623 (McCormick et al. 2018), as implemented in Kallisto (v0.48.0) (Bray et al. 2016). The output pseudoalignment bam files were converted to bigwig format using "bamCoverage" tool of deepTools (v3.5.0) (Ramirez et al. 2016). Differentially expressed genes (DEGs) were identified using DESeq2 v1.22 (Love et al. 2014), genes with TPM value higher than 1 in all the three replicates of each sample were regarded as expressed genes, genes with $|\log2 (Fold change)| > 1$, P adjust value < 0.05 and at least expressed in one sample were regarded as DEGs.

Analyses of ChIP-seq data.

High quality clean reads were aligned against the reference Sorghum_bicolor_NCBIv3 genome assembly using Bowtie2 (v2.4.4) (Langmead and Salzberg 2012). Aligned reads with MAPQ < 5 were filtered using SAMtools (v1.9) (Li et al. 2009). Duplicated alignments were removed using Picard (v2.23.9) (http://broadinstitute.github.io/picard/). Before peak calling, the extsize were predicted by phantompeakqualtools (v1.2.2) (Landt et al. 2012). Narrow peaks of H3K9ac, H3K27ac and H3K4me3 ChIP-seq reads densities were identified using MACS2 (v2.2.7.1) (Zhang et al. 2008) with parameters "-f BAM -p 1e-5 --mfold 2 20 --nomodel --extsize $spp_result --to-large". The input control were used by added parameter "-c". For broad peaks calling including H3K4me2, H3K27me3, H3K36me3 and KH2A.Z, the "--broad" parameter was added. The peaks from biological replicate samples were merged by BEDTools (v2.27) (Quinlan and Hall 2010), two replicate bam files were merged by SAMTools. We then extracted the reads from the merged bam files that mapped on the merged peaks using BEDtools, and converted the output new bam files to RPKM normalized bigwig files by deepTools. The metaplot and heat map profiles of epigenetic and genomic features were generated and plotted by deepTools.

The signal portion of tags (SPOT) and the fraction of reads in called peak regions (FRiP) value were calculated using Hotspot v4.1.0 (https://github.com/rthurman/hotspot) and commands in-house. To assess the repeatability between biological samples, we calculated the correlation coefficients between samples using deepTools. The visualization of reads coverage is available on http://crispr.hzau.edu.cn/cgi-bin/CRISPR-Cereal/gb2/gbrowse/Sorghum_bicolor/.

Visualization of overall signals.

All types of signals were visualized using Circos (version 0.69) (Krzywinski et al. 2009) with the commands available on github (https://github.com/hcph/Sorghum-omics-dataset-analysis).

Analyses of H3K27me3 regions.

The H3K27me3 peaks in bed files and the signal normalized bigwig files were downloaded from the ChIP-hub databases (Fu et al. 2022) using the sample ID of SRX7734769 and SRX7734770 for *Arabidopsis*, and SRX7426661 and SRX7426662 for rice. Peaks in the two replicates with 1 bp overlap (MG0) as well as overlap regions lower than 1,000 bp (MG1000) were merged using BEDTools. The number of H3K27me3 regions with different length distributedwere visualized. To investigate the distribution of H3K27m3 regions with different number of overlapped genes, we firstly found the overlapped genes for each H3K27me3 region, then the number of overlapped genes was calculated. The *Arabidopsis* and rice H3K27me3 regions longer than 10 kb were extracted and the H3K27me3 signal located in these regions were visualized using pyGenomeTracks v3.5 (Lopez-Delisle et al. 2021).

Detection of differentially histone modified genes (DMGs).

Reads mapped to genes were extracted as described before. The reads density on each genes was calculated by Subread v2.0.0 (Liao et al. 2013). The TPM value was used to quantified the reads density on each gene, followed by DMGs detection using DeSeq2. Genes with TPM value higher than 10 in the two replicates were regarded as modified genes, and genes with |log2 (Fold change)| > 0.75, *P* adjust value < 0.05 and at least being modified in one sample were regarded as DMGs. The pearson correlation coefficients for the differentially modification and expression levels was calculated. Long (>10 kb) H3K27me3 signals located in differentially H3K27me3 modified regions were extracted with BEDtools as described before, the extracted H3K27me3 signal and the corresponding gene expression signal were visualized with Circlize v0.4.15 (Gu et al. 2014) .

Analyses of relationship between different histone modifications.

The signal matrix of the seven histone modification marks was built by deepTools using the merged and RPKM normalized bigwig files as input. The signal of a single mark was used in turn as a template for clustering the signals of the remaining marks into three groups. The datasets in rice (Zheng et al. 2019; Lu et al. 2020; Zhao et al. 2020) and *Arabidopsis* (Carter et al. 2018; Xi et al. 2020) were downloaded from ChIP-hub and analyzed in the same way as

in sorghum. The signals located in long peaks (> 10 kb) were extracted and visualized as

described before.The number of DMGs with two differential types of histone modifications or

only differentially in one histone modifications were calculated and visualized.

GO and KEGG enrichment analysis.

GO and KEGG enrichment was performed using g:Profiler

(https://biit.cs.ut.ee/gprofiler/gost), the enriched GO terms and KEGG pathways were

visualized using R package ggplot2 v3.3.5.

Motif enrichment.

The promoter sequences (upstream 1,500 bp) of DMGs were extracted using BEDTools. The

sequences were submitted into the CentriMO (https://meme-suite.org/meme/tools/centrimo) online

analysis toolkit for motif enrichment analysis. The "anywhere" mode was selected to perform

local motif enrichment and the motif database was supported by CIS-BP v2.0 (Weirauch et al.

2014).

Declarations

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The ChIP-seq, MNase-seq and RNA-seq data generated in this article were deposited in the NCBI

Sequence Read Archive (SRA) (http://trace.ncbi.nlm.nih.gov/Traces/sra/) under the BioProject

accession number PRJNA952350.

Competing interests

The authors declare that they have no competing interests

Funding

This work was supported by Research Initiation Fund for High-level Talents of China Three

Gorges University.

Authors' contributions

615 The project was conceived by YH and supervised by XS. YH performed ChIP-seq,

616 MNase-seq and RNA-seq with help from QH, GZ, XQ and ZD. CH performed the

617 bioinformatics analyses with help from YS and SB. The manuscript was drafted by YH and

618 XS and revised by all the authors. ZH participate in the critical discussion. The authors read

619 and approved the final manuscript.

Acknowledgements

622 We sincerely thank the computing platform of the National Key Laboratory of Crop Genetic

623 Improvement in HZAU for providing the computational resources.

References

Barski A, Cuddapah S, Cui K, Roh TY, Schones DE, Wang Z, Wei G, Chepelev I, Zhao K. 2007. High-resolution profiling of histone methylations in the human genome. Cell 129 (4):823-837.

Blackledge NP, Klose RJ. 2021. The molecular principles of gene regulation by Polycomb repressive complexes. Nat Rev Mol Cell Biol 22 (12):815-833.

Bray NL, Pimentel H, Melsted P, Pachter L. 2016. Near-optimal probabilistic RNA-seq quantification. Nat Biotechnol 34 (5):525-527.

Carter B, Bishop B, Ho KK, Huang R, Jia W, Zhang H, Pascuzzi PE, Deal RB, Ogas J. 2018. The Chromatin Remodelers PKL and PIE1 Act in an Epigenetic Pathway That Determines H3K27me3 Homeostasis in Arabidopsis. Plant Cell 30 (6):1337-1352.

Cosio C, Dunand C. 2009. Specific functions of individual class III peroxidase genes. J Exp Bot 60 (2):391-408.

Dasgupta P, Prasad P, Bag SK, Chaudhuri S. 2022. Dynamicity of histone H3K27ac and H3K27me3 modifications regulate the cold-responsive gene expression in Oryza sativa L. ssp. indica. Genomics 114 (4):110433.

Ding Y, Ndamukong I, Xu Z, Lapko H, Fromm M, Avramova Z. 2012. ATX1-generated H3K4me3 is required for efficient elongation of transcription, not initiation, at ATX1-regulated genes. PLoS Genet 8 (12):e1003111.

Ernst J, Kheradpour P, Mikkelsen TS, Shoresh N, Ward LD, Epstein CB, Zhang X, Wang L, Issner R, Coyne M et al. 2011. Mapping and analysis of chromatin state dynamics in nine human cell types. Nature 473 (7345):43-49.

Finogenova K, Bonnet J, Poepsel S, Schafer IB, Finkl K, Schmid K, Litz C, Strauss M, Benda C, Muller J. 2020. Structural basis for PRC2 decoding of active histone methylation marks H3K36me2/3. Elife 9:e61964.

Fu LY, Zhu T, Zhou X, Yu R, He Z, Zhang P, Wu Z, Chen M, Kaufmann K, Chen D. 2022. ChIP-Hub provides an integrative platform for exploring plant regulome. Nat Commun 13 (1):3413.

Gates LA, Shi J, Rohira AD, Feng Q, Zhu B, Bedford MT, Sagum CA, Jung SY, Qin J, Tsai MJ et al. 2017. Acetylation on histone H3 lysine 9 mediates a switch from transcription

initiation to elongation. J Biol Chem 292 (35):14456-14472.

Gu Z, Gu L, Eils R, Schlesner M, Brors B. 2014. circlize Implements and enhances circular

visualization in R. Bioinformatics 30 (19):2811-2812.

Guenther MG, Levine SS, Boyer LA, Jaenisch R, Young RA. 2007. A chromatin landmark

and transcription initiation at most promoters in human cells. Cell 130 (1):77-88.

Heimann L, Horst I, Perduns R, Dreesen B, Offermann S, Peterhansel C. 2013. A Common

histone modification code on C4 genes in maize and its conservation in Sorghum and Setaria

italica. Plant Physiol 162 (1):456-469.

Hsu PL, Shi H, Leonen C, Kang J, Chatterjee C, Zheng N. 2019. Structural Basis of H2B

Ubiquitination-Dependent H3K4 Methylation by COMPASS. Mol Cell 76 (5):712-723 e714.

Hu Y, Chen X, Shen X. 2022. Regulatory network established by transcription factors

transmits drought stress signals in plant. Stress Biology 2 (1):1-22.

Hu Y, Lai Y, Chen X, Zhou DX, Zhao Y. 2020. Distribution pattern of histone marks

potentially determines their roles in transcription and RNA processing in rice. J Plant Physiol

249:153167.

Huang Y, Sicar S, Ramirez-Prado JS, Manza-Mianza D, Antunez-Sanchez J, Brik-Chaouche

R, Rodriguez-Granados NY, An J, Bergounioux C, Mahfouz MM et al. 2021.

Polycomb-dependent differential chromatin compartmentalization determines gene

coregulation in Arabidopsis. Genome Res 31 (7):1230-1244.

Iwamori N, Tominaga K, Sato T, Riehle K, Iwamori T, Ohkawa Y, Coarfa C, Ono E, Matzuk

MM. 2016. MRG15 is required for pre-mRNA splicing and spermatogenesis. Proc Natl Acad

Sci U S A 113 (37):E5408-5415.

Krzywinski M, Schein J, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA.

2009. Circos: an information aesthetic for comparative genomics. Genome Res 19

(9):1639-1645.

Landt SG, Marinov GK, Kundaje A, Kheradpour P, Pauli F, Batzoglou S, Bernstein BE,

Bickel P, Brown JB, Cayting P et al. 2012. ChIP-seq guidelines and practices of the ENCODE

and modENCODE consortia. Genome Res 22 (9):1813-1831.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nat Methods 9

(4):357-359.

Lauberth SM, Nakayama T, Wu X, Ferris AL, Tang Z, Hughes SH, Roeder RG. 2013.

H3K4me3 interactions with TAF3 regulate preinitiation complex assembly and selective gene

activation. Cell 152 (5):1021-1036.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R,

Genome Project Data Processing S. 2009. The Sequence Alignment/Map format and

SAMtools. Bioinformatics 25 (16):2078-2079.

Li Y, Dong XM, Jin F, Shen Z, Chao Q, Wang BC. 2017. Histone Acetylation Modifications

Affect Tissue-Dependent Expression of Poplar Homologs of C(4) Photosynthetic Enzyme

Genes. Front Plant Sci 8:950.

Li Y, Mukherjee I, Thum KE, Tanurdzic M, Katari MS, Obertello M, Edwards MB,

McCombie WR, Martienssen RA, Coruzzi GM. 2015. The histone methyltransferase SDG8

mediates the epigenetic modification of light and carbon responsive genes in plants. Genome

Biol 16 (1):79.

Liao Y, Smyth GK, Shi W. 2013. The Subread aligner: fast, accurate and scalable read

mapping by seed-and-vote. Nucleic Acids Res 41 (10):e108.

Liu B, Wei G, Shi J, Jin J, Shen T, Ni T, Shen WH, Yu Y, Dong A. 2016. SET DOMAIN

GROUP 708, a histone H3 lysine 36-specific methyltransferase, controls flowering time in

rice (Oryza sativa). New Phytol 210 (2):577-588.

Lopez-Delisle L, Rabbani L, Wolff J, Bhardwaj V, Backofen R, Gruning B, Ramirez F, Manke

T. 2021. pyGenomeTracks: reproducible plots for multivariate genomic datasets.

Bioinformatics 37 (3):422-423.

Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. Genome Biol 15 (12):550.

Lu Y, Tan F, Zhao Y, Zhou S, Chen X, Hu Y, Zhou DX. 2020. A

Chromodomain-Helicase-DNA-Binding Factor Functions in Chromatin Modification and

Gene Regulation. Plant Physiol 183 (3):1035-1046.

Margueron R, Justin N, Ohno K, Sharpe ML, Son J, Drury WJ, 3rd, Voigt P, Martin SR,

Taylor WR, De Marco V et al. 2009. Role of the polycomb protein EED in the propagation of

repressive histone marks. Nature 461 (7265):762-767.

Marzol E, Borassi C, Carignani Sardoy M, Ranocha P, Aptekmann AA, Bringas M,

Pennington J, Paez-Valencia J, Martinez Pacheco J, Rodriguez-Garcia DR et al. 2022. Class

III Peroxidases PRX01, PRX44, and PRX73 Control Root Hair Growth in Arabidopsis

thaliana. Int J Mol Sci 23 (10):5375.

McCormick RF, Truong SK, Sreedasyam A, Jenkins J, Shu S, Sims D, Kennedy M,

Amirebrahimi M, Weers BD, McKinley B et al. 2018. The Sorghum bicolor reference genome:

improved assembly, gene annotations, a transcriptome atlas, and signatures of genome

organization. Plant J 93 (2):338-354.

Offermann S, Danker T, Dreymuller D, Kalamajka R, Topsch S, Weyand K, Peterhansel C.

2006. Illumination is necessary and sufficient to induce histone acetylation independent of

transcriptional activity at the C4-specific phosphoenolpyruvate carboxylase promoter in

maize. Plant Physiol 141 (3):1078-1088.

Oksuz O, Narendra V, Lee CH, Descostes N, LeRoy G, Raviram R, Blumenberg L, Karch K,

Rocha PP, Garcia BA et al. 2018. Capturing the Onset of PRC2-Mediated Repressive Domain

Formation. Mol Cell 70 (6):1149-1162 e1145.

Quinlan AR, Hall IM. 2010. BEDTools: a flexible suite of utilities for comparing genomic

features. Bioinformatics 26 (6):841-842.

Ramirez F, Ryan DP, Gruning B, Bhardwaj V, Kilpert F, Richter AS, Heyne S, Dundar F,

Manke T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis.

Nucleic Acids Res 44 (W1):W160-165.

Ringel AE, Cieniewicz AM, Taverna SD, Wolberger C. 2015. Nucleosome competition

reveals processive acetylation by the SAGA HAT module. Proc Natl Acad Sci U S A 112

(40):E5461-5470.

Roudier F, Teixeira FK, Colot V. 2009. Chromatin indexing in Arabidopsis: an epigenomic

tale of tails and more. Trends Genet 25 (11):511-517.

Saluja M, Zhu F, Yu H, Walia H, Sattler SE. 2021. Loss of COMT activity reduces lateral root

formation and alters the response to water limitation in sorghum brown midrib (bmr) 12

mutant. New Phytol 229 (5):2780-2794.

Sani E, Herzyk P, Perrella G, Colot V, Amtmann A. 2013. Hyperosmotic priming of

Arabidopsis seedlings establishes a long-term somatic memory accompanied by specific

changes of the epigenome. Genome Biol 14 (6):R59.

Schmitges FW, Prusty AB, Faty M, Stutzer A, Lingaraju GM, Aiwazian J, Sack R, Hess D, Li

L, Zhou S et al. 2011. Histone methylation by PRC2 is inhibited by active chromatin marks.

Mol Cell 42 (3):330-341.

Schwartz YB, Kahn TG, Nix DA, Li XY, Bourgon R, Biggin M, Pirrotta V. 2006.

Genome-wide analysis of Polycomb targets in Drosophila melanogaster. Nat Genet 38

(6):700-705.

Sequeira-Mendes J, Araguez I, Peiro R, Mendez-Giraldez R, Zhang X, Jacobsen SE, Bastolla

U, Gutierrez C. 2014. The Functional Topography of the Arabidopsis Genome Is Organized in

a Reduced Number of Linear Motifs of Chromatin States. Plant Cell 26 (6):2351-2366.

Sorenson MR, Jha DK, Ucles SA, Flood DM, Strahl BD, Stevens SW, Kress TL. 2016.

Histone H3K36 methylation regulates pre-mRNA splicing in Saccharomyces cerevisiae. RNA

Biol 13 (4):412-426.

Suzuki S, Kato H, Suzuki Y, Chikashige Y, Hiraoka Y, Kimura H, Nagao K, Obuse C,

Takahata S, Murakami Y. 2016. Histone H3K36 trimethylation is essential for multiple

silencing mechanisms in fission yeast. Nucleic Acids Res 44 (9):4147-4162.

Tao Y, George-Jaeggli B, Bouteille-Pallas M, Tai S, Cruickshank A, Jordan D, Mace E. 2020.

Genetic Diversity of C(4) Photosynthesis Pathway Genes in Sorghum bicolor (L.). Genes

(Basel) 11 (7):806.

Taverna SD, Ilin S, Rogers RS, Tanny JC, Lavender H, Li H, Baker L, Boyle J, Blair LP,

Chait BT et al. 2006. Yng1 PHD finger binding to H3 trimethylated at K4 promotes NuA3

HAT activity at K14 of H3 and transcription at a subset of targeted ORFs. Mol Cell 24

(5):785-796.

Veluchamy A, Jegu T, Ariel F, Latrasse D, Mariappan KG, Kim SK, Crespi M, Hirt H,

Bergounioux C, Raynaud C et al. 2016. LHP1 Regulates H3K27me3 Spreading and Shapes

the Three-Dimensional Conformation of the Arabidopsis Genome. PLoS One 11

(7):e0158936.

Wang H, Fan Z, Shliaha PV, Miele M, Hendrickson RC, Jiang X, Helin K. 2023. H3K4me3

regulates RNA polymerase II promoter-proximal pause-release. Nature 615 (7951):339-348.

Weirauch MT, Yang A, Albu M, Cote AG, Montenegro-Montero A, Drewe P, Najafabadi HS,

Lambert SA, Mann I, Cook K et al. 2014. Determination and inference of eukaryotic

transcription factor sequence specificity. Cell 158 (6):1431-1443.

Widiez T, Symeonidi A, Luo C, Lam E, Lawton M, Rensing SA. 2014. The chromatin

landscape of the moss Physcomitrella patens and its dynamics during development and

drought stress. Plant J 79 (1):67-81.

Xi Y, Park SR, Kim DH, Kim ED, Sung S. 2020. Transcriptome and epigenome analyses of

vernalization in Arabidopsis thaliana. Plant J 103 (4):1490-1502.

Yang H, Howard M, Dean C. 2014. Antagonistic roles for H3K36me3 and H3K27me3 in the

cold-induced epigenetic switch at Arabidopsis FLC. Curr Biol 24 (15):1793-1797.

Yang T, Wang D, Tian G, Sun L, Yang M, Yin X, Xiao J, Sheng Y, Zhu D, He H et al. 2022.

Chromatin remodeling complexes regulate genome architecture in Arabidopsis. Plant Cell 34

(7):2638-2651.

Yano S, Ishiuchi T, Abe S, Namekawa SH, Huang G, Ogawa Y, Sasaki H. 2022. Histone

H3K36me2 and H3K36me3 form a chromatin platform essential for DNMT3A-dependent

DNA methylation in mouse oocytes. Nat Commun 13 (1):4440.

Yuan W, Xu M, Huang C, Liu N, Chen S, Zhu B. 2011. H3K36 methylation antagonizes

PRC2-mediated H3K27 methylation. J Biol Chem 286 (10):7983-7989.

Zhang X, Clarenz O, Cokus S, Bernatavichute YV, Pellegrini M, Goodrich J, Jacobsen SE.

2007. Whole-genome analysis of histone H3 lysine 27 trimethylation in Arabidopsis. PLoS

Biol 5 (5):e129.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM,

Brown M, Li W et al. 2008. Model-based analysis of ChIP-Seq (MACS). Genome Biol 9

(9):R137.

Zhao L, Xie L, Zhang Q, Ouyang W, Deng L, Guan P, Ma M, Li Y, Zhang Y, Xiao Q et al.

2020. Integrative analysis of reference epigenomes in 20 rice varieties. Nat Commun 11

(1):2658.

Zheng D, Wang L, Chen L, Pan X, Lin K, Fang Y, Wang XE, Zhang W. 2019.

Salt-Responsive Genes are Differentially Regulated at the Chromatin Levels Between

Seedlings and Roots in Rice. Plant Cell Physiol 60 (8):1790-1803.

Zilberman D, Coleman-Derr D, Ballinger T, Henikoff S. 2008. Histone H2A.Z and DNA

methylation are mutually antagonistic chromatin marks. Nature 456 (7218):125-129.

 Figure legends

 Figure 1 Distribution pattern of histone marks in sorghum.

 (a) Metaplots of 7 histone mark profiles of all genes in leaf and root before and after PEG

 treatment. **(b)** Heat maps of histone mark enrichment of intragenic region in leaf and root

 before and after PEG treatment. Black arrows indicate the TSS-enriched H3K27me3 present

 in leaf but not in root. Two-week-old sorghum seedlings were treated with 20% PEG6000 for

 6 hours. Then root and leaf of control (without PEG treatment) and PEG-treated plants were

 harvested for ChIP-seq. **(c)** Length distribution of H3K27me3 regions in leaf and root. **(d)**

 The number of genes covered by each H3K27me3 region in leaf and root. "MG1000"

 indicates that the adjacent H3K27me3 regions were joined when they were separated by less

 than 1 kb while "MG0" indicates that they were not jointed. **(e)** Genome browser screen shots

 of 4 H3K27me3 islands on Chromosomes 2, 5, 8. H3K27me3 islands were defined as the

 areas where a number of long H3K27me3 regions clustered.

 Figure 2 Analysis of overlapping enrichment of different histone marks in sorghum leaf.

 The enrichment of H3K9ac **(a)**, H3K27ac **(b)**, H3K4me3 **(c)**, H3K36me3 **(d)**, H2A.Z **(e)**,

 H3K4me2 **(f)**, and H3K27me3 **(g)** in intragenic region were clustered into 3 categories (C1,

 C2 and C3) respectively and were put on the far left. The other 6 histone mark enrichment and

 gene expression level were shown next to the clustered marks. **(h)** Genome browser screen

 shots showing part of the areas in 4 H3K27me3 islands where H3K36me3 and H3K27me3

 excluded from each other. **(i)** Genome browser screen shots showing some areas where

 H3K36me3 and H2A.Z excluded from each other.

 Figure 3 Correlation analysis of differential enrichment of histone marks and

 differential gene expression in leaf and root.

 (a) Metaplots of 7 histone mark profiles in differentially expressed genes (DEGs) between

 leaf and root. **(b)** Scatterplots of correlation between change of histone marks (log2FC-root vs

 leaf) and change of gene expression (log2FC-root vs leaf). Red dots represent the genes with

 both increased histone mark levels (log2FC > 0.75, p < 0.05) and increased expression levels

 (log2FC > 1, p < 0.05) and the genes with both decreased histone mark levels (log2FC < -0.75,

 p < 0.05) and decreased expression levels (log2FC < 1, p < 0.05) (Positive relationship). Blue

 dots represent the genes with increased histone mark levels and decreased expression levels

and the genes with decreased histone mark levels and increased expression levels (Negative

relationship). Gray dots represent the genes in which correlation between change of histone

marks and change of gene expression was not significant. The number of the co-related genes

was indicated. **(c)** The proportion of genes in which two histone marks changes between leaf

and root. "Former unique" means the genes in which the former one of the two compared

histone mark shown on the horizontal axis change but the latter one does not. "Latter unique"

means the genes in which the latter histone mark change but the former one does not.

"Common" means the genes in which both histone marks change. **(d)** Heat maps of

H3K27me3 and H3K36me3 enrichments in all the genes with changed H3K36me3 (left,

H3K36me3 increases in root; right, H3K36me3 increases in leaf). **(e)** Heat maps of H2A.Z

and H3K36me3 enrichments in all the genes with changed H3K36me3 (left, H3K36me3

increases in root; right, H3K36me3 increases in leaf).

Figure 4 Leaf-specific expression of photosynthesis genes is associated with dynamic

change of histone marks.

(a) KEGG analysis of the leaf-up-DEGs and root-up-DEGs with differential enrichment of

various histone marks in leaf and root. The pathways marked with pink background represent

leaf-up-DEGs enriched Photosynthesis-related pathways. The pathway marked with purple

background represents root-up-DEGs enriched Phenylpropanoid biosynthesis pathway. Heat

maps showing fold change (log2FC) of expression and histone marks of photosynthesis genes

(b) and C4 and non-C4 genes **(c)** in leaf compared to root. **(d)** Genome browser screen shots

of histone mark enrichment on C4 genes in leaf and root.

Figure 5 Root-specific expression of lignin biosynthesis and POD genes is associated with

dynamic change of histone marks.

(a) Schematic diagram of lignin synthesis. Heat maps showing fold change (log2FC) of

expression and histone marks of photosynthesis genes **(b)** and C4 and non-C4 genes **(c)** in

leaf compared to root. **(d)** Genome browser screen shots of histone mark enrichment on ten

POD genes with topmost expression change in leaf and root.

Figure 6 Correlation analysis of changes of histone marks and gene expression in

response to PEG treatment in leaf and root.

Metaplots of 7 histone mark profiles in differentially expressed genes (DEGs) in response to

PEG treatment in leaf **(a)** and root **(b)**. Scatterplots of correlation between change of histone

marks (log2FC) and change of gene expression (log2FC) in response to PEG treatment in leaf

(c) and root **(d)**. Red dots represent the genes with both increased histone mark levels

(log2FC > 0.75, $p < 0.05$) and increased expression levels (log2FC > 1, $p < 0.05$) and the

genes with both decreased histone mark levels (log2FC < -0.75, $p < 0.05$) and decreased

expression levels (log2FC < 1, $p < 0.05$) (Positive relationship). Blue dots represent the genes

with increased histone mark levels and decreased expression levels and the genes with

decreased histone mark levels and increased expression levels (Negative relationship). Gray

dots represent the genes in which correlation between change of histone marks and change of

gene expression was not significant. The number of the co-related genes was indicated.

Figure 7 Dynamic change of histone marks in the genes that were both differentially

expressed between leaf and root and in response to PEG treatment.

(a) Venn diagram of overlapping DEGs both between leaf and root and in response to PEG

treatment. Four groups (I, II, III, IV) of the overlapping DEGs, also designated as the

common DEGs, were identified. Root-up-DEGs: genes with increased expression in root

compared to leaf; Leaf-up-DEGs: genes with increased expression in leaf compared to root;

PEG-up-DEGs(leaf): genes with increased expression in response to PEG in leaf;

PEG-down-DEGs(leaf): genes with decreased expression in response to PEG in leaf;

PEG-up-DEGs(root): genes with increased expression in response to PEG in root;

PEG-down-DEGs(root): genes with decreased expression in response to PEG in root. **(b)** The

overlapping analysis of histone mark changes in four groups of the common DEGs identified

in **(a)** both between leaf and root and in response to PEG treatment. "Unique" represents

histone marks changed only in this condition but not the other while "common" means

histone marks changed in both conditions. The number of "unique" or "common" genes was

marked on the column. The total number of each group of the common DEGs was indicated.

Heat maps showing fold change (log2FC) of expression and histone marks of photosynthesis

genes **(c)** and C4 and non-C4 genes **(d)** before and after PEG treatment in leaf, and POD

genes **(e)** before and after PEG treatment in root.

Figure 8 Cis-elements analysis of genes with histone marks changing between leaf and

root (a) and in response to PEG treatment (b).

CL: Leaf control; PL: Leaf PEG treatment; CR: Root control; PR: Leaf PRG treatment. The

percentage (%) of the genes possessing the cis-element relative to the total genes with the

histone mark changing was marked on the column.

Figure S1. The quality of ChIP-seq and RNA-seq data obtained in this study.

Figure S2. The overview of genome-wide enrichment of histone marks in sorghum.

Figure S3. Long H3K27me3 regions in Arabidopsis and rice.

Figure S4. Analysis of overlapping enrichment of different histone marks in sorghum

root.

Figure S5. The relationship between H3K36me3, H3K27me3 and H2A.Z in Arabidopsis

and rice.

Figure S6. Long H3K27me3 peaks change between leaf and root

Figure S7 Gene Ontology (GO) analysis of the leaf-up-DEGs and root-up-DEGs with

differential enrichment of various histone marks in leaf and root.

Figure S8. The role of histone marks in the induction of PEG-responsive genes.

Figure S9. Genome browser screen shots of histone mark enrichment in some common

DEGs (identified in Figure 6) with 7 histone marks changing between leaf and root but

not in response to PEG treatment.

Figure S10 Heat maps showing fold change (log2FC) of expression and histone marks

of lignin biosynthesis genes

Table S1 Total reads and mapping rate of ChIP-seq data

Table S2 Changes of expression and histone marks of stress genes

(a)

(b)