Transcriptome Analysis Reveals Organ-Specific Effects of 2-Deoxyglucose Treatment in Healthy Mice

Ann E. Wells[1], John J. Wilson[1], Sarah E. Heuer[1,2], John D. Sears[1], Jian Wei[1], Raghav Pandey[1], Mauro W. Costa[1], Derry C. Roopenian[1], Chih-Hao Chang[1,2,3*], Gregory W. Carter[1,2,3*]

[1]The Jackson Laboratory, Bar Harbor, ME 04469, USA.

[2]Tufts University Graduate School of Biomedical Sciences, Boston, MA 02111, USA.

[3]Graduate School of Biomedical Sciences and Engineering, University of Maine, Orono, ME 04469, USA.

*Corresponding author

Abstract

Objective: Glycolytic inhibition via 2-deoxy-D-glucose (2DG) has potential therapeutic benefits

for a range of diseases, including cancer, epilepsy, systemic lupus erythematosus (SLE), and

rheumatoid arthritis (RA), and COVID-19, but the systemic effects of 2DG on gene function

across different tissues are unclear.

Methods: This study analyzed the transcriptional profiles of nine tissues from C57BL/6J mice

treated with 2DG to understand how it modulates pathways systemically. Principal component

analysis (PCA), weighted gene co-network analysis (WGCNA), analysis of variance, and

pathway analysis were all performed to identify modules altered by 2DG treatment.

Results: PCA revealed that samples clustered predominantly by tissue, suggesting that 2DG

affects each tissue uniquely. Unsupervised clustering and WGCNA revealed six distinct tissue-

specific modules significantly affected by 2DG, each with unique key pathways and genes. 2DG

predominantly affected mitochondrial metabolism in the heart, while in the small intestine, it

affected immunological pathways.

Conclusions: These findings suggest that 2DG has a systemic impact that varies across

organs, potentially affecting multiple pathways and functions. The study provides insights into

the potential therapeutic benefits of 2DG across different diseases and highlights the

importance of understanding its systemic effects for future research and clinical applications.

Abbreviations

2DG - 2-deoxy-D-glucose, RA - rheumatoid arthritis, SARS-CoV-2 - severe acute respiratory

syndrome coronavirus-2, SLE - systemic lupus erythematosus, ER - endoplasmic reticulum,

WGCNA - weighted gene co-network analysis, ANOVA - analysis of variance, PCA - Principal

component analysis, GSVA – Gene Set Variation Analysis, BCAA - branched-chain amino

acids, NAD+ - nicotinamide adenine dinucleotide

Keywords: 2-deoxy-D-glucose, C57BL/6J, metabolism, multi-tissue, transcriptomics

1. Introduction

2-deoxy-D-glucose (2DG) is a glucose analog that has garnered considerable interest in recent years as a potential therapeutic agent for various diseases characterized by abnormal glycolysis, including cancer[1-5], epilepsy[6], systemic lupus erythematosus (SLE)[7], rheumatoid arthritis (RA)[8], and COVID-19 [9-12]. 2DG is taken up into cells by glucose transporters and is converted by hexokinase into 2DG-6-phosphate which cannot be further broken down to yield energy, resulting in a reduction in the rate of glycolysis[13]. Similarly, by competing with mannose, 2DG disrupts the early steps of N-linked glycosylation, ultimately resulting in the misfolding of proteins and the onset of endoplasmic reticulum (ER) stress[14]. Despite an increasing understanding of these cellular mechanisms, there remains a critical need to delineate the specific mechanisms through which 2DG could be used to treat a range of diseases and the organs that would be most affected in this context.

Studies have shown that 2DG can modulate progression or attenuation in multiple disease paradigm. For instance, in anti-tumor effects, 2DG may act through energy restriction to prevent tumor growth while maintaining bodyweight, glucose levels, and immunity[2, 4, 14, 15]. Furthermore, 2DG has been shown to reduce ATP due to its glycolytic properties and protein synthesis through the AMPK/mTORC1 pathway, reducing translation and promoting autophagy[16]. In seizure models, 2DG has been shown to reduce epileptic seizures both acutely and chronically[6]. Additionally, in mice with traumatic brain injuries, cortical slices showed reduced excitatory neurons and prevented epileptiform activity when treated immediately with 2DG[6] . 2DG has also been shown to reduce amyloid precursor protein and amyloid-beta oligomers in an Alzheimer's mouse model[17], delaying the progression of these critical hallmarks of Alzheimer's disease. When treating autoimmune diseases, 2DG has attenuated symptoms of SLE, RA, and multiple sclerosis in mice[7, 18]. Additionally, 2DG has been shown to significantly extend the lifespan of SLE-prone mice compared to untreated

controls[7]. In these and several other examples, 2DG has shown early promise in both disease

prevention and attenuation.

The use of 2DG in cancer therapy has been limited and its therapeutic efficacy has been

inconsistent, but it has been shown to be well tolerated with no significant safety concerns

raised[5, 13]. Moreover, glycolytic inhibition has been shown to reduce viral replication[11, 19],

and 2DG is currently being investigated for its potential to treat severe acute respiratory

syndrome coronavirus-2 (SARS-CoV-2)[9-12]. In clinical trials, patients with moderate to severe

SARS-CoV-2 infection who were administered oral 2DG showed an improvement in their

symptoms and were taken off oxygen supplementation significantly earlier compared to patients

treated with standard of care therapies[12]. Although 2DG has shown the potential to ameliorate

multiple diseases, a better understanding of 2DG's systemic effects help to identify potential

targets for clinical trials in metabolic disorders.

Much of what is known concerning the effects of 2DG is based on studies of cell lines, selected

tissues, whole animal physiology, and disease. The current study, instead, sought to understand

the effects of 2DG administered systemically on major organs. Our goal was to understand how

2DG alters baseline metabolism of a genetically uniform population of young, healthy C57BL/6J

mice unencumbered by advanced age, disease, or manipulation. Our approach was to analyze

the transcriptomes of nine organs (heart, kidney, hippocampus, hypothalamus, prefrontal cortex,

skeletal muscle, small intestine, and spleen) from 2DG- and vehicle-treated C57BL/6J mice to

assess systemic responses to 2DG treatment (**Fig. 1**). Using a robust statistical filtering

strategy, which combined weighted gene co-network analysis (WGCNA), analysis of variance

(ANOVA), and correlation, our results show that 2DG altered metabolism, immunity, and

transcription in heart, small intestine, hypothalamus, prefrontal cortex, skeletal muscle, and liver

through a unique set of genes in each tissue. The small intestine presented an

immunomodulatory signature, while both heart and hypothalamus showed reduced

mitochondrial metabolism. Few pathways responded to 2DG in skeletal muscle and liver; and all

were down-regulated regardless of biological function. Pathways involved in RNA transcription

and endoplasmic reticulum (ER) stress in the prefrontal cortex were over and under-expressed

in response to 2DG, respectively. Our results provide a comprehensive understanding of the

systemic impact of 2DG on glycolysis and N-linked glycosylation and highlight the organ-specific

effects through which 2DG acts. Overall, our study lays the foundation for further research on

the therapeutic potential of targeting these metabolic pathways.

2. Materials and Methods

2.1 Mice, Treatment, and Tissue Isolation

C57BL/6J male mice (Jackson Laboratory, #000664), used in this study were bred and housed

at The Jackson Laboratory (Bar Harbor, Maine). Twenty mice at 8 and 20 mice at 11 weeks of

age were switched from a 6% (g/g) fat JL Mouse Breeder/Auto 6F (LabDiet® 5K52) diet to a

10% fat JL Mouse Breeder/Auto (LabDiet® 5K20) diet when moved into their mouse room,

provided with water *ad libitum,* and were housed on a 14-hour light, 10-hour dark cycle in a

specific pathogen-free room. 2-Deoxy-D-glucose (2DG, Thermo Fisher Scientific, category

number AC111980250, 99% purity) dissolved in drinking water at a concentration of 6 g/L was

provided to mice *ad libitum* for either 4-day or 28-day treatment times. As the average mouse

consumed ~ 3.5 ml of water per day, this equates to a dosage of 800 mg/kg/d. The rationale for

selecting this dosing regime is that it has proven to be therapeutically effective and safe in

treatment of autoimmune disease in several mouse models[18, 20]. After allometric scaling, this

dosage approximates an accepted human dosage of ~65mg/kg/d. The 4-day treatment cohort

was provided 2DG-water starting at 12 weeks of age, while the 28-day treatment cohort was

provided 2DG-water starting at 9 weeks of age (**Fig. 1**). Control mice received regular drinking

water with no additives. All mice, including age-matched control mice, were sacrificed at 13

weeks of age via cervical dislocation. Brains were removed and dissected to isolate the

hippocampus, hypothalamus, and pre-frontal cortex. Spleen, liver, heart, kidney, skeletal muscle

from left hind-leg and intestinal ileum were also harvested. Heart and intestine were flushed with

saline post-dissection to remove blood and feces, respectively. All harvested organs were

stored in RNAlater for subsequent RNA-seq analysis. The Jackson Laboratory Institutional

Animal Care and Use Committee (IACUC) approved all procedures.

2.2 RNA Sequencing

RNA-sequencing (RNA-seq) was performed by Omega Bioservices. Bulk RNA of the 9 excised

tissues (heart, hippocampus, hypothalamus, kidney, liver, prefrontal cortex, skeletal muscle,

small intestine, and spleen) (4 samples per treatment and time combination) was isolated with

the QIAGEN miRNeasy mini extraction kit (QIAGEN) and cDNA was synthesized with the High-

Capacity cDNA Reverse Transcription Kit (Applied Biosystems). RNA quality was assessed with

a Bioanalyzer 2100 (Agilent Technologies). Poly(A)-selected RNA-seq libraries were generated

using the Illumina TruSeq RNA Sample preparation kit v2. RNA-seq was performed in a 150-bp

paired-end format with a minimum of 40 million reads per sample on the Illumina HiSeq platform

according to the manufacturer's instructions. RNA-seq reads were filtered and trimmed for

quality scores >30 using a custom python script. The filtered reads were aligned to *Mus*

musculus GRCm38 using RSEM (v1.2.12) (58) with Bowtie2 (v2.2.0)[21] (command: rsem-

calculate-expression -p 12 --phred33-quals --seed-length 25 --forward-prob 0 --time --output-

genome-bam -- bowtie2). RSEM calculates expected counts and transcript per million (TPM).

The expected counts from RSEM were used in the Bioconductor edgeR 3.20.9 package[22-24]

to determine differentially expressed genes.

2.3 Statistical Analysis

All statistical analysis was performed in the language R (3.6.0/4.1.0)[25] equipped with RStudio.

The analytical strategy is summarized in Figure 1 and explained in subsequent method

sections.

2.3.1 Principal Component Analysis

Principal component analysis (PCA) was performed to identify whether samples clustered

based on treatment, time, or treatment-by-time interaction. The PCA algorithm was

implemented using the function prcomp() in the package stats[25]. Hotelling's T^2 ellipses were

used to identify the 95% confidence intervals for each cluster.

2.3.2 Weight Gene Co-network Analysis

To identify potential transcript networks Weighted Gene Co-Expression Network Analysis

(WGCNA) was performed using the `WGCNA' R package[26, 27]. Soft-thresholding power was

chosen so that the scale-free topology correlation hits as close to 0.9 as possible. The soft

thresholding powers chosen for heart, hippocampus, hypothalamus, kidney, liver, prefrontal

cortex, skeletal muscle, small intestine, and spleen were 5, 13, 9, 3, 3, 15, 13, and 20,

respectively. The adjacency matrix was created using type = "unsigned". Dynamic tree cutting

was used to cluster metabolites and generate modules with a minimum of 15 genes in each

module. Networks were identified using all genes quantified by RNA-seq without replacement.

Module colors were assigned arbitrarily and have no bearing on functional annotation.

2.3.3 Functional Pathway Analysis

Overrepresentation pathway analysis was performed using gProfiler2[28]. KEGG and reactome

databases were used as the references to compare genes in each module identified in WGCNA

analysis, using Ensembl 104 and Ensembl Genomes 51 databases. To determine statistical

significance of each pathway identified, Fisher's exact test was used. P-values were adjusted

using the Benjamini-Hochberg procedure to reduce type 1 error (FDR < 0.05).

2.3.4 Analysis of Variance using Aligned Rank Transformation

Since the data was not normally distributed aligned rank transformation was performed using

the ARTool package[29, 30] to identify significance for using the linear model:

$$\text{Module eigengene} = \text{intercept} + \text{treat} + \text{time} + \text{treat*time}$$

to determine the effects of treatment, time, or treatment-by-time interaction on each module

eigengene.

Since the change in diet composition and time points overlap exactly, neither were individually

assessed in this study but they were evaluated in the ANOVA models—at the module, gene,

and Gene Set Variation Analysis (GSVA) level—for interaction to identify their combined

influence on treatment. Only modules, GSVA, and genes significant for treatment but not time or

treatment dependent on time were discussed.

2.3.5 Mediation Analysis

The R package mediation[31] was used to perform mediation analysis. Eigengenes for

overrepresented pathways were created using GSVA[32] (explained in section 2.6) and

assessed across tissues to determine if pathways were acting as mediators on the response to

treatment using two models; the mediator model:

$$\text{Pathway 1} = \text{intercept} + \text{treatment} + \text{time} + \text{treatment*time}$$

and the outcome model:

$$\text{Pathway 2} = \text{intercept} + \text{pathway 1} + \text{treatment} + \text{time} + \text{treatment*time}$$

The models were then entered into the function mediate() and simulated 1000 times.

Confidence intervals and p-values were created using the Quasi-Bayesian Monte Carlo method.

2.4 Modules Identified Using Three Filtering Criteria

To identify modules related to treatment for each tissue each module had to meet three criteria:

1.) module was significantly correlated with treatment, 2.) module identified significantly

overrepresented pathways, and 3.) module was uniquely significant for treatment only,

according to ANOVA. Correlation was used to directly describe the relationship between each

module and treatment. ANOVA was used to assess the difference between main effects and

their interactions. This filtering strategy was chosen to identify modules that had the most robust

response to 2DG treatment. Modules that fit these criteria were further assessed for genes that

may be central to the networks identified.

2.5 Gene Set Variation Analysis

GSVA was performed, using the GSVA package[32], to identify enrichment of significantly

overrepresented pathways for each tissue module (see data resource for comprehensive

description of all modules, section 2.6). To identify enriched pathways a reference set of

pathways is needed. To create the reference set of pathways, all ensembl genes listed for *Mus*

musculus were compared against the KEGG and reactome databases in gProfiler2 using the

gost() function and the ensembl 104 and ensembl genomes 51 databases. This identified 2,030

pathways, which were used as the pathway gene set. The set of genes in each module was

then assessed against the pathway gene set to identify significantly overrepresented pathways.

To calculate overrepresentation, we used the Fisher's exact test and corrected for multiple

comparisons using the Benjamini-Hochberg procedure (FDR < 0.05). To summarize the gene

expression in each pathway, the method plage was used, which uses the coefficients of the first

right-singular vector from the singular value decomposition. Genes within these overrepresented

pathways were then summarized into a singular eigengene representing their expression for

each sample. The pathway eigengenes for each significantly overrepresented pathway were

then tested using ANOVA:

pathway eigengene = intercept + treat + time + treat*time

to identify resident pathways significantly altered by treatment. This identified which significantly

overrepresented pathways were also significantly enriched.

2.6 Data Resource Development for Reproducible and Transparent Data

The data resource was developed using R 4.1.0, rmarkdown, and bookdown[33]. Static html

files were output using knitr and collated using a yaml file that controls the navigation bar on the

website and the general theme and layout, including footer information, and links in the

navigation bar to the shiny app, git repository, figshare, and contact information. The theme

used for the website was cerulean and type was set to inverse to change the color to dark blue.

The table of contents for each page was set to float. The website is publicly hosted at:

https://storage.googleapis.com/bl6_2dg_rnaseq/index.html.

3. Results

3.1 Molecular changes are induced in a tissue-specific manner by 2-DG treatment

The impact of 2DG on transcriptomic profiles of nine different tissues was assessed to identify

genes that were differentially expressed for treatment only. The number of differentially

expressed genes varied across the tissues, with a minimum of 230 genes in spleen and a

maximum of 1152 genes in hippocampus. Interestingly, only three tissues (muscle (1), small

intestine (119), and prefrontal cortex (3)) exhibited overrepresented pathways, indicating that

the effect of 2DG on biological pathways may be more significant in these tissues than in the

other six tissues examined. Assessing 2DG's effect at a granular level, however, do not account

for potential gene interactions. These findings suggest that assessing the correlation structure of

genes within each tissue may provide a better biological understanding of 2DG's impact at the

tissue level.

To identify the most robust signature in each tissue and minimize noise, a novel filtering strategy

was devised (**Fig. 1**). Principal Component Analysis showed that samples clustered by tissue

based on their transcriptomic profiles (**Fig. 2**). We used WGCNA to group genes into modules

that share highly correlated expression patterns. Across tissues, the number of modules into

which genes were clustered ranged from 25 to 46 and contained anywhere from 22 to 10,127

genes within each module. Identifying modules that fit the three criteria in the filtering strategy

(significant for treatment, through correlation and ANOVA, and contained significantly

overrepresented pathways ($p < 0.05$, FDR < 0.05)), we narrowed our focus to six tissues —

heart, hypothalamus, liver, skeletal muscle, prefrontal cortex, and small intestine — and

identified one module for each of these tissues. This approach allowed us to elucidate the most

robust changes in gene expression in response to 2DG treatment in each tissue and highlight

the biological pathways that are most affected. More information about the analysis, including

the number of modules and genes per tissue, can be found in the Supplementary Information or

on our website (https://storage.googleapis.com/bl6_2dg_rnaseq/index.html).

3.2 2DG reduced expression of metabolic pathways in the heart

The expression of genes in heart muscle was analyzed and found to cluster into 25 distinct

modules (**Supplementary Fig. 1A**), with three of them showing significant correlation with 2DG

treatment (**Supplementary Fig. 1B**). Further analysis using ANOVA revealed that only the

darkgreen module was uniquely significant for the treatment ($p < 0.001$). This module contained

138 genes predominantly related to metabolism (**Supplementary Fig. 1C**), which summary

eigengenes revealed were decreased in expression in 2DG-treated mice compared to control

mice (**Supplementary Fig. 1D, E**). The darkgreen module was found to contain ten pathways

related to mitochondrial metabolism and the breakdown of branched-chain amino acids (BCAA)

(**Table 1**), which were significantly overrepresented ($p < 0.05$). Comparison of genes within

each pathway using a Jaccard similarity index revealed that each pathway consisted of a unique

set of genes (**Supplementary Fig. 2**). To assess pathway enrichment, gene set variation

analysis (GSVA) was performed. Of the ten significantly overrepresented pathways, seven

showed significant reduction in eigengene expression for the treatment group only ($p < 0.05$,

Supplementary Table 1), referred to as high confidence pathways from this point forward.

BCAA degradation was found to be inversely correlated with changes in mitochondrial

metabolism, consistent with previous findings of mitochondrial dysfunction leading to BCAA

accumulation[34]. Failure in both mitochondrial metabolism and BCAA degradation have been

linked to increase cardiovascular disease[35], and reduced BCAA degradation has been

identified as a signature in heart failure[36]. The inverse relationship of BCAA degradation and

mitochondrial metabolism identified in this study suggests that 2DG's differential effects on heart

metabolism may provide protection against heart failure.

278
279 **Table 1: Metabolic pathways are overrepresented in the darkgreen heart module**

Pathways	GSVA Significance for Treatment (ANOVA p-value)*	GSVA Expression Direction (2DG compared to Control
Metabolism	**0.0001**	Down
Mitochondrial translation	0.133	Down
Mitochondrial translation elongation	0.133	Down
Mitochondrial translation termination	0.133	Down
Propanoate metabolism	**0.033**	Down

Pyruvate metabolism	**0.033**	Down
Pyruvate metabolism and Citric Acid (TCA) cycle	**0.004**	Down
Regulation of pyruvate dehydrogenase (PDH) complex	**0.001**	Down
The citric acid (TCA) cycle and respiratory electron transport	**0.001**	Down
Valine, leucine, and isoleucine degradation	**0.002**	Up

280   * Significant p-values in bold
281

3.3 2DG suppressed expression of immunological pathways in the small intestine

The genes expressed in the small intestine were clustered into 46 modules (**Fig. 3A**). Three of

these modules were found to be significantly correlated for treatment (p < 0.05) and identified

significantly overrepresented pathways (p < 0.05), with the green4 module being uniquely

significant for treatment, according to ANOVA (p = 0.04, **Fig. 3B**). This module contained 467

genes (**Fig. 3E**) predominantly related to the immune system and summary eigengenes

revealed decreased expression in mice treated with 2DG compared to control mice (**Fig. 3C, D**).

Overrepresentation analysis identified 69 significant pathways (p < 0.05, **Supplementary Table**

2), containing a unique set of genes in each pathway according to the Jaccard similarity index

(**Supplementary Fig. 3**). GSVA results showed that 61 out of the 69 pathways were

significantly enriched for treatment only (p < 0.05, **Supplementary Table 2**) and therefore

indicated high confidence. Furthermore, when pathways were functionally annotated it was

noted that 39 of the 61 high confidence pathways were related to immunological function and

immunology related diseases. Thirteen pathways showed decreased gene expression in mice

treated with 2DG, while 26 pathways showed increased expression. The up-regulated pathways

were primarily involved in innate immunity and the down-regulated pathways were related to

adaptive immunity (**Supplementary Table 2**). CIBERSORT analysis was used to assess

immune cell types in the green4 module and found that nine cell types were determined to be

significantly different between the green4 module and all the genes identified within the small

intestine (referred to as all genes). Four of these cell types (eosinophils, macrophages, memory

B cells, and Th1 cells) were significantly increased in the green4 module compared to all genes

(P < 0.05). The expression of genes related to eosinophils and macrophages (innate immune

cells) increased with 2DG treatment, while genes related to memory B cells and Th1 cells

(adaptive immune cells) decreased, but not significantly. The expression of genes related to

immature dendritic cells was significantly different between the green4 module and all genes but

in a treatment-dependent manner, with decreased expression in the green4 module and mice

treated with 2DG compared to all genes and control mice (**Supplementary Table 3**).

Collectively, these results indicate that treatment with 2DG leads to changes in the gene

expression profile of the small intestine's immune system characterized by a reduction in

adaptive immunity, and potentially compensation for corresponding by an enrichment in innate

immune cells.

3.4 2DG reduced expression of nicotinate metabolism in the liver

A total of thirty-one modules were identified in liver (**Supplementary Fig. 4A**), but only the

darkolivegreen module, consisting of 280 genes that represented diverse functions met all three

filtering criteria (**Supplementary Fig. 4B, C**). The summary eigengenes revealed increased

expression in 2DG-treated mice compared to the control mice (**Supplementary Fig. 4D, E**). The

only significant pathway found was nicotinate metabolism, which was deemed to be of high

confidence through GSVA analysis. This pathway was overrepresented (p < 0.05,

Supplementary Table 4) and was significantly reduced in mice treated with 2DG compared to

the control group (p = 0.008). Nicotinate metabolism is important for maintaining levels of

nicotinamide adenine dinucleotide (NAD+), an important coenzyme necessary for energy

metabolism[37], which is primarily produced in the liver[38]. These findings suggest that 2DG

treatment may have an impact on NAD+ levels in the liver and thus on ATP production.

3.5 2DG reduced expression of protein metabolism in skeletal muscle

A comprehensive analysis of the skeletal muscle revealed the existence of 40 modules (**Supplementary Fig. 5A**), of which 32 were found to have significantly overrepresented pathways and three were significantly correlated with the treatment ($p < 0.05$, **Supplementary Fig. 5B**). Further analysis revealed that among these three modules, only the salmon2 module was determined to be unique and significant for treatment through ANOVA analysis ($p = 0.017$). This module contained 153 genes (**Supplementary Fig. 5C**) and showed an increase in summary eigengenes in mice treated with 2DG compared to the control mice (**Supplementary Fig. 5D, E**). Like the findings in liver, genes in this module were found to be associated with a myriad of functions, but only five pathways related to protein metabolism were significantly overrepresented ($p < 0.05$, **Supplementary Table 5**) and showed a reduction in expression after 2DG treatment. Further analysis using the Jaccard similarity index and GSVA revealed that the genes involved in each pathway were unique (**Supplementary Fig. 6**), and two of the five pathways were deemed uniquely significant for treatment and therefore high confidence ($p < 0.05$). As N-linked glycosylation is crucial for protein metabolism[39], the reduction in expression may lead to up-regulation of the unfolded protein response and down-regulation of protein synthesis. Overall, these data suggest that the treatment with 2DG has a significant impact on the expression of genes related to protein metabolism in the skeletal muscle.

3.6 2DG increased expression of fatty acid oxidation in the hypothalamus

The genes expressed in the hypothalamus grouped into 25 modules (**Supplementary Fig. 7A**). Four of these modules were found to be correlated with treatment (**Supplementary Fig. 7B**); however, only the bisque4 module was uniquely significant for treatment ($p = 0.008$). This module contained 289 genes predominantly related to metabolism and cell cycle (**Supplementary Fig. 7C**). The summary eigengenes of this module were decreased in mice

treated with 2DG compared to the control mice (**Supplementary Fig. 7D, E**). Nine pathways

were significantly overrepresented ($p < 0.05$), and the Jaccard similarity index showed that the

genes identified in each pathway were unique (**Supplementary Fig. 8**). Functional annotation

revealed that these nine pathways were related to fatty acid oxidation and chromosome health.

Two of the overrepresented pathways (mitochondrial fatty acid beta-oxidation and mitochondrial

fatty acid beta-oxidation of saturated fatty acids) were uniquely enriched for treatment based on

GSVA analysis, supporting their high confidence ($p < 0.05$, **Supplementary Table 6**). Both

pathways were related to fatty acid oxidation, which often increases as a result of reduced

glycolysis[40]. Although the other seven pathways were not significantly enriched using GSVA,

the eigengenes in the G2/M DNA damage checkpoint pathway were negatively correlated with

each of the other eight pathways (**Supplementary Table 6**). The G2/M DNA checkpoint showed

significantly reduced expression in mice treated with 2DG ($p = 0.02$), but also for time ($p =$

0.036, **Supplementary Table 6**). These results indicate that the hypothalamus increase fatty

acid oxidation in response to the blockade of glycolysis and protects DNA through upregulation

of pathways related to chromosome maintenance, potentially in response to the increase in

unfolded protein response[41]. Overall, these findings suggests that the hypothalamus may use

lipids as an alternative fuel source in response to 2DG.

3.7 Enhancement of RNA transcription pathways by 2DG in the prefrontal cortex

Thirty-six gene modules were identified in the prefrontal cortex (**Supplementary Fig. 9A**). Of

these, five modules showed a correlation with treatment ($p < 0.05$, **Supplementary Fig. 9B**)

and 26 modules contained significantly overrepresented pathways ($p < 0.05$), with only the

bisque2 module being uniquely significant for treatment, according to ANOVA ($p = 0.014$). The

bisque2 module consisted of 718 genes that represented many biological functions

(**Supplementary Fig. 9C**). The summary eigengenes were found to be elevated in 2DG-treated

mice compared to control mice (**Supplementary Fig. 9D, E**). Functional analysis revealed that

14 pathways were overrepresented in the bisque2 module (p < 0.05), with seven related to RNA

transcription and seven to protein metabolism. According to the Jaccard similarity index, the

genes in each pathway were found to be unique (**Supplementary Fig. 10**). Among these

pathways, 12 were considered high confidence pathways, as they showed significant

enrichment for treatment using GSVA (p < 0.05, **Supplementary Table 7**). 2DG increased the

overall expression of five high confidence pathways related to RNA transcription compared to

control mice. Conversely, the seven high confidence pathways related to protein metabolism

showed decreased expression in 2DG-treated mice compared to control mice. Although 2DG

decreased expression of the nucleotide excision repair pathways, it was negatively correlated

with other pathways that also showed decreased expression in response to 2DG. All pathways

related to RNA transcription were negatively correlated with pathways involved in protein

metabolism. These results indicate that in the prefrontal cortex protein metabolism and RNA

transcription both respond to 2DG. Their differential response to 2DG may protect the prefrontal

cortex from the accumulation of unfolded proteins caused by the disruption of N-linked

glycosylation.

3.8 Tissue-specific responses to 2DG treatment revealed through correlation analysis of

gene modules

Tissues act synergistically to maintain the health of the body, but each tissue expresses genes

at varying levels. In this study, the expression of the same genes was analyzed across tissues;

however, the comparison of the genes contained in each selected tissue module revealed

minimal gene overlap (**Fig. 4**). The highest number of genes shared across tissue modules,

although not significant (p = 0.117), was 18 between skeletal muscle and prefrontal cortex,

while the number of genes identified in each module ranged from 113 to 659. No gene was

shared by all tissues. Moreover, we also compared overrepresented pathways across tissues

and identified only one shared pathway between small intestine and prefrontal cortex

(**Supplementary Fig. 11**). These results indicate that the tissue-specific effects of 2DG are

generally unique at both the gene and pathway levels.

We also identified distinct, organ-specific responses to 2DG that can be interpreted as a

network of co-occurring transcriptional changes across the organism (**Fig. 5**). To investigate the

relationships between tissue modules, we performed correlation analysis, resulting in five

significant pairwise relationships out of 21 ($p < 0.05$, **Fig. 5**). To gain a deeper understanding of

these relationships, we used GSVA eigengenes to examine the relationship of each pathway

across each tissue module. Positive correlations were found between small intestine and heart

tissues ($r = .51$, $p = 0.08$). Further assessment at the pathway level revealed a complex

relationship between adaptive and innate immunity in the small intestine and BCAA degradation

in the heart (**Fig. 5, Supplementary Table 8**). The pathways predominantly related to adaptive

immunity were inversely correlated with BCAA degradation, while those predominantly related

to innate immunity were positively correlated. Although lymphocytes do not remain in the heart

for any length of time, BCAA are necessary for lymphocyte development and clonal

expansion[42-45]. To assess any causal relationship between BCAA degradation and pathways

in the small intestine we performed causal mediation analysis. BCAA degradation did not

mediate any pathway's response to 2DG in the small intestine. These results indicate that while

there is an association between adaptive and innate immunity in the small intestine and BCAA

degradation in heart there is not causing the response to 2DG in the small intestine.

Tissue modules of the prefrontal cortex and hypothalamus were inversely related ($r = -0.48$, $p =$

0.09, **Fig. 6**). The incorporation of 2DG into the oligosaccharide chain inhibits N-linked

glycosylation and can lead to the formation of unfolded glycoproteins and induce ER stress[19].

In response to ER stress, the cell cycle is arrested at the G2/M DNA damage checkpoint[46].

Our findings showed a positive correlation between the G2/M DNA checkpoint pathway in the

hypothalamus and pathways related to ER stress in the prefrontal cortex (**Fig. 7,**

Supplementary Table 9). To assess the potential causal relationship between the

hypothalamus and prefrontal cortex, we conducted a causal mediation analysis and observed

that the G2/M DNA damage checkpoint pathway had a direct effect across all pathways in the

prefrontal cortex ($p < 0.05$) but no causal mediated effects. This indicates that the activation of

the G2/M DNA damage checkpoint pathway may have been a secondary response to the ER

stress induced by 2DG treatment, rather than the primary cause of the differential response to

2DG. Overall, these results suggest that both the hypothalamus and prefrontal cortex respond

to the disruption of N-linked glycosylation by modulating the expression of pathways to

counteract ER stress and the accumulation of unfolded proteins.

3.9 Data Resource

While we highlighted results from six tissues that stood out based on our filtering strategy, the

data resource that accompanies this paper contains WGCNA and other analyses for each of the

nine tissues, all the tissues combined, the brain tissues combined, and spleen, skeletal muscle,

and small intestine combined. Additionally, analyses of all modules are present for the

investigation of genes and networks not discussed in this paper. To ensure transparency and

reproducibility a data resource was created to contain the analyses performed along with code,

plots, and analysis description. The data resource can be intuitively navigated to meet the

specific needs of individual researchers and is accessible through the link to the website

(https://storage.googleapis.com/bl6_2dg_rnaseq/index.html).

The website is organized in a user-friendly manner with each WGCNA analysis housed under

its own tab, classified by tissue or tissue combination. In addition to the WGCNA analysis, the

website includes pages dedicated to the distribution analysis, outlier assessment, and sample

information, as well as additional analyses performed for the paper. Researchers can access

the raw code repository via a link to figshare containing the entire project with data, including the

Shiny app, and a Shiny app for basic plots for any gene identified in this study.

4. Discussion

The systemic effects of 2DG across the body have applications for understanding the potential

downstream effects of 2DG therapy, which has shown promise in treating cancer, epilepsy,

polycystic kidney disease, autoimmune disease, and in delaying Alzheimer's disease onset[1, 2,

4, 6, 7, 17, 47-50], and is currently being used to treat SARS-CoV-2[9, 11, 12]. Although 2DG

therapy has shown promise in treating these conditions, there is lingering concerns

regarding its safety. To better understand the potential risks associated with this

treatment, our study investigated the systemic effects of 2DG on the transcriptomes of nine

tissues in healthy young mice. By developing a novel filtering strategy of multiple statistical tests

to identify robust genes and pathways affected by 2DG in multiple tissues, we found that

WGCNA identified between 25 and 46 modules for each tissue, but only assessing modules

with overrepresented pathways reduced the number of modules of interest for each tissue to

between 12 and 32. Modules correlated with treatment further reduced the number of modules

of interest to between zero and three for each tissue. Finally, ANOVA revealed that heart,

hypothalamus, prefrontal cortex, small intestine, skeletal muscle, and liver were significantly

associated with treatment and fit our filtering strategy. These six tissues were assessed for their

robust 2DG effects, independently and collectively. Kidney, hippocampus, and spleen did not

meet all the filtering criteria and therefore were not discussed (see data resource for their

analysis).

2DG uniquely altered pathways and genes within each tissue module; however, tissue

alterations could still be broadly identified as a result of glycolysis or N-linked glycosylation

disruption. Glycolysis is necessary for energy production while N-linked glycosylation is

necessary for protein function[39]. By analyzing the functions of our high confidence pathways,

four tissues appeared to be affected by inhibition of glycolysis: heart, small intestine,

hypothalamus, and liver. In contrast, we inferred that the remaining two tissues were affected by

disruption of N-linked glycosylation: the prefrontal cortex and skeletal muscle.

Our study showed that 2DG treatment reduced the expression of gene pathways related to ATP

production and glycolysis in the heart, both of which have been shown to play a role in heart

failure[35, 51, 52]. Our results are consistent with a previous study[9], demonstrating that 2DG

altered these pathways, but did not report any cardiac toxicity or significant changes in

pathways related to ER stress. Although we did not perform EKGs, we observe no signs of

heart trouble in the treated mice; moreover, patients with moderate to severe SARS-CoV-2 or

advanced tumors treated with 2DG in clinical trials also showed no significant adverse heart-

related effects due to this treatment[5, 10, 53]. Our findings suggest that there may be a strong

association between ATP production and glycolysis pathways and the incidence of heart failure.

Therefore, the reduction of these pathways by 2DG treatment may provide a new avenue for

developing therapies to target these metabolic processes and prevent or treat heart failure.

While 2DG inhibits glycolysis, leading to a decrease in ATP production, it may also stimulate

compensatory mechanisms that enhance oxidative phosphorylation, another pathway for ATP

production[3, 9]. These findings suggest that 2DG primarily modifies metabolism in the heart

and may have implications for developing new therapies for cardiovascular diseases, including

using 2DG to improve heart health.

The small intestine is a unique immunological tissue because as it is the site in which immune

cells, both sessile and recirculating, are continuously exposed to gut microfloral stimuli[54]. This

encounter results in intestinal lymphocytes and leukocytes in an activated state that is not found

in other organs of healthy mice. In agreement, we found that the small intestine was the only

tissue in which untreated mice showed a very strong immune signature. Treatment with 2DG, in

the small intestine resulted in a striking dampening of pathways associated with adaptive

immunity and enrichment of innate immune pathways. Our analysis using CIBERSORT

suggested that genes related to memory B cells and Th1 cells had decreased expression in

mice treated with 2DG, which is in line with other studies showing a reduction in B and T cells

and an increase in macrophages and eosinophils in the presence of 2DG[15, 55]. The increase

in gene expression related to innate immunity pathways in the small intestine suggests that 2DG

treatment may enhance the intestinal immune response against invading pathogens, while the

decrease in gene expression related to adaptive immunity pathways may weaken the immune

response against certain pathogens, potentially leading to increased susceptibility to infections.

Additionally, our findings have implications for the potential relationship between the immune

system and microbiome of the small intestine as the reduction in expression of memory B cell

genes in mice treated with 2DG may promote dysbiosis. We envision two possibilities to explain

the effects of 2DG on intestinal immune signature: 1) 2DG acts directly to eliminate on

activated adaptive lymphocytes or their antigen presenting cells; and 2), 2DG eliminates gut

bacteria that are the cause of immune stimulation. Our results emphasize the modulatory

effects of 2DG on immunity in the small intestine, which may have implications for the

development of new therapies for immune-related disorders.

Fatty acid oxidation is a critical process that helps to maintain energy homeostasis by providing

energy substrates for ATP production[56]. The increase in fatty acid oxidation pathways in the

hypothalamus with 2DG treatment may play a crucial role in the regulation of energy balance

and metabolism when glycolysis is limited[57]. Our findings suggest that an increase in fatty

acid oxidation pathways in the hypothalamus with 2DG may have positive implications for the

overall health of the mice and inform new therapeutic approaches for metabolic disorders, such

as obesity and diabetes, which are characterized by metabolic dysregulation[57]. This

therapeutic approach may result in a shift towards energy expenditure and may help to

counteract the effects of a high-fat diet or other metabolic disorders. However, further studies

are needed to investigate the long-term effects of 2DG treatment on metabolic disorders in

humans. Additionally, it is reported that treatment with 2DG in the 3xTgAD mouse model of

Alzheimer's disease increased expression of genes involved in ketone formation in the brain,

reduced pathology, and resulted in increased serum ketone bodies[17]. Our study has shown

that the hypothalamus switched to fatty acid oxidation in the presence of 2DG, suggesting a

propensity of this critical anatomical structure to compensate for a reduction in glycolysis by

utilizing fatty acids. Future studies could explore the potential benefits and limitations of

targeting fatty acid oxidation pathways in the hypothalamus for the treatment of metabolic and

neurodegenerative disorders.

Treatment with 2DG alters genes involved in protein metabolism in both the prefrontal cortex

and skeletal muscle, suggesting potential implications for various diseases. In the prefrontal

cortex, 2DG reduces gene expression in protein metabolism pathways in the ER compared to

controls, which is associated with brain function regulation[58]. Previous studies have shown

that by shutting down protein synthesis, 2DG can induce the unfolded protein response,

reducing ER stress[14]. This is of particular interest as the prefrontal cortex is susceptible to

dysregulation due to ER stress[59, 60], which is linked to neurological disorders such as

Alzheimer's and Parkinson's disease[61]. Further investigations into the potential applications of

altered protein metabolism pathways in the prefrontal cortex with 2DG treatment could be

crucial for developing effective therapies for neurological disorders. Similarly, in skeletal muscle,

prolonged ERK activation events were reduced in mice treated with 2DG compared to controls,

which may have implications for metabolic health. Although ERK1/2 activation is necessary for

maintenance of myofibers and neuromuscular synapses, prolonged activation can lead to

muscle weakness[62-64]. Moreover, ERK pathway overactivation in skeletal muscle can lead to

inflammation through activation of TNF-α, IL-6, IL-1ß, and contribute to metabolic disorders

such as type 2 diabetes and obesity[65, 66]. This agrees with another study, which reported that

2DG decreases ERK pathways, resulting in decreased inflammatory markers[65]. Targeting

ERK activation with 2DG in humans may have significant implications for the prevention and

treatment of various diseases. Furthermore, the reduction of prolonged ERK activation events in

the muscle and protein synthesis in the prefrontal cortex of mice treated with 2DG may also

reduce inflammation and protect against misfolded proteins, which could be important for

developing therapies for related diseases.

Our comparison between tissues showed that each tissue module was unique at the gene and

pathway level demonstrating that 2DG affected each tissue in a unique manner. Although

tissues responded uniquely, we did note potential relationships across tissues in response to

2DG. These findings demonstrate that the complex effects of 2DG on different tissues and

pathways in mice, suggesting that further research is needed to understand the implications of

these effects for overall health. However, our results also suggest that 2DG therapy has

promising potential for the treatment of a range of diseases that affect different tissues and

pathways in humans.

Although the present analysis sheds light on the effects of 2DG in male mice, there are several

avenues for further investigation to expand our understanding of its potential benefits. First, it

would be informative to explore whether female mice respond differently to 2DG. Second, due

to the change in diet composition the effects of dosage length could not be assessed

systemically. Third, while our filtering strategy identified robust tissues and module responses to

2DG, it is possible that other effects were not captured. Additionally, future research in disease

models and in human patients is needed to fully understand the potential implications for clinical

applications, and it should be noted that our study did not directly evaluate the safety of 2DG.

Nonetheless, our findings improve our understanding of 2DG's effects on healthy tissues, which

may have implications for its safety in the long term.

5. Conclusion

Given the increasing interest in therapeutic use of 2DG, the goal of this study was to provide a

systematic analysis of the transcriptomic consequences of glycolysis blockade by 2DG in major

organs and tissues of healthy mammals. As we found surprisingly little transcriptional overlap

among tissue types affected by 2DG treatment, the results are best explained by 2DG uniquely

affecting each tissue type. The results are most consistent with two mechanisms through which

2DG is considered to act: inhibition of glycolysis and of N-linked glycosylation. This study

provides a comprehensive resource on the effects of 2DG, a glycolytic inhibitor, on various

tissues in non-diseased C57BL/6J mice, including a complete dataset, code, plots, and analysis

descriptions for transparency and reproducibility. The website

(https://storage.googleapis.com/bl6_2dg_rnaseq/index.html) is user-friendly, enabling easy

navigation through all available analyses from all nine tissues. This system-wide

characterization of 2DG activity across multiple tissues has the potential to inform its targeted

therapeutic use, taking into account the mechanism relevant to each disease-affected tissue.

However, further studies are needed to fully understand the mechanisms underlying the

responses and determine their physiological significance.

Funding: This work was supported by NIH grant GM115518 to GWC., The Jackson Laboratory

Director's Innovation Fund (JAX-DIF 19000-18-19) and the Longevity Impetus Grant from Norn

Group to C-HC.

CRediT authorship contribution statement

AEW: Methodology, Writing-Original draft preparation, Software, Visualization, Formal analysis,

Data Curation; **JJW:** Conceptualization, Methodology, Writing-Review and Editing, Project

administration; **SEH:** Investigation, Writing-Review, and Editing; **JDS:** Investigation; **JW:**

Investigation; **RP:** Conceptualization; **MWC:** Conceptualization, Writing-Review and Editing;

DCR: Conceptualization, Methodology, Resources, Writing-Review and Editing; **C-HC:**

Conceptualization, Methodology, Supervision, Resources, Writing-Review and Editing, Project

administration, Funding acquisition; **GWC:** Supervision, Resources, Writing-Review and Editing,

Funding acquisition.

Declaration of Competing Interest

The authors declare no conflict of interest.

Acknowledgements

We would like to thank Selcan Aydin for her invaluable discussions regarding the statistical

approach and in proofreading the data resource. We thank Tamar Abel for proofreading the

data resource. We thank Grace Stafford for aligning the transcriptomes and providing the

processed RNA-seq data.

References

[1] Ben Sahra I, Tanti JF, Bost F. The combination of metformin and 2 deoxyglucose inhibits autophagy and induces AMPK-dependent apoptosis in prostate cancer cells. Autophagy. 2010;6:670-1.

[2] Aft RL, Zhang FW, Gius D. Evaluation of 2-deoxy-D-glucose as a chemotherapeutic agent: mechanism of cell death. Br J Cancer. 2002;87:805-12.

[3] Sottnik JL, Lori JC, Rose BJ, Thamm DH. Glycolysis inhibition by 2-deoxy-D-glucose reverts the metastatic phenotype in vitro and in vivo. Clin Exp Metastasis. 2011;28:865-75.

[4] Cay O, Radnell M, Jeppsson B, Ahren B, Bengmark S. Inhibitory effect of 2-deoxy-D-glucose on liver tumor growth in rats. Cancer Res. 1992;52:5794-6.

[5] Raez LE, Papadopoulos K, Ricart AD, Chiorean EG, Dipaola RS, Stein MN, et al. A phase I dose-escalation trial of 2-deoxy-D-glucose alone or combined with docetaxel in patients with advanced solid tumors. Cancer Chemother Pharmacol. 2013;71:523-30.

[6] Koenig JB, Cantu D, Low C, Sommer M, Noubary F, Croker D, et al. Glycolytic inhibitor 2-deoxyglucose prevents cortical hyperexcitability after traumatic brain injury. JCI Insight. 2019;5.

[7] Sproule TJ, Wilson J, Adkins E, Crocker BP, Morel L, Roopenian DC. AI-19 Metabolic inhibition by 2-deoxyglucose prevents and reverses lupus in mice. Lupus Science & Medicine. 2016;3:A10-A.

[8] Wang H, Zhang N, Fang K, Chang X. 2-Deoxy-D-glucose Alleviates Collagen-Induced Arthritis of Rats and Is Accompanied by Metabolic Regulation of the Spleen and Liver. Front Immunol. 2021;12:713799.

[9] Aiestaran-Zelaia I, Sanchez-Guisado MJ, Villar-Fernandez M, Azkargorta M, Fadon-Padilla L, Fernandez-Pelayo U, et al. 2 deoxy-D-glucose augments the mitochondrial respiratory chain in heart. Sci Rep. 2022;12:6890.

[10] Bhatt AN, Shenoy S, Munjal S, Chinnadurai V, Agarwal A, Vinoth Kumar A, et al. 2-deoxy-D-glucose as an adjunct to standard of care in the medical management of COVID-19: a proof-of-concept and dose-ranging randomised phase II clinical trial. BMC Infect Dis. 2022;22:669.

[11] Pajak B, Zielinski R, Manning JT, Matejin S, Paessler S, Fokt I, et al. The Antiviral Effects of 2-Deoxy-D-glucose (2-DG), a Dual D-Glucose and D-Mannose Mimetic, against SARS-CoV-2 and Other Highly Pathogenic Viruses. Molecules. 2022;27.

[12] Sahu KK, Kumar R. Role of 2-Deoxy-D-Glucose (2-DG) in COVID-19 disease: A potential game-changer. J Family Med Prim Care. 2021;10:3548-52.

[13] Pajak B, Siwiak E, Soltyka M, Priebe A, Zielinski R, Fokt I, et al. 2-Deoxy-d-Glucose and Its Analogs: From Diagnostic to Therapeutic Agents. Int J Mol Sci. 2019;21.

[14] Xi H, Kurtoglu M, Liu H, Wangpaichitr M, You M, Liu X, et al. 2-Deoxy-D-glucose activates autophagy via endoplasmic reticulum stress rather than ATP depletion. Cancer Chemother Pharmacol. 2011;67:899-910.

[15] Zhao Q, Chu Z, Zhu L, Yang T, Wang P, Liu F, et al. 2-Deoxy-d-Glucose Treatment Decreases Anti-inflammatory M2 Macrophage Polarization in Mice with Tumor and Allergic Airway Inflammation. Front Immunol. 2017;8:637.

[16] Wang Q, Liang B, Shirwany NA, Zou MH. 2-Deoxy-D-glucose treatment of endothelial cells induces autophagy by reactive oxygen species-mediated activation of the AMP-activated protein kinase. Plos One. 2011;6:e17234.

[17] Yao J, Chen S, Mao Z, Cadenas E, Brinton RD. 2-Deoxy-D-glucose treatment induces ketogenesis, sustains mitochondrial function, and reduces pathology in female mouse model of Alzheimer's disease. Plos One. 2011;6:e21788.

[18] Abboud G, Choi SC, Kanda N, Zeumer-Spataro L, Roopenian DC, Morel L. Inhibition of Glycolysis Reduces Disease Severity in an Autoimmune Model of Rheumatoid Arthritis. Front Immunol. 2018;9:1973.

[19] Wang Y, Li JR, Sun MX, Ni B, Huan C, Huang L, et al. Triggering unfolded protein response
by 2-Deoxy-D-glucose inhibits porcine epidemic diarrhea virus propagation. Antiviral Res.
2014;106:33-41.
[20] Choi SC, Titov AA, Abboud G, Seay HR, Brusko TM, Roopenian DC, et al. Inhibition of
glucose metabolism selectively targets autoreactive follicular helper T cells. Nat Commun.
2018;9:4369.
[21] Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9:357-9.
[22] McCarthy DJ, Chen Y, Smyth GK. Differential expression analysis of multifactor RNA-Seq
experiments with respect to biological variation. Nucleic Acids Res. 2012;40:4288-97.
[23] Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data. Bioinformatics. 2010;26:139-40.
[24] Chen Y, Lun AT, Smyth GK. From reads to genes to pathways: differential expression
analysis of RNA-Seq experiments using Rsubread and the edgeR quasi-likelihood pipeline.
F1000Res. 2016;5:1438.
[25] Team RC. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2022.
[26] Langfelder P, Horvath S. WGCNA: an R package for weighted correlation network analysis.
BMC Bioinformatics. 2008;9:559.
[27] Horvath PLS. Fast R Functions for Robust Correlations and Hierarchical Clustering. Journal
of Statistical Software. 2012;46:1-17.
[28] Raudvere LKU. gprofiler2: Interface to the 'g:Profiler' Toolset. R package version 0.2.0
ed2020.
[29] Kay ME, L; Higgins, J; Wobbrock, J. _ARTool: Aligned Rank Transform for Nonparametric
Factorial ANOVAs_. 2021.
[30] Wobbrock JF, L; Gergle, D; Higgins, J. The Aligned Rank Transform for Nonparametric
Factorial Analyses Using Only ANOVA Procedures. Proceedings of the ACM
Conference on Human Factors in Computing Systems (CHI '11). 2011:143-6.
[31] Imai DTTYKHLKK. mediation: R Package for Causal Mediation Analysis. Journal of
Statistical Software. 2014;59:1-38.
[32] Hänzelmann SC, R.; Guinney, A. GSVA: gene set variation analysis for microarray and
RNA-seq data. BMC Bioinformatics. 2013;14.
[33] Xie Y. bookdown: authoring Books and Technical Documents with R markdown: Chapman
and Hall/CRC; 2016.
[34] Ye Z, Wang S, Zhang C, Zhao Y. Coordinated Modulation of Energy Metabolism and
Inflammation by Branched-Chain Amino Acids and Fatty Acids. Front Endocrinol (Lausanne).
2020;11:617.
[35] Jiang M, Xie X, Cao F, Wang Y. Mitochondrial Metabolism in Myocardial Remodeling and
Mechanical Unloading: Implications for Ischemic Heart Disease. Front Cardiovasc Med.
2021;8:789267.
[36] Adeva-Andany MM, Gonzalez-Lucan M, Donapetry-Garcia C, Fernandez-Fernandez C,
Ameneiros-Rodriguez E. Glycogen metabolism in humans. BBA Clin. 2016;5:85-100.
[37] Xie N, Zhang L, Gao W, Huang C, Huber PE, Zhou X, et al. NAD(+) metabolism:
pathophysiologic mechanisms and therapeutic potential. Signal Transduct Target Ther.
2020;5:227.
[38] Liu L, Su X, Quinn WJ, 3rd, Hui S, Krukenberg K, Frederick DW, et al. Quantitative Analysis
of NAD Synthesis-Breakdown Fluxes. Cell Metab. 2018;27:1067-80 e5.
[39] Aebi M. N-linked protein glycosylation in the ER. Biochim Biophys Acta. 2013;1833:2430-7.
[40] Hue L, Taegtmeyer H. The Randle cycle revisited: a new head for an old hat. Am J Physiol
Endocrinol Metab. 2009;297:E578-91.

[41] Moncan M, Mnich K, Blomme A, Almanza A, Samali A, Gorman AM. Regulation of lipid
metabolism by the unfolded protein response. J Cell Mol Med. 2021;25:1359-70.
[42] Calder PC. Branched-chain amino acids and immunity. J Nutr. 2006;136:288S-93S.
[43] Baakhtari M, Imaizumi N, Kida T, Yanagita T, Ramah A, Ahmadi P, et al. Effects of
branched-chain amino acids on immune status of young racing horses. J Vet Med Sci.
2022;84:558-65.
[44] Yahsi B, Gunaydin G. Immunometabolism - The Role of Branched-Chain Amino Acids.
Front Immunol. 2022;13:886822.
[45] Mu WC, VanHoosier E, Elks CM, Grant RW. Long-Term Effects of Dietary Protein and
Branched-Chain Amino Acids on Metabolism and Inflammation in Mice. Nutrients. 2018;10.
[46] Lee D, Hokinson D, Park S, Elvira R, Kusuma F, Lee JM, et al. ER Stress Induces Cell
Cycle Arrest at the G2/M Phase Through eIF2alpha Phosphorylation and GADD45alpha. Int J
Mol Sci. 2019;20.
[47] Dwarakanath B, Jain V. Targeting glucose metabolism with 2-deoxy-D-glucose for
improving cancer therapy. Future Oncol. 2009;5:581-5.
[48] Ciavardelli D, Rossi C, Barcaroli D, Volpe S, Consalvo A, Zucchelli M, et al. Breast cancer
stem cells rely on fermentative glycolysis and are sensitive to 2-deoxyglucose treatment. Cell
Death Dis. 2014;5:e1336.
[49] Chiaravalli M, Rowe I, Mannella V, Quilici G, Canu T, Bianchi V, et al. 2-Deoxy-d-Glucose
Ameliorates PKD Progression. J Am Soc Nephrol. 2016;27:1958-69.
[50] Tailler M, Lindqvist LM, Gibson L, Adams JM. By reducing global mRNA translation in
several ways, 2-deoxyglucose lowers MCL-1 protein and sensitizes hemopoietic tumor cells to
BH3 mimetic ABT737. Cell Death Differ. 2019;26:1766-81.
[51] Sun H, Olson KC, Gao C, Prosdocimo DA, Zhou M, Wang Z, et al. Catabolic Defect of
Branched-Chain Amino Acids Promotes Heart Failure. Circulation. 2016;133:2038-49.
[52] Neinast MD, Jang C, Hui S, Murashige DS, Chu Q, Morscher RJ, et al. Quantitative
Analysis of the Whole-Body Metabolic Fate of Branched-Chain Amino Acids. Cell Metab.
2019;29:417-29 e4.
[53] Stein M, Lin H, Jeyamohan C, Dvorzhinski D, Gounder M, Bray K, et al. Targeting tumor
metabolism with 2-deoxyglucose in patients with castrate-resistant prostate cancer and
advanced malignancies. Prostate. 2010;70:1388-94.
[54] Wiertsema SP, van Bergenhenegouwen J, Garssen J, Knippels LMJ. The Interplay
between the Gut Microbiome and the Immune System in the Context of Infectious Diseases
throughout Life and the Role of Nutrition in Optimizing Treatment Strategies. Nutrients. 2021;13.
[55] Cai W, Cheng J, Zong S, Yu Y, Wang Y, Song Y, et al. The glycolysis inhibitor 2-
deoxyglucose ameliorates adjuvant-induced arthritis by regulating macrophage polarization in
an AMPK-dependent manner. Mol Immunol. 2021;140:186-95.
[56] Houten SM, Violante S, Ventura FV, Wanders RJ. The Biochemistry and Physiology of
Mitochondrial Fatty Acid beta-Oxidation and Its Genetic Disorders. Annu Rev Physiol.
2016;78:23-44.
[57] McFadden JW, Aja S, Li Q, Bandaru VV, Kim EK, Haughey NJ, et al. Increasing fatty acid
oxidation remodels the hypothalamic neurometabolome to mitigate stress and inflammation.
Plos One. 2014;9:e115642.
[58] Butterfield DA, Favia M, Spera I, Campanella A, Lanza M, Castegna A. Metabolic Features
of Brain Function with Relevance to Clinical Features of Alzheimer and Parkinson Diseases.
Molecules. 2022;27.
[59] Trinh MA, Kaphzan H, Wek RC, Pierre P, Cavener DR, Klann E. Brain-specific disruption of
the eIF2alpha kinase PERK decreases ATF4 expression and impairs behavioral flexibility. Cell
Rep. 2012;1:676-88.

[60] Kim P, Scott MR, Meador-Woodruff JH. Dysregulation of the unfolded protein response
(UPR) in the dorsolateral prefrontal cortex in elderly patients with schizophrenia. Mol Psychiatry.
2021;26:1321-31.
[61] Ajoolabady A, Lindholm D, Ren J, Pratico D. ER stress and UPR in Alzheimer's disease:
mechanisms, pathogenesis, treatments. Cell Death Dis. 2022;13:706.
[62] Shi H, Scheffler JM, Zeng C, Pleitner JM, Hannon KM, Grant AL, et al. Mitogen-activated
protein kinase signaling is necessary for the maintenance of skeletal muscle mass. Am J
Physiol Cell Physiol. 2009;296:C1040-8.
[63] Penna F, Costamagna D, Fanzani A, Bonelli G, Baccino FM, Costelli P. Muscle wasting
and impaired myogenesis in tumor bearing mice are prevented by ERK inhibition. Plos One.
2010;5:e13604.
[64] Seaberg B, Henslee G, Wang S, Paez-Colasante X, Landreth GE, Rimer M. Muscle-
derived extracellular signal-regulated kinases 1 and 2 are required for the maintenance of adult
myofibers and their neuromuscular junctions. Mol Cell Biol. 2015;35:1238-53.
[65] Francis R, Singh PK, Singh S, Giri S, Kumar A. Glycolytic inhibitor 2-deoxyglucose
suppresses inflammatory response in innate immune cells and experimental staphylococcal
endophthalmitis. Exp Eye Res. 2020;197:108079.
[66] Wen X, Zhang B, Wu B, Xiao H, Li Z, Li R, et al. Signaling pathways in obesity:
mechanisms and therapeutic interventions. Signal Transduct Target Ther. 2022;7:298.
786

Created with Biorender.com

787
788 **Figure 1:** Experimental Design and analysis flow. C57BL/6J mice were treated with 2DG (6g/L)
789 for 96 hours or 4 weeks. Nine tissues were harvested for bulk transcriptomics. Genes identified
790 for each tissue were clustered into modules using WGCNA. Effects of 2DG on each module
791 were determined by assessing each modules relationship with treatment through correlation,
792 ANOVA, and pathway analysis. Created with Biorender.com
793
794

795
796 **Figure 2:** Principal Component Analysis demonstrated clustering of samples based on tissue
797 type. Hotelling's T^2 ellipses were calculated, and the clustering patterns of all tissue were
798 contained within the 95% confidence intervals.
799
800

801
Figure 3: Analysis of modules-treatment association. A) A total of 46 modules were identified in
the small intestine, each containing between 22 and 2,259 genes. B) Correlation analysis
between time, treatment, and treatment in a time dependent manner revealed three modules
significantly correlated with treatment. C) The average eigengene expression of the green4
module shows that expression levels are lower in mice treated with 2DG compared to control
mice. D) Summary eigengene expression for each sample in the green4 module confirms lower
expression in mice treated with 2DG compared to control mice., and E) Individual genes across
samples clustered by treatment except for one control sample, which clustered with 2DG-
treated mice.

811
812

813
814
815 **Figure 4:** Number of identified genes shared between tissue modules. Each dot represents the
816 tissue or tissues being compared. Very few genes were shared across tissues and no gene was
817 shared across all tissues. Only tissue combinations with one or more shared genes are shown.
818
819

820
821 **Figure 5:** Correlation structure of gene module changes across tissues in 2DG-treated mice.
822 Six tissues, each represented by one high confidence module, demonstrated unique response
823 to 2DG treatment. The expression levels of genes and pathways within each tissue showed

824   varying degrees of over- and under-expression in response to 2DG. Created with
825   Biorender.com.
826
827

828
829
Figure 6: Correlations between pathway eigengenes in the heart (y-axis) and small intestine (x-
axis). Branched chain amino acid degradation was negatively correlated with pathways
predominantly related to the adaptive immunity and positively correlated with pathways
predominantly related to innate immunity in small intestine. All other metabolic pathways in
heart were negatively correlated with pathways related to innate immunity and positively
correlated with pathways related to adaptive immunity in small intestine.
836
837

838
Figure 7: Correlations between pathway eigengenes in the hypothalamus (y-axis) and
prefrontal cortex (x-axis). G2/M DNA damage checkpoint in the hypothalamus was negatively
correlated with pathways related to ER stress and positively correlated with pathways related to
RNA transcription in the prefrontal cortex. All other metabolic pathways in the hypothalamus
were positively correlated with pathways related to ER stress and negatively correlated with
pathways related to RNA transcription in the prefrontal cortex.