A multiplexed RT-PCR Assay for Nanopore Whole Genome Sequencing of Tilapia lake virus (TiLV)

Running title: *Whole genome sequencing of TiLV using Nanopore*

Jerome Delamare-Deboutteville[1#*], Watcharachai Meemetta[2#], Han Ming Gan[3], Chadag Vishnumurthy Mohan[1], Ha Thanh Dong[4], Saengchan Senapin[2,5*]

[1]WorldFish, Penang, Malaysia
[2]Fish Health Platform, Center of Excellence for Shrimp Molecular Biology and Biotechnology (Centex Shrimp), Faculty of Science, Mahidol University, Rama VI Rd., Bangkok, 10400, Thailand
[3]Patriot Biotech Sdn Bhd, Bandar Sunway, 47500, Selangor, Malaysia
[4] School of Environment, Resources and Development, Asian Institute of Technology, Pathum Thani, 12120, Thailand
[5]National Center for Genetic Engineering and Biotechnology (BIOTEC), National Science and Technology Development Agency (NSTDA), Pathum Thani 12120, Thailand

*Corresponding authors:
J. Delamare-Deboutteville, j.delamare@cgiar.org
S Senapin, senapin2010@gmail.com

# first authors with equal contributions

Acknowledgments

This research was undertaken as part of the CGIAR Initiatives on Aquatic Foods led by WorldFish, and the One CGIAR Initiative "Protecting Human Health Through a One Health Approach", supported by contributors to the CGIAR Trust Fund: https://www.cgiar.org/funders/. Funding provided support in the form of salary for authors [J.D.D; C.V.M], travels, laboratory consumables and analytical costs.

Data Availability Statement

The data that support the findings of this study are available at the following links: demultiplexed FastQ files for all ten samples can; be found under BioProject PRJNA957495 with the corresponding BioSample accession SAMN34257318 (B1-1), SAMN34257319 (B1-2), SAMN34257320 (FM2), SAMN34257321 (Nk), SAMN34257322 (A1-2), SAMN34257323 (A1-3), SAMN34257324 (D1-2), SAMN34257325 (Ri), SAMN34257326 (Cell_line), SAMN34257327 (RiverWater).

The Linux scripts used to generate raw fastq files, assembled genomes and phylogenetics tree along with the generated files are publicly available in the Zenodo.org dataset (https://zenodo.org/record/7851622)

Conflict of interest statement

The authors declare no conflict of interest.

Ethics approval statement

The authors confirm that the ethical policies of the journal, as noted on the journal's author guidelines page, have been adhered to. No ethical approval was required as no animals were used in this study. Virus sequences were generated from archived samples

Abstract

Tilapia lake virus (TiLV) is a highly contagious viral disease that affects tilapia, a globally significant and affordable source of fish protein. To combat the introduction and spread of TiLV and its impact, there is an urgent need for increased surveillance, improved biosecurity measures, and continuous development of effective diagnostic and sequencing methods. In this study, we have developed a multiplexed RT-PCR assay that can amplify all ten complete genomic segments of TiLV from various sources of isolation. The amplicons generated using this approach are immediately compatible with real-time sequencing on the Nanopore system. By using this approach, we have recovered and assembled 13 TiLV genomes from total RNA extracted from naturally TiLV-infected tilapia fish, concentrated tilapia rearing water, and other isolation sources. Our phylogenetic analysis, consisting of more than 36 TiLV genomes from both newly sequenced and publicly available TiLV genomes, provides new insights into the high phylogenetic diversity of TiLV from Thailand, suggesting multiple genetic origins. This work is an essential stepping stone towards integrating rapid and real-time Nanopore-based amplicon sequencing into routine genomic surveillance of TiLV, as well as future vaccine development.

Keywords: Red tilapia (*Oreochromis* spp.), Tilapia lake virus, one-step RT-PCR, multiplex RT-PCR, whole genome sequencing, Oxford Nanopore Technologies, diagnostic by sequencing, bioinformatics and phylogeny analyses

INTRODUCTION

Tilapia lake virus disease (TiLV) is a highly contagious viral disease that affects tilapia, an important and affordable fish protein source produced in aquaculture globally. TiLV has rapidly spread since, now reported in over 16 countries (Debnath et al., 2022), posing a significant threat to tilapia production and the livelihoods of farmers who rely on tilapia farming for income and food security (Hounmanou et al., 2018). To combat the introduction and spread of TiLV and its impacts, increased surveillance, improved biosecurity measures, farming practices and continuous development of effective diagnostic and sequencing methods are urgently needed.

The TiLV genome consists of 10 segments that complicates its genome sequencing process thus precluding high mass-scale genome sequencing efforts to be undertaken. The first TiLV genome was sequenced using a shotgun transcriptome approach on an Illumina sequencing platform (Bacharach et al., 2016). Recently, a similar approach using shotgun metagenomics was used to generate the near complete genome of a TiLV isolate causing mass-mortality in Tilapia farmed in Bangladesh (Chaput et al., 2020). Shotgun metagenomics involved the random sequencing of all RNA fragments, i.e. TiLV-positive tilapia liver samples without any enrichment of mRNA; followed by bioinformatics analysis to identify and assemble the 10-segments of TiLV genome present in the sample. This approach is not scalable as the RNA library preparation cost is higher and a majority of the sequencing data will belong to the host, requiring high sequencing depth to successfully assemble the TiLV-derived contigs. More importantly, Illumina technology requires significant investment of infrastructure, precluding rapid on-site deployment and real-time sequencing.

In recent years, Oxford Nanopore Technologies (ONT) have become more commonly used to sequence part(s) or whole genomes of pathogens affecting aquatic animals (Gallagher et al., 2018). Nanopore sequencing offers several advantages, including high throughput, portability, cost-effectiveness, and real-time sequencing, which can greatly facilitate the detection and sequencing of viral genomes in remote locations (Alathari et al., 2023). Rapid amplicon-based reverse transcription polymerase chain reaction (RT-PCR) assays coupled with Nanopore technologies can provide a sensitive and specific means of detecting and genotyping TiLV in field samples, allowing for fine epidemiological surveillance and timely management and control of outbreaks (Delamare-Deboutteville et al., 2021). To date, Nanopore has been used to sequence the genomes of non-segmented fish viruses such as infectious spleen and kidney necrosis virus (ISKNV) (Alathari et al., 2023), salmonid alphavirus (SAV) and infectious salmon anaemia virus (ISAV) (Gallagher et al., 2018).

Complete viral genomes provide valuable information on the genetic diversity, evolution, and potential virulence factors. Such methods can enhance our understanding of TiLV epidemiology, transmission dynamics, and evolution, and contribute to the development of effective control and management strategies.

In this study, for the first time, we designed a new method for TiLV whole genome sequencing using singleplex and multiplex amplicon-based RT-PCR protocols coupled with Minion Nanopore sequencing. These new tools can be used for real-time monitoring and characterization of TiLV for better surveillance and control measures in tilapia aquaculture.

MATERIALS AND METHODS

Primer design

TiLV primers targeting 10 complete genome segments (containing conserved sequences at 5' and 3' termini) were designed based on sequences of the Israel strain Til-4-2011 (GenBank accession no. KU751814 to KU751823) (Bacharach et al., 2016) (Table 1).

Sampling and total RNA extraction

RNA template for TiLV genome sequence amplification and analysis was prepared from tissues of TiLV infected Nile tilapia (*Oreochromis niloticus*), red tilapia (*Oreochromis* spp.), TiLV isolates propagated in E-11 cell culture, and concentrated water samples from fish ponds (Table 2). Extraction of total RNA used the Trizol reagent (Invitrogen) according to the manufacturer's instruction followed by spectrophotometry-based quantification measuring the absorbance at $OD_{260\,nm}$ and $OD_{280\,nm}$.

Development of singleplex one-step RT-PCR for the enrichment of TiLV genome

The efficiency of the designed TiLV primers and their optimal annealing temperatures (Ta) were investigated by one-step gradient RT-PCR assays with the range of Ta from 50 to 60°C. RT-PCR reaction mixture (25 µl) comprised of 0.5 µl of SuperScript III RT/Platinum Taq Mix (Invitrogen catalog no. 12574-018), 12.5 µl of 2X reaction mix, 2 µl of RNA template (100 ng/µl), 1 µl of 10 µM each primer, and distilled water. The temperature profile included a reverse transcription (RT) step at 50°C for 30 min, heat-inactivation of RT enzyme at 94°C for 2 min, 30 cycles of denaturation at 94°C for 30 sec, annealing step for 30 sec, extension 72°C for 1 min-2 min (1min/kb), and a final extension at 72°C for 2 min. Amplified products were analyzed by agarose gel electrophoresis

Development of multiplex RT-PCR to streamline the PCR enrichment of TiLV genome

Two multiplex PCR reactions were developed to reduce the number of PCR reactions from 10 to only two reactions per sample. The primers were divided into two sets based on their annealing temperatures similarity. Reaction 1 employs primers for segment 1, 2, 3, 4, 5 and 8 with Ta at 52°C while reaction 2 uses primers for segment 6, 7, 9 and 10 with Ta at 60°C Then, various PCR conditions were tested by varying the dNTPs (200 nM - 500 nM), MgSO4 (1.6 mM - 1.8 mM), enzyme (1 - 2.5 volume) and primer concentrations (100 - 300 nM) to obtain optimal PCR outcomes.

Nanopore sequencing

PCR products were pooled (10 PCR reactions and 2 PCR reactions were pooled for the singlex and multiplex protocol, respectively) followed by PCR clean up using NucleoSpin Gel and PCR Clean-up column (Macherey-Nagel) and quantification with Qubit dsDNA Broad Range kit (Invitrogen). Approximately 250 ng of the purified and pooled amplicons was used as the template for library preparation using the native barcoding expansion 1-12 kit (EXP-NBD104) according to the manufacturer's instructions. The prepared library was loaded onto a R9.4.1 Flongle and sequenced for 24 hours. Basecalling of the fast5 raw signals used Guppy v4.4.1 in super accuracy mode to generate the fastq sequences for subsequent bioinformatics analysis.

Reference-based genome assembly of TiLV samples

Raw reads were quality- and length-filtered using NanoFilt (qscore > 9 and length > 250bp). The raw and filtered read statistics were generated using seqkit v.2.1.0. Reference-based genome assembly of the TiLV was performed according to the ARTIC pipeline (https://github.com/artic-network/fieldbioinformatics) (Tyson et al., 2020). This pipeline is an open-source software that integrates a series of tools for base-calling, quality control, read trimming, reference-based mapping, variant calling, consensus sequence generation, and annotation. Briefly, the filtered reads were aligned to the reference TilV genome using Minimap2 v2.17 (Li, 2018) followed by variant calling using Medaka (r941_min_sup_g507) (https://github.com/nanoporetech/medaka). The variants identified were subsequently filtered based on several criteria, including the quality score, depth of coverage, strand bias, and frequency of occurrence. In addition, genomic regions with read depth of lower than 20x were masked prior to generating the final consensus sequence for each sample. Each assembled viral segment from each sample was analyzed with QUAST v5 (Gurevich et al., 2013) to calculate the percentage of the assembled viral genome that is represented by gaps (Ns), providing insights into the PCR and pooling efficiency.

Phylogenetic analysis

The assembled viral segments with less than 20% gap were selected and combined with publicly available TiLV genomes for phylogenetic analysis (Table 2). The DNA sequences of the viral genome segments from each sample were extracted and grouped based on their segment number followed by alignment with MAFFT v8 (--adjustdirection --maxiterate 1000 --localpair) (Katoh & Standley, 2013). All 10 individual alignments were subsequently concatenated and used to reconstruct a maximum likelihood tree using FastTree 2 (Price et al., 2010). The resulting tree was visualized and annotated using FigTree v1.4.4 (http://tree.bio.ed.ac.uk/software/figtree/).

RESULTS

Ten primer pairs for the recovery of complete TiLV genome from various isolation source

A total of 10 primer pairs were designed with their PCR condition optimized (Table 1 and Supplemental Table 1) to amplify the complete segment of one of the ten TiLV genomic segments. Intact and specific band corresponding to the respective size of the TiLV genomic segments could be obtained when the total RNA from TiLV-infected tilapia, TiLV-infected E-11 cell line, and concentrated pond water sample were used as the template for RT-PCR (Figure 1 and data not shown). However, the PCR band intensity for segment 5 (1,099 bp) is substantially lower compared to the other segments, requiring another round of PCR.

A streamlined two-tube multiplex RT-PCR for TiLV genome amplification

To avoid unnecessary human error associated with handling multiple singleplex (N=10) PCR reactions, a two-tube multiplex RT-PCR was designed (Supplemental Table 2). The addition of $MgSO_4$ (increased magnesium ion concentration) was crucial for improved sensitivity (stronger band intensity) while an increase in dNTP concentration does not improve PCR efficiency (Supplemental Figure 2). In addition, increasing the amount of RT/Taq enzyme mix was also shown to slightly improve over band intensity (Supplemental Figure 1). As a result, the concentration of $MgSO_4$ and RT/Taq enzyme mix was increased in further multiplex RT-PCR assays (Supplemental Figure 2). Unlike the singleplex PCR (Figure 1), the band intensity of the segment 5 amplicon was similar to that of other segment amplicons (Figure 2A). However, while the expected number of amplicons was observed in the cell-line derived sample, this is not the case for various naturally-infected samples (Figure 2A and 2B), particularly for the first multiplex RT-PCR targeting viral segments 1-5 and 8 (Figure 2A). It is worth noting that samples with inconsistent amplification displayed relatively higher Cq values (> 20). indicating lower viral load. Sample A1-2 that showed no observable bands in both multiplex RT-PCR reactions, resembling that of negative control (-ve) has a Cq of 37.46 and 32.06 for Segment 1 and Segment 9, respectively (Table 3). Additional RT-PCR amplification of other clinical samples with known Cq values (TiLV copies) based on qRT-PCR also showed that the Set 1 multiplex RT-PCR was indeed less sensitive showing inconsistent to no observable bands when the Cq value is more than 19 (Supplemental Figure 3).

Rapid and on-site Sequencing of the TiLV genome using Oxford Nanopore

A total of 413,379 demultiplex raw reads with an accumulative length of 238,983,973 bp were generated from the Flongle sequencing runs (Supplemental Table 3). After filtering (qscore > 9, length > 250 bp), only 194,564 reads (147,415,018 bp) remain. On average more than 35% data reduction was observed across the samples with sample A1-2 showing the largest reduction (67.7%, from 28 Mb to 9.2 Mb) in the amount of usable data. Overall, the Arctic-based reference genome assembly could successfully assemble the viral segments 6,7,8,9,10 for all samples with more than 99-100% completeness with the exception of samples A1-2_m and Nk_m that showed only a slightly lower completeness of 98% for segments 6 and 7 (Table 3). On the contrary, several segments from the first set of multiplex RT-PCR showed reduced completeness (high % of gaps in sequence) particularly for samples with high Cq. The reduced completeness is a direct result of the low read depth (< 20) observed for the viral segments in the respective samples. Generally, any viral segment with a read depth of more than 50x will produce a highly complete assembly that can be used in subsequent analysis (Table 3).

High phylogenetic diversity among Thai TiLV strains

The total alignment length after the concatenation of 10 individually aligned TiLV viral segments is 10,396 bp. Using a midpoint rooting approach, multiple clades with high SH-like support values were observed in the maximum likelihood tree (Figure 3). Nanopore-sequenced samples from either the pooled singleplex or multiplex amplicons were always placed in the same cluster, consistent with their identical sample origin (Table 2). TiLV strains from Peru, Ecuador, Israel and India were clustered together and this subclade subsequently formed a sister group with slightly lower support with two earlier TiLV strains from Thailand isolated in 2013 and 2014 to form Clade A (Figure 3). Clade B consisting entirely of Thai TiLV strains from 2015 to 2016 formed a sister group with Clades A. However, a majority of Thai TiLV strains that were reported in 2018 onwards showed yet another distinct clustering as indicated by their phylogenetic placement in Clades C and E with a majority of the sequences reported in this study belonging to subclade E1. On the other hand, subclade E2 consists of a mixture of Thai and USA TiLV strains. The currently sampled Bangladeshi TiLV strains consist of only single clade despite being isolated 2 years apart (2017 and 2019) while the Vietnamese strains are highly divergent even between themselves (Pairwise nucleotide similarity of only 92%), possibly representing novel strains of TiLV.

DISCUSSION

In this study, we report the successful recovery of the complete TiLV genome using a novel approach that combines multiplex PCR and Nanopore amplicon sequencing. Our method employs a primer binding region at the terminal 5' and 3' ends of each viral segment, which ensures maximal preservation of genetic information. Notably, since the maximum length of the viral segment is less than 1500 bp, our approach obviates the need for a PCR-tiling strategy typically used for recovering large non-segmented viruses such as ISKNV and SAR-CoV-2 (Lambisia et al., 2022) (Alathari et al., 2023). Moreover, our method offers the added advantage of visualizing PCR efficiency and specificity for each viral segment on a gel, as each fragment has a different size

Our current multiplex PCR appears to show lower efficiency for Multiplex Set 1 particularly in samples with high Cq values. To improve the multiplex RT-PCR amplification uniformity and efficiency, the Multiplex Set 1 reaction, which amplifies TiLV genomic segments 1-5 and 8, can be further split into two pools (e.g 1A and 1B) that will amplify an average of three viral segments each. In addition, the use of a more processive High-Fidelity Taq polymerase such as Q5 that was currently used for high-degree multiplex tiling PCR of the SAR-CoV19 and ISKNV viral genomes is also worth exploring (Tyson et al., 2020) (Alathari et al., 2023). It is also worth noting that despite the absence of visible bands for some of the samples, partial or even near-complete genome assembly was still attainable using our sequencing pipeline. It is possible that the amount of PCR product is below the detection limit of gel-staining dye at its loading concentration although it is in fact present in the gel. To streamline future work in high throughput sequencing of TiLV using this approach, gel visualization may be skipped once a lab can consistently reproduce the PCR outcome with evidence from sequencing data.

Nanopore sequencing is an attractive approach for viral amplicon sequencing due to its portability, convenience, and speed (Delamare-Deboutteville et al., 2021). Our method, which utilizes Nanopore sequencing, eliminates the need for additional fragmentation steps, allowing motor proteins to be directly ligated to amplicons for native sequencing. On the same day, tens

of samples can be prepared and sequenced, and the low computing requirements of the ARTIC protocol (Artic Network, n.d.) enable swift genome assembly on a laptop computer, without requiring access to a dedicated server. To further streamline TiLV genome sequencing on the Nanopore platform, we suggest designing multiplex primers that incorporate a partial adapter suitable for Nanopore sequencing (Ezpeleta et al., 2022). This enables cost-effective PCR-based barcoding that is both efficient and scalable. In cases of low data output, samples can be re-pooled and sequenced on a separate flow cell to achieve the necessary sequencing depth for genome assembly.

By utilizing R9.4.1 sequencing chemistry with super accuracy mode and implementing the ARTIC pipeline, we successfully recovered TiLV genomes that are highly suitable for phylogenetic inference. Our study revealed the presence of TiLV in both fish and environmental water samples from the same farm, which clustered together in Clade E1. Our approach, combining the previously reported water sample concentration method (Taengphu et al., 2022) with a multiplex RT-PCR amplicon-based Nanopore sequencing strategy, allowed for direct recovery of TiLV genomes from water samples. This innovative method has significant implications for environmental monitoring, as it eliminates the need for sacrificing fish for genomic analysis.

Our analysis indicates that, in addition to country of origin, host genetic background may also contribute to the clustering patterns observed within Clade E. Specifically, our results demonstrate distinct phylogenetic grouping of Thai TiLV strains (E1: red tilapia, E2: nile tilapia), suggestive of multiple introductions into the country. Furthermore, our findings contradict the initial hypothesis previously put forth on Tilapia trade movement, which was based on a small genome-based phylogenetic tree with poorly supported clustering of Bangladeshi and Thai TiLV strains (Chaput et al., 2020). Specifically, we found no grouping of Thai strains within the Bangladesh clade (Figure 3, Clade D), thereby reducing support for the previously proposed hypothesis.

Although viral whole genome sequencing of TiLV is now technically feasible, the current representation of its genome in public databases is limited, making it difficult to infer its evolutionary relationships. Given the significant impact of TiLV on the tilapia aquaculture industry, there is a critical need for more robust genomic surveillance to facilitate better management and tracking. Our proposed Nanopore-based amplicon sequencing approach offers a promising solution, as it enables cost-effective and high-throughput sequencing of TiLV virus genomes. We anticipate that our approach will provide a valuable resource for ongoing efforts to understand the molecular epidemiology and evolution of TiLV, with important implications for disease control and prevention.

References

Alathari, S., Chaput, D. L., Bolaños, L. M., Joseph, A., Jackson, V. L. N., Verner-Jeffreys, D., Paley, R., Tyler, C. R., & Temperton, B. (2023). A Multiplexed, Tiled PCR Method for Rapid Whole-Genome Sequencing of Infectious Spleen and Kidney Necrosis Virus (ISKNV) in Tilapia. *Viruses*, *15*(4), Article 4. https://doi.org/10.3390/v15040965

Al-Hussinee, L., Subramaniam, K., Ahasan, M. S., Keleher, B., & Waltzek, T. B. (2018). Complete Genome Sequence of a Tilapia Lake Virus Isolate Obtained from Nile Tilapia (Oreochromis niloticus). *Genome Announcements*, *6*(26), e00580-18. https://doi.org/10.1128/genomeA.00580-18

Bacharach, E., Mishra, N., Briese, T., Zody, M. C., Kembou Tsofack, J. E., Zamostiano, R., Berkowitz, A., Ng, J., Nitido, A., Corvelo, A., Toussaint, N. C., Abel Nielsen, S. C., Hornig, M., Del Pozo, J., Bloom, T., Ferguson, H., Eldar, A., & Lipkin, W. I. (2016). Characterization of a novel orthomyxo-like virus causing mass die-offs of tilapia. *MBio*, *7*(2), e00431-16. https://doi.org/10.1128/mBio.00431-16

Chaput, D. L., Bass, D., Alam, Md. M., Al Hasan, N., Stentiford, G. D., Van Aerle, R., Moore, K., Bignell, J. P., Haque, M. M., & Tyler, C. R. (2020). The segment matters: Probable reassortment of tilapia lake virus (TiLV) complicates phylogenetic analysis and inference of geographical origin of new isolate from Bangladesh. *Viruses*, *12*(3), 258. https://doi.org/10.3390/v12030258

Debnath, P. P., Delamare-Deboutteville, J., Jansen, M. D., Phiwsaiya, K., Dalia, A., Hasan, M. A., Senapin, S., Mohan, C. V., Dong, H. T., & Rodkhum, C. (2020). Two-year surveillance of tilapia lake virus (TiLV) reveals its wide circulation in tilapia farms and hatcheries from multiple districts of Bangladesh. *Journal of Fish Diseases*, *43*(11), 1381–1389. https://doi.org/10.1111/jfd.13235

Debnath, P. P., Dinh-Hung, N., Taengphu, S., Nguyen, V. V., Delamare-Deboutteville, J., Senapin, S., Vishnumurthy Mohan, C., Dong, H. T., & Rodkhum, C. (2022). Tilapia Lake Virus was not detected in non-tilapine species within tilapia polyculture systems of Bangladesh. *Journal of Fish Diseases*, *45*(1), 77–87. https://doi.org/10.1111/jfd.13537

Delamare-Deboutteville, J., Taengphu, S., Gan, H. M., Kayansamruaj, P., Debnath, P. P., Barnes, A., Wilkinson, S., Kawasaki, M., Vishnumurthy Mohan, C., Senapin, S., & Dong, H. T. (2021). Rapid genotyping of tilapia lake virus (TiLV) using Nanopore sequencing. *Journal of Fish Diseases*, *44*(10), 1491–1502. https://doi.org/10.1111/jfd.13467

Eyngor, M., Zamostiano, R., Kembou Tsofack, J. E., Berkowitz, A., Bercovier, H., Tinman, S., Lev, M., Hurvitz, A., Galeotti, M., Bacharach, E., & Eldar, A. (2014). Identification of a novel RNA virus lethal to tilapia. *Journal of Clinical Microbiology*, *52*(12), 4137–4146. https://doi.org/10.1128/JCM.00827-14

Gallagher, M. D., Matejusova, I., Nguyen, L., Ruane, N. M., Falk, K., & Macqueen, D. J. (2018). Nanopore sequencing for rapid diagnostics of salmonid RNA viruses. *Scientific Reports*, *8*(1), 16307. https://doi.org/10.1038/s41598-018-34464-x

Gurevich, A., Saveliev, V., Vyahhi, N., & Tesler, G. (2013). QUAST: Quality assessment tool for genome assemblies. *Bioinformatics*, *29*(8), 1072–1075. https://doi.org/10.1093/bioinformatics/btt086

Hounmanou, Y. M. G., Mdegela, R. H., Dougnon, T. V., Achoh, M. E., Mhongole, O. J., Agadjihouèdé, H., Gangbè, L., & Dalsgaard, A. (2018). Tilapia lake virus threatens tilapiines farming and food security: Socio-economic challenges and preventive measures in Sub-Saharan Africa. *Aquaculture*, *493*, 123–129.

https://doi.org/10.1016/j.aquaculture.2018.05.001

Katoh, K., & Standley, D. M. (2013). MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution*, *30*(4), 772–780. https://doi.org/10.1093/molbev/mst010

Li, H. (2018). Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics*, *34*(18), 3094–3100. https://doi.org/10.1093/bioinformatics/bty191

Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2 – Approximately Maximum-Likelihood Trees for Large Alignments. *PLOS ONE*, *5*(3), e9490. https://doi.org/10.1371/journal.pone.0009490

Pulido, L. L. H., Mora, C. M., Hung, A. L., Dong, H. T., & Senapin, S. (2019). Tilapia lake virus (TiLV) from Peru is genetically close to the Israeli isolates. *Aquaculture*, *510*, 61–65. https://doi.org/10.1016/j.aquaculture.2019.04.058

Subramaniam, K., Ferguson, H. W., Kabuusu, R., & Waltzek, T. B. (2019). Genome sequence of tilapia lake virus associated with syncytial hepatitis of tilapia in an Ecuadorian aquaculture facility. *Microbiology Resource Announcements*, *8*(18), e00084-19. https://doi.org/10.1128/MRA.00084-19

Surachetpong, W., Janetanakit, T., Nonthabenjawan, N., Tattiyapong, P., Sirikanchana, K., & Amonsin, A. (2017). Outbreaks of tilapia lake virus infection, Thailand, 2015–2016. *Emerging Infectious Diseases*, *23*(6), 1031–1033. https://doi.org/10.3201/eid2306.161278

Taengphu, S., Sangsuriya, P., Phiwsaiya, K., Debnath, P. P., Delamare-Deboutteville, J., Mohan, C. V., Dong, H. T., & Senapin, S. (2020). Genetic diversity of tilapia lake virus genome segment 1 from 2011 to 2019 and a newly validated semi-nested RT-PCR method. *Aquaculture*, *526*, 735423. https://doi.org/10.1016/j.aquaculture.2020.735423

Thawornwattana, Y., Dong, H. T., Phiwsaiya, K., Sangsuriya, P., Senapin, S., & Aiewsakun, P. (2021). Tilapia lake virus (TiLV): Genomic epidemiology and its early origin. *Transboundary and Emerging Diseases*, *68*(2), 435–444. https://doi.org/10.1111/tbed.13693

Tran, T. H., Nguyen, V. T. H., Bui, H. C. N., Tran, Y. B. T., Tran, H. T. T., Le, T. T. T., Vu, H. T. T., & Ngo, T. P. H. (2022). Tilapia Lake Virus (TiLV) from Vietnam is genetically distantly related to TiLV strains from other countries. *Journal of Fish Diseases*, *45*(9), 1389–1401. https://doi.org/10.1111/jfd.13669

Tyson, J. R., James, P., Stoddart, D., Sparks, N., Wickenhagen, A., Hall, G., Choi, J. H., Lapointe, H., Kamelian, K., Smith, A. D., Prystajecky, N., Goodfellow, I., Wilson, S. J., Harrigan, R., Snutch, T. P., Loman, N. J., & Quick, J. (2020). Improvements to the ARTIC multiplex PCR method for SARS-CoV-2 genome sequencing using nanopore. *BioRxiv*, 2020.09.04.283077. https://doi.org/10.1101/2020.09.04.283077

Table1. TiLV primers designed in this study. Underlines represent conserved TiLV genomic sequences at the 5' and 3' end of each segment.

Target gene	Sequence (5' →3')	Ta (°C)	Size (bp)	Reference
Segment 1	**CS1F;** GCAAATATTTCTCTCATTCGCCTAT **CS1R;** CCAAACGTTATCTCTTAATTACGCA	55	1,641	This study
Segment 2	**CS2F;** GCAAATCTTTCTCATTATTACCATA **CS2R;** CCAAATTTTACTCTCTATTACCAAA	55	1,471	This study
Segment 3	**CS3F;** GCAAATTTTTCCCATAATCCTCTAT **CS3R;** CCAAATATTACCCCTTAATCCTTAA	55	1,371	This study
Segment 4	**CS4F;** GCAAATCTTTCTCCAATTACCGTCT **CS4R;** CCAAAGTTTACTCCTATTACCCAGA	50	1,250	This study
Segment 5	**CS5F;** GCAAATTTTACTCTTTTTCTCAGTT **CS5R;** CCAAATGTTTCTCTTATCTCAGACT	58	1,099	This study
Segment 6	**CS6F;** GCAAATATTTCTCTCAATCAAGCAC **CS6R;** CCAAATTTTACCTCTCGCATGCATT	60	1,044	This study
Segment 7	**CS7F;** GCAAATCTTTCTCTCATGCTACCAT **CS7R;** CCAAATTTTACTCTCTTTGCATTGC	60	777	This study
Segment 8	**CS8F;** GCAAATTTTTCTCATCATTACACAA **CS8R;** CCAAATATTACCTCATCTACACTAA	58	657	This study
Segment 9	**CS9F;** GCAAATCTTTCTCACGTCCTTAAAG **CS9R;** CCAAATTTTACTCACAAGTCCGATT	60	548	This study
Segment 10	**CS10F;** GCAAATCTTTCCCTCTGACACC **CS10R;** CCAAATTTTAACCCTACTAACACCA	60	465	This study

Table 2. Background information of TiLV strains reported in study as well as strains with publicly available genomes.

#	Sequence name (strain)	Country of origin	Collection date	Fish host	Tissues	Accession number	Reference	Sequencing Technology
1	B1-1_m	TH	2021	Red tilapia	L, K, S	SRR24222694	This study	Nanopore
2	B1-2_m	TH	2021	Red tilapia	L, K, S	SRR24222693	This study	Nanopore
3	FM2_m	TH	2019	Nile tilapia	L, K, S	SRR24222689	This study	Nanopore
4	Nk_m	TH	2019	Nile tilapia	L, B	SRR24222688	This study	Nanopore
5	A1-2_m	TH	2021	Red tilapia	L, K, S	SRR24222687	This study	Nanopore
6	A1-3_s (TH-2021-A)	TH	2021	Red tilapia	L, K, S	SRR24222682	This study	Nanopore
7	A1-3_m (TH-2021-A)	TH	2021	Red tilapia	L, K, S	SRR24222686	This study	Nanopore
8	D1-2_s (TH-2021-B)	TH	2021	Red tilapia	L, K, S	SRR24222692	This study	Nanopore
9	D1-2_m (TH-2021-B)	TH	2021	Red tilapia	L, K, S	SRR24222685	This study	Nanopore
10	Ri_m	TH	2020	Nile tilapia	L	SRR24222684	This study	Nanopore
11	Cell_line_s (RB19)	BD	2019	Nile tilapia	E-11 cells	SRR24222690	This study	Nanopore
12	Cell_line_m (RB19)	BD	2019	Nile tilapia	E-11 cells	SRR24222683	This study	Nanopore
13	RiverWater_s	TH	2021	Red tilapia (water)	na	SRR24222691	This study	Nanopore
14	IL-2011 (Til-4-2011)	IL	2011	Tilapia	na	KU751814–823	(Eyngor et al., 2014)	Illumina
15	TH-2013	TH	2013	Nile tilapia	Fe	MN687685-695	(Taengphu et al., 2020)	Sanger
16	TH-2014	TH	2014	Nile tilapia	Fr	MN687695-704	(Thawornwattana et al., 2021)	Sanger
17	TH-2015	TH	2015	Nile tilapia	Fi	MN687705-714	(Thawornwattana et al., 2021)	Sanger
18	TH-2016-CU	TH	2016	Nile tilapia	Ju	MN687715-724	(Thawornwattana et al., 2021)	Sanger
19	TH-2016-CN	TH	2016	Red tilapia hybrid	Fi	MN687725-734	(Thawornwattana et al., 2021)	Sanger
20	TH-2016 (TV1)	TH	2016	Red tilapia	na	KX631921–930	(Surachetpong et al., 2017)	Sanger
21	TH-2017	TH	2017	Nile tilapia	na	MN687735-744	(Thawornwattana et al., 2021)	Sanger
22	TH-2018-N	TH	2018	Nile tilapia	Fi	MN687745-754	(Thawornwattana et al., 2021)	Sanger
23	TH-2018-K	TH	2018	Nile tilapia	Ju	MN687755-764	(Thawornwattana et al., 2021)	Sanger
24	TH-2018 (WVL18053-01A)	TH	2018	Nile tilapia	na	MH319378–387	(Al-Hussinee et al., 2018)	Illumina
25	TH-2019	TH	2019	Nile tilapia	Fi	MN687765-774	(Thawornwattana et al., 2021)	Sanger
26	EC-2012 (EC-2012)	EC	2012	Nile tilapia	na	MK392372–381	(Subramaniam et al., 2019)	Illumina
27	PE-2018 (F3-4)	PE	2018	Nile tilapia	na	MK425010–019	(Pulido et al., 2019)	Sanger
28	US-2019 (WVL19031-01A)	US	2018	Nile tilapia	na	MN193513–522	unpublished*	Illumina
29	US-2019 (WVL19054)	US	2019	Nile tilapia	na	MN193523–532	unpublished*	Illumina
30	BD-2017 (BD-2017)	BD	2017	Nile tilapia	na	MN939372–381	(Chaput et al., 2020)	Illumina
31	BD-2019-E1	BD	2019	Nile tilapia	na	MT466447-456	(Debnath et al., 2020)	Sanger
32	BD-2019-E3	BD	2019	Nile tilapia	na	MT466457-466	(Debnath et al., 2020)	Sanger
33	BD-2017-181	BD	2017	Nile tilapia	na	MT466437-446	(Debnath et al., 2020)	Sanger
34	IND-2018	IND	2017	Nile tilapia	na	MZ297923-932	unpublished*	Illumina
35	VN (HB196-VN-2020)	VN	2020	Nile tilapia	na	ON376572-581	(Tran et al., 2022)	Illumina; Sanger
36	VN (RIA2-VN-2019)	VN	2019	Nile tilapia	na	ON376582-591	(Tran et al., 2022)	Illumina; Sanger

Country of origin: TH, Thailand; IL, Israel; EC, Ecuador; PE, Peru; US, USA; BD, Bangladesh; IND, India; VN, Vietnam
Fish host: Red tilapia, *Oreochromis* sp.; Nile tilapia, *Oreochromis niloticus*; Hybrid tilapia *Oreochromis niloticus* x *Oreochromis aureus*
L, Liver; K, Kidney; S, Spleen; B, Brain; Fe, fertilized eggs; Fr, fry; Fi, fingerlings; Ju, juveniles; na, info not available; *, direct submission

Table 3. PCR outcome, sequencing and alignment statistics of 10 individual TiLV segments for each sample used in this study. Viral segments with similar color indicate that they were amplified together in a multiplex reaction. The Cq values from the qPCR detection of the segment 1 (Cq seg1) and segment 9 (Cq seg9) were shown below each sample. +/- in the mPCR (multiplex PCR) row indicates presence (+) or absence (-) of visible PCR band representing the respective viral segment. %COV (% coverage) indicates the percentage of bases in the assembled contig that consists of a non-ambiguous base.

		TiLV segments (Length)									
		seg01	seg02	seg03	seg04	seg05	seg06	seg07	seg08	seg09	seg10
		(1641)	(1471)	(1371)	(1250)	(1099)	(1044)	(777)	(657)	(548)	(465)
A1-2_m Cq seg1: 37.46 Cq seg9: 32.06	mPCR	-	-	-	-	-	-	-	-	-	-
	Depth	37	36	20	51	26	522	578	428	371	414
	%COV	42	62	37	99	95	98	98	99	99	100
A1-3_s Cq seg1: 20.57 Cq seg9: 15.32	mPCR	+	+	+	+	+	+	+	+	+	+
	Depth	890	628	586	836	744	875	1100	1113	1212	2442
	%COV	100	100	100	100	100	100	100	100	100	100
A1-3_m Cq seg1: 20.57 Cq seg9: 15.32	mPCR	+	+	+	+	+	+	+	+	+	+
	Depth	113	139	6	72	12	813	723	673	563	604
	%COV	66	100	0	100	11	100	100	100	100	100
B1-1_m Cq seg1: 19.17 Cq seg9: 15.22	mPCR	+	+	+	+	+	+	+	+	+	+
	Depth	227	125	19	160	38	2428	2000	2701	1357	910
	%COV	60	61	51	68	99	100	100	100	100	100
B1-2_m Cq seg1: 20.47 Cq seg9: 16.31	mPCR	+	+	+	+	+	+	+	+	+	+
	Depth	260	239	47	185	66	1800	2237	2304	1367	1319
	%COV	62	100	100	100	99	100	100	100	100	100
Cell_line_s Cq seg1: 28.87 Cq seg9: 25.12	mPCR	+	+	+	+	+	+	+	+	+	+
	Depth	1260	1046	744	907	888	365	930	1766	1717	1614
	%COV	100	100	100	100	100	100	100	100	99	100
Cell_line_m Cq seg1: 28.87 Cq seg9: 25.12	mPCR	+	+	+	+	+	+	+	+	+	+
	Depth	99	43	94	157	106	748	1303	1732	1138	1290
	%COV	100	99	99	100	100	100	100	100	99	100
D1-2_s Cq seg1: 17.95 Cq seg9: 13.99	mPCR	+	+	+	+	+	+	+	+	+	+
	Depth	731	610	550	511	541	715	834	753	1297	1235
	%COV	100	100	100	100	100	100	100	100	100	100
D1-2_m Cq seg1: 17.95 Cq seg9: 13.99	mPCR	+	+	+	+	+	+	+	+	+	+
	Depth	2555	2914	1468	3471	1684	11387	8037	5031	7085	5987
	%COV	100	100	100	100	100	100	100	100	100	100
FM2_m Cq seg1: 30.86 Cq seg9: 27.18	mPCR	+	+	+	-	+	+	+	+	+	+
	Depth	11	18	8	26	55	794	1328	1665	772	1674
	%COV	14	39	0	96	98	100	100	100	99	100
Nk_m Cq seg1: 35.40 Cq seg9: 31.06	mPCR	-	+	+	-	+	+	+	+	+	+
	Depth	10	14	3	9	8	161	217	316	497	872
	%COV	21	15	0	0	0	99	99	100	99	99
Ri_m Cq seg1: 27.02 Cq seg9: 22.78	mPCR	-	-	-	-	-	-	+	+	+	+
	Depth	28	24	13	14	16	2254	4647	1246	3436	3224
	%COV	33	48	0	18	9	100	100	100	100	100
RiverWater_s Cq seg1: Nd Cq seg9: Nd	mPCR	+	+	+	+*	+	+	+	+	+	+
	Depth	566	540	698	629	1827	1268	2246	905	1415	1306
	%COV	100	100	100	100	100	100	100	100	100	100

Notes: +, detectable; -, undetectable; +*, detectable from re-amplification; Nd, Not done; low Ct green all the way

Figure legends

Figure 1. Gel electrophoresis results of one-step RT-PCR amplification of 10 genomic segments of TiLV. A 1% agarose gel was used to visualize the PCR products, with expected band sizes indicated at the bottom of the gel. M, DNA marker (New England Biolabs).

Figure 2. Amplification results of multiplex PCR (mPCR) for TiLV segments. Two separate reactions (Reaction#1 and Reaction#2) were used to amplify 10 TiLV segments. Reaction#1 amplified segments 1, 2, 3, 4, 5, and 8, while Reaction#2 amplified segments 6, 7, 9, and 10. A 2-log DNA marker (New England Biolabs) was used to visualize the PCR products. -ve, no template control.

Figure 3. Maximum likelihood tree showing the evolutionary relationships of TiLV strains analyzed in this study. Thirteen strains (10 unique samples) with ONT-TH prefix and publicly available genomes were used. The blue colored tip labels indicate the TiLV strains reported in this study. SH-like local support values and branch length indicate the number of substitutions per site.

Supplemental Figure 1. Multiplex RT-PCR condition optimization. The original reaction (Non) and modified reactions with additions of dNTPs, MgSO4, dNTPs + MgSO4, and RT/Taq enzyme mix were compared. A DNA marker (New England Biolabs) was used to visualize the PCR products.

Supplemental Figure 2. Multiplex RT-PCR amplification of TiLV segments. Two sets of reactions were used to amplify 10 TiLV segments using TiLV RNA templates. Conditions C1 and C2 were used for Reaction 1 and Reaction 2, respectively. C1 amplified segments 1, 2, 3, 4, 5, and 8, while C2 amplified segments 6, 7, 9, and 10. A DNA marker (New England Biolabs) was used to visualize the PCR products.

Supplemental Figure 3. Multiplex RT-PCR amplification of 10 TiLV genomic segments from six clinical samples (samples 1-6). Set 1 and Set 2 reactions were used to amplify the 10 TiLV segments. Set 1 amplified segments 1, 2, 3, 4, 5, and 8, while Set 2 amplified segments 6, 7, 9, and 10. DNA markers (New England Biolabs), positive control (P), and no template control (N) were used to visualize and confirm the PCR products. TiLV loads of each sample and the amplification results are summarized on the left panel table.

A

B