**Genomes from 25 historical *Drosophila melanogaster* specimens illuminate adaptive and**

**demographic changes across more than 200 years of evolution**

Short title (70 char):  Museum Fly Genomes Reveal Recent Adaptive and Demographic History

Max Shpak[1], Hamid R. Ghanavi[2], Jeremy D. Lange[1,^], John E. Pool[1,@], Marcus C. Stensmyr[2,3,@]

[1]Laboratory of Genetics, University of Wisconsin–Madison, USA

[2]Department of Biology, Lund University, Sweden

[3]Max Planck Center on Next Generation Insect Chemical Ecology

[^]Present address: Genus PLC, Waunakee, WI, USA

[@]Corresponding authors:  jpool@wisc.edu, marcus.stensmyr@biol.lu.se

**ABSTRACT**

The ability to perform genomic sequencing on long-dead organisms is opening new frontiers in evolutionary research. These opportunities are especially profound in the case of museum collections, from which countless documented specimens may now be suitable for genomic analysis. Here, we report 25 newly sequenced genomes from museum specimens of the model organism *Drosophila melanogaster*, including the oldest extant specimens of this species. By comparing historical samples ranging from the early 1800s to 1933 against modern day genomes, we document evolution across thousands of generations, including time periods that encompass the species' initial occupation of northern Europe and an era of rapidly increasing human activity. At the genome-wide level, we find that historical flies from the same time and place show much greater evidence for relatedness than flies from modern collections, and some show evidence of inbreeding as well, potentially reflecting either much smaller local population sizes in the past or else the specific circumstances of the collections. We also find that the Lund, Sweden population underwent local genetic differentiation during the early 1800s to 1933 interval (potentially due to accelerated drift) but then became more similar to other European populations thereafter (potentially due to increased migration). Within each time period, our temporal sampling allows us to document compelling candidates for recent natural selection. In some cases, we gain insights regarding previously implicated selection candidates, such as *ChKov1*, for which our inferred timing of selection favors the hypothesis of antiviral resistance over insecticide resistance. Other candidates are novel, such as the circadian-related gene *Ahcy*, which yields a selection signal that rivals that of the DDT resistance gene *Cyp6g1*. These insights deepen our understanding of recent evolution in a model system, and highlight the potential of future museomic studies.

## INTRODUCTION

Museum collections around the world contain billions of specimens collected over the last centuries. These collections serve as a window back in time – to an era before industrialization and modern agricultural practices – and could as such provide insights into issues ranging from insecticide resistance to climate change. The analysis of so-called historical DNA (hDNA), *i.e.* extracted genetic material from the last centuries, from museum samples accordingly holds much promise (Card et al. 2021; Raxworthy & Smith 2021). For example, whole genome analysis of historical museum samples has been used to e.g. document temporal genomic erosion in the endangered Grauer's and mountain gorillas (van der Valk et al. 2019), and to decipher the evolutionary response of rapidly changing environmental conditions in montane chipmunks (Bi et al. 2019).

A handful of prior studies have targeted whole genome sequences from museum insect specimens. These have included proof-of-concept sequencing studies (Staats et al. 2013; Korlević et al. 2021), and investigations of the taxonomic status of museum specimens (Mikheyev et al. 2017; Grewe et al. 2021), with one study including butterfly specimens up to 150 years old (Cong et al. 2021). A few insect museomic studies have sequenced multiple individuals to examine potential targets of recent positive selection, such as focusing on the responses of honey bees to the introduction of a parasitic mite (Mikheyev et al. 2015; Parejo et al. 2020) and the emergence of the Colorado potato beetle as a crop pest (Cohen et al. 2022). In short, museomics has enabled previously unreachable research topics to be examined and brought new relevance to the myriad of specimens collected and carefully curated over the centuries in natural history collections worldwide. However, there remains much untapped potential in this area, especially in terms of sequencing larger numbers of specimens of greater antiquity, in order to extend the temporal horizon and the range of questions that museomic studies can address.

*D. melanogaster* has long been a primary model system for population genetics, including the identification of targets of selective sweeps. Recently, the availability of data from multiple time points has begun to improve the prospects for distinguishing natural selection

from neutral evolution in this system. Genome sequencing of wild-caught and laboratory-
maintained *Drosophila melanogaster* individuals has enabled the study of genomic evolution
across seasonal (*e.g.* Bergland et al. 2014; Machado et al. 2021) and decadal (Lange et al.
2022) time-scales, making the species an important model for temporal genomic study.
However, no previous study has investigated genome-scale variation from historical *D.*
*melanogaster* specimens, nor from any species of such a minute animal. We here set out to
sequence genomes from historical *D. melanogaster* specimens using minimally destructive
techniques. Our ultimate goal was to identify changes in genome-wide diversity and specific
genetic loci showing signs of having differentiated over time. Although not an agricultural
pest, nor a disease vector – and accordingly not an intentional target of control methods or a
casualty of altered agricultural practices – as a strict human commensal outside of its native
African range, the drastic changes brought upon the planet since the dawn of the
Anthropocene (here defined as the onset of the industrial revolution ca. 1780) should have left
its marks in the genome of the fly. The roughly two hundred year time-frame of our analysis
here should also encompass the earliest stages of this species' adaptation to a novel high
latitude environment (Keller 2007). Hence, museomic study of European *D. melanogaster*
offers the potential to reveal which genes contributed to adaptive evolution during this pivotal
time period. Furthermore, this interval also offers unique opportunities to investigate the
demographic unknowns of early versus recent northern European *D. melanogaster*
populations (such as the their local size and connectivity). Finally, for no other species do we
have as detailed understanding of the genetic organization, molecular function, and the
physiological and behavioral manifestations of most genes, making this fly an ideal subject for
temporal genomic investigations.

The insect collections held in Lund and Stockholm are among the oldest in the world
and contain many important samples, including the specimens used to erect the genus
*Drosophila* (Fallén, 1814-1826). The Swedish collections span the whole Anthropocene, from
the first steam engines to widespread industrialization and urbanization, to altered agricultural
practices and the introduction of mass-produced chemical insecticides, to globalization and
climate change (**Figure 1A**). The oldest material in the Swedish museums are four specimens

from the collection of Carl Fredrik Fallén (1764-1830; **Figure 1B**) – professor in natural history at Lund university and the authority behind genus *Drosophila*. These flies are kept in the natural history museum in Stockholm but were collected in Lund. No date is provided, nor is the collector specified, but the specimens were probably collected before 1810 (see below). From about the same time stems three specimens from the collection of Johan Wilhelm Zetterstedt (1785-1874; **Figure 1C**), which are kept in Lund. Zetterstedt was a student of Fallén, and from 1822 professor in natural history at Lund University. This material comprises the lectotype (not examined here) and two paralectotypes of *Drosophila approximata*, a junior synonym of *D. melanogaster*. The two paralectotypes we have examined are labelled Lund and Smol (*i.e.* Småland) respectively, the latter indicating the province north of Scania. No collector is specified. From the collection in Copenhagen there are two specimens from Passau, Germany (**Figure 1D**) collected by Joseph Waltl (1805-1888), physician and naturalist, and from 1835 a docent and later professor in natural history at the Philosophisch-Theologische Hochschule Passau. These flies are similarly not dated but likely collected in the 1850s. Precise collection locality is unknown. Also from Copenhagen, there is one specimen from the latter half of the 19[th] century collected by Rasmus William Traugott Schlick (1839-1916) on northeast Zealand, Denmark (**Figure 1E**). Again, no date is provided, but Schlick collected extensively in this region in the 1880s. Lastly, we have a set of 15 specimens from the collection in Lund (**Figure 1F**), collector unknown, labelled "Zootis 1933" – the informal name of the former Zoological Department at Lund University (**Figure 1G**).

**Figure 1: Historical context of the museum specimens and time interval encompassed by the present study.**

(A) Timeline of the Anthropocene showing the chronology of the analyzed fly samples, notable human events, and climate trends in Sweden (Temperature data from SMHI, Sweden).

(B) One of the four *D. melanogaster* specimens from the Fallén collection at the Swedish Natural History Museum (Stockholm). Insert shows the label (in Fallén's handwriting) accompanying the specimen. Scale bar: 1 mm. Photo: C. Fägerström.

(C) One of three *D. melanogaster* specimens from the Zetterstedt collection at the Biological

Museum (Lund). Insert depicts the attached label. Scale bar: 1 mm. Photo: C. Fägerström.

(D) *D. melanogaster* specimen from the Natural History Museum of Denmark (Copenhagen),

collected by Joseph Waltl in Passau, Germany. Scale bar: 1 mm. Photo: C. Fägerström.

(E) *D. melanogaster* specimen from the Natural History Museum of Denmark (Copenhagen),

collected by William Schlick on NE Själland, Denmark. Scale bar: 1 mm. Photo: C. Fägerström.

(F) 15 *D. melanogaster* specimens from Lund, kept in collections of the Biological Museum

(Lund). Scale bar: 5 mm. Photo: C. Fägerström.

(G) The old department of Zoology building at Lund University. Photo: Lund University.

(H) Lydbecksa Gården in Lund, where Fallén had his lodgnings until 1811. Photo: Lund

University.

(I) Sandgatan 5 in Lund, where Zetterstedt lived. Photo: Lund University.

(J) A representation of the 6 contemporary population samples included in the present

analysis. Photo: M. Stensmyr

The precise timing, location, and collector for the old Swedish material is not easy to

decipher. The Fallén samples are accompanied by a single (contemporary) note, in what looks

like Fallén's handwriting, stating (translated) "*Imago and larvae in sawdust [*unintelligible*] can*

*of raisins*". If the flies were caught by Fallén himself, a potential locality would then be his

lodgings at Lidbeckska gården in central Lund (**Figure 1H**), and that the material was collected

around 1807-1809, based on the known whereabouts of Fallén (who after 1811 spent

considerable time at his wife's family's mansion at Esperöd, near Simrishamn). From his

personal correspondence, the first mention of *Drosophila* can be found in a letter to Zetterstedt

dated 14 September 1810 (Zetterstedt archive, Lund University). The samples in Zetterstedt's

collection are likewise difficult to precisely date. From Zetterstedt's *Diptera Scandinaviae*

*disposita et descripta* we can read that Zetterstedt collected *D. approximata* in Lund (possibly

at his home at Sandgatan 5 in central Lund; **Figure 1I**). The specimens from Småland were

attributed to Sven Ingemar Ljungh (1757-1828), civil servant and prominent naturalist, who

resided at Skärsjö manor (Jönköping County, Småland – roughly 200 km from Lund). Whether

or not the specimens in the collection are the ones mentioned in his treatise on Scandinavian

Diptera we cannot know, but if so, the flies would likely have been caught within a decade or

so of Fallén's samples.  It should be noted that Zetterstedt kept many of Fallén's specimens in

his collection, and moreover, that given the considerable age of these collections, labels might

have been lost or details obscured over the years, and lastly, although both Fallén and

Zetterstedt kept type specimens, detailed labelling was evidently not considered all that

important.

All in all, the Swedish and Danish museum collections contain 26 *D. melanogaster*

specimens suitable for analysis.  The old Swedish specimens are to the best of our knowledge

the earliest *D. melanogaster* available in any collection, likely predating the Meigen holotypes

kept in Paris by two decades or more.  We here provide whole-genome analysis from these

flies, obtaining from most of them relatively complete and high quality genomes, and compare

them against modern fly genomes (**Figure 1J**).  This analysis may span more than 3,000 fly

generations (Turelli and Hoffmann 1995; Pool 2015), analogous to studying human evolution

across more than 75,000 years (approximately corresponding to the time period in which

modern humans first colonized Eurasia from Africa, e.g. Mallick et al  2016, Jouganos et al

2017, Yang and Fu 2018, Wohns et al 2022).  Examining changes in genomic diversity across

two roughly century-scale time intervals, we find that the relationship between the Lund

population and other European populations has changed over time.  We identify a variety of

strong candidates for the action of positive selection in each time interval, providing temporal

context for previously known cases of selection, while also identifying novel putative selection

targets.   These findings illustrate the potential of museomic studies to deepen our

understanding of recent evolution in a rapidly changing world.

**RESULTS**

**Most historical flies yielded genomes of comparable quality to contemporary flies**

For proof-of-principle, we first attempted to extract DNA from one of the 1933 "Zootis"

specimens (**Figure 1F**).  Briefly, the whole fly – still attached to the pin – was first transferred to

a lysate buffer containing Proteinase K and left at 56°C for 2 hours, after which the fly was

removed, washed, dehydrated, and returned to the collection (**Figure 2A**). The extraction

process left the fly largely unharmed, except for a slight loss in coloration and some shrinkage

of the abdomen (**Figure 2B**). The extracted DNA was subsequently prepared for Illumina short

read sequencing. As expected, the DNA was highly fragmented with an average fragment

length of about ~50 bp. The Illumina run yielded ~24 million reads, which we subsequently

mapped onto a *D. melanogaster* reference genome. The ensuing historic genome was 91%

complete with a 15.7X mean sequencing depth. In short, the employed method was indeed

able to extract enough DNA to piece together largely complete genomes, while doing minimal

harm to the specimen.

**Figure 2: Sequencing of historical *Drosophila* specimens yields genomes with slightly**

**reduced genomic coverage and some evidence of cytosine deamination.**

(A) Illustration of the process of extracting and sequencing DNA from pinned insect specimens.

Flies were submerged in a lysate buffer for two hours, then put through three washing steps

and dried, before being restored to the museum repository. Genomic libraries were then

prepared and sequenced using the Illumina NovaSeq 6000 platform. Further details appear

in the Methods section.

(B) Images of the appearance of a representative specimen (from 1933) before and after the

DNA extraction protocol, illustrating minor changes in coloration. Photo: C. Fägerström.

(C) The fraction of genomic coverage is plotted as a function of mean per-site read depth for

each historical fly genome, and for a representative set of contemporary inbred line genomes

from Lyon, France. For a given level of depth most historical fly genomes achieve levels of

genomic coverage only slightly lower than contemporary genomes.

(D) The relationship between depth and unique (singleton) variant-calling. Only the lowest

quality historical fly genomes show notable elevations in the rate of singleton variants called.

(E) The fraction of singleton variants called as adenine or thymine is elevated for historical fly

genomes, particularly those with greater singleton rates, potentially due to cytosine

deamination.

Having verified the method, we next proceeded to process the other specimens for

DNA extraction and sequencing. Out of the 25 specimens examined, we were able to generate

genomes from all but one. While the mean depth of coverage per site varies among the

historical fly genomes (**Figure 2C**), the median value among these genomes was

approximately 20X, which is comparable to typical *D. melanogaster* genomes generated from

contemporary source material (Lack et al. 2016). We find, however, that with similar mean

depth, genomic coverage tends to be slightly lower in historical versus contemporary

genomes (often with only 2-3% less of the genome covered; **Figure 2C**, **Table S1**).

The modest reduction in genomic coverage in most of the historical fly genomes may

relate to their consistently shorter insert sizes – typically only about 50 bp (**Table S1**), compared

to contemporary genomes which are typically sequenced from fragments of a few hundred bp

in length. The shorter DNA fragments from the old specimens are expected to limit our ability

to align into repetitive genomic regions, and hence such regions may not receive base calls in

the historical genomes, leading to somewhat reduced genomic coverage as observed above.

Genomes from the 1800s flies had mean insert sizes averaging 47.6 bp, while those from the

1933 collection averaged 50.8 bp. Although the distributions of mean insert sizes overlap

between time points, they nevertheless differ significantly ($P$ = 0.0188; Mann Whitney two-

tailed *U* test). There was a marginal correlation between fragment length and depth of

coverage (correlation $r$ = 0.40, $P$ = 0.055). A stronger and significant association was found

between fragment length and genomic coverage ($r$ = 0.63, $P$ = 0.0001).

The shorter length of historical fly DNA fragments may be due to DNA degradation,

which may also explain the reduced depth and genomic coverage obtained from a minority of

the examined specimens. Among the nine 1800s samples, four genomes were characterized

by a low average read depth of < 10 reads per site and/or a low genomic coverage of < 80%.

For the 1933 samples, one sample was immediately excluded from further analyses due to

extremely low coverage and read depth. Most samples with low coverage also had low mean

read depth (**Figure 2C**; **Table S1**), and the overall correlation between these quality metrics

was statistically significant ($r$ = 0.404, $P$ = 0.018). We note that all seven low-quality genomes

(as defined above) were from female flies (out of 14 total females), whereas all 11 males yielded

higher coverage/depth genomes. This sex difference is statistically significant (two-tailed

binomial $P = 0.0057$) and may be due to female flies being larger and potentially experiencing

greater decay before dehydration.

Regarding the six historical fly genomes retained with lower sequencing depth,

inferences of diploid genotype are not statistically robust at sites for which an individual has

few reads. Therefore, we treated the lower quality genomes as though they were haploid for

autosomal and female X chromosomes by assigning only a single, most frequent nucleotide at

each site. Consequently, there were effectively 14 rather than 18 autosome alleles for the

1800s samples and 28 rather than 30 for the 1933 autosomes. Similarly, treating each low-

quality female X as haploid reduced allele counts to 11 and 21 for 1800s and 1933, respectively

(from what would otherwise be X chromosome allele counts of 15 and 23).

The DNA degradation inferred above could also lead to incorrect base identification

via anomalous sample-specific variants at individual sites. Therefore, we assessed whether

singleton variants (unique to individual genomes) were enriched in samples with low depth

and coverage. Indeed, we found a higher relative incidence of singleton variants unique to

genomes with lower genomic coverage (two-tailed binomial $P < 0.01$; Figure 2B) and to a

lesser extent those with low depth as well ($P = 0.05$).

One of the most common types of nucleotide degradation is cytosine deamination,

which can lead to spurious thymine enrichment, which has been documented extensively in

ancient DNA samples (e.g. Dabney et al 2013). We therefore assessed the extent to which A/T

nucleotides are over-represented among singletons from each genome, and we found a

higher fraction of A/T sites in singletons among the historical samples than in modern genomes

(**Figure 2D, E; Table S1**). Whereas the modern genomes had singleton A/T frequencies of

~55%, many of the historical genomes' singletons were 60-74% AT, presumably due to

cytosine deamination. Although most historical genomes do not show meaningfully elevated

singleton rates, our findings regarding singleton AT% justify the exclusion of singleton alleles

for analyses that are sensitive to rare alleles and small frequency differences, such as

demographic inferences based on allele frequency change over time. In contrast, analyses

that search for window-scale outliers for elevated allele frequency differentiation between

samples should be little-affected by low rates of spurious singleton variants.

**The historical fly genomes show signs of relatedness and inbreeding**

A preliminary analysis of pairwise genetic distances among the genomes indicated that some

specific pairs of individuals from within the early 1800s and 1933 Lund samples had much

greater similarity than expected.  For example, as seen in **Table S2**, the raw (Hamming)

distances among X chromosomes within and among modern European populations are

typically 0.1-0.14.  In contrast, there are 28 pairs of historical genomes with pairwise distances

below 0.08 (as low as 0.038 in one case).  These results suggested relatedness among some

historical flies, which should lead to substantial tracts of Identity By Descent (IBD; **Figure 3A**).

Because the inclusion of related individuals in estimates of allele frequency generates

artifactual non-independent sampling, these redundancies should be identified and

eliminated from downstream population genetic analyses.

**Figure 3:  Significant levels of identity-by-descent (IBD) were observed within and**

**between many historical fly genomes, reflecting inbreeding and relatedness.**

(A) Illustration of the genealogical relationships that may result in IBD due to relatedness

between individuals (color1) or inbreeding within individuals (color2).

(B) For the 1800s Sweden-Lund and Germany-Passau samples, the proportions of each

genome detected as IBD due to apparent inbreeding (given along the diagonal) were low or

zero.  Above the diagonal, the proportions of each pairwise genomic comparison detected

as IBD due to apparent relatedness varied within the Lund sample from zero to over 75% (the

latter indicating a close familial relationship), while the two Passau genomes showed low-

level relatedness IBD.

(C) The same quantities are given for the Lund 1933 population sample, revealing some

genomes with high levels of inbreeding, and a broad spectrum of relatedness among

genomes.

Since the probability of IBD detection is different for diploid and haploid sequences, males
(with a single X chromsome) are given in italics. Lower quality genomes treated as haploids
are shown in bold. IBD regions were masked in subsequent analyses.

We identified IBD regions by partitioning each chromosome arm into windows of
approximately 100 kb length, and for each pair of individuals from the same population
sample, testing whether pairwise distance in a given window was closer to simplified
expectations for unrelated sequences (*i.e.* nucleotide diversity) or else IBD (*i.e.* zero for two
haploid sequences, or intermediate values for IBD involving diploid sequences; see Methods).
In line with the chromosome-wide distances cited above, we inferred many IBD windows
spanning various lengths, up to whole chromosome arms (**Table S2; Table S3**), while **Figure**
**3B and C** summarize the proportion of autosomal windows with IBD between every pair of
genomes in the 1800s and 1933 samples. These indicated up to sibling-level relatedness
among some flies in both the 1800s and 1933 Swedish samples, and a much lower level of
relatedness between the two specimens from 1840s Germany. The 1933 specimens, which
are all mounted together, were likely collected at the same time, at a guess from leftover fruits
on the third-floor lunch room. The six Swedish flies from the early 1800s also all show
relatedness, with some pairs even at sibling-level IBD. The most plausible explanation would
be that in spite of museum records tentatively linking them with three different scientists, these
specimens were all from the same Lund collection event (see Discussion).

In addition to IBD between genomes, we found within-genome IBD due to inbreeding
– several samples from 1933 had long regions of chromosomes with depleted heterozygosity
(**Table S2; Table S4**). Two 1933 Lund samples had especially extensive regions of near-zero
heterozygosity (covering 45% and 36% of their genomes; **Figure 3C**), consistent with mating
between close relatives, and several other historical genomes had homozygosity across 20-
40% of specific chromosome arms. Most of the genomes, however, showed no evidence of
inbreeding (**Figure 3B, C**),

The pairwise and within-genome IBD indicates that many of the individuals collected
for the historical were closely related and that (at least among the 1933 Lund flies) inbreeding

was prevalent.  These observations contrast with the low levels of relatedness observed from

contemporary population samples, even when multiple fly lines founded from the same trap

site were analyzed (Lack et al. 2016).  The elevated IBD of the historical fly genomes could have

resulted from sampling methodology and/or from a lower population density of flies in earlier

eras (see Discussion).

Following IBD and inbreeding masking, we were left with 9-11 autosomal alleles for the

combined 1800s samples on average rather than 14, and 10-14 rather than 28 for the 1933

samples.  The average sample size of X chromosome alleles was then 8 for the 1800s set and

11 for 1933, rather than 12 and 20.  Based on the mosaic pattern of window masking due to

intergenomic and intragenomic IBD across individuals on each chromosome arm, post-

filtering sample size varied around these averages, and downstream analyses included sample

size thresholds for site inclusion (see Methods).

**Inversion frequencies appear to increase with time**

We then sought to examine how within-population genetic diversity and between-population

genetic differentiation have changed across more than two centuries.  We first examined

frequencies of known polymorphic inversions (specifically *(2L)t, (2R)Ns, (3L)P, (3R)K,* and

*(3R)Mo)*, since inversions are known to significantly impact genomic diversity in this species

(*e.g.* Corbett-Detig and Hartl 2012; Kapun et al. 2016).  We used a previously reported list of

inversion-associated SNPs (Kapun et al. 2016), a majority of which appeared to be inversion-

associated in our analysis as well (see Methods and **Table S5**).  Our 24 historical flies comprise

a total of 192 sampled autosomal arms (as well as 37 X chromosomes), yet we inferred that

only a single autosomal arm from the 1933 sample carried an inversion (being heterozygous

for *(2L)t*), whereas no inversions were detected in any 1800s genome (**Table S5**).  Inversion

frequencies were however, also somewhat low in the modern Lund data, with only *(2L)t* (at

13%) giving a non-zero frequency estimate.  The inferred frequency changes in *(2L)t* were

statistically significant when modern Lund was compared to all historical genomes (Fisher exact

test *P*=0.048), though not when Lund was compared to either historical sample separately

(*P*=0.226 vs. 1800s and *P*=0.203 vs. 1933).   These results hint at increasing inversion

frequencies through time, but further temporal sampling will be needed to evaluate the

previous suggestion of a recent African origin for most inversions now present in European *D.*

*melanogaster* (Corbett-Detig and Hartl 2012; Pool et al. 2012).

**Genomic diversity suggests transiently elevated genetic differentiation**

We then focused our SNP-based analysis of genetic diversity and differentiation on the X

chromosome, in order to minimize the influence of inversions, which are very rare on this

chromosome in European populations (*e.g.* Kapun et al. 2016). Patterns of nucleotide diversity

($\pi$) in Lund appeared to show a decline with time (**Figure 4A**), which could reflect the action

of genetic drift. However, we note that damage-induced errors may inflate diversity estimates,

especially for the 1800s sample (as potentially indicated by its slightly elevated $D_{xy}$ values). In

addition, the modern Lund sample represents a published pool-seq data set, analyzed via a

distinct pipeline including the masking of rare variants, which may deflate its diversity estimate.

**Figure 4: Genetic differences among historical and modern fly populations suggest**

**transient differentiation in the Lund population.**

(A) Genetic differentiation between population samples is summarized by $F_{ST}$ (below

diagonal, red heat map), and by the rate of pairwise sequence differences $D_{xy}$ (above

diagonal, blue heat map). Along the diagonal, the rate of pairwise sequence differences

within populations (nucleotide diversity) is given. Data reflects the X chromosome only (to

avoid effects of inversions), with down-sampling to four alleles per population at each site.

The Lund population appears to show decreasing diversity with time, subject to caveats

regarding historical genome quality and the differentially-processed Lund 2010s pool-seq

data. When Lund allele frequencies are compared to other European populations (via $F_{ST}$),

differentiation increases between the 19[th] and 20[th] century Lund samples, then decreases

again between the 20[th] and 21[st] century samples.

(B) Prinicple Components Analysis (PCA) results are shown for the same individually-

sequenced population samples included in (A), in terms of loadings from the first two principle

components. Here again, the Lund 1933 population shows elevated differentiation, whereas

genomes from the three 1800s locations cluster with modern European genomes. These

results reflect an analysis of X chromosome variation, with each population sample

downsampled to five individuals, with non-inbred historical genomes lacking close relatedness

chosen for this analysis.

Patterns of genetic differentiation (as indicated by $F_{ST}$) revealed a curious temporal

trajectory (**Figure 4A**). When Lund samples are compared to modern samples from around

Europe, between the 1800s and 1900s sampling points, Lund becomes more differentiated

from other populations. In contrast, between the 1900s and 2000s sampling points, Lund

became more similar to other modern European populations. Concordant patterns were

observed from Principal Components Analysis (PCA) based on X-linked variation (**Figure 4B**;

**Table S6**). Here, 1800s Lund genomes were seen to cluster with modern European genomes,

whereas the 1933 Lund genomes form a distinct cluster. This indication of transiently increased

local genetic differentiation in Lund could indicate a history in which genetic drift led to

increased allele frequency divergence from more southerly European populations, but

subsequently elevated migration led to genetic homogenization between European

populations (see Discussion).

We further investigated a simple model of the demographic history of Lund by using

population genetic simulations to fit single-population models of genetic drift. We varied the

effective population size ($N_e$) for a given time interval and compared the mean difference in

allele frequency at non-singleton SNPs between a given pair of time points from simulated

versus empirical data, using a Bayesian approach to identify the best-fitting $N_e$ for each time

interval, for each chromosome arm separately. In contrast to longer-term estimates of $N_e$ in

this species, which are often on the order of one million (e.g. Sprengelmeyer et al. 2020), our

point estimates of Lund $N_e$ were only 2,000-2,500 diploid individuals for the 1800s-1933 time

interval, and 2,400-3,200 for the 1933-2010s interval (**Table S7**). These values are of the same

order of magnitude as an estimate of 9,500 from a northeast United States population between

1975 and 2015 (Lange et al. 2022). Estimates from both studies share however, a key limitation

in assuming that genetic drift (along with random sampling variance) is responsible for all

observed frequency differences. To address how well this assumption holds, we compared $\pi$

from coalescent simulations based on our estimated $N_e$ values to those from the empirical data,

using a demographic model previously estimated for a French population (Sprengelmeyer et

al. 2020) as a starting point for the pre-1800s history. While this history allowed us to match

Lund ~1809 $\pi$ reasonably well, our low estimates of $N_e$ yielded simulated values of $\pi$ much

lower than those observed in our empirical data. For example, **Table S7** shows that for arm

3L, simulated $\pi$ for 1933 and 2010s Lund was less than half of observed $\pi$. These results

indicated that drift-only models did not provide an accurate approximation of the Lund

population's demographic history, and that true $N_e$ values were probably greater than our

estimates. Instead, migration may have played an important role in shifting allele frequencies

without decreasing $\pi$, as suggested above. Estimating a reasonable spatiotemporal

demographic model that incorporates both local population sizes and geographic patterns of

genetic structure and gene flow may require more extensive sampling of population genomic

data across space and time.

**Genome scans to detect recent shifts in SNP and structural variant frequencies**

One principal goal of this study is to identify instances of elevated genetic differentiation

between time points that may reflect the action of recent positive selection. Our SNP-based

genome scan focused on Population Branch Excess (PBE; Yassin et al. 2016), a statistic that

quantifies elevated allele frequency differentiation in a focal population compared to two other

reference/outgroup populations. PBE builds upon the $F_{ST}$-based Population Branch Statistic

(PBS; Yi et al. 2010) but is more tailored to detecting selection that is specific to the focal

population. To search for selection between the early 1800s and 1933, we defined the 1800s

samples as the focal population and included the Lund 1933 and Lund 2010s samples as

outgroups. To focus on the 1933 to 2010s interval, Lund 1933 was the focal population and

Lund 2010s and France were the outgroups. PBE was evaluated in diversity-scaled windows

averaging 4.6 kb in length. In the absence of a suitable demographic model to provide a null

hypothesis for the extent of genetic differentiation, we focused our analysis on the top 1% of

window PBE values. However, we emphasize that any outlier-based genome scan may entail

both false positives and false negatives. Our SNP-based PBE scans revealed notable outlier

peaks reflecting elevated allele frequency change for each time period (**Figure 5A, B**). Overall,

we identified 190 outlier regions for the 1800s-1933 time interval (referred to below as scan A)

and 173 for the 1933-2010s interval (scan B), with some of these regions incorporating multiple

outlier windows (**Table S8**). Since not all outliers may represent true positive targets of recent

natural selection, we focus here on loci that showed the most extreme frequency changes

across each century, referring to outlier regions based on their ranked maximal window PBE

values.

**Figure 5: Allele frequency changes across centuries reveal candidates for positive**

**selection.**

(A) Allele frequency change for ~5 kb genomic windows, focusing on the 1800s-1933 time

interval (scan A), as quantified by Population Branch Excess (PBE). Genes within top outlier

regions that are discussed in the text are indicated.

(B) PBE is shown for the same genomic windows, focused on the 1933-2010s time interval

(scan B).

(C) Window-level PBE is shown for a roughly 50 kb region focusing on the top outlier region

from scan A, with the position of the known selection target *ChKov1* indicated.

(D) Window-level PBE is shown for a roughly 100 kb region focusing on the top outlier region

from scan B, with the position of the known selection target *Cyp6g1* indicated.

We complemented this SNP-based approach with a parallel statistical methodology to

search for copy number variants (CNVs). We treated each population sample's mean read

depth within a given window (scaled relative to its chromosome arm average) as a quantitative

trait, and calculated $Q_{ST}$ (a quantitative trait analog of $F_{ST}$; Lande 1992; Spitze 1993) based on

variation in read depth within and between two population samples. We then used the

pairwise $Q_{ST}$ values among a trio of populations to calculate an analog of PBE (here referred to

as QPBE), which will be elevated when the focal population has a scaled mean window

sequencing depth that is much different in comparison to the two outgroup populations,

potentially indicating a CNV at contrasting frequency in the focal population. It is possible that

locus-specific heterogeneity in DNA degradation might also influence depth and thus QPBE

values involving historical samples. Therefore, we highlight only the top result from each time

interval (with further top-1% outliers given in **Table S9**), in part because both of these top

outliers involve loci with previously-detected structural variants. For the 1800s to 1933 period,

our top depth QPBE window contained only the gene *Tequila*, a serine protease involved in

insulin/glucose regulation, lifespan, and memory (Colomb et al. 2009; Huang et al. 2015), for

which polymorphic transposon insertions have created new introns (Chakraborty et al. 2018).

For the 1933 to 2010s interval, the top window contained only the gene *Tenascin accessory*,

involved in synapse organization (Hong et al. 2012; Mosca et al. 2012), for which polymorphic

transposons are associated with expression differences (Rech et al. 2022) and balancing

selection has been suggested (Croze et al. 2017). Otherwise, we refer to QPBE outliers only

where they intersect with the SNP-based PBE outliers discussed below.

**Top SNP-based outliers inform classic examples of positive selection**

In both time intervals, the top window PBE value was also associated with the widest outlier

region, which is consistent with relatively strong positive selection. For the 1800s-1933

interval, the top outlier (labelled A1) was a 31 kb region that included the gene *CHKov1* (**Figure**

**5C**), which is thought to encode an ecdysteroid kinase (Scanlan *et al*. 2020). This gene is known

to be associated with a protein-truncating transposon insertion under recent positive selection

in non-African *D. melanogaster* populations, which was found to correlate with resistance to an

organophosphate pesticide (Aminetzach *et al*. 2005). Transposon insertion and gene

duplication at the *CHKov1/CHKov2* locus has, however, also been found to correlate with

resistance to infection by the *Drosophila melanogaster* sigmavirus (DmelSV) (Magwire *et al*.

2011). Given the lack of widespread insecticide usage during this period, the timing of

selection indicated by our analysis rather favor the antiviral role of this locus as a potential

selective advantage. Measurements of the substitution rate in DmelSV indicate *D.*

*melanogaster* acquired this rhabdovirus around 200 years ago (Carpenter *et al*. 2007), which

predicts that this time interval should have included key stages in the evolution of resistance

to DmelSV. Consistent with this hypothesis, we also found the gene *refractory to Sigma P*

(*ref(2)p*) within outlier region A20. Ref2p activates the Toll pathway and has been shown to

confer resistance to DmelSV infections (Bangham *et al.* 2007).

*CHKov1* and *ref(2)p* are of particular interest because mutational variants associated

with viral resistance have been previously identified (Wayne et al. 1996; Magwire et al. 2011).

In *CHKov1*, several SNPs have alleles in linkage disequilibrium with a transposable element

insertion in the gene (Aminetzach et al. 2005) associated with viral resistance (Magwire et al.

2011). Based on seven such SNPs scored in our data, we inferred that the resistant haplotype

increased from 16.7% in the 1800s sample to 100% in the 1933 and 2015 Lund samples. The

occurrence of three resistance-associated haplotypes (carried in heterozygous form by two

early 1800s Lund flies, as well as the later Zealand sample) could indicate that selection on this

allele began prior to the collection of our 1800s flies. The fixation of the resistant haplotype in

the 1933 and 2010s Lund samples suggests a complete (or nearly complete) sweep, mirroring

findings from some but not all recently-collected *D. melanogaster* population samples

(Aminetzach et al. 2005).

The outlier PBE score for *ref(2)p* for the 1800s time interval appeared not to be driven

by the short, complex deletion identified by Wayne et al. (1996). Based on inspecting reads,

this variant appeared to exist in two early 1800s Lund flies and two 1933 Lund flies, implying

modest frequency change between temporal samples. The highest SNP-level PBE score at this

gene was at a non-synonymous variant over 1 kb downstream from the complex deletion

(2L:19544138, a threonine/serine polymorphism).

The top outlier for the 1933-2010s interval (denoted region B1) spanned 75 kb and

included a known target of insecticide resistance evolution, *Cytochrome P450 6g1* (*Cyp6g1*,

shown in **Figure 5D**). This gene is associated with resistance to dichlorodiphenyl-

trichloroethane (DDT) and other insecticides in *D. melanogaster*. As with *ChKov1*, there is prior

evidence for positive selection associated with transposon insertions and gene duplication

(Daborn et al. 2002; Chung 2007; Schmidt et al. 2010; Battlay et al. 2016). Here, the novel and

widespread deployment of DDT, introduced in 1944, provides a clear hypothesis for a selective

pressure driving the observed frequency changes at SNPs linked to the locus in question.

**A subset of top outliers show narrow peaks of genetic differentiation**

Both of the above outlier regions were relatively broad, and some of the other top outliers also showed less-broad plateaus of elevated genetic differentiation between temporal samples. In contrast, several other top outliers showed more narrowly localized signals of elevated genetic differentiation. At least in spatial analyses of local adaptation, broader intervals of genetic differentiation are more likely to be associated with selection favoring a single initial haplotype (resulting in a hard sweep), whereas narrow signals of frequency change may result from selection favoring multiple initial haplotypes, resulting in a "soft sweep" (da Silva Ribeiro et al. 2022). Narrow differentiation signals are of particular interest based on their potential to indicate not just a specific gene but often a small set of candidate variants which may include the target of selection. **Figure 6A-D** depicts four narrow SNP-level patterns of allele frequency change focused on the 1800s-1933 time interval, whereas **Figure 7A-D** depicts four such examples for the 1933-2010s interval (all of which were among the top 11 regions for window PBE for their time interval). Focusing on the earlier period, a few SNPs upstream of *hector* (*hec*; in region A3), a calcitonin-like neuropeptide receptor involved in male courtship behavior (Li et al. 2011), showed elevated PBE values (**Figure 6A**). In region A8, a cluster of SNPs within a 5′ UTR intron showed the highest PBE values at *rutabaga* (*rut*; **Figure 6B**), a calcium-responsive adenyl cyclase that influences learning, memory, lifespan, circadian rhythm, ethanol tolerance, and response to heat and oxidative stress (Levin et al. 1992; Tong et al. 2007; Donlea et al. 2009; Qin and Dubnau 2010; Scheunemann et al. 2012). We also detected a narrow set of SNPs within region A7 at the lysozyme genes *LysC* and *LysD* (**Figure 6C**), a locus for which a gene duplication and inversion polymorphisms have been reported (Chakraborty et al. 2018), and which was found to be strongly differentiated between Swedish and Italian *D. melanogaster* populations (Mateo et al. 2018). In contrast, no function is known for *CG17032* (in region A5), which also featured a cluster of high PBE SNPs (**Figure 6D**), the most extreme of which was a synonymous variant (at R5 position 3L:15953092). Among other notable 1800s-1933 outliers, the A2 region encompassed *sturkopf*, a lipid droplet-associated protein that regulates growth and stress response (Werthebach et al. 2019). A polymorphic duplication of

this gene has been associated with low genetic diversity (Cardoso-Moreira et al. 2016) and this

window was detected by our depth qPBE scan as well (**Table S9**).  Region A4 included *cabut*

(*cbt*), a transcription factor that regulates metabolic and circadian responses to nutrition

(Bartok et al. 2015).

**Figure 6:  Narrow SNP-level intervals of differentiation indicate potential targets of**

**natural selection from the 1800s-focused scan.**

PBE for each SNP is plotted across selected top outlier regions that showed narrow peaks of

differentiation.  These peaks occurred in or near:

(A) *hector* (hec) in region A3,

(B) *rutabaga* (rut) in region A8,

(C) *Lysozyme C* (*LysC*) and *Lysozyme D* (*LysD*) in region A7, and

(D) *CG17032* in region A5.

For the 1933-2010s interval, a window focused on *Adenosylhomocysteinase* (*Ahcy*; in

region B2) yielded a PBE value nearly as extreme as that observed at *Cyp6g1*, but with a far

narrower genomic scale (**Figure 7A**).  Curiously, the top SNP PBE value occurs within a protein-

coding exon, but at a synonymous variant.  *Ahcy* regulates methionine metabolism, histone

methylation, and lifespan (Parkhitko et al. 2016).  It is preferentially expressed in circadian-

regulating pacemaker neurons (Rivas et al. 2021), and its role in circadian regulation has been

confirmed in mice (Greco et al. 2020).  Despite the strength of this signal, *Ahcy* does not appear

to have been detected as a top selection candidate from previously published scans of

contemporary genomic variation.  The SNP with the highest site PBE in *Ahcy* is associated with

a synonymous variant (at R5 position 3L:15953092).  We note that within the next-highest

region (B3, not plotted), *Lysine demethylase 4B* (*Kdm4B*) also impacts histone methylation and

circadian rhythm (Shalaby et al. 2019).  For the narrow signal at region B8, the highest PBE

SNPs were observed at the upstream end of *retinal degeneration B* (*rdgB*; **Figure 7B**), involved

in phototransduction, olfaction, and phospholipid transport (Woodard et al. 1992; Milligan et

al. 1999; Gardner et al. 2012).  Notably, the narrow PBE signal at region B11 (**Figure 7C**)

centered on a gene that was recently detected as a top outlier in the 1975-2015 US temporal

population genomic study cited above (Lange et al. 2022); *Hexosaminidase 2* (*Hexo2*) encodes

a sperm-associated protein that may play a role in fertilization (Cattaneo et al. 2006; Intra et al.

2017).  Whereas another narrowly-focused top outlier (region B5) centered on the non-coding

RNA gene *CR43835* (**Figure 7D**), for which no function is known.

**Figure 7:  Narrow SNP-level intervals of differentiation indicate potential targets of**

**natural selection from the 1900s-focused scan.**

PBE for each SNP is plotted across selected top outlier regions that showed narrow peaks of

differentiation.  These peaks occurred in or near:

(A) *Adenosylhomocysteinase* (*Ahcy*, or *Ahcy13*) in region B2,

(B) *retinal degeneration B* (*rdgB*) in region B8,

(C) *Hexosaminidase 2* (*Hexo2*) in region B11, and

(D) *ncRNA-CR43835* in region B5.

As hinted by the exclusively X-linked examples shown in **Figure 7**, for the 1933-2010s

scan specifically, there was an abundance of X-linked loci among the top outliers: although the

X chromosome constitutes less than one fifth of the analyzed genome, 12 of the top 15 outlier

regions were X-linked (**Table S8**).  Explanations of this enrichment could include greater X

chromosome effects of sampling variance (although this effect should have been stronger for

the 1800s scan), genetic drift, or positive selection (potentially facilitated by X chromosome

hemizygosity; Charlesworth et al. 2018).  However, neither scan was meaningfully enriched for

X-linked loci if the full list of outliers was considered (with the X contributing 21.7% of outliers

for scan A and 18.6% for scan B).  The excess of top outliers specifically for scan B could indicate

that more of the X chromosome's selection in this interval was from hard sweeps (as suggested

by Harris and Garud 2023) and therefore more readily detectable from window genetic

differentiation (da Silva Ribeiro et al. 2022).  However, the narrower peaks of differentiation

shown in **Figure 7** could indicate a role of X-linked soft sweeps as well.

**Further insights into previously implicated selection targets**

**Further insights into previously implicated selection targets**

Three other top outliers for the 1933-2010s interval included genes previously implicated in

recent positive selection in European *D. melanogaster*. *Prosap* (in region B6) is thought to

encode a structural component of the postsynaptic density (Leibl and Featherstone 2008) and

affects circadian rhythm (Harbison et al. 2019). This gene was previously found to display

signals of parallel adaptation across cold-adapted *D. melanogaster* populations (Pool et al.

2017), and it harbors a polymorphic transposon insertion associated with signals of selection

(Rech et al. 2022). Region B7 centered on *phantom* (*phm*), which encodes a cytochrome P450

protein involved in ecdysteroid biosynthesis (Warren et al. 2004). Variation at this gene

revealed a selective sweep signal in a European *D. melanogaster* population (Orengo and

Aguadé 2007), and the protein displays rapid evolution between species (Good et al. 2014).

Further down the list, the region B21 contained *polyhomeotic-proximal* (*ph-p*), a

developmental and chromatin-modifying gene (*e.g.* Wang et al. 2006; Enderle et al. 2011),

which also displayed evidence of a selective sweep in a European population (Beisswanger

and Stephan 2008), associated with altered thermosensitivity of gene expression (Voigt et al.

2015). For these outliers, our findings offer insights regarding the timing of positive selection.

**Gene Ontology analysis reveals shifts in membranes and metabolism**

We also applied a Gene Ontology (GO) enrichment analysis, using the permutation approach

initially described in Langley et al. (2012), separately to the outliers detected from each

temporal scan. From the 1800s-1933 scan, some of the most enriched categories related to

cell shape and morphogenesis, including actin-mediated contraction and membrane

invagination (**Table S10**). Enrichment was also observed for membrane (and vesicle)

organization, compatible with the existence of population differences in membrane lipid

composition (Cooper et al. 2014) and this time interval coinciding with the species' earliest

known occupation of northern Europe (Keller 2007). Curiously, "response to DDT" was also

enriched, due to three cytochrome P450 outlier genes other than *Cyp6g1*, potentially

reflecting pre-insecticide roles of insecticide-related genes (as with the *ChKov* locus described

above). For the 1933-2010s interval, insecticide-related GO categories were not enriched

among outliers. Instead, metabolic processes dominated the top results (including amide,

fatty acid, lipid, and peptide metabolism), and categories related to cell division were enriched

as well (**Table S10**). As with membrane composition, metabolism is also known to vary

significantly between *D. melanogaster* populations from contrasting thermal environments

(*e.g.* Cogni et al. 2015; Brown et al. 2018).

**DISCUSSION**

Here, we report sequenced genomes from more than two dozen historical specimens of

*Drosophila melanogaster*. Comparing these historical genomes to modern population

genomic data, we have quantified allele frequency changes across roughly three thousand

generations of evolution, and these analyses have yielded a range of insights regarding recent

evolution in this model species.

**Illuminating Known and Novel Candidates for Recent Adaptive Evolution**

Above, we describe genomic scans for outliers reflecting elevated SNP frequency changes

across two separate time intervals (early 1800s to 1933, and 1933 to 2010s). While we report

outliers reflecting the upper 1% of genomic windows for our focal statistic, we focus above on

the most extreme of these outlier signals. This focus is motivated by uncertainty regarding the

actual number of loci under selection in each time interval, which we do not attempt to estimate

due to limitations in the data and in our ability to identify a realistic neutral model. Among the

top outliers discussed above, none appeared within the full list of outliers for the alternate time

period. This lack of outlier overlap could indicate that selective pressures differed between

eras, or that adaptive events tended to complete within a given time period rather than

spanning between them.

The strongest signals of positive selection revealed by our study added to our

understanding of known selection targets, particularly with regard to the timing of their

evolution. The top outlier from the 1933 to 2010s interval, *Cyp6g1*, was known to be the target

of a particularly strong selective sweep related to DDT resistance, and thus provides a strong

proof of concept for our ability to detect loci under recent selection. However, our inferred

timing of selection for our top 1800s to 1933 outlier, *CHKov1*, served to favor one adaptive

hypothesis (resistance to Sigma virus) over another (insecticide resistance).  A second outlier,

*ref(2)p*, reinforced the potential importance of Sigma virus as a selective pressure during our

earlier time interval.  As cited above, a significant proportion of our other top outliers also

centered on previously reported selection targets.  In addition to the above three outliers,

these included *LysC/LysD* and *sturkopf* for the 1800s to 1933 time period, and *Hexo2*, *ph-p*,

*phm*, and *Prosap* for the 1933 to 2010s interval.  The number of our top outliers identified by

previous studies is a testament to the depth of previous population genomic studies in *D.*

*melanogaster*, particularly for European populations.

And yet, the majority of top outlier regions contained no previously known selection

candidate.  A prime example is *Ahcy* from the 1930s to 2010s scan, which showed a striking

level of window frequency change comparable to *Cyp6g1*, in spite of the more narrowly

focused peak at *Ahcy*.  The role of *Ahcy* in circadian regulation connects it with other top

outliers, especially the functionally related *Kdm4B* (the third highest outlier in this scan after

*Cyp6g1* and *Ahcy*).  Both *Ahcy* and *Kdm4B* alter histone methylation, an epigenetic mechanism

known to regulate fly circadian rhythm (*e.g.* Shalaby et al. 2018).  In addition, *Prosap* from this

same 1933 to 2010s time interval, and *cbt* and *rut* from the 1800s to 1933 scan, also represent

top outliers with circadian functions.  The evolution of circadian behavior in high latitude *D.*

*melanogaster* populations is well known (*e.g.* Kyriacou et al. 2008; Svetec et al. 2015).  Changes

in day length help regulate reproductive diapause (Saunders et al. 1989), a key adaptation in

*D. melanogaster* populations from strongly seasonal climates (Schmidt et al. 2005; Erickson et

al. 2020).  The northerly location of Sweden entails pronounced seasonal variation in day

length, which could heighten local selective pressures on daylight-responsive behaviors.

While each of the above five genes have been functionally linked to circadian regulation, to

our knowledge none have been linked to the evolution of circadian behavior (although *Prosap*

does have documented signatures of local adaptation; Pool et al. 2017).  Hence, these genes

represent strong candidates for future investigations into the molecular basis of behavioral

evolution.

In theory, alleles promoting adaptation to the local environment should have been

adaptive in the earlier time period, and yet three of these five circadian genes were only found

as outliers in the later interval.  It is possible that each local population initially possessed only

a subset of adaptive variants, and others arrived later via migration.  Alternatively, in the 1800s

to 1933 interval, the frequencies of causative variants at some circadian genes may have still

been early in their rise, poising them to rise more quickly in the 1933 to 2010s interval.  Another

possibility is contingency – some circadian variants may have only become beneficial after

other adaptive changes had occurred due to epistatic interactions among circadian genes.

**Temporal analyses reveal the complexity of demographic history**

Analyses involving our museum fly genomes also yielded novel insights regarding the

demographic history of a local *D. melanogaster* population from Lund, Sweden.  We found that

between the early 1800s and 1933, Lund genomic diversity became more differentiated from

other modern European populations with respect to allele frequencies, and yet this pattern

reversed between the 1933 and 2010s, with Lund regaining similarity to other European

populations.  This unexpected pattern of transient local differentiation suggests that distinct

evolutionary forces had predominant influences on genomic diversity between these time

points.   As elaborated below, a primary hypothesis for this shifting pattern is that important

changes in population dynamics occurred between these eras.

The increased genetic differentiation of the early 1800s to 1933 period could reflect an

important influence of genetic drift, which could have occurred for multiple reasons.  This

interval appears to reflect the earliest days of *D. melanogaster* occupying northern Europe

(Keller 2007), perhaps as shifting human activities only just permitted the species to survive in

this region, and hence we might predict that initial local population sizes were low.  The climate

during this period was also colder than at present (**Figure 1A**), which may have also limited

local population sizes.  In agreement with this hypothesis, Zetterstedt commented in his

treatise on Scandinavian Diptera with regard to *D. melanogaster*, "*habitat in Svecia meridionali*

*& media rarius*", *i.e.* in Southern and Middle Sweden rare.  Furthermore, the species is

conspicuously absent from Carl Linnaeus' *Fauna Svecica* (Linnaeus 1746; 1761), nor mentioned

in Johann Christian Fabricius' *Genera insectorum* (Fabricius 1776).  While these factors indicate

that local population size was likely low in the early 1800s, we cannot exclude an important

influence of natural selection in the observed increase in genome-wide population

differentiation, as European fly populations adapted to novel environments.

The hypothesized low historical population sizes of *D. melanogaster* in Lund could also

contribute to the inferred prevalence of relatedness among individuals within each historical

population sample, whereas flies from modern population samples generally show much less

relatedness, even when multiple individuals or lines obtained from the same fly trap are

analyzed (*e.g.* Lack et al. 2016).  Furthermore, a notable fraction of flies from the 1933 sample

show evidence of inbreeding, indicating that their parents were related to some degree.  Such

observations should be more likely in a relatively small and isolated local population.  However,

the precise circumstances of these collections are not entirely known, and details of sampling

methodology (*e.g.* if multiple flies eclosing from the same piece of fruit were collected, or if

flies were raised in captivity for observation) may also contribute to the relatedness and

inbreeding we infer.  Such uncertainties represent an inherent limitation in studying specimens

that are accompanied by finite documentation.

In contrast, the genomic homogenization between Lund and other European regions

that occurred between 1933 and the 2010s could reflect increased migration through time, in

conjunction with reduced genetic drift in fly populations that may have grown with time.  Both

of those demographic changes might be predicted based on concurrent changes in human

activity during the 20th century.  This time period featured a five-fold increase in the human

population of Lund and the surrounding region, along with a warming climate (**Figure 1A**),

both of which may have been conducive to population growth in this human commensal insect

(and hence reduced drift), even as new challenges such as insecticides emerged.  In parallel,

increased human transportation, particularly the expanded shipping of fruit and other

commodities, would be expected to facilitate increased long-distance dispersal of *D.*

*melanogaster*.  While these hypothesized demographic changes could be sufficient to explain

the genetic homogenization of the 1933 to 2010s time period, here again, we cannot exclude

a meaningful role for natural selection (such as selective sweeps spreading between regions)

in contributing to reduced differentiation.

The shifting population history potentially implied by our results raises questions about

the typical consideration of demographic processes in population genetic studies. Usually,

data is only available from one or more modern populations, and by necessity, fairly simple

neutral models of population history are fit to the data. However, we find that when genomes

from multiple temporal samples are available, we see hints of important historical changes in

population size and gene flow that conventional demographic inference would be unlikely to

capture. How well simple demographic models can reflect key aspects of complex

demographic models, and to what degree they can recapitulate the effects of complex

demographies on genomic diversity, is largely unknown. Regardless, it is clear that

ancient/historical genomes have strong potential to inform our understanding of the histories

of natural populations, just as they have for our own species (*e.g.* Yang and Fu 2018; Wohns et

al. 2022) and for other taxa (*e.g.* van der Valk et al. 2019; Cohen et al. 2022).

**Implications of sequencing historical *D. melanogaster* specimens**

This project contributes to the rapidly growing field of historical genomics and the study of

temporal population genomics in natural populations. To our knowledge, our samples include

the oldest pinned insects from which genome sequences have been reported (extending the

temporal horizon of insect museomics from the late 1800s back to the early 1800s). Overall,

96% of our specimens yielded analyzable genomes, and 73% yielded genomes of relatively

high quality. These results are encouraging for future museomic studies of small invertebrates.

To our knowledge, *D. melanogaster* is the smallest animal from which historical

genomes have been obtained. Surprisingly, that small size may have even been an advantage.

Male *D. melanogaster* are smaller than females (roughly 1 mg vs. 1.8 mg wet weight; Hillesheim

and Stearns 1991), and yet all 11 males in our study yielded high quality genomes, whereas

only half of the 14 females gave comparable outcomes. It is possible that less voluminous

insects such as male *D. melanogaster* experience a faster rate of desiccation relative to decay,

which should improve the quality and quantity of available DNA.

The genomes of these museum specimens have added to our knowledge of not only

recent evolution in *D. melanogaster*, but also the specimens themselves.  Beyond confirming

the sex of each fly, we inferred from relatedness that the Swedish early 1800s flies, which had

been linked to three different scientists and two different localities, were actually from the same

time and place.  These results may reflect widespread sharing of specimens among

contemporary scientists, and/or imperfect record keeping through time.  Whether these flies

were collected by Zetterstedt (who received his doctoral degree from Lund University in 1808)

or by his mentor Fallén (who was primarily active before 1810) is difficult to say.  We note that

all of these Swedish early 1800s flies – including the one labelled Småland – share the same

type of pin (similar to the those used by Fallén).  Intriguingly, the lectotype specimen (also

labelled Småland) – which we were not granted permission to sequence – is, however, mounted

with a distinctly different pin. The lectotype may accordingly be one of Ljungh's flies, as

described in Zetterstedt's *Diptera Scandinaviae*, unlike the paralectotype that evidently was

caught in Lund, despite the label stating otherwise.  The exact collection date of the sequenced

flies is unknown, although it may have occurred before 1810 (see Introduction).  Given the

apparent acquisition of one of these flies by Ljungh, his death in 1828 puts a latest possible

bound on the collection date.  These circumstances imply that the flies examined here are the

oldest surviving specimens of *D. melanogaster*, predating those described by Meigen (1830).

The expanse of existing knowledge regarding the biology and genome of *D.*

*melanogaster* has greatly enhanced the insights we have been able to draw from our museum

fly genomes.  In turn, our results have implications for the extensive *Drosophila* research

community.  We have identified various loci that represent likely targets of adaptive evolution

within specific recent time intervals, and in some cases, these genes have been found to impact

traits relevant to known selective pressures in the recent history of *D. melanogaster* (*e.g.*

circadian regulation).  Hence, our study provides a wealth of candidates for functional evolution

that can be investigated in subsequent laboratory studies, through leverage of the molecular,

genetic, and transgenic tools available in this model system.  Although museum specimens for

this (or any) species are finite, further historical genomic studies in *D. melanogaster* have the

potential to greatly inform the spatial and temporal dynamics of adaptive evolution and

population history, broadening the range of insights that can be obtained from genomic

variation in this population genetics model species.

**METHODS:**

**Samples and DNA extraction**

Historical specimens of *Drosophila melanogaster* were sampled from the entomological

collections of Biological Museum (Lund, Sweden: MZLU), Swedish Museum of Natural History

(Stockholm, Sweden: NRM), and National History Museum Denmark (Copenhagen, Denmark:

NHMD) (**Table S1**).  DNA was extracted from the sampled specimens using the QIAamp DNA

Micro kit (Qiagen).  Given the fragility of these old samples, special attention was given to

preserve their integrity.  The samples were placed in a microtube without removing the pin. In

cases where the pin was too long to fit in the microtube, or in the case the specimen was not

mounted near the end, the pin was shortened.  The lysis buffer was carefully added to avoid

damaging the specimens due to surface tension of the buffer prior to the addition of the

Proteinase K.  The microtubes were incubated at 56°C for 2 hours without shaking.  The buffer

containing the DNA was transferred to a new microtube, and 1 mL of MilliQ water was added

to the tube containing the fly to wash away the buffer.  After 30 min the water was transferred

to a new microtube and stored at -20°C for future DNA extraction if needed.  Then the fly was

washed in successive ethanol solutions of increasing concentrations (1 mL, 50%, 70%, 80%,

99%), again paying extra attention to not harm the specimen.  And finally, 500 µl of acetone

was added to the specimen, which was then placed to dry in a suspended position.  The rest

of the DNA extraction followed the protocol, except for the final elution which was performed

in two steps.  In the first step, 50 µl of elution buffer was added to the columns, which were left

to incubate at room temperature for 15 min prior to centrifugation.  For the second step, 50 µl

of elution buffer was added and the columns were incubated at 70°C for 15 min prior to

centrifugation.

**Library Prep and Sequencing**

The quality of the extracted DNA was visually investigated on an agarose gel (1.5%). cThe

library prep followed mainly the protocol of Twort *et al.* (2021).  Briefly, the fragmented DNA

was blunt-end repaired with T4 Polynucleotide Kinase (New England Biolabs) and then purified

with the MinElute Purification Kit (Qiagen).   Next followed adaptor ligation, reaction

purification, and adapter fill in.  The resulting products were subsequently indexed with unique

dual indexes. Indexing PCR was performed in ten independent reactions to reduce

amplification bias. Each PCR reaction consisted of 12-18 cycles depending on the

concentration. Indexing PCR reactions were pooled prior to purification with Sera-Mag

SpeedBeads™ carboxylate-modified hydrophilic (Sigma-Aldrich).   An initial bead

concentration of 0.5X was used to remove long fragments that were likely contaminants of

fresh DNA.  Libraries were selected using a 1.8X bead concentration to size select the intended

library range of ~300 bp.  The resulting libraries were quantified and quality checked using the

Quanti-iT™ PicoGreen™ dsDNA assay and Bioanalyzer 2100 (Agilent technologies). The final

indexed libraries were pooled together prior to sequencing on an Illumina Novaseq 6000

platform at the Swedish National Genomics Institute (NGI) in Stockholm (2x150 bp, S4 flow

cell).   After observing that sequenced fragments only averaged about 50 bp in length,

subsequent sequencing of most specimens with 2x50 bp reads was performed by the

University of Wisconsin Biotechnology Center using an Illumina NovaSeq 6000 (**Table S1**).

**Genome Alignment and Variant Calling**

The procedures for alignments and variant calling used to process the museum genomes

largely followed the *Drosophila* Genome Nexus (DGN) pipeline as described in Lack et al.

(2015).  Using this pipeline with the release 5.57 *D. melanogaster* reference genome allowed

newly-assembled museum genomes to be compared against published data from modern

population samples of individual genomes.  Modern genomes from France and Egypt were

reported in the subsequent DGN publication (Lack et al. 2016).  A sample from the Netherlands

was also aligned in that study, and originally sequenced by Grenier et al. (2015).  Population

samples from Sweden (Stockholm) and Italy were sequenced by Mateo et al. (2018) and

assembled using the DGN pipeline for this study.   The (distinct) processing of published

pooled sequencing (pool-seq) data from Lund, Sweden (Kapun et al. 2021) is described in the

next section.

Following read trimming using fastp (v0.21.0, Chen et al. 2018), we followed the DGN

pipeline to map, filter, and call genotypes from the sequenced museum genomes. While the

pipeline is described in detail in the previously-cited papers, we note its key feature of two

rounds of mapping prior to genotyping. The initial mapping of reads to the reference genome

is performed with bwa -aln v.0.5.9 (Li and Durbin 2010) and stampy (Lunter and Goodson. This

step is followed by indel and variant-calling using GATK (McKenna et al 2010, Depristo et al

2011). The SNPs and indels were mapped onto the (release 5) reference genome, to create a

more robust target reference for a second round of alignment of the reads with bwa -aln and

stampy. It is only after this second realignment and a shift of coordinates around indels to

match the reference genome that genotyping for further downstream analyses was performed.

Diploid genotyping was initially performed at all sites, except for the X chromosomes

from male flies. The sex of the flies in the museum samples was not identified prior to

sequencing. It was identified by calculating the ratios of average read depth in the X

chromosome versus the autosomes, which is expected to be approximately 1:1 in females and

1:2 in males. All perl scripts referenced in this section and elsewhere were executed on perl

v5.34.0, all python scripts were executed on python 3.8.6 (except for the stampy aligner, which

was run on python 2.7.18-3 to ensure compatibility), and the R scripts were run using R v.4.2.2.

For those museum genomes with a total coverage of < 80% and average per-site read

depth of < 10, we concluded that diploid genotype calls could not be confidently made. These

were instead treated as "pseudo-haploid" with only a single nucleotide called per site;

UnifiedGenotyper (GATK, McKenna et al 2010) was run in haploid mode for all chromosomes

for these low-quality genomes. When more than a single base was identified at a (pseudo-

haploid) site, we retained the most frequent allele (or selected one at random in cases of a tie).

For diploid genotypes, we required at least 25% of reads to represent a minor allele, otherwise

the site was classified as homozygous and assigned the more frequent allele.

**Processing and Analysis of Pool-seq Data**

Published pool-seq data (Kapun et al. 2021) was obtained from modern three Lund population

samples: one from summer 2014 (SRR 5647735) and two from summer 2015 (SRR8439151,

8439156). Each was derived from 40 wild-caught males. For each of these samples, alignment

and post-mapping processing was performed using the same procedures as for individual

genomes in the first round of the DGN pipeline, and the three pool-seq sample alignments

were then merged to a single bam file. Because of the aggregate, multi-individual nature of

these sequence data, frequency estimation was performed using PoolSNP (Kapun et al. 2020).

We imposed an additional constraint of requiring there to be at least 5 instances of a minor

allele for a site to be identified as polymorphic when filtering the VCFs, in order to minimize

mislabeling read base-calling errors as SNPs. Additionally (and obviously), because individual

genomes cannot be identified from pool-seq data and thus could not be used to create a

revised reference genome, only a single round of alignments and genotyping was performed

on the pool-seq data.

Pool-seq data introduces several statistical sampling artifacts that are not an issue with

individual sequencing. Specifically, the number of reads contributed by each individual varies,

so that some genomes are over-represented and others are under-represented. To deal with

this, we calculate an effective allele (lineage) count (Ferretti et al. 2013), i.e. the number of $j$

unique lineages at a site given $n_r$ reads and $n_c$ the number of chromosomes in the sample pool

($n_c$ = 240 for autosomes, $n_c$ = 120 for X chromosomes because all of the 120 flies sampled in

the pool-seq study were male). The effective count $j$ follows a distribution based on Stirling

numbers of the second kind $S(n_r,j)$, i.e. the number of ways in which $n_r$ reads can be partitioned

into $j$ lineages. The probability of $j$ independent lineages in the sample is then:

922
$$P(j|, n_c, n_r) = S(n_r, j) \frac{n_c!}{(n_c - j)! \ n_c^{n_r}}$$

923 The effective number of distinct lineages (i.e. de facto allele count per site) was estimated from

924 the expectation of this distribution. We found a maximum pool size of 183 for the autosomes

925 (median = 145) and of 95 for ChrX (median = 70). Allele frequencies at each site were

926 multiplied by the pool size to generate the appropriate down-sampled counts for population

927 genetic analysis.

**Assessment of Genomic Data Quality**

The total number of reads and the average read length was assessed from each bam file using CollectInsertSizeMetrics from the Picard suite of tools. Based on the depth and coverage criteria given above, the low-quality genomes that were treated as pseudo-haploid in the analysis pipeline were 18SL4, 18SL5, 18SL6, 18SL12 (from 1800s), 19SL3, and 19SL7 (from 1933). One sample from 1933, 19SL2, was of such low quality that it was excluded from any further processing or analysis.

To determine whether low quality sequences contributed a significant number of false positive variants in each population sample, we tallied the number of "singletons" - instances where a given genome had an allele otherwise not observed among at least five total alleles. The fraction of singleton alleles that were A or T bases was calculated to determine if cytosine deamination accounted for many of the singletons, and whether this phenomenon was correlated with sample quality. Singleton minor alleles were excluded from certain downstream analyses, such as demographic inference, while a total allele count of at least four was required for all analyses unless otherwise indicated. We also calculated the Pearson correlations between read depth and coverage (as well as between read depth and the fraction of singletons contributed by each genome, and between the fraction of singletons and the A/T content) to quantify the association between these quality metrics.

**Identification of Identity By Descent**

Identity by descent (IBD) among certain genomes in both the 1800s and 1933 Lund samples was initially suspected based on highly variable pairwise differences among genomes. To identify specific regions of individual genomes with IBD, we began by partitioning chromosomes into "windows" defined by having 25,000 SNPs in an ancestral range Zambia population sample (Lack et al. 2015), averaging approximately 100 kb in length.

We then sought to account for genome-specific differences in genetic distances, such as due to sample quality. For each chromosome window of the $i$th genome, a background genetic distance $d_i$ is the median Hamming distance between the $i$th genome and unrelated

modern Lyon samples. This genome-specific individual distance estimate was used to obtain

a distance correction factor $m_i$ that accounts for its tendency to yield higher or lower distances

than others from its population sample. The baseline population-specific distances $D$ for 1800s

and 1933 Lund were estimated as the median pairwise distance among individuals with no

obvious IBD based on the Hamming distance matrix. Pairwise distances were compared to this

baseline, scaled with respect to the Lyon outgroup distance, as follows.

For two genomes $i,j$ collected in the same location and time point, the chromosome

window Hamming distance is $d(i,j)$, while $m_i$, $m_j$ respectively are the scaling factors specific to

sequences $i,j$ – calculated from $d_i, d_j$ divided by the median pairwise distance between all of the

Lund sequences (of a given time point) and the Lyon sequences.

A.  For the rescaled distance $D'(i,j) = d(i,j)m_i m_j$, a chromosome window is considered to be

IBD between two diploid sequences (autosomes or female X chromosomes) if $D'(i,j) <$

$(7/8)D$. The reasoning is that if one allele from each individual is IBD, then ¼ of pairwise

allelic comparisons will yield zero distance due to this IBD, and hence the overall

expected distance is ¾ of that from a non-IBD pair. For haploid to haploid (male X

chromosome comparison) the same rule with a factor of ½ is applied, while for haploid

to diploid (male to female X chromosome comparison) a factor of ¾ is used.

B.  When pairwise IBD regions are identified, this window is masked in one genome of the

pair. The masking is prioritized so that low-quality genomes are preferentially masked

when compared to a higher quality genome in order to minimize data loss.

C.  To avoid false negatives due to IBD regions covering partial windows, when a window

was masked, the proximate half of the "upstream" and the "downstream" window was

also masked. While this practice may mask some non-IBD regions, it errs on the

conservative side for avoiding oversampling IBD alleles (since IBD covering more than

half of a flanking window would be likely to satisfy the above masking criteria and cause

the full window to be masked).

In addition to IBD among genomes, there was also evidence for depleted

heterozygosity as a consequence of inbreeding in a subset of genomes from both the 1800s

and (especially) the 1933 Lund populations, *i.e.* intragenomic IBD. In non-inbred genomes,

the heterozygosity should approximately equal nucleotide diveristy ($\pi$). Therefore, to identify

enrichment of homozygosity in inbred genomes, we compared the observed heterozygosity

per window to the median pairwise distance among non-IBD pairs from the same population

in that window. If the within-window heterozygosity of a sample was less than $\pi/2$, the window

was flagged as IBD and a randomly selected allele of the two in the window was masked.

We included samples believed to originate from other sites and time points (18DZ5

from Zealand, Denmark and 18SL6, which was labelled as being from Småland, Sweden) in the

IBD analyses, because the latter showed low divergence when compared to some of the 1800s

Lund samples. There was unambiguous evidence of close relatedness between 18SL6 and

18SL10, suggesting that the former was in fact from Lund and thus could be grouped with this

population in subsequent analyses. In contrast, 18DZ5 showed roughly 1% of its genome with

apparent IBD with one Lund genome, and lesser levels with two others. As these could be

false positives due to conservative filtering criteria, we continued to treat this sample as being

from Zealand rather than Lund in demographic analyses based on location, while at the same

time masking that small region of the genome (18SL12) that showed low, IBD-like divergence

with respect to 18DZ5.

1002

**Analysis of Genetic Differentiation**

Temporal and spatial divergence among different populations (defined by time and location)

was assessed with analyses of population genetic distances. Our analysis focused on X

chromosome sequences (to the exclusion of low recombination regions defined by Pool et al.

2017) because of the rarity of sex chromosome inversions in *Drosophila*, and only considered

polymorphic sites without missing data in any of the analyzed samples.

To avoid distortions due to sample sizes, the number of alleles taken from each

population (*i.e.* the 1933 Lund and the modern European and Egyptian samples) was down-

sampled to match the 4 alleles from the IBD-masked 1800s Lund allele counts. This entailed

selecting 4 random alleles at each site for the 1933 Lund population, the 2010s Lund pool-seq

population, and the other modern populations, sampling independently at each analyzed site.

This down-sampled data was used to calculate nucleotide diversity within each population, as
well as between-population genetic distance ($D_{xy}$) and $F_{ST}$ (Reynolds et al. 1983).

Genetic differentiation among the populations was also characterized using principal
components analysis (PCA). Down-sampling of the Lund 1933 and modern samples was
necessary to avoid over-weighting larger samples. However, because individual haplotypes
were projected into the PCA space, the pool-seq Lund sample had to be excluded. To increase
the number of sites with no missing data, we sampled non-masked X chromosomes that had
no detectable pairwise IBD from the Hamming distance matrices from the 1800s and 1933
populations. Five such unrelated X chromosomes were identified from the 1800s samples:
18SL11, 18SL13, 18DZ5, 18GP9, and 18GP25, which were matched by the 5 unrelated 1933
Lund samples 19SL8, 19SL15, 19SL16, 19SL20, and 19SL23. Five X chromosomes were
sampled from each of the modern European and Egyptian outgroup populations as well, for a
total of 35 chromosomes (5 from 7 populations). For each of the 25,094 polymorphic sites with
no missing data across all 35 samples in the analyzed regions of the X, the frequency of the
minor allele at each site was calculated. An R-mode PCA was then used to compute the 35
eigenvectors of the correlation matrix.

1030

**Inversion Polymorphism**

To assay for inversions, we used the list of inversion-linked SNP alleles identified by
Kapun (2016, 2020) for the following common autosomal inversions: *In(2L)t*, *In(2R)Ns*, *In(3L)P*,
*In(3R)K*, *In(3R)Mo*, and *In(3R)P*. The frequency of each inversion was estimated from the median
frequency of the inversion-associated alleles among their respective loci. For the Lund pool-
seq data, the requirement of > 4 minor alleles per site to call minor alleles was relaxed to
identify low-frequency inversions, *i.e.* even singleton minor alleles were included.

We used the median rather than the mean allele frequency because some alleles may
be absent due to missing data. Additionally, not all of the alleles identified in Kapun (2016) as
inversion-associated appeared to be fully linked to inversions in our data set. There were
several instances where a small fraction of inversion-linked alleles are present in the museum
samples or modern Lund (*e.g.* for *In(3R)P*, 4 of the 20 inversion-linked alleles were present at

low frequency in the modern Lund pool-seq data), while the absence of inversion-linked alleles

at the majority of sites suggested that the inversion itself is absent. In such cases, the use of

mean allele frequency as a proxy for inversions would result in a non-zero estimated inversion

frequency that is much lower than the reciprocal of the number of chromosomes sampled in

the population. The estimated frequencies for each inversion in modern Lund versus the

historical populations were compared using Fisher exact tests on karyotype counts, using R.

1049

**Demographic Estimation**

Using allele frequency data from 1800s, 1933, and 2010s Lund, we estimated local effective

population size ($N_e$) for the 1809-1933 and 1933-2015 intervals by implementing a simple drift-

only neutral model of evolution (thus assuming no effect of migration, selection, or other

processes on allele frequency changes). To estimate $N_e$ for these two time intervals, we

compared the mean changes in allele frequencies across observed segregating sites to

changes in allele frequencies simulated under a Fisher-Wright (FW) model of random genetic

drift in discrete, non-overlapping generations. Specifically, we implemented a Bayesian

approach to optimize $N_e$ that minimizes the difference between observed $\overline{\Delta p}$, the mean

absolute value difference in allele frequency across sites between 1809 and 1933 (and

between 1933 and 2015) and the $\overline{\Delta p}$ simulated under a FW model of genetic drift.

We initialized the simulations by calculating the allele frequencies at polymorphic loci

at each time point in the Lund populations. Regions of low recombination, as defined by Pool

et al. (2017) were excluded, as were singleton variants and non-Lund 1800s genomes. The

sample frequencies $p_i$ in the 1809 population (or 1933 for the second time interval) at the $i$th

locus were used to approximate the initial allele frequencies $f_i$ in the population using Bayesian

estimation. The prior of $f_i$ was based on a mutation-drift equilibrium with a site mutation rate

of 5.21e-10 per generation (Huang et al. 2016) and a candidate haploid effective population

size $N*$. The posterior probabilities of $f_i$ were calculated  prior and the observed frequencies

at each site. The initial allele frequencies in each simulation replicate were generated by

drawing $n$ alleles equal to the number in each sample. Using a uniform prior instead resulted

in very similar distributions of $f_i$.

Population allele frequencies were initialized to $f_i(0)$ at simulation start time,

representing 1809 and 1933 for the first and second time intervals, respectively. The

frequencies in generation t+1 were generated by drawing a binomial sample ~ $bin(f_i(t), N^*)$ at

each site, where $f_i = x_i/N$ for $x_i$ count of what was the minor allele at $t = 0$. The model assumes

a sufficiently large gene pool for sampling with replacement and treating each locus as

statistically independent (*i.e.* under linkage equilibrium). This sampling is repeated for 1860

(non-overlapping) generations for 1809-1933 and 1230 generations for 1933-2015, *i.e.* for 15

generations per year (Turelli and Hoffmann 1995; Pool 2015). In the final (census) generation,

an autosomal allele count $n = 14$ ($n = 9$ for ChrX) was drawn to represent the 1933 Lund sample,

giving terminal sample allele frequencies $f_i^*(T)$ based on ~ $bin(f_i(T), n)$.

Much the same method was used for estimating the best fit $N_e$ for the 1933-2015 time

interval, with an important exception. Because the 2010s Lund samples contributed pool-seq

data, the additional sample variance that is introduced by sampling of both individuals and

reads must be incorporated into the simulation. In the terminal (census) generation, the

simulations implemented a two-step sampling model consistent with the nature of pool-seq

data. Namely, we sampled 240 alleles (corresponding to the 120 flies) in the census generation

~ $bin(f_i(T), 240)$. This created a pool of 240 alleles where the frequency of the (initial) minor

allele frequency was $f_i'(T)$, with an expectation equal to $f_i(T)$. To simulate the final sample set,

we drew $n$ alleles from this pool-seq subset of individuals (where $n$ is the effective number of

alleles per site sampled in the pool-seq data, with median values of 145 for autosomes and 70

for ChrX), *i.e.* ~ $bin(f_i'(T), n)$. This final draw gave terminal sample allele frequencies $f_i^*(T)$.

The absolute differences between final and initial sample allele frequencies in the

simulations $\Delta p_i = |f_i^*(T) - p_i(0)|$ were averaged across polymorphic sites, giving $\overline{\Delta p}$. We

optimized our estimate of $N_e$ by running the FW simulations over 1000 replicates over a range

of $N^* = \{1000...20000\}$ in increments of 1000 and selecting the value minimizing the difference

between simulated and observed $\overline{\Delta p}$. This estimate was further refined by considering $N^*$ in

five increments of 100 on either side of the coarse-grained optimum.

We evaluated the accuracy our estimates of effective population size ($N_1$, $N_2$) for the

time intervals 1809-1933 and 1933-2015 by comparing $\pi$ observed in the 1800s, 1933, and

2010s Lund samples to that of simulated populations of sizes $N_1$, $N_2$ at these time points. The

simulations were initiated by sampling $N_1$ alleles from a source European population based on

the demographic history of *D. melanogaster* inferred in Sprengelmeyer et al. (2020). The size

$N_1$ Lund sample then experienced genetic drift (with no gene flow or natural selection) for 1860

generations (scaled in units of $4N_e$) until 1933, at which point the population size was set to $N_2$

and the simulations continued for a further 1230 generations. This neutral demographic model

was implemented using the program ms (Hudson 1990), with the mutation rate cited above,

along with a crossing-over rate of 1.09e-8, a gene conversion rate of 6.25e-8, and mean gene

conversion tract length of 518 bp Comeron et al. (2012). We focused ms simulations on

chromosome arm 3L because of the favorable sample sizes and relative rarity of inversion

polymorphisms on this arm, and the availability of a parameterized demographic model for its

evolutionary history in Europe.

1113

**Population Branch Excess Scan for Positive Selection**

To identify regions of the genome that may contain sites that may have experienced directional

selection in the time between 1800s, 1900s, and modern samples, we used Population Branch

Excess (PBE; Yassin et al. 2016), a statistic that aims to identify loci with particularly strong allele

frequency differentiation in a focal population, in light of data from two outgroup populations.

PBE is based on the Population Branch Statistic (PBS; Yi et al. 2010), which essentially estimates

the length of the focal population's branch (in terms of allele frequency differentiation). PBE is

determined by a comparison of the focal population's observed PBS value to that expected

based on the lengths of the non-focal populations' branches at the same locus, along with the

average lengths of the focal and non-focal branches across loci (Yassin et al. 2016). Compared

to PBS, PBE is more focused on population-specific differentiation, and will not return a large

value at loci where all population branches are long (*e.g.* due to background selection or

positive selection in all populations).

These statistics are based on log-transformed window $F_{ST}$ values $d = -\log(1-F_{ST})$

between each pair of populations. Here as in other studies, window $F_{ST}$ was calculated by

summing the numerator terms and summing the denominator terms of the Reynolds et al. 1983

$F_{ST}$ estimator, and obtaining their ratio.  Chromosome windows were defined to contain 250

Zambia non-singleton SNPs, corresponding to approximately 4.6 kb windows on average.  We

performed a separate scan for each time interval.  To focus on the 1800s, we made Lund 1800s

the focal population, and used Lund 1933 and Lund 2010s as the non-focal populations.  To

focus on the 1900s, we made Lund 1933 the focal population, and used Lund 2010s and Lyon

(2010) as the outgroups.  Although the Lund 2010s sample comes from (distinctly-processed)

pool-seq data, the presence of one other individually-sequenced population sample among

the non-focal populations should prevent any artefactual differences between individual and

pool-seq data from registering as differentiation specific to the focal population.

Given that a viable demographic model was not available to provide a distribution of

PBE values under a null hypothesis, outlier PBE values were defined based on the upper 1%

quantile of the window distribution (which contained 240 windows genome-wide).  Because

strong selection can create linkage disequilibrium that extends across multiple windows, we

aggregated adjacent or near-adjacent PBE-significant windows (separated by no more than 2

non-outlier windows).  Although SNP-level genetic differentiation can sometimes allow the

detection of soft sweeps that do not register as outlier in window-level scans, our sample sizes

of museum genomes were not sufficient to enable this approach (da Silva Ribeiro et al. 2022).

Instead, we used site PBE as a heuristic for identifying sites and genes of interest within

window-based PBE outlier regions.

1149

**Gene Ontology Enrichment**

We performed a gene ontology (GO) enrichment analysis on genes located PBE outlier regions

from the 1800s- and 1900s-focused genome scans, in order to establish whether certain

functional categories (*e.g.* key metabolic or developmental pathways) were disproportionately

over-represented among genes that may differ between the historical and modern

populations.  The GO analysis methods followed the approach of Pool et al. (2012), which

involves randomly permuting the genomic locations of outlier regions, to account for variation

in gene length and the genomic clustering of functionally related genes.  Raw p-values were

computed from 10,000 random permutations, based on the proportion of reprlicates with an

equal or greater number of outlier regions associated with a given GO category as observed
in the empirical data.

1161

**Copy Number Variation**

To identify potential copy number variation (CNV) with frequency differences among

populations, we adopted an approach analogous to the one used above for SNP-based branch

statistics. We searched for CNV differentiation among samples by comparing the mean read

depth per genomic window, on the assumption that duplication (or deletion) present at

elevated frequency in a given population sample will result in higher (or lower) depth of

coverage, all else being equal.

Because these analyses compare read depths in museum samples with short fragments

to read depths in modern samples with much longer fragments, we used trimmomatic (Bolger

2014) to reduce the read length from the modern Lyon and Stockholm samples to 50 bp. We

then reran the relevant portions of the DGN assembly pipeline, while treating the truncated

reads as unpaired in the mapping. We used the same genomic windows as in the SNP PBE

analysis, and for each population, we calculated the mean read depth for each window across

all sites in that interval. Because genomes vary in their mean depth of coverage, we corrected

for such differences by dividing the mean read depth of each window by the mean depth over

the entire chromosome arm. These rescaled window read depths were treated as a

quantitative trait. The quantitative trait analog to the fixation index $Q_{ST}$ (Lande 1992, Whitlock

1999), was computed for read depths among populations based on variances in read depth,

as $Q_{ST} = \sigma_{GB}^2/(\sigma_{GB}^2 + 2\sigma_{GW}^2)$, where $\sigma_{GW}^2$ is the within-population genetic trait variance and $\sigma_{GB}^2$ is

the among-sample genetic trait variance.

By further analogy with allele frequency-based branch statistics, we calculated the

quantitative trait equivalents of PBS and PBE using $d = $ -log($1-Q_{ST}$) as the $Q_{ST}$-based distance

metric. To distinguish these trait-based branch statistics from $F_{ST}$-based measures, we refer to

them as QPBS and QPBE. These branch statistics were calculated for the 1800s population

using 1933 Lund and modern Lyon as outgroups. For 1933 as the focal population, Lyon and

Stockholm were used as the modern outgroups (since this approach is not applicable to pool-

seq data such as the Lund 2010s sample).  As with SNP-based PBE, we identified outlier

windows in the upper 1% quantiles, and we aggregated these into regions by merging outlier

windows separated by no more than one non-outlier window.

1191

1192 **Data Availability Statement:**

1196

**Acknowledgments:**

We wish to thank kind assistance with obtaining samples from Biologiska Museet (Lund,

Sweden) and Naturhistoriska Museet (Stockholm, Sweden), and in particular Niklas Whalberg

and Rune Bygebjerg.  We also thank other members of our labs for providing valuable

suggestions as well as relevant data and code from previous projects.

Funding was provided from Vetenskapsrådet (2019-03536) and the Max Planck Center on Next

Generation Insect Chemical Ecology to MCS, and from NIH NIGMS award 144 AAH7544 to

JEP.

1205

1206 **Author contributions:**

1207 HRG and MCS collected data.  MS, JDL, HRG, and JEP analyzed data.  MS, JEP, and MCS wrote

1208 the manuscript.

1209

1210 **Declaration of interests:**

1211 All authors declare no financial interests.

1212

**REFERENCES:**

Aminentzach YT, Macpherson JM, Petrov DA. 2005. Pesticide resistance via transposition-mediated adaptive gene truncation in *Drosophila*. Science 209: 764-767.

Bangham J, Obbard DJ, Kim KW, Haddrill PR, Jiggins FM. 2007. The age and evolution of an antiviral resistance mutation in *Drosophila melanogaster*. Proceedings of the Royal Society B: Biological Sciences 274:2027-2034.

Bartok O, Teesalu M, Ashwall-Fluss R, Pandey V, Hanan M, Rovenko BM, Poukkula M, Havula E, Moussaieff A, Vodala S et al. 2015. The transcription factor Cabut coordinates energy metabolism and the circadian clock in response to sugar sensing. EMBO Journal 34: 1538-1553.

Battlay P, Schmidt JM, Fournier-Level A, Robin C. 2016. Genomic and transcriptomic association identify a new insecticide resistance phenotype for the selective sweep at the *Cyp6g1* locus of *Drosophila melanogaster*. G3 6:2573-2481.

Beisswanger S, Stephan W. 2008. Evidence that strong positive selection drives neofunctionalization in the tandomly duplicated polyhomeotic genes in *Drosophila*. Proceedings of the National Academy of Sciences USA 105:5447-5452.

Bergland AO, Behrman EL, O'Brien KR, Schmidt PS, Petrov DA. 2014. Genomic evidence of rapid and stable adaptive oscillations over seasonal time scales in *Drosophila*. PLoS Genetics 10:e1004775.

Bi KE, Linderoth T, Singhal S, Vanderpool D, Patton JL, Nielsen R, Moritz C, Good JM. 2019. Temporal genomic contrasts reveal rapid evolutionary responses in an alpine mammal during recent climate change. PLoS Genetics 15:e1008119.

1242

1243 Brown EB, Torres J, Bennick RA, Rozzo V, Kerbs A, DiAngelo JR, Keen AC. 2018. Variation in

1244 sleep and metabolic function is associated with latitude and average temperature in

1245 *Drosophila melanogaster*. Ecology and Evolution 8:4084-4097.

1246

1247 Card DC, Shapiro B, Giribet G, Moritz C, Edwards SV. 2021. Museum genomics. Annual Review

1248 of Genetics 55:633-659.

1249

1250 Cardosa-Moreira M, Arguello JR, S. Gottipati S, Harshman LG, Grenier JK, Clark AG. 2016.

1251 Evidence for the fixation of gene duplications by positive selection in *Drosophila*. Genome

1252 Research 26:787-798.

1253

1254 Carpenter JA, Obbard DJ, Maside X, Jiggins FM. 2007. The recent spread of a vertically

1255 transmitted virus through populations of *Drosophila melanogaster*. Molecular Ecology

1256 16:3947-3954.

1257

1258 Cattaneo F, Pasini ME, Intra J, Matsumoto M, Briani F, Hoshi M, Perotti ME. 2006. Identification

1259 and expression analysis of *Drosophila melanogaster* genes encoding beta-hexosaminidases of

1260 the sperm plasma membrane. Glycobiology 16:786-800.

1261

1262 Chakraborty M, VanKuren NW, Zhao R, Zhang X, Kalsow S, Emerson JJ. 2018. Hidden genetic

1263 variation shapes the structure of functional elements in *Drosophila*. Nature Genetics 50:20-25.

1264

1265 Charlesworth B, Campos JL, Jackson BC 2018. Faster-X evolution: theory and evidence from

1266 *Drosophila*. Molecular Ecology 27:3753-3771.

1267

1268 Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.

1269 Bioinformatics 34:i884-i890.

1270

1271    Chung H, Bogwitz MR, McCart C, Andrianopoulos A, Ffrench-Constant RH, Batterham P,

1272    Daborn PJ. 2007. Cis-regulatory elements in the accord retrotransposon result in tissue-

1273    specific expression of the *Drosophila melanogaster* insecticide resistance gene *Cyp6g1*.

1274    Genetics 175:1071-1077.

1275

1276    Cogni R, Kuczynski K, Lavington E, Koury S, Behrman EL, O'Brien KR, Schmidt PS, Eanes WF.

1277    2015. Variation in *Drosophila melanogaster* central metabolic genes appears driven by natural

1278    selection both within and between populations. Proceedings of the Royal Society of London B

1279    282:20142688.

1280

1281    Cohen ZP, François O, Schoville SD. 2022. Museum genomics of an agricultural super-pest, the

1282    Colorado Potato Beetle, Leptinotarsa decemlineata (Chrysomelidae), provides evidence of

1283    adaptation from standing variation. Integrative and Comparative Biology 62:1827-1837.

1284

1285    Colomb J, Kaiser L, Chabaud M-A, Preat T 2009. Parametric and genetic analysis of *Drosophila*

1286    appetitive long-term memory and sugar motivation. Genes Brain and Behavior 8:407-415.

1287

1288    Comeron JM, Ratnappan R, Bailin S. 2012. The many landscapes of recombination in

1289    *Drosophila melanogaster*. PloS Genetics 8(10):e1002905.

1290

1291    Cong Q, Shen J, Zhang J, Li W, Kinch LN, Calhoun JV, Warren AD, Grishin NV. 2021. Genomics

1292    reveals the origins of historical specimens. Molecular Biology and Evolution 38:2166-2176.

1293

1294    Cooper BS, Hammad LA, Montooth KL. 2014. Thermal adaptation of cellular membranes in

1295    natural populations of *Drosophila melanogaster*. Functional Ecology 28:886-894.

1296

1297    Corbett-Detig RB and Hartl DL 2012. Population genomics of inversion polymorphisms in

1298    *Drosophila melanogaster*. PLoS Genetis 8:e1003056.

1299

1300    Croze M, Wollstein A, Bozicevic V, Zivkovic D, Stephan W, Hutter S. 2017. A genome-wide scan

1301    for genes under balancing selection in *Drosophila melanogaster*. BMC Evolutionary Biology

1302    17:15

1303

1304    Dabney J, Meyer M, Paabo S. 2013. Ancient DNA damage. Cold Spring Harbor Perspectives

1305    in Biology 5:a012567.

1306

1307    Daborn PJ, Yen JL, Bogwitz MR, Le Goff G, Feil E, Jeffers S, Tijet N, Perry T, Heckel D, Batterham

1308    P et al. 2002. A single p450 allele associated with insecticide resistance in *Drosophila*. Science

1309    297:2253-2256.

1310

1311    da Silva Ribeiro T, Galvan JA, Pool JE. 2022. Maximum SNP $F_{ST}$ outperforms full-window

1312    statistics for detecting soft sweeps in local adaptation. Genome Biology and Evolution

1313    14:evac143

1314

1315    Depristo MA, Banks E, Poplin R, Garimella KV, Maguire JR, Hartl C, Philippakis AA, del Angel

1316    G, Rivas MA, Hanna M et al. 2011. A framework for variation discovery and genotyping using

1317    next-generation DNA sequencing data. Nature Genetics 43:491-498.

1318

1319    Donlea JM, Ramanan N, Shaw PJ. 2009. Use-dependent plasticity in clock neurons regulates

1320    sleep need in *Drosophila*. Science 324:105-108.

1321

1322    Enderle D, Beisel C, Stadler MB, Gerstung M, Ahtri P, Paro R. 2011. Polycomb preferentially

1323    targets stalled promoters of coding and noncoding transcripts. Genome Research 21:216-226.

1324

1325    Erickson PA, Weller CA, Song DY, Bangerter AS, Schmidt P, Bergland AO. 2020. Unique

1326    genetic signatures of local adaptation over space and time for diapause, an ecologically

1327    relevant complex trait, in *Drosophila melanogaster*. PLoS Genetics 16:e1009110.

1328

1329    Ferretti L, Ramos-Onsins SE, Perez-Enciso M. 2013. Population genomics from pool

1330    sequencing. Molecular Ecology 22:5561-5576.

1331

1332    Fabricius JC. 1776. Genera insectorum: eorumque characteres naturales secundam numerum,

1333    figuram, situm et proportionem, omnium partium oris adiecta mantissa specierum nuper

1334    detectarum. Litteris MF Bartschii.

1335

1336    Garner K, Hunt AN, Koster G, Somerharju P, Groves E, Li M, Raghu P, Holic R, Cockcroft S. 2012.

1337    Phosphatidylinositol transfer protein, cytoplasmic 1 (PITPNC1) binds and transfers

1338    phosphatidic acid. Journal of Biological Chemistry 287: 32263-32276.

1339

1340    Good RT, Gramzow L, Battlay P, Sztal T, Batterham P, Robin C. 2014. The molecular evolution

1341    of cytochrome P450 genes within and between *Drosophila* species. Genome Biology and

1342    Evolution 6:1118-1134.

1343

1344    Greco CM, Cervantes M, Fustin JM, Ito K, Ceglia N, Samad M, Shi J, Koronowski KB, Forne I,

1345    Ranjit S, Gaucher J. 2020. S-adenosyl-l-homocysteine hydrolase links methionine metabolism

1346    to the circadian clock and chromatin remodeling. Science Advances 16:eabc5629.

1347

1348    Grewe F, Kronforst MR, Pierce NE, Moreau CS. 2021. Museum genomics reveals the Xerces

1349    blue butterfly (*Glaucopsyche xerces*) was a distinct species driven to extinction. Biology Letters

1350    17:20210123.

1351

1352    Harbison ST, Kumar S, Huang W, McCoy LJ, Smith KR, Mackay TFC .2019. Genome-wide

1353    association study of circadian behavior in *Drosophila melanogaster*. Behavioral Genetics

1354    49:60-82.

1355

1356    Harris M, Garud NR. 2023. Enrichment of hard sweeps on the X chromosome in *Drosophila*

1357    *melanogaster*. Molecular Biology and Evolution 40:msac268

1358

1359 Hillesheim E, Stearns SC. 1991. The responses of *Drosophila melanogaster* to artificial selection

1360 on body weight and its phenotypic plasticity in two larval food environments. Evolution

1361 45:1909-1923.

1362

1363 Hong W, Mosca TJ, Luo L. 2012 Teneurins instruct synaptic partner matching in an olfactory

1364 map. Nature 484:201-207.

1365

1366 Huang CW, Wang H-D, Bai H, Wu M-S, Yen J-H, Tatar M, Fu T-F, Wang P-Y. 2015. Tequila

1367 regulates insulin-like signaling and extends lifespan in *Drosophila melanogaster*. Journal of

1368 Gerontology A Biological and Medical Sciences 70:1461-1469

1369

1370 Huang W, Lyman RF, Lyman RA, Carbone MA, Harbison ST, Magwire MM, Mackay TFC et al.

1371 2016. Spontaneous mutations and the origin and maintenance of quantitative genetic

1372 variation. eLife 5:e14625

1373

1374 Intra J, Veltri C, De Caro D, Perotti ME, Pasini ME. 2017. In vitro evidence for the participation

1375 of *Drosophila melanogaster* sperm Beta-N-actylglucasaminidases in the interactions with

1376 glycans carrying terminal N-acetylglucosamine residues on the egg's envelopes. Archives of

1377 Insect Biochemistry and Physiology 96:arch21403

1378

1379 Jouganous J, Long W, Ragsdale AP, Gravel S. 2017. Inferring the joint demographic history of

1380 multiple populations: beyond the diffusion approximation. Genetics 206:1549-1567.

1381

1382 Kapun M, Fabian DK, Goudet J, Flatt T. 2016. Genomic evidence for adaptive inversion clines

1383 in *Drosophila melanogaster*. Molecular Biology and Evolution 33: 1317-1336.

1384

1385 Kapun M, Barron MG, Staubach F, Obbard DJ, Wiberg AW, Vieira J, Goubert C, Rota-Stabelli

1386 O, Kankare M, Bogaerts-Marquez M, et al. 2020. Genomic analysis of European *Drosophila*

1387 *melanogaster* population reveals longitudinal structure, continent-wide selection, and
1388 previously unknown DNA viruses. Molecular Biology and Evolution 37:2661-2678

1389

1390 Kapun M, Nunez JCB, Bogaerts-Marequz M, Murga-Moreno J, Paris M, Outten J, Coronado-
1391 Zamora M, Tern C, Rota-Stabelli O, Garcia Guerreiro MP, et al. 2021. *Drosophila* evolution over
1392 space and time (DEST): a new population genomics resource. Molecular Biology and Evolution
1393 38(12):5782-5805.

1394

1395 Keller A. 2007. *Drosophila melanogaster*'s history as a human commensal. Current Biology 17:
1396 R77-81.

1397

1398 Korlević P, McAlister E, Mayho M, Makunin A, Flicek P, Lawniczak MK. 2021. A Minimally
1399 Morphologically Destructive Approach for DNA Retrieval and Whole-Genome Shotgun
1400 Sequencing of Pinned Historic Dipteran Vector Species. Genome Biology and Evolution
1401 13:evab226.

1402

1403 Kyriacou CP, Peixoto AA, Sandrelli F, Costa R, Tauber E. 2008. Clines in clock genes: fine-tuning
1404 circadian rhythms to the environment. Trends in Genetics 24:124-132.

1405

1406 Lack JB, Cardeno CM, Crepeau MW, Taylor W, Corbett-Degit RB, Stevens KA, Langley CH, Pool
1407 JE. 2015. The Drosophila Genome Nexus: a population genomic resource of 623 *Drosophila*
1408 *melanogaster* genomes, including 197 from a single ancestral range population. Genetics
1409 199:1229-1241

1410

1411 Lack JB, Lange JD, Tang AD, Corbett-Detig RB, Pool JE. 2016. A thousand fly genomes: an
1412 expanded *Drosophila* Genome Nexus. Molecular Biology and Evolution 33:3308-3313.

1413

1414 Lande R 1992. Neutral theory of quantative genetic variance in an island model with local
1415 extinction and colonization. Evolution 46:381-389.

1416

1417  Lange JD, Hastide H, Lack JB, Pool JE. 2022. A population genomic assessment of three

1418  decades of evolution in a *Drosophila* population. Molecular Biology and Evolution 39:msab368

1419

1420  Langley CH, Stevens S, Cardeno C, Lee YCG, Schrider DR, Pool JE, Langley SA, Suarez C,

1421  Corbett-Detig RB, Kolaczkowski B et al. 2012. Genomic variation in natural populations of

1422  *Drosophila melanogaster*. Genetics 192: 533-598.

1423

1424  Levin LR, Han PL, Huang PM, Feinstein PG, Davis RL Reed RR. 1992. The *Drosophila* learning

1425  and memory gene *rutabaga* encodes a Ca2+/Calmodulin-responsive adenylyl cyclase. Cell

1426  68:479-489.

1427

1428  Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler tranform.

1429  Bioinformatics 25:1754-1760.

1430

1431  Li Y, Hoxha V, Lama C, Dien BH, Vo CN, Dauwalder B. 2011. The hector G-protein coupled

1432  receptor is required in a subset of fruitless neurons for male courtship behavior. PLoS One

1433  6:e28269

1434

1435  Liebl FL, Featherstone DE. 2008. Identification and investigation of *Drosophila* postsynaptic

1436  density homologs. Bioinformatics and Biology Insights, 2:375-388

1437

1438  Linnaeus C. 1746. Fauna Svecica sistens animalia Sveciae regni: qvadrupedia, Aves, Amphibia,

1439  Pisces, Insecta, Vermes, distributa per classes and ordines, genera and species. Cum

1440  differentiis specierum, synonymis auctorum, nominibus incolarum, locis habitationum,

1441  descriptionibus Insectorum. Laurentius Salvius, Stockholm.

1442

1443  Linnaeus C. 1761. Fauna Svecica sistens animalia Sveciae regni: qvadrupedia, Aves, Amphibia,

1444  Pisces, Insecta, Vermes, distributa per classes and ordines, genera and species. Cum

1445 differentiis specierum, synonymis auctorum, nominibus incolarum, locis habitationum,

1446 descriptionibus Insectorum (2nd edition). Laurentius Salvius, Stockholm.

1447

1448 Lunter G, Goodson M. 2011. Stampy: a statistical algorithm for sensitive and fast mapping of

1449 Illumina sequence reads. Genome Research 21: 936-939.

1450

1451 Machado HE, Bergland AO, Taylor R, Tilk S, Behrman E, Dyer K, Fabian DK, Flatt T, González J,

1452 Karasov TL, Kim B, et al. 2021. Broad geographic sampling reveals the shared basis and

1453 environmental correlates of seasonal adaptation in *Drosophila*. eLife 10:e67577.

1454

1455 Magwire MM, Bayer F, Webster CL, Can C, Jiggins FM. 2011. Successive increases in the

1456 resistance of *Drosophila* to viral infection through a transposon insertion followed by

1457 duplication. PLoS Genetics 7:e1002337.

1458

1459 Mallick S, Li H, Lipson M, Gymrek M, Racimo F, Zhao M, Chennagiri N, Nordenfelt S, Tandon

1460 A, Skoglund P et al. 2016. The Simons Genome Diversity Project: 300 genomes from 142

1461 diverse populations. Nature 538:201-206.

1462

1463 Mateo L, Rech GE, Gonzalez J. 2018. Genome-wide patterns of local adaptation in Western

1464 European *Drosophila melanogaster* natural populations. Scientific Reports 8:16143

1465

1466 McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler

1467 D, Gabriel S, Daly M, DePristo M. 2010. The genome analysis toolkit: a mapreduce framework

1468 for analyzing next-generation DNA sequencing data. Genome Research 20: 1297-303.

1469

1470 Meigen JW. 1830. Systematische Beschreibung der Bekannten Europaischen Zweiflugeligen

1471 Insekten, Vol. 6. Theil Schulze, Vienna.

1472

1473    Mikheyev AS, Tin MM, Arora J, Seeley TD. 2015. Museum samples reveal rapid evolution by

1474    wild honey bees exposed to a novel parasite. Nature Communications. 6:7991.

1475

1476    Mikheyev AS, Zwick A, Magrath MJ, Grau ML, Qiu L, Su YN, Yeates D. 2017. Museum genomics

1477    confirms that the Lord Howe Island stick insect survived extinction. Current Biology 27:3157-

1478    3161.

1479

1480    Milligan SC, Alb JG, Elaine RB, Bankaitis VA, Hyde DR. 1997. The phosphatidylinositol transfer

1481    protein domain of *Drosophila* retinal degeneration B protein is essential for photoreceptor cell

1482    survival and recovery from light stimulation. Journal of Cell Biology 139: 351-363.

1483

1484    Mosca TJ, Hong W, Dani VS, Favaloro V, Luo L. 2012. Trans-synaptic teneurin signalling in

1485    neuromuscular synapse organization and target choice. Nature 484:237-241.

1486

1487    Orengo DJ, Aguade M. 2007. Genome scans of variation and adaptive change: extended

1488    analysis of a candidate locus close to the *phantom* gene region in *Drosophila melanogaster*.

1489    Molecular Biology and Evolution 24:1122-1129.

1490

1491    Parejo M, Wragg D, Henriques D, Charriere JD, Estonba A. 2020. Digging into the genomic

1492    past of Swiss honey bees by whole-genome sequencing museum specimens. Genome Biology

1493    and Evolution. 12:2535-2551.

1494

1495    Parkhito AA, Binari R, Zhang N, Asara JM, Demotis F, Perrimon N. 2016. Tissue-specific down-

1496    regulation of S-adenosyl-homocysteine via suppression of dAhcyL1/dAhcyL2 extends health

1497    span and life span in *Drosophila*. Genes and Development 30:1409-1422.

1498

1499    Pool JE, Corbett-Detig RB, Sugino RP, Stevens KA, Cardeno CM, Crepeau MW, Duchen P,

1500    Emerson JJ, Saelao P, Begun DJ, Langley CH. 2012. Population genomics of sub-Saharan

*Drosophila melanogaster*: African diversity and non-African admixture. PLoS Genetics 8:e1003080.

Pool JE. 2015. The mosaic ancestry of the *Drosophila* Genetic Reference Panel and the *D. melanogaster* reference genome reveals a network of epistatic fitness interactions. Molecular Biology and Evolution 32:3236-3251.

Pool JE, Braun DT, Lack JB. 2017. Parallel evolution of cold tolerance within *Drosophila melanogaster*. Molecular Biology and Evolution 34:349-360.

Raxworthy CJ, Smith BT 2021. Mining museums for historical DNA: advances and challenges in museomics. Trends in Ecology and Evolution 36:1049-1060.

Rech GE, Radio S, Guirao-Rico S, Aguilera L, Horvath V, Green L, Lindstadt H, Jamilloux V, Quesneville H, Gonzales J, et al. 2022. Population-scale short read sequencing uncovers transposable elements associated with gene expression variation and adaptive signatures in *Drosophila*. Nature Communications 13:1948.

Reynolds J, Weir BS, CC Cockerham CC. 1983. Estimation of the coancestry coefficient: basis for a short-term genetic distance. Genetics 105: 767-779.

Rivas GBS, Zhou J, Merlin C, Hardin PE. 2021. Clockwork orange promotes CLOCK-CYCLE activation via the putative *Drosophila* ortholog of clock interacting protein circadian. Current Biology 31:4207-4219.

Saunders DS, Henrich VC, Gilbert LI. 1989. Induction of diapause in *Drosophila melangoaster*: photoperiodic regulation and the impact of arrhythmic clock mutations on time measurement. Proceedings of the National Academy of Sciences (USA) 86:3748-3752.

1530    Scanlan JL, Glendhill-Smith RS, P Battley P, Robin C. 2020. Genomic and transcriptomic

1531    analyses in Drosophila suggest that the ecdysteroid kinase-like (EcKL) gene family encodes the

1532    'detoxification-by-phosphorylation' enzymes of insects. Insect Biochemistry and Molecular

1533    Biology 123: 103429

1534

1535    Scheunemann L, Jost E, Richlitzki A, Day JP, Sebastian S, Thum AS, Efetova M, Davies S-A,

1536    Schwarzel M. 2012. Consolidated and labile door memory are separately encoded in the

1537    *Drosophila* brain. Journal of Neuroscience 32:17163-17171

1538

1539    Schmidt JM, Good RT, Appleton B, Sherrad J, Raymart GC, Bogwitz MR, Martin J, Daborn PJ,

1540    Goddard ME, Batterham, Robin C. 2010. Copy number variation and transposable elements

1541    feature in recent ongoing adaptation at the *Cyp6g1* locus. PLoS Genetics 6:e1000998

1542

1543    Schmidt PS, Matzkin L, Ippolito M, Eanes WF. 2005. Geographic variation in diapause

1544    incidence, life-history traits, and climatic adaptation in *Drosophila melanogaster*. Evolution

1545    59:1721-1732.

1546

1547    Shalaby NA, Pinzon JH, Narayanan AS, Jin EG, Ritz MP, Dove RJ, Wolfenberg H, Rodan AR,

1548    Buszczak M, Routhenfluh A. 2018. JmjC domain proteins modulate circadian behaviors and

1549    sleep in *Drosophila*. Scientific Reports 8:815

1550

1551    Spitze K 1993. Population structure in *Daphnia obtusa*: quantitative genetic and allozyme

1552    variation. Genetics 135:367-374.

1553

1554    Sprengelmeyer QD, Mansourian S, Lange JD, Matute DR, Cooper BS, Jirle EV, Stensmyr MC,

1555    Pool JE. 2020. Recurrent collection of *Drosophila melanogaster* from wild African environments

1556    and genomic insights into species history. Molecular Biology and Evolution 37:2775.

1557

1558    Staats M, Erkens RH, van de Vossenberg B, Wieringa JJ, Kraaijeveld K, Stielow B, Geml J,

1559    Richardson JE, Bakker FT. 2013. Genomic treasure troves: complete genome sequencing of

1560    herbarium and insect museum specimens. PLOS ONE 8:e69189.

1561

1562    Svetec N, Zhao L, Saelo P, Chiu JC Begun DJ. 2015. Evidence that natural selection maintains

1563    genetic variation for sleep in *Drosophila melanogaster*. BMC Evolutionary Biology 15:316

1564

1565    Tong JJ, Schriner SE, McCleary D, Day BJ, Wallace DC 2007. Life extension through

1566    neurofibromin mitochondrial regulation and antioxidant therapy for neurofibromatosis-1 in

1567    *Drosophila melanogaster*. Nature Genetics 39:476-485.

1568

1569    Turelli M, Hoffmann AA. Cytoplasmic incompatibility in *Drosophila simulans*: dynamics and

1570    parameter estimates from natural populations. Genetics 140:1319-1338.

1571

1572    Van Der Valk T, Díez-del-Molino D, Marques-Bonet T, Guschanski K, Dalén L. 2019. Historical

1573    genomes reveal the genomic consequences of recent population decline in eastern gorillas.

1574    Current Biology 29:165-170.

1575

1576    Voigt S, Laurent S, M Litovchenko, W. Stephan W. 2015. Positive selection at the polyhomeotic

1577    locus led to decreased thermosensitivity of gene expression in temperate Drosophila

1578    melanogaster. Genetics 200:591-599.

1579

1580    Wang J, Lee CH, Lin S, Lee T. 2006. Steroid hormone-dependent transformation of

1581    polyhomeotic mutant neurons in the Drosophila brain. Development 133:1231-1240.

1582

1583    Warren JT, Petryk A, Marques G, Parvy J-P, Shinoda T, Itoyama K, Kobayashi J, Jarcho M, Li Y,

1584    O'Connor MB, Dauphin-Villemart C, Gilbert LI. 2004. Phantom encodes the 25-hydroxylase of

1585    *Drosophila melanogaster* and *Bombyx mori*: a P450 enzyme critical in ecdysone biosynthesis.

1586    Insect Biochemistry and Molecular Biology 34:991-1010.

1587

1588 Werthebach M, Stewart FA, Gahlen A, Mettler-Altmann T, Akhtar I, Maas-Enriquez K, Droste A,

1589 Eichmann TO, Poschmann G, Stuhler K, Beller M. 2019. Control of *Drosophila* growth and

1590 survival by the lipid droplet-associated protein CG9186/Sturkopf. Cell Reports 26:3726-3740.

1591

1592 Whitlock MC 1999. Neutral additive genetic variation in a metapopulation. Genetical Research

1593 74:215-221.

1594

1595 Wohns AW, Wong Y, Jeffery B, Akbari A, Mallick S, Pinhasi R, Patterson N, Reich D, Kelleher J,

1596 McVean G. 2022. A unified genealogy of modern and ancient genomes. Science

1597 375:eabi8264.

1598

1599 Woodard C, E Alcorta E, J Carlson 1992. The *rdgB* gene in *Drosophila*: a link between vision

1600 and olfaction. Journal of Neurogenetics 8: 17-31

1601

1602 Yang MA, Fu Q. 2018. Insights into modern human prehistory using ancient genomes. Trends

1603 in Genetics 34:184-96.

1604

1605 Yassin A, Debat V, Bastide H, Pool JE. 2016. Recurrent specialization on a toxic fruit in an island

1606 *Drosophila* population. Proceedings of the National Academy of Sciences (USA) 113:4771-

1607 4776.

1608

1609 Yi X, Liang Y, Huerta-Sanchez E, Jin X, Cuo ZXP, Pool JE, Xu X, Jian H, Vinckenbosch N,

1610 Korneliussen TS, et al. 2010. Sequencing fifty human exomes reveals adaptation to high

1611 altitude. Science 329: 75-79.

1612

1613 Zetterstedt JW. 1842-1860. Diptera Scandinaviae disposita et descripta. Lundae ex Officina

1614 Lundbergiana. Vol 4-14: sumtibus rebus

**Supplementary Information:**

**Table S1.  Sample Information and Genomic Sequencing Statistics.**  Available collection information is given for each museum specimen.  All early 1800s Sweden flies (including the one labelled Småland) were ultimately inferred to have come from the same collection event. Sequencing statistics including read count and lengths, depth of coverage, genomic coverage, singleton rate, and proportion of singletons with A or T are also given.  Sample 19SL2 was excluded from most analyses because of its low sequencing depth and genomic coverage. The sex of each specimen was inferred from the relative levels of sequencing depth on the X chromosome and the autosomes.  Genomic characteristics of a representative set of genomes from modern material (from Lyon, France - Lack et al. 2016 MBE) are provided, as depicted in Figure 1.

**Table S2: Evidence of identity-by-descent from relatedness and inbreeding and genomic masking proportions.**

**Tab A:**  Mean pairwise genetic distance between X chromosomes from museum and modern genomes (excluding sites with missing data from any individual).  Unusually low values are highlighted.  Mean distance against a panel of modern genomes is indicated at the bottom.

**Tab B:**  Proportion of each genome masked due to pairwise IBD with other genomes, given as a fraction of the total genome and each chromosome arm.

**Tab C:**  Proportion of each genome masked due to inbreeding IBD, as indicated by runs of minimal heterozygosity, for each full genome and each chromosome arm.

**Table S3.  Regions of each museum fly genome that were masked due to relatedness IBD.**  Release 5 positions are given in separate tabs for each chromosome arm.

**Table S4.  Regions of each museum fly genome that were masked due to inbreeding IBD.** Release 5 positions are given in separate tabs for each chromosome arm.

**Table S5. Results of Principle Components Analysis of Genetic Structure.** Loadings for each genome are given for all 35 principle components. The X chromomsome was analyzed to avoid the influence of autosomal inversions. Up to 5 apparently unrelated individuals were included per population sample.

**Table S6. The estimated frequency of each of the most common inversions in museum and modern populations.** Inversion frequency estimates and results from each SNP reported to be associated with each inversion (by Kapun et al. 2016) are given in separate tabs. Estimation of inversion frequency was based on the median of SNP frequencies, in order to avoid spurious non-zero estimates.

**Table S7. Effective population size estimates from drift-only models do not predict observed nucleotide diversity.**
**Top:** Mean frequency differences and haploid effective population size estimates from drift-only models for each Lund time interval.
**Bottom:** Comparison of nucleotide diversity from simulations (using estimated Ne) versus observed data, for arm 3L.

**Table S8. Outlier regions from read depth QPBE as a potential indicator of structural variants at differing frequencies between samples.**
**Tab A:** For the 1800s time interval (1800s as focal population, Lund 1933 and Lyon as outgroups), QPBE outlier regions based on the upper 1% quantile and overlapping genes are given.
**Tab B:** For the 1900s time interval (Lund 1933 as focal population, Lyon and Stockholm as outgroups), QPBE outlier regions based on the upper 1% quantile and overlapping genes are given.

**Table S9. Outlier regions from PBE reflecting windows with SNP frequencies at strongly differing frequencies between samples.**

1673     **Tab A:** For the 1800s time interval (1800s as focal population, Lund 1933 and Lund 2010s),

1674     PBE outlier regions based on the upper 1% quantile and overlapping genes are given.

1675     **Tab B:** For the 1900s time interval (Lund 1933 as focal population, Lund 2010s and Lyon as

1676     outgroups), PBE outlier regions based on the upper 1% quantile and overlapping genes are

1677     given.

1678

**Table S10. Gene Ontology (GO) Enrichment based on PBE Outlier Regions.**

**Tab A:** Gene ontology enrichment for the 1800s-focused PBE outliers.

**Tab B:** Gene ontology enrichment for the 1900s-focused PBE outliers.

For each GO category, the total number of genes in the analyzed regions is given, along with

the number of outlier regions associated with these genes, and the identities of these outlier-

overlapping genes. A permutation P value is also given for each GO category, which does not

attempt to correct for the number of GO categories tested.

# Figure 1.

**A**

| 1780 The steam engine | 1816 The year without summer | 1825 First railway line | 1827 Agricultural reform (Swe) | 1845-52 The great hunger | 1867 Paris green (first chemical insecticide) | 1870s Urbanization (Swe) | 1870s Industrialization (Swe) | 1886 Benz patents the motor car | 1903 Wright brothers first flight | 1944 DDT | 1945 Atomic bombs | 1948 Dieldrin | 1953 Allethrin | 1962 Rachel Carson *Silent spring* | 1972 DDT banned | 1979 Permethrin | 1980 First wind farm | 1994 Imidacloprid | 2018 Imidacloprid banned (EU) |

Avg temp Stockholm
3.2°C    9.0°C

**B** Fallén collection Lund *Early 1800s*

**C** Zetterstedt collection Lund *Early 1800s* Lund.

**D** Passau *1850s*

**E** NE Zealand *1880s*

**F** Lund *1933*

**G** Zoologiska instiutionen Lund *1915*

**H** Lilla torg Lund *1782* Sydberckska gården

**I** Sandgatan Lund *1832*

**J** Lund Stockholm Netherlands France Italy Egypt *Contemporary*

Figure 1

## Figure 2.

# Figure 3.

**A** Identity-By-Decent mapping

**Inbreeding and relatedness - 1800s genomes**

**B** Inbreeding and relatedness - 1933 genomes

Figure 3

## Figure 4.

**A**

| | Lund 1800s | Lund 1933 | Lund 2010s | Lyon | Stockholm | Netherlands | Italy | Egypt |
|---|---|---|---|---|---|---|---|---|
| Lund 1800s | **0.00319** | 0.00410 | 0.00387 | 0.00400 | 0.00425 | 0.00402 | 0.00410 | 0.00425 |
| Lund 1933 | 0.0585 | **0.00260** | 0.00353 | 0.00373 | 0.00392 | 0.00373 | 0.00374 | 0.00391 |
| Lund 2010s | 0.0455 | 0.0676 | **0.00234** | 0.00346 | 0.00420 | 0.00347 | 0.00402 | 0.00356 |
| Lyon | 0.0301 | 0.0651 | 0.0414 | **0.00263** | 0.00384 | 0.00365 | 0.00369 | 0.00381 |
| Stockholm | 0.0438 | 0.0632 | 0.0420 | 0.0399 | **0.00291** | 0.00385 | 0.00386 | 0.00391 |
| Netherlands | 0.0474 | 0.0792 | 0.0588 | 0.0527 | 0.0522 | **0.00255** | 0.00370 | 0.00377 |
| Italy | 0.0435 | 0.0581 | 0.0402 | 0.0400 | 0.0336 | 0.0568 | **0.00269** | 0.00378 |
| Egypt | 0.0886 | 0.0794 | 0.0701 | 0.0807 | 0.0549 | 0.0842 | 0.0652 | **0.00262** |

$F_{ST}$

Rate of pairwise sequence differenes ($D_{xy}$)
Genetic differentiation summarized by $F_{ST}$
**Diagonal:** Rate of pairwise sequence differences within populations (π)

**B**

Legend:
- Lund 1800s
- Germany 1850s
- Denmark 1880s
- Lund 1933
- Stockholm
- Netherlands
- France
- Italy
- Egypt

Figure 4

**Figure 5.**

# Figure 6.

**1800s-focused scan**

Figure 6

## Figure 7.

**1900s-focused scan**

Figure 7