Discovery and characterization of a specific inhibitor of serine-threonine kinase cyclin dependent kinase-like 5 (CDKL5) demonstrates role in hippocampal CA1 physiology

Anna Castano*
Department of Pharmacology, University of Colorado School of Medicine, Aurora, CO
Anna.Castano@cuanschutz.edu

Margaux Silvestre*
Kinases and Brain Development Laboratory, The Francis Crick Institute, 1 Midland Road, London, NW1 1AT UK
Margaux.silvestre@crick.ac.uk

Carrow I. Wells
Structural Genomics Consortium, UNC Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America
carrow.i.wells@gsk.com

Jennifer L. Sanderson
Department of Pharmacology, University of Colorado School of Medicine, Aurora, CO
JENNIFER.SANDERSON@CUANSCHUTZ.EDU

Carla A. Ferrer
Structural Genomics Consortium, UNC Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America
Calafe2@hotmail.com

Han Wee Ong
Structural Genomics Consortium, UNC Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America
onghw@live.unc.edu

Yi Liang
Structural Genomics Consortium, UNC Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America
liang_yi@wuxiapptec.com

William Richardson
Centre for Medicines Discovery, Nuffield Department of Medicine, University of Oxford, Oxford, OX3 7FZ, United Kingdom
william.richardson95@outlook.com

Josie A. Silvaroli
Division of Pharmaceutics and Pharmacology, College of Pharmacy and Comprehensive Cancer Center, The Ohio State University, Columbus, Ohio, 43210, United States of America
silvaroli.1@osu.edu

Frances M. Bashore
Structural Genomics Consortium, UNC Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America
Frances.Bashore@unc.edu

Jeffery L. Smith
Structural Genomics Consortium, UNC Eshelman School of Pharmacy, University of North Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America

55   jeffs@email.unc.edu
56
57   Isabelle M. Genereux
58 Structural Genomics Consortium, UNC Eshelman School of Pharmacy, University of North
59   Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America
60   genereux@live.unc.edu
61
62   Kelvin Dempster
63 Kinases and Brain Development Laboratory, The Francis Crick Institute, 1 Midland Road,
64   London, NW1 1AT UK
65   Kelvin.dempster@crick.ac.uk
66
David H. Drewry
Structural Genomics Consortium, UNC Eshelman School of Pharmacy, University of North
Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America; UNC
Lineberger Comprehensive Cancer Center, School of Medicine, University of North Carolina at
Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America
david.drewry@unc.edu
73
74   Navjot S. Pabla
75 Division of Pharmaceutics and Pharmacology, College of Pharmacy and Comprehensive
76 Cancer Center, The Ohio State University, Columbus, Ohio, 43210, United States of America
77   pabla.2@osu.edu
78
79   Alex N. Bullock
80 Centre for Medicines Discovery, Nuffield Department of Medicine, University of Oxford, Oxford,
81   OX3 7FZ, United Kingdom
82   alex.bullock@cmd.ox.ac.uk
83
84   Tim A. Benke#
85 Departments of Pediatrics, Pharmacology, Neurology and Otolaryngology, University of
86   Colorado School of Medicine, Aurora, CO
87   Tim.Benke@cuanschutz.edu
88
89   Sila K. Ultanir#
90 Kinases and Brain Development Laboratory, The Francis Crick Institute, 1 Midland Road,
91   London, NW1 1AT UK
92   Sila.ultanir@crick.ac.uk
93
94   Alison D. Axtman#
95 Structural Genomics Consortium, UNC Eshelman School of Pharmacy, University of North
96   Carolina at Chapel Hill, Chapel Hill, North Carolina, 27599, United States of America;
97   orcid.org/0000-0003-4779-9932
98   Alison.Axtman@unc.edu
99
100 *Co-first authors, #Co-senior authors
101
Corresponding author: Tim A. Benke, University of Colorado Denver, School of Medicine, 12800

E 19th, MS8102, Denver, CO 80045, tim.benke@cuanschutz.edu

Number of words
Abstract: 210
Introduction: 811
Methods: 2976
Results: 2194
Discussion: 810
Number of Figures: 10
Number of Tables: 1
Number of References: 81
Supplementary information: 10 Figures, 2 schema, 2 tables, text
115

Funding: NIH-NINDS NS112770 (all), Ponzio Family Chair in Neurology Research (AC, TAB),

The Structural Genomics Consortium (SGC) is a registered charity (number 1097737) that

receives funds from Bayer AG, Boehringer Ingelheim, the Canada Foundation for Innovation,

Eshelman Institute for Innovation, Genentech, Genome Canada through Ontario Genomics

Institute [OGI-196], EU/EFPIA/OICR/McGill/KTH/Diamond, Innovative Medicines Initiative 2

Joint Undertaking [EUbOPEN grant 875510], Janssen, Merck KGaA (aka EMD in Canada and

USA), Pfizer, the São Paulo Research Foundation-FAPESP, and Takeda (CIW, CAF, HWO, YL,

FMB, JLS, IMG, DHD, ADA). Research reported in this publication was supported in part by the

NC Biotechnology Center Institutional Support Grant 2018-IDG-1030 (DHD, ADA), NIH

U24DK116204 (DHD, ADA), and NIH 1R44TR001916 (DHD, ADA). This work was supported by

the Francis Crick Institute which receives its core funding from Cancer Research UK (CC2037),

the UK Medical Research Council (CC2037), and the Wellcome Trust (CC2037); Loulou

Foundation Project Grant (11015). Margaux Silvestre is a recipient of the Loulou foundation

junior fellowship award (2021). For the purpose of Open Access, the author has applied a CC

BY public copyright license to any Author Accepted Manuscript version arising from this

submission.

Acknowledgments: NanoBRET constructs for CDKL5, GSK3α, and GSK3β were kindly provided

to the SGC-UNC team by Promega. TREE*spot* kinase interaction mapping software

(http://treespot.discoverx.com) was used to prepare the kinome trees in Figure 3. The authors

thank the Diamond Light Source for beamtime (proposal mx28172) as well as the staff of

beamline i04 for their guidance.

Disclosures:

Benke – Consultancy for AveXis, Ovid, GW Pharmaceuticals, International Rett Syndrome

Foundation, Takeda, Taysha, CureGRIN, GRIN Therapeutics, Alcyone, Neurogene, and

Marinus; Clinical Trials with Acadia, Ovid, GW Pharmaceuticals, Marinus and RSRT; all

remuneration has been made to his department

Axtman – Advisor for Proteic Bioscience Inc.

Abstract

Pathological loss-of-function mutations in cyclin-dependent kinase-like 5 (*CDKL5*) cause

CDKL5 deficiency disorder (CDD), a rare and severe neurodevelopmental disorder associated

with severe and medically refractory early-life epilepsy, motor, cognitive, visual and autonomic

disturbances in the absence of any structural brain pathology. Analysis of genetic variants in

CDD have indicated that CDKL5 kinase function is central to disease pathology. *CDKL5*

encodes a serine-threonine kinase with significant homology to GSK3β, which has also been

linked to synaptic function. Further, *Cdkl5* knock-out rodents have increased GSK3β activity and

often increased long-term potentiation (LTP). Thus, development of a specific CDKL5 inhibitor

must be careful to exclude cross-talk with GSK3β activity. We synthesized and characterized

specific, high-affinity inhibitors of CDKL5 that do not have detectable activity for GSK3β. These

compounds are very soluble in water but blood-brain barrier penetration is low. In rat

hippocampal brain slices, acute inhibition of CDKL5 selectively reduces post-synaptic function

of AMPA-type glutamate receptors in a dose-dependent manner. Acute inhibition of CDKL5

reduces hippocampal LTP. These studies provide new tools and insights into the role of CDKL5

as a newly appreciated, key kinase necessary for synaptic plasticity. Comparisons to rodent

knock-out studies suggest that compensatory changes have limited the understanding of the

roles of CDKL5 in synaptic physiology, plasticity and human neuropathology.

Introduction

Pathological loss-of-function mutations in cyclin-dependent kinase-like 5 (*CDKL5*)[1, 2] cause CDKL5 deficiency disorder (CDD, OMIM 300203, 300672), a rare (incidence 1:40,000–60,000[3-5]), severe neurodevelopmental disorder associated with severe early-life epilepsy, motor, cognitive, visual and autonomic disturbances[2, 6-9, 10]. Epilepsy in CDD does not respond to *any* antiseizure medications[11]; a newly approved therapy provides modest improvements[12]. Hemizygous males and heterozygous females can be equally and severely affected[13, 14]. These severe symptoms are in absence of any structural brain pathology[15]. CDKL5 is a key mediator of synaptic and network development and physiology. Recent analysis of genetic variants in CDD have indicated that CDKL5 kinase function is central to disease pathology[13, 16]. In other words, loss of kinase function seems functionally equivalent to loss of the protein.

In rodent models, CDKL5 is expressed throughout the CNS primarily in neurons, increases post-natally and is stabilized at peak levels in adults[17-19], indicating roles during development and into adulthood. These features make it impossible to tease apart the precise role of altered CDKL5 function in mediating these symptoms in CDD. It remains unclear whether symptomatic abnormalities are due to chronic CDKL5 dysfunction, acute CDKL5 dysfunction during an earlier critical developmental time point, worsened by epilepsy or any combination. Adult rodent models of CDD are associated with abnormal behaviors, visual disturbances[20] and multiple abnormal signaling cascades[21-25]. CDKL5 is localized to the dendritic spines of excitatory synapses as well as the nucleus of neurons[17, 26, 27]. CDKL5 functions in the formation of excitatory synapses with partners that include PSD-95[28], NGL-1[29], Shootin1[30], and actin[17]. Previous work had suggested CDKL5 specific substrates including MeCP2, DNMT1, AMPH1, NGL-1 and HDAC4 but reliable antibodies have not been available[20]. Recent work has identified MAP1S and microtubule end binding protein 2 (EB2) as physiological CDKL5

substrates in brain [19]. A phosphospecific antibody for EB2-S222 has been used to report

CDKL5 activity in mouse models and human iPSCs [19, 31, 32]. CDKL5 is a binding partner of

both PSD-95 and gephyrin[33]; binding with PSD-95 is critical for spine development[21].

Studies have typically found that loss of CDKL5 leads to a global reduction in excitatory

synapse numbers[29, 34], reduced PSD-95[35, 36] and synapsin[35] with loss of AMPA-type

glutamate receptors (GluA2[37]) and increased NMDA-type glutamate receptors (GluN2B[38]).

Inhibitory synapses appear to be unaffected[29], although inhibitory synaptic currents are

affected in some CDKL5 mouse models[39]. Embryonic knock-out of *Cdkl5* rodents either

enhanced long-term potentiation (LTP) [38, 40] or did not affect LTP [37, 40] in an age

dependent fashion. Reports so far do not fully explain the presumed excitation/inhibition

imbalance in epilepsy. Further, immature CDKL5 deficient mice do not have seizures or

epilepsy[21, 41] and appear insensitive to pro-convulsants such as kainate [21]. This paradox

highlights a big gap in our understanding of CDKL5 function: early developmental onset of

medically resistant epilepsy in CDD contrasts with a complete lack in experimental models. The

reasons for this mismatch are unclear and include possible developmental or other rodent-

specific compensations, which typically involve transcriptional changes. Finally, the role of acute

CDKL5 kinase dysfunction at any developmental time point is unknown. To address this, we

sought to develop a sensitive and specific inhibitor of CDKL5.

CDKL5 is an X-linked gene encoding a 115 kDa serine-threonine kinase and member of the

CMGC family that includes cyclin dependent kinases (<u>C</u>DK), <u>M</u>AP-kinases, glycogen synthase

kinases (<u>G</u>SK) and cyclin-dependent kinase-like (<u>C</u>DKL) [42]. It has limited structural homology

to CDKL1, 2, 3 and 4 (http://mbv.broadinstitute.org/) but significant homology to GSK3β[43].

CDKL1-4 mRNA are expressed at much lower levels than *CDKL5* in mouse and human brain

(http://mouse.brain-map.org). Of these kinases, only GSK3β has been linked to brain synaptic

function[44].

CDKL5 appears to be linked to GSK3β function. *Cdkl5* knock-out mice have increased

GSK3β activity, as evidenced by *hypo*-phosphorylation of GSK3β at S-9 and decreased β-

catenin[22]. β-catenin, which is phosphorylated and destabilized by active GSK3β, is often used,

with S9-phospho-GSK3β, as a marker of GSK3β activity[45]. Inhibition of GSK3β in CDKL5

(young but not old) knock-out mice normalizes expression of β-catenin and S9-phospho-

GSK3β[22, 23]. While upstream signaling is also affected (mTOR and AKT)[21], it is completely

unclear how CDKL5 and GSK3β activities are linked in CDKL5 knock-out mice.

GSK3β mediates a yin-yang interaction of LTP and LTD. Activation of GSK3β is required for the

induction of GluN-LTD while inhibition occurs with the induction of LTP[44]. Thus, development

of a specific CDKL5 inhibitor must be careful to exclude crosstalk with GSK3β activity.

Here, we synthesized and characterized specific, high-affinity inhibitors of CDKL5 that do

not have detectable activity for GSK3β. These compounds are very soluble in water, but blood-

brain barrier penetration is low. When applied directly to rat hippocampal brain slices, acute

inhibition of CDKL5 selectively reduces post-synaptic function of AMPA-type glutamate

receptors in a dose-dependent manner and inhibits a key form of synaptic plasticity,

hippocampal LTP. These studies provide new insights into the role of CDKL5 in neuronal

function and pathology.

Methods

Synthetic Schemes. Compound **B1** was prepared as a TFA salt with >95% purity (by

HPLC) according to Scheme 1 (Supplementary Information) via a reductive coupling reaction in

the presence of sodium borohydride and ethanol followed by amide coupling and deprotection.

Compounds **B4** and **B12** were prepared as an HCl salt with >95% purity (by NMR) according to

Scheme 2 (Supplementary Information). An amide coupling reaction in the presence of

propylphosphonic anhydride (T3P) followed by nitro group reduction generated a common

intermediate. Subsequent amide coupling reaction followed by deprotection yielded **B4** and

B12. Full characterization of the intermediates and final products (Figure 2) are included as

Supplementary Information.

NanoBRET Assays. Human embryonic kidney (HEK293) cells obtained from ATCC

(Manassas, VA, USA) were cultured in Dulbecco's Modified Eagle's medium (DMEM, Gibco)

supplemented with 10% (v/v) fetal bovine serum (FBS, Corning). These cells were incubated at

37°C in 5% CO_2 and passaged every 72 hours with trypsin (Gibco) so that they never reached

confluency. Promega (Madison, WI, USA) kindly provided constructs for NanoBRET

measurements of CDKL5 (NLuc-CDKL5), GSK3α (NLuc-GSK3α), and GSK3β (NLuc-GSK3β)

included in Table S1 as well as Figures 3, S4, and S5. N-terminal NLuc orientations were used

for all three kinases. NanoBRET assays were executed in dose–response (12-pt curves) format

in HEK293 cells as previously reported [46-48]. Assays were executed as recommended by

Promega, using 0.31 μM of tracer K11 for CDKL5, 0.13 μM of tracer K8 for GSK3α, and 0.063

μM of tracer K8 for GSK3β.

Analysis of Kinome-Wide Selectivity. The broad selectivity of compounds **B1**, **B4**, and

B12 at 1 μM was evaluated via the Eurofins DiscoverX Corporation (Fremont, CA, USA)

*scan*MAX assay platform. In total, 403 wild-type (WT) human kinases were included in these

analyses. Percent of control (PoC) values were generated, which allowed for the calculation of

selectivity scores (S_{10}(1 μM)) noted in Figure 3 [43]. WT human kinases in the *scan*MAX panel

with PoC ≤20 (and GSK3β) for **B1**, with PoC <10 for **B4**, and PoC <35 for **B12** are also included

in Figure 3. The kinome tree diagrams in Figure 3 were created based on WT kinases with PoC

<10 for each compound.

Biochemical Assays. A homogenous competition binding assay for CDKL5 was

executed at Luceome Biotechnologies, LLC (Tuscon, AZ, USA) using their *KinaseSeeker*

technology [49]. Briefly, this is a luminescence-based assay that relies on displacement of an

active site dependent probe by an ATP-competitive test compound. Compounds **B1**, **B4**, and

B12 were evaluated in dose–response (12-pt curve) format in duplicate. Curves are included in

Figure S4 and IC_{50} values generated are embedded in tables within Figure 3.

Radiometric enzymatic assays were executed at Eurofins at the K_m value for ATP to

generate dose–response (9-pt) curves for all kinases listed in Figure 3 (except for CDKL5). A

detailed protocol for these assays, which includes the protein constructs, substrate, and controls

employed, is part of the Eurofins website: https://www.eurofinsdiscoveryservices.com.

Kinetic Solubility. The kinetic solubility of compound **B1** was evaluated using an aliquot

of a 10 mM DMSO stock solution dissolved in phosphate buffered saline solution (PBS) at pH

7.4. This analysis was done by Analiza, Inc (Cleveland, OH, USA) as described previously [50].

The reported solubility values in Table 1 have been corrected for background nitrogen present

in DMSO and the media.

Microsomal Stability. Mouse liver microsomal stability of compound **B1** was evaluated by

Analiza, Inc (Cleveland, OH, USA) as described previously [48]. This data generated by Analiza

is included in Table 1.

CDKL Family Thermal Shift Assays. CDKL1, CDKL2, CDKL3, and CDKL5 proteins were

produced using the construct boundaries previously used to generate 4AGU, 4AAA, 3ZDU, and

4BGQ, respectively. 4 µM of the kinase domain of CDKL1, CDKL2, CDKL3, or CDLK5 in 10 mM

HEPES-NaOH pH 7.4 and 500 mM NaCl was incubated with **B1** (12.5, 25, or 50 µM) in the

presence of 5× SyPRO orange dye (Invitrogen). Next, fluorescence was evaluated using a

Real-Time PCR Mx3005p machine (Stratagene). A previously reported protocol was followed to

run the Tm shift assays and assess melting temperatures[56]. Data is included in Figure S6.

CDKL5 Enzymatic Assays. These assays were executed as previously described[57,

58]. Briefly, FLAG-tagged WT or kinase dead (KD, CDKL5 K42R) constructs of human CDKL5

were subcloned into a pT7CFE1-CHis plasmid (Thermo Fisher). A HeLa cell lysate-based Kit (1-

Step Human Coupled IVT Kit—DNA, 88881, Life Technologies) enabled *in vitro* translation of

these constructs. His Pur cobalt spin columns (Thermo Scientific) were used to purify the *in*

vitro-translated proteins. To execute the *in vitro* kinase assays, myelin basic protein (Active

Motif, 31314) was employed as a substrate for recombinant CDKL5. These two components

were incubated in kinase buffer (Cell Signaling, 9802) supplemented with or without ATP (50

µM) for 30 minutes at 30 °C, followed by kinase assays run using the ADP-Glo Kinase Assay kit

(Promega). After the termination of kinase assay, NuPAGE LDS Sample Buffer (4X) was added

to the reaction mixture and the samples were heated at 100°C for 10 minutes. The protein

samples were run on Invitrogen Bis-tris gradient midi-gels and transferred to PVDF membrane

for western blot analysis. Primary antibody used for western blot analysis was from Cell

Signaling: FLAG (#14793) and was used at 1:1,000 dilution. Anti-Rabbit HRP-conjugated

secondary antibody was from Jackson Immunoresearch (#111-035-144) and used at 1:5,000

dilution. Immunoblot signals were detected using SignalFire ECL Reagent (#6883) and X-ray

films (Bioland #A03-02). Canon LiDE400 scanner was used for scanning the films. Precision

Plus Protein Dual Color Standard prestained protein marker (Biorad) was used to estimate the

molecular weight of sample proteins.

Snapshot PK Study. The pharmacokinetics of compound **B1** was evaluated by

Pharmaron (San Diego, CA, USA) following a single intraperitoneal administration to CD1 mice

(two males). A dose of 2.29 mg/kg of the TFA salt of compound **B1** in NMP/Solutol/PEG-

400/normal saline (v/v/v/v, 10:5:30:55) was prepared just before use. This dose was calibrated

due to the compound being a salt form. Plasma was sampled 0.5-, 1-, 2-, and 4-hours post-

dose. Bioanalytical assays were run using Prominence HPLC and AB Sciex Triple Quan 5500

LC/MS/MS instruments, and a HALO column (90A, C18, 2.7 µm, 2.1x50 mm). Snapshot PK

results are included in Figure S10. PK parameters were estimated by non-compartmental model

using WinNonlin 8.3 (Certara, Princeton, NJ, USA). The lower limit of quantification for plasma

sampling was 1.53 ng/mL.

Analysis of Brain/Plasma Concentration. The brain/plasma concentration of compound

B1 was evaluated by Pharmaron following a single intraperitoneal administration to CD1 mice

(three males per dose). Two doses (2.29 mg/kg and 7.63 mg/kg) of the TFA salt of compound

B1 in NMP/Solutol/PEG-400/normal saline (v/v/v/v, 10:5:30:55) were prepared just before use.

These doses were calibrated due to the compound being a salt form. Plasma and brain samples

were collected 1-hour post-dose. Brain samples were homogenized at a ratio of 1:3 with PBS

(W/V, 1:3) and then final brain concentrations corrected. Bioanalytical assays were run using

Prominence HPLC and AB Sciex Triple Quan 5500 LC/MS/MS instruments, and a HALO

column (90A, C18, 2.7 µm, 2.1x50 mm). Brain/plasma concentration results are included in

Table S2. The lower limit of quantification for plasma sampling was 1.53 ng/mL and 0.763

ng/mL for brain.

Animals-The Francis Crick Institute (TFCI). Rat handling and housing was performed

according to the regulations of the Animal (Scientific Procedures) Act 1986. Animal studies were

approved by the Francis Crick Institute ethical committee and performed under U.K. Home

Office project license (PPL P5E6B5A4B).

Rat Neuron Primary Culture-TFCI. Primary cortical cultures were prepared from

embryonic day (E) E18.5 embryos of Long Evans rats as described [59]. Pregnant females were

culled using cervical dislocation, embryos were removed from the uterus and the brains were

taken out. Cortices were dissected out, pooled from multiple animals, and washed three times

with HBSS. An incubation with 0.25 % trypsin for 15 minutes at 37 °C was followed by four

times washing with HBSS. Cells were dissociated and then counted using a hemocytometer.

Neurons were plated on 12-well culture plates at a density of 300,000 cells per well. The wells

were coated with 0.1 M borate buffer containing 60 µg/mL poly-D-lysine and 2.5 µg/mL laminin

and placed in the incubator overnight. Neurons were plated with minimum essential medium

(MEM) containing 10% fetal bovine serum (FBS), 0.5% dextrose, 0.11 mg/mL sodium pyruvate,

2 mM glutamine and penicillin/streptomycin. After 4 hours, cultures were transferred to

neurobasal medium containing 1 mL of B27 (Gibco), 0.5 mM glutamax, 0.5 mM glutamine, 12.5

µM glutamate and penicillin/streptomycin. Primary neuronal cultures were kept at 37 °C and 5%

CO_2. Every 3-4 days, 20-30% of the maintenance media was refreshed. Between DIV14-16,

neurons were treated with 5 nM, 50 nM, 500 nM and 5 µM of the different compounds for 1

hour. The compounds were added directly to the media and the plates were place at 37 °C for

the time of the treatment. DMSO was added to the well for the control condition.

Western Blotting-TFCI. After treatment, neuronal cultures were lysed in 300 µL of 1X

sample buffer (Invitrogen) containing 0.1 M DTT. Lysates were sonicated briefly twice and

denatured at 70 °C for 10 minutes. The samples were centrifuged at 13,300 rpm for 10 minutes

and ran on NuPage 4-12% Bis-Tris polyacrylamide gels (Invitrogen). Proteins were transferred

onto a Immobilon PVDF membrane (Millipore), which was then blocked in 4% milk in tris-

buffered saline containing 0.1% Tween-20 (TBST) for 30 minutes. Primary antibodies were

incubated at 4 °C overnight, and HRP-conjugated secondary antibodies at RT for 2 hours. The

following primary antibodies were used: rabbit anti-CDKL5 (1:1,000; Atlas HPA002847), rabbit

anti-pS222 EB2 (1:2,000; Covalab, from [19]), rat anti-EB2 (1:2,000; Abcam ab45767), mouse

anti-tubulin (1:100,000; Sigma T9026), rabbit anti-β-catenin (1:1,000; Cell Signalling 9562) and

rabbit anti phospho-β-catenin (Ser33/37 Thr41) (1:250, Cell Signalling 9561).The following

secondary antibodies were used at a concentration of 1:10,000: HRP-conjugated anti-rabbit

(Jackson 711-035-152), HRP-conjugated anti-mouse (Jackson 715-035-151) and HRP-

conjugated anti-rat (Jackson 712-035-153). The membrane was developed using ECL reagent

(Cytiva) and was visualized with an Amersham Imager 600 (GE Healthcare). Quantification of

Western blots was manually performed using Image Studio Lite Software (version 5.2). EB2

phosphorylation was measured relative to total EB2 and catenin phosphorylation to total

catenin. The other proteins were normalized to tubulin, if not indicated otherwise.

Animals-University of Colorado School of Medicine (UC-SOM). All studies conformed to

the requirements of the National Institutes of Health *Guide for the Care and Use of Laboratory*

Rats and were approved by the Institutional Animal Care and Use subcommittee of the

University of Colorado Anschutz Medical Campus (protocol 00411). Timed-pregnant Sprague

Dawley rats (Charles Rivers Labs, Wilmington, MA, USA) gave birth in-house. All rodents were

housed in micro-isolator cages with water and chow available *ad libitum*.

Hippocampal Slice Preparation and Electrophysiology-UC-SOM. As done previously [60-

62], following rapid decapitation and removal of the rat brain at post-natal day (P) 20–30,

sagittal hippocampal slices (400 µm) were made using a vibratome (Leica VT 1200, Buffalo

Grove, IL) in ice-cold sucrose artificial cerebral spinal fluid (sACSF: 206 mM sucrose, 2.8 mM

KCl, 1 mM $CaCl_2$, 3 mM $MgSO_4$, 1.25 mM NaH_2PO_4, 26 mM $NaHCO_3$, 10 mM D-glucose and

bubbled with 95%/5% O_2/CO_2). Following removal of CA3, slices were recovered in a

submersion type chamber perfused with oxygenated artificial cerebral spinal fluid (ACSF: 124

mM NaCl, 3 mM KCl, 1 mM $MgSO_4$, 2 mM $CaCl_2$, 1.2 mM NaH_2PO_4, 26 mM NaHCO3, 10 mM

D-glucose and bubbled with 95%/5% O_2/CO_2) at 28°C for at least 90 min and then submerged

in a recording chamber perfused with recirculated ACSF. All electrophysiology was performed in

the CA1 region at 28°C. Drugs were added to ACSF. Two twisted-tungsten bipolar stimulating

electrodes were offset in the CA1 to stimulate one or two independent Schaffer collateral-

commissural pathways using a constant current source (WPI, Sarasota, FL) with a fixed

duration (20 µs), each at a rate of 0.033 Hz. Field excitatory post-synaptic potentials (fEPSPs)

were recorded from the stratum radiatum (or stratum pyramidale region where indicated) region

of CA1 using a borosilicate glass (WPI, Sarasota, Fl) microelectrode (pulled (Sutter, Novato,

CA) to 6 to 9 MΩ when filled with ACSF), amplified 1000x (WPI, Sarasota, Fl and Warner,

Hamden, CT), and digitized (National Instruments, Austin, Texas) at 20 kHz using winLTP-

version 2.4[63] to follow fEPSP slope (averaged over 4 EPSPs), measured using 20% to 80%

rise times, expressed as percent of baseline, during the course of an experiment. To be sure

only "healthy" slices were included in our studies, responses had to meet several criteria: fiber

volleys less than 1/3 of response amplitude and peak responses larger than 0.5 mV; responses

and fiber volley must be stable (<5% drift). Following baseline stabilization of fEPSP slope at

~50% of maximal slope for at least 30 min, measurements of synaptic physiology were

performed. Stimulation was adjusted to make 4-6 measurements from ~10% to -90% maximal

slope to measure input-output (I/O) curves; recordings that deviated from linear slope were

rejected. Paired-pulse ratios of slopes and peaks were measured at ~50% of maximal slope.

For measurements of NMDA-receptor dependent population spikes [64], Mg^{2+} was omitted

from recording ACSF, the recording electrode was placed in stratum pyramidale, stimulation

was adjusted to evoke a first population spike > 2 mV, and stabilization of the 2nd NMDA-

receptor dependent population spike (measured from the coast-line of the fEPSP) for greater

than 30 minutes was allowed prior to measuring the effect of antagonists. Theta burst LTP was

induced at test strength of two trains, separated by 20 seconds, of 10 trains of 4 stimuli at

100Hz, separated by 200 msec. Potentiation was statistically compared over the last 5 minutes.

For whole-cell, voltage-clamp electrophysiological recordings of excitatory post-synaptic

currents (EPSCs), 350 µm horizontal hippocampal slices were prepared from rats at post-natal

day (P) 20-30 similar to previously[65, 66]. After 15 min recovery at 34°C, slices were

maintained at room temperature until recording in modified ACSF containing (in mM): 126 NaCl,

5 KCl, 2 $CaCl_2$, 1.25 NaH_2PO_4, 1 $MgSO_4$, 26 $NaHCO_3$, 10 glucose and 2 N-acetyl cysteine.

Slices were incubated in either CAF-382 (**B1**) (100 nM) or vehicle (water) in modified ACSF for

at least 1h prior to recording. Using infrared–differential interference contrast microscopy for

visualization, whole-cell recordings (Axopatch 200B amplifier and pClamp software (Molecular

Devices)) from CA1 pyramidal neurons (series resistance between 3 and 6 MΩ) were performed

at 29-30°C with an intracellular solution containing (in mM): 130 Cs gluconate, 1 CsCl, 1 $MgSO_4$,

10 HEPES, 1 EGTA, 4 MgATP, 0.4 MgGTP, spermine (0.01) and 2 QX-314, pH 7.3. AMPA-type

EPSCs, evoked using a bipolar tungsten-stimulating electrode as above, at a holding potential

of −65 mV were pharmacologically isolated using picrotoxin (50 μM; Tocris). EPSC peaks were

evoked every 20 seconds. After establishing whole-cell access, a 3 min baseline recording at

−65 mV was followed by LTP induction using a pairing protocol: cells were depolarized to 0 mV

for 90 sec and stimulated every 20 sec then followed by delivery of 1s x 100Hz high frequency

stimulation. Evoked EPSC amplitudes were subsequently monitored at -65 mV. Potentiation

was statistically compared over the last 5 minutes.

Western Blotting-UC-SOM. Hippocampal slices were prepared and recovered as for

electrophysiology with the addition of cuts to isolate CA1 from the remaining hippocampus; after

recovery, the slices from a given rat were incubated in inhibitor or vehicle (water) in ACSF for 2

hours. Following this, slices were suspended in STE buffer, sonicated, boiled for 5 minutes and

then frozen until further use. To minimize the effects of slice preparation[67] and to control for

slice quality, only slices that came from a preparation that met electrophysiological criteria were

used. All concentrations were quantified with BCA and then loaded in duplicate on 10%

polyacrylamide gel and a five-point dilution series of naive rat hippocampal homogenate was

included on each gel as a standard curve for quantification of immunoreactivity/ μg loaded

protein[68]. Following transfer to PolyScreen PVDF transfer membrane (Genie, Idea Scientific

Company, Minneapolis, MN, USA), blots were blocked in BSA or Carnation nonfat dry milk for 1

h and incubated either 1h at room temperature or overnight at 4°C with antibodies.   The

following primary antibodies were used: sheep anti-CDKL5 (1:2,000; University of Dundee, Anti-

CDKL5(350-650)), rabbit anti-pS222 EB2 (1:1,000; Covalab, from [19]), rat anti-EB2 (1:1,000;

Abcam ab45767). Blots were then subjected to three 10-min washings in Tris-buffered saline

(140 mm NaCl, 20 mm Tris pH 7.6) plus 0.1% Tween 20 (TTBS), before being incubated with

anti-sheep, anti-rat or anti-rabbit secondary antibody (1:5000) in 1% BSA or milk for 1 h at room

temperature, followed by three additional 10-min washes preformed with TTBS.

Immunodetection was accomplished using a chemiluminescent substrate kit (SuperSignal West

Femto Maximum Sensitivity Substrate; Pierce) and the Alpha Innotech (Alpha Innotech, San

Leandro, CA, USA) imaging system. Quantification was performed using AlphaEase software

(Alpha Innotech, San Leandro, CA, USA) and Excel (Microsoft, Redmond, WA, USA).

Immunoreactivity was reported as the density of sample bands relative to the standard curve.

For phospho-proteins, ratios of immunoreactivity/μg to totals are reported without

standardization. Only values falling within the standard curve generated from the dilution series

included on each gel were incorporated into the final analysis. Some of the blots were then

stripped (Restore PLUS Western Blot Stripping Buffer; Thermo Scientific, Rockford, IL) and

reblotted if needed.

Statistical Analysis-OSU. Data were analyzed via GraphPad Prism 9. Each assay was

run in triplicate and mean values are graphed in Figure 4 with error bars showing standard

deviation (SD). In Figure 4, statistical analysis was done using one-way ANOVA with Dunnett's

multiple comparisons test. Thresholds for significance (p-values) were set at ***$p < 0.0001$ and

non-significant statistics were not included.

Statistical Analysis-TFCI. Data were analyzed using GraphPad Prism 9. Exact values of

n and statistical methods are mentioned in the figure legends. Each concentration was

compared to the control using a one-way ANOVA test. A p-value higher than 0.05 was not

considered as statistically significant. Thresholds for significance were placed at *$p \leq 0.05$, **$p \leq$

0.01, ***p ≤ 0.001 and ****p≤ 0.0001. All errors bars in the figures are standard deviation (SD)

where indicated. Non-significant statistics were not included.

Statistical Analysis-UCSOM. Data are expressed as mean ± SEM with n = number of

rats for a given treatment, unless otherwise stated. Data are plotted using Sigmaplot 12.5

(Systat, Chicago, IL, USA). Mann-Whitney Rank Sum, one-way, two-way and repeated

measures (RM) ANOVA (Holm-Sidak post-hoc testing) and Student's t-tests were used, where

indicated and appropriate, for statistical comparisons for electrophysiological and biochemical

using SigmaPlot 12.5 (Systat, Chicago, IL). Significance is reported at $p < 0.05$ unless otherwise

stated. All errors bars in the figures are standard error (SE) where indicated. Non-significant

statistics were not included.

Data and materials availability. Data has been uploaded to Dryad

(https://doi.org/10.5061/dryad.sn02v6x88). Newly synthesized compounds are available from

the senior authors (Alison.Axtman@unc.edu).

Results

Identification of CDKL5 inhibitors that lack GSK3β activity. SNS-032 and AT-7519 were

initially identified as potent CDK inhibitors [43, 69]. We profiled our extensive library of SNS-032

and AT-7519 analogs using the CDKL5 NanoBRET assay. In parallel, the CDKL5 actives (IC_{50}

<1 µM) were profiled using the GSK3β NanoBRET assay. Based on the data generated, a plate

of 25 compounds was assembled. Criteria for inclusion on this plate included an IC_{50} <400 nM in

the CDKL5 NanoBRET assay and >20-fold difference between the CDKL5 NanoBRET and the

GSK3β NanoBRET IC_{50} values (Table S1).

A targeted screen of CDKL5 inhibitors in neurons. A set of 20 CDKL5 inhibitors were

selected based on their selectivity properties for CDKL5 versus GSK3α/β for testing their

inhibition of CDKL5 in rat primary cortical neuron cultures. We tested if we could detect a dose-

dependent reduction of EB2 pSer222 following 1 hour incubation in culture media. We

normalized pS222 EB2 with total EB2 levels to assess a specific reduction in phosphorylation as

opposed to a loss of total protein levels. Almost all inhibitors showed a pSer222 EB2 reduction

at 500 nM; most inhibitors showed a reduction at 50 nM (Figure 1). Three inhibitors caused a

significant reduction in pSer222 EB2 at 5 nM without a change in total EB2 levels: CAF-382

(**B1**), HW2-013 (**B4**) and LY-213 (**B12**) (Figure 1). These represented different chemical parent

backbones, so we decided to study these inhibitors further.

Potency and selectivity analyses of lead compounds. As shown in Figure 2, **B1** is an

SNS-032 analog, while **B4** and **B12** are analogs of AT-7519. SNS-032 and AT-7519 are

published inhibitors of several members of the cyclin-dependent kinase (CDK) and cyclin-

dependent kinase-like (CDKL) families [43]. Before using these compounds as tools for

dissecting CDKL5 biology, we needed to first understand their kinome-wide selectivity. We

analyzed **B1**, **B4**, and **B12** at 1 µM using the Eurofins DiscoverX *scan*MAX panel, which

assesses binding to 403 wild-type (WT) human as well as several non-human and mutant

kinases. Percent of control (PoC) values are generated for each kinase contained within the

panel [43].

For **B1**, an analog of SNS-032, pan CDK inhibition was maintained (Figure 3A). All

kinases with PoC ≤20 in the DiscoverX *scan*MAX panel plus GSK3β were profiled using

orthogonal binding or enzymatic assays. While **B1** has a good selectivity score (S_{10}(1 µM) =

0.017), several CDKs (CDK9, PCTK1/CDK16, PCTK2/CDK17, PCTK3/CDK18) are potently

inhibited (IC_{50} ≤100 nM) (colored green in the nested table in Figure 3A). Weaker inhibition of

CDK7 as well as both GSK3α and GSK3β was noted. Curves corresponding to the affinity data

for CDKL5 and GSK3 are included in Figures S4 and S5A, respectively. As the binding pockets

share a high degree of similarly, the selectivity of **B1** within the CDKL family was further profiled

via thermal shift assays (Figure S6). The highest change in melting temperature (ΔTm, >5 $^{\circ}$C)

was noted for CDKL5, while a modest ΔTm of 3–4 $^{\circ}$C was observed for CDKL2. As a PoC value

of 48 for CDKL2 was observed when **B1** was screened at 1 µM in the DiscoverX *scan*MAX

panel, follow-up enzymatic studies were not executed for this kinase. We propose that only at

higher concentrations (>10 µM) would inhibition of this kinase become significant.

Compounds **B4** and **B12**, which are analogs of AT-7519, also demonstrated good

kinome-wide selectivity with S_{10}(1 µM) scores of 0.02 and 0.01, respectively (Figure 3).

Orthogonal profiling of **B4** confirmed high affinity binding to CDKL5, GSK3α, and GSK3β (green

in nested table in Figure 3B, curves in Figures S4 and S5B), with slightly weaker affinity for

GSK3α/β than CDKL5 evaluated via the respective cellular target engagement (NanoBRET)

assays (Figures 3, S4, and S5). Kinases with PoC ≤10 in the DiscoverX *scan*MAX panel were

profiled using orthogonal binding or enzymatic assays. An additional nine kinases demonstrated

PoC <35. Orthogonal profiling of **B12** confirmed high affinity binding to CDKL5, GSK3α, and

GSK3β (green in nested table in Figure 3C, curves in Figures S4 and S5C). This compound is a

more effective inhibitor of GSK3α when compared to GSK3β. As was the case for **B4**, **B12**

bound with greater affinity to CDKL5 than GSK3α/β in the respective NanoBRET assays

(Figures 3, S4, and S5C). Aside from GSK3α/β, a large (>100-fold) window exists between

CDKL5 and the next most potently inhibited kinase, DYRK2, when considering the enzymatic

and/or binding data.

Binding and orthogonal assay results in combination with Western blots confirmed that in

cells, **B1** is a CDKL5 and multi-CDK inhibitor with much weaker GSK3α/β affinity (>1.8 µM) and

inhibitory activity. Compounds **B4** and **B12** exhibited similar inhibition profiles and are potent

inhibitors of CDKL5, GSK3α, and GSK3β. Despite its potency, compound **B4** demonstrated

sub-optimal selectivity when compared with **B1** and **B12**. This was determined based on the

number of WT human kinases with PoC <35 in the DiscoverX *scan*MAX panel when profiled at

1 µM: **B1** = 12, **B4** = 17, **B12** = 9.

In vitro kinase activity. We employed *in vitro* kinase assays to evaluate the impact of **B1**

on CDKL5 activity (Figure 4). After preparing human WT and kinase dead (KD, CDKL5 K42R

[57, 58]) recombinant proteins, equal amounts were employed in CDKL5 activity assays (Figure

4B). We observed that, in the presence of ATP (50 µM, Figure 4A), WT CDKL5 but not KD

CDKL5 elicited kinase activity, as measured by turnover of ATP to ADP via a luminescent

reagent (ADP-Glo assay). Next, the activity of untreated WT CDKL5 was compared with WT

CDKL5 treated with vehicle (DMSO) or kinase inhibitors (10 or 100 nM). **B1**, AST-487, and

lapatinib were the inhibitors chosen. AST-487 is a confirmed, but less selective inhibitor of

CDKL5 (positive control) [51]. Lapatinib is a relatively selective kinase inhibitor that does not

inhibit CDKL5 (negative control). The difference between DMSO, lapatinib, and untreated

CDKL5 was not significant at 10 or 100 nM (Figure 4C, D). **B1** and AST-487, however, robustly

inhibited CDKL5 in a dose-dependent manner (Figure 4C, D). **B1** was confirmed to be a potent

inhibitor of CDKL5 kinase activity using this assay.

Kinase domains of GSK3β and CDKL5 have high similarity, to test the crosstalk of these

inhibitors with GSK3β, we measured phospho-β-catenin (Ser33/37/Thr41), which is directly

phosphorylated by GSK3β [19]. Phospho-β-catenin (Ser33/37/Thr41) is reduced upon exposure

to GSK3β inhibitor CHIR 99021 while EB2 pSer222, is not affected by GSK3β inhibition (Figure

S7). We measured pSer222 EB2 levels normalized to total EB2 at increasing concentrations of

inhibitors in primary neurons and compared this to phospho-β-catenin normalized to total

β–catenin (Figure 5). We found that CAF-382 (**B1**) showed a significant reduction in pSer222

EB2/total EB2 at 500 nM concentration (Figure 5A). There was a trend of pSer222 EB2/total

EB2 reduction from 5 nM, which was not significant (Figure 5A). Phospho-β-catenin/ total β-

catenin was not changed over the entire concentration range, indicating that CAF-382 (**B1)** does

not inhibit GSK3β (Figure 5A). HW2-013 (**B4)** was effective in reducing EB2 pSer222 at 500 nM

but also inhibited GSK3β significantly at 500 nM with a trend of inhibition from 5 nM, indicating

that this inhibitor affects GSK3β (Figure 5B). Finally, LY-213 (**B12**) reduced pSer222 EB2 at 500

nM but also seemed to decrease β-catenin phosphorylation at this concentration. LY-213 (**B12**)

showed comparable trends of inhibition for GSK3β and CDKL5 (Figure 5C). There were no

significant differences in total CDKL5, EB2 or β-catenin levels during these treatments (Figure

S8). We concluded that CAF-382 (**B1**) can inhibit CDKL5 activity without affecting GSK3 activity

in neurons.

*Acute CDKL5 inhibition by CAF-382 (**B1**) in rat hippocampal slices.* To determine if CAF-

382 (**B1**) could inhibit CDKL5 activity within intact brain tissue, we prepared acute hippocampal

slices from P20–30 rats and incubated them with CAF-382 (**B1**) for 2h as described. As for

primary cultures, CAF-382 (**B1**) reduced EB2 phosphorylation in a dose dependent fashion

(Figure 6). Similarly, SNS-032, the parent compound of CAF-382 (**B1**), also reduced EB2

phosphorylation in hippocampal slices (Figure S9).

Previous studies found that genetic knock-down of *Cdkl5* in primary rodent brain cultures

reduced glutamatergic excitatory synaptic transmission associated with loss of dendritic spines

[29]. In this approach, the time-course of loss of CDKL5 function is expected to occur over

several days. To clarify the more acute role of CDKL5 in regulating excitatory synaptic

transmission, we applied CAF-382 (**B1**) (10–100 nM) to hippocampal slices while measuring

fEPSPs in the CA1 hippocampus. While varying the stimulation strength applied to pre-synaptic

Schaffer-collateral pathway inputs, we were able to measure the input-output responsiveness

through measuring the input (fiber volley size) versus output (fEPSP slope). The slope of this

relationship was reduced in a dose-response relationship by CAF-382 (**B1**) within 30 minutes of

application (Figure 7A). In other words, for a similar input (fiber volley amplitude), the post-

synaptic responsiveness (fEPSP slope) was reduced (Figure 7B). This suggests that the effect

of acute CDKL5 inhibition by CAF-382 (**B1**) on glutamatergic synaptic transmission is primarily

post-synaptic. Determination that amphiphysin-1, a pre-synaptic protein, is phosphorylated by

CDKL5 [70] suggested a role of CDKL5 in presynaptic function. However, *Cdkl5* knock-out

studies showed that amphiphysin-1 was not phosphorylated by CDKL5 *in vivo*, suggesting an

alternative amphiphysin-1-independent mechanism is responsible for observed presynaptic

changes [Kontaxi et al Mike Cousin, biorxiv]. To investigate the presynaptic role of CDKL5

kinase function, the paired-pulse ratio across several paired-pulse intervals was measured

(Figure 8). CAF-382 (**B1**) (100 nM x 1 hour) had no effect on the paired-pulse ratio, confirming

the acute effects of CAF-382 (**B1**) are restricted to the post-synapse. fEPSP slopes are

mediated by AMPA-type glutamate receptors with NMDA-type glutamate receptors largely silent

under these conditions. In order to visualize the impact of CAF-382 (**B1**) on NMDA-type

glutamate receptor mediated responses, extracellular Mg^{2+} was removed from the recording

medium [64]. Under these conditions, a NMDA-receptor mediated field potential (population

spike 2) becomes apparent; this was not affected by CAF-382 (**B1**) (Figure 9) but was

completely blocked by the NMDA-type glutamate receptor antagonist D-APV (50 μM). This

result suggests that the acute effect of CAF-382 (**B1**) mediated CDKL5 inhibition is to selectively

reduce AMPA-type glutamate receptor-mediated responses post-synaptically.

A previous study found that germ-line knock-out of *Cdkl5* in the mouse enhanced long-

term potentiation (LTP) at 7–13 week old (wo) hippocampal Schaffer Collateral to CA1

synapses without alteration of the input-output relationship [38]. In another study, germ-line

knock-out of *Cdkl5* in a mouse at 4–5 wo did not affect stable LTP at this synapse [37]. In a rat

Cdkl5 knock-out, LTP at this synapse was selectively increased at 3–4 wo but not at later ages

[40]. To clarify the more acute role of CDKL5 in regulating LTP at this synapse, we incubated 3–

4 wo hippocampal slices with CAF-382 (**B1)** (100 nM) for at least 30 minutes and, following

stable baseline measurements of fEPSP slope, induced LTP via theta burst to Schaffer

Collateral to CA1 synapses in the continuous presence of CAF-382 (**B1)**. Compared to

interleaved recordings from control slices, CAF-382 (**B1)** significantly reduced LTP at 60

minutes post theta-burst (Figure 10). To investigate this further, LTP at these synapses of

isolated AMPA-type glutamate receptors via a pairing protocol was measured using whole-cell

patch-clamp. In this case, hippocampal slices were incubated with CAF-382 (**B1)** (100 nM) for

only one hour prior to, but not during, recording. Compared to interleaved recordings from

control slices, CAF-382 (**B1)** significantly reduced LTP at 30 minutes post pairing (Figure 11).

The induction and expression of LTP includes the contributions of pre-synaptic and post-

synaptic mechanisms at both excitatory and inhibitory synapses[71, 72]. Since we do not see a

presynaptic effect of CAF-382 (**B1)** (Figure 8), and the approach (Fig. 11) removed the

contribution of inhibitory synapses, this supports the direct role of CDKL5 in the post-synaptic

mediation of LTP at this synapse at this age.

Evaluation of suitability of CDKL5 lead for in vivo studies. As our studies were all

executed *in vitro*, the suitability of **B1** for use *in vivo* was unknown. To gauge whether **B1** could

be dosed in animals using a water-based formulation, we first analyzed its aqueous kinetic

solubility (Table 1). **B1** demonstrated good solubility (196 µM), supporting that it readily

dissolves in aqueous buffer without observed precipitation. Mouse liver microsomal stability data

was also collected (Table 1). **B1** proved to be very tolerant to microsomal exposure and

demonstrated excellent stability (>85%) after 30 minutes. Since compound **B1** was found to

have the required aqueous solubility and mouse liver microsomal stability, it was submitted for

in vivo characterization. Pharmacokinetic analysis and brain exposure of mice following

intraperitoneal administration with **B1** were next examined. Plasma was sampled 0.5-, 1-, 2-,

and 4-hours post-dose with 2.29 mg/kg of a salt form of **B1** (Figure S10). No abnormal clinical

symptoms were observed in dosed animals. The half-life of **B1** was found to be <1h, with the

highest concentration of drug in the plasma found 0.5 h after dosing (Cmax = 508 ng/mL) and

total system exposure (AUCinf) of 646 h*ng/mL (Figure S10). The same dose (2.29 mg/kg) as

well as a 7.63 mg/kg dose were used to analyze the brain penetration of **B1** in mice 1h after

administration. As shown in Table S2, brain and plasma sampling indicated that brain

penetration of **B1** was generally low.

Discussion

CDD is a rare, devastating, genetically mediated neurological disorder with symptomatic

onset in very early infancy. Though there are newly approved therapies that provide some relief

for epilepsy [12], the additional features that are impactful to caregivers (Jeff Neul, personal

communication) of motor, cognitive, visual and autonomic disturbances [2, 6-9, 10] are not

currently addressed. CDKL5 protein [73] and *CDKL5* gene [74] replacement approaches are

advancing   through   pre-clinical   studies   to   potential   first-in-human   trials

(https://www.ultragenyx.com/our-research/pipeline/ux055-for-cdd/). While CDKL5 is a key

mediator of synaptic and network development and physiology, the precise role of this kinase as

explored through genetic manipulations in rodent models remains unclear. This knowledge gap

prevents a full understanding of the role of CDKL5 across development and has implications for

determining and interpreting when protein and gene replacement strategies may be most

efficacious. As CDKL5 kinase function is central to disease pathology [13, 16], our development

of a specific CDKL5 kinase inhibitor aims to address this gap.

Following identification of existing inhibitors [69], detailed analysis of the activity of

analogs of these inhibitors in a range of assays identified three compounds with significant

selectivity over GSK3β, a key requirement due to structural homology [43] and linkage of

GSK3β to both synaptic physiology [44] and CDKL5-related signaling [22, 23]. The lead

compound, **CAF-382 (B1**), demonstrated potent activity in multiple assays, including target-

based *in vitro* biochemical assays and functional assays utilizing both rodent neuronal cultures

and brain slices. The results across all assays are consistent, and align with the hypothesis that

compound **B1** inhibits CDKL5 and blocks phosphorylation of a known CDKL5 substrate, EB2

[19, 31, 32]. Several cyclin dependent kinases (CDKs) that are potently (IC_{50} ≤100 nM) inhibited

(CDK9 [75], PCTK1/CDK16 [76], PCTK2/CDK17 [77], and PCTK3/CDK18 [78]) by **B1** are

expressed in the brain (http://mouse.brain-map.org). However, none of these have been

demonstrated to have synaptic activity or localization; each are largely involved in replication

and nuclear functions. At issue, is whether observations of CDKL5 activity, either through

genetic manipulations or interference with the kinase, are specific and not potentially subsumed

by the activity of other kinases. Given the reduction of phosphorylation of the specific substrate,

EB2, this is less likely here.

The biochemical activity of **CAF-382** (**B1**) against CDKL5 is <5 nM, based on *in vitro*

activity studies (Figure 4), and for studies investigating cellular target engagement as well as

activity in neuronal cultures and brain slices at the specific substrate, EB2, and synaptic

physiology we observe a 10-fold shift in potency (~50 nM). A loss in potency, albeit modest,

when moving from a biochemical to cell-based assay is commonly observed, mostly due to

suboptimal cell penetrance of the small molecule, and has been observed for several orthogonal

chemical series of kinase inhibitors[47, 48, 50]. As the *in vitro* assay eliminates the contribution

of phosphatases (and the phosphatases specific for EB2 are currently unknown), the apparent

mismatch could also be due to residual presence of phosphatase-dependent and previously

phosphorylated EB2 balanced with active phosphorylation by CDKL5. Nevertheless, CDKL5

kinase function is clearly necessary for the induction and/or expression long-term potentiation,

adding to the very short list of kinases required for this essential process linked to learning and

memory[71, 72, 79] and finally clarifying previous genetic studies. The increased LTP seen in

genetic studies suggests that observations of increased LTP in these models is compensatory

and not directly due to loss of CDKL5. One interpretation of those prior studies is that CDKL5

provided a limiting "brake" on LTP, with genetic loss resulting in more LTP which is in

contradistinction to our findings. The role of CDKL5 at inhibitory synapses was not specifically

examined. Nevertheless, synaptic function at the developmental age studied (P20–30) requires

CDKL5 activity for synaptic plasticity, suggesting that strategies that boost function at this age-

equivalent in humans may be clinically important. Selective stability and plasticity dependent

insertion of AMPA-type glutamate receptors at CA1 hippocampal synapses likely require a

balance between active CDKL5 kinase and phosphatases. Further studies are needed to

determine this balance more specifically and continue to clarify CDKL5 function at pre-

synapses[80].

While no clinical abnormalities were observed with *in vivo* administration of CAF-382

(**B1**), this is likely due to low brain penetration. Due to low brain penetration of CAF-382 (**B1**),

strategies that optimize brain penetration or different administration routes could be considered

in future studies investigating the role of CDKL5 *in vivo*. These studies, behavioral and perhaps

with electrical assessments of seizure activity, may clarify the apparent mismatch of rodent

knock-out with clinical observations. Using CDKL5 inhibitors *in vivo* across developmental

stages can help ascertain possible therapeutic windows for CDD. More broadly, CDKL5

inhibition is beneficial for cell survival upon ischemia or nephrotoxin induced kidney injury[58,

81], suggesting that the inhibitors we characterized here may prove useful to evaluate the

promise of CDKL5 inhibition in other disease models.

Contributions: TAB, SKU and ADA conceived the project. CAF, HWO, YL, FMB, IMG, and

DHD designed and synthesized compounds. WR and ANB executed CDKL family thermal shift

assays. AC, MS, CIW, JAS, JLS, KD, DHD, NSP, ANB, TAB, SKU, and ADA conducted

experiments and analyses. TAB, SKU and ADA drafted the paper, and all contributed to

revisions.

References

1. Weaving, L.S., et al., *Mutations of CDKL5 cause a severe neurodevelopmental disorder with infantile spasms and mental retardation.* Am.J Hum.Genet., 2004. **75**(6): p. 1079-1093.

2. Bahi-Buisson, N., et al., *Recurrent mutations in the CDKL5 gene: genotype-phenotype relationships.* Am.J Med.Genet.A, 2012. **158A**(7): p. 1612-1619.

3. Lindy, A.S., et al., *Diagnostic outcomes for genetic testing of 70 genes in 8565 patients with epilepsy and neurodevelopmental disorders.* Epilepsia, 2018. **59**(5): p. 1062-1071.

4. Kothur, K., et al., *Diagnostic yield of targeted massively parallel sequencing in children with epileptic encephalopathy.* Seizure, 2018. **59**: p. 132-140.

5. Symonds, J.D., et al., *The Genetic and Autoimmune Childhood Epilepsy (GACE) Study*, in *American Epilpesy Society.* 2017: Washington, D.C.

6. Bahi-Buisson, N., et al., *Key clinical features to identify girls with CDKL5 mutations.* Brain, 2008. **131**(Pt 10): p. 2647-2661.

7. Nemos, C., et al., *Mutational spectrum of CDKL5 in early-onset encephalopathies: a study of a large collection of French patients and review of the literature.* Clin.Genet., 2009. **76**(4): p. 357-371.

8. Castren, M., et al., *Epilepsy caused by CDKL5 mutations.* Eur J Paediatr Neurol, 2011. **15**(1): p. 65-9.

9. Melani, F., et al., *CDKL5 gene-related epileptic encephalopathy: electroclinical findings in the first year of life.* Developmental Medicine and Child Neurology, 2011. **53**(4): p. 354-360.

10. Fehr, S., et al., *There is variability in the attainment of developmental milestones in the CDKL5 disorder.* J Neurodev Disord, 2015. **7**(1): p. 2.

11. Muller, A., et al., *Retrospective evaluation of low long-term efficacy of antiepileptic drugs and ketogenic diet in 39 patients with CDKL5-related epilepsy.* Eur J Paediatr Neurol, 2016. **20**(1): p. 147-51.

12. Knight, E.M.P., et al., *Safety and efficacy of ganaxolone in patients with CDKL5 deficiency disorder: results from the double-blind phase of a randomised, placebo-controlled, phase 3 trial.* Lancet Neurol, 2022. **21**(5): p. 417-427.

13. Demarest, S.T., et al., *CDKL5 deficiency disorder: Relationship between genotype, epilepsy, cortical visual impairment, and development.* Epilepsia, 2019.

14. Wong, K., et al., *Factors influencing the attainment of major motor milestones in CDKL5 deficiency disorder.* Eur J Hum Genet, 2022.

15. Pini, G., et al., *Variant of Rett syndrome and CDKL5 gene: clinical and autonomic description of 10 cases.* Neuropediatrics, 2012. **43**(1): p. 37-43.

16. Hector, R.D., et al., *CDKL5 variants: Improving our understanding of a rare neurologic disorder.* Neurol Genet, 2017. **3**(6): p. e200.

17. Chen, Q., et al., *CDKL5, a protein associated with rett syndrome, regulates neuronal morphogenesis via Rac1 signaling.* J Neurosci, 2010. **30**(38): p. 12777-86.

18. Hector, R.D., et al., *Characterisation of CDKL5 Transcript Isoforms in Human and Mouse.* PLoS One, 2016. **11**(6): p. e0157758.

19. Baltussen, L.L., et al., *Chemical genetic identification of CDKL5 substrates reveals its role in neuronal microtubule dynamics.* EMBO J, 2018.

20. Zhu, Y.C. and Z.Q. Xiong, *Molecular and Synaptic Bases of CDKL5 Disorder.* Dev Neurobiol, 2018.

21. Wang, I.T., et al., *Loss of CDKL5 disrupts kinome profile and event-related potentials leading to autistic-like phenotypes in mice.* Proc.Natl.Acad.Sci.U.S.A, 2012. **109**(52): p. 21516-21521.

22. Fuchs, C., et al., *Inhibition of GSK3beta rescues hippocampal development and learning*
in a mouse model of CDKL5 disorder. Neurobiol Dis, 2015. **82**: p. 298-310.
23. Fuchs, C., et al., *Loss of CDKL5 impairs survival and dendritic growth of newborn*
neurons by altering AKT/GSK-3beta signaling. Neurobiol Dis, 2014. **70**: p. 53-68.
24. Schroeder, E., et al., *Neuron-Type Specific Loss of CDKL5 Leads to Alterations in*
mTOR Signaling and Synaptic Markers. Mol Neurobiol, 2019. **56**(6): p. 4151-4162.
25. Ren, E., et al., *Functional and Structural Impairments in the Perirhinal Cortex of a Mouse*
Model of CDKL5 Deficiency Disorder Are Rescued by a TrkB Agonist. Front Cell
Neurosci, 2019. **13**: p. 169.
26. Rusconi, L., et al., *CDKL5 expression is modulated during neuronal development and its*
subcellular distribution is tightly regulated by the C-terminal tail. J Biol Chem, 2008.
283(44): p. 30101-11.
27. Oi, A., et al., *Subcellular distribution of cyclin-dependent kinase-like 5 (CDKL5) is*
regulated through phosphorylation by dual specificity tyrosine-phosphorylation-regulated
kinase 1A (DYRK1A). Biochem Biophys Res Commun, 2017. **482**(2): p. 239-245.
28. Zhu, Y.C., et al., *Palmitoylation-dependent CDKL5-PSD-95 interaction regulates*
synaptic targeting of CDKL5 and dendritic spine development. Proc Natl Acad Sci U S A,
2013. **110**(22): p. 9118-23.
29. Ricciardi, S., et al., *CDKL5 ensures excitatory synapse stability by reinforcing NGL-1-*
PSD95 interaction in the postsynaptic compartment and is impaired in patient iPSC-
derived neurons. Nat.Cell Biol., 2012. **14**(9): p. 911-923.
30. Nawaz, M.S., et al., *CDKL5 and Shootin1 Interact and Concur in Regulating Neuronal*
Polarization. PLoS One, 2016. **11**(2): p. e0148634.
31. Terzic, B., et al., *Temporal manipulation of Cdkl5 reveals essential postdevelopmental*
functions and reversible CDKL5 deficiency disorder-related deficits. J Clin Invest, 2021.
131(20).
32. Di Nardo, A., et al., *Phenotypic characterization of Cdkl5-knockdown neurons*
establishes elongated cilia as a functional assay for CDKL5 Deficiency Disorder.
Neurosci Res, 2022. **176**: p. 73-78.
33. Uezu, A., et al., *Identification of an elaborate complex mediating postsynaptic inhibition.*
Science, 2016. **353**(6304): p. 1123-9.
34. Della Sala, G., et al., *Dendritic Spine Instability in a Mouse Model of CDKL5 Disorder Is*
Rescued by Insulin-like Growth Factor 1. Biol Psychiatry, 2016. **80**(4): p. 302-11.
35. Negraes, P.D., et al., *Altered network and rescue of human neurons derived from*
individuals with early-onset genetic epilepsy. Mol Psychiatry, 2021. **26**(11): p. 7047-
7068.
36. Lupori, L., et al., *Site-specific abnormalities in the visual system of a mouse model of*
CDKL5 deficiency disorder. Hum Mol Genet, 2019. **28**(17): p. 2851-2861.
37. Yennawar, M., R.S. White, and F.E. Jensen, *AMPA Receptor Dysregulation and*
Therapeutic Interventions in a Mouse Model of CDKL5 Deficiency Disorder. J Neurosci,
2019. **39**(24): p. 4814-4828.
38. Okuda, K., et al., *CDKL5 controls postsynaptic localization of GluN2B-containing NMDA*
receptors in the hippocampus and regulates seizure susceptibility. Neurobiol Dis, 2017.
106: p. 158-170.
39. Tang, S., et al., *Loss of CDKL5 in Glutamatergic Neurons Disrupts Hippocampal*
Microcircuitry and Leads to Memory Impairment in Mice. J Neurosci, 2017. **37**(31): p.
7420-7437.
40. de Oliveira, L.S., et al., *Enhanced hippocampal LTP but typical NMDA receptor and*
AMPA receptor function in a novel rat model of CDKL5 deficiency disorder. bioRxiv,
2022: p. 2022.06.29.497927.

41. Amendola, E., et al., *Mapping pathological phenotypes in a mouse model of CDKL5 disorder.* PLoS One, 2014. **9**(5): p. e91613.
42. Manning, G., et al., *The protein kinase complement of the human genome.* Science, 2002. **298**(5600): p. 1912-34.
43. Davis, M.I., et al., *Comprehensive analysis of kinase inhibitor selectivity.* Nat Biotechnol, 2011. **29**(11): p. 1046-51.
44. Peineau, S., et al., *The role of GSK-3 in synaptic plasticity.* Br.J Pharmacol, 2008. **153 Suppl 1**: p. S428-S437.
45. Wada, A., *Lithium and neuropsychiatric therapeutics: neuroplasticity via glycogen synthase kinase-3beta, beta-catenin, and neurotrophin cascades.* J Pharmacol Sci, 2009. **110**(1): p. 14-28.
46. Wells, C., et al., *SGC-AAK1-1: A Chemical Probe Targeting AAK1 and BMP2K.* ACS Med Chem Lett, 2020. **11**(3): p. 340-345.
47. Wells, C.I., et al., *Development of a potent and selective chemical probe for the pleiotropic kinase CK2.* Cell Chem Biol, 2021. **28**(4): p. 546-558 e10.
48. Drewry, D.H., et al., *Identification and Utilization of a Chemical Probe to Interrogate the Roles of PIKfyve in the Lifecycle of beta-Coronaviruses.* J Med Chem, 2022. **65**(19): p. 12860-12882.
49. Jester, B.W., et al., *A coiled-coil enabled split-luciferase three-hybrid system: applied toward profiling inhibitors of protein kinases.* J Am Chem Soc, 2010. **132**(33): p. 11727-35.
50. Drewry, D.H., et al., *Identification of Pyrimidine-Based Lead Compounds for Understudied Kinases Implicated in Driving Neurodegeneration.* J Med Chem, 2022. **65**(2): p. 1313-1328.
51. Canning, P., et al., *CDKL Family Kinases Have Evolved Distinct Structural Features and Ciliary Function.* Cell Rep, 2018. **22**(4): p. 885-894.
52. Winter, G., C.M. Lobley, and S.M. Prince, *Decision making in xia2.* Acta Crystallogr D Biol Crystallogr, 2013. **69**(Pt 7): p. 1260-73.
53. Winter, G., et al., *DIALS: implementation and evaluation of a new integration package.* Acta Crystallogr D Struct Biol, 2018. **74**(Pt 2): p. 85-97.
54. Liebschner, D., et al., *Macromolecular structure determination using X-rays, neutrons and electrons: recent developments in Phenix.* Acta Crystallogr D Struct Biol, 2019. **75**(Pt 10): p. 861-877.
55. Emsley, P. and K. Cowtan, *Coot: model-building tools for molecular graphics.* Acta Crystallogr D Biol Crystallogr, 2004. **60**(Pt 12 Pt 1): p. 2126-32.
56. Fedorov, O., et al., *A systematic interaction map of validated kinase inhibitors with Ser/Thr kinases.* Proc Natl Acad Sci U S A, 2007. **104**(51): p. 20523-8.
57. Lin, C., B. Franco, and M.R. Rosner, *CDKL5/Stk9 kinase inactivation is associated with neuronal developmental disorders.* Hum Mol Genet, 2005. **14**(24): p. 3775-86.
58. Kim, J.Y., et al., *A kinome-wide screen identifies a CDKL5-SOX9 regulatory axis in epithelial cell death and kidney injury.* Nat Commun, 2020. **11**(1): p. 1924.
59. Rosianu, F., et al., *Loss of NDR1/2 kinases impairs endomembrane trafficking and autophagy leading to neurodegeneration.* Life Sci Alliance, 2023. **6**(2).
60. Cornejo, B.J., et al., *A single episode of neonatal seizures permanently alters glutamatergic synapses.* Ann Neurol, 2007. **61**(5): p. 411-426.
61. Bernard, P.B., et al., *Necessary, but not sufficient: Insights into the mechanisms of mGluR mediated long-term depression from a rat model of early life seizures.* Neuropharmacology, 2014. **84**: p. 1-12.
62. Bernard, P.B., et al., *Phosphorylation of FMRP and alterations of FMRP complex underlie enhanced mLTD in adult rats triggered by early life seizures.* Neurobiol.Dis., 2013. **59**: p. 1-17.

63.   Anderson, W.W. and G.L. Collingridge, *The LTP program: a data acquistion program for on-line analysis of long-term potentiation and other synaptic events.* J Neurosci Methods, 2001. **108**: p. 71-83.

64.   Coan, E.J. and G.L. Collingridge, *Magnesium ions block an N-methyl-D-aspartate receptor-mediated component of synaptic transmission in rat hippocampus.* Neuroscience Letters, 1985. **53**: p. 21-26.

65.   Sanderson, J.L., et al., *beta-Amyloid disruption of LTP/LTD balance is mediated by AKAP150-anchored PKA and Calcineurin regulation of Ca(2+)-permeable AMPA receptors.* Cell Rep, 2021. **37**(1): p. 109786.

66.   Sanderson, J.L., J.A. Gorski, and M.L. Dell'Acqua, *NMDA Receptor-Dependent LTD Requires Transient Synaptic Incorporation of Ca(2)(+)-Permeable AMPARs Mediated by AKAP150-Anchored PKA and Calcineurin.* Neuron, 2016. **89**(5): p. 1000-15.

67.   Osterweil, E.K., et al., *Hypersensitivity to mGluR5 and ERK1/2 leads to excessive protein synthesis in the hippocampus of a mouse model of fragile X syndrome.* J Neurosci, 2010. **30**(46): p. 15616-15627.

68.   Grosshans, D.R. and M.D. Browning, *Protein kinase C activatio induces tyrosine phosphorylation of the NR2A and NR2B subunits of the NMDA receptor.* J Neurochem, 2001. **76**: p. 737-744.

69.   Vasta, J.D., et al., *Quantitative, Wide-Spectrum Kinase Profiling in Live Cells for Assessing the Effect of Cellular ATP on Target Engagement.* Cell Chem Biol, 2018. **25**(2): p. 206-214 e11.

70.   Sekiguchi, M., et al., *Identification of amphiphysin 1 as an endogenous substrate for CDKL5, a protein kinase associated with X-linked neurodevelopmental disorder.* Arch Biochem Biophys, 2013. **535**(2): p. 257-67.

71.   Collingridge, G.L. and W.C. Abraham, *Glutamate receptors and synaptic plasticity: The impact of Evans and Watkins.* Neuropharmacology, 2022. **206**: p. 108922.

72.   Nicoll, R.A., *A Brief History of Long-Term Potentiation.* Neuron, 2017. **93**(2): p. 281-290.

73.   Trazzi, S., et al., *CDKL5 protein substitution therapy rescues neurological phenotypes of a mouse model of CDKL5 disorder.* Hum Mol Genet, 2018. **27**(9): p. 1572-1592.

74.   Gao, Y., et al., *Gene replacement ameliorates deficits in mouse and human models of cyclin-dependent kinase-like 5 disorder.* Brain, 2020. **(in press)**.

75.   Yu, D.S. and D. Cortez, *A role for CDK9-cyclin K in maintaining genome integrity.* Cell Cycle, 2011. **10**(1): p. 28-32.

76.   Mikolcevic, P., J. Rainer, and S. Geley, *Orphan kinases turn eccentric: a new class of cyclin Y-activated, membrane-targeted CDKs.* Cell Cycle, 2012. **11**(20): p. 3758-68.

77.   Liu, M., et al., *The Identification of Key Genes and Pathways in Glioma by Bioinformatics Analysis.* J Immunol Res, 2017. **2017**: p. 1278081.

78.   de Oliveira Pepino, R., et al., *Overview of PCTK3/CDK18: A Cyclin-Dependent Kinase Involved in Specific Functions in Post-Mitotic Cells.* Curr Med Chem, 2021. **28**(33): p. 6846-6865.

79.   Giese, K.P. and K. Mizuno, *The roles of protein kinases in learning and memory.* Learn Mem, 2013. **20**(10): p. 540-52.

80.   Kontaxi, C., et al., *Epilepsy-related CDKL5 deficiency slows synaptic vesicle endocytosis in central nerve terminals.* J Neurosci, 2023.

81.   Kim, J.Y., et al., *Involvement of the CDKL5-SOX9 signaling axis in rhabdomyolysis-associated acute kidney injury.* Am J Physiol Renal Physiol, 2020. **319**(5): p. F920-F929.

Figure 1. Screening of CDKL5 inhibitors in rat primary neurons using Western blotting. Western blots showing expression of total EB2 and levels of Ser222 EB2 phosphorylation in DIV14-16 rat primary neurons upon treatment of 1 hour with 5 nM, 50 nM, 500 nM and 5000 nM of selected CDKL5 inhibitors.

919 **Figure 2.** Structures of CDKL5 inhibitor leads and corresponding parent compounds.

A

B1

$S_{10}(1\ \mu M) = 0.017$

Kinase	DiscoverX PoC value	Assay format	IC$_{50}$ value (nM)
GSK3A	0.7	Enzymatic and NB	470 and 1900
CDKL5	0.7	Binding and NB	6.7 and 11
PCTK2	1.3	Enzymatic	89
CDK7	1.9	Enzymatic	300
PCTK1	5.5	Enzymatic	62
CDC2L5	5.9	Enzymatic	>10000
CDK9	8.3	Enzymatic	20
YSK4	14	Enzymatic	>10000
PCTK3	17	Enzymatic	100
CDK4	20	Enzymatic	1900
GSK3B	51	Enzymatic and NB	2200 and 4500

B

B4

$S_{10}(1\ \mu M) = 0.02$

Kinase	DiscoverX PoC value	Assay format	IC$_{50}$ value (nM)
GSK3A	0	Enzymatic and NB	11 and 120
CDKL5	0.2	Binding and NB	6.4 and 17
GSK3B	2.1	Enzymatic and NB	38 and 250
DYRK2	4.4	Enzymatic	240
CDK7	4.4	Enzymatic	>10000
PCTK1	6.3	Enzymatic	1200
DYRK1A	9.1	Enzymatic	900
PCTK2	9.5	Enzymatic	1600

C

B12

$S_{10}(1\ \mu M) = 0.01$

Kinase	DiscoverX PoC value	Assay format	IC$_{50}$ value (nM)
GSK3A	0	Enzymatic and NB	16 and 190
CDKL5	0.6	Binding and NB	8.7 and 23
GSK3B	1.7	Enzymatic and NB	48 and 1200
DYRK2	4.6	Enzymatic	920
CDK7	17	Enzymatic	>10000
PCTK1	19	Enzymatic	1700
RPS6KA4	27	Enzymatic	>10000
DYRK1A	29	Enzymatic	4800
DYRK1B	34	Enzymatic	1700

Figure 3. Kinome-wide selectivity data for CDKL5 lead compounds. Kinome tree diagrams illustrate the selectivity of these
compounds when profiled against 403 wild-type (WT) human kinases at 1 µM at Eurofins DiscoverX in their *scan*MAX panel. The
percent control legend shows red circles of different sizes corresponding with percent control value each kinase binds the small
molecule in this large binding panel. Also included are selectivity scores ($S_{10}(1\ \mu M)$), which were calculated using the PoC values for
WT human kinases in the *scan*MAX panel only. The S_{10} score is a way to express selectivity that corresponds with the percent of the
kinases screened that bind with a PoC value <10. In the embedded tables kinases are listed by their gene names and ranked by their
percent of control (PoC) value generated in the *scan*MAX panel. Rows colored green demonstrate enzymatic IC$_{50}$ values within a 30-
fold window of the CDKL5 binding IC$_{50}$ value. Correlating assay format used to generate data in the last column of nested table is
listed in the preceding column, with NB = NanoBRET.

Figure 4. B1 potently inhibits the kinase activity of CDKL5. (A) Representative graph from a CDKL5 kinase assay demonstrating that purified WT human CDKL5 retains kinase activity, while the kinase dead (KD, CDKL5 K42R) human protein is functionally inactive (n = 3). (B) Representative western blot illustrating that equal expression of WT and KD proteins was observed in the kinase assay experiments. (C and D) Kinase assays using purified WT human CDKL5 in the presence of 10 or 100 nM of the indicated compounds (n = 3). One-way ANOVA with Dunnett's multiple comparison test used. *** = p <0.0001 and non-significant comparisons not shown.

Figure 5. CAF-382 (**B1**), HW2-013 (**B4**), and LY-213 (**B12**) compounds reduce CDKL5 activity in rat primary neurons. HW2-013 (**B4**) and LY-213 (**B12**) also downregulate GSK3 activity. (A) Western blot and quantification showing expression of EB2 phosphorylation and β-catenin phosphorylation in DIV14-15 rat primary neurones after an hour treatment with different concentrations of CAF-382 (**B1**). (B) Western blot and quantification showing expression of EB2 phosphorylation and β-catenin phosphorylation in DIV14-15 rat primary neurons after an hour treatment with different concentrations of HW2-013 (**B4**). (C) Western blot and quantification showing expression of EB2 phosphorylation and β-catenin phosphorylation in DIV14-15 rat primary neurons after an hour treatment with different concentrations of LY-213 (**B12**). Each concentration was compared to the control using a Kruskal-Wallis test. n = 3 biological replicates with 2 repetitions. *p≤0.05. **p≤0.01. ***p≤0.001. Error bars are SD.

952
Figure 6. CAF-382 (**B1**) reduced phosphorylation of EB2 in CA1 hippocampal slices. (A)
Example blots demonstrate no alterations in CDKL5 expression across treatments (stats in text)
and suggest relative CAF-382 (**B1**) dose-dependent differences in EB2 phosphorylation. (B)
Normalized quantification of EB2-phosphorylation (density of EB2-phosphorylation bands/
density of EB2-total bands). CAF 382 (**B1**) (control: 1±0.10 versus 100 nM: 0.49±0.02, n = 8, p
= 0.002, RM-ANOVA) (control: 1±0.08 versus 45 nM: 0.16±0.03, n = 10, p < 0.001, RM-ANOVA)
(control: 1± 0.14 versus 10 nM: 0.62± 0.11, n=7, p = 0.003, RM-ANOVA) reduced EB2
phosphorylation at all concentrations.
961

Figure 7. CAF-382 (**B1**) reduced post-synaptic fEPSP responsiveness, as measured by input-output curves. (A) Sample input (fiber volley) / output (fEPSP slope) (I/O) curves at baseline, 30 minutes and 60 minutes after CAF-382 (**B1**) (100nM) in a hippocampal slice. Slopes of I/O curves were obtained at baseline, and 30 minutes and 60 minutes after treatment with CAF-382 (**B1**). Asterisks indicate sample traces (B) from treated slice; stimulus artifact (arrow) has been removed for clarity. Initial negative deflection after the artifact is the fiber volley, followed by the fEPSP. (C) RM-ANOVA of I/O slopes of CAF-382 (**B1**) (100 nM and 45 nM) were significantly decreased after 30 minutes (100 nM: 1.91±0.23, n = 9, p = 0.006; 45 nM: 1.58±0.24, n=10, p = 0.013) and 60 minutes (100 nM: 2.00±0.30, n = 9, p = 0.001; 45 nM: 1.57±0.22, n=10, p = 0.009) compared to baseline (100 nM: 3.24±0.52, n = 9; 45 nM: 2.29±0.23, n=10). Baseline I/O slope was used to normalize each recording to allow comparisons across all treatments. CAF-382 (**B1**) (10 nM) did not alter I/O slopes after 30 or 60 minutes (RM-ANOVA, n = 9, p = 0.07). Error bars are SE.

Figure 8. CAF-382 (**B1**) did not alter presynaptic release as measured by paired-pulse ratio. (A)
Paired-pulse ratios (PPR) of fEPSPs (Slope of fEPSP2/Slope of fEPSP1), reflective of pre-
synaptic release, were unaltered across a range of stimulus intervals (RM-ANOVA, n=7) by
CAF-382 (**B1**) (100nM) for 60 minutes compared to baseline. (B) Sample trace for 50 ms
interval. Reduction of initial fEPSP slopes were consistent with Figure 7. Stimulus artifacts
(arrows) have been removed for clarity. Error bars are SE.
983

Figure 9. CAF-382 (**B1**) did not alter NMDAR-mediated component of fEPSPs. Maximal fEPSPs recorded in stratum pyramidale of CA1 with reduced extracellular Mg^{2+} demonstrate multiple population spikes; previous studies have demonstrated that secondary population spikes in these conditions are NMDAR-dependent[64]. CAF-382 (**B1**) (100 nM) did not alter population spike 1 (control -4.057±0.584mV, CAF-382 (**B1**) 3.929±0.736mV, n=10, p=0.742, RM-ANOVA) or NMDAR-mediated population spike 2 (control -2.093±0.568mV, CAF-382 (**B1**) -1.656±0.4442, n=10, p=0.867, RM-ANOVA). For comparison, the selective NMDAR antagonist D-APV (50 µM) completely blocked population spike 2, as previously reported [64]. Stimulation artifact has been removed (arrow) for clarity.

995
Figure 10. CAF-382 (**B1**) acutely reduces the expression of LTP in CA1 hippocampus.
Hippocampal slices were initially incubated in CAF-382 (**B1**) for at least 30 minutes and then
continuously perfused with CAF-382 (**B1**) (100 nM). fEPSP slope was normalized to initial
baseline, expressed as 100% (dotted line); mean±SEM shown. Following stable baseline of
fEPSP slope (> 20 minutes), LTP was induced by two theta-burst (TBS) trains. Compared to
interleaved control slices, CAF-382 (**B1**) significantly reduced LTP as measured by fEPSP slope
at 56-60 minutes post theta-burst (control: 115.6±2.2 n = 13 slices, 12 rats; CAF-382 (**B1**):
106.4±3.5, n = 5 slices, 4 rats; p = 0.029, 2-way ANOVA, Holm-Sidak post-hoc). Sample traces
for control (black filled circle) and CAF-382 (**B1**) (red filled circle) before (1, solid trace) and after
LTP (2, dotted trace) are shown to the right. Error bars are SE.

1006

1007
Figure 11. CAF-382 (**B1**) acutely reduces the expression of LTP mediated by AMPA-type
glutamate receptors in CA1 hippocampus. Hippocampal slices were initially incubated in CAF-
382 (**B1**) (100 nM) for at least 60 minutes prior to recording. Peak negative current of AMPA-
type glutamate receptor mediated synaptic responses were normalized to initial baseline post
break-in (time = 0), expressed as 100% (dotted line); mean±SEM shown. Following baseline,
LTP was induced by a pairing protocol. Compared to interleaved control slices, CAF-382 (**B1**)
significantly reduced LTP at 27-31 minutes post break-in (control: 197.5±8.4 n = 14 slices, 14
rats; CAF-382 (**B1**): 123.14±9.0, n = 12-13 slices, 13 rats; p < 0.001, 2-way ANOVA, Holm-
Sidak post-hoc). Sample traces for control (black filled circle) and CAF-382 (**B1**, red filled circle)
before (1, solid trace) and after LTP (2, dotted trace) are shown to the right. Error bars are SE.

1018

1019

1020 **Table 1**. Solubility and microsomal stability data for **B1**

Compound	Kinetic solubility		Mouse liver microsomal stability (%)		
	µM	µg/mL	T = 0 min	T = 30 min	T = 30 min (-NADPH)
B1	196.8[a]	72.1[a]	100	86.1	87.5

1021 [a] Measured solubility is >75% dose concentration, actual solubility may be higher.
1022

1023
1024

 Supplementary Information

Synthesis of key compounds. Chemistry General Information. Reagents were purchased

from reputable commercial vendors and used in accordance with safety data sheets. A rotary

evaporator was employed to remove solvent(s) under reduced pressure (*in vacuo*). Thin layer

chromatography (TLC) as well as LC–MS were used to monitor reaction progress. The following

abbreviations are used in experimental procedures: mmol (millimoles), μmol (micromoles), μL

(microliters), mg (milligrams), equiv (equivalent(s)), min (minutes), and h (hours). ^{1}H NMR and

^{13}C NMR spectra as well as microanalytical data were collected for a key intermediate and final

compound **B1** to confirm their identity and evaluate their purity prior to initiating cell- and animal-

based studies. ^{1}H and ^{13}C NMR spectra were collected in DMSO-d_6 using Bruker

spectrometers. Magnet strength is indicated in each corresponding line listing. Peak positions

are included in parts per million (ppm) and calibrated versus the shift of DMSO-d_6; coupling

constants (*J* values) are noted in hertz (Hz); and multiplicities are included as follows: singlet

(s), doublet (d), pentet of doublets (pd), and multiplet (m).

1040

1041   **Int 1**   **B1**

1042 **Scheme 1**. Preparation of compound **B1**. a) EtOH, NaBH$_4$, acetone, 80 °C, 1 h, 57% yield; b)

1043 HATU, DIPEA, 1-(*tert*-butoxycarbonyl)piperidine-4-carboxylic acid, DMF; c) TFA, CH$_2$Cl$_2$, 36%

1044 yield over two steps.

1045

*5-(((5-isopropyloxazol-2-yl)methyl)thio)thiazol-2-amine (**Int 1**)*. To a flask was added 5-

thiocyanatothiazol-2-amine (400 mg, 1.0 equiv, 2.5 mmol) in EtOH (24 mL), followed by addition

of NaBH$_4$ (183 mg, 1.9 equiv, 4.8 mmol) portion-wise at room temperature. The mixture was

stirred for 1 h at room temperature then acetone (12 mL) was added. After 1 hr, a solution of 2-

(chloromethyl)-5-isopropyloxazole (412 mg, 1.0 equiv, 2.6 mmol) in EtOH (4mL) was added,

and the reaction was heated at 80 °C for 1 h. The resulting mixture was cooled, concentrated *in*

vacuo, and then partitioned between EtOAc and brine. The organic phase was separated, dried

with MgSO$_4$, and concentrated *in vacuo* to give a crude solid. The crude material was triturated

with diethyl ether/hexane to provide the desired product 4-(((5-isopropyloxazol-2-

yl)methyl)thio)thiazol-2-amine as a red oil (368 mg, 57 %).

N-(5-(((5-isopropyloxazol-2-yl)methyl)thio)thiazol-2-yl)piperidine-4-carboxamide   *2,2,2-*

*trifluoroacetate (**B1**).* To a flask was added 1-(*tert*-butoxycarbonyl)piperidine-4-carboxylic acid

(99 mg, 1.1 equiv, 431 μmol) and DIPEA (205 μL, 3 equiv, 1.2 mmol) and HATU (194 mg, 1.3

equiv, 509 μmol) and the flask was stirred at room temperature for approximately 15 min. Next,

5-(((5-isopropyloxazol-2-yl)methyl)thio)thiazol-2-amine (100 mg, 1.0 equiv, 392 μmol) was

added and the reaction was stirred at room temperature for 16 h. The reaction mixture was

concentrated and purified by column chromatography (SiO$_2$, MeOH 0–10% in CH$_2$Cl$_2$) to yield

the desired product as a white solid to yield the intermediate *tert*-butyl 4-((5-(((5-

isopropyloxazol-2-yl)methyl)thio)thiazol-2-yl)carbamoyl)piperidine-1-carboxylate.   To   this

intermediate was added 20% TFA and CH$_2$Cl$_2$ (5 mL) and the reaction stirred for 1 h. The

reaction was concentrated *in vacuo* and after addition of MeOH a white precipitate crashed out.

The solid was filtered under vacuum to yield the desired product *N*-(5-(((5-isopropyloxazol-2-

yl)methyl)thio)thiazol-2-yl)piperidine-4-carboxamide 2,2,2-trifluoroacetate as a white solid (20

mg). The remaining crude material was purified by preparative HPLC (MeOH 10–100% in H$_2$O

(+0.05% TFA)) to yield the desired product *N*-(5-(((5-isopropyloxazol-2-yl)methyl)thio)thiazol-2-

yl)piperidine-4-carboxamide 2,2,2-trifluoroacetate (**B1**) as a white solid (32 mg). Yield over two

steps   *N-(5-(((5-isopropyloxazol-2-yl)methyl)thio)thiazol-2-yl)piperidine-4-carboxamide*   *2,2,2-*

trifluoroacetate (52 mg, 36 %). [1]H NMR (400 MHz, DMSO-*d$_6$*) δ 7.38 (s, 1H), 6.72 (d, *J* = 1.1 Hz,

1H), 4.03 (s, 2H), 3.21 (d, *J* = 12.8 Hz, 2H), 2.93 – 2.85 (m, 1H), 2.84 – 2.75 (m, 2H), 2.70 (s,

1H), 1.89 (d, *J* = 13.6 Hz, 2H), 1.77 – 1.63 (m, 2H), 1.12 (d, *J* = 6.9 Hz, 6H). [13]C NMR (214

MHz, DMSO-*d$_6$*) δ 172.45, 161.00, 158.76, 158.31, 145.12, 120.91, 119.04, 42.48, 38.65, 34.03,

25.29, 24.83, 20.48. HPLC purity: >95%. HRMS Calculated for [M+H]$^+$ C$_{16}$H$_{23}$N$_4$O$_2$S$_2$: 367.1184;

observed [M+H]$^+$: 367.1248.

1079

Figure S1. Characterization of N-(5-(((5-isopropyloxazol-2-yl)methyl)thio)thiazol-2-yl)piperidine-4-carboxamide 2,2,2-trifluoroacetate (**B1**) A) ^{1}H Spectra and chemical structure (inset). B) ^{13}C Spectra C) HPLC-UV Trace D) HRMS Trace.

1083

1084

1085

1086

1087

1088

1089

CAF-382 #509-518 RT: 2.20-2.24 AV: 10 NL: 5.45E8
T: FTMS + c ESI Full ms [150.0000-2000.0000]

1090

1091

1092

1093

1094

Scheme 2. Preparation of compounds **B4** and **B12**. a) Propylphosphonic anhydride (T3P),

DIPEA, THF, 0°C, 1 h, 93% yield; b) Palladium on carbon (Pd/C), MeOH, THF, 25°C, 16 h, 99%

yield; c) 4-methoxybenzoic acid, HATU, DIPEA, THF, 25°C, 17 h, 72% yield; d) HCl, dioxane,

25°C, 1h, 58% yield; e) 3-cyanobenzoic acid, T3P, DIPEA, THF, 25°C, 16 h; f) HCl, dioxane,

25°C, 1h, 39% yield over two steps.

General procedure for preparation **B4** and **B12**: To a flask was added 4-nitro-1*H*-pyrazole-3-
carboxylic acid (1.00 g, 1.0 equiv, 6.4 mmol), 1-Boc-4-aminopiperidine (1.28 g, 1.0 equiv, 6.4
mmol), and DIPEA (4.21 mL, 4.0 equiv, 25.5 mmol) in THF (20 mL). The reaction mixture was
cooled to 0°C, and then T3P (6.08 mL, 1.5 equiv, 9.5 mmol) was slowly added. After stirring for
1 h at 0°C, the reaction mixture was concentrated *in vacuo*, and then partitioned between EtOAc
and brine. The organic phase was separated, dried with $MgSO_4$, and concentrated *in vacuo* to
give a crude solid. The crude material was purified using an automated purification system
(SiO_2) 90%/10% hexanes/EtOAc to 100% EtOAc to give *tert*-butyl 4-(4-nitro-1*H*-pyrazole-3-
carboxamido)piperidine-1-carboxylate (**Int 2**) as an amorphous solid (2.00 g, 93% yield).

Palladium on carbon (5%, 31.3 mg, 0.1 equiv, 0.295 mmol) was added to *tert*-butyl 4-(4-
nitro-1*H*-pyrazole-3-carboxamido)piperidine-1-carboxylate (**Int 2**, 1.0 g, 1.0 equiv, 2.95 mmol) in
a mixture of methanol (10 mL) and THF (3.0 mL). The solution was placed under an
atmosphere of H_2 at room temperature. After 16h, the solution was filtered through SiO_2 and the
eluent was concentrated to afford *tert*-butyl 4-(4-amino-1*H*-pyrazole-3-carboxamido)piperidine-
1-carboxylate (**Int 3**) as an amorphous solid (900 mg, 99% yield).

To a flask was added *tert*-butyl 4-(4-amino-1*H*-pyrazole-3-carboxamido)piperidine-1-
carboxylate (**Int 3**, 1 equiv) and the corresponding carboxylic acid (1.2 equiv) and DIPEA (4.0
equiv) in THF (0.5 M) at room temperature. The reaction mixture was then treated slowly with
T3P (3.0 equiv). After stirring for 16–17 h, the reaction mixture was concentrated *in vacuo*, and
then partitioned between EtOAc and brine. The organic phase was separated, dried with
$MgSO_4$, and concentrated *in vacuo* to give a crude solid. The crude material was purified using
an automated purification system (SiO_2) 90%/100% hexanes/EtOAc to 100% EtOAc) to give an
intermediate, which was stirred in HCl/dioxane (3.3 mL) for 1 h at room temperature. Solvent
was removed and the crude material was purified using an automated purification system (SiO_2)
90%/100% hexanes/EtOAc to 100% EtOAc) to give the desired product (**B4** or **B12**) as an
amorphous solid (39–42% yield over 2 steps).

4-(4-methoxybenzamido)-N-(piperidin-4-yl)-1H-pyrazole-3-carboxamide (**B4**) [1]H NMR
(850 MHz, Methanol-d_4) δ 8.30 (s, 1H), 7.89 (d, *J* = 8.8 Hz, 2H), 7.07 (d, *J* = 8.8 Hz, 2H), 4.22
(tt, *J* = 10.8, 4.1 Hz, 1H), 3.89 (s, 3H), 3.48 (dt, *J* = 13.4, 3.9 Hz, 2H), 3.18 (td, *J* = 12.7, 3.1 Hz,
2H), 2.22 (dd, *J* = 14.3, 3.9 Hz, 2H), 1.96 – 1.85 (m, 2H). [13]C NMR (214 MHz, Methanol-d_4) δ
165.66, 165.58, 164.53, 134.00, 130.01, 126.71, 124.57, 121.57, 115.24, 56.06, 45.26, 44.25,
29.54. [1]H NMR purity: >95%. HRMS Calculated for [M + H]$^+$ $C_{17}H_{22}N_5O_3$: 344.1723; observed
[M+H]$^+$: 344.1717.

Figure S2. Characterization of 4-(4-methoxybenzamido)-N-(piperidin-4-yl)-1H-pyrazole-3-
carboxamide (**B4**) A) ^{1}H Spectra and chemical structure (inset). B*)* ^{13}C Spectra C) HRMS Trace.
1135

1136

1137

1138

1139

1140

4-(3-cyanobenzamido)-N-(piperidin-4-yl)-1H-pyrazole-3-carboxamide (**B12**) [1]H NMR

(850 MHz, Methanol-d_4) δ 8.33 (s, 1H), 8.26 (t, *J* = 1.7 Hz, 1H), 8.22 (dt, *J* = 7.9, 1.5 Hz, 1H),

7.97 (dt, *J* = 7.7, 1.4 Hz, 1H), 7.75 (t, *J* = 7.8 Hz, 1H), 4.22 (tt, *J* = 10.9, 4.1 Hz, 1H), 3.49 (dt, *J* =

14.1, 3.5 Hz, 2H), 3.18 (td, *J* = 12.9, 3.1 Hz, 2H), 2.22 (dd, *J* = 14.2, 3.7 Hz, 2H), 1.95 – 1.88 (m,

2H). [13]C NMR (214 MHz, Methanol-d_4) δ 165.57, 163.51, 136.50, 136.14, 134.35, 132.50,

131.84, 131.32, 124.01, 122.04, 118.88, 114.39, 45.33, 44.29, 29.53. [1]H NMR purity: >95%.

HRMS Calculated for [M + H]$^+$ C$_{17}$H$_{19}$N$_6$O$_2$: 339.1569; observed [M+H]$^+$: 339.1563.

1148

1149

1150　**Figure S3.** Characterization of 4-(3-cyanobenzamido)-N-(piperidin-4-yl)-1H-pyrazole-3-
1151　carboxamide (**B12**) A) ¹H Spectra and chemical structure (inset). B) ¹³C Spectra C) HRMS
1152　Trace.

1153

1154

1155

B.

1156

1157

1158

1159

1160

1161

1162

1163

1164   **Table S1.** NanoBRET data corresponding to compounds selected for initial study.

Compound	Parent	CDKL5 NB IC$_{50}$ (nM)	GSK3β NB IC$_{50}$ (nM)	fold
A01	AT-7519	363	>10,000	27.5
A02	AT-7519	178	4135	23.2
A03	AT-7519	19	404.8	21.3
A04	SNS-032	108	5688	52.7
A05	AT-7519	12	308.2	25.7
A06	SNS-032	221	7787	35.2
A07	AT-7519	6	192	32.0
A08	AT-7519	56	1374	24.5
A09	AT-7519	21	481	22.9
A10	SNS-032	424	>10,000	23.6
A11	AT-7519	9	3699	411.0
A12	AT-7519	20	524.8	26.2
B01	SNS-032	8	>10,000	1250.0
B02	SNS-032	190	9100	47.9
B03	SNS-032	10	>10,000	1000.0
B04	AT-7519	14	553.9	39.6
B05	SNS-032	388	8000	20.6
B06	AT-7519	155	8193	52.9
B07	AT-7519	36	3256	90.4
B08	AT-7519	161	>10,000	62.1
B09	AT-7519	37	3627	98.0
B10	AT-7519	90	4945	54.9
B11	AT-7519	141	3101	22.0
B12	AT-7519	69	3631	52.6
C01	SNS-032	323	>10,000	31.0

1165

1166

1167

1168 **Figure S4.** Curves corresponding with CDKL5 affinity measurements in Figure 2. (A) Curves
1169 generated in CDKL5 split-luciferase assay for B1, B4, and B12. (B) Curves generated in CDKL5
1170 NanoBRET assay for B1, B4, and B12.

1171

1172 **Figure S5.** Curves corresponding with GSK3 NanoBRET measurements in Figure 2. (A) Curves
1173 generated in GSK3α and GSK3β NanoBRET assays for B1. (B) Curves generated in GSK3α
1174 and GSK3β NanoBRET assays for B4. (C) Curves generated in GSK3α and GSK3β NanoBRET
1175 assays for B12.

1176

1177

1178

1179

1180   **Figure S6.** CDKL family selectivity evaluation via thermal shift assays.

1181

1182

1183

1184

1185 **Figure S7.** Phosphorylation of β-catenin, a substrate of GSK, is reduced upon treatment with a
1186 GSK inhibitor CHIR 99021 (Tocris) but EB2 phosphorylation is not changed. Western blot
1187 showing expression of total EB2, pS222 EB2, total β-catenin, phospho-β-catenin and tubulin in
1188 DIV14-15 rat primary neurones after an hour treatment with a GSK inhibitor.

1189

1190

1191

Figure S8. CDKL5, total EB2 and β-catenin expression in rat primary neurons after treatment with CAF-382, HW2-013 and LY-213 compounds. A, D, G, Quantification of CDKL5 expression in DIV14-15 rat primary neurones after an hour treatment with different concentrations of CAF-382, HW2-013 and LY-213 compounds respectively. Each concentration was compared to the control using a one-way ANOVA test. N = 3 biological replicates with 2 repetitions. Error bars are SD. B, E, H, Quantification of total EB2 expression in DIV14-15 rat primary neurones after an hour treatment with different concentrations of CAF-382, HW2-013 and LY-213 compounds respectively. Each concentration was compared to the control using a one-way ANOVA test. N = 3 biological replicates with 2 repetitions. Error bars are SD. C, F, I, Quantification of total β-

1201 catenin expression in DIV14-15 rat primary neurones after an hour treatment with different
1202 concentrations of CAF-382, HW2-013 and LY-213 compounds respectively. Each concentration
1203 was compared to the control using a one-way ANOVA test. N = 2 biological replicates with 2
1204 repetitions. Error bars are SD.

1205

1206

1207 **Figure S9.** SNS-032 (Selleck Chemicals) inhibits phosphorylation of EB2 in rat hippocampal
1208 slices in a time and concentration-dependent fashion. Effect of SNS-032 is compared to control
1209 slices with similar concentration of DMSO vehicle under three different conditions. Compare to
1210 Figure 6.

1211

1212

A	Time (h)	Concentration (ng/mL)
	0.5	508
	1	317
	2	79.1
	4	17.3

C	PK parameters	Unit	Mean
	T1/2	h	0.738
	Tmax	h	0.500
	Cmax	ng/mL	508
	AUC_{last}	h*ng/mL	628
	AUC_{Inf}	h*ng/mL	646
	$AUC_{\%Extrap}$_obs	%	2.85
	MRT_{Inf_obs}	h	1.18
	AUC_{last}/D	h*mg/mL	274
	F	%	NA

B — Mean plasma concentration vs time profile for B1 after 2.29 mg/kg in CD1 mice

1213
1214 **Figure S10.** Snapshot pharmacokinetic (PK) results for B1. (A) Plasma concentration
1215 measurements at timepoints post-administration. (B) Plot of mean plasma concentration over
1216 the time course of the PK study. (C) Summary of B1 PK properties (n = 2).

1217

1218

A	IP dose: 2.29 mg/kg		
Time (h)	Brain Conc (ng/g)	Plasma (ng/mL)	Ratio (Brain/plasma)
1	15.6	340	0.0459

B	IP dose: 7.63 mg/kg		
Time (h)	Brain Conc (ng/g)	Plasma (ng/mL)	Ratio (Brain/plasma)
1	158	750	0.210*

1219

1220

1221 **Table S2.** Brain exposure results for B1. (A) Brain/plasma concentration-time data after 2.29

1222 mg/kg dose. (B) Brain/plasma concentration-time data after 7.63 mg/kg dose. *One animal

1223 artificially drove up blood/plasma ratio. N = 3 per dose.

1224

1225

Figure 1. Screening of CDKL5 inhibitors in rat primary neurons using Western blotting. Western blots showing expression of total EB2 and levels of Ser222 EB2 phosphorylation in DIV14-16 rat primary neurons upon treatment of 1 hour with 5 nM, 50 nM, 500 nM and 5000 nM of selected CDKL5 inhibitors.

Figure 2. Structures of CDKL5 inhibitor leads and corresponding parent compounds.

A

B1

$S_{10}(1 \mu M) = 0.017$

Percent Control
- 0%
- 0.1%
- 0.1-1%
- 1-5%
- 5-10%
- 10-35%
- > 35%

Kinase	DiscoverX PoC value	Assay format	IC$_{50}$ value (nM)
GSK3A	0.7	Enzymatic and NB	470 and 1900
CDKL5	0.7	Binding and NB	6.7 and 11
PCTK2	1.3	Enzymatic	89
CDK7	1.9	Enzymatic	300
PCTK1	5.5	Enzymatic	62
CDC2L5	5.9	Enzymatic	>10000
CDK9	8.3	Enzymatic	20
YSK4	14	Enzymatic	>10000
PCTK3	17	Enzymatic	100
CDK4	20	Enzymatic	1900
GSK3B	51	Enzymatic and NB	2200 and 4500

B

B4

$S_{10}(1 \mu M) = 0.02$

Percent Control
- 0%
- 0.1%
- 0.1-1%
- 1-5%
- 5-10%
- 10-35%
- > 35%

Kinase	DiscoverX PoC value	Assay format	IC$_{50}$ value (nM)
GSK3A	0	Enzymatic and NB	11 and 120
CDKL5	0.2	Binding and NB	6.4 and 17
GSK3B	2.1	Enzymatic and NB	38 and 250
DYRK2	4.4	Enzymatic	240
CDK7	4.4	Enzymatic	>10000
PCTK1	6.3	Enzymatic	1200
DYRK1A	9.1	Enzymatic	900
PCTK2	9.5	Enzymatic	1600

C

B12

$S_{10}(1 \mu M) = 0.01$

Percent Control
- 0%
- 0.1%
- 0.1-1%
- 1-5%
- 5-10%
- 10-35%
- > 35%

Kinase	DiscoverX PoC value	Assay format	IC$_{50}$ value (nM)
GSK3A	0	Enzymatic and NB	16 and 190
CDKL5	0.6	Binding and NB	8.7 and 23
GSK3B	1.7	Enzymatic and NB	48 and 1200
DYRK2	4.6	Enzymatic	920
CDK7	17	Enzymatic	>10000
PCTK1	19	Enzymatic	1700
RPS6KA4	27	Enzymatic	>10000
DYRK1A	29	Enzymatic	4800
DYRK1B	34	Enzymatic	1700

Figure 3. Kinome-wide selectivity data for CDKL5 lead compounds. Kinome tree diagrams illustrate the selectivity of these compounds when profiled against 403 wild-type (WT) human kinases at 1 µM at Eurofins DiscoverX in their *scan*MAX panel. The percent control legend shows red circles of different sizes corresponding with percent control value each kinase binds the small molecule in this large binding panel. Also included are selectivity scores ($S_{10}(1 \mu M)$), which were calculated using the PoC values for WT human kinases in the *scan*MAX panel only. The S_{10} score is a way to express selectivity that corresponds with the percent of the kinases screened that bind with a PoC value <10. In the embedded tables kinases are listed by their gene names and ranked by their percent of control (PoC) value generated in the *scan*MAX panel. Rows colored green demonstrate enzymatic IC$_{50}$ values within a 30-fold window of the CDKL5 binding IC$_{50}$ value. Correlating assay format used to generate data in the last column of nested table is listed in the preceding column, with NB = NanoBRET.

Figure 4. B1 potently inhibits the kinase activity of CDKL5. (A) Representative graph from a CDKL5 kinase assay demonstrating that purified WT human CDKL5 retains kinase activity, while the kinase dead (KD, CDKL5 K42R) human protein is functionally inactive (n = 3). (B) Representative western blot illustrating that equal expression of WT and KD proteins was observed in the kinase assay experiments. (C and D) Kinase assays using purified WT human CDKL5 in the presence of 10 or 100 nM of the indicated compounds (n = 3). One-way ANOVA with Dunnett's multiple comparison test used. *** = p <0.0001 and non-significant comparisons not shown.

Figure 5. CAF-382 (**B1**), HW2-013 (**B4**), and LY-213 (**B12**) compounds reduce CDKL5 activity in rat primary neurons. HW2-013 (**B4**) and LY-213 (**B12**) also downregulate GSK3 activity. (A) Western blot and quantification showing expression of EB2 phosphorylation and β-catenin phosphorylation in DIV14-15 rat primary neurones after an hour treatment with different concentrations of CAF-382 (**B1**). (B) Western blot and quantification showing expression of EB2 phosphorylation and β-catenin phosphorylation in DIV14-15 rat primary neurons after an hour treatment with different concentrations of HW2-013 (**B4**). (C) Western blot and quantification showing expression of EB2 phosphorylation and β-catenin phosphorylation in DIV14-15 rat primary neurons after an hour treatment with different concentrations of LY-213 (**B12**). Each concentration was compared to the control using a Kruskal-Wallis test. n = 3 biological replicates with 2 repetitions. *p≤0.05. **p≤0.01. ***p≤0.001. Error bars are SD.

Figure 6. CAF-382 (**B1**) reduced phosphorylation of EB2 in CA1 hippocampal slices. (A) Example blots demonstrate no alterations in CDKL5 expression across treatments (stats in text) and suggest relative CAF-382 (**B1**) dose-dependent differences in EB2 phosphorylation. (B) Normalized quantification of EB2-phosphorylation (density of EB2-phosphorylation bands/ density of EB2-total bands). CAF 382 (**B1**) (control: 1±0.10 versus 100 nM: 0.49±0.02, n = 8, p = 0.002, RM-ANOVA) (control: 1±0.08 versus 45 nM: 0.16±0.03, n = 10, p < 0.001, RM-ANOVA) (control: 1± 0.14 versus 10 nM: 0.62± 0.11, n=7, p = 0.003, RM-ANOVA) reduced EB2 phosphorylation at all concentrations.

Figure 7. CAF-382 (**B1**) reduced post-synaptic fEPSP responsiveness, as measured by input-output curves. (A) Sample input (fiber volley) / output (fEPSP slope) (I/O) curves at baseline, 30 minutes and 60 minutes after CAF-382 (**B1**) (100nM) in a hippocampal slice. Slopes of I/O curves were obtained at baseline, and 30 minutes and 60 minutes after treatment with CAF-382 (**B1**). Asterisks indicate sample traces (B) from treated slice; stimulus artifact (arrow) has been removed for clarity. Initial negative deflection after the artifact is the fiber volley, followed by the fEPSP. (C) RM-ANOVA of I/O slopes of CAF-382 (**B1**) (100 nM and 45 nM) were significantly decreased after 30 minutes (100 nM: 1.91±0.23, n = 9, p = 0.006; 45 nM: 1.58±0.24, n=10, p = 0.013) and 60 minutes (100 nM: 2.00±0.30, n = 9, p = 0.001; 45 nM: 1.57±0.22, n=10, p = 0.009) compared to baseline (100 nM: 3.24±0.52, n = 9; 45 nM: 2.29±0.23, n=10). Baseline I/O slope was used to normalize each recording to allow comparisons across all treatments. CAF-382 (**B1**) (10 nM) did not alter I/O slopes after 30 or 60 minutes (RM-ANOVA, n = 9, p = 0.07). Error bars are SE.

Figure 8. CAF-382 (**B1**) did not alter presynaptic release as measured by paired-pulse ratio. (A) Paired-pulse ratios (PPR) of fEPSPs (Slope of fEPSP2/Slope of fEPSP1), reflective of pre-synaptic release, were unaltered across a range of stimulus intervals (RM-ANOVA, n=7) by CAF-382 (**B1**) (100nM) for 60 minutes compared to baseline. (B) Sample trace for 50 ms interval. Reduction of initial fEPSP slopes were consistent with Figure 7. Stimulus artifacts (arrows) have been removed for clarity. Error bars are SE.

Figure 9. CAF-382 (**B1**) did not alter NMDAR-mediated component of fEPSPs. Maximal fEPSPs recorded in stratum pyramidale of CA1 with reduced extracellular Mg^{2+} demonstrate multiple population spikes; previous studies have demonstrated that secondary population spikes in these conditions are NMDAR-dependent[64]. CAF-382 (**B1**) (100 nM) did not alter population spike 1 (control -4.057±0.584mV, CAF-382 (**B1**) 3.929±0.736mV, n=10, p=0.742, RM-ANOVA) or NMDAR-mediated population spike 2 (control -2.093±0.568mV, CAF-382 (**B1**) -1.656±0.4442, n=10, p=0.867, RM-ANOVA). For comparison, the selective NMDAR antagonist D-APV (50 μM) completely blocked population spike 2, as previously reported [64]. Stimulation artifact has been removed (arrow) for clarity.

Figure 10. CAF-382 (**B1**) acutely reduces the expression of LTP in CA1 hippocampus. Hippocampal slices were initially incubated in CAF-382 (**B1**) for at least 30 minutes and then continuously perfused with CAF-382 (**B1**) (100 nM). fEPSP slope was normalized to initial baseline, expressed as 100% (dotted line); mean±SEM shown. Following stable baseline of fEPSP slope (> 20 minutes), LTP was induced by two theta-burst (TBS) trains. Compared to interleaved control slices, CAF-382 (**B1**) significantly reduced LTP as measured by fEPSP slope at 56-60 minutes post theta-burst (control: 115.6±2.2 n = 13 slices, 12 rats; CAF-382 (**B1**): 106.4±3.5, n = 5 slices, 4 rats; p = 0.029, 2-way ANOVA, Holm-Sidak post-hoc). Sample traces for control (black filled circle) and CAF-382 (**B1**) (red filled circle) before (1, solid trace) and after LTP (2, dotted trace) are shown to the right. Error bars are SE.

Figure 11. CAF-382 (**B1**) acutely reduces the expression of LTP mediated by AMPA-type glutamate receptors in CA1 hippocampus. Hippocampal slices were initially incubated in CAF-382 (**B1**) (100 nM) for at least 60 minutes prior to recording. Peak negative current of AMPA-type glutamate receptor mediated synaptic responses were normalized to initial baseline post break-in (time = 0), expressed as 100% (dotted line); mean±SEM shown. Following baseline, LTP was induced by a pairing protocol. Compared to interleaved control slices, CAF-382 (**B1**) significantly reduced LTP at 27-31 minutes post break-in (control: 197.5±8.4 n = 14 slices, 14 rats; CAF-382 (**B1**): 123.14±9.0, n = 12-13 slices, 13 rats; p < 0.001, 2-way ANOVA, Holm-Sidak post-hoc). Sample traces for control (black filled circle) and CAF-382 (**B1**, red filled circle) before (1, solid trace) and after LTP (2, dotted trace) are shown to the right. Error bars are SE.

Figure S7. Phosphorylation of β-catenin, a substrate of GSK, is reduced upon treatment with a GSK inhibitor CHIR 99021 (Tocris) but EB2 phosphorylation is not changed. Western blot showing expression of total EB2, pS222 EB2, total β-catenin, phospho-β-catenin and tubulin in DIV14-15 rat primary neurones after an hour treatment with a GSK inhibitor.