A guide to the *in vitro* reconstitution of Cdc42 GTPase activity and its regulation

Sophie Tschirpke[1], **Frank van Opstal**[1], **Ramon van der Valk**[1], **Werner K-G. Daalman**[1], **Liedewij Laan**[1,*]

[1]Bionanoscience Department, Delft University of Technology, Delft, the Netherlands; [*]For correspondence: L.Laan@tudelft.nl

Abstract Cdc42 is a small Rho-type GTPase and the main regulator of cell division in eukaryotes. It is surrounded by a large network of regulatory proteins. To understand the processes around cell division, in-depth understanding of Cdc42 and its regulation is required. *In vitro* reconstitutions are a suitable tool for such detailed mechanistic studies, as they allow a high level of control over the conditions and components used and. For these Cdc42 and its regulators need to be expressed, purified, and tested for their activity. There are many methods described for this, but their details, possible difficulties, and points of failure are rarely discussed. This makes *in vitro* studies on Cdc42 less accessible to scientists that have a background different from biochemistry. We here present our experience with working with Cdc42 *in vitro*. We describe the recombinant expression and purification behaviour of 12 Cdc42, six Cdc42-mNeonGreen^{SW} and four Cdc42-sfGFP^{SW} constructs in *E. coli*. We explore Cdc42 dimerisation *in vitro* and assess its activity using GTPase Glo assays and Flag-pulldown assays. GTPase Glo assays turn out to be a reliable tool to quantitatively asses GTPase activities, wheareas pulldown experiments are more error prone. We find that most Cdc42 constructs, with the exception of those with an N-terminal Twin-Step-tag, show a similar GTPase activity and interaction with the GDP/GTP exchange factor Cdc24. We close with using enterokinase and TEV protease to generate untagged Cdc42. Enterokinase also cuts Cdc42 in an undesired position. TEV protease leads to the desired product, which retains its GTPase activity but shows a reduced Cdc24 interaction. The work presented here acts as a guide for scientists desiring to work with Cdc42 *in vitro* through describing Cdc42's properties in detail and examining assays that can be used to study its behaviour or act as activity checks.

Graphical abstract:

Introduction

Cellular processes arise through complex interactions of many components. In order to enhance our understanding of these interactions, research on the cellular components (e.g. DNA, RNA, proteins, lipids) and their interplay is required. *In vivo* investigations give insights into the overall cellular function of a component, and *in vitro* assays elucidate the specifics of single interactions and component functions. Bridging the gap between highly controlled single to a few component *in vitro* assays

and highly complex *in vivo* processes are *in vitro* reconstitutions (for example [1, 2, 3, 4]). Here the interplay between several components, synergies, and emergent behaviours can be characterised. Factors such as concentrations, ratios, and temperature can be manipulated and their effect on the overall system can be observed - giving rise to an an in-depth understanding of the system and a base towards building a synthetic cell. With increasing amount of added components the complexity of the system increases - it moves towards a cell-like state - but it also becomes experimentally more challenging [5]; for example because an altered property of a single component can change the overall behaviour of the entire system. Thus, a high level of control and knowledge over the components is needed.

The system surrounding the Rho-type GTPase Cdc42 is an interesting case for reconstitutions. Cdc42 is the main regulator of polarity establishment and cell division in eukaryotes [6]. It is part of a complex network of polarity proteins, including GDP/GTP exchange factors (GEFs), GTPase activating proteins (GAPs), guanine nucleotide dissociation inhibitors (GDIs), scaffold proteins, and other regulatory proteins, that all interact with each other and with Cdc42 in particular [7, 8, 9]. In order to understand the regulation of cell division detailed understanding Cdc42 and of its complex protein network is needed, making it a suitable target for *in vitro* reconstitutions. For these Cdc42 constructs (with purification tags) need to be designed, expressed, purified, stored until use, and tested for their activity. One of the first decisions that thus needs to be made is that of native or recombinant protein expression/purification. In native purification the protein of interest (POI) is expressed in its native organism, in the recombinant approach a different organism is used. Native purification gives the most biologically relevant protein, but expression and purification protocols are usually less established and more difficult to follow, and the obtained POI batch can contain small amounts of co-purified POI binding partners and consists of a mixture of POI with various forms of post-translational modifications (for example, for *Saccharomyces cerevisiae* Cdc42: ubiquitinylation [10, 11], phophorylation [12], farnesylation and geranyl-geranylation [13]. The latter, however, can also be an advantage of a native purification: if the post-translational modification is an essential feature of the POI, its presence is required. Previous *in vitro* studies on Cdc42 obtained the protein using (1) *Escherichia coli* based expression systems [14], (2) insect cell expression systems [15, 16, 17, 18, 19, 20, 21], or (3) through purification of membrane-bound Cdc42 from yeast [22, 23]. We chose to purify *S. cerevisiae* Cdc42 recombinantly from *Escherichia coli* due to the method's advantages: (1) Recombinant purification from *E. coli* is widely used and protocols are established and easy to follow. *E. coli* grows fast and reliably and protein can be over-expressed in a high yield. (2) As Cdc42 is not native in *E. coli*, it will have little to no binding-partners that it can co-purify. One can thus be more certain that the obtained Cdc42 batch is only Cdc42. This is especially important if Cdc42 ought to be used to quantitatively study its interaction with its binding partners. If the Cdc42 batch contains already small amounts of binding partners, claims about the effect of a binding partner on Cdc42 and the concentration-dependence of this effect are less accurate. We like to point out that Cdc42 purified in this way does not possess its C-terminal prenyl group, which is responsible for anchoring Cdc42 to the membrane and binding to the GDI Rdi1 [24, 25]. This group would need to be added later in an *in vitro* reaction [14, 26].

There are many frequently used methods for protein expression, purification, and assays that assess protein activity and their interactions, but their details, possible difficulties, and points of failure are rarely discussed. This makes *in vitro* studies, and reconstitutions in particular, with Cdc42 less accessible to scientist that have a background different from biochemistry. Here we set out to make the *in vitro* use of (*Saccharomyces cerevisiae*) Cdc42 more accessible to a broad audience, through describing our work and experience with various Cdc42 constructs. The goal of this paper is not to map out all properties of all constructs used, but to showcase what steps and difficulties working with Cdc42 *in vitro* entails. We include positive as well as negative results and point out heterogeneous behaviours of Cdc42. We start by describing our Cdc42 construct design in detail. We then show the recombinant expression and purification behaviour of Cdc42 and Cdc42-sfGFP and -mNeongreen sandwich fusions in *E. coli* and discuss Cdc42 dimerisation *in vitro*. We then used GTPase and pulldown assays to (1) examine our Cdc42 constructs on their functionality - their GTPase activity and interaction with two binding partners, and (2) to assess how reliable and straightforward these assays are to test protein properties. We close with exploring how suitable two cleavage enzymes, enterokinase and TEV protease, are for cleaving tags off Cdc42.

Results

Cdc42 constructs

Purification of proteins for *in vitro* studies is generally done by attaching an N- or C-terminal purification tag to the POI. The rational behind the placement of purification tags and their effect on the protein's expression/purification behaviour and properties are rarely discussed. This contributes to making *in vitro* studies less accessible for non-biochemists. Further, tag cleavage is not always easy or possible. To cleave off a tag a recognition site for a cleavage enzyme needs to be placed in-between the POI and

purification tag. Cleavage enzymes have a relatively high sequence specificity, but can also cleave within the POI causing degradation products. The enzyme's cleavage behaviour and efficiency need to be tested for every POI to ensure proper matching. After the cleavage reaction the tagged, untagged, degraded POI species and cleavage enzyme need to be separated, adding another undesirable and yield-reducing purification step. In other cases the cleavage of the purification tag is not possible. For example when the tag is used to bind the POI to beads or a modified surface to study protein-protein interactions.

Cdc42 is a small Rho-type GTPase of 191 amino acids (AAs)/21 kDa, consisting mainly of a Rho-domain, that is responsible for its catalytic activity, and a short flexible N-terminal region made of the polybasic region (PBR) directly followed by the CAAX box. The PBR is a five AA long unstructured region of mostly positively charged amino acids that has been linked to dimerisation and supports binding to negatively charged membranes [27]. The CAAX-box is a four AA sequence at the C-terminus, at which the protein gets post-translationally modified: a prenyl group (geranylgeranyl or farnesyl) gets appended, allowing the protein to bind to membranes [24].

So-far, *in vitro* studies with 6His-tagged [17, 28, 19, 21, 14, 29], GST-tagged [16, 30, 19, 22], and 6His-Strep-II-tagged [23] Cdc42 were conducted. Of these references, only one reported to have cleaved off the tag [17]. All of these constructs have the purification tag placed on the N-terminus. It is likely the case because *in vivo* Cdc42 is prenylated at its C-terminus. Even though not all mentioned studies used prenylated Cdc42, all left the C-terminus unmodified. We set out to explore the effect of N- and C-terminal additions, including various purification tags, on Cdc42. We constructed three types of constructs:

- Type H: Here only a single 6His-tag is directly appended to the N-terminus of Cdc42 (Fig. 1). This construct has the least amount of additions and therefore is expected to be least influenced by them.
- Type S: Here only N-terminal additions are added; a purification tag followed by a thrombin site, T7 tag, and enterokinase site (Fig. 1). Thrombin and enterokinase sites allow the cleavage of the N-terminal region, thus facilitating for the removal of the purification (and T7) tag, giving the possibility to generate an untagged protein. The T7 tag is an 11-residue peptide from the leader sequence of the T7 bacteriophage gene 10 [31] and aids protein expression in *E. coli*.
- Type D: Here both the N-terminal additions from type S plus C-terminal additions are added. On the C-terminus a sortase site (sortase A from *Staphylococcus aureus*) followed by a second purification tag is added (Fig. 1). The sortase site allows the removal of the C-terminal tag and the ligation of a peptide probe to the protein in a single reaction step [26]. With this, a protein prenylation moity or other modification could be added. The C-terminal purification tag can be used for an additional affinity chromatography step during protein purification, and to separate labelled from unlabelled protein after a sortase-mediated labelling reaction. To asses C-terminal Cdc42 additions further, we designed a Cdc42 construct with an alternative membrane-binding domain (Cdc42-BC). Here the first two basic cluster (BC) regions of the yeast protein Bem1 (51 AA) got inserted into the protein's C-terminus in between its PBR and CAAX box. The BCs are a 23 to 74 AAs long unstructured region of mostly positively charged AAs that were shown to anchor Bem1 to negatively charged membranes *in vitro* [32]. We designed this construct to further asses how unstructured (linker) regions on Cdc42's C-terminus affect its behaviour. We did not add structured regions, as the addition of a C-terminal amphipathic helix from Rit to Cdc42 resulted in folding issues of the protein *in vitro* (unpublished data by P. Schwille group (MPI Martinsried, Germany)). (The addition was in reference to Bendezu *et al.*, which showed that this alternative membrane binding domain leads to viable cells *in vivo* [33].).

Figure 1. Schematic overview of the protein constructs. (A) Illustration of the general size and outline of double-tagged (type D) and single-tagged constructs (type S) and of constructs to which only a single 6His-tag got added (type H). (B) Zoom-in of the N- and C-terminal tag regions from type S and type D and illustration of the size difference of the purification tags. All constructs are drawn roughly up to scale, in terms of their number of amino acids (AA). The position of domains of interest are indicated. An overview of the specific constructs is given in Tab. 2.

Table 1. Purification tag specifications. For more information regarding the Strep-II®-tag and Twin-Strep®-tag see [34], and regarding the Flag®-tag see [35].

Tag	Amino acid sequence	Length	Size
H: 6His	HHHHHH	6 AA	0.8 kDa
F: Flag®	DYKDHDGDYKDHDIDYKDDDDK	22 AA	2.7 kDa
S: Strep-II®	WSHPQFEK	8 AA	1.1 kDa
SS: Twin-Strep®	WSHPQFEKGGGSGGGSGGGSWSHPQFEK	28 AA	2.9 kDa
H-SS: 6His + Twin-Strep®	HHHHHHWSHPQFEKGGGSGGGSGGGSWSHPQFEK	34 AA	3.7 kDa

As a standard we added a 6His-tag (H), placed N- and C-terminally, to all our constructs. The 6His-tag it is a small often-used tag that facilitates His-affinity chromatography (His-AC) - an easy, fast, and cheap purification method. Type D constructs contained additionally a Flag- (F), Strep-II- (S), or Twin-Strep- (SS) tag. All of these are also rather small in size and can be used for affinity chromatography. The Strep-II- and Twin-Strep-tag differ only in so far from each other as the Twin-Strep-tag is made from two repeats of the Strep-II-tag that are spaced with a linker (Tab. 1). The Twin-Strep-tag binds by an order of magnitude tighter to Strep-Tactin and is therefore more effective when used for purification purposes [34]. Sequences and sizes of these purification tags are given in Tab. 1.

Cdc42 expression in *E. coli* does not result in prenylated Cdc42, as *E. coli* lacks the machinery responsible for it. If needed, such a group can still be added in an *in vitro* reaction: Cdc42 constructs type H and S contain a C-terminal CAAX box, which can be prenylated using purified farnesyltransferase or geranylgeranyl-transferase type I [14]. Type D constructs contain a sortase site with which a prenyl group can be ligated to the protein's C-terminus [26].

To expand out investigations on Cdc42 further, we designed fluorescent Cdc42 fusions. In principle, a fluorophore could be attached to either Cdc42's N- or C-terminus. Cdc42 fusions with N-terminal fluorophores have been controversial: N-terminal fusions of Cdc42 with fluorescent proteins have been shown to lead to disfunctional proteins *in vivo* [33], but two studies used such constructs successfully *in vitro* [36, 29]. C-terminal additions on Cdc42 are not well-studied. We are only aware of the unsuccessful attempt to add a C-terminal amphipathic helix. However, it has been shown that fast-folding fluorophores can be inserted into a solvent-exposed loop of Cdc42 [33]. We used this approach to create sandwich fusions of Cdc42 and sfGFP or mNeonGreen (Fig. 1), two proteins that are known to fold quickly [37, 38].

An overview of the herein used constructs and their abbreviations is given in Tab. 2.

Table 2. List of protein constructs and specification of their attributes. Protein sequences are stated in Supplement S5. If not stated otherwise, Cdc42 constructs contain the proteins natural CAAX sequence (CAIL). All 'pRV' plasmids were created as part of this work. [1]: pDM272 was received from D. McCusker (University of Bordeaux) and was published in [23]. Abbreviations: EK: enterokinase.

Abbreviation	Type	N-term. tag	C-term. tag	Other modification	Plasmid	Protein size
Cdc42						
H-Cdc42	H	6His		CAAX sequence = CAIA	pRV035	22 kDa
H-Cdc42:CAIA	S	6His		CAAX sequence = CAIA	pRV050	26 kDa
H-Cdc42:CTIS	S	6His		CAAX sequence = CTIS	pRV086	26 kDa
H*-Cdc42:CTIS	S	6His		TEV instead of EK site, CAAX sequ. = CTIS	pRV104	26 kDa
H-Cdc42-F	D	6His	Flag		pRV007	29 kDa
F-Cdc42-H	D	Flag	6His		pRV073	29 kDa
H-Cdc42-S	D	6His	Strep-II		pRV024	28 kDa
H-Cdc42-SS	D	6His	Twin-Strep		pRV068	29 kDa
S-Cdc42-H	D	Strep-II	6His		pRV030	28 kDa
SS-Cdc42-H	D	Twin-Strep	6His		pRV058	29 kDa
H-SS-Cdc42-F	D	6His + Twin-Strep	Flag		pRV087	32 kDa
S-Cdc42-BC-H	D	Strep-II	6His	+ BC from Bem1	pRV051	33 kDa
Fluorescent Cdc42						
H-Cdc42-mNeon-F	D	6His	Flag	+ mNeonGreen^{SW}	pRV010	55 kDa
F-Cdc42-mNeon-H	D	Flag	6His	+ mNeonGreen^{SW}	pRV074	55 kDa
H-Cdc42-mNeon-S	D	6His	Strep-II	+ mNeonGreen^{SW}	pRV052	54 kDa
H-Cdc42-mNeon-SS	D	6His	Twin-Strep	+ mNeonGreen^{SW}	pRV067	56 kDa
S-Cdc42-mNeon-H	D	Strep-II	6His	+ mNeonGreen^{SW}	pRV057	54 kDa
S-Cdc42-mNeon-BC-H	D	Strep-II	6His	+ BC from Bem1 + mNeonGreen^{SW}	pRV093	61 kDa
H-SS-Cdc42-mNeon-F	D	6His + Twin-Strep	Flag	+ mNeonGreen^{SW}	pRV090	58 kDa
F-Cdc42-sfGFP-H	D	Flag	6His	+ sfGFP^{SW}	pRV101	57 kDa
H-Cdc42-sfGFP-SS	D	6His	Twin-Strep	+ sfGFP^{SW}	pRV100	56 kDa
S-Cdc42-sfGFP-H	D	Strep-II	6His	+ sfGFP^{SW}	pRV099	55 kDa
H-SS-Cdc42-sfGFP-F	D	6His + Twin-Strep	Flag	+ sfGFP^{SW}	pRV102	59 kDa
Cdc24 and Bem1						
Cdc24-H	H		6His		pDM272 [1]	98 kDa
H-Cdc24-SS	D	6His	Twin-Strep		pRV065	103 kDa
S-Cdc24-H	D	Strep-II	6His		pRV031	105 kDa
SS-Cdc24-H	D	Twin-Strep	6His		pRV060	105 kDa
H-SS-Cdc24-F	D	6His + Twin-Strep	Flag		pRV088	108 kDa
H-Bem1-F	D	6His	Flag		pRV009	70 kDa
S-Bem1-H	D	Strep-II	6His		pRV032	68 kDa
SS-Bem1-H	D	Twin-Strep	6His		pRV061	70 kDa
H-SS-Bem1-F	D	6His + Twin-Strep	Flag		pRV089	73 kDa

Figure 2. Cdc42 constructs type S and D, but not type H, express in a high amount and show dimer bands on SDS-Page. (A) Cdc42 constructs type S and D, but not type H, express in all three conditions ('f': 3 h 37°C 1.0 mM IPTG, 's': 18 h 18°C 0.2 mM IPTG, 'AI': 3 h 37°C + 18 h 18°C [39], Expression levels of Cdc42 were assessed by anti-His Western blotting. (B) Cdc42 expresses well, independent of which N- or C-terminal tags were added. Cdc24 and Bem1 with an N-terminal Twin-Strep-tag do not express. Expression levels were assessed by anti-His and anti-Cdc42 Western blotting. Cdc42 constructs were expressed in condition 'f' and Cdc24 and Bem1 constructs were expressed in condition 's'. (C) Cdc42 shows dimer bands on SDS-Page, but dimers can not be detected in SEC-MALS. Molecular weight of purified Cdc42 constructs determined by SEC-MALS in comparison to the expected mass based on the amino acid sequence ('Theory'). Anti-His and anti-Cdc42 Western blots of the peak fractions from SEC-MALS runs.

Cdc42 expresses robustly and can be purified in a high yield using His-AC

First, we examined expression levels of Cdc42. We assed Cdc42 constructs which only are 6His-tagged (type H), have a larger additional N-terminal region (type S), and contain both N- and C-terminal additions (type D) (Fig. 1). Cdc42 was placed under an IPTG inducible promotor so that it's expression can be induced through addition of that chemical. We tested three expression conditions:

'f': a strong and fast expression at elevated temperatures, induced by a high amount of IPTG (3 h at 37°C with 1 mM IPTG).

's': a low and slow expression at lower temperatures, induced by a smaller amount of IPTG (18 h at 18°C with 0.2 mM IPTG).

'AI': a self-inducing combined approach, called auto-induction (3 h at 37°C + 18 h at 18°C) [39].

All Cdc42 constructs contained a 6His-tag, therefore expression levels could be analysed by anti-His Western blotting (Fig. 2a-c). Both type S and D expressed at all conditions in roughly equal amounts (Fig. 2a). The presence of a C-terminal tag or variations in Cdc42's CAAX box did not alter the expression levels. Further, the position of the tags in type D did not affect Cdc42 expression (Fig. 2b). Type H did not express in any condition (Fig. 2a). The N-terminal region of constructs of type S and D, but not of type H, contains a thrombin cut site, T7 tag, and enterokinase cut site. If the direct placement of the 6His-tag onto Cdc42's N-terminus interferes with protein folding, the additional elements could act as a spacer, circumventing the issue. The T7 tag could also directly be responsible for facilitating Cdc42 expression, as it aids protein expression in *E. coli*.

We then purified Cdc42 type S and D constructs. All constructs could be purified in a high yield in a single step His-AC (3-7 mg per 1 L expression volume) (Fig. 3). After purification only one construct, SS-Cdc42-H, was highly unstable and precipitated even at low concentrations (0.5 mg/mL). This was only the case for SS-Cdc42-H, all other constructs (including S-Cdc42-H, H-Cdc42-SS, and H-SS-Cdc42-F) showed no precipitation and were stable up to at least 2 mg/mL. This suggests that the N-terminal Twin-Strep-tag destabilises Cdc42. We observed a similar trend for the expression of two other yeast proteins; Cdc24 and Bem1. Cdc24 and Bem1 constructs, that were double-tagged in the same fashion as described for Cdc42, expressed when tagged with an N-terminal Strep-II-tag, C-terminal Twin-Strep-tag, or N-terminal 6His-Twin-Strep-tag, but did not, or only to greatly reduced level, express when tagged with an N-terminal Twin-Strep-tag (Fig. 2b). We suspect that the negative effect of the N-terminal Twin-Strep-tag is thus not Cdc42-specific, but inherent to the properties it exhibits when placed at a protein's N-terminus.

We further observed that Flag-tagged protein constructs of type D run slightly higher than their expected size on SDS-Page (Fig. 2b, 3). We can imagine that this is an artefact of how the Flag-tag moves through an SDS-Page gel: the Flag-tag is filled with

Figure 3. Cdc42 can be purified in a high yield in a single His-affinity chromatography step. (A) Purification profile of H-SS-Cdc42-F, purified by His-AC. The protein was eluted in a gradient of His-AC washing buffer (50 mM Tris-HCl (pH=8.0), 1 M NaCl, 5 mM imidazole, 1 mM 2-mercaptoethanol) and His-AC elution buffer (50 mM Tris-HCl (pH=8.0), 100 mM NaCl, 500 mM imidazole, 1mM 2-mercaptoethanol). The purification profiles of Cdc42 constructs shown in (B) looked alike. (B) SDS-Page of purified Cdc42 after a single step His-AC. The band of the full-length protein is indicated by an orange arrow. The asterix indicates the height of Cdc42 dimers.

charged residues and contains very few hydrophobic residues. The lack of hydrophobic residues (and the presence of charged residues) makes it difficult for SDS to bind to the tag. This could allow the Flag-tag to fold back into its native structure. A folded structure will migrate differently through the gel, and in the case of the Flag-tag, slower that one would expect for a peptide of a similar size.

Taken together, Cdc42 expresses highly in a broad range of conditions and it can easily and in a high yield be purified in a single-step His-AC. The majority of constructs with N- and N- plus C-terminal additions (type S and D) behave equally well. The only exceptions were H-Cdc42 and SS-Cdc42-H. H-Cdc42 did not express at all, suggesting that a linker region between 6His-tag and Cdc42, or the T7 lead specifically, is required. The N-terminal Twin-Strep-tag caused Cdc42 precipitation and expression issues in Cdc24 and Bem1, indicating that the observed issues are tag-related.

Cdc42 shows dimer bands on SDS-Page

We assessed Cdc42 expression levels of various constructs. Next to the expected Cdc42 band at 25-30 kDa, we observed a 50-60 kDa band for almost every construct in the anti-His and anti-Cdc42 Western blots (Fig. 2b). Its presence in the anti-Cdc42 Western blot confirms that this band is also indeed Cdc42, suggesting that Cdc42 forms dimers. Cdc42 belongs to the group of small GTPases, of which some have been shown to dimerise or oligomerise. Dimerisation has been linked to the PBR and (partial) removal of the PBR was shown to impede dimerisation. An illustration of this is given in Supplement S1 Tab. 1. Different GTPases have similar, yet slightly distinct PBR sequences. *S. cerevisiae* Cdc42 has the PBR sequence that has neither been linked to dimerisation nor to its absence. As the direct relation between the PBR sequence and protein dimerisation is still largely unknown, it is difficult to say, based on the PBR sequence, if Cdc42 dimerises. *In vivo* data suggests that it does not [40], but the absence of Cdc42 dimers *in vivo* does not necessarily exclude their existence *in vitro*: Cdc42 occupies a central role in the yeast cell cycle and has a lot of binding partners [9], making it part of a lot of protein complexes. Cdc42 dimerisation was assessed though the fluorescence signal of two Cdc42-YFP fusions with YFP truncations: One Cdc42 copy was fused to the N-terminal part of YFP, the other Cdc42 copy to the C-terminal part. Fluorescence appears when both YFP fragments are brought together through association of two Cdc42 molecules [40]. Hence, only dimeric Cdc42 leads to a YFP signal, monomeric Cdc42 and Cdc42 in hetero-complexes does not. If only a small fraction of Cdc42 dimers are Cdc42-Cdc42 homodimers (compared to Cdc42-effector protein heterodimers), not enough YFP signal is generated to observe these homodimers.

To further investigate this possibility, we ran purified Cdc42 sample on a size-exclusion chromatography (SEC) column and used

Figure 4. Most Cdc42-sfGFP^{SW} and -mNeonGreen^{SW} constructs express, but all show degradation. Expression levels ('f': 3 h 37°C 1.0 mM IPTG, 's': 18 h 18°C 0.2 mM IPTG) were analysed by anti-His Western blotting. The black arrow indicates the size of monomeric Cdc42 and the green arrow indicates that of the fluorescent Cdc42 fusions. Bands annotated with an asterisk (lane 22) are not from the protein in that lane, but bands of the protein standard that migrated to this lane during the blotting process.

multi-angle light scattering (MALS) to determine the molecular weight of the protein in each peak (Fig. 2c). Cdc42 ran in one peak and the SEC-MALS molecular weight corresponded to that of a monomer, independent of construct type or added nucleotide. We subsequently analysed the protein of these peak fractions by anti-His and anti-Cdc42 Western blotting. Again, the blots showed bands of the size of a dimer. Interestingly, the presence of dimers on SDS-Page was again influenced by the Strep-tag. Constructs with an N-terminal Twin-Strep-tag (SS-Cdc42-H, H-SS-Cdc42-F) formed no dimers (Fig. 2b), and S-Cdc42-H showed dimers in the anti-His, but not in the anti-Cdc42 Western blot (both in the expression test (Fig. 2b) and SEC-MALS samples (Fig. 2c)). C-terminal tags did not induce or influence dimer formation; type S constructs, and type D constructs with all possible C-terminal tags (6His, Flag, Strep-II, Twin-Strep) formed dimers (Fig. 2b). Thus, N-terminal Strep-II and Twin-Strep-tags seem to interfere with Cdc42 dimerisation under denaturing conditions. The origin of this remains elusive.

Taken together, we conclude that Cdc42 does not form stable long-lasting dimers *in vitro*. From this data we can not exclude that it forms transient and weakly bound dimers, as such complexes would not sustain themselves under the constant flow under which SEC is performed. Our data shows that it dimerises under denaturing conditions. This can be an artefact that has no translation to the behaviour of the folded protein, or mean that Cdc42 has some, still unexplored, potential to dimerise.

Cdc42-mNeongreen^{SW}/-sfGFP^{SW} shows a more varied expression and purification behaviour than Cdc42

We explored the expression and purification behaviour of fluorescent Cdc42 fusions. Double-tagged Cdc42 was mostly unaffected by its specific N- and C-terminal tags (with the exception of an N-terminal Twin-Strep-tag). We assessed if this was also true for sandwich fusions of Cdc42 and sfGFP or mNeonGreen [33]. We conducted expression tests (condition 'f' and 's') and analysed them by anti-His Western blotting (Fig. 4). We tested combinations of N- and C-terminal 6His-, Flag-, Strep-, and C-terminal Twin-Strep-tags. We did not conduct tests with constructs with an N-terminal Twin-Strep-tag, as they already lead to issues with Cdc42 alone.

Most fusion-constructs showed bands at the expected size of 55 kDa (Fig. 4, green arrow). Only H-Cdc42-mNeon-F and H-Cdc42-mNeon-S did not express the full-size protein but only parts of it (Fig. 4 lane 2, 3, 10, 11). This is surprising as their Cdc42 equivalents showed no expression problems at all (Fig. 4 black arrow, lane 1 and 9). In contrast to Cdc42, most Cdc42-mNeon^{SW}/-sfGFP^{SW} fusions showed additional lower bands, that can originate from degradation or translation processes that terminated prematurely. This is not surprising, as the fluorophore sequence was inserted directly into the Cdc42 sequence, potentially making the fusion more susceptible to degradation.

In most cases both Cdc42-mNeonGreen^{SW} and Cdc42-sfGFP^{SW} fusions expressed with roughly the same amount and to same level of degradation, through significantly less strong than Cdc42. An exception is F-Cdc42-sfGFP-H (Fig. 4 lane 7 and 8), that expressed at a much higher amount than its mNeonGreen equivalent (Fig. 4 lane 5 and 6). It also shows higher amounts of degradation, even though it is possible that the degradation bands are only more pronounced due to the generally higher expression levels.

In most cases no big difference between the slow ('s') and fast ('f') expression condition could be observed; the fast condition lead to slightly more protein than the slow condition, but also showed a proportionally higher amount of degradation bands.

Figure 5. F-Cdc42-sfGFP-H can be purified in a single His-affinity chromatography step. The protein was eluted in a gradient of His-AC washing buffer (50 mM Tris-HCl (pH=8.0), 1 M NaCl, 5 mM imidazole, 1 mM 2-mercaptoethanol) and His-AC elution buffer (50 mM Tris-HCl (pH=8.0), 100 mM NaCl, 500 mM imidazole, 1mM 2-mercaptoethanol). SDS-Page of the elution peak, the band of the full-length protein is indicated by an orange arrow.

An exception is S-Cdc42-sfGFP-H (Fig. 4 lane 20 and 21) where less degradation is present in the fast compared to the slow condition. The opposite is true for the mNeonGreen equivalent (Fig. 4 S-Cdc42-mNeon-H, lane 18 and 19). Here the slow condition expressed roughly the same amount of protein with less degradation.

In contrast to Cdc42, the expression behaviour of fluorescent Cdc42 fusions is very heterogeneous. Full-size protein can be produced with several, but not all, N- and C-terminal tag combinations. Degradation bands are observed in all cases but to different degrees, depending on the combination of (1) fluorophore used (sfGFP or mNeonGreen), (2) expression condition ('f' or 's'), and (3) N- and C-terminal tags. From our data no apparent relation between these factors could be drawn.

Next, we purified a subset of the constructs used in the expression test. We chose the constructs based on our applications and did not do entire screens. The data shown here therefore acts as exemplary evidence. We purified F-Cdc42-sfGFP-H, H-Cdc42-sfGFP-SS, F-Cdc42-mNeon-H, and S-Cdc42-mNeon-BC-H.

F-Cdc42-sfGFP-H could be purified in a high yield (5 mg per 1 L expression volume) in a single step His-AC (Fig. 5). The other constructs required multiple chromatography techniques and gave a very low yield (<0.1 mg per 1 L expression volume). Addition of detergents to the buffers increased the yield, but not to a substantial degree. We suspect that the high expression levels of F-Cdc42-sfGFP-H, which are already visible in the expression test (Fig. 4 lane 7,8), lead to the high yield. An extended discussion of the purification of these constructs is given in Supplement S2.

We further observed for all constructs that the lysis step was incomplete (about 30-50% of cells lysed), even though the same protocol resulted in 100% lysed cells if a non-fluorescent Cdc42 construct was used. The lysis could be slightly improved when the number of lysis rounds with the high pressure homogenizer were doubled from five to ten. The origin of this remains elusive.

Based on this data, we suggest to use F-Cdc42-sfGFP-H due to its easy purification procedure which results in a high yield. If another construct is required, we advise to screen for one that has the highest yield after His-AC, as we found this to be the limiting factor.

Cdc42's GTPase activity and interaction with the GEF Cdc24 can be reliably assessed using GTPase-Glo assays

Cdc42 is a small GTPase and can therefore hydrolyse GTP. To test how active the various Cdc42 constructs are, we performed GTPase assays using the Promega GTPase Glo™ assay. Here serial dilutions of Cdc42 were incubated with GTP for a certain time, after which the reactions were stopped and the amount of remaining GTP was measured (see materials and methods). To compare the GTPase activity of different Cdc42 constructs, we determined GTP hydrolysis cycling rates k. These rates encompass the entire GTPase cycle, which can be described in three steps (Fig. 6a): (1) Cdc42 binds to a free GTP. (2) GTP gets hydrolysed by Cdc42. (3) Cdc42 releases GDP.

If one construct shows decreased rates k it indicates that at least one of these steps is occuring at a slower rate, suggesting that this batch has less active Cdc42 or a higher ratio of inactive to active protein. If batches of a certain construct show this behaviour repeatedly, the N- and/or C-terminal additions are likely interfering with its functionality.

Figure 6. Cdc42's GTPase activity and interaction with the GEF Cdc24 can be quantified using GTPase Glo assays (Promega). (A) Schematic illustration of the GTPase cycle. (B) GTPase Glo assay: The amount of remaining GTP declines exponentially over time. Dashed lines are fits (Eq. 1). (C) H-SS-Cdc42-F shows cooperative behaviour (high k_2 and $k_1 \approx 0$) and S-Cdc42-H exhibits almost no cooperativity (high k_1 and $k_2 \approx 0$). (D) The overall GTP hydrolysis cycling rate scales non-linearly with the Cdc24 concentration.

First, we examined how the GTP concentration changes for different Cdc42 concentrations over time. We performed experiments to measure the amount of remaining GTP after incubation times of 1.5 h, 3 h, and 5 h. The graph of the amount of remaining GTP over time (Fig. 6b) shows that the GTP hydrolysis process can be described by an exponential decline (Eq. 1). The assay data of all three incubation times gave the same fit (Fig. 6b), suggesting that one data set from one assay, i.e. one incubation time, is sufficient. In this assay we used Cdc42 concentrations of 0.75 to 2.00 µM. For assays comparing the GTPase activity of various Cdc42 constructs we used wider concentration ranges (1 - 9 µM Cdc42 in combination with incubation times of 1 -1.5 h) to ensure a better fit quality.

We determined the GTP hydrolysis cycling rates of Cdc42 using

$$[GTP]_t = [GTP]_{t_0} \exp(-Kt)$$
$$\text{using } [GTP]_{t_0} = 100\% \tag{1}$$
$$\text{and } K = k_1[Cdc42] + k_2[Cdc42]^2$$

We will refer to K as the 'overall GTP hydrolysis cycling rate' in the following. In this equation k_1 describes the GTP hydrolysis cycling rate of a single Cdc42 molecule and k_2 includes any effects due to Cdc42 dimerisation or cooperativity. Introduction of this two-parameter fit was needed as some Cdc42 constructs showed cooperative behaviour (high k_2 and $k_1 \approx 0$), whereas others showed almost none (high k_1 and $k_2 \approx 0$) (Fig. 6c).

Next to Cdc42's GTPase activity, we also investigated its interaction with the GEF Cdc24. As a GEF Cdc24 boosts the release of GDP from Cdc42 (GTPasy cycle step (3), Fig. 6a) and thereby increases the cycling speed of the GTPase cycle.

Assays with Cdc24 dilutions show that Cdc24 is highly active; addition of sub-µM concentrations of Cdc24 to 1 µM Cdc42 were sufficient to boost the reaction cycle by a factor of about three (Fig. 6d).

Further, the data showed that the GTP decline does not depend linearly on the Cdc24 concentration, but is better approximated by a quadratic term. This suggests cooperativity. Previous work showed that Cdc24 has the capability to oligomerise via its DH domain [41]. We expect dimers and oligomers to have an increased GEF activity. This could, for example, be facilitated by Cdc24's C-terminal PB1-domain, which has been suggested to reduce Cdc24's GEF activity through intramolecular interactions [42]. Cdc24 oligomerisation could interfere with this self-interaction and thereby increase the proteins GEF activity. We note that previous this assumption could be seen to contradict previous *in vitro* work: Cdc24 peptides, consisting only of Cdc24's

DH and PH domain, showed that these peptides exhibit GEF activity. The GEF activity was not changed when oligomerisation was inhibited (through mutations) or amplified [41]. These findings would exclude that Cdc24 oligomers exhibit an increased GEF activity. However, one has to be careful when applying these findings to full-size Cdc24: For one, not full size protein, but only peptides were used. Other domains that are not directly involved in oligomerisation or GEF funtion can still affect these properties. For example, the PB1 domain was suggested to reduce Cdc24 GEF activity in a self-inhibitory fashion [42]. Next, samples representing a heightened oligomerisation state were produced through addition of an additional oligomerisation domain to the peptides. This domain was not related to Cdc24 and could be triggered to oligomerise through the addition of a chemical. It is thus questionable if the GEF activity of these oligomerised peptides relates to the GEF activity of oligomeric Cdc24.

To determine Cdc42-Cdc24 interaction rates k_3, we incubated Cdc42:Cdc24 mixtures, as well as samples containing only Cdc42, with GTP for 1-1.5 h and measured the amount of remaining GTP. We had observed that rates for Cdc42 can vary slightly between assays (Fig. 7a). To account for this variance, we introduced the parameter c_{corr}, that maps all factors that lead to variations between assays onto the Cdc42 concentration. The assay data, including samples containing only Cdc42 and Cdc42 - effector protein mixtures, was fitted with

$$[\text{GTP}]_t = [\text{GTP}]_{t_0} \exp(-Kt)$$
$$\text{using } [\text{GTP}]_{t_0} = 100\%$$
$$\text{and } K = k_1 c_{corr}[\text{Cdc42}] + k_2(c_{corr}([\text{Cdc42}])^2 + k_3 c_{corr}[\text{Cdc42}][\text{Cdc24}]^2$$

(2)

to determine c_{corr} and k_3 (using k_1 and k_2 values determined earlier (Eq. 1)). Values of c_{corr} are 0.7 - 1.6, with the majority close to 1.0 (see Supplement S3).

We used these assays to examine the variation (1) within assays, (2) between purification batches of the same Cdc42 construct, and (3) between Cdc42 constructs.

The variation between assays is generally small

We conducted several assays with H-Cdc42-F. In all assays the overall GTP hydrolysis cycling rate K increases similarly with Cdc42 concentration; data points of fits from three assays overlay (Fig. 6a). Variations observed can be due to small concentration differences introduced though pipetting of small volumes (as are required for this assay), small temperature and shaker speed fluctuations during the GTPase reaction, and/or intrinsic changes in the Cdc42 protein due to other external conditions. Fitting each measurement data with k_1 and k_2 results in values that vary a bit (Fig. 7b, blue dots). However, they describe a similar GTPase behaviour: We used the rates and Eq. 1 to calculate how much GTP would remain in a simulated GTPase assay with equimolar amounts of Cdc42. All three k_1 and k_2 pairs lead to similar amounts of predicted remaining GTP (Fig. 7c, blue bars with blue dot). We assume that the variation in k_1 and k_2 is due to fitting quality. Measurements using more data points, or restricting the fit to either k_1 (no cooperativity) or k_2 (cooperativity) would result in less variation. As the aim of this assay was to compare different constructs, and as constructs showed a heterogeneous tendency towards cooperativity (Fig. 6c), we considered the observed variations in k_1 and k_2 unideal but necessary. We will use them as a baseline variability to compare purification and construct variations with.

We further used H-Cdc42-F to asses the variability of the Cdc42-Cdc24 interaction rate k_3 between assays. Variations in k_3 were bigger than those in k_1/k_2: For H-Cdc42-F, values of 2.2 to 4.5 $\mu M^{-3} h^{-1}$ were observed (Fig. 7d, blue dots). As Cdc24 boosts the GTPase cycle in a strong fashion that depends non-linearly (quadratically in our fits) on its concentration (Fig. 6d), small activity changes of Cdc24 and dilution inaccuracies between assays affect k_3 quadratically. For example, if the intended Cdc24 concentration was 0.2 μM, but in fact 0.1 μM was present in the assay, k_3 will decrease by a factor of four. To accurately determine k_3 one thus needs to perform several assays. To compare different Cdc42 constructs, one ideally would add them all to the same assay. If this is not feasible, addition of a reference Cdc42 construct to all assays is advantageous, as then k_3 values can be normalised/ compared to k_3 of the reference construct. If this was not done, the degree of k_3 variation needs at least to be considered. Constructs can then only be considered to have a stronger/ weaker GEF interaction, if their k_3's are significantly above or below the observed variation.

Some purification batches are not functional

We assessed two purification batches for H-Cdc42-F and S-Cdc42-H, and three purifications for F-Cdc42-H (Fig. 7b-d, indicated by distinct symbols of the same colour). All batches were expressed and purified using the same protocol, and we expected to see only minor differences.

Figure 7. Cdc42 purification batches can show variations in their GTPase activity and GEF interaction or be non-functional. (A) Seperate GTPase assays show a similar GTPase activity for Cdc42, but purification batches can vary in their activity. (B) GTP hydrolysis cycling rates k_1 and k_2 for different Cdc42 constructs and purification batches. (C) Predicted amount of remaining GTP for Cdc42 constructs (5 μM) after an incubation time of 1.5 h, calculated using rates k_1, k_2 shown in (B). Symbols indicate if rates of a specific purification batch were used. (D) Cdc42-Cdc24 interaction rates k_3. (A-D): GTP hydrolysis cycling rates k_1 and k_2 and Cdc24 interaction rates k_3 for different Cdc42 constructs. Data points of the same colour refer to the rates obtained from separate measurements/assays. Data points of the same colour but a distinct shape (circle, triangle, rhombus) represent different purification batches of the same construct. Summary graphs of the data, values, and overview figures are given in Supplement S3.

Surprisingly, one purification batch of H-Cdc42-F showed a hugely reduced GTPase activity (Fig. 7a). We assume that this batch contained either protein with a reduced activity or a substantial portion of inactive protein. We discarded this batch from further analysis.

One of three purification batches of F-Cdc42-H had a reduced GTPase activity (Fig. 7b,c, green triangle, and Supplement S3). It was less severe than the reduction observed for H-Cdc42-F. Its GEF interaction rate k_3 was also similar to that of the more active F-Cdc42-H batch (Fig. 7d), given the observed variability of k_3. We conclude that this batch is a bit less active that it ideally could be, but still shows sufficient activity. We also observed a reduced GTPase activity for one purification of S-Cdc42-H (Fig. 7b,c, orange triangle, and Supplement S3), and the reduction was similar to that in F-Cdc42-H. In contrast to F-Cdc42-H, this Cdc42 batch did not interact with Cdc24 (Fig. 7d), suggesting that Cdc42's GTPase activity is independent from its ability to interact with Cdc24. One can thus not deduce full protein functionality from GTPase activity data alone.

The decrease in GTPase activity of certain purifications is visible in this data set as a drop in k_2: k_1 of the less active batch is within error of the more active batch, but k_2 drops close to zero (Fig. 7b). A lack of cooperative behaviour could explain the reduced overall activity; Cdc42 molecules in these batches would still possess their baseline GTPase activity, but lack the ability to interact and enhance each others GTPase activity, thus leading to an overall reduced activity. On the other hand, the observed drop in k_2 could also be a fitting artefact due to the decreased Cdc42 activity; a drop in activity in the used Cdc42 concentration range can make the data appear more linear than it actually is (Supplement S1). Use of a wider concentration range for these constructs would show if they have indeed less cooperative behaviour or if the perceived drop is an artefact. The data we collected is limited and of exemplary nature, and further experiments are required to claim that is is indeed the case.

All Cdc42 constructs, with the exception of SS-Cdc42-H, are active

Finally, the different constructs can now be compared. GTPase data (Supplement S3) were fitted to determine GTP hydrolysis cycling rates k_1 and k_2 (Fig. 8a) and GEF interaction rates k_3 (Fig. 8c). The amount of remaining GTP in a simulated GTPase assay with equimolar amounts of each construct using rates k_1 and k_2 is shown in Fig. 8b.

Most double-tagged constructs (type D: H-Cdc42-F, F-Cdc42-H, H-Cdc42-SS, S-Cdc42-H, S-Cdc42-BC-H) show similar GTPase activities and a weak cooperativity (both k_1 and k_2 are unequal to zero) (Fig. 8a,b), and GEF interaction rates that fall within the assay variability observed for H-Cdc42-F (Fig. 8c). Both GTPase activity as well as GEF interaction are similar to the only N-terminally tagged H-Cdc42:CTIS (type S). Even S-Cdc42-BC-H, which contained further C-terminal modifications, does not show any reduced activity (Fig. 8). This suggests that C-terminal additions do not hinder Cdc42's functionality (GTPase activity and GEF interaction). SS-Cdc42-H has almost no GTPase activity and does not interact with Cdc24 (Fig. 8 red dots/bar). SS-Cdc42-H showed already precipitation issues during the purification process and this assay confirms that the protein is barely functional. (We show only one data point/one assay for this construct. However, several additional assays were performed to ensure that the observed behaviour is not an artefact. The same behaviour was observed repeatedly and in most assays no GTPase activity was observed.

Figure 8. All Cdc42 constructs of type D, with the exception of SS-Cdc42-H and H-SS-Cdc42-F, show a similar GTPase activity and GEF interaction. (A): GTP hydrolysis cycling rates k_1 and k_2 of different Cdc42 constructs. Data points of the same colour refer to the rates obtained from separate measurements/assays. Data points of the same colour but a distinct shape (circle, triangle, rhombus) represent different purification batches of the same construct. (B): Predicted amount of remaining GTP for Cdc42 constructs (5 µM) after an incubation time of 1.5 h, calculated using rates k_1, k_2 shown in (A). Symbols indicate if rates of a specific purification batch were used. (C) Cdc42-Cdc24 interaction rates k_3. (A-C): Summary graphs of the data, values, and overview figures are given in Supplement S3.

The rates could only be extract from one assay where a significantly longer incubation time was used.) H-SS-Cdc42-F also has a lower GTPase activity and a lower GEF interaction ($0.5 - 1.5\,\mu M^{-3} h^{-1}$) (Fig. 8, purple symbols). It is more active than SS-Cdc42-H but less than the other constructs. This is true for all three purification batches tested. In the same assay that we used to assess the GEF interaction of H-SS-Cdc42-F, we had also added H-Cdc42-F as a control. In this assay H-Cdc42-F had a GEF interaction rate of $4.5\,\mu M^{-3} h^{-1}$. The low k_3 of H-SS-Cdc42-F is thus not due to assay variation, but describes a reduced or weaker interaction with Cdc24. This data shows that SS-Cdc42-H is not active and that H-SS-Cdc42-F has a reduced activity. The 6His-tag upstream of the Twin-Strep-tag seems to restore Cdc42 functionality thus partially, but not fully.

Taken together, our data shows that the GTPase activity of Cdc42 can be assessed with the GTPase Glow assay (Promega). The variation between assays containing only Cdc42 is small. Cdc42 is heterogeneous in its tendency cooperativity, and a lack of any cooperativity might be a potential indication for a less functional protein. We observe a stronger assay variation for assays assessing the Cdc42 Cdc24 interaction. To effectively compare different Cdc42 constructs, one thus has to account for this assay variation through assessing it, adding a reference to every assay, or by only comparing constructs, which rates were measured in the same assay. We further see that not all purification batches give fully functional protein and that constructs can have a GTPase activity but show no GEF interaction. Thus, every purification batch has to be assessed for its activity. Our data is not sufficient to provide quantitative measures that aid the decision process of if a construct is sufficiently active, but acts as exemplary evidence to guide the process and to highlight outcomes that can be observed. Our data suggests that mostly unstructured C-terminal additions to Cdc42 do not interfere with its GTPase activity and GEF interaction, but that the N-terminal Twin-Strep tag does. Placement of a 6His-tag prior to the Twin-Strep-tag partially, but not fully, restores Cdc42 functionality.

Pulldown assays are less robust to study weakly bound Cdc42 effector interactions as the Cdc42-Bem1 complex

We then wanted to explore other Cdc42 - effector interactions. The GTPase assay worked well to investigate the Cdc42 Cdc24 interaction, but is limited to only those interactors of Cdc42 that affect its GTPase cycle. Pulldown assays, which directly assess protein - protein binding, can be used to study a wider range of interactions. Here protein A is tethered to a matrix using its affinity tag (e.g. a Flag-tag). Protein B, which does not contain this affinity tag, is added. The matrix is washed several times. Then protein A is eluted. If protein B is in this elution as well, it suggests that it is bound to protein A (Fig. 9a). We conducted Flag-pulldown experiments with Bem1 and Cdc42, using Flag-tagged Bem1 (H-Bem1-F) and Cdc42 constructs that do not contain a Flag-tag.

We observed that binding between Bem1 and Cdc42 occurred, but that it was so weak that Cdc42 could only be detected using Western blotting (Fig. 9c). In comparison, pulldown experiments with the same Bem1 construct and Cdc24, another binding partner of Bem1, showed so high amounts of Cdc24 in the elution fraction that it was visible on SDS-Page, indicating stronger binding. Cdc42 alone did not bind to the anti-Flag affinity gel. The Bem1 - Cdc24 interaction was also specific: Ovalbumin, a protein not interacting with Bem1, was not present in the elution fraction (Fig. 9b).

Figure 9. Flag-pulldown experiments are suitable to study strongly binding proteins (Cdc24-Bem1), but less reliable for weakly interacting proteins (Cdc42-Bem1). (A) Schmematic illustration of Flag-pulldown experiments. (B) SDS-Page of elution fractions of Flag-pulldown experiments with H-Bem1-F, Cdc24-H, and Ovalbumin. (C) Anti-His Western blots of elution fractions of three Flag-pulldown experiments with H-Bem1-F and Cdc42 preloaded with no nucleotide, GDP (D), or GTPγS (T).

We used several Strep-tagged Cdc42 versions (Fig. 9c): S-Cdc42-H, H-Cdc42-SS, and SS-Cdc42-H. All of these Cdc42 constructs bound to Bem1. The blotting signal for H-Cdc42-SS was weaker in comparison to S-Cdc42-H and SS-Cdc42-H. This is surprising, as we expected the barely functional SS-Cdc42-H to not, or only very weakly, bind to Cdc42. SS-Cdc42-H either still shows some remaining functionality (ability to bind to Bem1), or all our binding events are due to nonspecific interactions (and thus only detectable with Western blots). We tested if Strep-tagged Cdc42 (S-Cdc42-H) would also stick to the anti-Flag affinity gel, thus causing the presence of Cdc42 in all elution samples. In absence of H-Bem1-F, a small amount of S-Cdc42-H was binding to the anti-Flag affinity gel. Its signal in the Western blot however was significantly weaker than when Cdc42 was incubated with anti-Flag affinity gel in presence of H-Bem1-F (Fig. 9c). Out data suggests that there is weak binding between Cdc42 and Bem1, but we can not exclude that this is due to non-specific interactions between both proteins. In order to test this, one would need to add another control in which Bem1 and another His-tagged protein, that is not a binding partner of Bem1, are used. We did conduct this control due to the lack of a suitable His-tagged protein.

Bem1 was shown to specifically bind to GTP-bound, but not GDP-bound, Cdc42 [30]. We conducted two separate Flag-pulldown experiments in which 1 nmol Cdc42 got pre-loaded with no nucleotide, 100 nmol GDP, or 100 nmol GTPγS (a non-hydrolysable variant of GTP). The intensities of the Western blot signal for Cdc42 varied with the nucleotide type, but also per experiment: In one experiment more Cdc42-GDP (than Cdc42-GTP or nucleotide free Cdc42) bound to Bem1, but in another experiment it was nucleotide-free Cdc42 (Fig. 9c). We thus could not observe a significant effect of the nucleotide state on the Cdc42-Bem1 interaction. This further points to the conclusion that something in our Cdc42 - Bem1 interaction assays is not properly functioning. We also spotted several small experimental differences between the pulldown assays shown here and those conducted by Bose *et al.*: Bose *et al.* conducted similar pulldown experiments as shown here, but used GST-tagged (26 kDa) instead of a Flag-tagged (3 kDa) proteins (and subsequently GST agarose instead of anti-Flag affinity gel). Given the size of the GST-tag, it can not be excluded that this tag is destabilising the Bem1-Cdc42 interaction, thereby exaggerating the effect of a conformational state of Cdc42 on Bem1 binding. Pulldowns with (1) GST-Cdc42 and Bem1, and (2) GST-Bem1 and Cdc42, both showed the same result. Thus, the GST-tag would need to be positioned in both cases in such a way that it destabilise the interaction, questioning this hypothesis. Another experimental difference is the way in which Cdc42 was locked in one nucleotide state: We pre-loaded Cdc42 with excess GDP or GTPγS, Bose *et al.* used Cdc42-mutants that are locked in a GDP- or GTP-state. One interpretation is that GDP/GTP-loaded Cdc42 and Cdc42 mutants are structurally distinct in a way that affects Cdc42-Bem1 binding. Alternatively, it could indicate that our pre-loading was insufficient and that it did not bring the majority of Cdc42 into its GDP-bound conformation.

In conclusion, we find that pulldown assays are a great tool to study the interaction of strongly interacting proteins. Here binding

can be analysed by SDS-Page, and controls for non-specific binding can easily be added. In contrast, weaker protein - protein interactions, as the Cdc42 - Bem1 interaction, are more difficult to study using pulldown experiments. We observed a very weak, but potentially non-specific, interaction between Cdc42 and Bem1, that also did not depend on the nucleotide state of Cdc42 (contradicting [30]. Western blots are very sensitive and can easily be overinterpreted. Controls to correct for signal from Cdc42 sticking to the matrix, and from non-specific protein-protein binding (Bem1 plus a protein that is not a binding partner of Bem1) are required. With sufficient preparation they can be a suitable tool, but are, in our experience, much more difficult and error-prone than GTPase assays.

TEV protease, but not enterokinase, is suitable to cleave tags off Cdc42

We created Cdc42 constructs with both an N- and C-terminal tag (type D), and one which only contains an N-terminal tag (type S). The effect of C-terminal additions (sortase site and purification tag) can be determined by comparing type D with type S. To asses the the effect of the N-terminal additions (incl. purification tag), an Cdc42 version lacking that region needs to be created. To generate such a Cdc42 version, we used the existing enterokinase site in H-Cdc42:CTIS (type S) to cleave the N-terminal region. We conducted a condition screen varying amounts of enterokinase used, reaction time and temperature (Supplement S4 Fig. 1a). In all conditions the majority of the cleavage products are smaller than the expected product; bands at ~14 kDa and ~8 kDa show a stronger intensity than the band of the expected cleavage product at ~20 kDa (Supplement S4 Fig. 1a). The enterokinase cleavage sequence is 'Asp-Asp-Asp-Asp-Lys', but it was reported that it can also cleave after a site of an acidic amino acid followed by a basic amino acid [43, 44]. We found two sequences that fit those requirements and that can explain the observed fragments (Supplement S4 Fig. 2).

As enterokinase showed an undesirable cleavage behaviour, we designed another Cdc42 construct in which the enterokinase site was replaced with a TEV cleavage site (H*-Cdc42:CTIS). We tested two commercially available enzymes (TEV (1): TEV protease (Sigma Aldrich), TEV (2): AcTEV (Invitrogen)) and one in-house purified TEV protease (TEV (3): pSF1818). We again conducted a condition screen varying amounts of protease used, reaction time and temperature (Supplement S4 Fig. 1b). In all conditions only the desired cleavage product was observed. Increasing the amount of protease used resulted in more cleavage product in the case of TEV (1) and TEV (2), but not with TEV (3). Prolonged reaction times did only marginally increase the yield, and reactions at room temperature and at 4℃ showed equal amounts of product (Supplement S4 Fig. 1b).

We cleaved the N-terminal 6His-tag from H*-Cdc42:CTIS in two large-scale reactions (using TEV (1) or TEV (3)), and used His-AC to separate cleavage product from uncleaved Cdc42 and protease. To ensure all uncleaved Cdc42 binds to the column, the sample/flow-through got loaded thrice onto the column. The flow-through contained the cleavage product (Fig. 10a). Anti-His Western blots show that a tiny fraction of uncleaved Cdc42 is still present in the flow-through (Fig. 10b). In accordance to the observation of Cdc42 dimers in expression tests before (Fig. 2), we again observed higher bands, correspond to Cdc42 dimers, in the Western blots (Fig. 10b). Both Cdc42 with the N-terminal region and Cdc42 without showed dimers on the blots. These dimers are thus due to properties of Cdc42 and not due to N- or C-terminal additions.

We tested the cleavage educts and products on their GTPase activity and ability to interact with the GEF Cdc24. H*-Cdc42:CTIS, containing an N-terminal TEV site, is similar in both of these two properties to H-Cdc42:CTIS, which contains an N-terminal enterokinase site (Fig. 10c-e). Thus, the cleavage site sequence (enterokinase vs. TEV protease cut site) does not influence protein function. The cleavage products of both reactions with TEV protease (*-Cdc42:CTIS) had a slightly reduced GTPase activity (Fig. 10c,d), though the magnitude of this reduction lies within assay variability and is thus likely not due to protein properties (see Supplement S1). It's interaction with Cdc24, however, was significantly reduced ($k_{3,}$*-Cdc42:CTIS $= 1.4-1.5\,\mu M^{-3}h^{-1}$, Fig. 10e). We measured the GEF interaction of H-Cdc42-F in the same assay: $k_{3,}$H-Cdc42-F $= 4.5\,\mu M^{-3}h^{-1}$. The low k_3 of *-Cdc42:CTIS is thus not due to assay variation, but describes a reduced or weaker interaction with Cdc24. It is possible that the N-terminal additions boost Cdc42's ability to interact with Cdc24 or that during the cleavage reaction the protein got damaged.

In conclusion, enterokinase is not a suitable enzyme for cleaving additions/tags off Cdc42. TEV protease might a better option, as it does not lead to undesired cleavage products and works robustly in many conditions. The cleavage product can easily be isolated using His-AC, even though a very small fraction of 6His-tagged Cdc42 remains mixed with the cleavage product after His-AC. However, the cleaved product shows a significantly reduced Cdc24 interaction. This indicates that during both cleavage reactions the protein got damaged or that the N-terminal additions to Cdc42 aid GEF interaction. If the protein got damaged, it questions if TEV protease is indeed a suitable cleavage enzyme for Cdc42.

Figure 10. N-terminal tags can be cleaved of Cdc42 with TEV protease: the cleavage product maintains its GTPase activity but shows a reduced interaction with the GEF Cdc24. (A) N-terminal tags can be cleaved off Cdc42 with TEV protease and the cleavage product can be purified using His-AC (FT: flow through). (B) SDS-Page and Western blots of the purified cleavage reaction product of reactions shown in (A). (C) GTP hydrolysis cycling rates (k_1, k_2) of different Cdc42 constructs. (D) Predicted amount of remaining GTP for Cdc42 constructs (5 µM) after an incubation time of 1.5 h, calculated using rates k_1, k_2 shown in (C). Symbols indicate if a specific protein batch was used. (E) Cdc42-Cdc24 interaction rates k_3. (c-E) Summary graphs of the data, values, and overview figures are given in Supplement S3.

Conclusion and Discussion

Here, we shared our experiences of working with *S. cerevisiae* Cdc42 *in vitro*, to make its *in vitro* use more accessible to a broader audience. We discussed our construct design and the expression and purification of Cdc42 and Cdc42-mNeonGreen^{SW}/-sfGFP^{SW} constructs. We explored two assays for Cdc42 activity checks (GTPase Glo assay (Promega), and Flag-pulldown assay), and investigated tag cleavage with enterokinase and TEV protease.

Most Cdc42 constructs expressed in several conditions and could be purified in a high yield. The exceptions were H-Cdc42 (construct type H, where only a 6His-tag was added N-terminally) and SS-Cdc42-H (type D). H-Cdc42 did not express at all, and we suspect that either a linker region between an N-terminal tag and Cdc42, or the T7 tag in particular, is required for Cdc42 expression in *E. coli*. SS-Cdc42-H expressed well, but precipitated after purification even at low concentrations and showed almost no GTPase activity. As neither S-Cdc42-H, F-Cdc42-H nor H-Cdc42-SS exhibited these issues, we conclude that the N-terminal Twin-Strep-tag is responsible for this. We also observed negative effects of the N-terminal Twin-Strep-tag for Cdc24 and Bem1 constructs, which did not express well or at all. These two protein are larger and less tightly folded than Cdc42, making them more susceptible to degradation and/or folding issues. We think that these early and easily detectable expression issues already indicate a general problem with the construct design: in the case of the bigger and less stable Cdc24 and Bem1 it already showed up on the expression level. In the case of the smaller and more stable Cdc42 it was only observable after purification. We suggest to not use an N-terminal Twin-Strep-tag on (any) protein constructs. Our data shows that even a stable and easy to work with protein (as Cdc42 is) can have a few conditions where it does not behave (express or be stable), creating unpredictable barriers of entry for its *in vitro* use.

These issues were exacerbated for Cdc42-mNeonGreen^{SW} and -sfGFP^{SW}. Here additional constructs (H-Cdc42-mNeon-F, H-Cdc42-mNeon-S) did not express the full-size protein, even though their Cdc42 counterparts did. It seems that purification tags influence the expression and degradation levels for those constructs. Cdc42 is a rather small and tightly folded protein. The sandwich fusions of Cdc42 and sfGFP and mNeonGreen [33] consist of two folded proteins parts. Because the sequence of sfGFP/mNeonGreen is inserted into that of Cdc2, Cdc42 can only completely fold if the fluorophore is fully folded. As long as both protein parts are not full folded, the entire construct is more susceptible to degradation. It is therefore not surprising to

see almost no degradation bands in the Cdc42 expression tests, but many degradation bands in the Cdc42-mNeonGreen^{SW}/-sfGFP^{SW} expression tests. However, the reason why the two tag combinations, that hinder expression of the full-size protein, exactly have their observed effect remains elusive to us.

After purifying a protein, it ought to be tested for its activity. We used the GTPase Glo assay to investigate the GTPase activity of Cdc42 constructs and their interaction with the GEF Cdc24, and assesd Flag-pulldown assays on their usability to study the interaction of Cdc42 with its other interactors.

The GTPase assay is a suitable assay to test the GTPase activity of Cdc42; it includes all steps in the GTPase cycle and can measure the effect of GEFs and GAPs as well. Our data showed that there was little variation between assays assessing the same Cdc42 purification batch. There can, however, be a large variation between purification batches of the same Cdc42 construct (even though the same purification protocol and conditions were used). This means that: (1) One should test all Cdc42 purification batches on their activity, as purification batches can vary a lot. (2) When comparing different Cdc42 constructs, one should purify several batches of each construct and test each. Otherwise construct differences can be confused with what are in fact purification batch differences. (3) One should be critical/reserved in assuming that kinetic rates and other protein-specific parameters derived from other Cdc42 batches are exactly applicable to one's protein. The exact numbers of such rates can be both context specific (e.g. depend on buffer components, temperature, ...) and purification batch specific.

We also used the GTPase assay to study the Cdc42 - Cdc24 (GEF) interaction. Cdc24 was highly active, and sub-micromolar additions boosted Cdc42 GTPase cycle drastically. We suspect that this leads to the larger variation of Cdc24's effect between assays; small dilution errors and activity changes of Cdc24 affect the entire GTPase cycle stronger than similar errors and changes in Cdc42. To quantify the Cdc42 - Cdc24 interaction, one should thus conduct the assays a sufficient numberr of times to get a sense of the 'normal' variation. When using this assay to quantitatively compare the GEF interaction of several Cdc42 constructs, one should thus add all Cdc42 constructs to the same assay. If this is not feasible, the addition of a reference Cdc42 construct to all assays is advantageous, as then rate values can be normalised to the rate of the reference construct in each assay.

Considering these assay and purification batch variations, we conclude that most Cdc42 constructs showed a similar GTPase activity and GEF interaction. Surprisingly, all C-terminal additions did not impede Cdc42's function. So even though so-far no C-terminally tagged Cdc42 was used in *in vitro* studies, these seem to be acceptable additions. On the contrary, Cdc42's N-terminus seemed to be more delicate: Constructs with an N-terminal Twin-Strep-tag were not active at all and those with an N-terminal 6His-Twin-Strep-tag showed both a reduced GTPase activity and GEF interaction. Additionally, Cdc42 construct type H did not express at all, and Cdc42 where the N-terminal tag got cleaved showed also a reduced GEF interaction. When designing new Cdc42 constructs, we thus advise to be more careful when placing N-terminal additions, and to check literature for evidence that the intended additions are not interfering with Cdc42's functionality. If such data is not available, we advise to design several variations of this construct and test them in parallel.

In contrast to GTPase assays, we found Flag-pulldown assays to be potentially more difficult for studying protein functionality. They are versatile for studying any protein - protein interaction, as only two differently tagged proteins are needed. In our experience they also gave clear results when working with two strongly interacting proteins (e.g. Cdc24 and Bem1). This must, however, not be true for Cdc42 - effector interactions. We tested the Cdc42 - Bem1 interaction, and could not reproduce previous results [30]. The Western blot data showed blotting intensities that varied per experiment and would be easy to misinterpret, when not done repeatedly. We think this assay can be a useful tool, but requires several controls and replicas to ensure genuine outcomes.

Lastly, we tested two cleavage enzymes to cleave the N-terminal tag of Cdc42 to produce Cdc42 without additions. Our data showed that enterokinase is not a suitable enzyme, as it cleaves also within Cdc42. TEV protease showed a better cleavage behaviour. The cleavage product, however, showed a reduced GEF interaction. This reduction could have several causes: (1) Due to external circumstances Cdc42 misfolded (as we saw during the purification of one S-Cdc42-H construct) during the cleavage reaction. (2) The reaction with TEV protease lead to the changes in Cdc42, and the TEV enzyme might not be a good choice for cleaving tags off Cdc42. (3) The N-terminal additions on Cdc42 enhance its ability to interact with Cdc24, facilitating the reaction. From our data we can not identify which of these causes is most likely and additional experiments are needed. These include repeating the cleavage with TEV protease, using another cleavage enzyme, and expressing only C-terminally tagged Cdc42. Depending on the cause, this speaks for or against cleaving N-terminal additions off Cdc42: If Cdc42's N-terminus is sensitive and changes/ modifications easily result in protein misfolding and a reduced activity, N-terminal tags ought to be left unchanged, thus avoiding the possibility of damaging the protein. If N-terminal tags/additions on Cdc42 facilitate its GEF interaction, they should be cleaved. This would also mean that all previously conducted *in vitro* studies using N-terminally tagged Cdc42 potentially show an effect facilitated by additions/tags, as other Cdc42 - effector interactions could be equally effected by the tag.

Studies comparing and parameterising Cdc42 constructs with N- and/or C-terminal tags would thus be beneficial to the entire community, setting the foundation for all other *in vitro* studies. It would also be interesting to see if only C-terminally tagged Cdc42 expresses and if it is functional. Such a construct would avoid all possible issues of the N-terminus and can be a great choice if post-translational modifications are not needed.

To make working with Cdc42, and any protein in general, more accessible for a broader spectrum of scientists, it will also be hugely beneficial if publications state and explain the used construct designs and their effects on the protein behaviour, as well as show data of constructs and experiments that "failed".

Materials and Methods

Plasmid construction

Genes of interest were obtained from the genome of *Saccharomyces cerevisiae* W303, or in the case of mNeonGreen and sfGFP from plasmids, and were amplified through PCR. The target vector was also amplified through PCR. Additionally, each PCR incorporated a small homologous sequences needed for Gibson assembly [45]. After Gibson assembly, the resulting mixture was used to transform chemically competent Dh5α and BL21 DE::3 pLysS cells and plated out onto a Petri dish containing Lysogeny broth agar and the correct antibiotic marker. The primers used for each PCR can be found in Supplement S6 Tab. 1. Gibson assembly resulted in plasmids found Tab. 2 (of which protein sequences are stated in Supplement S5) and Tab. 3.

Buffer composition

If not mentioned otherwise, the buffers are of the composition stated in Tab. 4.

Protein expression tests

Proteins were expressed in Bl21::DE3 pLysS cells, which carry the gene for the bacteriophage T7 RNA polymerase under the regulation of a Lactose dependent promoter. Expression of the T7 RNA polymerase, which for example can be induced by IPTG, results in the transcription and therefore expression of the genes of interest that are placed under the T7 promoter. This system is of advantage, as the T7 RNA polymerase transcribes 5-10× faster than E. coli RNA polymerase [31, 47]. Four conditions were used:

1. Cells were grown in Lysogeny broth at 37°C until an OD$_{600}$ of 0.7, the expression was induced through addition of $1.0\,\mathrm{mM}$ IPTG, after which cells were grown for 3 h at 37°C.
2. Cells were grown in Lysogeny broth at 37°C until an OD$_{600}$ of 0.7, the expression was induced through addition of $0.2\,\mathrm{mM}$ IPTG, after which cells were grown for 18 h at 18°C.
3. Cells were grown in Studier Induction ZYP-5052 Medium for 3 h at 37°C, followed by 18 h at 18°C, and in accordance to the recommended protocol [39].

Cells were harvested through centrifugation. Cell pellets were resuspended in SDS loading buffer (Laemmli buffer, [48] and expression levels were analysed through SDS-Page and Western blotting.

Protein purification

The expression and purification protocols for Cdc24, Cdc42, and Bem1 described here are based, with modifications, on the Cdc24 purification protocol described previously [23].

Table 3. List of additional protein constructs/plasmids used throughout this publication.

Plasmid	Description	Source
pWKD011a	Cdc42-sfGFP^{SW}	this work
pWKD017	Cdc42-mNeonGreen^{SW}	this work
pET28a-His-mcm10-Sortase-Flag	plasmid template for N-term. 6His- and C-term. Flag-tagged proteins	received from N. Dekker (TU Delft), based on pBP6 [46]
pSF1818	TEV protease, referred to as TEV (3) in this publication	purified protein received from C. Dekker (TU Delft)

Table 4. Buffer composition.

Buffer	Composition
Lysis buffer	5 mM Tris-HCl (pH=8.0), 1 M NaCl, 5 mM imidazole, 1 mM 2-mercaptoethanol (Sigma Aldrich), supplemented with EDTA-free Protease inhibitor cocktail (Roche) and 1 mM freshly prepared PMSF.
His-AC washing buffer	50 mM Tris-HCl (pH=8.0), 1 M NaCl, 5 mM imidazole, 1 mM 2-mercaptoethanol (Sigma Aldrich).
His-AC elution buffer	50 mM Tris-HCl (pH=8.0), 100 mM NaCl, 500 mM imidazole, 1mM 2-mercaptoethanol (Sigma Aldrich).
Strep-AC washing buffer	50 mM Tris-HCl (pH=8.0), 100 mM NaCl, 1mM 2-mercaptoethanol (Sigma Aldrich).
Strep-AC elution buffer	50 mM Tris-HCl (pH=8.0), 100 mM NaCl, 10mM desthiobiotin (Sigma Aldrich), 1mM 2-mercaptoethanol (Sigma Aldrich).
SEC buffer	50 mM Tris-HCl (pH=7.5), 100 mM NaCl, 10 mM $MgCl_2$, 1 mM 2-mercaptoethanol (Sigma Aldrich).

The methods used are described in the following and protein-specific purification routes are stated thereafter.

Protein expression

The proteins were expressed in BI21::DE3 pLysS cells. Cells were grown in Lysogeny broth at 37°C until an OD_{600} of 0.7. One of these two induction conditions was chosen:

1. Cells were grown in Lysogeny broth at 37°C until an OD_{600} of 0.7, the expression was induced through addition of 1.0 mM IPTG, after which cells were grown for 3 h at 37°C.
2. Cells were grown in Lysogeny broth at 37°C until an OD_{600} of 0.7, the expression was induced through addition of 0.2 mM IPTG, after which cells were grown for 18 h at 18°C.

Cells were harvested through centrifugation.

Lysis and His affinity chromatography (His-AC)

Cell pellets were resuspended in lysis buffer and lysed with a high pressure homogenizer (French press cell disruptor, CF1 series Constant Systems) at 4°C, using 5-10 rounds of exposing the sample to pressurisation. The cell lysate was centrifuged at 37000× g for 30 min and the supernatant was loaded onto a HisTrap™ excel column (Cytiva)[1]. After several rounds of washing with His-AC washing buffer, the protein was eluted in a gradient of His-AC washing buffer and His-AC elution buffer.

Strep affinity chromatography (Strep-AC)

For some proteins Strep-AC was chosen for further purification. The His-AC elution peak was loaded onto a StrepTrap™ HP column (Cytiva). After several rounds of washing with Strep-AC washing buffer, the protein was eluted in a gradient of Strep-AC washing buffer and Strep-AC elution buffer.

Size exclusion chromatography (SEC)

Some proteins were further purified by SEC using SEC buffer and a HiPrep 16/60 Sephacryl S-300 HR (Cytiva) column.

Dialysis and storage

All proteins were dialysed twice in SEC buffer. After the addition of 10% glycerol, samples were flash frozen in liquid nitrogen and kept at -80°C for storage.

Expression and purification conditions per protein

H-Cdc42-F, F-Cdc42-H, H-SS-Cdc42-F, S-Cdc42-H, S-Cdc42-BC-H, H-Cdc42-SS, H-Cdc42:CTIS, H*-Cdc42:CTIS, and F-Cdc42-sfGFP-H were expressed using expression condition (1) and purified in a single step His-AC using an AKTA Pure (Cytiva) machine. The purification profile of H-SS-Cdc42-F, as an example for Cdc42 purification, is shown in Fig. 3a and that of F-Cdc42-sfGFP-H in

---

[1] Newer columns (bought 2020 or later) required a higher amount of imidazole in the lysis and washing buffer, as indicated by the recommendation 'use 20-40 mM imidazole in sample and binding buffer for highest purity' on the column package. For these columns the amount of imidazole in lysis and His-AC washing buffer was increased to 50 mM.

Figure 11. SDS-Page of purified Cdc24, Bem1, and fluorescent Cdc42. Purified Cdc42 is shown in Fig. 3b.

Fig. 5.

H-Cdc42-sfGFP-SS was expressed using expression condition (1) and purified using His-AC followed by Strep-AC. Here lysis buffer, His-AC washing buffer, and His-AC elution buffer were supplemented with 0.1 v/v% Tween-20 (Sigma Aldrich), 0.1 v/v% NP-40 (Thermo Fischer Scientific), and 0.1 v/v% Triton-X-100 (Sigma Aldrich). The purifications were done using an AKTA Pure (Cytiva) machine. Purification profiles are shown in Supplement S2 Fig. 1.

F-Cdc42-mNeon-H was expressed using expression condition (1) and purified using His-AC followed by SEC using an AKTA Pure (Cytiva) machine. Purification profiles are shown in Supplement S2 Fig. 2.

S-Cdc42-mNeon-BC-H was expressed using expression condition (1) and purified using His-AC followed by SEC. Here lysis buffer, His-AC washing buffer, and SEC buffer were supplemented with 0.1 v/v% Tween-20 (Sigma Aldrich), 0.1 v/v% NP-40 (Thermo Fischer Scientific), and 0.1 v/v% Triton-X-100 (Sigma Aldrich). The purifications were done using an AKTA Pure (Cytiva) machine. Purification profiles are shown in Supplement S2 Fig. 3.

Cdc24-H was expressed using expression condition (2) and purified using His-AC followed by SEC using an AKTA Pure (Cytiva) machine.

H-Bem1-F was expressed using expression condition (2) and purified in a single-step His-AC using an AKTA Pure (Cytiva) machine. Fig. 3b and 11 shows all purified proteins on SDS-Page.

GTPase activity assay

GTPase activity was measured using the GTPase-Glo^{TM} assay (Promega) as described previously [49]. In brief, 5 µL protein in SEC buffer (Tab. 4) was mixed with 5 µL of a GTP-solution (10 µM GTP, 50 mM Tris-HCl (pH=7.5), 100 mM NaCl, 10 mM MgCl$_2$, 1 mM 2-mercaptoethanol (Sigma Aldrich), 1 mM dithiothreitol (VWR) in 384-well plates (Corning) to initiate the reaction. The reaction mixture got incubated at 30°C on an Innova 2300 platform shaker (New Brunswick Scientific) (120 rpm), before the addition of 10 µL Glo buffer and another 30 min incubation. The Glo buffer contains a nucleoside-diphosphate kinase that convert remaining GTP to ATP. Addition of 20 µL detection reagent, containing a luciferase/luciferin mixture, makes the ATP luminescent, which was read on a Synergy HTX plate reader (BioTek) in luminescence mode. The amount of hydrolysed GTP inversely correlates with the measured luminescence. Wells without protein ('buffer') were used for the normalization and represent 0% GTP hydrolysis (Eq. 3). Reactions were carried out with at least 4 replicates (wells) per assay, and the average ('Lum.') and standard deviation ('ΔLum.') of each set was used to calculate the activity and error of each set.

$$\text{hydrolysed GTP} = 1 - \text{remaining GTP} = \left(1 - \frac{\text{Lum. protein}}{\text{Lum. buffer}}\right) \times 100\% \tag{3}$$

Error bars were calculated by using error propagation:

$$\Delta\text{hydrolysed GTP} = \sqrt{\left(\frac{\Delta\text{Lum. protein}}{\text{Lum. protein}}\right)^2 + \left(\frac{\Delta\text{Lum. buffer}}{\text{Lum. buffer}}\right)^2} \times \frac{\text{Lum. protein}}{\text{Lum. buffer}} \times 100\% \tag{4}$$

The amount of remaining GTP at the time of reaction termination ($t_{term.}$) was calculated by

$$[GTP]_{t_{term.}} = \left(\frac{\text{Lum. protein}}{\text{Lum. buffer}} \right) \times 100\% \tag{5}$$

We developed a Cdc42 GTPase activity model for determining the GTPase cycling rates k. It is described in the following:
The GTP hydrolysis process involves three steps: (1) A GTP molecule from solution binds to Cdc42. (2) Cdc42 hydrolyses GTP. (3) Cdc42 releases GDP.

$$[GTP] + [Cdc42] \rightleftharpoons [Cdc42-GTP]$$

$$[Cdc42-GTP] \longrightarrow [Cdc42-GDP]$$

$$[Cdc42-GDP] \rightleftharpoons [Cdc42] + [GDP]$$

It can further be upregulated by effector proteins: GAPs have been shown to enhance GTP hydrolysis by Cdc42 (step 2), GEFs enhance the release of GDP from Cdc42 (step 3) [50, 51] and the scaffold Bem1 enhances the GEF activity of Cdc24 [23].
To quantitatively describe the GTPase reaction cycle, we coarse-grained the GTPase reaction steps with

$$[GTP] + n\,[Cdc42] \longrightarrow [GDP] + n\,[Cdc42]$$

To account for possible Cdc42 dimerisation and cooperativity, we included the following reactions into the model:
(1) We assume that Cdc42 can dimerise, as other small GTPases have been shown to dimerise [17, 18, 52, 40]:

$$2\,[mCdc42] \rightleftharpoons [dCdc42]$$

and both monomeric and dimeric Cdc42 can contribute to the overall GTP hydrolysis with different rates:

$$[GTP] + [mCdc42] \xrightarrow{k_1'} [GDP] + [mCdc42]$$

$$[GTP] + [dCdc42] \xrightarrow{k_2'} [GDP] + [dCdc42]$$

Assuming that the majority of Cdc42 is in its monomeric form ($[mCdc42] < C_d$, with C_d as the concentration at which half of the total Cdc42 is dimeric), we can approximate

$$[dCdc42] = \frac{[mCdc42]^2}{2C_d}$$

$$[mCdc42] \approx [Cdc42] - \frac{[Cdc42]^2}{C_d} \tag{6}$$

(2) Next to cooperativity from dimerisation, cooperativity can also emerge when Cdc42 proteins come in close contact with each other - they can affect each other's behaviour without forming a stable homodimer, effectively functioning as an effector protein for themselves:

$$[GTP] + 2\,[mCdc42] \xrightarrow{k_3'} [GDP] + 2\,[mCdc42]$$

$$[GTP] + [mCdc42] + [dCdc42] \xrightarrow{k_4'} [GDP] + [mCdc42] + [dCdc42]$$

(3) Effector proteins, such as GAPs and GEFs, affect the speed of the GTP hydrolysis cycle:

$$[GTP] + [Cdc42][X] \xrightarrow{k_5'} [GDP] + [Cdc42][X]$$

Here X is an effector protein.

Our data showed that the amount of remaining GTP follows an exponential decline over time:

$$[GTP]_t = [GTP]_{t_0} \exp(-Kt)\,, \text{ using } [GTP]_{t_0} = 100\% \tag{7}$$

Considering reactions (1) - (3), we can thus define K in Eq. 7 as

$$K = k_1'[mCdc42] + k_2'[dCdc42] + k_3'[mCdc42]^2 + k_4'[mCdc42][dCdc42] + k_5'[Cdc42][X] \tag{8}$$

Using Eq. 6, and considering only up to second-order terms, results in

$$K = k_1'[\text{Cdc42}] + \left(\frac{k_2'}{2C_d} + k_3' - \frac{k_1'}{C_d} \right) [\text{Cdc42}]^2 + k_5'[\text{Cdc42}][X]$$

$$= k_1[\text{Cdc42}] + k_2[\text{Cdc42}]^2 + k_{3,X}[\text{Cdc42}][X]$$

(9)

where k_1 refers to GTP hydrolysis cycling rates of monomeric Cdc42, k_2 includes effects of cooperativity and dimerisation and k_3 represents the rate of Cdc42 - effector interaction. We will refer to K as 'overall GTP hydrolysis rate' in the following.

We used Eq. 9 with [X]=0 to determine the rates of Cdc42 alone. We then conducted assays with Cdc42 and an effector protein to determine k_3. While doing so we needed to account for assay variability, i.e. for the observation that the rates for Cdc42 can vary between assays. Possible reasons for this include small concentration differences introduced though pipetting of small volumes (as are required for this assay), temperature and shaker speed fluctuations during the incubation step, and/or intrinsic changes in the protein activities due to other external conditions. To account for this variance, we introduced the parameter c_{corr}. It maps all factors that lead to variations between assays onto the Cdc42 concentration.
The assay data, including samples containing only Cdc42 and Cdc42 - (effector) protein mixtures, was fitted with

$$K = k_1 c_{corr}[\text{Cdc42}] + k_2 (c_{corr}([\text{Cdc42}])^2 + k_{3,X} c_{corr}[\text{Cdc42}][X]$$

(10)

to determine c_{corr} and $k_{3,X}$ (using k_1 and k_2 determined earlier).

Our data suggested that the GTP decline in assays Cdc42-Cdc24 mixtures does not depend linearly on the Cdc24 concentration (Eq. 10), but is better approximated by a quadratic term:

$$K = k_1 c_{corr}[\text{Cdc42}] + k_2 (c_{corr}([\text{Cdc42}])^2 + k_{3,Cdc24} c_{corr}[\text{Cdc42}][\text{Cdc24}]^2$$

(11)

The GTPase assays conducted here involve many pipetting steps and can, in done without a lot of attention, be prone to errors. These result in assays with lots of variability and big error bars. Most GTPase assays resulted in data with small error bars. We used the following criteria to exclude assays from our analysis:

- The error of buffer wells (ΔLum. buffer) is above 10%. This generally means that the assay and all assay data has big error bars.
- The variability of the Cdc42 activity between assays using only Cdc42 and Cdc42 - Cdc24 mixtures is high, i.e. $> c_{corr} < 0.5$ or $c_{corr} > 1.6$. Assays of only Cdc42 dilutions showed a low variability between assays. In Cdc42 - Cdc24 assays only one Cdc42 concentration was used. If it doesnt show the same GTPase activity as in the assays with only Cdc42, c_{corr} deviates from one. This then affects the other fitting parameters and leads to more variability in k_3. Ideally only data of c_{corr} close to one would be included. As we did not intent to provide a detailed parametrisation of the constructs used, but to use this assay to compare them and to point ot points of failure when working with Cdc42, we included a wider range of data. Most assays, nontheless, had values of c_{corr} close to one (see Supplement S3).

Additionally, data points that showed less than 5% remaining GTP were excluded from the fit, as they correspond to almost no measurable luminescence.

Flag pulldown assay
In this assay, Flag-tagged Bem1 (H-Bem1-F) and another protein, that does not contain a Flag-tag, was mixed. For each experiment, 0.2 nmol Bem1, and 1.0 nmol Cdc42 or 0.6-1.4 nmol Ovalbumin (as a neg. control, Gel Filtration HMW Calibration Kit (Cytiva)) were used. If Cdc42 was used, 1.0 nmol got first pre-loaded with either no nucleotide, 100 nmol GDP (Jena Bioscience), or 100 nmol GTPγS (Jena Bioscience) for 30 min at room temperature. Two proteins were incubated for 1 h at 30 °C. 100 µL anti-Flag® M2 Affinity gel (Sigma Aldrich) was added and incubated for another 30 min at room temperature. The reaction mixture was added onto a Poly-Prep Chromatography column (Biorad) and was washed 3× with 1 mL of assay buffer (25 mM Tris-HCl (pH=7.2), 300 mM NaCl, 10% Glycerol, 0.01 v/v% NP-40 (Thermo Fischer Scientific); supplemented either with no nucleotide, 10-100 µM GDP, or 10-100 µM GTPγS). Proteins were eluted with 200 µL assay buffer supplemented with 0.6 mg/mL 3× Flag® peptide (Sigma Aldrich), and analysed by Western blotting.

Blot	Primary antibody	Secondary antibody
anti-His	His Tag Antibody, Mouse (OAEA00010, Aviva Systems Biology) (dilution: 1:4000)	IgG2b Antibody HRP conjugated (Goat Anti-Mouse) (OASA06620, Aviva Systems Biology) (dilution: 1:1000)
anti-Cdc42	Anti-CDC42 antibody, Rabbit (ab64533, abcam) (dilution: 1:2000)	Polyclonal Goat Anti-Rabbit Immunoglobulins/HRP (Goat Anti-Rabbit) (P0448, Agilent Dako) (dilution: 1:500)

Table 5. Antibodies used for Western blotting.

Reactions with enterokinase and TEV protease

For enterokinase screening reactions, 50 µg H-Cdc42:CTIS was mixed with 5, 10, or 20 U of enterokinase (16 U/µL) (P8070, New England Biolabs) in SEC buffer, supplemented with 2 mM $CaCl_2$. After 1-3 h at room temperature or 24-48 h at 4°C reaction samples were taken and analysed by SDS-Page.

For TEV protease screening reactions, tree enzymes were assessed: TEV (1): 50 µg H*-Cdc42:CTIS was mixed with 5 or 50 U of TEV protease (10 U/µL) (T4455, Sigma-Aldrich) in SEC buffer. TEV (2): 50 µg H*-Cdc42:CTIS was mixed with 5 or 50 U of AcTEV^{TM} protease (10 U/µL) (Invitrogen) in SEC buffer. TEV (3): 50 µL H*-Cdc42:CTIS (63 µM) was mixed with 1, 5, or 10 µL of purified TEV protease (125 µM, pSF1818) in SEC buffer. The enzyme was received as a gift from C. Dekker lab (TU Delft). After 1-3 h at room temperature or 24-48 h at 4°C reaction samples were taken and analysed by SDS-Page.

For cleavage reactions, H*-Cdc42:CTIS (63 µM) was mixed with either TEV (1) (TEV protease, 10 U/µL, T4455, Sigma-Aldrich) in a 10 U/100 µg TEV/Cdc42 ratio, or with TEV (3) (125 µM, pSF1818) in a 5 µL/100 µL TEV/Cdc42 ratio, in SEC buffer. After an incubation of 45 h at 4°C, the reaction was stopped. The sample was loaded thrice onto a HisTrap^{TM} excel column (Cytiva). The flow-through contains Cdc42 of which the N-terminal 6His-tag got cleaved. Samples were analysed by SDS-Page and Western blotting.

SDS-Page

SDS-Page gels (12-15% acrylamide) were prepared freshly. In brief, a solution of 375 mM Tris-HCl (pH=8.8), 30-37.5 v/v% 40% acrylamide solution (Biorad), 0.2 w/v% sodium dodecyl sulfate (Sigma Aldrich), 0.5 v/v% 2,2,2-Trichloroethanol (Sigma Aldrich), 0.1 w/v% ammonium persulfate (Sigma Aldrich), and 0.1 v/v% N,N,N',N'-tetramethyl ethylenediamine (Sigma Aldrich) was prepared and casted into 1.00 mm mini-protean glass plates (Biorad), filling them up to 80%. To protect the gel surface from drying, a layer of isopropanol (Sigma Aldrich) was added. The gel was let solidify for 20 min, after which the isopropanol layer was removed. A solution of 155 mM Tris-HCl (pH=6.5), 10 v/v% 40% acrylamide solution (Biorad), 0.2 w/v% sodium dodecyl sulfate (Sigma Aldrich), 0.1 w/v% ammonium persulfate (Sigma Aldrich), and 0.1 v/v% N,N,N',N'-tetramethyl ethylenediamine (Sigma Aldrich) was prepared and added to the existing gel layer, after which a well comb (Biorad) was added. The gel was let solidify for 20 min.

Cell and protein samples were mixed with SDS loading buffer (Laemmli buffer, [48]. Before loading onto the SDS-Page gels, they were kept for 5 min at 95°C. Gels were run for 5 min at 130 V followed by 55min min at 180 V (PowerPac Basic Power Supply (Biorad)). Imaging was done on a ChemiDoc MP (Biorad) using the 'Stain-free gels' feature and automatic exposure time determination. Precision Plus Protein Unstained standard (Biorad) was used as a protein standard.

Western blotting

After SDS-Page, the sample was transferred from the SDS-Page to a blotting membrane (Trans-Blot Turbo Transfer Pack, Bio-rad) using the 'Mixed MW' program of the Trans-Blot Turbo Transfer System (Bio-rad). The blotting membrane was incubated with Immobilon signal enhancer (Millipore) at 4°C for 18 h. The blotting membrane was incubated with primary antibody, diluted in Immobilon signal enhancer (see Tab. 5), at room temperature for 1 h. It was washed thrice with TBS-T (10 mM Tris-HCl (pH=7.5), 150 mM NaCl, 0.1 v/v% Tween-20 (Sigma Aldrich)). For each washing step the blotting membrane was incubated with TBS-T at room temperature for 20 min. The blotting membrane was incubated with primary antibody, diluted in Immobilon signal enhancer (see Tab. 5), at room temperature for 1 h, after which it was again washed thrice with TBS-T. SuperSignal West Pico Mouse IgG Detection Kit (Thermo Scientific) was used for activation. Imaging was done on a ChemiDoc MP (Biorad) using the 'Chemi Sensitive' feature and automatic exposure time determination. Precision Plus Protein Unstained standard (Biorad) and Precision Plus Protein All Blue Standard (Biorad) were used as protein standards.

Acknowledgements and contributions

R. van der Valk constructed the plasmids of protein constructs. Proteins were expressed and purified by S. Tschirpke and F. van Opstal. S. Tschirpke conducted GTPase assays, Flag-pulldown experiments, expression tests, Western blots, and analysed the data thereof. The modelling and fitting of the GTPase data was done by W. Daalman. F. van Opstal conducted expression tests and Western blots. Figure preparation and writing of this publication was done by S. Tschirpke.

We thank C. de Agrela Pinto for conducting SEC-MALS with our samples and analysing the data thereof. We thank D. McCusker (University of Bordeaux) for the plasmid pDM272, N. Dekker (TU Delft) for the plasmid pET28a-His-mcm10-Sortase-Flag, and C. Dekker lab (TU Delft) for a sample of purified TEV protease (pSF1818).

References

 [1] Bieling P, Li TD, Weichsel J, McGorty R, Jreij P, Huang B, et al. Force Feedback Controls Motor Activity and Mechanical Properties of Self-Assembling Branched Actin Networks. Cell. 2016;164(1-2):115–127. doi:10.1016/j.cell.2015.11.057.

 [2] Roth S, Gârlea IC, Vleugel M, Mulder BM, Dogterom M. Reconstitution of basic mitotic spindles in cell-like confinement. bioRxiv. 2019; p. 770602.

 [3] Bezeljak U, Loya H, Kaczmarek B, Saunders TE, Loose M. Stochastic activation and bistability in a Rab GTPase regulatory network. Proceedings of the National Academy of Sciences. 2020;117(12):6540–6549. doi:10.1073/pnas.1921027117.

 [4] Kohyama S, Merino-Salomón A, Schwille P. In vitro assembly, positioning and contraction of a division ring in minimal cells. Nature Communications. 2022;13(1). doi:10.1038/s41467-022-33679-x.

 [5] Vendel KJA, Tschirpke S, Shamsi F, Dogterom M, Laan L. Minimal in vitro systems shed light on cell polarity. Journal of Cell Science. 2019;132(4):1–21. doi:10.1242/jcs.217554.

 [6] Diepeveen ET, Gehrmann T, Pourquié V, Abeel T, Laan L. Patterns of conservation and diversification in the fungal polarization network. Genome Biology and Evolution. 2018;10(August):evy121–evy121. doi:10.1093/gbe/evy121.

 [7] Gao JT, Guimera R, Li H, Pinto IM, Sales-Pardo M, Wai SC, et al. Modular coherence of protein dynamics in yeast cell polarity system. Proceedings of the National Academy of Sciences. 2011;108(18):7647–7652. doi:10.1073/pnas.1017567108.

 [8] Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, Tan G, et al. A global genetic interaction network maps a wiring diagram of cellular function. Science. 2016;353(6306):aaf1420–aaf1420. doi:10.1126/science.aaf1420.

 [9] Daalman WKG, Sweep E, Laan L. The Path towards Predicting Evolution as Illustrated in Yeast Cell Polarity. Cells. 2020;9(12). doi:10.3390/cells9122534.

[10] Swaney DL, Beltrao P, Starita L, Guo A, Rush J, Fields S, et al. Global analysis of phosphorylation and ubiquitylation cross-talk in protein degradation. Nature Methods. 2013;10(7):676–682. doi:10.1038/nmeth.2519.

[11] Back S, Gorman AW, Vogel C, Silva GM. Site-Specific K63 Ubiquitinomics Provides Insights into Translation Regulation under Stress. Journal of Proteome Research. 2019;18(1):309–318. doi:10.1021/acs.jproteome.8b00623.

[12] Lanz MC, Yugandhar K, Gupta S, Sanford EJ, Faça VM, Vega S, et al. In-depth and 3-dimensional exploration of the budding yeast phosphoproteome. EMBO reports. 2021;22(2). doi:10.15252/embr.202051121.

[13] Caplin BE, Hettich LA, Marshall MS. Substrate characterization of the saccharomyces cerevisiae protein farnesyltransferase and type-I protein geranylgeranyltransferase. Biochimica et Biophysica Acta (BBA)/Protein Structure and Molecular. 1994;1205(1):39–48. doi:10.1016/0167-4838(94)90089-2.

[14] Golding AE, Visco I, Bieling P, Bement WM. Extraction of active RhoGTPases by RhoGDI regulates spatiotemporal patterning of RhoGTPases. eLife. 2019;8:1–26. doi:10.7554/eLife.50471.

[15] Zheng Y, Cerione R, Bender A. Control of the yeast bud-site assembly GTPase Cdc42. Catalysis of guanine nucleotide exchange by Cdc24 and stimulation of GTPase activity by Bem3. J Biol Chem. 1994;269(4):2369–2372.

[16] Zheng Y, Bender A, Cerione RA. Interactions among proteins involved in bud-site selection and bud-site assembly in Saccharomyces cerevisiae. Journal of Biological Chemistry. 1995;270(2):626–630.

[17] Zhang B, Zheng Y. Negative regulation of Rho family GTPases Cdc42 and Rac2 by homodimer formation. Journal of Biological Chemistry. 1998;273(40):25728–25733. doi:10.1074/jbc.273.40.25728.

[18] Zhang B, Zhang Y, Collins CC, Johnson DI, Zheng Y. A built-in arginine finger triggers the self-stimulatory GTPase- activating activity of Rho family GTPases. Journal of Biological Chemistry. 1999;274(5):2609–2612. doi:10.1074/jbc.274.5.2609.

[19] Kozminski KG, Beven L, Angerman E, Tong AHY, Boone C, Park HO. Interaction between a Ras and a Rho GTPase Couples Selection of a Growth Site to the Development of Cell Polarity in Yeast. Molecular Biology of the Cell. 2003;14(12):4958–4970. doi:10.1091/mbc.E03.

[20] Johnson JL, Erickson JW, Cerione RA. New Insights into How the Rho Guanine Nucleotide Dissociation Inhibitor Regulates the Interaction of Cdc42 with Membranes. Journal of Biological Chemistry. 2009;284(35):23860–23871. doi:10.1074/jbc.M109.031815.

[21] Johnson JL, Erickson JW, Cerione RA. C-terminal Di-arginine motif of Cdc42 protein is essential for binding to phosphatidylinositol 4,5-bisphosphate-containing membranes and inducing cellular transformation. Journal of Biological Chemistry. 2012;287(8):5764–5774. doi:10.1074/jbc.M111.336487.

[22] Das A, Slaughter BD, Unruh JR, Bradford WD, Alexander R, Rubinstein B, et al. Flippase-mediated phospholipid asymmetry promotes fast Cdc42 recycling in dynamic maintenance of cell polarity. Nature Cell Biology. 2012;14(3):304–310. doi:10.1038/ncb2444.

[23] Rapali P, Mitteau R, Braun C, Massoni-Laporte A, Ünlü C, Bataille L, et al. Scaffold-mediated gating of Cdc42 signalling flux. eLife. 2017;6:1–18. doi:10.7554/eLife.25257.

[24] Cox AD, Der CJ. Protein prenylation: more than just glue? Current Opinion in Cell Biology. 1992;4(6):1008–1016. doi:10.1016/0955-0674(92)90133-W.

[25] Koch G, Tanaka K, Masuda T, Yamochi W, Nonaka H, Takai Y. Association of the Rho family small GTP-binding proteins with Rho GDP dissociation inhibitor (Rho GDI) in Saccharomyces cerevisiae. Oncogene. 1997;15(4):417–422. doi:10.1038/sj.onc.1201194.

[26] Popp MWL, Ploegh HL. Making and breaking peptide bonds: Protein engineering using sortase. Angewandte Chemie - International Edition. 2011;50(22):5024–5032. doi:10.1002/anie.201008267.

[27] Williams CL. The polybasic region of Ras and Rho family small GTPases: a regulator of protein interactions and membrane association and a site of nuclear localization signal sequences. Cellular Signalling. 2003;15(12):1071–1080. doi:10.1016/S0898-6568(03)00098-6.

[28] Zhang B, Zhang Y, Wang ZX, Zheng Y. The role of Mg2+ cofactor in the guanine nucleotide exchange and GTP hydrolysis reactions of Rho family GTP-binding proteins. Journal of Biological Chemistry. 2000;275(33):25299–25307. doi:10.1074/jbc.M001027200.

[29] Sonal LK, Heermann T, Schwille P. One-pot synthesis of prenylated proteins utilizing E. coli cell-free expression. BioRxiv. 2022;.

[30] Bose I, Irazoqui JE, Moskow JJ, Bardes ES, Zyla TR, Lew DJ. Assembly of scaffold-mediated complexes containing Cdc42p, the exchange factor Cdc24p, and the effector Cla4p required for cell cycle-regulated phosphorylation of Cdc24p. J Biol Chem. 2001;276(10):7176–7186. doi:10.1074/jbc.M010546200.

[31] Studier FW, Moffatt BA. Use of bacteriophage T7 RNA polymerase to direct selective high-level expression of cloned genes. Journal of Molecular Biology. 1986;189(1):113–130. doi:10.1016/0022-2836(86)90385-2.

[32] Meca J, Massoni-Laporte A, Martinez D, Sartorel E, Loquet A, Habenstein B, et al. Avidity-driven polarity establishment via multivalent lipid– GTPase module interactions. The EMBO Journal. 2019;38(3):1–19. doi:10.15252/embj.201899652.

[33] Bendezú FO, Vincenzetti V, Vavylonis D, Wyss R, Vogel H, Martin SG. Spontaneous Cdc42 Polarization Independent of GDI-Mediated Extraction and Actin-Based Trafficking. PLoS Biology. 2015;13(4):1–30. doi:10.1371/journal.pbio.1002097.

[34] Schmidt TGM, Batz L, Bonet L, Carl U, Holzapfel G, Kiem K, et al. Development of the Twin-Strep-tag® and its application for purification of recombinant proteins from cell culture supernatants. Protein Expression and Purification. 2013;92(1):54–61. doi:10.1016/j.pep.2013.08.021.

[35] Hopp TP, Prickett KS, Price VL, Libby RT, March CJ, Pat Cerretti D, et al. A Short Polypeptide Marker Sequence Useful for Recombinant Protein Identification and Purification. Bio/Technology. 1988;6(10):1204–1210. doi:10.1038/nbt1088-1204.

[36] Sartorel E, Unlu C, Jose M, Aurélie ML, Meca J, Sibarita JB, et al. Phosphatidylserine and GTPase activation control Cdc42 nanoclustering to counter dissipative diffusion. Molecular Biology of the Cell. 2018;29(11):1299–1310. doi:10.1091/mbc.E18-01-0051.

[37] Pédelacq JD, Cabantous S, Tran T, Terwilliger TC, Waldo GS. Engineering and characterization of a superfolder green fluorescent protein. Nature Biotechnology. 2006;24(1):79–88. doi:10.1038/nbt1172.

[38] Shaner NC, Lambert GG, Chammas A, Ni Y, Cranfill PJ, Baird MA, et al. A bright monomeric green fluorescent protein derived from Branchiostoma lanceolatum. Nature Methods. 2013;10(5):407–409. doi:10.1038/nmeth.2413.

[39] Studier FW. Protein production by auto-induction in high density shaking cultures. Protein expression and purification. 2005;41(1):207–234. doi:10.1016/j.pep.2005.01.016.

[40] Kang PJ, Béven L, Hariharan S, Park HO. The Rsr1/Bud1 GTPase interacts with itself and the Cdc42 GTPase during bud-site selection and polarity establishment in budding yeast. Molecular Biology of the Cell. 2010;21(17):3007–3016. doi:10.1091/mbc.E10-03-0232.

[41] Mionnet C, Bogliolo S, Arkowitz RA. Oligomerization regulates the localization of Cdc24, the Cdc42 activator in Saccharomyces cerevisiae. Journal of Biological Chemistry. 2008;283(25):17515–17530. doi:10.1074/jbc.M800305200.

[42] Shimada Y, Wiget P, Gulli MP, Bi E, Peter M. The nucleotide exchange factor Cdc24p may be regulated by auto-inhibition. EMBO Journal. 2004;23(5):1051–1062. doi:10.1038/sj.emboj.7600124.

[43] Shahravan SH, Qu X, Chan IS, Shin JA. Enhancing the specificity of the enterokinase cleavage reaction to promote efficient cleavage of a fusion tag. Protein Expression and Purification. 2008;59(2):314–319. doi:10.1016/j.pep.2008.02.015.Enhancing.

[44] Light A, Savithri HS, Liepnieks JJ. Specificity of bovine enterokinase toward protein substrates. Analytical Biochemistry. 1980;106(1):199–206. doi:10.1016/0003-2697(80)90138-4.

[45] Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, Smith HO. Enzymatic assembly of DNA molecules up to several hundred kilobases. Nature Methods. 2009;6(5):343–345. doi:10.1038/nmeth.1318.

[46] Douglas ME, Diffley JFX. Recruitment of Mcm10 to sites of replication initiation requires direct binding to the minichromosome maintenance (MCM) complex. Journal of Biological Chemistry. 2016;291(11):5879–5888. doi:10.1074/jbc.M115.707802.

[47] Dubendorf JW, Studier FW. Controlling basal expression in an inducible T7 expression system by blocking the target T7 promoter with lac repressor. Journal of Molecular Biology. 1991;219(1):45–59. doi:10.1016/0022-2836(91)90856-2.

[48] Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. Nature. 1970;227:680–685.

[49] Mondal S, Hsiao K, Goueli SA. A Homogenous Bioluminescent System for Measuring GTPase, GTPase Activating Protein, and Guanine Nucleotide Exchange Factor Activities. Assay and Drug Development Technologies. 2015;13(8):444–455. doi:10.1089/adt.2015.643.

[50] Martin SG.   Spontaneous cell polarization: Feedback control of Cdc42 GTPase breaks cellular symmetry.   BioEssays. 2015;37(11):1193–1201. doi:10.1002/bies.201500077.

[51] Chiou Jg, Balasubramanian MK, Lew DJ. Cell Polarity in Yeast. Annual Review of Cell and Developmental Biology. 2017;33:77– 101. doi:10.1146/annurev.cellbio.15.1.365.

[52] Zhang B, Gao Y, Moon SY, Zhang Y, Zheng Y. Oligomerization of Rac1 GTPase Mediated by the Carboxyl-terminal Polybasic Domain. Journal of Biological Chemistry. 2001;276(12):8958–8967. doi:10.1074/jbc.M008720200.

Supporting information captions

S1 Table. Relationship between polybasic regions and protein dimerisation.

S2 Appendix. Purification of Cdc42-sfGFP^{SW} and -mNeonGreen^{SW}.

S3 Figures and Tables. GTPase activity data and fits for Cdc42 constructs and summary graphs of Cdc42's GTPase activity and interaction with Cdc24.

S4 Figures. TEV protease, but not enterokinase, is a suitable cleavage enzyme for Cdc42.

S5 Appendix. Amino acid sequences of proteins.

S6 Table. Primer overview.

Supplement S1

S1 Table 1. Relationship between polybasic regions and protein dimerisation. Amino acid (AA) sequences of the C-terminal region of selected small GTPases in relation to the ability of these GTPases to dimerise/ oligomerise. The PBR and other basic AA (K: lysine, R: arginine) are highlighted in bold. Abbreviations: *S.c.*: *S. cerevisia*, *H.s.*: *H. sapiens*, WT: wild-type, M: mutant.

Protein	Type	C-term. AA sequ.	State	Exp. condition	Reference
S.c. Cdc42	WT	PVI **KKSKK** CAIL	unknown	*in vitro*	
			monomer	*in vivo*	[40]
H.s. Cdc42	WT	PEP **KKSRR** CVLL	dimer	*in vitro*	[17]
H.s. Cdc42	M	PEP **KKSKR** CVLL	dimer	*in vitro*	[18]
H.s. Cdc42	M	PEP **KK**	monomer	*in vitro*	[17]
H.s. Rac1	WT	PV**K KRKRK** CLLL	oligomer	*in vitro*	[52]
H.s. Rac1	M	PVP **KKSRR** CVLL	oligomer	*in vitro*	[52]
H.s. Rac1	M	PV**K QQQQQ** CVLL	monomer	*in vitro*	[52]
H.s. Rac1	M	PV**K**	monomer	*in vitro*	[52]
H.s. Rac2	WT	PT**R QQKRA** CSLL	dimer	*in vitro*	[17]
H.s. RhoA	WT	**RR**G **KKKSG** CLVL	dimer	*in vitro*	[17]
H.s. RhoA	M	**RR**G **KKKRG** CLVL	dimer	*in vitro*	[18]
H.s. RhoA	M	**RR**G **K**	monomer	*in vitro*	[17]
S.c. Rsr1	WT	SQQ **KKKKK** NASTCTIL	dimer	*in vivo*	[40]
S.c. Rsr1	M	SQQ **SSSSS** NASTCTIL	monomer	*in vivo*	[40]

Supplement S2

We purified F-Cdc42-sfGFP-H, H-Cdc42-sfGFP-SS, F-Cdc42-mNeon-H, and S-Cdc42-mNeon-BC-H. F-Cdc42-sfGFP-H could be puri-fied in a high yield (30 mg per 6 L expression) in a single step His-AC (Fig. 5), and the other constructs required further purification and gave a significantly lower yield. An overview of the expression and purification conditions used is given in S2 Tab. 1.

H-Cdc42-sfGFP-SS was purified using His-AC followed by Strep-AC (S2 Fig. 1). The final product was pure, but the yield was low. We observed that the addition of detergents doubled the yield (S2 Tab. 1).

F-Cdc42-mNeon-H was purified using His-AC followed by SEC-AC (S2 Fig. 2). After His-AC F-Cdc42-mNeon-H was not pure, in addition to the 50 kDa protein band a ~70 kDa impurity is visible on SDS-Page (S2 Fig. 2c). The protein was further cleaned by SEC. Here the 50 kDa protein started to elute before a 70 kDa impurity. The peak overlapped strongly with the 70 kDa impurity. Much of the protein fractions therefore had to be discarded, resulting in a low yield.

For S-Cdc42-mNeon-BC-H a similar route was chosen (S2 Fig. 3). After His-AC the elution peak fraction showed also a impurity at ~70 kDa. In SEC the 70 kDa impurity eluted before the 50 kDa S-Cdc42-mNeon-BC-H. We suspect that this time the order changed to the expected one because detergents were added to the buffer. Because the amount of protein in the His-AC elu-tion peak was little, this purification also resulted in a small yield.

Based on this data we suggest to use detergents in the buffers, as this increases the yield during affinity chromatography and could help in SEC. However, the addition of detergents does not lead to the yield obtained for F-Cdc42-sfGFP-H.

In general, if a protein ought to be purified that does not express in a high yield and requires SEC, we advise to significantly scale-up the expression volume and and use large purification columns: We used up to 12 L medium per expression, and puri-fied it in a single His-AC using a 5 mL column (HisTrap™ excel column (Cytiva)). The bigger column volume allows for faster flow rates and reduces the overall purification time, even when large sample volumes are used. Elution using a steep gradient (or even step elution) concentrates all protein into a small volume, which can immediately be used for SEC. If ideal SEC conditions are unknown, a smaller SEC column (e.g. Superdex 200 Increase 10/300 GL (Cytiva), maximum sample volume of 500 µL) can be used to screen conditions. Smaller columns accommodate less sample volume, but have shorter run-times. Once a good condition is found, we advise to switch to a larger SEC column (e.g. HiPrep 16/60 Sephacryl S-300 HR (Cytiva), maximum sample volume of 5 mL), so that the entire, or at least half of the, His-AC elution peak can be loaded onto the SEC column without any in-termediate concentration steps. If the sample after SEC is low in concentration, or samples from several SEC ought to be pooled and concentrated, they can be loaded onto a His-AC column again and eluted in a step elution. This strategy avoids intermediate concentration steps, which, in our experience, reduce the yield up to 30% due to protein sticking to the spin-concentrator.

S2 Table 1. Overview of expression conditions, volumes, purification steps, and final yields of protein for fluorescent Cdc42 constructs. 'X' indicates that the respective purification method was used, (det.) indicates that the buffers were supplemented with a detergent mixture (0.1% Tween-20, 0.1% NP-40, 0.1% Triton-X-100). 'f' refers to a strong and fast expression at elevated temperatures, induced by a high amount of IPTG (3 h at 37°C with 1 mM IPTG).

Construct	Expression condition	Expression volume	His-AC	Strep-AC	SEC	Final yield
F-Cdc42-sfGFP-H	f	6 L	X			~30 mg
H-Cdc42-sfGFP-SS	f	12 L	X	X		~0.5 mg
H-Cdc42-sfGFP-SS	f	12 L	X (det.)	X (det.)		~1 mg
F-Cdc42-mNeon-H	f	6 L	X		X	~0.5 mg
S-Cdc42-mNeon-BC-H	f	6 L	X (det.)		X (det.)	~0.5 mg

S2 Figure 1. Purification profile of H-Cdc42-sfGFP-SS. H-Cdc42-sfGFP-SS was purified by His-affinity chromatography (A) followed by Strep-affinity chromatography (B). (C) SDS-Page of the elution peaks, the band of the full-length protein is indicated by an orange arrow.

S2 Figure 2. Purification profile of F-Cdc42-mNeon-H. F-Cdc42-mNeon-H was purified by His-affinity chromatography (A) followed by size exclusion chromatography (B). (C) SDS-Page of the elution peaks, the band of the full-length protein is indicated by an orange arrow.

S2 Figure 3. Purification profile of S-Cdc42-mNeon-BC-H. S-Cdc42-mNeon-BC-H was purified by His-affinity chromatography (A) followed by size exclusion chromatography (B). (C) SDS-Page of the elution peaks. The band of the full-length protein is indicated by an orange arrow.

◇ data point used for fitting

✕ data point not used for fitting (remaining GTP < 5%)

S3 Figure 1. GTPase activity data and fits for Cdc42 constructs.

S3 Figure 2. Summary graphs of Cdc42's GTPase activity and interaction with Cdc24. (A) GTP hydrolysis cycling rates k_1 and k_2. (B) Predicted amount of remaining GTP for Cdc42 constructs (5 µM) after an incubation time of 1.5 h, calculated using rates k_1, k_2 shown in (A). (C) correction factors c_{corr}. (D) Cdc42-Cdc24 interaction rate k_3. (A-D): Data points of the same colour refer to the rates obtained from separate measurements/assays. Data points of the same colour but a distinct shape (circle, triangle, rhombus) represent different purification batches of the same construct.

S3 Table 1. GTP hydrolysis cycling rates k_1 and k_2 of Cdc42. Purification batches are indicated in brackets.

Cdc42 construct	Experiment number	k_1 [$\times 10^{-2}$ μM^{-1} h^{-1}]	k_1 std. err.	k_2 [$\times 10^{-2}$ μM^{-2} h^{-1}]	k_2 std. err.
H-Cdc42-F (triangle)	E11	5.79	2.79	0.38	0.22
H-Cdc42-F (circle)	E32B	6.23	0.56	4.41	0.15
H-Cdc42-F (circle)	E41C	7.94	1.57	3.27	0.28
H-Cdc42-F (circle)	E41D	10.54	1.68	3.12	0.31
S-Cdc42-H (triangle)	E18	14.80	1.54	0.34	0.37
S-Cdc42-H (circle)	E32B	15.94	2.29	4.26	0.58
S-Cdc42-BC-H	E34A	12.17	4.60	5.21	0.87
H-Cdc42-SS	E32B	14.73	1.52	1.63	0.39
F-Cdc42-H (triangle)	E39A	6.72	1.21	1.46	0.19
F-Cdc42-H (rhombus)	E35A	12.85	0.45	2.09	0.08
F-Cdc42-H (circle)	E32B	6.59	5.13	3.27	1.52
SS-Cdc42-H	E33D	4.03	1.53	-0.39	0.30
H-SS-Cdc42-F (triangle)	E39A	1.58	0.70	0.97	0.10
H-SS-Cdc42-F (rhombus)	E42A	0.00	0.00	2.15	0.05
H-SS-Cdc42-F (circle)	E42A	0.00	0.00	2.45	0.13
H-Cdc42:CTIS	E39A	3.65	3.55	3.52	0.56
H*-Cdc42:CTIS	E42B	1.97	1.54	3.00	0.24
*-Cdc42:CTIS (triangle)	E42B	1.21	0.95	2.27	0.16
*-Cdc42:CTIS (circle)	E42B	0.00	0.00	2.78	0.06

S3 Table 2. Correction factors c_{corr} and Cdc42-Cdc24 interaction rate k_3. Purification batches are indicated in brackets.

Cdc42 construct	Experiment number	c_{corr}	k_3 [$\mu M^{-3}\,h^{-1}$]	k_3 std. err.
SS-Cdc42-H	E33D	1.00	0.10	0.05
H-Cdc42-F (circle)	E39B	1.41	2.18	
H-Cdc42-F (circle)	E33C	1.49	4.16	
H-Cdc42-F (circle)	E33F	1.46	3.36	0.68
H-Cdc42-F (circle)	E42D	1.06	4.54	0.00
S-Cdc42-H (triangle)	E21	0.71	0.01	
S-Cdc42-H (circle)	E33C	1.01	2.13	
S-Cdc42-BC-H	E34B	1.15	2.53	0.39
H-Cdc42-SS	E33C	0.94	3.56	
F-Cdc42-H (circle)	E33C	1.00	4.49	
F-Cdc42-H (triangle)	E39B	1.43	2.32	
H-SS-Cdc42-F (triangle)	E39B	1.33	1.46	
H-SS-Cdc42-F (rhombus)	E42D	1.16	0.49	
H-SS-Cdc42-F (circle)	E42D	1.09	1.43	
H-Cdc42:CTIS	E39B	1.57	3.13	
H*-Cdc42:CTIS	E42C	0.92	3.04	0.89
H*-Cdc42:CTIS	E42D	1.10	3.11	
*-Cdc42:CTIS (triangle)	E42D	1.09	1.54	
*-Cdc42:CTIS (circle)	E42D	1.19	1.38	

Supplement S4

A H-Cdc42:CTIS (●): Condition screen with enterokinase

Enterokinase [U]	5	10	20		5	10	20		20
Cdc42 (50 µg)	X	X	X	X	X	X	X	X	

time and temp.	24h at RT			48h at RT				

Enterokinase [U]	5	10	20		5	10	20		20
Cdc42 (50 µg)	X	X	X	X	X	X	X	X	

time and temp.	24h at 4°C			48h at 4°C				

B H*-Cdc42:CTIS (◉): Condition screen with TEV protease

TEV (1) [U]									
TEV (2) [U]									
TEV (3) [µL]		1	5	10		1	5	10	3
Cdc42 (50 µL)	X	X	X	X	X	X	X	X	

time [h], at RT	1			3				

time [h], at 4°C	24			48				

TEV (1) [U]				5	50	5	50	10					
TEV (2) [U]									5	50	5	50	10
TEV (3) [µL]													
Cdc42 (50 µg)	X	X	X	X	X	X		X	X	X	X	

time [h], at RT	1	3	1		3			1		3	

time [h], at 4°C	24	48	24		48			24		48	

S4 Figure 1. TEV protease, but not enterokinase, is a suitable cleavage enzyme for Cdc42. (A) Cleavage reactions of enterokinase with Cdc42 (H-Cdc42:CTIS) lead mainly to undesired cleavage products. The images shown are SDS-Page. (B) Cleavage reactions of TEV protease with Cdc42 (H*-Cdc42:CTIS) lead to the desired cleavage products. Images shown are SDS-Page of cleavage reaction products of reactions using two commercially available TEV proteases (TEV (1), TEV (2)) and one enzyme that was purified in-house (TEV (3), 125 µM) and H*-Cdc42:CTIS (63 µM).

```
        10          20          30          40          50          60
MGSSHHHHHH SSGLVPRGSH MASMTGGQQM GRGSEFDDDD KMQTLKCVVV GDGAVGKTCL

        70          80          90         100         110         120
LISYTTNQFP ARYVPTVFDN YAVTVMIGDE PYTLGLFDTA GQEDYDRLRP LSYPSTDVFL

       130         140         150         160         170         180
VCFSVISPPS FENVKEKWFP EVHHHCPGVP CLVVGTQIDL RDDKVIIEKL QRQRLRPITS

       190         200         210         220         230
EQGSRLAREL KAVKYVECSA LTQRGLKNVF DEAIVAALEP PVIKKSKKCT IS
```

S4 Figure 2. Sequence of H-Cdc42:CTIS (25.7 kDa). Acidic AAs are shown in red and basic AAs are shown in blue. Potential cleavage sites for Enterokinase are highlighted in yellow. Cleavage after '…DDDDK' (site 1) results in the desired products of 4.4 and 21.3 kDa, cleavage after '…EDYDR' (site 2) results in fragments of 11.7 and 14.0 kDa, and cleavage after '…DDK' (site 3) results in 18.2 and 7.5 kDa pieces. Cleavage at site 2 in combination with cleavage at site 3 results in fragments of sizes 18.2, 14.0, 11.7, 7.5, and 6.5 kDa, which partially matches observed fragments on SDS-Page (S4 Fig. 1a).

Supplement S5

H-Cdc42, pRV035:

```
MHHHHHHMQTLKCVVVGDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVT
VMIGDEPYTLGLFDTAGQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENV
KEKWFPEVHHHCPGVPCLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGS
RLARELKAVKYVECSALTQRGLKNVFDEAIVAALEPPVIKKSKKCAIA
```

H-Cdc42:CAIA, pRV050:

```
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGSRLARELKAVKYVECSA
LTQRGLKNVFDEAIVAALEPPVIKKSKKCAIA
```

H-Cdc42:CTIS, pRV086:

```
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGSRLARELKAVKYVECSA
LTQRGLKNVFDEAIVAALEPPVIKKSKKCTIS
```

H*-Cdc42:CTIS, pRV104:

```
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFENLYFQGMQTLKCV
VVGDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFD
TAGQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPG
VPCLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGSRLARELKAVKYVEC
SALTQRGLKNVFDEAIVAALEPPVIKKSKKCTIS
```

H-Cdc42-F, pRV007:

```
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGSRLARELKAVKYVECSA
LTQRGLKNVFDEAIVAALEPPVIKKSKKCAILLPETGGDYKDHDGDYKDH
DIDYKDDDDK
```

F-Cdc42-H, pRV073:

```
MGSSDYKDHDGDYKDHDIDYKDDDDKSSGLVPRGSHMASMTGGQQMGRGS
EFDDDDKMQTLKCVVVGDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVT
VMIGDEPYTLGLFDTAGQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENV
KEKWFPEVHHHCPGVPCLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGS
RLARELKAVKYVECSALTQRGLKNVFDEAIVAALEPPVIKKSKKCAILLP
ETGGHHHHHH
```

H-Cdc42-S, pRV024:

```
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGSRLARELKAVKYVECSA
LTQRGLKNVFDEAIVAALEPPVIKKSKKCAILGGGGSLPETGGWSHPQFEK
```

H-Cdc42-SS, pRV068:
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGSRLARELKAVKYVECSA
LTQRGLKNVFDEAIVAALEPPVIKKSKKCAILLPETGGWSHPQFEKGGGS
GGGSGGGSWSHPQFEK

S-Cdc42-H, pRV030:
MGSSWSHPQFEKSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCV
VVGDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFD
TAGQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPG
VPCLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGSRLARELKAVKYVEC
SALTQRGLKNVFDEAIVAALEPPVIKKSKKCAILGGGGSLPETGGHHHHHH

SS-Cdc42-H, pRV058:
MAGWSHPQFEKGGGSGGGSGGGSWSHPQFEKSSGLVPRGSHMASMTGGQQ
MGRGSEFDDDDKMQTLKCVVVGDGAVGKTCLLISYTTNQFPADYVPTVFD
NYAVTVMIGDEPYTLGLFDTAGQEDYDRLRPLSYPSTDVFLVCFSVISPP
SFENVKEKWFPEVHHHCPGVPCLVVGTQIDLRDDKVIIEKLQRQRLRPIT
SEQGSRLARELKAVKYVECSALTQRGLKNVFDEAIVAALEPPVIKKSKKC
AILGGGGSLPETGGHHHHHH

H-SS-Cdc42-F, pRV087:
MGSSHHHHHHWSHPQFEKGGGSGGGSGGGSWSHPQFEKSSGLVPRGSHMA
SMTGGQQMGRGSEFDDDDKMQTLKCVVVGDGAVGKTCLLISYTTNQFPAD
YVPTVFDNYAVTVMIGDEPYTLGLFDTAGQEDYDRLRPLSYPSTDVFLVC
FSVISPPSFENVKEKWFPEVHHHCPGVPCLVVGTQIDLRDDKVIIEKLQR
QRLRPITSEQGSRLARELKAVKYVECSALTQRGLKNVFDEAIVAALEPPV
IKKSKKCAILLPETGGDYKDHDGDYKDHDIDYKDDDDK

S-Cdc42-BC-H, pRV051:
MGSSWSHPQFEKSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCV
VVGDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFD
TAGQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPG
VPCLVVGTQIDLRDDKVIIEKLQRQRLRPITSEQGSRLARELKAVKYVEC
SALTQRGLKNVFDEAIVAALEPPVIKKSKKMLKNFKLSKRDSNGSKGRIT
SADISTPSHDNGSVIKHIKTVPVRYLSSSSTCAILLPETGGHHHHHH

H-Cdc42-mNeon-F, pRV010:
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLSGGSAMASLPATHELHIFGSINGVD
FDMVGQGTGNPNDGYEELNLKSTKGDLQFSPWILVPHIGYGFHQYLPYPD
GMSPFQAAMVDGSGYQVHRTMQFEDGASLTVNYRYTYEGSHIKGEAQVKG
TGFPADGPVMTNSLTAADWCRSKKTYPNDKTIISTFKWSYTTGNGKRYRS
TARTTYTFAKPMAANYLKNQPMYVFRKTELKHSKTELNFKEWQKAFTDVM
GMDELYKCSGPPGRPITSEQGSRLARELKAVKYVECSALTQRGLKNVFDE
AIVAALEPPVIKKSKKCAILLPETGGDYKDHDGDYKDHDIDYKDDDDK

F-Cdc42-mNeon-H, pRV074:
MGSSDYKDHDGDYKDHDIDYKDDDDKSSGLVPRGSHMASMTGGQQMGRGS
EFDDDDKMQTLKCVVVGDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVT
VMIGDEPYTLGLFDTAGQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENV
KEKWFPEVHHHCPGVPCLVVGTQIDLRDDKVIIEKLQRQRLSGGSAMASL
PATHELHIFGSINGVDFDMVGQGTGNPNDGYEELNLKSTKGDLQFSPWIL
VPHIGYGFHQYLPYPDGMSPFQAAMVDGSGYQVHRTMQFEDGASLTVNYR
YTYEGSHIKGEAQVKGTGFPADGPVMTNSLTAADWCRSKKTYPNDKTIIS
TFKWSYTTGNGKRYRSTARTTYTFAKPMAANYLKNQPMYVFRKTELKHSK
TELNFKEWQKAFTDVMGMDELYKCSGPPGRPITSEQGSRLARELKAVKYV
ECSALTQRGLKNVFDEAIVAALEPPVIKKSKKCAILLPETGGHHHHHH

H-Cdc42-mNeon-S, pRV052:
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLSGGSAMASLPATHELHIFGSINGVD
FDMVGQGTGNPNDGYEELNLKSTKGDLQFSPWILVPHIGYGFHQYLPYPD
GMSPFQAAMVDGSGYQVHRTMQFEDGASLTVNYRYTYEGSHIKGEAQVKG
TGFPADGPVMTNSLTAADWCRSKKTYPNDKTIISTFKWSYTTGNGKRYRS
TARTTYTFAKPMAANYLKNQPMYVFRKTELKHSKTELNFKEWQKAFTDVM
GMDELYKCSGPPGRPITSEQGSRLARELKAVKYVECSALTQRGLKNVFDE
AIVAALEPPVIKKSKKCAILLPETGGWSHPQFEK

H-Cdc42-mNeon-SS, pRV067:
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLSGGSAMASLPATHELHIFGSINGVD
FDMVGQGTGNPNDGYEELNLKSTKGDLQFSPWILVPHIGYGFHQYLPYPD
GMSPFQAAMVDGSGYQVHRTMQFEDGASLTVNYRYTYEGSHIKGEAQVKG
TGFPADGPVMTNSLTAADWCRSKKTYPNDKTIISTFKWSYTTGNGKRYRS
TARTTYTFAKPMAANYLKNQPMYVFRKTELKHSKTELNFKEWQKAFTDVM
GMDELYKCSGPPGRPITSEQGSRLARELKAVKYVECSALTQRGLKNVFDE
AIVAALEPPVIKKSKKCAILLPETGGWSHPQFEKGGGSGGGSGGGSWSHP
QFEK

S-Cdc42-mNeon-H, pRV057:
MGSSWSHPQFEKSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCV
VVGDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFD
TAGQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPG
VPCLVVGTQIDLRDDKVIIEKLQRQRLSGGSAMASLPATHELHIFGSING
VDFDMVGQGTGNPNDGYEELNLKSTKGDLQFSPWILVPHIGYGFHQYLPY
PDGMSPFQAAMVDGSGYQVHRTMQFEDGASLTVNYRYTYEGSHIKGEAQV
KGTGFPADGPVMTNSLTAADWCRSKKTYPNDKTIISTFKWSYTTGNGKRY
RSTARTTYTFAKPMAANYLKNQPMYVFRKTELKHSKTELNFKEWQKAFTD
VMGMDELYKCSGPPGRPITSEQGSRLARELKAVKYVECSALTQRGLKNVF
DEAIVAALEPPVIKKSKKCAILLPETGGHHHHHH

S-Cdc42-mNeon-BC-H, pRV093:
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV

```
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLSGGSAMASLPATHELHIFGSINGVD
FDMVGQGTGNPNDGYEELNLKSTKGDLQFSPWILVPHIGYGFHQYLPYPD
GMSPFQAAMVDGSGYQVHRTMQFEDGASLTVNYRYTYEGSHIKGEAQVKG
TGFPADGPVMTNSLTAADWCRSKKTYPNDKTIISTFKWSYTTGNGKRYRS
TARTTYTFAKPMAANYLKNQPMYVFRKTELKHSKTELNFKEWQKAFTDVM
GMDELYKCSGPPGRPITSEQGSRLARELKAVKYVECSALTQRGLKNVFDE
AIVAALEPPVIKKSKKMLKNFKLSKRDSNGSKGRITSADISTPSHDNGSV
IKHIKTVPVRYLSSSSTCAILLPETGGWSHPQFEKGGGSGGGSGGGSWSH
PQFEK
```

H-SS-Cdc42-mNeon-F, pRV090:
```
MGSSHHHHHHWSHPQFEKGGGSGGGSGGGSWSHPQFEKSSGLVPRGSHMA
SMTGGQQMGRGSEFDDDDKMQTLKCVVVGDGAVGKTCLLISYTTNQFPAD
YVPTVFDNYAVTVMIGDEPYTLGLFDTAGQEDYDRLRPLSYPSTDVFLVC
FSVISPPSFENVKEKWFPEVHHHCPGVPCLVVGTQIDLRDDKVIIEKLQR
QRLSGGSAMASLPATHELHIFGSINGVDFDMVGQGTGNPNDGYEELNLKS
TKGDLQFSPWILVPHIGYGFHQYLPYPDGMSPFQAAMVDGSGYQVHRTMQ
FEDGASLTVNYRYTYEGSHIKGEAQVKGTGFPADGPVMTNSLTAADWCRS
KKTYPNDKTIISTFKWSYTTGNGKRYRSTARTTYTFAKPMAANYLKNQPM
YVFRKTELKHSKTELNFKEWQKAFTDVMGMDELYKCSGPPGRPITSEQGS
RLARELKAVKYVECSALTQRGLKNVFDEAIVAALEPPVIKKSKKCAILLP
ETGGDYKDHDGDYKDHDIDYKDDDDK
```

F-Cdc42-sfGFP-H, pRV101:
```
MGSSDYKDHDGDYKDHDIDYKDDDDKSSGLVPRGSHMASMTGGQQMGRGS
EFDDDDKMQTLKCVVVGDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVT
VMIGDEPYTLGLFDTAGQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENV
KEKWFPEVHHHCPGVPCLVVGTQIDLRDDKVIIEKLQRQRLSGGSAMSKG
EELFTGVVPILVELDGDVNGHKFSVRGEGEGDATNGKLTLKFICTTGKLP
VPWPTLVTTLTYGVQCFSRYPDHMKRHDFFKSAMPEGYVQERTISFKDDG
TYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLEYNFNSHNVYITA
DKQKNGIKANFKIRHNVEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQ
SVLSKDPNEKRDHMVLLEFVTAAGITHGMDELYKSGPPGRPITSEQGSRL
ARELKAVKYVECSALTQRGLKNVFDEAIVAALEPPVIKKSKKCAILLPET
GGHHHHHH
```

H-Cdc42-sfGFP-SS, pRV100:
```
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCVVV
GDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFDTA
GQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPGVP
CLVVGTQIDLRDDKVIIEKLQRQRLSGGSAMSKGEELFTGVVPILVELDG
DVNGHKFSVRGEGEGDATNGKLTLKFICTTGKLPVPWPTLVTTLTYGVQC
FSRYPDHMKRHDFFKSAMPEGYVQERTISFKDDGTYKTRAEVKFEGDTLV
NRIELKGIDFKEDGNILGHKLEYNFNSHNVYITADKQKNGIKANFKIRHN
VEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSVLSKDPNEKRDHMVL
LEFVTAAGITHGMDELYKSGPPGRPITSEQGSRLARELKAVKYVECSALT
QRGLKNVFDEAIVAALEPPVIKKSKKCAILLPETGGWSHPQFEKGGGSGG
GSGGGSWSHPQFEK
```

S-Cdc42-sfGFP-H, pRV099:
MGSSWSHPQFEKSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMQTLKCV
VVGDGAVGKTCLLISYTTNQFPADYVPTVFDNYAVTVMIGDEPYTLGLFD
TAGQEDYDRLRPLSYPSTDVFLVCFSVISPPSFENVKEKWFPEVHHHCPG
VPCLVVGTQIDLRDDKVIIEKLQRQRLSGGSAMSKGEELFTGVVPILVEL
DGDVNGHKFSVRGEGEGDATNGKLTLKFICTTGKLPVPWPTLVTTLTYGV
QCFSRYPDHMKRHDFFKSAMPEGYVQERTISFKDDGTYKTRAEVKFEGDT
LVNRIELKGIDFKEDGNILGHKLEYNFNSHNVYITADKQKNGIKANFKIR
HNVEDGSVQLADHYQQNTPIGDGPVLLPDNHYLSTQSVLSKDPNEKRDHM
VLLEFVTAAGITHGMDELYKSGPPGRPITSEQGSRLARELKAVKYVECSA
LTQRGLKNVFDEAIVAALEPPVIKKSKKCAILLPETGGHHHHHH

H-SS-Cdc42-sfGFP-F, pRV102:
MGSSHHHHHHWSHPQFEKGGGSGGGSGGGSWSHPQFEKSSGLVPRGSHMA
SMTGGQQMGRGSEFDDDDKMQTLKCVVVGDGAVGKTCLLISYTTNQFPAD
YVPTVFDNYAVTVMIGDEPYTLGLFDTAGQEDYDRLRPLSYPSTDVFLVC
FSVISPPSFENVKEKWFPEVHHHCPGVPCLVVGTQIDLRDDKVIIEKLQR
QRLSGGSAMSKGEELFTGVVPILVELDGDVNGHKFSVRGEGEGDATNGKL
TLKFICTTGKLPVPWPTLVTTLTYGVQCFSRYPDHMKRHDFFKSAMPEGY
VQERTISFKDDGTYKTRAEVKFEGDTLVNRIELKGIDFKEDGNILGHKLE
YNFNSHNVYITADKQKNGIKANFKIRHNVEDGSVQLADHYQQNTPIGDGP
VLLPDNHYLSTQSVLSKDPNEKRDHMVLLEFVTAAGITHGMDELYKSGPP
GRPITSEQGSRLARELKAVKYVECSALTQRGLKNVFDEAIVAALEPPVIK
KSKKCAILLPETGGDYKDHDGDYKDHDIDYKDDDDK

Cdc24-H, pDM272:
MAIQTRFASGTSLSDLKPKPSATSISIPMQNVMNKPVTEQDSLFHICANI
RKRLEVLPQLKPFLQLAYQSSEVLSERQSLLLSQKQHQELLKSNGANRDS
SDLAPTLRSSSISTATSLMSMEGISYTNSNPSATPNMEDTLLTFSMGILP
ITMDCDPVTQLSQLFQQGAPLCILFNSVKPQFKLPVIASDDLKVCKKSIY
DFILGCKKHFAFNDEELFTISDVFANSTSQLVKVLEVVETLMNSSPTIFP
SKSKTQQIMNAENQHRHQPQQSSKKHNEYVKIIKEFVATERKYVHDLEIL
DKYRQQLLDSNLITSEELYMLFPNLGDAIDFQRRFLISLEINALVEPSKQ
RIGALFMHSKHFFKLYEPWSIGQNAAIEFLSSTLHKMRVDESQRFIINNK
LELQSFLYKPVQRLCRYPLLVKELLAESSDDNNTKELEAALDISKNIARS
INENQRRTENHQVVKKLYGRVVNWKGYRISKFGELLYFDKVFISTTNSSS
EPEREFEVYLFEKIIILFSEVVTKKSASSLILKKKSSTSASISASNITDN
NGSPHHSYHKRHSNSSSSNNIHLSSSSAAAIIHSSTNSSDNNSNNSSSSS
LFKLSANEPKLDLRGRIMIMNLNQIIPQNNRSLNITWESIKEQGNFLLKF
KNEETRDNWSSCLQQLIHDLKNEQFKARHHSSTSTTSSTAKSSSMMSPTT
TMNTPNHHNSRQTHDSMASFSSSHMKRVSDVLPKRRTTSSSFESEIKSIS
ENFKNSIPESSILFRISYNNNSNNTSSSEIFTLLVEKVWNFDDLIMAINS
KISNTHNNNISPITKIKYQDEDGDFVVLGSDEDWNVAKEMLAENNEKFLN
IRLYLEHHHHHH

H-Cdc24-SS, pRV065:
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMAIQTRFAS
GTSLSDLKPKPSATSISIPMQNVMNKPVTEQDSLFHICANIRKRLEVLPQ
LKPFLQLAYQSSEVLSERQSLLLSQKQHQELLKSNGANRDSSDLAPTLRS
SSISTATSLMSMEGISYTNSNPSATPNMEDTLLTFSMGILPITMDCDPVT
QLSQLFQQGAPLCILFNSVKPQFKLPVIASDDLKVCKKSIYDFILGCKKH

```
FAFNDEELFTISDVFANSTSQLVKVLEVVETLMNSSPTIFPSKSKTQQIM
NAENQHRHQPQQSSKKHNEYVKIIKEFVATERKYVHDLEILDKYRQQLLD
SNLITSEELYMLFPNLGDAIDFQRRFLISLEINALVEPSKQRIGALFMHS
KHFFKLYEPWSIGQNAAIEFLSSTLHKMRVDESQRFIINNKLELQSFLYK
PVQRLCRYPLLVKELLAESSDDNNTKELEAALDISKNIARSINENQRRTE
NHQVVKKLYGRVVNWKGYRISKFGELLYFDKVFISTTNSSSEPEREFEVY
LFEKIIILFSEVVTKKSASSLILKKKSSTSASISASNITDNNGSPHHSYH
KRHSNSSSSNNIHLSSSSAAAIIHSSTNSSDNNSNNSSSSSSLFKLSANEP
KLDLRGRIMIMNLNQIIPQNNRSLNITWESIKEQGNFLLKFKNEETRDNW
SSCLQQLIHDLKNEQFKARHHSSTSTTSSTAKSSSMMSPTTTMNTPNHHN
SRQTHDSMASFSSSHMKRVSDVLPKRRTTSSSFESEIKSISENFKNSIPE
SSILFRISYNNNSNNTSSSEIFTLLVEKVWNFDDLIMAINSKISNTHNNN
ISPITKIKYQDEDGDFVVLGSDEDWNVAKEMLAENNEKFLNIRLYGGGGS
LPETGGWSHPQFEKGGGSGGGSGGGSWSHPQFEK
```

S-Cdc24-H, pRV031:
```
MGSSWSHPQFEKSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMAIQTRF
ASGTSLSDLKPKPSATSISIPMQNVMNKPVTEQDSLFHICANIRKRLEVL
PQLKPFLQLAYQSSEVLSERQSLLLSQKQHQELLKSNGANRDSSDLAPTL
RSSSISTATSLMSMEGISYTNSNPSATPNMEDTLLTFSMGILPITMDCDP
VTQLSQLFQQGAPLCILFNSVKPQFKLPVIASDDLKVCKKSIYDFILGCK
KHFAFNDEELFTISDVFANSTSQLVKVLEVVETLMNSSPTIFPSKSKTQQ
IMNAENQHRHQPQQSSKKHNEYVKIIKEFVATERKYVHDLEILDKYRQQL
LDSNLITSEELYMLFPNLGDAIDFQRRFLISLEINALVEPSKQRIGALFM
HSKHFFKLYEPWSIGQNAAIEFLSSTLHKMRVDESQRFIINNKLELQSFL
YKPVQRLCRYPLLVKELLAESSDDNNTKELEAALDISKNIARSINENQRR
TENHQVVKKLYGRVVNWKGYRISKFGELLYFDKVFISTTNSSSEPEREFE
VYLFEKIIILFSEVVTKKSASSLILKKKSSTSASISASNITDNNGSPHHS
YHKRHSNSSSSNNIHLSSSSAAAIIHSSTNSSDNNSNNSSSSSSLFKLSAN
EPKLDLRGRIMIMNLNQIIPQNNRSLNITWESIKEQGNFLLKFKNEETRD
NWSSCLQQLIHDLKNEQFKARHHSSTSTTSSTAKSSSMMSPTTTMNTPNH
HNSRQTHDSMASFSSSHMKRVSDVLPKRRTTSSSFESEIKSISENFKNSI
PESSILFRISYNNNSNNTSSSEIFTLLVEKVWNFDDLIMAINSKISNTHN
NNISPITKIKYQDEDGDFVVLGSDEDWNVAKEMLAENNEKFLNIRLYGGG
GSLPETGGHHHHHH
```

SS-Cdc24-H, pRV060:
```
MAGWSHPQFEKGGGSGGGSGGGSWSHPQFEKSSGLVPRGSHMASMTGGQQ
MGRGSEFDDDDKMAIQTRFASGTSLSDLKPKPSATSISIPMQNVMNKPVT
EQDSLFHICANIRKRLEVLPQLKPFLQLAYQSSEVLSERQSLLLSQKQHQ
ELLKSNGANRDSSDLAPTLRSSSISTATSLMSMEGISYTNSNPSATPNME
DTLLTFSMGILPITMDCDPVTQLSQLFQQGAPLCILFNSVKPQFKLPVIA
SDDLKVCKKSIYDFILGCKKHFAFNDEELFTISDVFANSTSQLVKVLEVV
ETLMNSSPTIFPSKSKTQQIMNAENQHRHQPQQSSKKHNEYVKIIKEFVA
TERKYVHDLEILDKYRQQLLDSNLITSEELYMLFPNLGDAIDFQRRFLIS
LEINALVEPSKQRIGALFMHSKHFFKLYEPWSIGQNAAIEFLSSTLHKMR
VDESQRFIINNKLELQSFLYKPVQRLCRYPLLVKELLAESSDDNNTKELE
AALDISKNIARSINENQRRTENHQVVKKLYGRVVNWKGYRISKFGELLYF
DKVFISTTNSSSEPEREFEVYLFEKIIILFSEVVTKKSASSLILKKKSST
SASISASNITDNNGSPHHSYHKRHSNSSSSNNIHLSSSSAAAIIHSSTNS
SDNNSNNSSSSSSLFKLSANEPKLDLRGRIMIMNLNQIIPQNNRSLNITWE
```

SIKEQGNFLLKFKNEETRDNWSSCLQQLIHDLKNEQFKARHHSSTSTTSS
TAKSSSMMSPTTTMNTPNHHNSRQTHDSMASFSSSHMKRVSDVLPKRRTT
SSSFESEIKSISENFKNSIPESSILFRISYNNNSNNTSSSEIFTLLVEKV
WNFDDLIMAINSKISNTHNNNISPITKIKYQDEDGDFVVLGSDEDWNVAK
EMLAENNEKFLNIRLYGGGGSLPETGGHHHHHH

H-SS-Cdc24-F, pRV088:
MGSSHHHHHHWSHPQFEKGGGSGGGSGGGSWSHPQFEKSSGLVPRGSHMA
SMTGGQQMGRGSEFDDDDKMAIQTRFASGTSLSDLKPKPSATSISIPMQN
VMNKPVTEQDSLFHICANIRKRLEVLPQLKPFLQLAYQSSEVLSERQSLL
LSQKQHQELLKSNGANRDSSDLAPTLRSSSISTATSLMSMEGISYTNSNP
SATPNMEDTLLTFSMGILPITMDCDPVTQLSQLFQQGAPLCILFNSVKPQ
FKLPVIASDDLKVCKKSIYDFILGCKKHFAFNDEELFTISDVFANSTSQL
VKVLEVVETLMNSSPTIFPSKSKTQQIMNAENQHRHQPQQSSKKHNEYVK
IIKEFVATERKYVHDLEILDKYRQQLLDSNLITSEELYMLFPNLGDAIDF
QRRFLISLEINALVEPSKQRIGALFMHSKHFFKLYEPWSIGQNAAIEFLS
STLHKMRVDESQRFIINNKLELQSFLYKPVQRLCRYPLLVKELLAESSDD
NNTKELEAALDISKNIARSINENQRRTENHQVVKKLYGRVVNWKGYRISK
FGELLYFDKVFISTTNSSSEPEREFEVYLFEKIIILFSEVVTKKSASSLI
LKKKSSTSASISASNITDNNGSPHHSYHKRHSNSSSSNNIHLSSSSAAAI
IHSSTNSSDNNSNNSSSSSLFKLSANEPKLDLRGRIMIMNLNQIIPQNNR
SLNITWESIKEQGNFLLKFKNEETRDNWSSCLQQLIHDLKNEQFKARHHS
STSTTSSTAKSSSMMSPTTTMNTPNHHNSRQTHDSMASFSSSHMKRVSDV
LPKRRTTSSSFESEIKSISENFKNSIPESSILFRISYNNNSNNTSSSEIF
TLLVEKVWNFDDLIMAINSKISNTHNNNISPITKIKYQDEDGDFVVLGSD
EDWNVAKEMLAENNEKFLNIRLYGGGGSLPETGGDYKDHDGDYKDHDIDY
KDDDDK

H-Bem1-F, pRV009:
MGSSHHHHHHSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMLKNFKLSK
RDSNGSKGRITSADISTPSHDNGSVIKHIKTVPVRYLSSSSTPVKSQRDS
SPKNRHNSKDITSPEKVIKAKYSYQAQTSKELSFMEGEFFYVSGDEKDWY
KASNPSTGKEGVVPKTYFEVFDRTKPSSVNGSNSSSRKVTNDSLNMGSLY
AIVLYDFKAEKADELTTYVGENLFICAHHNCEWFIAKPIGRLGGPGLVPV
GFVSIIDIATGYATGNDVIEDIKSVNLPTVQEWKSNIARYKASNISLGSV
EQQQQQSITKPQNKSAKLVDGELLVKASVESFGLEDEKYWFLVCCELSNG
KTRQLKRYYQDFYDLQVQLLDAFPAEAGKLRDAGGQWSKRIMPYIPGPVP
YVTNSITKKRKEDLNIYVADLVNLPDYISRSEMVHSLFVVLNNGFDREFE
RDENQNNIKTLQENDTATFATASQTSNFASTNQDNTLTGEDLKLNKKLSD
LSLSGSKQAPAQSTSGLKTTKIKFYYKDDIFALMLKGDTTYKELRSKIAP
RIDTDNFKLQTKLFDGSGEEIKTDSQVSNIIQAKLKISVHDIRLGGGGSL
PETGGDYKDHDGDYKDHDIDYKDDDDK

S-Bem1-H, pRV032:
MGSSWSHPQFEKSSGLVPRGSHMASMTGGQQMGRGSEFDDDDKMLKNFKL
SKRDSNGSKGRITSADISTPSHDNGSVIKHIKTVPVRYLSSSSTPVKSQR
DSSPKNRHNSKDITSPEKVIKAKYSYQAQTSKELSFMEGEFFYVSGDEKD
WYKASNPSTGKEGVVPKTYFEVFDRTKPSSVNGSNSSSRKVTNDSLNMGS
LYAIVLYDFKAEKADELTTYVGENLFICAHHNCEWFIAKPIGRLGGPGLV
PVGFVSIIDIATGYATGNDVIEDIKSVNLPTVQEWKSNIARYKASNISLG
SVEQQQQQSITKPQNKSAKLVDGELLVKASVESFGLEDEKYWFLVCCELS

NGKTRQLKRYYQDFYDLQVQLLDAFPAEAGKLRDAGGQWSKRIMPYIPGP
VPYVTNSITKKRKEDLNIYVADLVNLPDYISRSEMVHSLFVVLNNGFDRE
FERDENQNNIKTLQENDTATFATASQTSNFASTNQDNTLTGEDLKLNKKL
SDLSLSGSKQAPAQSTSGLKTTKIKFYYKDDIFALMLKGDTTYKELRSKI
APRIDTDNFKLQTKLFDGSGEEIKTDSQVSNIIQAKLKISVHDIRLGGGG
SLPETGGHHHHHH

SS-Bem1-H, pRV061:
MAGWSHPQFEKGGGSGGGSGGGSWSHPQFEKSSGLVPRGSHMASMTGGQQ
MGRGSEFDDDDKMLKNFKLSKRDSNGSKGRITSADISTPSHDNGSVIKHI
KTVPVRYLSSSSTPVKSQRDSSPKNRHNSKDITSPEKVIKAKYSYQAQTS
KELSFMEGEFFYVSGDEKDWYKASNPSTGKEGVVPKTYFEVFDRTKPSSV
NGSNSSSRKVTNDSLNMGSLYAIVLYDFKAEKADELTTYVGENLFICAHH
NCEWFIAKPIGRLGGPGLVPVGFVSIIDIATGYATGNDVIEDIKSVNLPT
VQEWKSNIARYKASNISLGSVEQQQQQSITKPQNKSAKLVDGELLVKASV
ESFGLEDEKYWFLVCCELSNGKTRQLKRYYQDFYDLQVQLLDAFPAEAGK
LRDAGGQWSKRIMPYIPGPVPYVTNSITKKRKEDLNIYVADLVNLPDYIS
RSEMVHSLFVVLNNGFDREFERDENQNNIKTLQENDTATFATASQTSNFA
STNQDNTLTGEDLKLNKKLSDLSLSGSKQAPAQSTSGLKTTKIKFYYKDD
IFALMLKGDTTYKELRSKIAPRIDTDNFKLQTKLFDGSGEEIKTDSQVSN
IIQAKLKISVHDIRLGGGGSLPETGGHHHHHH

H-SS-Bem1-F, pRV089:
MGSSHHHHHHWSHPQFEKGGGSGGGSGGGSWSHPQFEKSSGLVPRGSHMA
SMTGGQQMGRGSEFDDDDKMLKNFKLSKRDSNGSKGRITSADISTPSHDN
GSVIKHIKTVPVRYLSSSSTPVKSQRDSSPKNRHNSKDITSPEKVIKAKY
SYQAQTSKELSFMEGEFFYVSGDEKDWYKASNPSTGKEGVVPKTYFEVFD
RTKPSSVNGSNSSSRKVTNDSLNMGSLYAIVLYDFKAEKADELTTYVGEN
LFICAHHNCEWFIAKPIGRLGGPGLVPVGFVSIIDIATGYATGNDVIEDI
KSVNLPTVQEWKSNIARYKASNISLGSVEQQQQQSITKPQNKSAKLVDGE
LLVKASVESFGLEDEKYWFLVCCELSNGKTRQLKRYYQDFYDLQVQLLDA
FPAEAGKLRDAGGQWSKRIMPYIPGPVPYVTNSITKKRKEDLNIYVADLV
NLPDYISRSEMVHSLFVVLNNGFDREFERDENQNNIKTLQENDTATFATA
SQTSNFASTNQDNTLTGEDLKLNKKLSDLSLSGSKQAPAQSTSGLKTTKI
KFYYKDDIFALMLKGDTTYKELRSKIAPRIDTDNFKLQTKLFDGSGEEIK
TDSQVSNIIQAKLKISVHDIRLGGGGSLPETGGDYKDHDGDYKDHDIDYK
DDDDK

pSF1818, TEV (3):
MSKHHHHSGHHHTGHHHHSGSHHHTGESLFKGPRDYNPISSSICHLTNES
DGHTTSLYGIGFGPFIITNKHLFRRNNGTLLVQSLHGVFKVKDTTTLQQH
LVDGRDMIIIRMPKDFPPFPQKLKFREPQREERICLVTTNFQTKSMSSMV
SDTSCTFPSSDGIFWKHWIQTKDGQCGSPLVSTRDGFIVGIHSASNFTNT
NNYFTSVPKNFMELLTNQEAQQWVSGWRLNADSVLWGGHKVFMNKPEEPF
QPVKEATQLMN

Supplement S6

S6 Table 1. Primer overview. List of primers used to generate plasmids of Tab 2 and 3. Abbreviations: ins: insertion; add: addition; fw: forward; rev: reverse.

	Designed purpose	Primer	Sequence
oRV39	ins. of Cdc42	gene fw	Gcaaatggt cgcggatccg aattcGATGA CGACGATAAA ATGCAAACGC TAAAGTGTGT TGTTG
oRV40	ins. of Cdc42	gene rev	CACCGTCGTG GTCCTTGTAG TCACCGCCGG TTCCGGTAA CAAAATTGCA CATTTTTAC TTTTCTTGAT AACAGG
oRV41	ins. of Cdc42	vector fw	TTACCGGAAA CCGGCGGT
oRV42	ins. of Cdc42	vector rev	TTTATCGTCG TCATCgaatt cggatcc
oRV43	ins. of Cdc24	gene fw	caGcaaatgg gtcgcggatc cgaattcGAT GACGACGATA AAATGGCGAT CCAAACCCG
oRV44	ins. of Cdc24	gene rev	CTTGTAGTCA CCGCCGGTT CCGGTAAGCT TCCTCCGCCA CCATACAGAC GAATGTTCAA GAATTTCTCA TTG
oRV45	ins. of Cdc24	vector fw	GGTGGCGGAG GAAGCT
oRV46	ins. of Cdc24	vector rev	TTTATCGTCG TCATCgaatt cggat
oRV47	ins. of Bem1	gene fw	caGcaaatgg gtcgcgatc cgaattcGAT GACGACGATA AAATGCTGAA AAACTTCAAA CTCTCAAAAA GAGA
oRV48	ins. of Bem1	gene rev	CTTGTAGTCA CCGCCGGTT CCGGTAAGCT TCCTCCGCCA CCGAGTCTAA TATCGTGAAC GGAAATTTTC AGT
oRV49	ins. of Bem1	vector fw	GGTGGCGGAG GAAGCT
oRV50	ins. of Bem1	vector rev	TTTATCGTCG TCATCgaatt cggatcc
oRV102	add. of mNeongreen to Cdc42	gene fw	GGTAATCATC GAGAAGTTGC AAAGACAAAG ATTAAGCGGC GGTTCTGCTA TGG
oRV103	add. of mNeongreen to Cdc42	gene rev	CCTTGTTCTG ATGTAATCGG ACGACCAGGA GGCCCACTAC ATTTG
oRV104	add. of mNeongreen to Cdc42	vector fw	CAAATGTAGT GGGCCTCCTG GTCGTCCGAT TACATCAGAA CAAGG
oRV105	add. of mNeongreen to Cdc42	vector rev	CCATAGCAGA ACCGCCGCTT AATCTTTGTC TTTGCAACTT CTCGATGATT ACC
oRV145	add. of a C-term. 6His tag	vector fw	catcatcatc atcatcacTA Agaattcgag ctccgtcgac aagc
oRV146	add. of a N-term. Strep-II tag	vector rev	TTTTTCAAAC TGCGGATGGG ACCAgctgct gcccatggta tatctccT
oRV147	add. of a N-term. Strep-II tag	gene fw	TGGTCCCATC CGCAGTTTGA AAAAagcagc ggcctggtgc cg
oRV148	add. of a C-term. 6His tag	gene rev	gtgatgatga tgatgatgAC CGCCGGTTTC CGG
oRV149	add. of a C-term. Strep-II tag	gene rev	TGGTCCCATC CGCAGTTTGA AAAATAAgaa ttcgagctcc gtcgacaagc
oRV150	add. of a C-term. Strep-II tag	vector fw	TTTTTCAAAC TGCGGATGGG ACCAACCGCC GGTTCCGGT AAGC
oRV224	add. of a N-term. Twin-Strep tag	gene fw	TTCGAAAAAG GTGGAGGTTC CGGAGGTGGA TCGGGAGGTG GATCGTGGAG CCACCCGCAG TTCGA-GAAAa gcagcggcct ggtgccg
oRV225	add. of a N-term. Twin-Strep tag	vector rev	CACCUCCGGA ACCUCCACCU UUUUCGAACU GCGGGUGGCU CCAGCCCGCC AUgguauauc uccUucuuaa aguuaaaca a
oRV227	add. of a C-term. Twin-Strep tag	gene rev	ggtggctcca cgatccacct ccgatccac cttccggaacc tccacctttt tcgaactgcg ggttggctcca ACCGCCGGTT TCCGGTAA
oRV228	add. of a C-term. Twin-Strep tag	vector fw	ggaggttccg gaggtggatc gggaggtgga tcgttggagcc acccgcagtt cgagaaataa gaattcgagc tccgtcgaca agcttgcgg
oRV250	add. of N-term. 6His-Twin-Strep tag	gene fw	tggaggttcc ggaggtggat cgggaggtgg atcgtggagc caccgcagt tcgagaaaag cagcggcctg gtgcc
oRV251	add. of N-term. 6His-Twin-Strep tag	vector rev	Ccgatccacc tccggaacct ccacctttt cgaactgcgg gtggctccag gtgatgatgat gatgatggct gctg
oRV258	add. of sfGFP to Cdc42	vector fw	CACTCACGGC ATGGATGAGC TTTATAAGAG TGGGCCTCCT GGTCGTCC

oRV259	add. of sfGFP to Cdc42	vector rev	CCTGTGAACA GCTCTTCTCC TTTTGACATA GCAGAACCGC CGCTTAATCT TTGTC
oRV260	add. of sfGFP to Cdc42	gene fw	GACAAAGATT AAGCGGCGGT TCTGCTATGT CAAAAGGAGA AGAGCTGTTC ACAGG
oRV261	add. of sfGFP to Cdc42	gene rev	GGACGACCAG GAGGCCCACT CTTATAAAGC TCATCCATGC CGTGAGTG