Epistasis between mutator alleles contributes to germline mutation spectra variability in laboratory mice

This manuscript was automatically generated on October 25, 2023.

Authors

- **Thomas A. Sasani** (iD) 0000-0003-2317-1374 · tomsasani · tomsasani
  Department of Human Genetics, University of Utah

- **Aaron R. Quinlan** (iD) 0000-0003-1756-0859 · aaronquinlan Department of Human Genetics, University of Utah; Department of Biomedical Informatics, University of Utah · Funded by NIH/NHGRI R01HG012252

- **Kelley Harris** (iD) 0000-0003-0302-2523 · Kelley__Harris Department of Genome Sciences, University of Washington · Funded by NIH/NIGMS R35GM133428; Burroughs Wellcome Career Award at the Scientific Interface; Searle Scholarship; Pew Scholarship; Sloan Fellowship; Allen Discovery Center for Cell Lineage Tracing

✉ — Correspondence possible via GitHub Issues or email to Aaron R. Quinlan <aquinlan@genetics.utah.edu>, Kelley Harris <harriske@uw.edu>.

Abstract

Maintaining germline genome integrity is essential and enormously complex. Although many proteins are involved in DNA replication, proofreading, and repair [1], *mutator alleles* have largely eluded detection in mammals.

DNA replication and repair proteins often recognize sequence motifs or excise lesions at specific nucleotides. Thus, we might expect that the spectrum of *de novo* mutations — the frequencies of C>T, A>G, etc. — will differ between genomes that harbor either a mutator or wild-type allele. Previously, we used quantitative trait locus mapping to discover candidate mutator alleles in the DNA repair gene *Mutyh* that increased the C>A germline mutation rate in a family of inbred mice known as the BXDs [2,3].

In this study we developed a new method to detect alleles associated with mutation spectrum variation and applied it to mutation data from the BXDs. We discovered an additional C>A mutator locus on chromosome 6 that overlaps *Ogg1*, a DNA glycosylase involved in the same base-excision repair network as *Mutyh* [4]. Its effect depended on the presence of a mutator allele near *Mutyh*, and BXDs with mutator alleles at both loci had greater numbers of C>A mutations than those with mutator alleles at either locus alone. Our new methods for analyzing

33   mutation spectra reveal evidence of epistasis between germline mutator alleles and may be

34   applicable to mutation data from humans and other model organisms.

Introduction

Germline mutation rates reflect the complex interplay between DNA proofreading and repair pathways, exogenous sources of DNA damage, and life-history traits. For example, parental age is an important determinant of mutation rate variability; in many mammalian species, the number of germline *de novo* mutations observed in offspring increases as a function of paternal and maternal age [5,6,7,8,9]. Rates of germline mutation accumulation are also variable across human families [6,10], likely due to either genetic variation or differences in environmental exposures. Although numerous protein-coding genes contribute to the maintenance of genome integrity, genetic variants that increase germline mutation rates, known as *mutator alleles*, have proven difficult to discover in mammals.

The dearth of observed germline mutators in mammalian genomes is not necessarily surprising, since alleles that lead to elevated germline mutation rates would likely have deleterious consequences and be purged by negative selection if their effect sizes are large [11]. Moreover, germline mutation rates are relatively low, and direct mutation rate measurements require whole-genome sequencing data from both parents and their offspring. As a result, large-scale association studies — which have been used to map the contributions of common genetic variants to many complex traits — are not currently well-powered to investigate the polygenic architecture of germline mutation rates [12].

Despite these challenges, less traditional strategies have been used to identify a small number of mutator alleles in humans, macaques [13], and mice. By focusing on families with rare genetic diseases, a recent study discovered two mutator alleles that led to significantly elevated rates of *de novo* germline mutation in human genomes [14]. Other groups have observed mutator phenotypes in the germlines and somatic tissues of adults who carry cancer-predisposing inherited mutations in the POLE/POLD1 exonucleases [15,16]. Candidate mutator loci were also found by identifying human haplotypes from the Thousand Genomes Project with excess counts of derived alleles in genomic windows [17].

In mice, a germline mutator allele was recently discovered by sequencing a large family of inbred mice [2]. Commonly known as the BXDs, these recombinant inbred lines (RILs) were derived from either F2 or advanced intercrosses of C57BL/6J and DBA/2J, two laboratory strains that exhibit significant differences in their germline mutation spectra [3,18]. The BXDs were maintained via brother-sister mating for up to 180 generations, and each BXD therefore accumulated hundreds or thousands of germline mutations on a nearly-homozygous linear mosaic of parental *B* and *D* haplotypes. Due to their husbandry in a controlled laboratory setting, the BXDs were largely free from confounding by environmental heterogeneity, and the effects of selection on *de novo* mutations were attenuated by strict inbreeding [19].

In this previous study, whole-genome sequencing data from the BXD family were used to map a quantitative trait locus (QTL) for the C>A mutation rate [2]. Germline C>A mutation rates were nearly 50% higher in mice with *D* haplotypes at the QTL, likely due to genetic variation in the DNA glycosylase *Mutyh* that reduced the efficacy of oxidative DNA damage repair. Pathogenic variants of *Mutyh* also appear to act as mutators in human germline and somatic tissues [16,20]. Importantly, the QTL did not reach genome-wide significance in a scan for variation in overall

germline mutation rates, which were only modestly higher in BXDs with *D* alleles, demonstrating
the utility of mutation spectrum analysis for mutator allele discovery. Close examination of the
mutation spectrum is likely to be broadly useful for detecting mutator alleles, as genes involved
in DNA proofreading and repair often recognize particular sequence motifs or excise specific
types of DNA lesions [21]. Mutation spectra are usually defined in terms of k-mer nucleotide
context; the 1-mer mutation spectrum, for example, consists of 6 mutation types after collapsing
by strand complement (C>T, C>A, C>G, A>T, A>C, A>G), while the 3-mer mutation spectrum
contains 96 (each of the 1-mer mutations partitioned by trinucleotide context).

Although mutation spectrum analysis can enable the discovery of mutator alleles that affect the
rates of specific mutation types, early implementations of this strategy have suffered from a few
drawbacks. For example, performing association tests on the rates or fractions of every k-mer
mutation type can quickly incur a substantial multiple testing burden. Since germline mutation
rates are generally quite low, estimates of k-mer mutation type frequencies from individual
samples can also be noisy and imprecise. In populations of RILs, inbreeding duration can also
vary substantially; for example, some BXDs were inbred for only 20 generations, while others
were inbred for nearly 200. As a result, the variance of individual k-mer mutation rate estimates
in those populations will be higher than if all samples were inbred for the same duration. We
were therefore motivated to develop a statistical method that could overcome the sparsity of *de*
novo mutation spectra, eliminate the need to test each k-mer mutation type separately, and
enable sensitive detection of alleles that influence the germline mutation spectrum.

Here, we present a new mutation spectrum association test, called "aggregate mutation
spectrum distance," that minimizes multiple testing burdens and mitigates the challenges of
sparsity in *de novo* mutation datasets. We leverage this method to re-analyze germline mutation
data from the BXD family and find compelling evidence for a second mutator allele that was not
detected using previous approaches. The new allele appears to interact epistatically with the
mutator that was previously discovered in the BXDs, further augmenting the C>A germline
mutation rate in a subset of inbred mice. Our observation of epistasis suggests that mild DNA
repair deficiencies can compound one another, as mutator alleles chip away at the redundant
systems that collectively maintain germline integrity.

Results

A novel method for detecting mutator alleles

We developed a statistical method, termed "aggregate mutation spectrum distance" (AMSD), to
detect loci that are associated with mutation spectrum variation in recombinant inbred lines
(RILs) (Figure 1; *Materials and Methods*). Our approach leverages the fact that mutator alleles
often leave behind distinct and detectable impressions on the *mutation spectrum*, even if they
increase the overall mutation rate by a relatively small amount. Given a population of
haplotypes, we assume that each has been genotyped at the same collection of biallelic loci and
that each harbors *de novo* mutations which have been partitioned by k-mer context (Figure 1).
At every locus, we calculate a cosine distance between the aggregate mutation spectra of
haplotypes that inherited either parental allele. Using permutation tests, we then identify loci at
which those distances are larger than what we'd expect by random chance. To account for
polygenic effects on the mutation process that might be shared between BXDs, we also regress
the cosine distance at each marker against the genetic similarity between haplotype groups,
and assess significance using the fitted residuals (which we call the "adjusted" cosine distances)
(*Materials and Methods*).

Using simulated data, we find that our method's power is primarily limited by the initial mutation
rate of the k-mer mutation type affected by a mutator allele and the total number of *de novo*
mutations used to detect it (Figure 1-figure supplement 1). Given 100 haplotypes with an
average of 500 *de novo* germline mutations each, AMSD has approximately 90% power to
detect a mutator allele that increases the C>A *de novo* mutation rate by as little as 20%.
However, the approach has less than 20% power to detect a mutator of identical effect size that
augments the C>G mutation rate, since C>G mutations are expected to make up a smaller
fraction of all *de novo* germline mutations to begin with. Simulations also demonstrate that our
approach is well-powered to detect large-effect mutator alleles (e.g., those that increase the
mutation rate of a specific k-mer by 50%), even with a relatively small number of mutations per
haplotype (Figure 1-figure supplement 1). Both AMSD and traditional quantitative trait locus
(QTL) mapping have similar power to detect alleles that augment the rates of individual 1-mer
mutation types (Figure 1-figure supplement 2), but AMSD has a number of potential advantages
for mutator allele discovery. For example, we find that AMSD is better-powered than QTL
mapping when the number of simulated *de novo* mutations is allowed to vary (by a factor of 20)
across haplotypes (Figure 1-figure supplement 3) and when mutator allele frequencies are less
than 50% (Figure 1-figure supplement 4). However, we also caution that many of the
parameters used in our simulations are specific to the BXD mice (e.g., numbers of haplotypes,
average numbers of mutations, expected allele frequencies at markers), and do not necessarily
reflect the power of AMSD on other populations.

*Figure 1: **Overview of aggregate mutation spectrum distance method for discovering***
***mutator alleles. a)** A population of four haplotypes has been genotyped at three informative*
markers (g_1 through g_3); each haplotype also harbors unique de novo germline mutations. In
practice, de novo mutations are partitioned by k-mer context; for simplicity in this toy example,
de novo mutations are simply classified into two possible mutation types (grey squares
*represent C>(A/T/G) mutations, while grey triangles represent A>(C/T/G) mutations). **b)** At each*
informative marker g_n, we calculate the total number of each mutation type observed on
haplotypes that carry either parental allele (i.e., the aggregate mutation spectrum) using all
genome-wide de novo mutations. For example, haplotypes with A (orange) genotypes at g_1
carry a total of three "triangle" mutations and five "square" mutations, and haplotypes with B
(green) genotypes carry a total of six triangle and two square mutations. We then calculate the
cosine distance between the two aggregate mutation spectra, which we call the "aggregate
mutation spectrum distance." Cosine distance can be defined as $1 - cos(\theta)$, where θ is the
angle between two vectors; in this case, the two vectors are the two aggregate spectra. We
*repeat this process for every informative marker g_n. **c)** To assess the significance of any*
distance peaks in b), we perform permutation tests. In each of N permutations, we shuffle the
haplotype labels associated with the de novo mutation data, run a genome-wide distance scan,
and record the maximum cosine distance encountered at any locus in the scan. Finally, we
calculate the $1 - p$ percentile of the distribution of those maximum distances to obtain a
genome-wide cosine distance threshold at the specified value of p.

162 Re-identifying a mutator allele on chromosome 4 in the BXDs

We applied our aggregate mutation spectrum distance method to 117 BXDs (*Materials and*
Methods) with a total of 65,552 *de novo* germline mutations [2]. Using mutation data that were
partitioned by 1-mer nucleotide context, we discovered a locus on chromosome 4 that was

significantly associated with mutation spectrum variation (Figure 2a; maximum adjusted cosine
distance of 1.20e-2 at marker ID `rs27509845`; position 118.28 Mbp in GRCm38/mm10
coordinates; 90% bootstrap confidence interval from 114.79 - 118.75 Mbp).

Figure 2: **Results of aggregate mutation spectrum distance scans in the BXDs. a)** *Adjusted cosine distances between aggregate 1-mer de novo mutation spectra on BXD haplotypes (n = 117 haplotypes; 65,552 total mutations) with either D or B alleles at 7,128 informative markers.*

Using quantitative trait locus (QTL) mapping, we previously identified a nearly-identical locus on
chromosome 4 that was significantly associated with the C>A germline mutation rate in the
BXDs [2]. This locus overlapped 21 protein-coding genes that were annotated by the Gene
Ontology as being involved in "DNA repair," but only one of those genes contained
nonsynonymous differences between the two parental strains: *Mutyh*. *Mutyh* encodes a protein
involved in the base-excision repair of 8-oxoguanine (8-oxoG), a DNA lesion caused by
oxidative damage, and prevents the accumulation of C>A mutations [4,22,23]. C>A germline
mutation fractions are nearly 50% higher in BXDs that inherit *D* genotypes at marker ID
`rs27509845` (the marker at which we observed the highest adjusted cosine distance on
chromosome 4) than in those that inherit *B* genotypes (Figure 3) [2].

An additional germline mutator allele on chromosome 6

After confirming that AMSD could recover the mutator locus overlapping *Mutyh*, we tested its
ability to identify additional mutator loci in the BXDs. To eliminate potential confounding of the
mutation spectrum landscape by the large-effect mutator locus on chromosome 4, we
performed AMSD scans that were conditional on the presence of either *D* or *B* alleles at
`rs27509845`. We hypothesized that such conditioning might reveal epistatic interactions
between alleles at the chromosome 4 locus and mutator alleles elsewhere in the genome.
Specifically, we divided the BXDs into those with either *D* (n = 66) or *B* (n = 44) genotypes at
`rs27509845` (n = 7 BXDs were heterozygous) and ran an aggregate mutation spectrum
distance scan using each group separately (Figure 2b-c). We excluded the BXD68 RIL from
these scans, since we previously found that BXD68 harbors a strain-private C>A mutator allele
of even larger effect than that of a *D* allele at `rs27509845` [2].

Using the BXDs with *D* genotypes at `rs27509845`, we identified a locus on chromosome 6 that
was significantly associated with mutation spectrum variation (Figure 2b; maximum adjusted
cosine distance of 3.69e-3 at marker `rs46276051`; position 111.27 Mbp in GRCm38/mm10
coordinates; 90% bootstrap confidence interval from 95.01 - 114.02 Mbp). This signal was
specific to BXDs with *D* genotypes at the `rs27509845` locus, as we did not observe any new
mutator loci after performing an AMSD scan using BXDs with *B* genotypes at `rs27509845`
(Figure 2c). The peak markers on chromosome 4 and 6 did not exhibit strong linkage
disequilibrium (R^2 = 4e-5). We also performed QTL scans for the fractions of each 1-mer
mutation type using the same mutation data, but none produced a genome-wide significant log-
odds score at any locus (Figure 2-figure supplement 1; *Materials and Methods*).

We queried the region surrounding the top marker on chromosome 6 (+/- the 90% bootstrap
confidence interval) and discovered 64 protein-coding genes, of which four were annotated with
a Gene Ontology (GO) [24,25] term related to "DNA repair": *Fancd2*, *Ogg1*, *Setmar*, and *Rad18*.
None of the remaining genes were annotated with a cellular function that would obviously
contribute to a germline mutator phenotype; however, many of these GO annotations are
imperfect and/or incomplete. Although we focus our analysis on DNA repair genes, it remains
possible that other genes within the confidence interval could underlie the C>A mutator
phenotype we identified in the BXDs.

Of the annotated DNA repair genes within the confidence interval, two harbored
nonsynonymous differences between the parental C57BL/6J and DBA/2J strains (Table 1).
Ogg1 encodes a key member of the base-excision repair response to oxidative DNA damage (a
pathway that also includes *Mutyh*), and in mice *Setmar* encodes a SET domain-containing
histone methyltransferase; both *Ogg1* and *Setmar* are expressed in mouse gonadal cells.
Because the bootstrap can exhibit poor coverage in QTL mapping studies [26], we also scanned
an interval +/- 5 Mbp from the peak AMSD marker on chromosome 6 for additional candidate
genes. Although the choice of a 10 Mbp interval is somewhat arbitrary, the interval does contain
a plausible candidate: *Mbd4*, a protein-coding gene involved in base excision repair that also
harbors a non-synonymous difference between the BXD parental strains and is expressed in
mouse gonads (Table 1).

Table 1: Nonsynonymous mutations in DNA repair genes near the chr6 peak

Gene name	Ensembl transcript name	Nucleotide change	Amino acid change	Position in GRCm38/mm10 coordinates	PhyloP conservation score	SIFT prediction
Setmar	ENSMUST00000049246	C>T	p.Leu103Phe	chr6:108,075,853	0.422	0.0 (intolerant/deleterious)
Setmar	ENSMUST00000049246	T>G	p.Ser273Arg	chr6:108,076,365	-0.355	0.3 (tolerant/benign)
Ogg1	ENSMUST00000032406	A>G	p.Thr95Ala	chr6:113,328,510	-0.016	0.84 (tolerant/benign)
Mbd4	ENSMUST00000032469	C>T	p.Asp129Asn	chr6:115,849,644	2.28	0.02 (intolerant/deleterious)

We also considered the possibility that expression quantitative trait loci (eQTLs), rather than
nonsynonymous mutations, could contribute to the C>A mutator phenotype associated with the
locus on chromosome 6. Using GeneNetwork [27] we mapped eQTLs for the four
aforementioned DNA repair genes (as well as *Mbd4*) in a number of tissues, though we did not
have access to expression data from germline cells. Notably, *D* alleles near the cosine distance
peak on chromosome 6 were significantly associated with decreased *Ogg1* expression in
kidney, liver, hippocampus, and gastrointestinal tissues (Table supplement 1). Although these
cis-eQTLs are challenging to interpret (given their tissue specificity and our lack of access to
germline expression data), the presence of strong-effect cis-eQTLs for *Ogg1* suggests that the

C>A mutator phenotype observed in the BXDs may be mediated by regulatory, rather than
protein-altering, variants.

Finally, we queried a dataset of structural variants (SVs) identified via high-quality, long-read
assembly of inbred laboratory mouse strains [28] and found 176 large insertions or deletions
(>100 bp) within the 90% bootstrap confidence interval around the cosine distance peak on
chromosome 6; none overlapped the exonic sequences of protein-coding genes. One protein-
coding gene involved in DNA repair (*Rad18*) harbored an intronic deletion within the interval on
chromosome 6 (chr6:112,629,618-112,636,619); however, additional experimental evidence will
be needed to probe the functional impact of this structural variant.

Evidence of epistasis between germline mutator alleles

Next, we more precisely characterized the effects of the chromosome 4 and 6 mutator alleles on
mutation spectra in the BXDs. To pinpoint the mutation type(s) that underlied the significant
cosine distance peak on chromosome 6, we compared the aggregate counts of each 1-mer
mutation type (plus CpG>TpG) on BXD haplotypes with *D* genotypes at rs27509845 and
either *D* or *B* genotypes at rs46276051. We found that C>A mutations were significantly
enriched on BXD haplotypes with *D* genotypes at the chromosome 6 mutator locus, relative to
those with *B* genotypes (χ^2 statistic = 85.36, p = 2.48e-20). On average, C>A germline mutation
fractions were significantly higher in BXDs with *D* alleles at both mutator loci than in BXDs with
D alleles at either locus alone (Figure 3a and 3-figure supplement 1). Among BXDs with *B*
alleles at the locus overlapping *Mutyh*, those with *D* alleles on chromosome 6 did not exhibit
significantly elevated C>A mutation fractions (Figure 3a). After controlling for inbreeding
duration, we observed that C>A *de novo* mutation counts were always highest in BXDs with *D*
alleles at both mutator loci (Figure 3b). After 100 generations of inbreeding, BXDs with *D* alleles
at both mutator loci were predicted to have 238.9 C>A mutations (95% CI: 231.4 - 246.4), about
20% more than the 199.0 mutations (95% CI: 193.3 - 204.7) predicted in those with *D* and *B*
alleles at the chromosome 4 and chromosome 6 loci, respectively (Figure 3b).

We also used SigProfilerExtractor [29] to assign the germline mutations in each BXD to single-
base substitution (SBS) mutation signatures from the COSMIC catalog [30]. Mutation signatures
often reflect specific exogenous or endogenous sources of DNA damage, and the proportions of
mutations attributable to particular SBS signatures can suggest a genetic or environmental
etiology. The SBS1, SBS5, and SBS30 mutation signatures were active in nearly all BXDs,
regardless of genotypes at the chromosome 4 and 6 mutator loci (Figure 3c). However, the
SBS18 signature, which is dominated by C>A mutations and likely reflects unrepaired DNA
damage from reactive oxygen species, was almost exclusively active in mice with *D* alleles at
the chromosome 4 locus; the highest SBS18 activity was observed in mice with *D* alleles at both
mutator loci (Figure 3c). SBS18 activity was lowest in mice with *D* alleles at the chromosome 6
mutator locus alone (Figure 3c), further demonstrating that *D* alleles at this locus are not
sufficient to cause a mutator phenotype.

To more formally test for statistical epistasis, we fit a generalized (Poisson) linear model
predicting counts of C>A mutations in each BXD as a function of genotypes at rs27509845
and rs46276051 (the markers with the largest adjusted cosine distance at the two mutator

loci); the model also accounted for differences in inbreeding duration and sequencing coverage
between the BXDs (*Materials and Methods*). A model that included an interaction term between
genotypes at the two markers fit the data significantly better than a model including only additive
effects (p = 7.92e-7; *Materials and Methods*), indicating that the combined effects of *D*
genotypes at both loci exceeded the sum of marginal effects of *D* genotypes at either locus
alone.

*Figure 3: **BXD mutation spectra are affected by alleles at both mutator loci. a)** C>A de novo*
germline mutation fractions in BXDs with either D or B genotypes at markers `rs27509845`
(chr4 peak) and `rs46276051` *(chr6 peak). Distributions of C>A mutation fractions were*
compared with two-sided Mann-Whitney U-tests; annotated p-values are uncorrected. `B-B`
vs. `B-D` *comparison: U-statistic = 149.0, p = 7.58e-2;* `B-D` *vs* `D-D` *comparison: U-statistic = 21.0,*
p = 2.61e-8; `D-B` *vs* `D-D` *comparison: U-statistic = 232.5, p = 6.99e-5.* ***b)** The count of C>A de*
novo germline mutations in each BXD was plotted against the number of generations for which
it was inbred. Lines represent predicted C>A counts in each haplotype group from a generalized
linear model (Poisson family, identity link), and shading around each line represents the 95%
confidence interval. ***c)** Germline mutations in each BXD were assigned to COSMIC SBS*
mutation signatures using SigProfilerExtractor [29]. After grouping BXDs by their genotypes at
`rs27509845` *and* `rs46276051`, *we calculated the fraction of mutations in each group that was*
attributed to each signature. The proposed etiologies of each mutation signature are: SBS1

(spontaneous deamination of methylated cytosine nucleotides at CpG contexts), SBS5
(unknown, clock-like signature), SBS18 (damage by reactive oxygen species, related to SBS36
and defective base-excision repair due to loss-of-function mutations in MUTYH), and SBS30
(defective base-excision repair due to NTHL1 mutations).

To explore the effects of the two mutator loci in other inbred laboratory mice, we also compared
the germline mutation spectra of Sanger Mouse Genomes Project (MGP) strains [31]. Dumont
[18] previously identified germline mutations that were private to each of the 29 MGP strains;
these private variants likely represent recent *de novo* mutations (Figure 3—figure supplement
2). Only two of the MGP strains possess *D* genotypes at both the chromosome 4 and
chromosome 6 mutator loci: DBA/1J and DBA/2J. As before, we tested for epistasis in the MGP
strains by fitting two linear models predicting C>A mutation counts as a function of genotypes at
the two mutator loci. A model incorporating an interaction term did not fit the MGP data
significantly better than a model with additive effects alone (p = 0.806), so we are unable to
confirm the signal of epistasis; however, this may be due to the smaller number of MGP strains
with *de novo* germline mutation data.

320 **Some of the candidate mutator alleles are segregating in wild**
321 **mice**

To determine whether the candidate mutator alleles on chromosome 6 were segregating in
natural populations, we queried previously published sequencing data generated from 67 wild-
derived mice [32]. Although these data are not a comprehensive sampling of the genetic
diversity in wild mice, they include three subspecies of *Mus musculus*, as well as the outgroup
Mus spretus. We found that the *Ogg1 D* allele was segregating at an allele frequency of 0.259 in
Mus musculus domesticus, the species from which C57BL/6J and DBA/2J derive the majority of
their genomes [33], and was fixed in *Mus musculus musculus*, *Mus musculus castaneus*, and
the outgroup *Mus spretus* (Figure 3-figure supplement 3). The *Setmar* p.Ser273Arg *D* allele was
also present at an allele frequency of 0.37 in *Mus musculus domesticus*, while *D* alleles at the
Setmar p.Leu103Phe variant were not observed in any wild *Mus musculus domesticus* animals.
D alleles at the *Mbd4* p.Asp129Asn variant were also absent from all wild mouse populations
(Figure 3-figure supplement 3).

Discussion

Epistasis between germline mutator alleles

We have identified a locus on chromosome 6 that amplifies a C>A germline mutator phenotype in the BXDs, a family of inbred mice derived from the laboratory strains DBA/2J and C57BL/6J. DBA/2J (*D*) alleles at this locus have no significant effect on C>A mutation rates in mice that also harbor "wild-type" C57BL/6J (*B*) alleles at a previously discovered mutator locus on chromosome 4 [2]. However, mice with *D* alleles at *both* loci have even higher mutation rates than those with *D* alleles at the chromosome 4 mutator locus alone (Figure 3). Epistatic interactions between mutator alleles have been previously documented in yeast [34] and in human cell lines [35], but never to our knowledge in a whole-animal context.

Importantly, we discovered epistasis between germline mutator alleles in an unnatural population of model organisms that have been inbred by brother-sister mating in a highly controlled laboratory environment [3]. This breeding setup has likely attenuated the effects of natural selection on all but the most deleterious alleles [19], and may have facilitated the fixation of large-effect mutator alleles that would be less common in wild mice. Without fine-mapping the chromosome 6 mutator allele, however, we are unable to trace its origin to either a captive breeding colony of laboratory mice or a wild, outbreeding *Mus musculus* population. If the mutator allele on chromosome 6 has even a weak deleterious fitness effect, there might be a greater likelihood that it persisted in captivity. Indeed, if purifying selection is required to keep mutation rates low, mutational pressure might cause mutation rates to rise in just a few generations of relaxed selection. This dynamic may explain the recent discovery of a large-effect mutator allele in a rhesus macaque research colony [13], as well as the observation that domesticated animals tend to have higher mutation rates than those in the wild [36]. Although we have not conclusively fine-mapped the chromosome 6 mutator locus to a causal variant, we argue that nonsynonymous or regulatory variants in the DNA glycosylase *Ogg1* are the best candidates.

Protein-coding genes that may underlie the chromosome 6 mutator locus

Five protein-coding genes involved in DNA repair overlap the C>A mutator locus on chromosome 6: *Ogg1*, a glycosylase that excises the oxidative DNA lesion 8-oxoguanine (8-oxoG) [4], *Setmar*, a histone methyltransferase involved in non-homologous end joining (NHEJ) of double-stranded breaks (DSBs) [37,38], *Fancd2*, and *Rad18*. One other DNA repair gene, *Mbd4*, lies just outside of the 90% bootstrap confidence interval on chromosome 6 (but within a 10 Mbp interval around the peak AMSD marker). We are unable to conclusively determine that one or more of these genes harbors a causal variant underlying the observed C>A mutator phenotype, but we believe that *Ogg1* is the most plausible candidate. *Ogg1* is a member of the same base-excision repair pathway as *Mutyh* (the gene that likely underlies the chromosome 4 mutator locus), contains a nonsynonymous fixed difference between the C57BL/6J and DBA/2J parental strains, and appears to be regulated by cis-eQTLs across a number of tissues within the BXD cohort.

The C57BL/6J and DBA/2J *Setmar* coding sequences differ by two missense variants (Table 1),
one of which is predicted to be deleterious by *in silico* tools. The primate *SETMAR* ortholog is
involved in NHEJ of double-strand breaks, but its role in DNA repair appears to depend on the
function of both a SET methyltransferase domain and a *Mariner*-family transposase domain
[38,39,40]. Since the murine *Setmar* ortholog lacks the latter element, and because primate
SETMAR is involved in a DNA repair process that is not expected to affect the rate of C>A
mutations, we believe it is unlikely to underlie the epistatic interaction between the chromosome
4 and 6 mutator loci in the BXDs (*Appendix 1*). Moreover, we did not observe any significant cis-
eQTLs for *Setmar* across a variety of tissues in the BXD cohort (Table supplement 1). None of
the remaining DNA repair genes (*Fancd2* or *Rad18*) contains a nonsynonymous fixed difference
between the C57BL/6J and DBA/2J parental strains, and none appear to be regulated by cis-
eQTLs that would feasibly lead to a germline C>A mutator phenotype (Table supplement 1); the
only significant cis-eQTLs we observed for genes other than *Ogg1* were those at which *D* alleles
actually led to *increased* expression.

An *Ogg1* mutator allele might impair the excision of 8-oxoguanine lesions

Ogg1 is a member of the same base-excision repair (BER) pathway as *Mutyh*, the protein-
coding gene we previously implicated as harboring mutator alleles at the locus on chromosome
4 [4]. Each of these genes has a distinct role in the BER response to oxidative DNA damage,
and thereby the prevention of C>A mutations [41,42]. Following damage by reactive oxygen
species, *Ogg1* is able to recognize and remove 8-oxoguanine lesions that are base-paired with
cytosine nucleotides; once 8-oxoG is excised, other members of the BER pathway are
mobilized to restore a proper G:C base pair at the site. If an 8-oxoG lesion is not removed
before the cell enters S-phase, adenine can be mis-incorporated opposite 8-oxoG during DNA
replication [41]. If this occurs, *Mutyh* can excise the mispaired adenine, leaving a one-nucleotide
gap that is processed and filled with a cytosine by other BER proteins. The resulting C:8-oxoG
base pair can then be "returned" to *Ogg1* for excision and repair. Defects in the BER response
to oxidative damage lead to significantly elevated rates of C>A mutation. For example, triple-
knockout (KO) mice lacking *Ogg1*, *Mutyh*, and *Mth1* (which encodes an enzyme that prevents 8-
oxo-dGTP from being incorporated during DNA synthesis [43]) accumulate a 100-fold excess of
8-oxoG in their gonadal cells [42]. Almost 99% of *de novo* germline mutations in the
Ogg1/Mutyh/Mth1 triple KO mice are C>A transversions, demonstrating the clear role of 8-oxoG
repair in preventing C>A mutation. Additionally, missense mutations and loss-of-heterozygosity
in *Ogg1* have been associated with increased risk of human cancer [44,45], and copy-number
losses of either *Ogg1* or *Mutyh* are linked to elevated rates of spontaneous C>A mutation in
human neuroblastoma [46].

Nonsynonymous mutations may underlie the chromosome 6 mutator phenotype

412 The p.Thr95Ala *Ogg1* missense variant is not predicted to be deleterious by the *in silico* tool
413 SIFT [47], and occurs at a nucleotide that is not particularly well-conserved across mammalian
414 species (Table 1). We also observe that the *D* allele at p.Thr95Ala is segregating at an allele
415 frequency of approximately 26% among wild-derived *Mus musculus domesticus* animals, and is

fixed in other wild populations of *Mus musculus musculus*, *Mus musculus castaneus*, and *Mus spretus* . Although we would expect *a priori* that *Ogg1* deficiency should lead to increased 8-oxoG accumulation and elevated C>A mutation rates, these lines of evidence suggest that p.Thr95Ala is not highly deleterious on its own, and might only exert a detectable effect on the BER gene network when *Mutyh* function is also impaired (or *Mutyh* expression is decreased). It is also possible that *D* alleles at *Ogg1* lead to a very subtle increase in C>A mutation rates, and we are simply underpowered to detect such a small mutation rate effect in the BXDs.

No indication of causal structural variation or mobile element insertions near the chromosome 6 mutator locus

Although we argue above that *Ogg1* is likely the the best candidate gene to explain the new BXD C>A mutator phenotype, we cannot conclusively determine that the p.Thr95Ala missense mutation is a causal allele. We previously hypothesized that *Mutyh* missense mutations on *D* haplotypes were responsible for the large-effect C>A mutator phenotype we observed in the BXDs [2]. However, subsequent long-read assemblies of several inbred laboratory mouse strains revealed that this mutator phenotype might be caused by a ~5 kbp mobile element insertion (MEI) within the first intron of *Mutyh* [28], which is associated with significantly reduced expression of *Mutyh* in embryonic stem cells. We queried the new high-quality assemblies for evidence of mobile elements or other large structural variants (SVs) in the region surrounding the mutator locus on chromosome 6, but found no similarly compelling evidence that either SVs or MEIs might underlie the mutator phenotype described in this study.

Expression quantitative trait loci (eQTLs) might mediate germline mutator phenotypes in the BXDs

We observed strong-effect cis-eQTLs for *Ogg1* expression across a number of tissues in the BXDs (Table supplement 1). In each of these tissue types, *D* genotypes were associated with decreased expression of *Ogg1*. As mentioned above, new evidence from long-read genome assemblies has demonstrated that an intronic mobile element insertion in *Mutyh* may be responsible for decreased *Mutyh* expression, and therefore higher C>A mutation rates, in BXDs with *D* haplotypes at the chromosome 4 mutator locus [28]. Taken together, these results raise the exciting possibility that the mutator loci on both chromosome 4 and chromosome 6 lead to increased C>A mutation rates by lowering the expression of DNA repair genes in the same base-excision repair network. Moreover, the effect of the mutator locus on chromosome 6 may be conditional on the expression of the mutator near *Mutyh*, suggesting that complex genetic interactions can underlie germline mutator phenotypes in mammalian genomes.

Mbd4 may buffer the effects of *Mutyh* mutator alleles by triggering apoptosis

As mentioned in the *Results*, *Mbd4* lies just outside of the 90% bootstrap confidence interval on chromosome 6, but within a 10 Mbp window surrounding the AMSD peak. Due to the uncertainties associated with bootstrap confidence intervals in QTL mapping [26], we have included a discussion of the evidence supporting *Mbd4* as a causal gene below.

Unlike the *Ogg1* p.Thr95Ala mutation, the p.Asp129Asn variant in *Mbd4* resides within an
annotated protein domain (the *Mbd4* methyl-CpG binding domain), occurs at a nucleotide and
amino acid residue that are both well-conserved, and is predicted to be deleterious by SIFT [47]
(Table 1). A missense mutation that affects the homologous amino acid in humans
(p.Asp142Gly in GRCh38/hg38) is also present on a single haplotype in the Genome
Aggregation Database (gnomAD) [48] and is predicted by SIFT and Polyphen [49] to be
"deleterious" and "probably_damaging" in human genomes, respectively.

One puzzling observation is that loss-of-function mutations in *Mbd4* are not typically associated
with C>A mutator phenotypes. Instead, *Mbd4* deficiency is usually implicated in C>T
mutagenesis at CpG sites, and we did not detect an excess of C>T mutations in BXDs with *D*
alleles at the chromosome 6 mutator locus (Figure 3-figure supplement 1). However, loss of
function mutations in *Mbd4* have also been shown to exacerbate the effects of exogenous DNA
damage agents. For example, mouse embryonic fibroblasts that harbor homozygous *Mbd4*
knockouts fail to undergo apoptosis following treatment with a number of chemotherapeutics
and mutagenic compounds [50]. Most of these exogenous mutagens cause DNA damage that is
normally repaired by mismatch repair (MMR) machinery, but murine intestinal cells with biallelic
Mbd4 LOF mutations also showed a reduced apoptotic response to gamma irradiation, which is
repaired independently of the MMR gene *Mlh1* [51]. Homozygous loss of *Mbd4* function also
leads to accelerated intestinal tumor formation in mice that harbor an *Apc* allele that
predisposes them to intestinal neoplasia [52], and mice with biallelic truncations of the *Mbd4*
coding sequence exhibit modestly increased mutation rates in colon cancer cell lines, including
increased C>A mutation rates in certain lines [53].

Together, these lines of evidence suggest that *Mbd4* can modulate sensitivity to many types of
exogenous mutagens, potentially through its role in determining whether cells harboring DNA
damage should undergo apoptosis [50,51]. We speculate that in mice with deficient 8-
oxoguanine repair — caused by a mutator allele in *Mutyh*, for example — reactive oxygen
species (ROS) could cause accumulation of DNA damage in the germline. If those germ cells
harbor fully functional copies of *Mbd4*, they might be able to trigger apoptosis and partially
mitigate the effects of a *Mutyh* mutator allele. However, mice with reduced activity of both *Mbd4*
and *Mutyh* may have a reduced ability to initiate cell death in response to DNA damage; as a
result, their germ cells may accumulate even higher levels of ROS-mediated damage, leading to
substantially elevated germline C>A mutation rates.

We anticipate that future experimental work will be able to more conclusively establish a
mechanistic explanation for the epistatic interaction between mutator loci described in this
paper.

490 Strengths and limitations of the aggregate mutation spectrum
491 distance approach

Our aggregate mutation spectrum distance (AMSD) approach was able to identify a mutator
allele that escaped notice using quantitative trait locus (QTL) mapping. To more systematically
compare the power of AMSD and QTL mapping, we performed simulations under a variety of
possible parameter regimes. Overall, we found that AMSD and QTL mapping have similar

power to detect mutator alleles on haplotypes that each harbor tens or hundreds of *de novo*
germline mutations (Figure 1-figure supplement 2). Nonetheless, only AMSD was able to
discover the mutator locus on chromosome 6 in the BXDs, demonstrating that it outperforms
QTL mapping in certain experimental systems. For example, simulations demonstrate that
AMSD enjoys greater power than QTL mapping when haplotypes carry highly variable numbers
of mutations that can be leveraged for mutator mapping (Figure 1-figure supplement 3).
Because the BXDs were generated in six breeding epochs over a period of nearly 40 years, the
oldest lines have accumulated orders of magnitude more mutations than the youngest lines;
these younger BXDs have much noisier mutation spectra as a result. While approaches for QTL
mapping typically weight the phenotypic measurements of each sample equally, AMSD
compares the *aggregate* mutation spectra of haplotypes at every locus, a property that likely
increased its power to detect mutators in the BXD dataset.

Another benefit of the AMSD approach is that it obviates the need to perform separate
association tests for every possible k-mer mutation type, and therefore the need to adjust
significance thresholds for multiple tests. Since AMSD compares the complete mutation
spectrum between haplotypes that carry either allele at a site, it would also be well-powered to
detect a mutator allele that exerted a coordinated effect on multiple k-mer mutation types (e.g.,
increased the rates of both C>T and C>A mutations).

However, the AMSD method suffers a handful of drawbacks when compared to QTL mapping.
Popular QTL mapping methods (such as R/qtl2 [54]) use linear models to test associations
between genotypes and phenotypes, enabling the inclusion of additive and interactive
covariates, as well as kinship matrices, in QTL scans. Although we have developed methods to
account for inter-sample relatedness in the AMSD approach (*Materials and Methods*), they are
not as flexible as similar methods in QTL mapping software. Additionally, the AMSD method
assumes that mutator alleles affect a subset of k-mer mutation types; if a mutator allele
increased the rates of all mutation types equally on haplotypes that carried it, AMSD would be
unable to detect it.

Discovering mutator alleles in other experimental systems

Our discovery of a second BXD mutator allele underscores the power of recombinant inbred
lines (RILs) as a resource for dissecting the genetic architecture of germline mutation rates.
Large populations of RILs exist for many model organisms, and we anticipate that as whole-
genome sequencing becomes cheaper and cheaper, the AMSD method could be useful for
future mutator allele discovery outside of the BXDs. At the same time, RILs are a finite resource
that require enormous investments of time and labor to construct. If germline mutator alleles are
only detectable in these highly unusual experimental populations, we are unlikely to discover
more than a small fraction of the mutator alleles that may exist in nature.

In natural, outbreeding populations, selection on germline mutator alleles will likely prevent
large-effect mutators from reaching high allele frequencies, but a subset may be detectable by
sequencing a sufficient number of human trios [56]. Since germline mutators often seem to
exert their effects on a small number of mutation types, mutation spectrum analyses may have

greater power to detect the genes that underlie heritable mutation rate variation, even if each
gene has only a modest effect on the overall mutation rate per generation.

Thousands of human pedigrees have been sequenced in an effort to precisely estimate the rate
of human *de novo* germline mutation [5,6,55], and as family sequencing has become a more
common step in the diagnosis of many congenital disorders, these datasets are growing on a
daily basis. AMSD could potentially be applied to these large cohorts of two- or three-generation
families. We propose that by pooling sparse mutation counts across many individuals who share
the same candidate mutator allele, even a subtle mutator signal might rise above the noise of *de*
novo germline mutation rate estimates. We note that the aggregate mutation spectrum distance
approach will require modification before it can be successfully applied to cohorts of outbred,
sexually-reproducing individuals. AMSD assumes that individuals harbor one of two possible
genotypes at each marker, and does not yet account for heterozygous genotypes. As a result,
our method is currently applicable only to resources like the BXD RILs, in which individuals have
been inbred for sufficiently long that effectively all genotypes are homozygous. However, as
human trio datasets become larger and larger in the coming years, rare mutator alleles may
eventually be represented in as many individuals as the high-frequency variants present in the
BXDs, thereby enabling more systematic mapping of elusive mutator phenotypes.

Materials and Methods

Identifying *de novo* germline mutations in the BXDs

The BXD resource currently comprises a total of 152 recombinant inbred lines (RILs). BXDs were derived from either F2 or advanced intercrosses, and subsequently inbred by brother-sister mating for up to 180 generations [3]. BXDs were generated in distinct breeding "epochs," which were each initiated with a distinct cross of C57BL/6J and DBA/2J parents; epochs 1, 2, 4, and 6 were derived from F2 crosses, while epochs 3 and 5 were derived from advanced intercrosses [3]. Previously, we analyzed whole-genome sequencing data from the BXDs and identified candidate *de novo* germline mutations in each line [2]. A detailed description of the methods used for DNA extraction, sequencing, alignment, and variant processing, as well as the characteristics of the *de novo* mutations, are available in a previous manuscript [2].

Briefly, we identified private single-nucleotide mutations in each BXD that were absent from all other BXDs, as well as from the C57BL/6J and DBA/2J parents. We required each private variant to be meet the following criteria:

- genotyped as either homozygous or heterozygous for the alternate allele, with at least 90% of sequencing reads supporting the alternate allele

- supported by at least 10 sequencing reads

- Phred-scaled genotype quality of at least 20

- must not overlap regions of the genome annotated as segmental duplications or simple repeats in GRCm38/mm10

- must occur on a parental haplotype that was inherited by at least one other BXD at the same locus; these other BXDs must be homozygous for the reference allele at the variant site

A new approach to discover germline mutator alleles

Calculating aggregate mutation spectrum distance

We developed a new approach to discover loci that affect the germline *de novo* mutation spectrum in biparental RILs (Figure 1).

We assume that a collection of haplotypes has been genotyped at informative markers, and that *de novo* germline mutations have been identified on each haplotype.

At each informative marker, we divide haplotypes into two groups based on the parental allele that they inherited. We then compute a k-mer mutation spectrum using the aggregate mutation counts in each haplotype group. The k-mer mutation spectrum contains the frequency of every possible k-mer mutation type in a collection of mutations, and can be represented as a vector of

size $6 \times 4^{k-1}$ after collapsing by strand complement. For example, the 1-mer mutation spectrum
is a 6-element vector that contains the frequencies of C>T, C>G, C>A, A>G, A>T, and A>C
mutations. Since C>T transitions at CpG nucleotides are often caused by a distinct mechanism
(spontaneous deamination of methylated cytosine), we expand the 1-mer mutation spectrum to
include a separate category for CpG>TpG mutations [57].

At each marker, we then compute the cosine distance between the two aggregate spectra. The
cosine distance between two vectors **A** and **B** is defined as

593
$$D^C = 1 - \frac{\mathbf{A} \cdot \mathbf{B}}{\parallel \mathbf{A} \parallel \parallel \mathbf{B} \parallel}$$

where $\parallel \mathbf{A} \parallel$ and $\parallel \mathbf{B} \parallel$ are the L^2 (or Euclidean) norms of **A** and **B**, respectively. The cosine
distance metric has a number of favorable properties for comparing mutation spectra. Since it
adjusts for the magnitude of the two input vectors, cosine distance can be used to compare two
spectra with unequal total mutation counts (even if those total counts are relatively small).
Additionally, by calculating the cosine distance between mutation spectra, we avoid the need to
perform separate comparisons of mutation counts at each individual k-mer mutation type.

Inspired by methods from QTL mapping [54,58], we use permutation tests to establish genome-
wide cosine distance thresholds. In each of N permutation trials, we randomly shuffle the per-
haplotype mutation data such that haplotype labels no longer correspond to the correct mutation
counts. Using the shuffled mutation data, we perform a genome-wide scan as described above,
and record the maximum cosine distance observed at any locus. After N permutations (usually
10,000), we compute the $1 - p$ percentile of the distribution of maximum statistics, and use that
percentile value as a genome-wide significance threshold (for example, at $p = 0.05$).

Estimating confidence intervals around AMSD peaks

If we identified an adjusted cosine distance peak on a particular chromosome, we used a
bootstrap resampling approach [59] to estimate confidence intervals. In each of $N = 10,000$
trials, we resampled the mutation spectrum data and corresponding marker genotypes (on the
chromosome of interest) with replacement. Using those resampled spectra and genotypes, we
performed an aggregate mutation spectrum distance scan on the chromosome of interest and
recorded the position of the marker with the largest adjusted cosine distance value. We then
defined a 90% confidence interval by finding two marker locations between which 90% of all N
bootstrap samples produced a peak cosine distance value. In other words, we estimated the
bounds of the 90% confidence interval by finding the markers that defined the 5th and 95th
percentiles of the distribution of maximum adjusted cosine distance values across N bootstrap
trials. We note, however, that the bootstrap can exhibit poor performance in QTL mapping
studies [26]; namely, bootstrap confidence intervals tend to be larger than those estimated using
either a "LOD drop" method or a Bayes credible interval, and can exhibit poorer-than-expected
coverage (a measure of whether the confidence interval contains the true QTL location).

Accounting for relatedness between strains

We expect each BXD to derive approximately 50% of its genome from C57BL/6J and 50% from DBA/2J. As a result, every pair of BXDs will likely have identical genotypes at a fraction of markers. Pairs of more genetically similar BXDs may also have more similar mutation spectra, potentially due to shared polygenic effects on the mutation process. Therefore, at a given marker, if the BXDs that inherited *D* alleles are more genetically dissimilar from those that inherited *B* alleles (considering all loci throughout the genome in our measurement of genetic similarity), we might expect the aggregate mutation spectra in the two groups to also be more dissimilar.

We implemented a simple approach to account for these potential issues of relatedness. At each marker g_i, we divide BXD haplotypes into two groups based on the parental allele they inherited. As before, we first compute the aggregate mutation spectrum in each group of haplotypes and calculate the cosine distance between the two aggregate spectra (D_i^C). Then, within each group of haplotypes, we calculate the allele frequency of the *D* allele at every marker along the genome to obtain a vector of length n, where n is the number of genotyped markers. To quantify the genetic similarity between the two groups of haplotypes, we calculate the Pearson correlation coefficient r_i between the two vectors of marker-wide *D* allele frequencies.

Put another way, at every marker g_i along the genome, we divide BXD haplotypes into two groups and compute two metrics: D_i^C (the cosine distance between the two groups' aggregate spectra) and r_i (the correlation between genome-wide *D* allele frequencies in the two groups). To control for the potential effects of genetic similarity on cosine distances, we regress $\left(D_1^C, D_2^C, \dots D_n^C\right)$ on $(r_1, r_2, \dots r_n)$ for all n markers using an ordinary least-squares model. We then use the residuals from the fitted model as the "adjusted" cosine distance values for each marker. If genome-wide genetic similarity between haplotypes perfectly predicts cosine distances at each marker, these residuals will all be 0 (or very close to 0). If genome-wide genetic similarity has no predictive power, the residuals will simply represent the difference between the observed cosine distance at a single marker and the marker-wide mean of cosine distances.

Accounting for BXD population structure due to breeding epochs

The current BXD family was generated in six breeding "epochs." As discussed previously, each epoch was initiated with a distinct cross of C57BL/6J and DBA/2J parents; BXDs in four of the epochs were generated following F2 crosses of C57BL/6J and DBA/2J, and BXDs in the other two were generated following advanced intercrosses. Due to this breeding approach the BXD epochs differ from each other in a few important ways. For example, BXDs derived in epochs 3 and 5 (i.e., from advanced intercross) harbor larger numbers of fixed recombination breakpoints than those from epochs 1, 2, 4, and 6 [3]. Although the C57BL/6J and DBA/2J parents used to initialize each epoch were completely inbred, they each possessed a small number unique *de novo* germline mutations that were subsequently inherited by many of their offspring. A number of these "epoch-specific" variants have also been linked to phenotypic variation observed between BXDs from different epochs [3,60,61,62].

To account for potential population structure, as well as these epoch-specific effects, we
introduced the ability to perform stratified permutation tests in the aggregate mutation spectrum
distance approach. Normally, in each of N permutations we shuffle the per-haplotype mutation
spectrum data such that haplotype labels no longer correspond to the correct mutation spectra
(i.e., shuffle mutation spectra *across* epochs). In the stratified approach, we instead shuffle per-
haplotype mutation data *within* epochs, preserving epoch structure while still enabling mutation
spectra permutations. We used this epoch-aware approach for all permutation tests presented
in this manuscript.

Implementation and source code

The aggregate mutation spectrum distance method was implemented in Python, and relies
heavily on the following Python libraries: `numpy`, `pandas`, `matplotlib`, `scikit-learn`,
`pandera`, `seaborn`, and `numba` [63,64,65,66,67,68,69].

The code underlying AMSD, as well as documentation of the method, is available on GitHub
(https://github.com/quinlan-lab/proj-mutator-mapping). We have also deposited a reproducible
Snakemake [70] workflow for running reproducing all analyses and figures presented in the
manuscript.

Simulations to assess the power of the aggregate mutation spectrum distance approach

681 We performed a series of simple simulations to estimate our power to detect alleles that affect
682 the germline mutation spectrum using the aggregate mutation spectrum distance method.

Simulating genotypes

First, we simulate genotypes on a population of haplotypes at a collection of sites. We define a
matrix G of size (s, h), where s is the number of sites and h is the number of haplotypes. We
assume that every site is biallelic, and that the minor allele frequency at every site is 0.5. For
every entry $G_{i,j}$, we take a single draw from a uniform distribution in the interval [0.0,1.0]. If the
value of that draw is less than 0.5, we assign the value of $G_{i,j}$ to be 1. Otherwise, we assign the
value of $G_{i,j}$ to be 0.

Defining expected mutation type probabilities

691 Next, we define a vector of 1-mer mutation probabilities:

$$P = (0.29,\ 0.17,\ 0.12,\ 0.075,\ 0.1,\ 0.075,\ 0.17)$$

These probabilities sum to 1 and roughly correspond to the expected frequencies of C>T,
CpG>TpG, C>A, C>G, A>T, A>C, and A>G *de novo* germline mutations in mice, respectively
[9]. If we are simulating the 3-mer mutation spectrum, we modify the vector of mutation
probabilities P to be length 96, and assign every 3-mer mutation type a value of $\frac{P_c}{16}$, where P_c is

the probability of the "central" mutation type associated with the 3-mer mutation type. In other
words, each of the 16 possible NCN>NTN 3-mer mutation types would be assigned a mutation
probability of $\frac{P_c}{16} = \frac{0.46}{16} = 0.02875$. We then generate a vector of lambda values by scaling the
mutation probabilities by the number of mutations we wish to simulate (m):

$$\lambda = Pm$$

We also create a second vector of lambda values (λ'), in which we multiply the λ value of a
single mutation type by the mutator effect size e.

Rather than simulate the same mean number of mutations (m) on every haplotype, we also
performed a series of simulations in which the mean number of mutations on each haplotype
was allowed to vary. The BXD RILs were inbred for variable numbers of generations, and each
BXD therefore accumulated a variable number of *de novo* germline mutations [2]. To more
closely approximate the BXD haplotypes, we performed simulations in which the number of
mutations (m) on each haplotype was drawn from a uniform distribution from m to $20m$. In other
words, we created a vector of mutation counts M containing h evenly-spaced integers from m to
$20m$, where h is the number of simulated haplotypes. Thus, if we simulated between 100 and
2,000 mutations on 50 haplotypes, the ith entry of M would be $100 + \frac{(2,000-100)}{50}i$. Each
haplotype's mean number of mutations was then assigned by looking up the haplotype's index i
in M.

In our simulations, we assume that genotypes at a single site (the "mutator locus") are
associated with variation in the mutation spectrum. That is, at a single site s_i, all of the
haplotypes with 1 alleles should have elevated rates of a particular mutation type and draw their
mutation counts from λ', while all of the haplotypes with 0 alleles should have "wild-type" rates
of that mutation type and draw their mutation counts from λ. We therefore pick a random site s_i
to be the "mutator locus," and identify the indices of haplotypes in G that were assigned 1 alleles
at s_i. We call these indices h_{mut}.

Simulating mutation spectra

To simulate the mutation spectrum on our toy population of haplotypes, we define a matrix C of
size (h, n), where $n = 6 \times 4^{k-1}$ (or if $k = 1$ and we include CpG>TpG mutations, $6 \times 4^{k-1} + 1$).

Then, we populate the matrix C separately for *mutator* and *wild-type* haplotypes. For every row i
in the matrix (i.e., for every haplotype), we first ask if i is in h_{mut} (that is, if the haplotype at
index i was assigned a 1 allele at the "mutator locus"). If so, we set the values of C_i to be the
results of a single Poisson draw from λ'. If row i is not in h_{mut}, we set the values of C_i to be the
results of a single Poisson draw from λ.

Assessing power to detect a simulated mutator allele using AMSD

For each combination of parameters (number of simulated haplotypes, number of simulated
markers, mutator effect size, etc.), we run 100 independent trials. In each trial, we simulate the
genotype matrix G and the mutation counts C. We calculate a "focal" cosine distance as the

cosine distance between the aggregate mutation spectra of haplotypes with either genotype at
s_i (the site at which we artificially simulated an association between genotypes and mutation
spectrum variation). We then perform an aggregate mutation spectrum distance scan using $N =$
1,000 permutations. If fewer than 5% of the N permutations produced a cosine distance greater
than or equal to the focal distance, we say that the approach successfully identified the mutator
allele in that trial.

Assessing power to detect a simulated mutator allele using quantitative trait locus (QTL) mapping

Using simulated data, we also assessed the power of traditional quantitative trait locus (QTL)
mapping to detect a locus associated with mutation spectrum variation. As described above, we
simulated both genotype and mutation spectra for a population of haplotypes under various
conditions (number of mutations per haplotype, mutator effect size, etc.). Using those simulated
data, we used R/qtl2 [54] to perform a genome scan for significant QTL as follows; we assume
that the simulated genotype markers are evenly spaced (in physical Mbp coordinates) on a
single chromosome. First, we calculate the fraction of each haplotype's *de novo* mutations that
belong to each of the $6 \times 4^{k-1}$ possible k-mer mutation types. We then convert the simulated
genotypes at each marker to genotype probabilities using the `calc_genoprob` function in
R/qtl2, with `map_function = "c-f"` and `error_prob = 0`. For every k-mer mutation type,
we use genotype probabilities and per-haplotype mutation fractions to perform a scan for QTL
with the `scan1` function; to make the results more comparable to those from the AMSD method,
we do not include any covariates or kinship matrices in these QTL scans. We then use the
`scan1perm` function to perform 1,000 permutations of the per-haplotype mutation fractions and
calculate log-odds (LOD) thresholds for significance. We consider the QTL scan to be
"successful" if it produces a LOD score above the significance threshold (defined using $\alpha = \frac{0.05}{7}$)
for the marker at which we simulated an association with mutation spectrum variation.

In our simulations, we augment the mutation rate of a single k-mer mutation type on haplotypes
carrying the simulated mutator allele. However, in an experimental setting, we would not expect
to have *a priori* knowledge of the mutation type affected by the mutator. Thus, by using an alpha
threshold of 0.05 in our simulations, we would likely over-estimate the power of QTL mapping
for detecting the mutator. Since we would need to perform 7 separate QTL scans (one for each
1-mer mutation type plus CpG>TpG) in an experimental setting, we calculate QTL LOD
thresholds at a Bonferroni-corrected alpha value of $\alpha = \frac{0.05}{7}$.

Applying the aggregate mutation spectrum distance method to the BXDs

We downloaded previously-generated BXD *de novo* germline mutation data from the GitHub
repository associated with our previous manuscript, which was also archived at Zenodo
[2,71,72], and downloaded a CSV file of BXD genotypes at ~7,300 informative markers from
GeneNetwork [27,73]. We also downloaded relevant metadata about each BXD from the

manuscript describing the updated BXD resource [3]. These files are included in the GitHub
repository associated with this manuscript.

As in our previous manuscript [2], we included mutation data from a subset of the 152 BXDs in
our aggregate mutation spectrum distance scans. Specifically, we removed BXDs that were
backcrossed to a C57BL/6J or DBA/2J parent at any point during the inbreeding process
(usually, in order to rescue that BXD from inbreeding depression [3]). We also removed BXD68
from our genome-wide scans, since we previously discovered a hyper-mutator phenotype in that
line; the C>A germline mutation rate in BXD68 is over 5 times the population mean, likely due to
a private deleterious nonsynonymous mutation in *Mutyh* [2]. In our previous manuscript, we
removed any BXDs that had been inbred for fewer than 20 generations, as it takes
approximately 20 generations of strict brother-sister mating for an RIL genome to become >98%
homozygous [74]. As a result, any potential mutator allele would almost certainly be either fixed
or lost after 20 generations; if fixed, the allele would remain linked to any excess mutations it
causes for the duration of subsequent inbreeding. In other words, the *de novo* mutations
present in the genome of a "young" BXD (i.e., a BXD that was inbred for fewer than 20
generations) would not reflect a mutator allele's activity as strongly as the mutations present in
the genome of a much older BXD. This presented a challenge when we used quantitative trait
locus mapping to discover mutator alleles in our previous manuscript, since the phenotypes
(i.e., C>A mutation rates) of young and old BXDs were weighted equally; thus, we simply
removed the younger BXDs from our analysis to avoid using their especially noisy mutation
spectra. Since AMSD computes an *aggregate* mutation spectrum using all BXDs that inherited a
particular allele at a locus, and can overcome the sparsity and noise of individual mutation
spectra, we chose to include these younger BXDs in our genome-wide scans in this study.

In total, we included 117 BXDs in our genome-wide scans.

Identifying candidate single-nucleotide mutator alleles overlapping the chromosome 6 peak

We investigated the region implicated by our aggregate mutation spectrum distance approach
on chromosome 6 by subsetting the joint-genotyped BXD VCF file (European Nucleotide
Archive accession PRJEB45429 [75]) using `bcftools` [76]. We defined the candidate interval
surrounding the cosine distance peak on chromosome 6 as the 90% bootstrap confidence
interval (extending from approximately 95 Mbp to 114 Mbp). To predict the functional impacts of
both single-nucleotide variants and indels on splicing, protein structure, etc., we annotated
variants in the BXD VCF using the following `snpEff` [77] command:

```
805  java -Xmx16g -jar /path/to/snpeff/jarfile GRCm38.75 /path/to/bxd/vcf
806  > /path/to/uncompressed/output/vcf
```

807 and used `cyvcf2` [78] to iterate over the annotated VCF file in order to identify nonsynonymous
808 fixed differences between the parental C57BL/6J and DBA/2J strains.

Identifying candidate structural variant alleles overlapping the chromosome 6 peak

We downloaded summary VCFs containing insertion, deletion and inversion structural variants (identified via high-quality, long-read assembly of inbred laboratory mouse strains [28]) from the Zenodo link associated with the Ferraj et al. manuscript: https://doi.org/10.5281/zenodo.7644286.

We then downloaded a TSV file containing RefSeq gene predictions in GRCm39/mm39 from the UCSC Table Browser [79], and used the `bx-python` library [80] to intersect the interval spanned by each structural variant with the intervals spanned by the `txStart` and `txEnd` of every RefSeq entry. We queried all structural variants within the 90% bootstrap confidence interval on chromosome 6.

Extracting mutation signatures

We used SigProfilerExtractor (v.1.1.21) [30] to extract mutation signatures from the BXD mutation data. After converting the BXD mutation data to the "matrix" input format expected by SigProfilerExtractor, we ran the `sigProfilerExtractor` method as follows:

```
# install the mm10 mouse reference data
genInstall.install('mm10')

# run mutation signature extraction
sig.sigProfilerExtractor(
    'matrix',
    /path/to/output/directory,
    /path/to/input/mutations,
    maximum_signatures=10,
    nmf_replicates=100,
    opportunity_genome="mm10",
)
```

Comparing mutation spectra between Mouse Genomes Project strains

We downloaded mutation data from a previously published analysis [18] (Supplementary File 1, Excel Table S3) that identified strain-private mutations in 29 strains that were originally whole-genome sequenced as part of the Sanger Mouse Genomes (MGP) project [31]. When comparing counts of each mutation type between MGP strains that harbored either *D* or *B* alleles at the chromosome 4 or chromosome 6 mutator loci, we adjusted mutation counts by the number of callable A, T, C, or G nucleotides in each strain as described previously [2].

Querying GeneNetwork for eQTLs at the mutator locus

We used the online GeneNetwork resource [27], which contains array- and RNA-seq-derived expression measurements in a wide variety of tissues, to find *cis*-eQTLs for the DNA repair genes we implicated under the cosine distance peak on chromosome 6. On the GeneNetwork homepage (genenetwork.org), we selected the "BXD Family" **Group** and used the **Type** dropdown menu to select each of the specific expression datasets described in Table 2. In the **Get Any** text box, we then entered the listed gene name and clicked **Search**. After selecting the appropriate trait ID on the next page, we used the **Mapping Tools** dropdown to run Hayley-Knott regression [81] with default parameters: 1,000 permutations, interval mapping, no cofactors, and WGS-based genotypes (2022).

If we discovered a significant cis-eQTL for the gene of interest (that is, a locus on chromosome 6 with an LRS greater than or equal to the "significant LRS" genome-wide threshold), we then performed a second genome-wide association test for the trait of interest using GEMMA [82] with the following parameters: WGS-based marker genotypes, a minor allele frequency threshold of 0.05, and leave-one-chromosome-out (LOCO). By using both Haley-Knott regression and GEMMA, we could first discover loci that exceeded a genome-wide LRS threshold, and then more precisely estimate the effect of those loci on gene expression [83].

The exact names of the expression datasets we used for each tissue are shown in Table 2 below:

Table 2: Names of gene expression datasets used for each tissue type on GeneNetwork

Tissue name	Complete name of GeneNetwork expression data
Kidney	Mouse kidney M430v2 Sex Balanced (Aug06) RMA
Gastrointestinal	UTHSC Mouse BXD Gastrointestinal Affy MoGene 1.0 ST Gene Level (Apr14) RMA
Hematopoetic stem cells	UMCG Stem Cells ILM6v1.1 (Apr09) transformed
Spleen	UTHSC Affy MoGene 1.0 ST Spleen (Dec10) RMA
Liver	UTHSC BXD Liver RNA-Seq Avg (Oct19) TPM Log2
Heart	NHLBI BXD All Ages Heart RNA-Seq (Nov20) TMP Log2 **
Hippocampus	Hippocampus Consortium M430v2 (Jun06) RMA

Calculating the frequencies of candidate mutator alleles in wild mice

To determine the frequencies of the *Ogg1* and *Setmar* nonsynonymous mutations in other populations of mice, we queried a VCF file containing genome-wide variation in 67 wild-derived mice from four species of *Mus* [32]. We calculated the allele frequency of each nonsynonymous mutation in each of the four species or subspecies (*Mus musculus domesticus*, *Mus musculus*

870 *musculus*, *Mus musculus castaneus*, and *Mus spretus*), including genotypes that met the
871 following criteria:

- • supported by at least 10 sequencing reads

- • Phred-scaled genotype quality of at least 20

Testing for epistasis between the two mutator loci

875 To test for statistical epistasis between the mutator loci on chromosome 4 and chromosome 6,
876 we modeled C>A mutation rates in the BXDs as a function of genotypes at either locus:

877
$$\log \lambda^i = \beta_0 + \beta_1 X_1^i + \beta_2 X_2^i + \log A^i$$

Here, λ^i represents the count of C>A mutations in the ith BXD. We assume that the count of
C>A mutations follows a Poisson distribution with a mean equal to the C>A mutation rate
(expressed per base pair, per generation) multiplied by product of two terms: the total number of
generations of inbreeding and the total number of base pairs accessible to variant calling. The
BXDs differ in both their durations of inbreeding and the proportions of their genomes that were
sequenced to sufficient depth, which influences the number of mutations we observe in each
BXD. We therefore model C>A mutation counts as *rates* by including a $log\,A^i$ term – sometimes
referred to as an "offset" – which represents the product of the number of "callable"
cytosine/guanine nucleotides in each BXD (i.e., the total number of cytosines/guanines covered
by at least 10 sequencing reads) and the number of generations for which the BXD was inbred.
The offset is assumed to have a coefficient of 1. The X_1^i and X_2^i terms represent BXD genotypes
in the ith BXD at markers `rs27509845` and `rs46276051` (the markers with peak cosine
distances on chromosomes 4 and 6 in the two aggregate mutation spectrum distance scans); *B*
genotypes are coded as 0 and *D* genotypes are coded as 1. We limited our analysis to the *n* =
108 BXDs that were homozygous at both loci, allowing us to model genotypes at either locus as
binary categorical variables. In the R statistical language, we defined the model as follows:

894
```
m1 <- glm(Count ~ offset(log(ADJ_AGE)) + Genotype_A + Genotype_B, data
895  = data, family=poisson())
```

In this model, `Count` is the count of C>A *de novo* mutations observed in each BXD. The
`Genotype_A` and `Genotype_B` terms are analogous to the X_1 and X_2 terms in the
mathematical model above, and `ADJ_AGE` is analogous to the A term.

To explicitly test if the effects of *D* genotypes at the chromosome 6 locus depended on the
presence of *D* genotypes at the chromosome 4 locus, we fit a second model incorporating a
multiplicative interaction between X_1 and X_2:

902
$$\log \lambda^i = \beta_0 + \beta_1 X_1^i + \beta_2 X_2^i + \beta_{12} X_1^i X_2^i + log\,A^i$$

Here, the β_{12} term capture the interaction effect between genotypes at the two mutator loci. If
both X_1^i and X_2^i are equal to 1 (i.e., genotypes at both the chromosome 4 and chromosome 6

loci are *D*), the β_{12} term will capture the resulting effect on C>A mutation counts. We can
interpret a significantly non-zero β_{12} value as evidence for a non-additive effect of *D* genotypes
at both loci.

We defined this model in the R statistical language as follows:

```
m2 <- glm(Count ~ offset(log(ADJ_AGE)) + Genotype_A * Genotype_B, data
910 = data, family=poisson())
```

911 An identical model can also be written by explicitly specifying both the additive and interaction
912 effects:

913 ```
m2 <- glm(Count ~ offset(log(ADJ_AGE)) + Genotype_A + Genotype_B +
914 Genotype_A:Genotype_B, data = data, family=poisson())
```

915 Using analysis of variance (ANOVA), we compared the model incorporating an interaction effect
916 to the one with only additive effects:

917 ```
anova(m2, m1, test="Chisq")
```

918 If model `m2` is a significantly better fit to the data than `m1`, we can reject the null hypothesis that
919 the effect of *D* genotypes at both markers is equal to the sum of the marginal effects of *D*
920 genotypes at either `rs27509845` or `rs46276051`. In other words, if `m2` is a better fit than `m1`,
921 then the combined effect of *D* genotypes at both markers is non-additive, and indicative of
922 statistical epistasis.

923 We tested for statistical epistasis in the Sanger Mouse Genomes Project (MGP) strains using a
924 nearly identical approach. In this analysis, we fit two models as follows:

925 ```
m1 <- glm(Count ~ offset(log(CALLABLE_C)) + Genotype_A * Genotype_B,
926 data = data, family=poisson())
927
928 m2 <- glm(Count ~ offset(log(CALLABLE_C)) + Genotype_A + Genotype_B,
929 data = data, family=poisson())
```

930 where `Count` is the count of strain-private C>A mutations observed in each MGP strain [18].
931 The `CALLABLE_C` term represents the total number of cytosine and guanine nucleotides that
932 were accessible for mutation calling in each strain, and the `Genotype_A` and `Genotype_B`
933 terms represent MGP genotypes at the chromosome 4 and chromosome 6 mutator loci,
934 respectively. We compared the two models using ANOVA as described above.

## Acknowledgments

We thank Robert W. Williams (University of Tennessee Health Sciences Center) and Don F. Conrad (Oregon Health & Science University) for very helpful comments and feedback on a draft of this manuscript.

# References

1. **Mechanisms of DNA damage, repair, and mutagenesis.** Nimrat Chatterjee, Graham C Walker *Environmental and molecular mutagenesis* (2017-05-09) https://www.ncbi.nlm.nih.gov/pubmed/28485537 DOI: 10.1002/em.22087 · PMID: 28485537 · PMCID: PMC5474181

2. **A natural mutator allele shapes mutation spectrum variation in mice.** Thomas A Sasani, David G Ashbrook, Annabel C Beichman, Lu Lu, Abraham A Palmer, Robert W Williams, Jonathan K Pritchard, Kelley Harris *Nature* (2022-05-11) https://www.ncbi.nlm.nih.gov/pubmed/35545679 DOI: 10.1038/s41586-022-04701-5 · PMID: 35545679 · PMCID: PMC9272728

3. **A platform for experimental precision medicine: The extended BXD mouse family.** David G Ashbrook, Danny Arends, Pjotr Prins, Megan K Mulligan, Suheeta Roy, Evan G Williams, Cathleen M Lutz, Alicia Valenzuela, Casey J Bohl, Jesse F Ingels, … Robert W Williams *Cell systems* (2021-01-19) https://www.ncbi.nlm.nih.gov/pubmed/33472028 DOI: 10.1016/j.cels.2020.12.002 · PMID: 33472028 · PMCID: PMC7979527

4. **Base-excision repair of oxidative DNA damage.** Sheila S David, Valerie L O'Shea, Sucharita Kundu *Nature* (2007-06-21) https://www.ncbi.nlm.nih.gov/pubmed/17581577 DOI: 10.1038/nature05978 · PMID: 17581577 · PMCID: PMC2896554

5. **Parental influence on human germline de novo mutations in 1,548 trios from Iceland.** Hákon Jónsson, Patrick Sulem, Birte Kehr, Snaedis Kristmundsdottir, Florian Zink, Eirikur Hjartarson, Marteinn T Hardarson, Kristjan E Hjorleifsson, Hannes P Eggertsson, Sigurjon Axel Gudjonsson, … Kari Stefansson *Nature* (2017-09-20) https://www.ncbi.nlm.nih.gov/pubmed/28959963 DOI: 10.1038/nature24018 · PMID: 28959963

6. **Large, three-generation human families reveal post-zygotic mosaicism and variability in germline mutation accumulation.** Thomas A Sasani, Brent S Pedersen, Ziyue Gao, Lisa Baird, Molly Przeworski, Lynn B Jorde, Aaron R Quinlan *eLife* (2019-09-24) https://www.ncbi.nlm.nih.gov/pubmed/31549960 DOI: 10.7554/elife.46922 · PMID: 31549960 · PMCID: PMC6759356

7. **De novo Mutations in Domestic Cat are Consistent with an Effect of Reproductive Longevity on Both the Rate and Spectrum of Mutations.** Richard J Wang, Muthuswamy Raveendran, RAlan Harris, William J Murphy, Leslie A Lyons, Jeffrey Rogers, Matthew W Hahn *Molecular biology and evolution* (2022-07-02) https://www.ncbi.nlm.nih.gov/pubmed/35771663 DOI: 10.1093/molbev/msac147 · PMID: 35771663 · PMCID: PMC9290555

8. **A comparison of humans and baboons suggests germline mutation rates do not track cell divisions.** Felix L Wu, Alva I Strand, Laura A Cox, Carole Ober, Jeffrey D Wall, Priya Moorjani, Molly Przeworski *PLoS biology* (2020-08-17) https://www.ncbi.nlm.nih.gov/pubmed/32804933 DOI: 10.1371/journal.pbio.3000838 · PMID: 32804933 · PMCID: PMC7467331

9. **Similarities and differences in patterns of germline mutation between mice and humans.** Sarah J Lindsay, Raheleh Rahbari, Joanna Kaplanis, Thomas Keane, Matthew E Hurles *Nature communications* (2019-09-06) https://www.ncbi.nlm.nih.gov/pubmed/31492841
DOI: 10.1038/s41467-019-12023-w · PMID: 31492841 · PMCID: PMC6731245

10. **Timing, rates and spectra of human germline mutation.** Raheleh Rahbari, Arthur Wuster, Sarah J Lindsay, Robert J Hardwick, Ludmil B Alexandrov, Saeed Al Turki, Anna Dominiczak, Andrew Morris, David Porteous, Blair Smith, … Matthew E Hurles *Nature genetics* (2015-12-14) https://www.ncbi.nlm.nih.gov/pubmed/26656846 DOI: 10.1038/ng.3469 · PMID: 26656846 · PMCID: PMC4731925

11. **Genetic drift, selection and the evolution of the mutation rate.** Michael Lynch, Matthew S Ackerman, Jean-Francois Gout, Hongan Long, Way Sung, WKelley Thomas, Patricia L Foster *Nature reviews. Genetics* (2016-10-14)
https://www.ncbi.nlm.nih.gov/pubmed/27739533 DOI: 10.1038/nrg.2016.104 · PMID: 27739533

12. **De novo mutations across 1,465 diverse genomes reveal mutational insights and reductions in the Amish founder population.** Michael D Kessler, Douglas P Loesch, James A Perry, Nancy L Heard-Costa, Daniel Taliun, Brian E Cade, Heming Wang, Michelle Daya, John Ziniti, Soma Datta, … Timothy D O'Connor *Proceedings of the National Academy of Sciences of the United States of America* (2020-01-21) https://www.ncbi.nlm.nih.gov/pubmed/31964835
DOI: 10.1073/pnas.1902766117 · PMID: 31964835 · PMCID: PMC7007577

13. **A naturally occurring variant of<i>MBD4</i>causes maternal germline hypermutation in primates** Alexandra M Stendahl, Rashesh Sanghvi, Samuel Peterson, Karina Ray, Ana C Lima, Raheleh Rahbari, Donald F Conrad *Cold Spring Harbor Laboratory* (2023-03-29) https://doi.org/gr3ghz DOI: 10.1101/2023.03.27.534460

14. **Genetic and chemotherapeutic influences on germline hypermutation.** Joanna Kaplanis, Benjamin Ide, Rashesh Sanghvi, Matthew Neville, Petr Danecek, Tim Coorens, Elena Prigmore, Patrick Short, Giuseppe Gallone, Jeremy McRae, … Matthew Hurles *Nature* (2022-05-11) https://www.ncbi.nlm.nih.gov/pubmed/35545669 DOI: 10.1038/s41586-022-04712-2 · PMID: 35545669 · PMCID: PMC9117138

15. **Increased somatic mutation burdens in normal human cells due to defective DNA polymerases.** Philip S Robinson, Tim HH Coorens, Claire Palles, Emily Mitchell, Federico Abascal, Sigurgeir Olafsson, Bernard CH Lee, Andrew RJ Lawson, Henry Lee-Six, Luiza Moore, … Michael R Stratton *Nature genetics* (2021-09-30)
https://www.ncbi.nlm.nih.gov/pubmed/34594041 DOI: 10.1038/s41588-021-00930-y · PMID: 34594041 · PMCID: PMC8492474

16. **Germline de novo mutations in families with Mendelian cancer syndromes caused by defects in DNA repair.** Kitty Sherwood, Joseph C Ward, Ignacio Soriano, Lynn Martin, Archie Campbell, Raheleh Rahbari, Ioannis Kafetzopoulos, Duncan Sproul, Andrew Green, Julian R Sampson, … Ian Tomlinson *Nature communications* (2023-06-19)
https://www.ncbi.nlm.nih.gov/pubmed/37336879 DOI: 10.1038/s41467-023-39248-0 · PMID: 37336879 · PMCID: PMC10279637

17.    **Inference of Candidate Germline Mutator Loci in Humans from Genome-Wide**
**Haplotype Data.** Cathal Seoighe, Aylwyn Scally *PLoS genetics* (2017-01-17)
https://www.ncbi.nlm.nih.gov/pubmed/28095480 DOI: 10.1371/journal.pgen.1006549 · PMID:
28095480 · PMCID: PMC5283766

18.    **Significant Strain Variation in the Mutation Spectra of Inbred Laboratory Mice.**
Beth L Dumont *Molecular biology and evolution* (2019-05-01)
https://www.ncbi.nlm.nih.gov/pubmed/30753674 DOI: 10.1093/molbev/msz026 · PMID:
30753674 · PMCID: PMC6501876

19.    **Spontaneous Mutation Accumulation Studies in Evolutionary Genetics** Daniel L
Halligan, Peter D Keightley *Annual Review of Ecology, Evolution, and Systematics* (2009-12-01)
https://doi.org/dvrjz8 DOI: 10.1146/annurev.ecolsys.39.110707.173437

20.    **Inherited MUTYH mutations cause elevated somatic mutation rates and distinctive**
**mutational signatures in normal human cells.** Philip S Robinson, Laura E Thomas, Federico
Abascal, Hyunchul Jung, Luke MR Harvey, Hannah D West, Sigurgeir Olafsson, Bernard CH
Lee, Tim HH Coorens, Henry Lee-Six, … Michael R Stratton *Nature communications* (2022-07-
08) https://www.ncbi.nlm.nih.gov/pubmed/35803914 DOI: 10.1038/s41467-022-31341-0 · PMID:
35803914 · PMCID: PMC9270427

21.    **Inferring evolutionary dynamics of mutation rates through the lens of mutation**
**spectrum variation.** Jedidiah Carlson, William S DeWitt, Kelley Harris *Current opinion in*
*genetics & development* (2020-06-30) https://www.ncbi.nlm.nih.gov/pubmed/32619789 DOI:
10.1016/j.gde.2020.05.024 · PMID: 32619789 · PMCID: PMC7646088

22.    **A Specific Mutational Signature Associated with DNA 8-Oxoguanine Persistence**
**in MUTYH-defective Colorectal Cancer.** Alessandra Viel, Alessandro Bruselles, Ettore
Meccia, Mara Fornasarig, Michele Quaia, Vincenzo Canzonieri, Eleonora Policicchio, Emanuele
Damiano Urso, Marco Agostini, Maurizio Genuardi, … Margherita Bignami *EBioMedicine* (2017-
04-13) https://www.ncbi.nlm.nih.gov/pubmed/28551381 DOI: 10.1016/j.ebiom.2017.04.022 ·
PMID: 28551381 · PMCID: PMC5478212

23.    **Mutational signature analysis identifies MUTYH deficiency in colorectal cancers**
**and adrenocortical carcinomas.** Camilla Pilati, Jayendra Shinde, Ludmil B Alexandrov,
Guillaume Assié, Thierry André, Zofia Hélias-Rodzewicz, Romain Ducoudray, Delphine Le
Corre, Jessica Zucman-Rossi, Jean-François Emile, … Pierre Laurent-Puig *The Journal of*
*pathology* (2017-03-29) https://www.ncbi.nlm.nih.gov/pubmed/28127763 DOI:
10.1002/path.4880 · PMID: 28127763

24.    **Gene ontology: tool for the unification of biology. The Gene Ontology Consortium.**
M Ashburner, CA Ball, JA Blake, D Botstein, H Butler, JM Cherry, AP Davis, K Dolinski, SS
Dwight, JT Eppig, … G Sherlock *Nature genetics* (2000-05)
https://www.ncbi.nlm.nih.gov/pubmed/10802651 DOI: 10.1038/75556 · PMID: 10802651 ·
PMCID: PMC3037419

25. **The Gene Ontology resource: enriching a GOld mine.** *Nucleic acids research* (2021-01-08) https://www.ncbi.nlm.nih.gov/pubmed/33290552 DOI: 10.1093/nar/gkaa1113 · PMID: 33290552 · PMCID: PMC7779012

26. **Poor performance of bootstrap confidence intervals for the location of a quantitative trait locus.** Ani Manichaikul, Josée Dupuis, Saunak Sen, Karl W Broman *Genetics* (2006-06-18) https://www.ncbi.nlm.nih.gov/pubmed/16783000 DOI: 10.1534/genetics.106.061549 · PMID: 16783000 · PMCID: PMC1569776

27. **GeneNetwork: A Toolbox for Systems Genetics.** Megan K Mulligan, Khyobeni Mozhui, Pjotr Prins, Robert W Williams *Methods in molecular biology (Clifton, N.J.)* (2017) https://www.ncbi.nlm.nih.gov/pubmed/27933521 DOI: 10.1007/978-1-4939-6427-7_4 · PMID: 27933521 · PMCID: PMC7495243

28. **Resolution of structural variation in diverse mouse genomes reveals chromatin remodeling due to transposable elements** Ardian Ferraj, Peter A Audano, Parithi Balachandran, Anne Czechanski, Jacob I Flores, Alexander A Radecki, Varun Mosur, David S Gordon, Isha A Walawalkar, Evan E Eichler, … Christine R Beck *Cell Genomics* (2023-05) https://doi.org/gr3t7b DOI: 10.1016/j.xgen.2023.100291 · PMID: 37228752

29. **Uncovering novel mutational signatures by** SMAshiqul Islam, Marcos Díaz-Gay, Yang Wu, Mark Barnes, Raviteja Vangara, Erik N Bergstrom, Yudou He, Mike Vella, Jingwei Wang, Jon W Teague, … Ludmil B Alexandrov *Cell genomics* (2022-11-09) https://www.ncbi.nlm.nih.gov/pubmed/36388765 DOI: 10.1016/j.xgen.2022.100179 · PMID: 36388765 · PMCID: PMC9646490

30. **COSMIC: the Catalogue Of Somatic Mutations In Cancer.** John G Tate, Sally Bamford, Harry C Jubb, Zbyslaw Sondka, David M Beare, Nidhi Bindal, Harry Boutselakis, Charlotte G Cole, Celestino Creatore, Elisabeth Dawson, … Simon A Forbes *Nucleic acids research* (2019-01-08) https://www.ncbi.nlm.nih.gov/pubmed/30371878 DOI: 10.1093/nar/gky1015 · PMID: 30371878 · PMCID: PMC6323903

31. **Mouse genomic variation and its effect on phenotypes and gene regulation.** Thomas M Keane, Leo Goodstadt, Petr Danecek, Michael A White, Kim Wong, Binnaz Yalcin, Andreas Heger, Avigail Agam, Guy Slater, Martin Goodson, … David J Adams *Nature* (2011-09-14) https://www.ncbi.nlm.nih.gov/pubmed/21921910 DOI: 10.1038/nature10413 · PMID: 21921910 · PMCID: PMC3276836

32. **Genomic resources for wild populations of the house mouse, Mus musculus and its close relative Mus spretus.** Bettina Harr, Emre Karakoc, Rafik Neme, Meike Teschke, Christine Pfeifle, Željka Pezer, Hiba Babiker, Miriam Linnenbrink, Inka Montero, Rick Scavetta, … Diethard Tautz *Scientific data* (2016-09-13) https://www.ncbi.nlm.nih.gov/pubmed/27622383 DOI: 10.1038/sdata.2016.75 · PMID: 27622383 · PMCID: PMC5020872

33. **On the subspecific origin of the laboratory mouse.** Hyuna Yang, Timothy A Bell, Gary A Churchill, Fernando Pardo-Manuel de Villena *Nature genetics* (2007-07-29) https://www.ncbi.nlm.nih.gov/pubmed/17660819 DOI: 10.1038/ng2087 · PMID: 17660819

34.     **Negative epistasis between natural variants of the Saccharomyces cerevisiae MLH1 and PMS1 genes results in a defect in mismatch repair.** Julie Akiko Heck, Juan Lucas Argueso, Zekeriyya Gemici, Richard Guy Reeves, Ann Bernard, Charles F Aquadro, Eric Alani *Proceedings of the National Academy of Sciences of the United States of America* (2006-02-21) https://www.ncbi.nlm.nih.gov/pubmed/16492773 DOI: 10.1073/pnas.0510998103 · PMID: 16492773 · PMCID: PMC1413905

35.     **Mechanisms of APOBEC3 mutagenesis in human cancer cells.** Mia Petljak, Alexandra Dananberg, Kevan Chu, Erik N Bergstrom, Josefine Striepen, Patrick von Morgen, Yanyang Chen, Hina Shah, Julian E Sale, Ludmil B Alexandrov, … John Maciejowski *Nature* (2022-07-20) https://www.ncbi.nlm.nih.gov/pubmed/35859169 DOI: 10.1038/s41586-022-04972-y · PMID: 35859169 · PMCID: PMC9329121

36.     **Evolution of the germline mutation rate across vertebrates.** Lucie A Bergeron, Søren Besenbacher, Jiao Zheng, Panyi Li, Mads Frost Bertelsen, Benoit Quintard, Joseph I Hoffman, Zhipeng Li, Judy St Leger, Changwei Shao, … Guojie Zhang *Nature* (2023-03-01) https://www.ncbi.nlm.nih.gov/pubmed/36859541 DOI: 10.1038/s41586-023-05752-y · PMID: 36859541 · PMCID: PMC9995274

37.     **Methylation of histone H3 lysine 36 enhances DNA repair by nonhomologous end-joining.** Sheema Fnu, Elizabeth A Williamson, Leyma P De Haro, Mark Brenneman, Justin Wray, Montaser Shaheen, Krishnan Radhakrishnan, Suk-Hee Lee, Jac A Nickoloff, Robert Hromas *Proceedings of the National Academy of Sciences of the United States of America* (2010-12-27) https://www.ncbi.nlm.nih.gov/pubmed/21187428 DOI: 10.1073/pnas.1013571108 · PMID: 21187428 · PMCID: PMC3021059

38.     **The SET domain protein Metnase mediates foreign DNA integration and links integration to nonhomologous end-joining repair.** Suk-Hee Lee, Masahiko Oshige, Stephen T Durant, Kanwaldeep Kaur Rasila, Elizabeth A Williamson, Heather Ramsey, Lori Kwan, Jac A Nickoloff, Robert Hromas *Proceedings of the National Academy of Sciences of the United States of America* (2005-12-06) https://www.ncbi.nlm.nih.gov/pubmed/16332963 DOI: 10.1073/pnas.0503676102 · PMID: 16332963 · PMCID: PMC1312370

39.     **The DDN catalytic motif is required for Metnase functions in non-homologous end joining (NHEJ) repair and replication restart.** Hyun-Suk Kim, Qiujia Chen, Sung-Kyung Kim, Jac A Nickoloff, Robert Hromas, Millie M Georgiadis, Suk-Hee Lee *The Journal of biological chemistry* (2014-02-25) https://www.ncbi.nlm.nih.gov/pubmed/24573677 DOI: 10.1074/jbc.m113.533216 · PMID: 24573677 · PMCID: PMC4036204

40.     **Biochemical characterization of metnase's endonuclease activity and its role in NHEJ repair.** Brian D Beck, Sung-Sook Lee, Elizabeth Williamson, Robert A Hromas, Suk-Hee Lee *Biochemistry* (2011-04-27) https://www.ncbi.nlm.nih.gov/pubmed/21491884 DOI: 10.1021/bi200333k · PMID: 21491884 · PMCID: PMC3388547

41.     **Not breathing is not an option: How to deal with oxidative DNA damage.** Enni Markkanen *DNA repair* (2017-09-22) https://www.ncbi.nlm.nih.gov/pubmed/28963982 DOI: 10.1016/j.dnarep.2017.09.007 · PMID: 28963982

42.   **8-oxoguanine causes spontaneous de novo germline mutations in mice.** Mizuki
Ohno, Kunihiko Sakumi, Ryutaro Fukumura, Masato Furuichi, Yuki Iwasaki, Masaaki Hokama,
Toshimichi Ikemura, Teruhisa Tsuzuki, Yoichi Gondo, Yusaku Nakabeppu *Scientific reports*
(2014-04-15) https://www.ncbi.nlm.nih.gov/pubmed/24732879 DOI: 10.1038/srep04689 · PMID:
24732879 · PMCID: PMC3986730

43.   **Cloning and expression of cDNA for a human enzyme that hydrolyzes 8-oxo-**
**dGTP, a mutagenic substrate for DNA synthesis.** K Sakumi, M Furuichi, T Tsuzuki, T
Kakuma, S Kawabata, H Maki, M Sekiguchi *The Journal of biological chemistry* (1993-11-05)
https://www.ncbi.nlm.nih.gov/pubmed/8226881 PMID: 8226881

44.   **Novel mutations of OGG1 base excision repair pathway gene in laryngeal cancer**
**patients.** Ishrat Mahjabeen, Nosheen Masood, Ruqia Mehmood Baig, Maimoona Sabir, Uzma
Inayat, Faraz Arshad Malik, Mahmood Akhtar Kayani *Familial cancer* (2012-12)
https://www.ncbi.nlm.nih.gov/pubmed/22829015 DOI: 10.1007/s10689-012-9554-2 · PMID:
22829015

45.   **Mutations in OGG1, a gene involved in the repair of oxidative DNA damage, are**
**found in human lung and kidney tumours.** S Chevillard, JP Radicella, C Levalois, J Lebeau,
MF Poupon, S Oudard, B Dutrillaux, S Boiteux *Oncogene* (1998-06-11)
https://www.ncbi.nlm.nih.gov/pubmed/9662341 DOI: 10.1038/sj.onc.1202096 · PMID: 9662341

46.   **Defects in 8-oxo-guanine repair pathway cause high frequency of C > A**
**substitutions in neuroblastoma** Marlinde L van den Boogaard, Rurika Oka, Anne Hakkert,
Linda Schild, Marli E Ebus, Michael R van Gerven, Danny A Zwijnenburg, Piet Molenaar, Lieke
L Hoyng, MEmmy M Dolman, … Jan J Molenaar *Proceedings of the National Academy of*
*Sciences* (2021-09-03) https://doi.org/grtcs9 DOI: 10.1073/pnas.2007898118 · PMID: 34479993
· PMCID: PMC8433536

47.   **SIFT: Predicting amino acid changes that affect protein function.** Pauline C Ng,
Steven Henikoff *Nucleic acids research* (2003-07-01)
https://www.ncbi.nlm.nih.gov/pubmed/12824425 DOI: 10.1093/nar/gkg509 · PMID: 12824425 ·
PMCID: PMC168916

48.   **The mutational constraint spectrum quantified from variation in 141,456 humans.**
Konrad J Karczewski, Laurent C Francioli, Grace Tiao, Beryl B Cummings, Jessica Alföldi,
Qingbo Wang, Ryan L Collins, Kristen M Laricchia, Andrea Ganna, Daniel P Birnbaum, …
Daniel G MacArthur *Nature* (2020-05-27) https://www.ncbi.nlm.nih.gov/pubmed/32461654 DOI:
10.1038/s41586-020-2308-7 · PMID: 32461654 · PMCID: PMC7334197

49.   **A method and server for predicting damaging missense mutations.** Ivan A
Adzhubei, Steffen Schmidt, Leonid Peshkin, Vasily E Ramensky, Anna Gerasimova, Peer Bork,
Alexey S Kondrashov, Shamil R Sunyaev *Nature methods* (2010-04)
https://www.ncbi.nlm.nih.gov/pubmed/20354512 DOI: 10.1038/nmeth0410-248 · PMID:
20354512 · PMCID: PMC2855889

50. **The base excision repair enzyme MED1 mediates DNA damage response to antitumor drugs and is associated with mismatch repair system integrity.** Salvatore Cortellino, David Turner, Valeria Masciullo, Filippo Schepis, Domenico Albino, Rene Daniel, Anna Marie Skalka, Neal J Meropol, Christophe Alberti, Lionel Larue, Alfonso Bellacosa *Proceedings of the National Academy of Sciences of the United States of America* (2003-11-12) https://www.ncbi.nlm.nih.gov/pubmed/14614141 DOI: 10.1073/pnas.2334585100 · PMID: 14614141 · PMCID: PMC299910

51. **MBD4 deficiency reduces the apoptotic response to DNA-damaging agents in the murine small intestine.** Owen James Sansom, Joanna Zabkiewicz, Stefan Mark Bishop, Jackie Guy, Adrian Bird, Alan Richard Clarke *Oncogene* (2003-10-16) https://www.ncbi.nlm.nih.gov/pubmed/14562041 DOI: 10.1038/sj.onc.1206850 · PMID: 14562041

52. **Enhanced CpG mutability and tumorigenesis in MBD4-deficient mice.** Catherine B Millar, Jacky Guy, Owen J Sansom, Jim Selfridge, Eilidh MacDougall, Brian Hendrich, Peter D Keightley, Stefan M Bishop, Alan R Clarke, Adrian Bird *Science (New York, N.Y.)* (2002-07-19) https://www.ncbi.nlm.nih.gov/pubmed/12130785 DOI: 10.1126/science.1073354 · PMID: 12130785

53. **A human cancer-associated truncation of MBD4 causes dominant negative impairment of DNA repair in colon cancer cells.** SA Bader, M Walker, DJ Harrison *British journal of cancer* (2007-02-06) https://www.ncbi.nlm.nih.gov/pubmed/17285135 DOI: 10.1038/sj.bjc.6603592 · PMID: 17285135 · PMCID: PMC2360052

54. **R/qtl2: Software for Mapping Quantitative Trait Loci with High-Dimensional Data and Multiparent Populations.** Karl W Broman, Daniel M Gatti, Petr Simecek, Nicholas A Furlotte, Pjotr Prins, Śaunak Sen, Brian S Yandell, Gary A Churchill *Genetics* (2018-12-27) https://www.ncbi.nlm.nih.gov/pubmed/30591514 DOI: 10.1534/genetics.118.301595 · PMID: 30591514 · PMCID: PMC6366910

55. **An analytical framework for whole-genome sequence association studies and its implications for autism spectrum disorder.** Donna M Werling, Harrison Brand, Joon-Yong An, Matthew R Stone, Lingxue Zhu, Joseph T Glessner, Ryan L Collins, Shan Dong, Ryan M Layer, Eirene Markenscoff-Papadimitriou, … Stephan J Sanders *Nature genetics* (2018-04-26) https://www.ncbi.nlm.nih.gov/pubmed/29700473 DOI: 10.1038/s41588-018-0107-y · PMID: 29700473 · PMCID: PMC5961723

56. **The impact of genetic modifiers on variation in germline mutation rates within and among human populations.** William R Milligan, Guy Amster, Guy Sella *Genetics* (2022-07-30) https://www.ncbi.nlm.nih.gov/pubmed/35666194 DOI: 10.1093/genetics/iyac087 · PMID: 35666194 · PMCID: PMC9339295

57. **Understanding what determines the frequency and pattern of human germline mutations.** Norman Arnheim, Peter Calabrese *Nature reviews. Genetics* (2009-07) https://www.ncbi.nlm.nih.gov/pubmed/19488047 DOI: 10.1038/nrg2529 · PMID: 19488047 · PMCID: PMC2744436

58.     **Empirical threshold values for quantitative trait mapping.** GA Churchill, RW Doerge
*Genetics* (1994-11) https://www.ncbi.nlm.nih.gov/pubmed/7851788 DOI:
10.1093/genetics/138.3.963 · PMID: 7851788 · PMCID: PMC1206241

59.     **Confidence intervals in QTL mapping by bootstrapping.** PM Visscher, R Thompson,
CS Haley *Genetics* (1996-06) https://www.ncbi.nlm.nih.gov/pubmed/8725246 DOI:
10.1093/genetics/143.2.1013 · PMID: 8725246 · PMCID: PMC1207319

60.     Alexandra M Stafford, Cheryl Reed, Harue Baba, Nicole Ar Walter, John Rk Mootz,
Robert W Williams, Kim A Neve, Lev M Fedorov, Aaron J Janowsky, Tamara J Phillips *eLife*
(2019-07-09) https://www.ncbi.nlm.nih.gov/pubmed/31274109 DOI: 10.7554/elife.46472 · PMID:
31274109 · PMCID: PMC6682400

61.     **Identification of a Functional Non-coding Variant in the GABA**  Megan K Mulligan,
Timothy Abreo, Sarah M Neuner, Cory Parks, Christine E Watkins, MTrevor Houseal, Thomas
M Shapaker, Michael Hook, Haiyan Tan, Xusheng Wang, … Robert W Williams *Frontiers in*
*genetics* (2019-03-29) https://www.ncbi.nlm.nih.gov/pubmed/30984232 DOI:
10.3389/fgene.2019.00188 · PMID: 30984232 · PMCID: PMC6449455

62.     **Investigating a downstream gene of**  Ye Lu, Diana Zhou, Hong Lu, Fuyi Xu, Junming
Yue, Jianping Tong, Lu Lu *Molecular vision* (2019-04-23)
https://www.ncbi.nlm.nih.gov/pubmed/31057322 PMID: 31057322 · PMCID: PMC6478243

63.     **Array programming with NumPy** Charles R Harris, KJarrod Millman, Stéfan J van der
Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, Julian Taylor, Sebastian
Berg, Nathaniel J Smith, … Travis E Oliphant *Nature* (2020-09-16) https://doi.org/ghbzf2 DOI:
10.1038/s41586-020-2649-2 · PMID: 32939066 · PMCID: PMC7759461

64.     **pandas-dev/pandas: Pandas** The Pandas Development Team *Zenodo* (2023-08-30)
https://doi.org/ggt8bh DOI: 10.5281/zenodo.3509134

65.     **Matplotlib: A 2D Graphics Environment** John D Hunter *Computing in Science &*
*Engineering* (2007) https://doi.org/drbjhg DOI: 10.1109/mcse.2007.55

66.     **Scikit-learn: Machine Learning in Python** Fabian Pedregosa, Gaël Varoquaux,
Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, Peter
Prettenhofer, Ron Weiss, Vincent Dubourg, … Édouard Duchesnay *Journal of Machine*
*Learning Research* (2011) http://jmlr.org/papers/v12/pedregosa11a.html

67.     **pandera: Statistical Data Validation of Pandas Dataframes** Niels Bantilan
*Proceedings of the Python in Science Conference* (2020) https://doi.org/grr54q DOI:
10.25080/majora-342d178e-010

68.     **seaborn: statistical data visualization** Michael Waskom *Journal of Open Source*
*Software* (2021-04-06) https://doi.org/gjqn3g DOI: 10.21105/joss.03021

69.    **Numba** Siu Kwan Lam, Antoine Pitrou, Stanley Seibert *Proceedings of the Second*
*Workshop on the LLVM Compiler Infrastructure in HPC* (2015-11-15) https://doi.org/gf3nks DOI:
10.1145/2833157.2833162

70.    **Sustainable data analysis with Snakemake** Felix Mölder, Kim Philipp Jablonski, Brice
Letcher, Michael B Hall, Christopher H Tomkins-Tinch, Vanessa Sochat, Jan Forster, Soohyun
Lee, Sven O Twardziok, Alexander Kanitz, … Johannes Köster *F1000Research* (2021-01-18)
https://doi.org/gjjkwv DOI: 10.12688/f1000research.29032.1 · PMID: 34035898 · PMCID:
PMC8114187

71.    **A natural mutator allele shapes mutation spectrum variation in mice** Tom Sasani
(2023-01-24) https://github.com/tomsasani/bxd_mutator_manuscript

72.    **tomsasani/bxd_mutator_manuscript: Final figure generation updates prior to**
**publication** Tom Sasani *Zenodo* (2022-02-01) https://doi.org/grrwv8 DOI:
10.5281/zenodo.5941048

73.    **BXD Genotype / WebQTL** https://gn1.genenetwork.org/dbdoc/BXDGeno.html

74.    **Genetics and Probability in Animal Breeding Experiments**
https://link.springer.com/book/10.1007/978-1-349-04904-2

75.    **ENA Browser** https://www.ebi.ac.uk/ena/browser/view/PRJEB45429

76.    **Twelve years of SAMtools and BCFtools.** Petr Danecek, James K Bonfield, Jennifer
Liddle, John Marshall, Valeriu Ohan, Martin O Pollard, Andrew Whitwham, Thomas Keane,
Shane A McCarthy, Robert M Davies, Heng Li *GigaScience* (2021-02-16)
https://www.ncbi.nlm.nih.gov/pubmed/33590861 DOI: 10.1093/gigascience/giab008 · PMID:
33590861 · PMCID: PMC7931819

77.    **A program for annotating and predicting the effects of single nucleotide**
**polymorphisms, SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118;**
**iso-2; iso-3.** Pablo Cingolani, Adrian Platts, Le Lily Wang, Melissa Coon, Tung Nguyen, Luan
Wang, Susan J Land, Xiangyi Lu, Douglas M Ruden *Fly* (2012)
https://www.ncbi.nlm.nih.gov/pubmed/22728672 DOI: 10.4161/fly.19695 · PMID: 22728672 ·
PMCID: PMC3679285

78.    **cyvcf2: fast, flexible variant analysis with Python.** Brent S Pedersen, Aaron R
Quinlan *Bioinformatics (Oxford, England)* (2017-06-15)
https://www.ncbi.nlm.nih.gov/pubmed/28165109 DOI: 10.1093/bioinformatics/btx057 · PMID:
28165109 · PMCID: PMC5870853

79.    **The UCSC Table Browser data retrieval tool.** Donna Karolchik, Angela S Hinrichs,
Terrence S Furey, Krishna M Roskin, Charles W Sugnet, David Haussler, WJames Kent *Nucleic*
*acids research* (2004-01-01) https://www.ncbi.nlm.nih.gov/pubmed/14681465 DOI:
10.1093/nar/gkh103 · PMID: 14681465 · PMCID: PMC308837

80. **bx-python** Taylor Lab at Johns Hopkins University (2023-10-22) https://github.com/bxlab/bx-python

81. **A simple regression method for mapping quantitative trait loci in line crosses using flanking markers.** CS Haley, SA Knott *Heredity* (1992-10) https://www.ncbi.nlm.nih.gov/pubmed/16718932 DOI: 10.1038/hdy.1992.131 · PMID: 16718932

82. **Study on a new chromogenic substrate for the prothrombin time determination.** A Girolami, A Santarossa, S Martinelli, MT Sartori, I Visentin *Folia haematologica (Leipzig, Germany : 1928)* (1987) https://www.ncbi.nlm.nih.gov/pubmed/2453419 PMID: 2453419

83. **GeneNetwork: a continuously updated tool for systems genetics analyses** Pamela M Watson, David G Ashbrook *Cold Spring Harbor Laboratory* (2020-12-24) https://doi.org/gsnpr5 DOI: 10.1101/2020.12.23.424047

84. **Birth of a chimeric primate gene by capture of the transposase gene from a mobile element.** Richard Cordaux, Swalpa Udit, Mark A Batzer, Cédric Feschotte *Proceedings of the National Academy of Sciences of the United States of America* (2006-05-03) https://www.ncbi.nlm.nih.gov/pubmed/16672366 DOI: 10.1073/pnas.0601161103 · PMID: 16672366 · PMCID: PMC1472436

85. **The roles of the human SETMAR (Metnase) protein in illegitimate DNA recombination and non-homologous end joining repair.** Michael Tellier, Ronald Chalmers *DNA repair* (2019-06-19) https://www.ncbi.nlm.nih.gov/pubmed/31238295 DOI: 10.1016/j.dnarep.2019.06.006 · PMID: 31238295 · PMCID: PMC6715855