Time-course of antipredator behavioral changes induced by the helminth *Pomphorhynchus laevis* in its intermediate host *Gammarus pulex*: the switch in manipulation according to parasite developmental stage differs between behaviors

Thierry Rigaud [orcid](): 0000-0002-0163-6639 *, Aude Balourdet and Alexandre Bauer

Université de Bourgogne, Laboratoire Biogéosciences, UMR CNRS 6282, équipe Ecologie Evolutive, 6 boulevard Gabriel, 21000 Dijon

*Corresponding author, [thierry.rigaud@u-bourgogne.fr](),

Key-words: parasite-induced behavioral manipulation, acanthocephalan, freshwater amphipod

Abstract:

Many trophically transmitted parasites with complex life cycles manipulate their intermediate host's antipredatory defenses in ways facilitating their transmission to final

host by predation. Some parasites also protect the intermediate host from predation when noninfective during its ontogeny. The acanthocephalan *Pomphorynchus laevis*, a fish

intestinal helminth, infecting freshwater gammarid amphipods as intermediate hosts, is using such a strategy of protection-then-exposure to predation by the definitive host.

However, the whole time-course of this sequence of behavioral switch is not yet known, and only one antipredator behavior has been studied to date. Here we show that the protective

part of this manipulation begins quite late during the parasite ontogeny, suggesting that the advantages overpass the costs induced by this protective manipulation only at this stage. We

confirmed that the refuge use behavior is showing a switch in the few days following the stage mature for the definitive host (switching from overuse to underuse). However, such a

switch was not observed for the gammarids activity rate, a behavior also known to make the host less conspicuous to predators when weak. While we predicted a low activity during

early development stages, then a switch to high activity, we observed general decrease in activity during the parasite ontogeny. The possible causes for this discrepancy between

behaviors are discussed. All these behavioral changes were observed mostly when animals were tested in water scented by potential predators (here brown trout), suggesting

condition-dependent manipulation.

Introduction

Heteroxenous parasites require successive hosts to complete their life cycle (Poulin, 2007). A successful transfer from one host to another is critical to the fitness of such parasites, and numerous parasite species rely on trophical transmission to achieve this goal. In this process, downstream infected host(s) – the intermediate host(s) – must be consumed by the upstream host – the definitive host – where the parasite reproduce. Parasite-induced phenotypic alterations (PIPA) may facilitate the completion of the parasite's life cycle (Moore, 2002; Poulin, 2010). Alterations of host behavior are particularly frequent in trophically-transmitted parasites, with infected hosts showing behaviors increasing their chances to be predated by their parasite's final host e.g. (Bethel & Holmes, 1977; Moore, 1983; Lagrue et al., 2007), reviewed, for example, in Poulin (2010). These parasites generally alter the antipredator behaviors of their intermediate hosts (Svensson et al., 2022), and some parasites are even able to reverse their intermediate host's fear of predator cues (Berdoy et al., 2000; Perrot-Minnot et al., 2007). These host behavioral changes favoring parasite transmission are adaptive for parasites but detrimental to the host. They are therefore seen as an extended phenotype of parasites and are referred to as parasitic manipulation (Hughes et al., 2012).

The understanding of the adaptive significance of PIPA was reinforced by studies highlighting variations in manipulations depending on the ontogeny of the parasite. Parker et al. (2009) predicted that the trophically-transmitted parasites showing the strongest selective advantage would be those able to protect from predation when not infective to the definitive host and then switch to an increase of exposure to predation when infectious. This phenomenon might have been selected because trophic transmission occurring before the parasite has reached the infective stage would result in zero fitness (Parker et al., 2009). Ideally, then, an optimal temporal sequence of manipulation could consist of a 'protect-then-expose' strategy against predation. Accordingly, reinforced antipredatory behaviors were observed during the early stages of the ontogeny in a number of parasite species (e.g. Hammerschmidt et al., 2009; Médoc & Beisel, 2011; Bailly et al., 2018) and direct testing for predation rate of uninfected versus infected hosts confirmed the existence of this switch in predation rates (Dianne et al., 2011; Weinreich et al., 2013). This phenomenon has been

74   studied in detail in a few model organisms only. An iconic model is the tapeworm
Schistocephalus solidus, where infection in its copepod first intermediate host firstly reduces

76   the host activity and recovery time after a disturbance mimicking a predator attack
(Hammerschmidt et al., 2009), resulting in a decrease in predation (Weinreich et al., 2013).

78   This manipulation occurs rapidly after the establishment of the infection within the copepod
body (Benesh, 2019). After infectivity for the next host is reached, the behavioral pattern

80   reverses, infected copepods becoming more active and conspicuous to predators
(Hammerschmidt et al., 2009). In this system, the intensity of host manipulation varied

82   mainly according to parasite populations, and responded to selection on parasites,
confirming that this trait can be a parasite extended phenotype (Hafer, 2018; Hafer-

84   Hahmann, 2019), rather than a trait of the host also favoring the parasite fitness (Jensen et
al. 2023).

86   Another well-studied parasite-host system for switching in behavioral parasitic manipulation
is the acanthocephalan *Pomphorhynchus laevis* infecting the amphipod *Gammarus pulex*.

88   Here, the infection by the non-infective stage (the acanthella) protects the host from
predation by increasing the rate of refuge use in the amphipod (Dianne et al., 2011), a trait

90   protecting gammarids from predation by fish (Abjornsson et al., 2004). A switch then occurs
few days after the parasite reached the cystacanth stage (infective for the next host),

92   exposing the infected host to fish predation (Dianne et al., 2011). In this system, no selection
was made on parasites to confirm this extended phenotype, but the intensity of

94   manipulation depends on the parasite strains (Franceschi, Bollache, et al., 2010) or
populations (Franceschi, Cornet, et al., 2010; Labaude et al., 2015), suggesting that

96   manipulation is a parasite trait. However, here, a number of data are still lacking to further
understand the 'protect-then-expose' strategy. For example, it is not known if *P. laevis*

98   invests in host protection early in its ontogeny, knowing that the development of the
acanthella stage necessitates quite a long time, i.e. between 2 and 3 months depending on

100   temperature (Labaude et al., 2020). No data are available on the course of behavioral
changes during the whole duration of the acanthellae stage. Indeed, measures of refuge use

102   or predation rates were made at the end of this acanthella stage, few days before the
parasite reached the infective stage (Dianne et al., 2011, 2014). However, a long time spent

104   in a host would necessitate a long investment in protection. Dianne et al. (2014) identified a

cost of the protective manipulation in terms of host food intake. Such a cost, if imposed on a

106   long time, could impact the survival of the host, therefore limiting a long protective

manipulation during parasite ontogeny. The life-cycle of *Schistocephalus solidus*, much

108   shorter (around two weeks before being infective in the copepod host), probably allows

early investment in the protective manipulation (Hammerschmidt et al., 2009). In addition, it

110   is known that manipulation of gammarids by *P. laevis* is multidimentional, i.e. that multiple

traits are changed in the hosts by the parasite infection (Perrot-Minnot et al., 2014; Fayard

112   et al., 2020). However, we still do not know if multiple antipredatory traits are changed

during protective manipulation. The activity rate, for example, known to be reduced as an

114   antipredator defense in gammarids (Williams & Moore, 1985; Bollache et al., 2006) should

be reduced during the acanthella stage, and then switch to an increase at the cystacanth

116   stage, but this has never been tested. Finally, while Dianne et al. (2011) found that

acanthellae induced an increased use of refuges only in the presence of predator cues (fish

118   scent), Dianne et al. (2014) found contradictorily that this protective manipulation in *P.*

laevis was observed both in presence or absence of predatory cues, suggesting that the

120   protective manipulation is not condition-dependent. However, again, no data is available on

the entire ontogeny of the acanthella.

122   In this study, we analyzed the behavioral changes induced by *P. laevis* in *G. pulex*, for the

first time during the whole ontogeny of parasites. Two anti-predatory behaviors, namely the

124   intensity of host refuge use and the host activity, were measured after experimental

infections (as recommended by Poulin & Maure, 2015) and under two conditions: with or

126   without predation cues. Three main questions were addressed: (i) is the protective

manipulation present during early acanthella stages of parasite's larval development? (ii) is

128   this manipulation condition(predator)-dependent during the whole larval ontogeny? (iii) are

these two different anti-predatory behaviors affected in the same way by *P. laevis* during its

130   ontogeny?

132   **Material and Methods**

Animals and experimental infections

134   Collection, maintenance and infection procedure were as described in Franceschi et al.

(2008). In brief, uninfected *G. pulex* amphipods were collected in February 2012 from a small

136   tributary of the Suzon River, Burgundy, eastern France (47°23'56.19''N, 4°50'31.13''E). Only

male gammarids were kept for the experiment to avoid any interference of the female

138   reproductive cycle on the outcome of parasite infection (Franceschi *et al.* 2008). They were

acclimated in the laboratory for 3 weeks before experimental infections under a 12:12

140   light:dark cycle and fed during the experiment with conditioned elm leaves (*Ulmus laevis*).

Adult *P. laevis* parasites were removed from the intestine of naturally infected chub

142   (*Squalius cephalus*) caught in February 2012 in the Vouge River, France (47°09'34.36''N,

5°09'02.50''E) and characterized molecularly as described in Franceschi et al. (2008). Parasite

144   eggs were collected from six females of *P. laevis* sampled from four different fish. Male

gammarids were allowed to feed for 48 h on a 1 cm^2 piece of elm leaf, on which about 200

146   eggs were deposited. A total of 190 *G. pulex* males were exposed to parasite eggs. After

exposure, gammarids were maintained in isolation in cups (6 cm diameter) filled with c.a. 50

148   mL of aerated water. Dead elm leaves were provided as food *ad libitum* and water was

changed every week. Twenty control, unexposed, gammarids were handled and maintained

150   under the same conditions as the exposed gammarids.

152   *Behavioral measurements and survival*

Survival was controlled every day post-exposure, and dead gammarids were dissected within

154   the 24 hours post-death in order to control for parasite presence. Due to small size of

acanthella, we were unable to detect this presence before 21 days in the 14 *G. pulex* that

156   died (personal observations). Since we were unable to assign hosts to the infected or

uninfected categories, the comparisons of refuge use and survival were made after the 21th

158   day post-infection. The experiment was stopped after 83 days post-exposure.

We measured the rate of refuge use every week. Gammarids were introduced individually

160   into a 10.5 × 16 cm rectangular box filled with 330 mL of either control or fish-scented water,

with each gammarid kept in the same water type during the whole experiment. Ten of the

162   20 unexposed control gammarids were tested under fish-scented water, 10 in control water.

To produce predator cues, six young brown trouts (*Salmo trutta*), each ~ 20 cm long, were

164   kept in the laboratory at 15 ± 1 °C under a 12:12 light:dark cycle, in a tank filled with 80 L of

tap water previously dechlorinated, oxygenated and UV-treated. Trouts were acclimated for

166   2 weeks prior to the experiments and fed with a mix of frozen chironomid larvae and alive

gammarids in order to strengthen the predation signal (Wudkevich et al., 1997). The water

168   from a similar tank, treated as previously described but without trout scent, was used as a

control water for behavioral experiments. Individuals were randomly assigned to these

170   water treatments. An opaque refuge, consisting of half a terracotta saucer (8.5 diameter) in

which a 1 cm^2 opening was made, was placed at one end of each aquarium, covering

172   approximately 18% of the total aquarium area. Refuge use was recorded after an acclimation

period of 30 min. The gammarid's position in the box was then checked every 5 min over a

174   period of 120 min. A score of 1 was given if the gammarid was inside the refuge and a score

of 0 if it was found outside. For each gammarid, the total score of refuge use therefore

176   ranged between 0 (gammarid always outside the refuge) and 24 (gammarid always inside

the refuge). After behavioral measure, the gammarid returned to its cup, where new aerated

178   water (without fish scent) and food were provided.

The activity of gammarids was recorded the day after the refuge use measurement, using a

180   ViewPoint device and software (©Viewpoint Life Sciences, Inc. – 2010, France). Gammarids

were placed individually in Petri dishes (diameter 8.5 cm, height 1 cm) filled with either

182   control water or scented water, this water treatment being the same as for refuge use

measurements for each gammarid. After 2 min of acclimation, gammarid activity was

184   recorded with an infrared camera for 5 min (continuous recording), as the proportion of

time spent inactive. Inactivity was defined as movements at a speed below 10 mm s^{-1}. This

186   speed threshold was determined based on preliminary tests, showing that below 10 mm s^{-1}

discrimination is not possible between gammarids moving at very low speed (crawling) and

188   gammarids moving their pleopods for respiration. The 20 control unexposed gammarids

were not measured for this trait.

190   The infection procedure allowed synchronized development of acanthellae. Around 55 days

post-exposure, the presence of *P. laevis* acanthellae was unambiguously detectable through

192   the host cuticle (translucent light-orange and shapeless larval stages). Therefore, at that

time, all gammarids were assigned to exposed-infected or exposed-uninfected categories

194   (either dead gammarids showed an infection after dissection, or the infection became

detectable in living gammarids). It means that behavioral tests were made in blind for the

196   infection status before 55 days post-exposure. The cystacanth stage was reached ten days

later (appearing through the cuticle as small bright orange opaque balls).

Statistical analysis

200   The survival of gammarids was analyzed using Cox regressions, where the effects of infection

status (in a first analysis: unexposed vs exposed-uninfected vs exposed-infected individuals;

202   in a second analysis: uninfected vs infected), water type (scented vs control) and their

interaction were analyzed using the 'survival' R package (V 3.2). Similar procedures were

204   used in infected individuals only to test the effect of the number of parasites on survival.

Scores for refuge use and activity rate were analyzed as repeated measures using the

206   'nparLD' R software package (V 2.1), with the F2 LD F1 model. This function enables

nonparametric analyses of right-censored longitudinal data, allowing the decrease in sample

208   size along time, due to individuals' death (Noguchi et al., 2012). The effects of water type

(scented vs control), infection status (uninfected vs infected) and their interaction were

210   investigated along time, considered here as an ordinal variable. ANOVA-type statistics (AT

Statistics), as recommended for small sample sizes (n < 8) were performed (Noguchi et al.,

212   2012), the statistical analysis of time-series with nparLD being based on rank-order of the

observed data (see Noguchi et al., 2012 for details). We also report Cliff's δ with their 95%

214   confidence intervals (Cliff, 1993) as measures of non parametric effect size for the difference

in refuge scores or activity rate between infected vs uninfected gammarids at each time

216   step, for each water type. This parameter measures the frequency of values in a given

distribution that are larger than the values in another distribution. The values of $\delta \geq 0.11$ are

218   considered as small, those ≥ 0.28 as medium, and those ≥ 0.43 as large (Kraemer & Kupfer,

2006).

220   All statistical analyses were performed using R version 4.1.0 software (R Foundation for

Statistical Computing, Vienna, Austria).

Results

We were able to detect unambiguously infections through the cuticle of the hosts at the late acanthella stage around the 55th day post-exposure. Before this stage, parasite infections were only detectable after dissection of dead individuals from day 21. Out of the 176 surviving individuals exposed to parasite eggs, 132 were finally infected (75%). Among them, 60 (45.5%) were infected by only one parasite, 39 (29.5%) were infected by two parasites and 33 (25.0%) were infected by more than two parasites.

The experiment was stopped after 83 days post-infection and all surviving animals were dissected to confirm their infection status and count the parasites. Exposed-uninfected animals did not differ significantly from control unexposed gammarids in either survival or shelter use (supplementary informations). Therefore, in order to increase the power of the behavioral analyses at the end of the experiment, we merged unexposed and exposed-uninfected animals to create the uninfected or not infected category (terminology used thereafter).

The analysis of survival did not fulfill the conditions of proportional hazards throughout the whole period (χ^2 = 20.9775; 3 d.f.; p = 0.0001). We therefore ran two analyses according to the two main stages of the larval developmental cycle, first during the acanthellae stage, then after the cystacanth stage has been reached. Analyses of the survival during these two periods fulfilled the conditions of proportional hazards (χ^2 = 2.67542; 3 d.f.; p = 0.44 and χ^2 = 3.9257; 3 d.f.; p = 0.27, respectively).

Animals survived equally through the acanthella stage regardless of their infection status, but the uninfected individuals survived better than the infected ones after the parasites reached the cystacanth stage (Table 1, Figure 1). The water condition had no significant effect on survival (Table1).

Table 1: Effect of water type, infection status by *Pomphorhynchus laevis*, and their interaction on the survival of *Gammarus pulex* individuals, during the acanthella stage development (a) and after the cystacanth stage has

been reached (b).

Source	d.f.	z	p-value
(a)			
Water type	1	1.840	0.0658
Infection	**1**	1.039	0.2988
Water type*infection	1	-1.486	0.1374
(b)			
Water type	1	1.433	0.1517
Infection	**1**	**-2.562**	**0.0104**
Water type*infection	1	-1.141	0.2540

Figure 1: Cumulative survival of *Gammarus pulex* after experimental exposure to *Pomphorhynchus laevis* according to their infection status, during the acanthella stage development (a) and after the cystacanth stage

has been reached (b). Animals were maintained in water that was (or not) signed with fish scent. After cystacanth stage was reached, infected *G.pulex* suffer higher mortality.

Among the infected individuals, to analyze the effect of the infection intensity on survival,

the number of parasites was categorized as "1 parasite" or "more than 1 parasite". The number of individuals infected by one parasite were 29 in control water and 31 in scented

water, while 37 individuals with more than one parasite were found in control water and 35

in scented water. Neither the parasite intensity nor the water type influenced significantly

268   the survival (z = -1.556, p = 0.120 and z = 1.051, p = 0.293, respectively). The interaction was

       also non significant (z = 1.149, p =0.251).

        The refuge use by gammarids varied considerably with time (Figure 2, Table 2), being most

272   of the time moderate at the beginning of our survey, becoming progressively more intensive

       until reaching high values around 30 to 40 days.

274   **Table 2:** Results of the model from the nparLD analysis, testing for the effects of water type (scented vs.
       control), infection status (infected with *P. laevis* vs. uninfected) and date of measurement on the scores of
276   refuge use by *G. pulex* individuals.

	AT statistics	df	P-value
Water type	**7.8568**	**1**	**0.0051**
Infection	0.9645	1	0.3261
date	**12.9330**	**5.7522**	**< 0.00001**
Water type * Infection	0.4299	1	0.5120
Water type * date	**2.9745**	**5.7522**	**0.0075**
Infection * date	**11.1301**	**5.7522**	**< 0.00001**
Water type * Infection * date	7.5111	5.7522	0.6030

Figure 2: Scores of refuge use by gammarids, according to their infection status by *Pomphorhynchus laevis*,
280   between day 21 and day 83 post-experimental exposure to parasite eggs. Gammarids were tested in control
water (a), or water signed with fish scent (b). Scores range from 0 (individuals always outside the refuge) to 24
282   (individuals always inside the refuge). Thick lines are the medians, boxes are the upper and lower quartiles,
whiskers are the upper and lower 1.5*interquartile range, and dots represent outliers. The vertical black line
284   denotes the transition between acanthella and cystacanth stages. Sample sizes are given in italics below each
plot. Refuge use increase during acanthealla stages (this phenomenon was stronger under signed water) and
286   then decreased after cyctacanth stage is reached.

288   This phenomenon was found regardless of the infection status and the water type, but was

found to be more rapid under scented water than control water (Figure 2), the interaction

290 between water type and date being statistically significant (Table 2). The interaction

between date and infection was also highly significant (Table 2).

This reflects the observation that, after the cystacanth stage being reached, the refuge score

294 became weaker for infected individuals, while during the acanthella stage, the tendency was

that infected individuals used more refuges than uninfected ones (Figure 2).

296 This observation is reinforced by the analysis of the effect sizes, but here differences

appeared between scented and control water. Indeed, at acanthella stage, under non-

298 scented water, the difference in refuge scores between infected and uninfected animals was

always close to zero, indicating negligible effect of the infection (Figure 3). Under scented

300 water, however, the values of δ were often between -0.11 and -0.4, i.e. a range where effect

sizes are considered as small to medium (Vargha & Delaney, 2000; Kraemer & Kupfer, 2006).

302 Nevertheless, only one value was significantly different from zero (95% C.I. did not overlap

with zero), at day 55, i.e. at the late acanthella stage. After the cystacanth stage was

304 reached, and here whatever the type of water, the effect size reversed, becoming positive,

and can be considered as large, with values of δ exceeding 0.4 (Figure 3).

Figure 3: Effect sizes of refuge use of *Gammarus pulex* infected with *Pomphorhynchus laevis* compared to uninfected individuals, according to the date when the tests were made, separated by the water type in which they were tested. Points represent Cliff 's δ and error bars represent 95% confidence intervals. Positive values mean that the uninfected individuals used the refuge more than infected individuals, and negative values mean a lower use. The white, blue, yellow and red colors corresponded to areas where the effect size are considered as trivial, small, medium and large, respectively, using the comparisons between Cliff 's δ and Cohen's d effect size made by (Kraemer & Kupfer, 2006). These areas must be seen as indicative. The dotted line denotes the transition between acanthella and cystacanth stages.

The effect of infection intensity was explored in infected animals only for the two dates where the analysis above suggested reversed effects of the infection, and where sample size was high enough to guarantee reasonable power, i.e. at days 55 and 69 post-exposure. At day 55, no effect of parasite number was seen on refuge use, nor on day 69. Indeed, after categorization as "1 parasite" and "more than 1 parasite", Wilcoxon tests were not significant (z = -0.8144, p = 0.4097 and z = -1.3688, p = 0.1675, respectively). The same result was found if parasite number was treated as a discrete numerical variable since the correlations between parasite number and refuge use scores were non-significant for the two dates (Spearman's ρ = 0.0595, p = 0.6379, n = 65 and Spearman's ρ = 0.2333, p = 0.1229, n = 45, respectively).

The analysis of levels of activity revealed an effect of water type close to statistical significance, with significantly more time spent inactive in infected gammarids compared to uninfected ones, but also a highly significant increased inactivity with time (Figure 4, Table 3). None of the interactions were significant (Table 3).

Table 3: Results of the model from the nparLD analysis, testing for the effects of water type (scented vs. control), infection status (infected with *P. laevis* vs. uninfected) and date of measurement on the activity rate of *G. pulex* individuals.

	AT statistics	df	P-value
Water type	3.3850	1	0.0658
Infection	**6.1807**	**1**	**0.0129**
date	**17.5550**	**4.8886**	**< 0.00001**
Water type * Infection	2.5109	1	0.1131
Water type * date	1.1737	4.8886	0.3193
Infection * date	1.1683	4.8886	0.3221
Water type * Infection * date	1.1111	4.8886	0.3517

Figure 4: Proportion of time spent inactive in gammarids, according to their infection status by
340 *Pomphorhynchus laevis*, between day 21 and day 69 post experimental exposure to parasite eggs. Gammarids
were tested for 5 minutes in control water (a), or water signed with fish scent (b). Thick lines are the medians,
342 boxes are the upper and lower quartiles, whiskers are the upper and lower 1.5*interquartile range, and dots
represent outliers. The vertical black line denotes the transition between acanthella and cystacanth stages.
344 Sample sizes are given in italics below each plot. Infection increased the gammarid's inactivity whatever the
water type and the developmental stage.

Analysis of effect sizes (Figure 5) confirmed but visually refined the analysis, with infected

348   individuals being significantly more inactive than uninfected animals in the scented water.

350   **Figure 5:** Effect sizes of time spent inactive in *Gammarus pulex* infected with *Pomphorhynchus laevis* compared
to uninfected individuals, according to the dates when the tests were made and the water type. Points are
352   Cliff's δ and error bars represent 95% confidence intervals. Negative values mean that the infected individuals
are more inactive than uninfected animals. The white, blue, yellow and red colors corresponded to areas where
354   the effect size are considered as trivial, small, medium and large, respectively (see Figure 3). The dotted line
denotes the transition between acanthella and cystacanth stages.

Discussion

358   Our longitudinal survey of antipredatory behaviors in *Gammarus pulex* infected by

Pomphorhynchus laevis, made in repeated measures, first confirmed previous observations

360   that a shift occurred in refuge use between acanthella and cystacanth stages, i.e. between

parasite stages non-infective and infective for the next host in the cycle (Dianne et al., 2011,

362   2014). This shift was nevertheless not observed for the activity rate, another behavior

involved in predation avoidance (Andersson et al., 1986; Bollache et al., 2006). In addition,

364   our study brings new information on this anti-predatory behavior.

First, confirming observations of Dianne et al. (2011), we found a difference in the

366   'protective' behavioral patterns of refuge use and activity rate according to the presence or

absence of predatory cues. Indeed, while almost no difference between infected and

368   uninfected animals was found when tests were made in control water, *G. pulex* infected by

370 acanthellae use more the refuges at the end of their parasite development and are more
inactive than uninfected ones in fish-scented water. Fish scent is definitely a source of stress
372 for gammarids: it generally increases anti-predatory behavior when gammarids perceive
predatory cues (e.g. they are repulsed by the chemical cues originating from fish (Perrot-
Minnot et al., 2007) and reduce their activity (Andersson et al., 1986; Bollache et al., 2006).
374 The infection by acanthellae would therefore strengthen this behavior. In another
acanthocephalan parasite, *Polymorphus minutus*, the protective behavior has been observed
376 in absence of predatory cues (Bailly et al., 2018). This parasite nevertheless shows a different
nature of behavioral changes, probably in link with the difference in life cycle (*P. minutus* is
378 using birds as definitive hosts), with probable different mechanisms involved (Tain et al.,
2006; Perrot-Minnot et al., 2016).

380 The second information is that protective behaviors associated with the infection were
significant – in scented water – at variable times during parasite ontogeny, depending on the
382 behavior considered. This means that reinforcing the host's antipredatory behavior
outweighs the potential costs (Parker et al., 2009) differently depending on the behavior.
384 The reduced activity rate in infected animals, on the one hand, appeared quite early during
acanthella growth (see a longer discussion on this trait in the next paragraph). The increase
386 of refuge use, on the other hand, was found when acanthellae are growing old and was only
significant at 55 days. Costs imposed by this behavioral change, as the one identified by
388 Dianne et al. (2014), if imposed during a long time, may impact the survival of the host and
may therefore limit in time the expression of protective manipulation during parasite
390 ontogeny. In addition, for long-developing parasites, energy spent during growth may not be
available for parasite manipulation (Poulin, 1994). It results that the moment when the
392 expression of protection is the most optimal is when growth is complete, because energy
then becomes available again. Therefore, the strategy of investing this energy in anti-
394 predator behavior to safeguard past investments in growth can be selected. Interestingly,
here, we observed that the time at which the protective component of manipulation is
396 significant in *P. laevis* is when acanthella growth is complete, an observation compatible
with such a strategy. Mechanistically, such a time course may be explained by chemicals
398 secreted by parasites, that have the potential to be involved in host behavioral change
(Berger et al., 2021, and references therein). However, we observed some variation (in

400   scented water) on the differences between infected and uninfected gammarids: the effect

size fluctuates between negligible values to medium values. This variation may be due to the

402   fact that even uninfected gammarids did not use refuges constantly through time, a

phenomenon that may interfere with the parasite's consequences on host behavior. Indeed,

404   from day 21 to day 55, there was a general trend of *G. pulex* using refuge more intensively as

time is passing, and this independently from their infection status. After four behavioral

406   measurements, all gammarids spent most of their time under the refuge, and this

phenomenon is more intense in scented water. Gammarids, like other crustaceans are

408   sensitive to stress, and one response to stress is to increase the intensity of anti-predatory

behaviors (Fossat et al., 2014; Perrot-Minnot et al., 2017). Regular handling of shrimps

410   increase their stress and modify the animal's behavior (Takahashi, 2022). Therefore, we may

infer that our experimental procedure was a source of stress for gammarids. Indeed,

412   individuals were isolated for several weeks (gammarids are gregarious animals and isolation

have consequences on their behavior, e.g. (Labaude et al., 2017) and were regularly handled

414   to take them out of their isolation box and transfer them to a new environment (the

aquarium or petri dish for behavioral tests). The observation that this phenomenon is more

416   rapid under predatory cues is not surprising since predators are a major source of stress.

Whatever its mechanistic basis, this general increase in refuge use interferes with our ability

418   to observe behavioral differences between gammarids infected or not by acanthellae,

because it becomes less easy to find a difference in refuge use when this usage is already

420   intense. After the parasite reached the cystacanth stage, we observed the shift in refuge use,

and infected *G. pulex* are using the refuge less intensively than uninfected animals, as

422   repeatedly observed in cases of infection with *P. laevis* (Franceschi, Bollache, et al., 2010;

Dianne et al., 2011; Bauer & Rigaud, 2015; Fayard et al., 2020). The effect size of this

424   exposure manipulation appeared higher than the one of protective manipulation in the

present experiment, which was not the case in Dianne et al. (2011). However, in this later

426   study, only one time point at acanthella and cystacanth stages each were measured, and the

acanthella time-point was the one closest to cystacanth stage. If we compare only the

428   absolute values of effect size at 55 and 69 days in our study, in the scented water, we will

find no clear difference. However, by considering the whole ontogeny, it seems that the

430   effect size of manipulation is greater at cystacanth stage than acanthella stage. Poulin (2010)

noted that the relationship between the intensity of behavioral manipulation and the

432   increase of predation rate (supposed to be due to manipulation) was not always

straightforward, so we cannot really conclude about the efficiency of protection and

434   exposure in our parasite-host system. However, since increase in mortality was mainly

observed after the cystacanth stage being reached (as also observed by Labaude et al.,

436   2020), it would make sense that the exposure manipulation is stronger than the protective

manipulation because the selective pressure of reaching the definitive host is strong at that

438   stage.

The third information of our study is that, as predicted, a second antipredatory behavior,

440   namely the decreased activity rate, was also strengthened by infection during acanthella

stage. Here again the differences in activity rate between infected and uninfected hosts

442   were found mostly under predatory threat. This behavioural change appeared earlier in

ontogeny than refuge use, suggesting that, if this change in behavior is adaptive for parasite,

444   the costs of reducing activity are lower than increasing refuge use (Parker et al., 2009).

Indeed, in gammarids, a reduced activity is associated with limited metabolic expenditures

446   rather than with costs (e.g. Normant et al., 2007), while increased refuge use induce costs on

food intake (Dianne et al. 2014). However, and contrary to predictions, no shift in behavioral

448   modification was observed for this behavior between acanthella and cystacanth stages, as

the one observed for refuge use, the infected gammarids even being less active at

450   cystacanth stages than at acanthella stages. Two alternative and exclusive hypotheses may

be proposed for this observation. The first one is that the decreased activity rate in infected

452   animals is a non-adaptive pathogenic byproduct of the infection, as sometimes suggested for

some behavioral changes induced by parasites (Levri, 1999; Cézilly et al., 2010). Under this

454   hypothesis, the progressive decrease according to time could just be interpreted as the long

time spent by parasite in the host, decreasing progressively its mobility capacity. This could

456   even be considered as a host strategy since here the interests of the parasite and the host

are aligned during acanthella stage (reduced activity reduce predation risk, which is

458   advantageous for both the parasite and the host, and save energy, which is advantageous

for the host). Such hypotheses nevertheless explain neither why such a byproduct of the

460   infection is not observed during acanthella stage in control water, nor the fact that the host

and parasite interests became unaligned after the cystacanth stage is reached. Indeed, here,

462   the reduced risk of predation due to low activity is advantageous for the host, but not for

the parasite. The second hypothesis would be to consider that decreasing activity is adaptive

464   for the parasite at both stages of development. During acanthella stage, this low activity rate

would make the gammarid less conspicuous towards predators, as classically interpreted

466   (Williams & Moore, 1985; Bollache et al., 2006). Notably, high rate of activity allows

gammarids to initiate swimming and consequently enter the river current (drift) where

468   predators are present (Williams & Moore, 1985; Schaffer et al., 2013). During the cystacanth

stage, the animal's tendency to search light (Franceschi et al., 2008) and avoid refuges (this

470   study, Dianne et al., 2011) would already bring them into the river drift (McCahon et al.,

1991; Lagrue et al., 2007). A low level of activity would therefore accentuate the predation

472   exposure by reducing the swimming capacity of gammarids, which could be easily

transported passively by the current of the river. Indeed, Lagrue et al. (2007) observed that

474   the presence of *G. pulex* in the river drift is increased by 25-30 times when infected by *P.
laevis*, the very same proportion of gammarids being found in the stomach of Bullhead

476   predators. We could have here an example of parasite's manipulation of a single host

behavior that could be beneficial at all parasite stages, a characteristic that could have

478   facilitated its selection.

To conclude, the first component of the 'protective' behavioral manipulation by the

480   acanthocephalan *Pomphorhynchus laevis* in its host *Gammarus pulex*, namely the increased

refuge use, was significant in late acanthella stage, only when predator cues were present in

482   the environment. This antipredator behavior enhancement was not significant early in the *P.
laevis* ontogeny, perhaps because of costs due to the long development time of parasites.

484   The refuge use then became weaker when the cystacanth stage was reached, illustrating the

'exposure' component of this parasitic manipulation. Another anti-predator behavior, the

486   decrease in activity rate, was also expressed more strongly after the parasite infection, but

showed no reversal during parasite ontogeny as observed for use of refuges. Differences in

488   the intensity and direction of these two behavioral changes may be due to differences in

their benefits relative to the developmental stage of the parasite: reduced host activity may

490   be beneficial for transmission of the parasite to all parasite stages, while high and low refuge

use are beneficial to the acanthella and cystacanth stages, respectively.

Funding: This study was funded by the French Agence Nationale de la Recherche (ANR),

494   ANR-13-BSV7-0004-01

Conflict of interest disclosure: The authors declare they have no financial conflicts of

496   interest in relation to the content of the article.

Author contributions: TR: Funding acquisition, Conceptualization, Experiments, Data

498   curation, Statistical analysis, Writing – original draft, Writing – review and editing; ABal:

Experiments, Data curation, Writing – review and editing; ABau: Statistical analysis, Writing –

500   review and editing.

Data access: https://doi.org/10.57745/AKA33Z

502   **Supplementary information :** found at https://doi.org/10.1101/2023.04.25.538244.
Includes comparisons between unexposed and uninfected-exposed animals and R codes

504   used for analyses.

506   **References**

Abjornsson K, Hansson L-A, Brönmark C (2004) Responses of prey from habitats with different
508   predator regimes: local adaptation and heritability. *Ecology*, **85**, 1859–1866.
https://doi.org/10.1890/03-0074

510   Andersson K, Brönmark C, Herrmann J, Malmqvist B, Otto C, Sjörström P (1986) Presence of sculpins
(*Cottus gobio*) reduces drift activity of *Gammarus pulex* (Amphipoda). *Hydrobiologia*, **133**,
512   209–215. https://doi.org/10.1007/BF00005592

Bailly Y, Cézilly F, Rigaud T (2018) Stage-dependent behavioural changes but early castration induced
514   by the acanthocephalan parasite *Polymorphus minutus* in its *Gammarus pulex* intermediate
host. *Parasitology*, **145**, 260–268. https://doi.org/10.1017/S0031182017001457

516   Bauer A, Rigaud T (2015) Identifying a key host in an acanthocephalan-amphipod system.
Parasitology, **142**, 1588–1594. https://doi.org/10.1017/S0031182015001067

518   Benesh DP (2019) Tapeworm manipulation of copepod behaviour: parasite genotype has a larger
effect than host genotype. *Biology Letters*, **15**, 20190495.
520   https://doi.org/10.1098/rsbl.2019.0495

Berdoy M, Webster JP, Macdonald DW (2000) Fatal attraction in rats infected with *Toxoplasma*
522   *gondii*. *Proceedings of the Royal Society B: Biological Sciences*, **267**, 1591–1594.

Berger CS, Laroche J, Maaroufi H, Martin H, Moon K-M, Landry CR, Foster LJ, Aubin-Horth N (2021)
524   The parasite *Schistocephalus solidus* secretes proteins with putative host manipulation
functions. *Parasites & Vectors*, **14**, 436. https://doi.org/10.1186/s13071-021-04933-w

526 Bethel WM, Holmes JC (1977) Increased vulnerability of amphipods to predation owing to altered behavior induced by larval acanthocephalans. *Canadian Journal of Zoology*, **55**, 110–115.

528 https://doi.org/10.1139/z77-013

Bollache L, Kaldonski N, Troussard J-P, Lagrue C, Rigaud T (2006) Spines and behaviour as defences

530 against fish predators in an invasive freshwater amphipod. *Animal Behaviour*, **72**, 627–633. https://doi.org/10.1016/j.anbehav.2005.11.020

532 Cézilly F, Thomas F, Médoc V, Perrot-Minnot M-J (2010) Host-manipulation by parasites with complex life cycles: adaptive or not? *Trends in Parasitology*, **26**, 311–317.

534 https://doi.org/10.1016/j.pt.2010.03.009

Cliff N (1993) Dominance Statistics: Ordinal Analyses to Answer Ordinal Questions. *Psychological*

536 *Bulletin*, **113**, 494–509. https://doi.org/10.1037/0033-2909.114.3.494

Dianne L, Perrot-Minnot M-J, Bauer A, Gaillard M, Léger E, Rigaud T (2011) Protection first then

538 facilitation: a manipulative parasite modulates the vulnerability to predation of its intermediate host according to its own developmental stage. *Evolution*, **65**, 2692–2698.

540 https://doi.org/10.1111/j.1558-5646.2011.01330.x

Dianne L, Perrot-Minnot M-J, Bauer A, Guvenatam A, Rigaud T (2014) Parasite-induced alteration of

542 plastic response to predation threat: increased refuge use but lower food intake in *Gammarus pulex* infected with the acanothocephalan *Pomphorhynchus laevis*. *International*

544 *Journal for Parasitology*, **44**, 211–216. https://doi.org/10.1016/j.ijpara.2013.11.001

Fayard M, Dechaume-Moncharmont F, Wattier R, Perrot-Minnot M (2020) Magnitude and direction

546 of parasite-induced phenotypic alterations: a meta-analysis in acanthocephalans. *Biological Reviews*, brv.12606. https://doi.org/10.1111/brv.12606

548 Fossat P, Bacqué-Cazenave J, De Deurwaerdère P, Delbecque J-P, Cattaert D (2014) Anxiety-like behavior in crayfish is controlled by serotonin. *Science*, **344**, 1293–1297.

550 https://doi.org/10.1126/science.1248811

Franceschi N, Bauer A, Bollache L, Rigaud T (2008) The effects of parasite age and intensity on

552 variability in acanthocephalan-induced behavioural manipulation. *International Journal for Parasitology*, **38**, 1161–1170. https://doi.org/10.1016/j.ijpara.2008.01.003

554 Franceschi N, Bollache L, Cornet S, Bauer A, Motreuil S, Rigaud T (2010) Co-variation between the intensity of behavioural manipulation and parasite development time in an

556 acanthocephalan-amphipod system: Co-variation between the intensity of behavioural manipulation and parasite development. *Journal of Evolutionary Biology*, **23**, 2143–2150.

558 https://doi.org/10.1111/j.1420-9101.2010.02076.x

Franceschi N, Cornet S, Bollache L, Dechaume-Moncharmont F-X, Bauer A, Motreuil S, Rigaud T

560 (2010) Variation between populations and local adaptation in acanthocephalan-induced parasite manipulation: variation in parasite-induced behavioral manipulation. *Evolution*, no-

562 no. https://doi.org/10.1111/j.1558-5646.2010.01006.x

Hafer N (2018) Differences between populations in host manipulation by the tapeworm

564 *Schistocephalus solidus* – is there local adaptation? *Parasitology*, **145**, 762–769. https://doi.org/10.1017/S0031182017001792

566   Hafer-Hahmann N (2019) Experimental evolution of parasitic host manipulation. *Proceedings of the Royal Society B: Biological Sciences*, **286**, 20182413. https://doi.org/10.1098/rspb.2018.2413

568   Hammerschmidt K, Koch K, Milinski M, Chubb JC, Parker GA (2009) When to go: optimization of host switching in parasites with complex life cycles. *Evolution*, **63**, 1976–1986.
570   https://doi.org/10.1111/j.1558-5646.2009.00687.x

Hughes DP, Brodeur J, Thomas F, UK P by RD University of Oxford (Eds.) (2012) *Host Manipulation by*
572   *Parasites*. Oxford University Press, Oxford, New York.

Jensen, C. H., Weidner, J., Giske, J.,Jørgensen, C., Eliassen, S., & Mennerat, A. (2023). Adaptive host
574   responses to infection can resemble parasitic manipulation. *Ecology and Evolution*, **13**, e10318. https://doi.org/10.1002/ece3.10318

576   Kraemer HC, Kupfer DJ (2006) Size of Treatment Effects and Their Importance to Clinical Research and Practice. *Biological Psychiatry*, **59**, 990–996.
578   https://doi.org/10.1016/j.biopsych.2005.09.014

Labaude S, Cézilly F, De Marco L, Rigaud T (2020) Increased temperature has no consequence for
580   behavioral manipulation despite effects on both partners in the interaction between a crustacean host and a manipulative parasite. *Scientific Reports*, **10**, 11670.
582   https://doi.org/10.1038/s41598-020-68577-z

Labaude S, Cézilly F, Tercier X, Rigaud T (2015) Influence of host nutritional condition on post-
584   infection traits in the association between the manipulative acanthocephalan *Pomphorhynchus laevis* and the amphipod *Gammarus pulex*. *Parasites & Vectors*, **8**, 403.
586   https://doi.org/10.1186/s13071-015-1017-9

Labaude S, Rigaud T, Cézilly F (2017) Additive effects of temperature and infection with an
588   acanthocephalan parasite on the shredding activity of *Gammarus fossarum* (Crustacea: Amphipoda): the importance of aggregative behavior. *Global Change Biology*, **23**, 1415–
590   1424. https://doi.org/10.1111/gcb.13490

Lagrue C, Kaldonski N, Perrot-Minnot MJ, Motreuil S, Bollache L (2007) Modification of hosts'
592   behavior by a parasite: field evidence for adaptive manipulation. *Ecology*, **88**, 2839–2847. https://doi.org/10.1890/06-2105.1

594   Levri EP (1999) Parasite-induced change in host behavior of a freshwater snail: parasitic manipulation or byproduct of infection? *Behavioral Ecology*, **10**, 234–241.
596   https://doi.org/10.1093/beheco/10.3.234

McCahon CP, Maund SJ, Poulton MJ (1991) The effect of the acanthocephalan parasite
598   (*Pomphorhynchus laevis*) on the drift of its intermediate host (*Gammarus pulex*). *Freshwater Biology*, **25**, 507–513. https://doi.org/10.1111/j.1365-2427.1991.tb01393.x

600   Médoc V, Beisel J-N (2011) When trophically-transmitted parasites combine predation enhancement with predation suppression to optimize their transmission. *Oikos*, **120**, 1452–1458.
602   https://doi.org/10.1111/j.1600-0706.2011.19585.x

Moore J (1983) Responses of an Avian Predator and Its Isopod Prey to an Acanthocephalan Parasite.
604   *Ecology*, **64**, 1000–1015. https://doi.org/10.2307/1937807

Moore J (2002) *Parasites and the Behavior of Animals*. Oxford University Press.

606   Noguchi K, Gel YR, Brunner E, Konietschke F (2012) nparLD: An R Software Package for the
       Nonparametric Analysis of Longitudinal Data in Factorial Experiments. *Journal of Statistical*
608   *Software*, **50**, 1–23. https://doi.org/10.18637/jss.v050.i12

       Normant M, Dziekonski M, Drzazgowski J, Lamprecht I (2007) Metabolic investigations of aquatic
610   organisms with a new twin heat conduction calorimeter. *Thermochimica Acta*, **458**, 101–106.
       https://doi.org/10.1016/j.tca.2007.01.025

612   Parker GA, Ball MA, Chubb JC, Hammerschmidt K, Milinski M (2009) When should a trophically
       transmitted parasite manipulate its host? *Evolution*, **63**, 448–458.
614   https://doi.org/10.1111/j.1558-5646.2008.00565.x

       Perrot-Minnot M-J, Banchetry L, Cézilly F (2017) Anxiety-like behaviour increases safety from fish
616   predation in an amphipod crustacea. *Royal Society Open Science*, **4**, 171558.
       https://doi.org/10.1098/rsos.171558

618   Perrot-Minnot M-J, Kaldonski N, Cézilly F (2007) Increased susceptibility to predation and altered
       anti-predator behaviour in an acanthocephalan-infected amphipod. *International Journal for*
620   *Parasitology*, **37**, 645–651. https://doi.org/10.1016/j.ijpara.2006.12.005

       Perrot-Minnot M, Maddaleno M, Cézilly F (2016) Parasite-induced inversion of geotaxis in a
622   freshwater amphipod: a role for anaerobic metabolism? *Functional Ecology*, **30**, 780–788.
       https://doi.org/10.1111/1365-2435.12516

624   Perrot-Minnot M-J, Sanchez-Thirion K, Cézilly F (2014) Multidimensionality in host manipulation
       mimicked by serotonin injection. *Proceedings of the Royal Society B: Biological Sciences*, **281**,
626   20141915. https://doi.org/10.1098/rspb.2014.1915

       Poulin R (1994) The evolution of parasite manipulation of host behaviour: a theoretical analysis.
628   *Parasitology,* **109**, S109-S118. doi:10.1017/s0031182000085127

       Poulin R (2007) *Evolutionary Ecology of Parasites: (Second Edition)*. Princeton University Press.
630   https://doi.org/10.2307/j.ctt7sn0x

       Poulin R (2010) Parasite Manipulation of Host Behavior. In: *Advances in the Study of Behavior*, pp.
632   151–186. Elsevier. https://doi.org/10.1016/S0065-3454(10)41005-0

       Poulin R, Maure F (2015) Host manipulation by parasites: A look back before moving forward. *Trends*
634   *in Parasitology*, **31**, 563-570. https://doi.org/10.1016/j.pt.2015.07.002

       Schaffer M, Winkelmann C, Hellmann C, Benndorf J (2013) Reduced drift activity of two benthic
636   invertebrate species is mediated by infochemicals of benthic fish. *Aquatic Ecology*, **47**, 99–
       107. https://doi.org/10.1007/s10452-013-9428-1

638   Svensson PA, Eghbal R, Eriksson R, Nilsson E (2022) How cunning is the puppet-master? Cestode-
       infected fish appear generally fearless. *Parasitology Research*, **121**, 1305–1315.
640   https://doi.org/10.1007/s00436-022-07470-2

       Tain L, Perrot-Minnot M-J, Cézilly F (2006) Altered host behaviour and brain serotonergic activity
642   caused by acanthocephalans: evidence for specificity. *Proceedings of the Royal Society B:*
       Biological Sciences, **273**, 3039–3045. https://doi.org/10.1098/rspb.2006.3618

644   Takahashi K (2022) Changes in the anxiety-like and fearful behavior of shrimp following daily
        threatening experiences. *Animal Cognition*, **25**, 319–327. https://doi.org/10.1007/s10071-
646   021-01555-8

    Vargha A, Delaney HD (2000) A Critique and Improvement of the CL Common Language Effect Size
648   Statistics of McGraw and Wong. *Journal of Educational and Behavioral Statistics*, **25**, 101–
        132.

650   Weinreich F, Benesh DP, Milinski M (2013) Suppression of predation on the intermediate host by two
        trophically-transmitted parasites when uninfective. *Parasitology*, **140**, 129–135.
652   https://doi.org/10.1017/S0031182012001266

    Williams DD, Moore KA (1985) The Role of Semiochemicals in Benthic Community Relationships of
654   the Lotic Amphipod *Gammarus pseudolimnaeus*: A Laboratory Analysis. *Oikos*, **44**, 280.
        https://doi.org/10.2307/3544701

656   Wudkevich K, Wisenden BD, Chivers DP, Smith RJF (1997) Reactions of *Gammarus lacustris* to
        Chemical Stimuli from Natural Predators and Injured Conspecifics. *Journal of Chemical
658   Ecology*, **23**, 1163–1173. https://doi.org/10.1023/B:JOEC.0000006393.92013.36