Leaf excision has minimal impact on photosynthetic parameters across crop functional types

John N. Ferguson[1,2*], Tamanna Jithesh[1†*], Tracy Lawson[2] and Johannes Kromdijk[1,3]

[1]Department of Plant Sciences, University of Cambridge, Cambridge, Cambridgeshire, CB2 3EA, UK

[2]School of Life Sciences, University of Essex, Wivenhoe Park, Colchester, Essex CO4 3SQ, UK

[3]Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, Illinois, 61801, USA

[†]Current address: Centre for Crop and Environmental Science, Harper Adams University, Edgmond, Shropshire TF10 8NB, UK

[*]Equal contribution

Highlight

Across the main photosynthesis functional types (C_3-dicot, C_3-monot, and C_4-monot) of major crops (tomato, barley, and maize), measurements of photosynthetic parameters demonstrate few, but important, differences when measured on excised relative to intact leaves.

Abstract

Photosynthesis is increasingly becoming a recognised target for crop improvement. Phenotyping photosynthesis-related traits on field-grown material is a key bottleneck to progress here due to logistical barriers and short measurement days. Many studies attempt to overcome these challenges by phenotyping excised leaf material in the laboratory. To date there are no demonstrated examples of the representative nature of photosynthesis measurements performed on excised leaves relative to intact leaves in crops. Here, we tested whether standardised leaf excision on the day prior to phenotyping impacted a range of common photosynthesis-related traits across crop functional types using tomato (C3-dicot), barley (C3-monocot), and maize (C4-monocot). Potentially constraining aspects of leaf physiology that could be forecasted to impair photosynthesis in excised leaves, namely leaf water potential and abscisic acid accumulation, were not different between intact and excised leaves. We also observed non-significant differences in spectral reflectance and chlorophyll fluorescence traits between the treatments across the three species. However, we did observe some significant differences between gas exchange and photosynthetic capacity associated traits across all three species. This study represents a useful reference for those who perform measurements of this nature and the differences reported should be considered in associated experimental design and statistical analyses.

Keywords

Leaf excision, photosynthesis, leaf reflectance, chlorophyll fluorescence, *Solanum lycopersicum*, *Hordeum vulgare*, *Zea mays*

Abbreviations

40	ABA	Abscisic acid
41	A_N	Net photosynthetic CO_2 assimilation
42	c_i	Intracellular CO_2 concentration
43	F_m	Dark adapted maximum fluorescence
44	F_m'	Maximum fluorescence
45	F_v/F_m	Maximum photosystem II efficiency
46	g_s	Stomatal conductance
47	$iWUE$	Intrinsic water use efficiency
48	J_{max}	Maximum rate of electron transport for RuBP regeneration
49	LWI	Leaf Water Index
50	MSI	Moisture Stress Index
51	NDNI	Normalised Difference Nitrogen Index
52	NPQ	Non-photochemical quenching
53	PAR	Photosynthetically active radiation
54	RE-NDVI	Red Edge-Normalised Difference Vegetation Index
55	V_{cmax}	Maximum rate of Rubisco carboxylation
56	V_{pmax}	Maximum rate of PEPc carboxylation
57	V_{max}	Maximum rate of photosynthesis

Introduction

The global demand for food is expected to double by the middle of this century. Human population growth means that the rate of increase in food demand outpaces the annual rate of increase in crop productivity (Zhu *et al.*, 2010; Kromdijk and Long, 2016; Lenaerts *et al.*, 2019). This mismatch highlights the necessity to adopt novel crop improvement targets to achieve food security (Gojon *et al.*, 2023). As the primary determinant of biomass accumulation, photosynthesis represents a sensible target to this end. Evidence from free air CO_2 enrichment experiments has highlighted how increasing CO_2 assimilation can lead to improved crop yields (Ainsworth and Long, 2021). Similarly, transgenic improvements to independent aspects of photosynthetic biochemistry have demonstrated that increasing carbon fixation is a realistic strategy for increasing yield (Driever *et al.*, 2017; Yoon *et al.*, 2020; De Souza *et al.*, 2022).

Targeting photosynthesis in crop improvement efforts necessitates the ability to screen many hundreds or thousands of genotypes for photosynthesis-associated traits. This is important for quantifying variation to incorporate in selection models. Additionally, it is necessary for facilitating forward genetics to identify novel genes or genetic regions underlying said variation to target through molecular approaches, such as marker-assisted breeding or direct manipulation through genome editing. This is challenging because the gold-standard methods through which many photosynthesis-associated traits are phenotyped are often logistically challenging to perform in field trial-like environments and they are time consuming (Long, 2003; Walker *et al.*, 2018). Standard methodologies, such as the commonly performed photosynthesis-CO_2 response (A-c_i) measurement used for modelling photosynthetic capacity can take up to 45-60 minutes to perform per sample. This is problematic because it limits the number of measurements that can be performed per day. This is especially true in the field relative to controlled conditions, where diurnal weather effects on physiological patterns, such as decreasing leaf water potential, photosystem II efficiency, and chloroplastic inorganic phosphate concentration, can transiently impact photosynthetic activity and the length of the measurement day (Murchie *et al.*, 1999; Leakey *et al.*, 2006). Logistical challenges arise, for example, due to specialised equipment not being suitable for use within crop canopies and because of the lack of access to power supplies.

A potential means through which to overcome some of these confounding effects and

logistical challenges is to excise leaf material from plants of interest and phenotype those

leaves in a more amenable and stable laboratory setting. Studies on woody species have

shown that phenotyping photosynthetic- and reflectance-associated traits on the leaves of

excised branches generates data that is highly comparable to that generated through

phenotyping leaves on branches still attached to the tree (Richardson and Berlyn, 2002;

Miyazawa *et al.*, 2011; Akalusi *et al.*, 2021). Over the past decade, a similar approach has

routinely been employed for multiple crop species (e.g., Driever *et al.*, 2014; Ferguson *et al.*,

2021; Montes *et al.*, 2022). Here, leaves, or tillers harbouring a leaf of interest, are excised

from plants and the cut end is placed into water in an attempt to maintain hydraulic

conductivity. For photo-physiological phenotyping, the maintenance of hydraulic efficiency is

important for replacing water lost via transpiration (Müllers *et al.*, 2022), which is

instrumental to keep stomata open for CO_2 uptake. It is possible that the demonstrable

success of employing this approach with woody species is at least in part due to the presence

of secondary xylem. Secondary xylem facilitates water transport, but it is also critical in

protecting the primary xylem from mechanical damage (Miodek *et al.*, 2021), such as that

that might ensue after branch excision. This presence of secondary xylem is a key anatomical

difference that distinguishes the tree species where this approach has been demonstrated to

work as effectively as *in-situ* phenotyping and the herbaceous annual crop species for which

this technique is now regularly employed. Due to this key difference and others, it is plausible

that the mechanistic basis that underpins the success of performing photosynthesis

measurements on foliage of detached branches of woody species does not necessarily

translate to herbaceous species. Consequently, a key knowledge gap exists here with respect

to understanding the extent to which measurements made on excised leaves are

representative of those made on intact leaves for crop species.

Addressing the above-described knowledge gap is a high-priority due to the rapid uptake of

this approach for phenotyping photosynthesis in crop species. Variations on this technique

have been reported in studies involving many key crop species, however we are not aware of

any studies that employ this approach and show that the data they obtain are representative

of what would have been obtained from intact material. Information to this end will be key

for emphasising the value of this approach. Maize is perhaps the species with which this

approach has been most employed. For example, Markelz et $al.$, (2011) excised leaves from maize plants growing in a free air CO_2 enrichment (FACE) experiment and under variable soil nitrogen availabilities to perform A-c_i response measurements to test interactive effects of CO_2 and nitrogen supply on photosynthetic capacity. Similarly, Liao et $al.$, (2022) phenotyped excised maize leaves from plants growing under full- and deficit-irrigation to again perform A-c_i response measurements to test how water availability influenced photosynthetic capacity and stomatal limitations to photosynthesis. As well as phenotyping small numbers of maize genotypes to test environmental interactions (See also: Leakey et $al.$, 2006) and Choquette et $al.$, (2019)), this approach has also been employed for maize to facilitate screening of large populations for forward genetics purposes. For example, Xie et $al.$, (2021) used this approach to phenotype a mapping population consisting of 197 recombinant inbred lines (RILs) for light-saturated gas exchange. By utilising this approach, this study was able to identify genetic regions underpinning variation in photosynthesis and associated traits. An identical approach has also been utilised for phenotyping genetic diversity in sorghum. Here, this approach enabled by-far the largest survey of genetic variation in photosynthesis, where over 800 genetically distinct sorghum varieties were characterised enabling genome- and transcriptome-wide association mapping (GWAS and TWAS) to pinpoint precise genes underlying variation in photosynthesis (Ferguson et $al.$, 2021). Phenotyping photosynthesis of excised leaves has also been used in other C_4 species, such as Miscanthus, where this approach has facilitated testing of the adaptation of photosynthesis to self-shading (Collison et $al.$, 2020), and sugarcane, where it has facilitated testing of the response of photosynthesis to the application of sugars (McCormick et $al.$, 2008).

Despite C_4 crops typically having more physiologically robust leaves, measuring photosynthesis in excised material has also been performed on studies focusing on monocot and dicot C_3 crops. For example, excised tillers were used to phenotype flag-leaves for a suite of photosynthesis-related traits in a diversity set of 64 wheat accessions at different developmental stages (Driever et $al.$, 2014; Carmo-Silva et $al.$, 2017). This approach has also been used in barley where the leaf of interest was directly excised at its base rather than harvesting the entire tiller to test how direct nitrate feeding impacted photosynthesis and sugar synthesis (Morcuende et $al.$, 2005). In rice, excised leaf material has also been used to measure chlorophyll fluorescence across distinct varieties to facilitate GWAS of fluorescence

parameters (Ferguson *et al.*, 2020). Similar fluorescence experiments on excised leaf

segments of wheat provided some preliminary evidence for an impact of the time between

excision and measurement, where a minimal waiting time may be needed to properly relate

observations to differences in photosynthetic efficiency (McAusland *et al.*, 2019). To the best

of our knowledge, utilisation of excised leaf material in C_3 dicots is constrained to studies on

soybean. For example, Morgan *et al.*, (2004) cut petioles of fully expanded leaves of soybean

plants growing in an open-air elevated ozone experiment to perform chlorophyll

fluorescence, A-c_i and photosynthesis light-response (AQ) measurements to understand the

photosynthetic response to ozone stress. The same technique was operated by Montes *et al.*,

(2022) to perform *A*-c_i measurements and concurrent measurements of hyperspectral

reflectance to build models for predicting photosynthetic capacity in the field and for

facilitating GWAS.

Excision plausibly elicits an abscisic acid (ABA) biosynthesis response in leaves in a manner

similar to that which promotes the stabilisation of leaf water status (Beardsell and Cohen,

1975) and/or resistance to herbivory (Vos *et al.*, 2013, 2019; Nguyen *et al.*, 2016). This is an

important consideration here since ABA accumulation within leaves also serves to close

stomatal apertures and reduce transpirational water loss in response to drought, which

consequently impairs carbon uptake (Blatt, 2000; Schroeder *et al.*, 2001). Further to this,

there may well be a myriad of minor-to-major alterations of leaf properties, e.g., changes to

xylem tension, cell turgor, leaf water potential, pigmentation, etc. (Govindjee *et al.*, 1981;

Carter, 1991; Canny, 1997; Richardson and Berlyn, 2002; Tyree and Cochard, 2003), that could

impact the photosynthetic activity of excised leaves. Due to the reality of these potential

pitfalls and because of our reliance on performing photosynthesis measurements on excised

leaf material in crop research, it is critical to understand whether leaf excision impacts

photosynthesis in crop species. To this end and with this study we tested how standardised

leaf excision on the day before phenotyping impacted light-saturated photosynthesis,

photosynthetic capacity, chlorophyll fluorescence, hyperspectral reflectance, ABA

accumulation and leaf water potential. For this we employed reference genotypes of tomato,

barley, and maize, which represent the main photosynthesis functional types of crops, i.e., C_3

monocot, C_3 dicot and C_4 monocot.

Materials and methods

Plant Material and Experimental Design

Reference genotypes for tomato, barley and maize were used for this study. These genotypes were M83 (tomato), Golden Promise (barley), and B73 (maize). All plants were grown at the Plant Growth Facility of the University of Cambridge. Seeds from all species were sown into modular trays before being transplanted into larger pots once the seedlings were established. Barley and tomato were grown in 1.5L sized pots. Maize was grown in 6L sized pots with two plants per plot. Barley was grown in multipurpose compost (M3 Compost, Levington Ltd., UK). Tomato was grown in a seedling and cuttings compost (F2 Compost, Levington Ltd., UK) and fertilised with specialised tomato feed (Tomorite, Levington Ltd., UK). Maize was grown in a 2:1 mix of these two compost types with added perlite. Tomato and maize were grown in the same growth room where the photoperiod was set to 14 hours. The light level was set to 600 μmol m^{-2} s^{-1} photosynthetically active radiation (PAR). The humidity was set to 65% relative humidity (RH). The air temperature was set to 28$°$C in the daytime and 20$°$C in the night-time. The growth room of the barley was set to a 16-hour photoperiod. The light level was set to 400 μmol m^{-2} s^{-1} PAR. The humidity was set to 60% RH. The air temperature was set to 22$°$C in the daytime and night-time.

Prior to any given measurement day, plants were transferred from the Plant Growth Facility to the laboratory at the Department of Plant Sciences (University of Cambridge) at 14.00 h. All plants were checked to be well watered and left on the laboratory bench. At 17.00 h, those plants that were to be measured as excised were cut. For tomato, the youngest fully expanded leaf was cut at the petiole below the two leaflets adjacent to the terminal leaflet. The petiole was then immediately placed under water and recut before being transferred to a 15 ml falcon tube under water. For barley, the youngest fully expanded leaf was excised at its base. The leaf was then recut just above the initial excision under water and transferred under water to a 15ml falcon tube. For maize, the youngest fully expanded leaf was excised at its base, recut just above the initial excision under water and transferred under water to a 50ml falcon tube. We ensured that all falcon tubes contained sufficient water so that the excision point was consistently submerged. Excised leaves were left on the same lab bench as their intact counterparts.

On the measurement day, excised and intact leaves were subject to a series of parallel
measurements as highlighted in Supplemental Figure S1, with details provided below. Tomato
plants were measured eight weeks post sowing across a two-day period. Barley plants were
measured across two separate two-day periods one week apart, four and five weeks post
sowing. Maize plants were measured across two separate two-day periods two weeks apart,
six and eight weeks post sowing.

Leaf-level gas exchange

Leaf-level gas exchange measurements were performed using two LI-COR 6400XT infra-red
gas analysers fitted with 6400-40 fluorometer LED light sources (LI-COR Biosciences, Lincoln,
NE). To measure light-saturated gas exchange, conditions within the gas exchange chamber
were set as follow: 25°C block temperature; 400 μmol s^{-1} air flow; 65-75% relative humidity
(RH); 400 μmol s^{-1} reference CO_2 concentration; 1500 μmol m^{-2} s^{-1} PAR. Once A_N and g_s were
stable, light saturated rates of gas exchange were logged before initiating a CO_2 response
curve as described below.

For the tomato and barley CO_2 response curves, all conditions were kept as described above
with gas exchange being measured at a series of reference CO_2 concentrations as follows:
400, 300, 200, 100, 50, 400, 400, 700, 1000, 1300, and 1800. Rates of gas exchange were
logged at each point according to stability criteria waiting a minimum of 90 seconds and a
maximum of 120 seconds at each point.

For the maize CO_2 response curves, all conditions were again kept as described for the light
saturated gas exchange. Reference CO_2 was then increased in the following steps: 400, 600,
800, 1000, and 1250. Reference CO_2 was then returned to 400 and rates of A_N and g_s were
allowed to re-stabilise before reducing reference CO_2 in the following sequence: 400, 300,
250, 200, 100, 75, and 25. Rates of gas exchange were logged at each point according to
stability criteria waiting a minimum of 60 seconds and a maximum of 90 seconds at each
point.

For tomato and barley, A_N-c_i data from the response curves were fit according to the FvCB model (Farquhar *et al.*, 1980) using the fitacis() function from the R package plantecophys (Duursma, 2015). This was achieved using the bilinear method to estimate the transition point. From this, estimates of the maximum rate of Rubisco carboxylation (V_{cmax}) and the maximum rate of electron transport for RuBP regeneration (J_{max}) on a C_i basis were obtained. For maize, A_N-c_i data from the response curves were fit using a custom R function following von Caemmerer, (2000). Here, the initial relationship between A_N and c_i was used to estimate the maximum rate of carboxylation by PEPC (V_{pmax}). Additionally, the horizontal asymptote of a four-parameter non-rectangular hyperbola was used as an estimate of CO_2- and light-saturated photosynthesis (V_{max}).

Hyperspectral reflectance

Hyperspectral reflectance measurements were performed using an ASD FieldSpec 4 Standard-Res Spectroradiometer equipped with a leaf-clip (Malvern Panalytical, Malvern, UK). The light source of the FieldSpec was allowed to warm up for 45 minutes prior to measurements. Hyperspectral reflectance was always measured on the same area of the leaf that gas exchange was performed on and always with the light source pointing towards the adaxial leaf surface. For each leaf, three technical replicates of hyperspectral reflectance were performed within the spot used for gas exchange measurements. Reflectance at each wavelength was then averaged across the three technical replicates for further data analyses. These measurements were performed immediately following the CO_2-response gas exchange measurements.

Chlorophyll fluorescence

Chlorophyll fluorescence measurements were performed using both the previously described LICOR 6400XT leaf chamber fluorometer and using a closed chlorophyll fluorescence imaging system (FluorCam FC 800-C, PSI, Czech Republic). For both, a similar program was designed to measure the induction of non-photochemical quenching (NPQ) in response to actinic light and the subsequent relaxation of NPQ following the actinic light being switched off.

Following the afternoon hyperspectral reflectance measurements, a portion of leaf was dark

adapted using aluminium foil for 45 minutes. After dark adaption, leaves were clamped within

the gas exchange chamber where conditions were as described previously except the light

source was switched off. Following estimation of dark-adapted maximum fluorescence (F_m),

the light source was switched on to 1500 µmol m^{-2} s^{-1} PAR for 10 minutes and a series of 12

saturating pulses were performed to estimate maximal fluorescence (F_m') at different time-

points. The light source was then subsequently switched off for 12 minutes and a further

series of eight saturating pulses were performed again to estimate F_m'. NPQ at each point of

measurement of chlorophyll fluorescence was then calculated as (F_m-F_m')/F_m'.

Samples for measurements of chlorophyll fluorescence in the closed system were first

analysed for leaf water potential using a Scholander pressure chamber (see below). For barley

and maize, leaf strips approximately 3cm x 5cm were taken from the middle portion of the

measured leaf and for tomato the same sized strip was from the measured terminal leaflet.

Leaf strips were arranged onto damp filter paper and encased between non-reflective glass

as per Ferguson *et al.*, (2020). These samples were then dark adapted overnight and

measured for NPQ induction and relaxation within the closed chlorophyll fluorescence system

following the same sequence of steps as described when using the LI-COR 6400 leaf chamber

fluorometer.

Using the NPQ data generated by both the leaf chamber fluorometer and the closed

fluorescence system, separate exponential models were fit to the light induction (Equation 1)

and dark relaxation (Equation 2) components.

$$\text{Equation 1:} \quad y = a * (1 - exp(-b * t)$$

$$\text{Equation 2:} \quad y = a * \left(exp(-b * t)\right) + c$$

For both equations a represents the amplitude of the induction or relaxation response (a_ind

and a_rel), b is the rate constant for the induction or relaxation of NPQ (b_ind and b_rel), c is

an offset to account for a non-zero intercept (relaxation only), t is the measurement time

point, and y is NPQ.

Leaf water potential

Measurements of leaf water potential were performed using a Scholander pressure chamber (Model 600, PMS Instrument Company, OR, USA) following manufacturer instructions. For intact leaves, a clean excision was made with a razor blade towards the base of the leaf for maize and barley or the petiole for tomato before immediately sealing the leaf into the chamber to measure the leaf water potential. For excised leaves, a new additional excision was made 3cm above the cut from the previous day before immediately measuring leaf water potential in the same manner.

ABA analysis

Separate plants were grown for analyses of leaf ABA concentration. Here, the plants were grown, and leaves were excised in exactly the same manner as described in the "Plant Material and Experimental Design" section. Barley and maize plants were measured five weeks post sowing and tomato plants were measured eight weeks post sowing. Samples for ABA concentration measurements were taken in the morning (10am). Each sample was taken from the middle portion of the leaf or terminal leaflet taking care to avoid the midrib in barley and maize. Samples were immediately flash frozen and subsequently analysed for ABA concentration using a competitive ELISA kit (Phytodetek Immunoassay Kit for ABA, Agdia, Elkhart, IN) following the manufacturer's instructions. The area of each sample was measured before flash freezing, so that ABA concentration could be expressed on leaf area basis.

Data processing and statistical analyses

All statistical analyses were performed within the R environment (Reference). All figures were generated using ggplot2 (Valero-Mora, 2010) and gridExtra with some pre-processing in Affinity designer (Serif, Nottingham, UK).

We performed two-way repeated measures analysis of variance (ANOVA) comparison of means tests to test for significance of the treatment (excised or intact), the time of measurement (AM or PM), and their interaction (treatment by time) for all gas exchange-associated traits. We performed one-sample t-tests to test for differences in leaf water potential and the chlorophyll fluorescence associated traits between excised and intact leaves

for each species independently. The chlorophyll fluorescence traits measured using the gas

exchange leaf chamber fluorometer and the closed chlorophyll fluorescence system were

analysed independently as instrumental differences in sample conditioning, excitation

wavelength as well as detection fore optics preclude direct comparisons.

The asdreader R package was used for managing the hyperspectral reflectance data following

best practise outlined in Burnett *et al.*, (2021). Hyperspectral reflectance data were initially

analysed through a spectra-wide approach where one-way ANOVA comparison of means

tests were performed to test for differences in reflectance between excised and intact leaves

at each wavelength from 250-3500nm. Separate tests were performed for each species and

each time point, i.e., AM and PM. We calculated a Bonferroni significance threshold ($\alpha = 0.05$)

to avoid identification of false positive results. It is worth noting that since there is covariation

between wavelengths of a given spectra, this threshold is likely to be somewhat stringent. For

visualisation purposes, we transformed the p-value associated with each wavelength by -

$\log_{10}$.

We additionally used the same hyperspectral reflectance data to calculate multiple

reflectance indices (Supplemental Table S1). We used these indices to perform repeated

measures two-way ANOVA comparison of means tests separately for each species and as

described previously to test for effects due to treatment, timepoint, and their interaction.

Results

Leaf water potential and ABA accumulation

At the end of the measurement day, we measured leaf water potential of all measured leaves

(Figure 1, Supplemental Figure S1). For tomato and maize, we did not observe a significant

effect of leaf excision on leaf water potential. However, for barley the leaf water potential of

excised leaves showed a small but significant difference with that of the intact leaves (-0.193

and -0.237 MPa, respectively; Figure 1b).

Leaf ABA concentration in excised and intact leaves was measured in the morning after the

excision. For all three species, leaf excision did not result in a significant up- or down-

regulation in ABA accumulation relative to intact leaves at this timepoint (Figure 2).

Light saturated gas exchange

Light-saturated gas exchange was logged prior to the initiation of the A_N-c_i response

measurements in the morning and the afternoon (Supplemental Figure S1, Figure 3). Across

all three species, leaf excision did not have a significant effect on A_N. The timing of the

measurement did have a significant effect on A_N for barley and maize, but not tomato. Here,

A_N reduced in the afternoon for barley and maize. This reduction in A_N was independent of

the treatment, where A_N declined in both intact and excised leaves (Figure 3e, i). Neither

treatment, time, nor their interaction had a significant effect on g_s for tomato (Figure 3b).

Time had a significant effect on g_s for barley, where it declined into the afternoon as per A_N

(Figure 3f). The excision treatment did have a significant effect on gs for maize, where it was

reduced in excised leaves. Although the treatment by time interactive effect for g_s was not

significant, the difference between excised and intact leaves did appear to increase into the

afternoon (Figure 3j). We did not detect a significant effect of treatment, time, or their

interaction on c_i for barley (Figure 3g). However, for tomato, c_i increased significantly in the

afternoon for excised and intact leaves (Figure 3c). For maize, all tested independent variables

had a significant effect on c_i. Intact leaves demonstrated higher c_i than excised leaves in the

morning and the afternoon, but the difference increased in the afternoon (Figure 3k).

Significant effects on *iWUE* mirrored those observed for c_i, such that it significantly declined

in the afternoon for both tomato and maize, and it was significantly increased in excised

leaves of maize (Figure 5d, l). With respect to the diurnal effect on *iWUE* and c_i for maize, it is

worth noting that the changes into the afternoon are much stronger in intact leaves, i.e.,

iWUE and c_i are more stable across the day in excised leaves.

Photosynthetic capacity

We modelled A_N-c_i response measurements to estimate photosynthetic capacity in the

morning and the afternoon of the three species (Supplemental Figure S1, Figure 4). Across

the three species, time had a significant effect on all modelled parameters except V_{pmax} in

maize (Figure 4k). In all other cases, the photosynthetic capacity parameters declined into the

afternoon regardless of whether the leaf being measured was excised or intact (Figure 4c, d,

g, h, l). We only detected one significant treatment effect, where V_{cmax} was significantly

reduced in excised barley leaves, however the reduction appeared relatively minor (Figure

4g). In addition, the p-value associated with the treatment effect for V_{pmax} in maize was close

to the traditional 5% significance threshold (Figure 4k). For all traits and species, we did not

observe a significant interaction effect of treatment and time, suggesting that any observed

effects were temporally consistent (Figure 4).

Leaf reflectance

Hyperspectral reflectance was measured following the A_N-c_i response measurements in the

morning and the afternoon (Supplemental Figures S1, S2). None of the specific wavelengths

showed any difference in reflectance between excised and intact leaves in the morning or the

afternoon for any species according to the Bonferroni significance threshold (Figure 5).

Despite the lack of significant wavelength-treatment associations, it is still interesting to

observe the distinct species-specific patterns in spectra-wide reflectance differences between

the treatments which can be interpreted based on the dominant factors that are known to

control leaf reflectance across the hyperspectral range. The pattern of p-values across the

spectral range were similar in the morning and the afternoon for tomato (Figure 5a-b) and

maize (Figure 5e-f), however the pattern was distinctly different across these time points for

barley (Figure 5c-d). For tomato, p-values showed a general trend of decreasing (inverse for -

$\log_{10}$(p-values)) from 300 to 1800nm before increasing between 1800 and 1900nm. After this,

they decreased to roughly what they were before the increase before increasing again toward

the end of the spectral range (Figure 5a-b). This pattern suggests potential differences in leaf

water status between the excised and intact tomato leaves. For maize, the pattern of p-values

across the spectral range partially-resembled the pattern of reflectance (Figure 5e-f;

Supplemental Figure S2). Here, there was a broad peak of low p-values between 600 and

1400nm and a narrower peak between 1500 and 1800nm. Here, the initial peak may reflect

differences in leaf structure and the secondary peak is often associated with differences in

leaf water status. For barley in the morning, the pattern of p-values across the spectral range

was similar to that of maize reflecting potential effects of excision on leaf structure and water

absorption (Figure 5c, e, f). However, the initial portion of the range was also characterised

by comparatively reduced p-values, especially around 400nm (Figure 5c), which may suggest

an effect on the absorption of light by chlorophyll b. In the afternoon, the broad peak of low
p-values between 600 and 1400nm disappeared, but there were still minor peaks between
600 and 700nm and between 1800 and 2000nm (Figure 5d), suggesting potential minor
effects on chlorophyll absorption and water status respectively.

Using the hyperspectral reflectance data (Supplemental Figure S2), we additionally calculated
reflectance indices related to nitrogen status, chlorophyll content and leaf water status
(Supplemental Table S1). We examined how these were affected by leaf excision and time of
measurement, i.e., AM and PM (Supplemental Figure S1), via two-way repeated measures
ANOVA tests as per the previously described photosynthesis traits (Supplemental Figures S4-
6). Although this approach did find some significant differences, the observed effects were
minor in all cases.

For tomato, we did not observe any significant treatment, time, or treatment by time
interactions for any of the reflection indices. (Supplemental Figures S3). For barley, we
observed a significant time effect for the Datt Index, which is a proxy of total chlorophyll
content, where it was higher in the afternoon (Supplemental Figure S4b). For barley, we also
observed a significant interactive effect for the moisture stress index (MSI), which is an
inverted proxy for water content, where excised leaves had a higher MSI in the morning only,
which suggest their water content was reduced compared to intact leaves at this point but
not again in the afternoon (Supplemental Figure S4d). For maize, the only significant effect
observed was a treatment effect for the Datt Index, which was marginally higher in intact
leaves compared to excised leaves suggesting chlorophyll content was reduced following leaf
excision (Figure S5b) .

Chlorophyll fluorescence
Chlorophyll fluorescence was measured using the leaf chamber fluorometer that was paired
to the gas exchange platform and on the subsequent day using the closed chlorophyll
fluorescence system (Supplemental Figure S1). We tested for significant differences between
samples taken from excised and intact leaves for F_v/F_m, the rates of induction and relaxation
of NPQ (b_ind and b_rel), and maximum NPQ. For the samples measured using the closed
chlorophyl fluorescence system, we did not observe a significant difference for any of these

four parameters (Figure 6). This was true for all of these same parameters as measured in the

leaf chamber fluorometer, expect for tomato b_ind, which was observed to be significantly

faster in intact versus excised leaves (Supplemental Figure S6b). It is worth noting that the p-

values associated with the one-sample t-tests comparing excised and intact tomato leaves for

F_v/F_m and maximum NPQ were close to the traditional 0.05 significance threshold (0.07 and

0.09 respectively), where intact leaves appeared to have higher photosystem II maximum

efficiency and light adapted NPQ (Supplemental Figure S5b).

Discussion

Screening variation in photosynthetic traits across large numbers of genotypes is a key

requirement for targeting photosynthesis for crop improvement. Large-scale screening to this

end is fraught with logistical challenges that are commonly circumvented via phenotyping

excised leaf material (See: Introduction). A common limitation of these studies is the lack of

evidence highlighting the representative nature of such phenotyping, which accordingly

represents a key knowledge gap for this field. To this end, the principal aim of this study was

to ascertain if photosynthetic parameters measured on excised leaves were representative of

the same parameters measured on intact leaves in tomato, barley, and maize. There are at

least three studies that have attempted to quantify potential differences in non-crop species.

Using the Japanese stone oak (*Lithocarpus edulis*) evergreen tree species, Miyazawa *et al.*,

(2011) showed that multiple photosynthetic parameters demonstrated no significant

differences when measured on excised leaves relative to intact leaves. Here, the only trait

that did demonstrate a significant difference was stomatal conductance, which was

significantly higher in excised leaves. Despite this difference, the authors showed that it was

possible to use data from excised leaves to parametrise an ecophysiological model to

reproduce diurnal patterns of *in situ* gas exchange. In a similar study, Akalusi *et al.*, (2021)

showed there to be negligible differences in light saturated rates of gas exchange and

modelled estimates of photosynthetic capacity between excised and intact leaves of balsam

fir (*Abies balsamea*). Moreover, the authors also demonstrated that the non-significant

differences between these measurement types carried through across the growing season. In

contrast to these two studies, Richardson and Berlyn (2002) focused on how spectral

reflectance and chlorophyll fluorescence parameters were impacted by leaf excision in four

temperate tree species. They observed that spectral and chlorophyll fluorescence changes
were only minor at 12-hours post excision. Furthermore, they demonstrated that with
appropriate handling to maintain leaf water status, representative data could be obtained
from excised leaves as late as three days-post excision depending on the species. These
studies have demonstrated the effectiveness of phenotyping traits related to photosynthesis
on detached leaves of perennial woody species. As we highlight in the introduction, a similar
approach is now commonly employed for many crop studies. The numerous differences that
distinguish herbaceous annuals from trees suggest that the success of this approach may not
carry over to crop species, thereby highlighting the timeliness of this present study. Our
findings here largely provide long overdue experimental justification for applying this
approach on three major crop species, representing the major monocots and dicot groups as
well as C4 and C3 photosynthesis. The lack of large effects due to excision, as well as the
occurrence of some small but significant effects will be discussed in more detail in the next
sections.
511
Stabilised leaf water potential and ABA accumulation underlie the utility of phenotyping
photosynthesis in excised leaves
The most important crop species for staple foods are herbaceous annuals. A key anatomical
difference between woody and herbaceous species is the composition of xylem tissue.
Primary xylem is the only form of xylem for most herbaceous plants, including cereals (Růžička
et al., 2015), although tomato can demonstrate a small amount of secondary xylem (Qi *et al.*,
2020). While primary xylem comprises a relatively thin central zone within stems and leaves,
secondary xylem grows out and differentiates from the vascular cambium (Kirst *et al.*, 2004).
Secondary xylem serves multiple purposes beyond facilitating water transport, including
biomechanical support and protection against physical damage (Sperry, 2003; Morris *et al.*,
2016; Miodek *et al.*, 2021). Avoiding xylem damage is critical for maintaining hydraulic
conductivity and protecting photosynthesis. Breakage of the water column through xylem
cavitation can disrupt this. Our results are supportive of the notion that the transpiration
stream of excised tomato, barley, and maize leaves is maintained following recutting them
underwater as we did not observe negative effects on leaf water status measured as leaf
water potential (Figure 1). The utility of performing photosynthesis on excised leaves from
woody species may be supported by the presence of secondary xylem tissue, since these

measurements are typically performed by excising woody branches attached to leaves of
interest. Furthermore, vulnerability segmentation work has shown that branches are more
resistant to xylem cavitation than terminal leaves (Zhu *et al.*, 2016). Stem and leaf hydraulic
conductance are coordinated (Pivovaroff *et al.*, 2014), therefore excising leaves may remove
the association of leaf and stem hydraulic conductance. It is plausible that the reduction in
leaf water potential of intact barley leaves (Figure 1b) is a function of this and that by
detaching the leaf from the stem and immersing it in water, this process increases water
availability by removing the hydraulic resistance imposed by the stem. Alternatively, this
difference may suggest that the plants attached to the measured leaves were marginally
water stressed, however this is unlikely given the absolute water potential values were
greater than -0.25MPa since we ensured that all plants were well-watered throughout
growing and on the day before measurements. Moreover, the barley light-saturated gas
exchange data did not demonstrate a signal of water stress in the intact leaves relative to the
excised leaves (as discussed below). In general, the values for water potential obtained here
and the minor difference between excised and intact leaves suggests a lack of water stress.
544
It has been suggested that roots are essential for stimulating ABA biosynthesis in leaves
(Takahashi *et al.*, 2018), however many studies that involve work with excised leaves have
demonstrated that ABA levels can increase independently of roots (Bauer *et al.*, 2013;
Sussmilch *et al.*, 2017; McAdam and Brodribb, 2018). It is therefore plausible that any
treatment effects observed as a result of leaf excision could be imparted due to foliar ABA
accumulation. Depending on the site and method of excision, the act of cutting a leaf or a
tiller from a plant may elicit a wounding-type response. ABA can accumulate in response to
wounding as a regulator of downstream defense mechanisms (Cui *et al.*, 2013; Savatin *et al.*,
2014). Additionally, the process of excision is likely to impair xylem tension influencing the
flow rate of water to the leaf. This may also result in ABA accumulation in the leaf to stabilise
leaf water status by reducing transpiration, however there no evidence presently exists to
support this notion. Moreover, it is likely that the subsequent recutting of leaves underwater
as per this study and other studies will quickly re-establish xylem tension by mitigating against
cavitation. The role of ABA in regulating ion transport into guard cells to induce stomatal
closure is well characterised (Schroeder *et al.*, 2001), consequently if leaf excision were to
induce leaf ABA accumulation through any basis, we could expect to see associated

reductions in gas exchange as result of stomatal closure. To this end, our results

demonstrated that in the morning following leaf excision there was no significant difference

in the leaf ABA concentration between intact and excised leaves in either of three species

(Figure 2). Performing leaf excision on the day prior to phenotyping was premeditated as we

hypothesised that this would allow sufficient time for degradation of any additionally

accumulated ABA. So, while our results do not allow us to determine whether excision

stimulates an immediate ABA response, they are supportive of a lack of differential

accumulation of ABA in leaves that are excised on the day prior to measurements.

Consequently, we recommend adopting this approach when phenotyping excised leaves to

avoid any confounding effects that may potentially arise due to ABA accumulation.

Differences between excised and intact leaves for gas exchange traits are limited, but

important and species-specific

Our methodology of leaf excision mirrors a widespread approach utilised by many research

groups for performing photosynthesis-related phenotyping of material that is difficult to

measure *in-situ* for a multitude of reasons. This approach is ultimately necessary and defines

the ability to perform large scale screening of diversity that is important to characterise.

Despite this, leaf excision may represent a significant shock to the leaf and plausibly elicit

substantial wounding responses. Prior to this study, little data existed to support the

efficiency of this approach for generating data matching that obtained from intact leaves. To

this end, perhaps the most critical finding of our study is that we did not observe a significant

effect of leaf excision on A_N. This was true for all three species tested and across the morning

and afternoon measurements (Figure 3a, e, i). This is an important finding and goes a long

way to highlight the utility of this approach for assessing light-saturated photosynthesis at

ambient CO_2. While we did not observe significant treatment effects on A_N, we did observe

significant time of measurement effects for barley and maize, where A_N declined into the

afternoon (Figure 1e, i). This is indicative of the diurnal pattern of photosynthesis where, in

many species, there is a decline in A_N into the afternoon (Singsaas *et al.*, 2000; Mohotti and

Lawlor, 2002; Ainsworth *et al.*, 2007; Koester *et al.*, 2016; Vialet-Chabrand *et al.*, 2017). This

can be attributed to multiple factors including declining electron transport and changes to

the apparent sink strength due to increasing leaf carbohydrate content (Singsaas *et al.*, 2000;

Koester *et al.*, 2016). So, while this excision approach is supportive of phenotyping A_N, our

findings suggest that interpreting and comparing data obtained across an extended diurnal

period necessitates caution regardless of whether it was collected from intact or excised

leaves.

A further, and curious, significant time of day effect was detected for c_i in tomato (Figure 3c).

Here, c_i increased into the afternoon for both treatments despite no significant changes in A_N

or g_s that could be presupposed to lead to increases in c_i. This result may be a function of

lower Rubisco activity, which is suggested by declining V_{cmax} in tomato for both treatments

into the afternoon (Figure 4c). Here, lower carboxylation by Rubisco may lead to an increase

in CO_2 in the intracellular air space of the leaf in the afternoon compared to the morning,

especially since g_s was not observed to change across these time periods. This result highlights

the importance of quantifying photosynthetic capacity to generate a broader picture of the

photo-physiological state of the leaf, as opposed to relying on light-saturated measurements

made under ambient CO_2. Declining V_{cmax} in the afternoon for tomato here may be a function

of the diurnal expression patterns of Rubisco activase (Rca). Rca is the molecular chaperone

that defines Rubisco activity, and its expression has been demonstrated to decline

significantly into the afternoon (Perdomo *et al.*, 2021). The afternoon decline in tomato *iWUE*

(Figure 3d) mirrors the increase in c_i (Leakey *et al.*, 2019) and is a function of the combination

of the decline in A_N (Figure 3b) and increase in g_s (Figure 3c), despite those changes being

non-significant in isolation.

One of the more surprising results from our study was that maize was the only one of the

three species where we detected a treatment effect on g_s (Figure 3j). Here, the excision

process resulted in a significant decline in g_s compared to intact leaves. This result is

somewhat surprising since maize leaves are larger and appear more physically robust than

barley leaves and the petiole of tomato leaves. Nevertheless, we ascribe confidence to this

result since it was consistent across the two time points, therein highlighting the impairing

influence of leaf excision on stomatal opening in maize. The C_4 carbon concentrating

mechanism that defines the relative de-coupling of g_s and A_N in C_4 species relative to C_3

species (Leakey *et al.*, 2019) may underly why we do not observe a concurrent treatment

effect on A_N in maize. Namely, the decrease in g_s observed in the excised leaves was not

sufficient to decrease c_i enough to lose CO_2 saturation of A_N (Figure k) as evident from the

A_N/c_i responses (Figure 3j). The treatment effect of leaf excision on g_s in maize but not on A_N,

led to expected results for c_i, where it decreased significantly at both time points due to

decreased capacity for CO_2 uptake (Figure 3k). Concurrently, the stomatal resistance to water

loss was greater, such that *iWUE* was higher at both time points and more so again in the

afternoon (Figure 3l). Taken together, these results show that the time-of-day effects on light-

saturated rate of gas exchange are stronger in intact maize leaves, thereby highlighting the

potential for longer measurement days when working with excised leaves.

Despite the above-described differences observed in maize, we did not observe any

significant effects in terms of treatment or time on the photosynthetic capacity traits, namely

V_{pmax} and V_{max} (Figure 2k, l). This suggests that reduced g_s at the starting point of the A_N-c_i

response measurements, i.e., ambient CO_2, in excised leaves did not compromise the ability

to assess the rate of photosynthetic induction from low-to-ambient CO_2 concentrations or the

rates of photosynthesis when g_s is minimal at high CO_2 concentrations. Our results are

therefore supportive of performing A_N-c_i response curves in excised maize leaves to model

photosynthetic capacity. In general, our results are also supportive of performing A_N-c_i

response curves on excised leaves of the C_3 species. We did detect significant reductions in

V_{cmax} (barley; Figure 4g) but the difference was minor and not exaggerated by the interaction

with time.

Reflectance and chlorophyll fluorescence appear relatively unchanged between intact and

excised leaves

There were no statistically significant differences detected for leaf reflectance between the

treatments for any of the species on a wavelength-by-wavelength basis (Figure 5). Despite

this, the pattern of p-values across the spectrum were surprisingly unique between the

species and different between the time points for barley. In general, if we consider the

dominant factors that influence leaf reflectance (Kokaly *et al.*, 2003), the p-value patterns

suggest the most excision-sensitive aspects of reflectance were due to leaf water content in

tomato and light scattering in the mesophyll in barley and maize (Figure 5). The results from

barley and maize here are similar to those observed across certain tree species by Richardson

and Berlyn, (2002), where the authors and others suggest reductions in near-infra red spectra

reflectance in excised leaves are related to leaf structural changes at the cellular level. Here,

declining leaf water status results in cytoplasmic shrinkage, such that light scattering alters at

the interface between cell walls and water (Carter, 1991; Aldakheel and Danson, 1997;

Richardson and Berlyn, 2002). However, we do not observe perturbed leaf water status in

excised leaves (Figure 1), consequently this may reflect why the patterns we highlight in terms

of differences in reflectance between the treatments do not pass significance thresholds

(Figure 5). The study of Richardson and Berlyn, (2002) tested differences between excised and

intact leaves over several days post-excision and did not begin to detect significant differences

in reflectance indices and chlorophyll fluorescence until 48 hours after excision.

Consequently, it is possible that we may have observed differences in reflectance and other

traits had we measured for a further day, however this is not common practice with crop

species, and we would advise against this.

To resolve any potential reflectance differences between intact and excised leaves, we

calculated five commonly employed reflectance indices (Supplemental Table S1,

Supplemental Figures S3-5). In doing so, we were only able to detect two statistically

significant differences between intact and excised leaves. For barley, the MSI was significantly

higher in the morning in excised leaves suggesting a reduction in leaf water content, however

this effect was not apparent in the afternoon (Supplemental Figure S4d). In maize, a

significant treatment effect on the Datt index, which is positively correlated to chlorophyll

content (Lu *et al.*, 2015), suggested that chlorophyll content was reduced in excised leaves

(Supplemental Figure S5b). Here, it could be hypothesised that ABA accumulation would be

an the underlying mechanism resulting in this difference, since there is also some evidence

linking leaf yellowing and reductions in leaf chlorophyll to ABA, however that is evidently not

the situation in our study (Figure 2) and this is typically a longer-term effect (Wang *et al.*,

2018). In general, our results here suggest that reflectance changes between excised and

intact leaves are slight, which is similar to what has previously been shown in woody species

(Richardson and Berlyn, 2002).

Our results also mirror those of Richardson and Berlyn, (2002) in that we saw no differences

in F_v/F_m in either of methods through which it was assessed in this study (Figure 6 and

Supplemental Figure S6). This highlights how electron flow in PSII is not significantly impacted

by this approach, so there is no impairment on the photosynthetic apparatus. In our study we

689 also demonstrated no significant differences in the dynamics of NPQ, expect for a minor

690 reduction in the rate of induction in excised tomato leaves when measured using the gas

691 exchange leaf chamber fluorometer (Supplemental Figure S6a). The fact that this was not

692 detected the following day using the closed chlorophyll fluorescence system suggests that

693 this is more likely to result from insufficient time allowed for dark adaption, especially since

694 F_v/F_m was also bordering on being significantly reduced in excised leaves when measured in

695 the leaf chamber fluorometer also (p = 0.07; Supplemental Figure S6a).

697 ***Conclusion***

698 Screening large amounts of diversity for photosynthesis-related traits is only realistically

699 possible with excised leaves. Therefore, the evidence presented in this study that validates

700 this approach, in terms of its affiliation to intact-leaf measurements, is welcomed. Despite

701 this, our study does highlight the existence of specifies-specific treatment effects, e.g., g_s

702 reduction in maize, that the community should be aware of. Moreover, this should encourage

703 those who are planning on adopting this approach to validate if it is suitable for their species

704 of interest. One salient point that our study does not address is the potential for differential

705 responses to excision on an intra-specific basis. An obvious further extension to this study

706 would be to test how genetically distinct genotypes within a species perform following leaf

707 excision as this has the potential to have large consequences for studies of this nature.

Supplemental Material

710 Supplemental Figure S1. Overview of experimental workflow.

712 Supplemental Figure S2. Mean hyperspectral reflectance data. The solid line represents the

713 mean reflectance, and the shaded area represents the standard error of the mean. (a) Tomato

714 AM (b) Tomato PM (c) Barley AM (d) Barley PM (e) Maize AM (f) Maize PM

716 Supplemental Figure S3. Hyperspectral reflectance derived indices for tomato. (a) Red Edge-

717 Normalised Difference Vegetation Index (RE-NDVI), (b) Datt Index (c) Normalised Difference

718 Nitrogen Index (NDNI), (d) Moisture Stress Index (MSI), and (e) Leaf Water Index (LWI)

Acknowledgments

This work was supported by the European Union's Horizon2020 research and innovation

programme (No.862201) project CAPITALISE. We thank Dr Cristina Sales and Dr Angie Burnett

for assistance in generating the ABA data. We thank Dr Sara Lopez-Gomollon for providing

the tomato seed. We thank Katie Shaw for assistance in performing the modelling of the

chlorophyll fluorescence data and Prof Howard Griffiths and Dr Jessica Royles for assistance

with the leaf water potential measurements. For the purpose of open access, the authors

have applied a Creative Commons Attribution (CC BY) licence to any Author Accepted

Manuscript version arising from this submission.

Author contributions

744   JNF, TL and JK designed the study. JNF and TJ carried out the experiments. JNF and TJ

745   performed the data analyses. All authors helped to interpret the results. JNF and JK wrote the

746   paper with input from all the authors.

Conflict of interest

749   The authors have no conflicts of interest to declare.

References

Ainsworth EA, Long SP. 2021. 30 years of free-air carbon dioxide enrichment (FACE): What have we learned about future crop productivity and its potential for adaptation? Global Change Biology **27**, 27–49.

Ainsworth EA, Rogers A, Leakey ADB, Heady LE, Gibon Y, Stitt M, Schurr U. 2007. Does elevated atmospheric [CO2] alter diurnal C uptake and the balance of C and N metabolites in growing and fully expanded soybean leaves? Journal of Experimental Botany **58**, 579–591.

Akalusi ME, Meng F, P.-A. Bourque C. 2021. Photosynthetic parameters and stomatal conductance in attached and detached balsam fir foliage. Plant-Environment Interactions **2**, 206–215.

Aldakheel YY, Danson FM. 1997. Spectral reflectance of dehydrating leaves: Measurements and modelling. International Journal of Remote Sensing **18**, 3683–3690.

Bauer H, Ache P, Lautner S, *et al.* 2013. The stomatal response to reduced relative humidity requires guard cell-autonomous ABA synthesis. Current biology: CB **23**, 53–57.

Beardsell MF, Cohen D. 1975. Relationships between Leaf Water Status, Abscisic Acid Levels, and Stomatal Resistance in Maize and Sorghum. Plant Physiology **56**, 207–212.

Blatt MR. 2000. Cellular signaling and volume control in stomatal movements in plants. Annual Review of Cell and Developmental Biology **16**, 221–241.

Burnett AC, Anderson J, Davidson KJ, Ely KS, Lamour J, Li Q, Morrison BD, Yang D, Rogers A, Serbin SP. 2021. A best-practice guide to predicting plant traits from leaf-level hyperspectral data using partial least squares regression (T Lawson, Ed.). Journal of Experimental Botany **72**, 6175–6189.

von Caemmerer S. 2000. *Biochemical Models of Leaf Photosynthesis*. CSIRO Publishing.

Canny M. 1997. Vessel contents of leaves after excision - a test of Scholander's assumption. American Journal of Botany **84**, 1217.

Carmo-Silva E, Andralojc PJ, Scales JC, Driever SM, Mead A, Lawson T, Raines CA, Parry MAJ. 2017. Phenotyping of field-grown wheat in the UK highlights contribution of light response of photosynthesis and flag leaf longevity to grain yield. Journal of Experimental Botany **68**, 3473–3486.

Carter GA. 1991. Primary and Secondary Effects of Water Content on the Spectral Reflectance of Leaves. American Journal of Botany **78**, 916–924.

Chen D, Wang S, Xiong B, Cao B, Deng X. 2015. Carbon/Nitrogen Imbalance Associated with Drought-Induced Leaf Senescence in Sorghum bicolor. PLOS ONE **10**, e0137026.

Choquette NE, Ogut F, Wertin TM, Montes CM, Sorgini CA, Morse AM, Brown PJ, Leakey ADB, McIntyre LM, Ainsworth EA. 2019. Uncovering hidden genetic variation in

photosynthesis of field-grown maize under ozone pollution. Global Change Biology **25**, 4327–4338.

Collison RF, Raven EC, Pignon CP, Long SP. 2020. Light, Not Age, Underlies the Maladaptation of Maize and Miscanthus Photosynthesis to Self-Shading. Frontiers in Plant Science **11**, 783.

Cui F, Brosché M, Sipari N, Tang S, Overmyer K. 2013. Regulation of ABA dependent wound induced spreading cell death by MYB 108. New Phytologist **200**, 634–640.

De Souza AP, Burgess SJ, Doran L, Hansen J, Manukyan L, Maryn N, Gotarkar D, Leonelli L, Niyogi KK, Long SP. 2022. Soybean photosynthesis and crop yield are improved by accelerating recovery from photoprotection. Science **377**, 851–854.

Driever SM, Lawson T, Andralojc PJ, Raines CA, Parry MAJ. 2014. Natural variation in photosynthetic capacity, growth, and yield in 64 field-grown wheat genotypes. Journal of Experimental Botany **65**, 4959–4973.

Driever SM, Simkin AJ, Alotaibi S, Fisk SJ, Madgwick PJ, Sparks CA, Jones HD, Lawson T, Parry MAJ, Raines CA. 2017. Increased SBPase activity improves photosynthesis and grain yield in wheat grown in greenhouse conditions. Philosophical Transactions of the Royal Society B: Biological Sciences **372**, 20160384.

Duursma RA. 2015. Plantecophys - An R Package for Analysing and Modelling Leaf Gas Exchange Data. PLOS ONE **10**, e0143346.

Farquhar GD, von Caemmerer S, Berry JA. 1980. A biochemical model of photosynthetic CO_2 assimilation in leaves of C 3 species. Planta **149**, 78–90.

Ferguson JN, Fernandes SB, Monier B, *et al.* 2021. Machine learning-enabled phenotyping for GWAS and TWAS of WUE traits in 869 field-grown sorghum accessions. Plant Physiology **187**, 1481–1500.

Ferguson JN, McAusland L, Smith KE, Price AH, Wilson ZA, Murchie EH. 2020. Rapid temperature responses of photosystem II efficiency forecast genotypic variation in rice vegetative heat tolerance. The Plant Journal: For Cell and Molecular Biology **104**, 839–855.

Gojon A, Nussaume L, Luu DT, *et al.* 2023. Approaches and determinants to sustainably improve crop production. Food and Energy Security **12**.

Govindjee, Downton WJS, Fork DC, Armond PA. 1981. Chlorophyll A fluorescence transient as an indicator of water potential of leaves. Plant Science Letters **20**, 191–194.

Kirst M, Myburg AA, De León JPG, Kirst ME, Scott J, Sederoff R. 2004. Coordinated Genetic Regulation of Growth and Lignin Revealed by Quantitative Trait Locus Analysis of cDNA Microarray Data in an Interspecific Backcross of Eucalyptus. Plant Physiology **135**, 2368–2378.

Koester RP, Nohl BM, Diers BW, Ainsworth EA. 2016. Has photosynthetic capacity increased with 80 years of soybean breeding? An examination of historical soybean cultivars. Plant, Cell & Environment **39**, 1058–1067.

Kokaly RF, Despain DG, Clark RN, Livo KE. 2003. Mapping vegetation in Yellowstone National Park using spectral feature analysis of AVIRIS data. Remote Sensing of Environment **84**, 437–456.

Kromdijk J, Long SP. 2016. One crop breeding cycle from starvation? How engineering crop photosynthesis for rising CO_2 and temperature could be one important route to alleviation. Proceedings of the Royal Society B: Biological Sciences **283**, 20152578.

Leakey ADB, Ferguson JN, Pignon CP, Wu A, Jin Z, Hammer GL, Lobell DB. 2019. Water Use Efficiency as a Constraint and Target for Improving the Resilience and Productivity of C3 and C4 Crops. Annual Review of Plant Biology **70**, 781–808.

Leakey ADB, Uribelarrea M, Ainsworth EA, Naidu SL, Rogers A, Ort DR, Long SP. 2006. Photosynthesis, Productivity, and Yield of Maize Are Not Affected by Open-Air Elevation of CO2 Concentration in the Absence of Drought. Plant Physiology **140**, 779–790.

Lenaerts B, Collard BCY, Demont M. 2019. Review: Improving global food security through accelerated plant breeding. Plant Science **287**, 110207.

Liao Q, Gu S, Kang S, Du T, Tong L, Wood JD, Ding R. 2022. Mild water and salt stress improve water use efficiency by decreasing stomatal conductance via osmotic adjustment in field maize. Science of The Total Environment **805**, 150364.

Long SP. 2003. Gas exchange measurements, what can they tell us about the underlying limitations to photosynthesis? Procedures and sources of error. Journal of Experimental Botany **54**, 2393–2401.

Lu S, Lu X, Zhao W, Liu Y, Wang Z, Omasa K. 2015. Comparing vegetation indices for remote chlorophyll measurement of white poplar and Chinese elm leaves with different adaxial and abaxial surfaces. Journal of Experimental Botany **66**, 5625–5637.

Makino A, Shimada T, Takumi S, Kaneko K, Matsuoka M, Shimamoto K, Nakano H, Miyao-Tokutomi M, Mae T, Yamamoto N. 1997. Does Decrease in Ribulose-1,5-Bisphosphate Carboxylase by Antisense RbcS Lead to a Higher N-Use Efficiency of Photosynthesis under Conditions of Saturating CO2 and Light in Rice Plants? Plant Physiology **114**, 483–491.

Markelz RJC, Strellner RS, Leakey ADB. 2011. Impairment of C4 photosynthesis by drought is exacerbated by limiting nitrogen and ameliorated by elevated [CO2] in maize. Journal of Experimental Botany **62**, 3235–3246.

McAdam SAM, Brodribb TJ. 2018. Mesophyll Cells Are the Main Site of Abscisic Acid Biosynthesis in Water-Stressed Leaves. Plant Physiology **177**, 911–917.

McAusland L, Atkinson JA, Lawson T, Murchie EH. 2019. High throughput procedure utilising chlorophyll fluorescence imaging to phenotype dynamic photosynthesis and photoprotection in leaves under controlled gaseous conditions. Plant Methods **15**, 109.

McCormick AJ, Cramer MD, Watt DA. 2008. Changes in Photosynthetic Rates and Gene Expression of Leaves during a Source–Sink Perturbation in Sugarcane. Annals of Botany **101**, 89–102.

Miodek A, Gizińska A, Włoch W, Kojs P. 2021. What do we know about growth of vessel
elements of secondary xylem in woody plants? Biological Reviews **96**, 2911–2924.

Miyazawa Y, Tateishi M, Komatsu H, Kumagai T, Otsuki K. 2011. Are measurements
from excised leaves suitable for modeling diurnal patterns of gas exchange of intact leaves?
Hydrological Processes **25**, 2924–2930.

Mohotti AJ, Lawlor DW. 2002. Diurnal variation of photosynthesis and photoinhibition in
tea: effects of irradiance and nitrogen supply during growth in the field. Journal of
Experimental Botany **53**, 313–322.

Montes CM, Fox C, Sanz-Sáez Á, *et al.* 2022. High-throughput characterization,
correlation, and mapping of leaf photosynthetic and functional traits in the soybean (*Glycine*
max) nested association mapping population (T Juenger, Ed.). Genetics, iyac065.

Morcuende R, Kostadinova S, Pérez P, Martínez-Carrasco R. 2005. Fructan synthesis is
inhibited by phosphate in warm-grown, but not in cold-treated, excised barley leaves. The
New Phytologist **168**, 567–574.

Morgan PB, Bernacchi CJ, Ort DR, Long SP. 2004. An in vivo analysis of the effect of
season-long open-air elevation of ozone to anticipated 2050 levels on photosynthesis in
soybean. Plant Physiology **135**, 2348–2357.

Morris H, Plavcová L, Cvecko P, *et al.* 2016. A global analysis of parenchyma tissue
fractions in secondary xylem of seed plants. New Phytologist **209**, 1553–1565.

Müllers Y, Postma JA, Poorter H, van Dusschoten D. 2022. Stomatal conductance tracks
soil-to-leaf hydraulic conductance in faba bean and maize during soil drying. Plant
Physiology **190**, 2279–2294.

Murchie EH, Chen Y, Hubbart S, Peng S, Horton P. 1999. Interactions between
Senescence and Leaf Orientation Determine in Situ Patterns of Photosynthesis and
Photoinhibition in Field-Grown Rice1. Plant Physiology **119**, 553–564.

Nguyen D, Rieu I, Mariani C, van Dam NM. 2016. How plants handle multiple stresses:
hormonal interactions underlying responses to abiotic stress and insect herbivory. Plant
Molecular Biology **91**, 727–740.

Perdomo JA, Buchner P, Carmo-Silva E. 2021. The relative abundance of wheat Rubisco
activase isoforms is post-transcriptionally regulated. Photosynthesis Research **148**, 47–56.

Pivovaroff AL, Sack L, Santiago LS. 2014. Coordination of stem and leaf hydraulic
conductance in southern C alifornia shrubs: a test of the hydraulic segmentation hypothesis.
New Phytologist **203**, 842–850.

Qi X, Takahashi H, Kawasaki Y, *et al.* 2020. Differences in xylem development between
Dutch and Japanese tomato (Solanum lycopersicum) correlate with cytokinin levels in
hypocotyls. Annals of Botany **126**, 315–322.

Richardson AD, Berlyn GP. 2002. Changes in foliar spectral reflectance and chlorophyll
fluorescence of four temperate species following branch cutting. Tree Physiology **22**, 499–
506.

Růžička K, Ursache R, Hejátko J, Helariutta Y. 2015. Xylem development – from the
cradle to the grave. New Phytologist **207**, 519–535.

Savatin DV, Gramegna G, Modesti V, Cervone F. 2014. Wounding in the plant tissue: the
defense of a dangerous passage. Frontiers in Plant Science **5**.

Schroeder JI, Allen GJ, Hugouvieux V, Kwak JM, Waner D. 2001. GUARD CELL
SIGNAL TRANSDUCTION. Annual Review of Plant Physiology and Plant Molecular
Biology **52**, 627–658.

Singsaas EL, Ort DR, DeLUCIA EH. 2000. Diurnal regulation of photosynthesis in
understory saplings: Diurnal regulation of photosynthesis. New Phytologist **145**, 39–49.

Sperry JS. 2003. Evolution of Water Transport and Xylem Structure. International Journal of
Plant Sciences **164**, S115–S127.

Sussmilch FC, Brodribb TJ, McAdam SAM. 2017. Up-regulation of NCED3 and ABA
biosynthesis occur within minutes of a decrease in leaf turgor but AHK1 is not required.
Journal of Experimental Botany **68**, 2913–2918.

Takahashi F, Suzuki T, Osakabe Y, Betsuyaku S, Kondo Y, Dohmae N, Fukuda H,
Yamaguchi-Shinozaki K, Shinozaki K. 2018. A small peptide modulates stomatal control
via abscisic acid in long-distance signalling. Nature **556**, 235–238.

Tegeder M, Masclaux-Daubresse C. 2018. Source and sink mechanisms of nitrogen
transport and use. The New Phytologist **217**, 35–53.

Tyree MT, Cochard H. 2003. Vessel contents of leaves after excision: a test of the
Scholander assumption. Journal of Experimental Botany **54**, 2133–2139.

Valero-Mora PM. 2010. **ggplot2:** Elegant Graphics for Data Analysis. Journal of Statistical
Software **35**.

Vialet-Chabrand SRM, Matthews JSA, McAusland L, Blatt MR, Griffiths H, Lawson
T. 2017. Temporal Dynamics of Stomatal Behavior: Modeling and Implications for
Photosynthesis and Water Use. Plant Physiology **174**, 603–613.

Vos I, Verhage A, Schuurink R, Watt L, Pieterse C, Van Wees S. 2013. Onset of
herbivore-induced resistance in systemic tissue primed for jasmonate-dependent defenses is
activated by abscisic acid. Frontiers in Plant Science **4**.

Vos IA, Verhage A, Watt LG, Vlaardingerbroek I, Schuurink RC, Pieterse CM, Wees
SCV. 2019. Abscisic acid is essential for rewiring of jasmonic acid-dependent defenses
during herbivory. , 747345.

Walker BJ, Busch FA, Driever SM, Kromdijk J, Lawson T. 2018. Survey of Tools for
Measuring In Vivo Photosynthesis. In: Covshoff S, ed. Methods in Molecular Biology.
Photosynthesis: Methods and Protocols. New York, NY: Springer, 3–24.

Wang M, Lee J, Choi B, Park Y, Sim H-J, Kim H, Hwang I. 2018. Physiological and
Molecular Processes Associated with Long Duration of ABA Treatment. Frontiers in Plant
Science **9**.

Xie J, Fernandes SB, Mayfield-Jones D, Erice G, Choi M, E Lipka A, Leakey ADB.
2021. Optical topometry and machine learning to rapidly phenotype stomatal patterning traits
for maize QTL mapping. Plant Physiology **187**, 1462–1480.

Xie Z, Jiang D, Dai T, Jing Q, Cao W. 2004. Effects of exogenous ABA and cytokinin on
leaf photosynthesis and grain protein accumulation in wheat ears cultured in vitro. Plant
Growth Regulation **44**, 25–32.

Yoon D-K, Ishiyama K, Suganami M, *et al.* 2020. Transgenic rice overproducing Rubisco
exhibits increased yields with improved nitrogen-use efficiency in an experimental paddy
field. Nature Food **1**, 134–139.

Zhu S-D, Liu H, Xu Q-Y, Cao K-F, Ye Q. 2016. Are leaves more vulnerable to cavitation
than branches? Functional Ecology **30**, 1740–1744.

Zhu X-G, Long SP, Ort DR. 2010. Improving Photosynthetic Efficiency for Greater Yield.
Annual Review of Plant Biology **61**, 235–261.

Figure Legends

Figure 1. Boxplots showing differences in leaf water potential between intact and excised

leaves for (a) tomato, (b) barley, and (c) maize. The p-values associated with the

corresponding one-sample t-test are inset in each panel.

Figure 2. Boxplots showing differences in leaf ABA concentration between intact and excised

leaves for (a) tomato, (b) barley, and (c) maize. The p-values associated with the

corresponding one-sample t-test are inset in each panel.

Figure 3. Boxplots showing differences in light-saturated gas exchange between excised and

intact leaves in the morning (AM) and afternoon (PM) for tomato (a-d), barley (e-h), and maize

(i-l). Net photosynthesis (A_N), stomatal conductance (g_s), the concentration of intracellular

CO_2 (c_i), and intrinsic water use efficiency (*iWUE*) are shown. The p-values associated with the

corresponding two-way analysis of variance (ANOVA) are inset in each panel.

Figure 4. Differences in photosynthetic capacities modelled via photosynthesis-CO_2 response

measurements between intact and excised leaves. (a-b) The response of photosynthesis (A_N)

to increasing intracellular CO2 concentration (c_i) in the (a) AM and (b) PM for tomato. The

solid line represents the mean fit of the FvCB model and the shaded area represents the

standard error of the mean. (c) The maximum rate of Rubisco carboxylation (V_{cmax}) for tomato

in the AM and PM. (d) The maximum rate of electron transport for RuBP regeneration (J_{max})

for tomato in the AM and PM. (e-f) The response of A_N to increasing c_i in the (e) AM and (f)

PM for barley. The solid line represents the mean fit of the FvCB model and the shaded area

represents the standard error of the mean. (g) V_{cmax} for barley in the AM and PM. (h) J_{max} for

barley in the AM and PM. (i-j) The response of A_N to increasing c_i in the (i) AM and (j) PM for

maize. The solid line represents the mean fit of a four-parameter non-rectangular hyperbola

and the shaded area represents the standard error of the mean. (k) The maximum rate of

carboxylation by PEPc (V_{pmax}) for maize in the AM and PM. (l) The maximum rate of

photosynthesis (V_{max}) for maize in the AM and PM.

Figure 5. The effect of leaf excision on hyperspectral reflectance in the morning (AM) and

afternoon (PM) for tomato, barley, and maize. (a) AM tomato. (b) PM tomato. (c) AM barley.

(d) PM barley. (e) AM maize. (f) PM maize. Each panel shows the -$\log_{10}$ transformed p-value

from a one-way analysis of variance (ANOVA) comparing reflectance of intact and excised

leaves at each incremental wavelength from 350-2500nm. The dashed red lines indicate a

Bonferroni significance threshold ($\alpha = 0.05$).

Figure 6. Boxplots showing variation in chlorophyll fluorescence-associated traits between

excised and intact leaves measured in a closed chlorophyll fluorescence system. (a) Maximum

efficiency of photosystem II (F_v/F_m) in tomato. (b) The rate of induction of non-photochemical

quenching (NPQ) in tomato. (c) The rate of relaxation of NPQ in tomato. (d) Maximum NPQ

in tomato. (e) F_v/F_m in barley. (f) The rate of induction of NPQ in barley. (g) The rate of

relaxation of NPQ in barley. (h) Maximum NPQ in barley. (i) F_v/F_m in maize. (j) The rate of

induction of NPQ in maize. (k) The rate of relaxation of NPQ in maize. (l) Maximum NPQ in

maize. The p-values associated with the corresponding one-sample t-test are inset in each

panel.

Figure 1. Boxplots showing differences in leaf water potential between intact and excised leaves for (a) tomato, (b) barley, and (c) maize. The p-values associated with the corresponding one-sample t-test are inset in each panel.

Figure 2. Boxplots showing differences in leaf ABA concentration between intact and excised leaves for (a) tomato, (b) barley, and (c) maize. The p-values associated with the corresponding one-sample t-test are inset in each panel.

Figure 3. Boxplots showing differences in light-saturated gas exchange between excised and intact leaves in the morning (AM) and afternoon (PM) for tomato (a-d), barley (e-h), and maize (i-l). Net photosynthesis (A_N), stomatal conductance (g_s), the concentration of intracellular CO_2 (c_i), and intrinsic water use efficiency (*iWUE*) are shown. The p-values associated with the corresponding two-way analysis of variance (ANOVA) are inset in each panel.

Figure 4. Differences in photosynthetic capacities modelled via photosynthesis-CO_2 response measurements between intact and excised leaves. (a-b) The response of photosynthesis (A_N) to increasing intracellular CO2 concentration (c_i) in the (a) AM and (b) PM for tomato. The solid line represents the mean fit of the FvCB model and the shaded area represents the standard error of the mean. (c) The maximum rate of Rubisco carboxylation (V_{cmax}) for tomato in the AM and PM. (d) The maximum rate of electron transport for RuBP regeneration (J_{max}) for tomato in the AM and PM. (e-f) The response of A_N to increasing c_i in the (e) AM and (f) PM for barley. The solid line represents the mean fit of the FvCB model and the shaded area represents the standard error of the mean. (g) V_{cmax} for barley in the AM and PM. (h) J_{max} for barley in the AM and PM. (i-j) The response of A_N to increasing c_i in the (i) AM and (j) PM for maize. The solid line represents the mean fit of a four-parameter non-rectangular hyperbola and the shaded area represents the standard error of the mean. (k) The maximum rate of

carboxylation by PEPc (V_{pmax}) for maize in the AM and PM. (l) The maximum rate of photosynthesis (V_{max}) for maize in the AM and PM.

Figure 5. The effect of leaf excision on hyperspectral reflectance in the morning (AM) and afternoon (PM) for tomato, barley, and maize. (a) AM tomato. (b) PM tomato. (c) AM barley. (d) PM barley. (e) AM maize. (f) PM maize. Each panel shows the -log$_{10}$ transformed p-value from a one-way analysis of variance (ANOVA) comparing reflectance of intact and excised leaves at each incremental wavelength from 350-2500nm. The dashed red lines indicate a Bonferroni significance threshold (α = 0.05).

Figure 6. Boxplots showing variation in chlorophyll fluorescence-associated traits between excised and intact leaves measured in a closed chlorophyll fluorescence system. (a) Maximum efficiency of photosystem II (F_v/F_m) in tomato. (b) The rate of induction of non-photochemical quenching (NPQ) in tomato. (c) The rate of relaxation of NPQ in tomato. (d) Maximum NPQ in tomato. (e) F_v/F_m in barley. (f) The rate of induction of NPQ in barley. (g) The rate of relaxation of NPQ in barley. (h) Maximum NPQ in barley. (i) F_v/F_m in maize. (j) The rate of induction of NPQ in maize. (k) The rate of relaxation of NPQ in maize. (l) Maximum NPQ in maize. The p-values associated with the corresponding one-sample t-test are inset in each panel.