Dorsal peduncular cortex activity modulates affective behavior and fear extinction in mice

Justin J. Botterill[1,2*], Abdessattar Khlaifia[1*], Ryan Appings[1], Jennifer Wilkin[1], Francesca Violi[1], Hanista Premachandran[1], Arely Cruz-Sanchez[1,3], Anna Elisabete Canella[1], Ashutosh Patel[1], S. Danyal Zaidi[1] and Maithe Arruda-Carvalho[1,3†]

1. Department of Psychology, University of Toronto Scarborough, Toronto, ON, M1C1A4, Canada
2. Department of Anatomy, Physiology, and Pharmacology, University of Saskatchewan, Saskatoon, SK, S7N 5E5, Canada.
3. Department of Cell and Systems Biology, University of Toronto, Toronto, ON, M5S3G5, Canada.

*JJB and AK contributed equally to this work.

[†]*Corresponding Author:*
Dr. Maithe Arruda-Carvalho
Department of Psychology
University of Toronto Scarborough
Toronto, Ontario, M1C1A4, Canada
Email: m.arrudacarvalho@utoronto.ca

Short/Running Title: Dorsal peduncular cortex in affective behaviors

Keywords: Prefrontal cortex, anxiety, depression, memory, stress, dorsal tenia tecta,

Abstract

The medial prefrontal cortex (mPFC) is critical to cognitive and emotional function and underlies many

neuropsychiatric disorders, including mood, fear and anxiety disorders. In rodents, disruption of mPFC

activity affects anxiety- and depression-like behavior, with specialized contributions from its subdivisions.

The rodent mPFC is divided into the dorsomedial prefrontal cortex (dmPFC), spanning the anterior

cingulate cortex (ACC) and dorsal prelimbic cortex (PL), and the ventromedial prefrontal cortex (vmPFC),

which includes the ventral PL, infralimbic cortex (IL), and in some studies the dorsal peduncular cortex

(DP) and dorsal tenia tecta (DTT). The DP/DTT have recently been implicated in the regulation of stress-

induced sympathetic responses via projections to the hypothalamus. While many studies implicate the

PL and IL in anxiety-, depression-like and fear behavior, the contribution of the DP/DTT to affective and

emotional behavior remains unknown. Here, we used chemogenetics and optogenetics to bidirectionally

modulate DP/DTT activity and examine its effects on affective behaviors, fear and stress responses in

C57BL/6J mice. Acute chemogenetic activation of DP/DTT significantly increased anxiety-like behavior in

the open field and elevated plus maze tests, as well as passive coping in the tail suspension test.

DP/DTT activation also led to an increase in serum corticosterone levels and facilitated auditory fear

extinction learning and retrieval. Activation of DP/DTT projections to the dorsomedial hypothalamus

(DMH) acutely decreased freezing at baseline and during extinction learning, but did not alter affective

behavior. These findings point to the DP/DTT as a new regulator of affective behavior and fear extinction

in mice.

Introduction

The medial prefrontal cortex (mPFC) is a key brain region involved in high-order functions such as attention [1,2], decision making [3,4], working memory [5,6], social behavior [7–10], mood [11,12] and anxiety [7,13,22–30,14–21]. The mPFC integrates inputs from cortical sensory and motor systems, as well as subcortical brain structures [31–36] toward top-down control of output structures such as the amygdala, thalamus, hypothalamus and bed nucleus of the stria terminalis (BNST) [31,34,36–38]. Despite a lack of unified consensus [39], many studies consider the rodent mPFC to be anatomically subdivided into the anterior cingulate cortex (ACC), prelimbic cortex (PL), and the infralimbic cortex (IL) [40], with some dividing the mPFC into two main regions with differences in their connectivity pattern and functional properties [40–43]: the dorsal mPFC (dmPFC), which includes the ACC and dorsal PL, and the ventral mPFC (vmPFC), which comprises the ventral PL, IL and the dorsal peduncular cortex (DP) and dorsal tenia tecta (DTT). Altered mPFC activity underlies many neuropsychiatric disorders, including mood, fear and anxiety disorders [38,44–52]. In rodents, disruption of mPFC activity through lesions, pharmacology, optogenetics or chemogenetics affects anxiety- [29,53–55] and depression-like [56–59] behavior, with evidence supporting specialized contributions from PL and IL subdivisions [21,23–27], as well as their subprojections [58,60]. Similarly, the PL and IL mPFC subdivisions exert distinct roles in auditory fear processing [61–66]. Although there is a considerable body of work implicating the mPFC as a key regulator of affective behavior and emotional learning, most studies focus on its main subdivisions, ACC, PL and IL, leaving subregions such as the DP/DTT understudied.

The DP/DTT is located at the ventral limit of the vmPFC [67], and plays a crucial role in driving thermoregulatory and cardiovascular sympathetic responses during stress [68–70]. The DP/DTT receives inputs from stress and emotion-related brain regions like the thalamus, amygdala, insular cortex and piriform cortex [68–70], and sends glutamatergic excitatory input to the dorsomedial hypothalamus (DMH), a central hub driving sympathetic responses to various stimuli, including psychosocial stress [68,70]. Interestingly, while dmPFC activation abrogates stress-related autonomic and neuroendocrine responses [71–74], DP/DTT activation mimics sympathetic responses to psychosocial stress [68], suggesting a distinct role for the DP/DTT in driving sympathetic stress responses. Given the high

prevalence of sympathetic system impairments in anxiety disorders [75–78] and the well-established role of the mPFC in modulating anxiety-, depression-like and cognitive behavior in rodents, we investigated the contribution of the DP/DTT to affective behavior and emotional learning in mice. Our results show that chemogenetic activation of DP/DTT with Designer Receptors Exclusively Activated by Designer Drugs (DREADDs) in C57BL/6J mice increases anxiety-like behaviors in the open field (OFT) and elevated plus maze (EPM) tests, passive coping in the tail suspension test (TST) and facilitates fear extinction learning and extinction retrieval. DP/DTT activation was accompanied by an increase in serum corticosterone levels. Activation of DP/DTT projections to the dorsomedial hypothalamus (DMH) acutely suppressed freezing. These findings uncover a novel role for the DP/DTT in promoting anxiety-like and fear extinction behaviors in mice.

Results

To explore the contribution of the DP/DTT to anxiety-, depression-like and fear behavior in mice, we first bidirectionally modulated DP/DTT activity across a series of behavioral tasks with excitatory and inhibitory DREADDs. AAV vectors encoding excitatory (hM3D), inhibitory (hM4D) DREADDs or the control fluorophore mCherry were injected into the DP/DTT of C57BL/6J mice. Viral expression was strongly localized to the DP/DTT across each treatment condition (**Supplementary Figure S1**).

Chemogenetic activation of DP/DTT increases anxiety-like behavior

We first examined the impact of manipulating DP/DTT activity on anxiety-like behavior in freely behaving mice in the OFT (**Figure 1A-F; Supplementary Figure S2A-B**). Mice expressing hM3D, hM4D or mCherry constructs in the DP/DTT region were injected with C21 one hour prior to OF testing (**Figure 1A-B**). Chemogenetic activation of DP/DTT neurons (hM3D mice) decreased the time spent in the center zone of the OF arena compared to mCherry and hM4D mice (**Figure 1C**; One-way ANOVA, $F_{2,38} = 7.795$, $p=0.002$, Tukey's post hoc tests, hM3D versus mCherry $p=0.002$; hM3D versus hM4D $p=0.011$; **Supplementary Figure S2B**). In contrast, hM3D mice spent significantly more time in the outer zone of the open field apparatus compared to both mCherry control and hM4D mice (**Figure 1D**, one way ANOVA, $F_{2,38}=7.283$, $p=0.002$; Tukey's post hoc tests, hM3D versus mCherry, $p=0.007$ and hM3D

versus hM4D, $p=0.006$). Furthermore, the total distance travelled was similar across groups (**Figure 1E**,

one way ANOVA, $F_{2,38}=2.292$, $p=0.115$), suggesting that the effect of C21-mediated activation of DP/DTT

neurons on anxiety-like behavior was not driven by changes in locomotor activity levels. These data

show that chemogenetic activation of the DP/DTT induces anxiety-like behavior in the OFT (**Figure 1F**).

Next, we examined the effects of manipulating DP/DTT activity on anxiety-like behavior using the EPM

test (**Figure 1G-L; Supplementary Figure S2C-F**). Consistent with the anxiogenic effect of DP/DTT

activation observed in the OFT, hM3D mice spent less time in the open arms of the EPM apparatus

compared to mCherry control and hM4D mice (**Figure 1I**; one-way ANOVA, $F_{2,38}=5.87$, $p=0.006$, Tukey's

post hoc tests, hM3D versus mCherry, $p=0.010$, hM3D versus hM4D, $p=0.025$; **Supplementary Figure**

S2D). Additionally, hM3D mice displayed fewer entries to the open arms of the EPM compared to hM4D

mice (**Figure 1J**, one-way ANOVA, $F_{2,38}=4.86$, $p=0.013$, Tukey's post hoc test, hM3D versus hM4D,

$p<0.011$). Moreover, hM3D mice spent significantly more time in the closed arms of the EPM compared

to mCherry control and hM4D mice (**Supplementary Figure S2E**). We saw no group differences in the

number of entries to the closed arm (**Supplementary Figure S2F**), middle platform ($F_{2,38}=2.130$,

$p=0.133$; data not shown), or in the time spent in the middle platform ($F_{2,38}=1.749$, $p=0.187$; data not

shown). All three experimental groups showed equivalent total distance travelled in the EPM apparatus

(**Figure 1K**). We found a significant effect of sex on the distance travelled in the EPM, with hM4D males

travelling significantly less distance compared to hM4D females ($F_{1,35}=8.23$, $p=0.007$; Sidak's multiple

comparisons test $p>0.109$; no main effect of treatment or interaction. For a discussion of sex differences,

see supplemental materials). Importantly, chemogenetic activation of the IL did not affect performance in

the OFT or in the EPM (**Figure 1M-R**). Taken together, our data suggest that acute activation of DP/DTT,

but not IL neurons results in increased anxiety-like responses in the OFT and EPM.

Chemogenetic activation of DP/DTT increases passive coping behavior

To examine whether chemogenetic modulation of DP/DTT activity affects passive coping behavior, we

next subjected mice to the TST (**Figure 2A-B**). The hM3D mice spent significantly more time immobile

compared to mCherry and hM4D groups following C21 injection (**Figure 2C-D**, one-way ANOVA,

$F_{2,38}$=5.07, p=0.11; Tukey's post hoc tests, hM3D vs mCherry, p=0.041, hM3D vs hM4D, p=0.017;

Supplementary Figure S2G-H), consistent with increased passive coping in the hM3D group. All three

groups showed equivalent length of the longest bout of immobility, and number of transitions from

mobility to immobility states (**Supplementary Figure S2I-J**). Chemogenetic activation of IL did not affect

immobility in the TST (**Figure 2E-F**).

To expand our findings of increased passive coping behavior in the TST following DP/DTT activation, we

conducted the FST (**Figure 2G-H**). C21 treatment did not affect time spent immobile (**Figure 2I-J**),

longest bout of immobility or the number of transitions between mobility and immobility states

(**Supplementary Figure 2K-N**) in any of the experimental groups in the FST. When we analyzed sex

differences, we found that hM3D females spent significantly less time immobile than hM3D males and

hM4D females (Two-way ANOVA, $F_{1,35}$=6.561, p=0.015; Sidak's multiple comparisons test, hM3D

females versus hM3D males p=0.0093; hM3D females versus hM4D females p=0.0482). Taken together,

our results demonstrate that activation of DP/DTT neurons promotes passive coping behavior in the TST,

but did not alter behavior in the FST.

Chemogenetic activation of DP/DTT activity facilitates auditory fear extinction

We next sought to explore whether manipulating DP/DTT activity might affect fear memory. To test this,

we first inhibited or activated the DP/DTT one hour prior to training mice to associate an auditory tone

with an electric foot shock (**Figure 3A**). During fear acquisition, chemogenetic manipulation of DP/DTT

activity did not affect baseline freezing (one-way ANOVA, $F_{2,38}$=2.367, p=0.107; data not shown), or the

mean percent freezing across all tones (**Figure 3B**). However, we found that hM4D mice displayed a

right shift in their fear learning curve, with significantly higher freezing during tone 4 and 6 of the training

protocol compared to hM3D mice (**Figure 3C**, two-way rmANOVA, significant effect of tone $F_{5,190}$=100.6,

p<0.001; Tukey's post hoc tests, minute 4 hM3D versus hM4d, p=0.012, and minute 6 hM3D versus

hM4D, p=0.012) suggesting a slight facilitation of auditory fear acquisition upon DP/DTT inhibition. We

found a significant main effect of sex on average freezing to tone (Two-way ANOVA, $F_{1,35}$=7.786,

p=0.009, see supplemental materials).

Auditory fear memory retrieval was evaluated 24h later in a different context (**Figure 3A**). All three experimental groups showed similar baseline freezing (one-way ANOVA, $F_{2,38}=0.153$, p=0.858; data not shown), and equivalent freezing to the tones (**Figure 3D-E**). Overall, these data suggest that manipulating DP/DTT activity has no major impact on auditory fear acquisition, except for a marginal facilitation of fear acquisition following DP/DTT inhibition.

To test whether DP/DTT manipulation might affect fear extinction and retrieval, we infused AAV vectors encoding either hM3D, hM4D or mCherry into the DP/DTT of a separate cohort of mice, and trained them in auditory fear conditioning as previously (**Figure 3F-H**). The next day, we injected C21 one hour prior to extinction training. During extinction training, hM3D animals froze less than hM4D and mCherry animals (**Figure 3I**: One-way ANOVA $F_{2,31}=12.91$, p<0.0001, mCherry versus hM3D $p=0.0003$, hM3D versus hM4D $p=0.0002$; 3J: Two-way rmANOVA main effect of treatment $F_{2,31}=12.91$, $p<0.0001$ and tone $F_{11,341}=7.869$, $p<0.0001$ only; mCherry versus hM3D $p=0.0003$, hM3D versus hM4D $p=0.0002$), suggesting that DP/DTT activation impairs auditory fear retrieval and facilitates within session extinction. The next day, animals underwent extinction retrieval in the absence of C21. At extinction retrieval, hM3D animals showed less freezing than both hM4D and mCherry animals (**Figure 3K**: One-way ANOVA $F_{2,31}=5.853$, p=0.0070, mCherry versus hM3D p=0.0078, hM3D versus hM4D p=0.0245; 3L: Two-way rmANOVA significant main effect of treatment $F_{2,31}=4.526$, $p=0.0189$ and tone $F_{4,124}=3.021$, $p=0.0204$ only; mCherry versus hM3D $p=0.0255$, hM3D versus hM4D $p=0.0385$). Overall, these data show that DP/DTT activation leads to reduced fear recall and a facilitation of fear extinction learning and extinction retrieval.

Chemogenetic activation of DP/DTT increases serum corticosterone levels

Given the important role of the mPFC in regulating the hypothalamic-pituitary-adrenal (HPA) axis response to emotional stress [72,79–81], and the regulation of stress-induced sympathetic responses by the DP/DTT [68], we next tested the effects of DP/DTT chemogenetic modulation on serum levels of the stress hormone corticosterone. We injected C21 into a subset of mCherry, hM3D, and hM4D mice that remained in their home cage, and subsequently extracted blood to evaluate serum corticosterone

(**Figure 3M**). Notably, these mice had not undergone any previous behavioral testing. Interestingly,

hM3D mice displayed higher serum corticosterone levels compared to mCherry control mice (**Figure 3N**,

one-way ANOVA, $F_{2,12}$=6.325, p=0.013, Tukey's post hoc tests, hM3D versus mCherry, p=0.011).

Chemogenetic inhibition of DP/DTT activity did not affect baseline serum corticosterone levels (**Figure**

3N). These data indicate that activation of DP/DTT neurons is sufficient to trigger a neuroendocrine

stress response by increasing serum corticosterone levels. We validated our chemogenetic

manipulations using cFos immunohistochemistry and slice electrophysiology (**Supplementary Figure**

S3).

186
Neurochemical profile and mapping of DP/DTT downstream projections

To explore potential downstream effectors mediating the effects of DP/DTT activation on affective, fear

behavior and neuroendocrine responses, we first analyzed brain-wide patterns of DP/DTT axonal

innervation with the help of the mCherry tag in AAV-injected mice (**Figure 4A-C**). Projections from the

DP/DTT were observed in the lateral septum (medial and lateral nuclei), indusium griseum (IG), BNST,

dorsal endopiriform nucleus (DEn), thalamus (nucleus reuniens, RE; paraventricular nuclei, PVT;

submedius nucleus, Sub), hypothalamus (DMH; lateral hypothalamus, LH; paraventricular nucleus, PVN;

posterior hypothalamus, PH; and retromammillary nuclei, RM) and brainstem (periaqueductal gray, PAG)

(**Figure 4A-C**). Posterior brainstem sections containing the raphe nuclei also showed DP/DTT mCherry+

projections (data not shown).

To determine which DP/DTT neuronal subpopulations and downstream targets are actively recruited

during behavior, we next used viral tracing in Fos[2A-iCreER] (TRAP2) transgenic mice [82,83] to tag DP/DTT

neurons active during the OFT. We infused an AAV expressing Cre-dependent eYFP (AAV-DIO-eYFP)

into the DP/DTT of TRAP2 mice and, following recovery, injected animals with tamoxifen immediately

after OFT testing to tag the soma and axons of activated DP/DTT neurons through YFP expression

(**Figure 4D-E**). We found that 67.9% of DP/DTT neurons active during OFT testing co-labeled with a

glutamatergic marker, 13.9% with a GABAergic marker, and 2.5% of labeled cells were positive for both

markers (**Figure 4F**). Overall, this points to OFT activating a heterogeneous but predominantly

glutamatergic DP/DTT cell population.

206
Optogenetic activation of the DP/DTT-DMH pathway acutely suppresses freezing but does not

alter affective behavior

We next wanted to test whether one of the main downstream pathways revealed by our previous

analysis, the DP/DTT-DMH pathway [68], might underlie our effects on affective and fear extinction

behavior. We optogenetically targeted the DP/DTT-DMH pathway by expressing the excitatory opsin

channelrhodopsin (ChR2) or control GFP into the DP/DTT and implanting an optic fiber over the DMH,

and tested the animals in the OFT, EPM and auditory fear (**Figure 5A, Supplementary Figure S4A-B**).

Optical activation of the DP/DTT-DMH pathway did not alter behavior in the OFT (**Figure 5B-D**) or EPM

tasks (**Figure 5E-G**). In contrast, activation of the DP/DTT-DMH during tone exposure at extinction

learning significantly suppressed freezing responses (**Figure 5H-I**: t_{10}=7.119, p<0.0001; **Figure 5J**: Two-

way rmANOVA significant main effect of stimulation $F_{1,10}$=50.68, p<0.0001; tone $F_{3.961,39.61}$=3.026,

p=0.0291, and interaction $F_{11,110}$=1.883, p=0.0491; Tone 1 p=0.0042, Tone 2 p=0.0461) without affecting

extinction retrieval (**Figure 5K-L**). To test whether this suppression of freezing was specific to the tone-

shock association, we fear conditioned a separate cohort of mice and activated the DP/DTT-DMH

pathway in the alternate context in the absence of tones (**Figure 5M**). DP/DTT-DMH activation led to a

reduction in baseline freezing behavior (**Figure 5N**; Two-way rmANOVA significant main effect of

stimulation $F_{1,11}$=15.88, p=0.0021 and interaction $F_{1,11}$=9.375, p=0.0108 only; ChR2 p=0.0011) and an

increase in rearing (**Figure 5O**; Two-way rmANOVA significant main effect of stimulation $F_{1,11}$=16.30,

p=0.002 and interaction $F_{1,11}$=11.28, p=0.006 only; ChR2 p=0.0008). These data suggest that activation

of the DP/DTT-DMH pathway acutely suppresses freezing at baseline and during extinction training

without affecting extinction retrieval.

Discussion

Here we examined the effects of modulating the activity of DP/DTT neurons to anxiety-, depression-like

and fear behavior in stress-naïve mice. We found that chemogenetic activation of the DP/DTT reduced

the time spent in the center zone of the OFT and in the open arms of the EPM. DP/DTT activation also

increased passive coping in the TST, but did not affect behavior in the FST. DP/DTT activation facilitated

fear extinction training and extinction retrieval, whereas DP/DTT-DMH activation acutely suppressed

freezing. Interestingly, DP/DTT activation led to an increase in serum corticosterone levels, suggesting

that the DP/DTT also modulates HPA axis activity. DP/DTT projection mapping showed strong

innervation of lateral septum, thalamus, hypothalamus, and brainstem, consistent with tracing studies

[33,67,84,85].

Our data showed that activation of the DP/DTT increased anxiety-like behavior in the OFT and EPM. A

considerable body of work implicates the rodent mPFC in the regulation of anxiety-like behavior

[7,13,22–29,86,14–21], with contradictory findings supporting anxiogenic [18–23,53–55] or anxiolytic

[24–27] roles. Manipulations selective to the vmPFC/IL also see conflicting effects on EPM and open

field performance [21,24,26,60]. Recently, Chen and colleagues found that optogenetic activation of IL

soma drives anxiety-like behavior in the EPM and open field [60], similar to our data with DP/DTT

activation. In contrast, they found that optical stimulation of the DP increased entries to the open arms in

the EPM and time spent in the center of the open field [60], contrary to our findings. Our data support an

anxiogenic role for the DP/DTT, which is in line with findings from Kataoka and colleagues showing that

inhibition of the DP/DTT pathway abolishes stress-induced avoidance behavior [68], and our finding of

increased serum corticosterone following DP/DTT activation. Importantly, the viral spread over the DP in

the study by Chen and colleagues is more dorsal (spreading to IL, but not DTT) compared to ours, and

targets glutamatergic neurons under the αCaMKII promoter [60], in contrast to the pan-neuronal

synapsin and EF1a promoters used here and by Kataoka and colleagues [68]. These discrepancies favor

additional cell-type and subregion specialization across the dorso-ventral axis of the vmPFC in regulating

anxiety-like behavior and neuroendocrine responses.

DP/DTT activation increased immobility time in the TST, but not in the FST. Given the differential

implications of animal weight on water-based (FST) and suspension (TST) task demands, it is possible

that the relatively high weight of our animals at the time of testing (>30g for males) contributed to the

differences in performance between these tasks. The mPFC has a well-established role in modulating

depression-like behavior [87] in both stress-naïve [56–59,88,89] and stress conditions [88,90–95],

including IL-specific manipulations [57,59]. Interestingly, Warden and colleagues identified mPFC

neuronal population coding for FST active and inactive behavioral states, which reflected specialization

within mPFC downstream projections to the dorsal raphe nucleus and lateral habenula [96]. Accordingly,

somatic activation of the mPFC as a whole did not affect behavior in the FST [96], suggesting that further

probing of specific DP/DTT projections could lead to modulation of FST behavior despite our negative

findings.

Brain-wide mapping of DP/DTT projections showed strong downstream connectivity with the

hypothalamus, thalamus, lateral septum and brainstem. Importantly, many of the upstream brain regions

projecting to the DP [68] also regulate anxiety-like behavior in rodents, including the mediodorsal (MD)

[97–100] and paraventricular (PVT) [101–105] thalamus, insular cortex (IC) [105–110], nucleus reuniens

[111] and amygdala [14,112–116]. While we found that the majority of OFT activity-tagged DP/DTT

neurons is glutamatergic, the precise patterns of afferent and efferent innervation of DP/DTT neuronal

subpopulations and their modulation of anxiety-like behavior remains unknown.

Our data showed minimal effects of DP/DTT soma manipulation on auditory fear acquisition, but a

marked reduction in freezing when the DP/DTT was activated during fear extinction, which persisted

during extinction retrieval in the absence of DP/DTT manipulation. This points to the DP/DTT as sharing

a similar function to IL [61,129–132, but see 133] in the modulation of fear inhibition. In contrast, a recent

study elegantly showed that activation and inhibition of DP neurons tagged during auditory fear

conditioning increases and reduces freezing at retrieval, respectively [122]. The opposing direction of

these findings in relation to ours could be driven by a differential contribution of the whole DP/DTT (our

manipulations) compared to the selective activation of DP neurons responding to fear acquisition [122],

whose connectivity is presently unknown. This is also underscored by a lack of activity signatures by

these fear-trapped DP neurons during extinction [122], whereas we saw an effect of DP/DTT soma

manipulation on extinction learning and retrieval.

We found that DP/DTT-DMH stimulation had no effect on affective behavior in the OFT or EPM, but

acutely suppressed freezing. Notably, in contrast to our DP/DTT soma manipulation, DP/DTT-DMH-

mediated suppression of freezing at extinction learning did not affect extinction retrieval, suggesting that

the DP/DTT effects on extinction retrieval might be mediated by a distinct downstream target.

Furthermore, the increased rearing seen in the DP/DTT-DMH ChR2 animals at baseline post-

conditioning might also signal stress-related behavior [123], consistent with this pathway's role in

mediating sympathetic responses and social avoidance following social defeat stress in rats [68].

Alternatively, DP/DTT-DMH activation after foot shock might be triggering flight-like behavior as recently

reported for the DP-central amygdala (CeA) pathway [124]. In this comprehensive study, inhibition of the

DP-CeA pathway increased freezing during fear conditioning [124], similar to our trends with DP soma

inhibition. It is also important to note that the small magnitude of our fear acquisition findings might have

been affected by previous exposure to stressful manipulations such as the FST. Interestingly, stimulation

of the DP-CeA pathway in a safe context after conditioning reduces freezing [124], similar to our results

with DP/DTT and DP/DTT-DMH stimulation. Importantly, the authors reported an anxiolytic effect of

inhibition of the DP-CeA pathway in the OFT and EPM, but no effect of activation [124], suggesting that

our effects of DP soma manipulation on affective behavior might be independent of the CeA. This is in

line with the absence of a strong projection to CeA with our viral strategy. Finally, Chen and colleagues

demonstrated that IL downstream projections to the lateral septum and central amygdala (CeA) promote

and inhibit anxiety-like behavior, respectively [60], suggesting a high degree of specialization within the

vmPFC for the modulation of anxiety-like behavior.

We found that chemogenetic activation of the DP/DTT was sufficient to trigger an increase in serum

corticosterone levels. This is consistent with the finding that optogenetic stimulation of the DP/DTT-DMH

pathway in stress-naïve animals drives sympathetic responses mimicking those triggered by social

defeat stress [68], suggesting that the DP/DTT might transduce stress responses from upstream

corticolimbic regions toward downstream autonomic, behavioral/motor [68] and endocrine responses.

Importantly, while most of the literature supports an inhibitory effect of the mPFC over HPA secretory

responses following stress [71,125–128] (but see [129]), Radley and colleagues saw a dichotomy

between the dorsal and ventral mPFC in modulating stress-induced HPA activation [72]. Specifically,

while dmPFC lesions increased corticosterone levels following acute restraint stress, vmPFC lesions

decreased corticosterone levels [72]. This suggests that mPFC inhibition of the HPA axis response is

mostly driven by the dorsal mPFC, with the ventral portion of the mPFC (possibly spanning the DP/DTT)

exerting the opposite effect, consistent with our findings. Importantly, Kataoka and colleagues saw no

evidence of DP/DTT inhibition affecting basal autonomic homeostasis [68], consistent with our lack of

effects of DP/DTT inhibition in stress-naïve animals. Similarly, inhibition of the ventral portion of the

mPFC did not affect baseline corticosterone levels [72].

While we observed that the DP/DTT can modulate neuroendocrine responses, the downstream targets

mediating this response are unclear. The hypothalamus is necessary for the effects of frontal cortex

stimulation on HPA activity [130], and mediates the autonomic effects of DP/DTT activation [68]. It is

hypothesized that (dorsal) mPFC inhibition of HPA activity occurs through stimulation of GABAergic

projections from DMH to the paraventricular nucleus of the hypothalamus (PVN) [131]. Consistent with

what has been described for the vmPFC [126], we saw DP/DTT projections to hypothalamic nuclei such

as the PVN, DMH and lateral hypothalamus (LH). While Radley and colleagues proposed that the

stimulating effects of ventral mPFC on HPA activity might be mediated through connections with BNST

[72], studies suggest an inhibitory effect of the BNST over HPA activity following acute stress [132], and

fail to see recruitment of mPFC-BNST projections following stress [128,133]. Altough we see DP/DTT

projections to the BNST, their contribution to the regulation of DP/DTT effects on HPA secretory activity

is unknown.

Overall, our data expand an emerging literature examining the contributions of the DP/DTT to the

modulation of psychosocial stress and sympathetic responses by revealing the DP/DTT as a regulator of

anxiety-like, passive coping and fear extinction behaviors. As its topography allows for the integration of

higher order information toward autonomic and sympathetic effectors, whose dysfunction is often a

feature of anxiety disorders, the DP/DTT may be uniquely placed to mediate the effects of stress on

affective function. Our findings support further specialization across the dorsoventral axis of the mPFC in

the processing of anxiety- and depression-like behavior and stress responsivity, in line with rising

evidence of pathway-specific modulation of anxiety-like behavior among mPFC subregions.

Acknowledgements:

We thank Hathairat Chanphao, Tejnarine Persaud, Bebhinn Treanor, and Christina Guzzo for their

generous and fundamental assistance with the corticosterone ELISA experiment and use of the Guzzo

lab plate reader. We would also like to thank Unza Mumtaz and Mehreen Inayat for their help with animal

colony management. Some figure diagrams were created with the assistance of BioRender.com.

Author Contributions:

Designed Research: JJB, AK, MAC.

Performed Research: JJB, AK, RA, JW, FV, HP, ACS, AEC, AP, SDZ, MAC

Analyzed/Interpreted Data: JJB, AK, JW, FV, AEC, AP, MAC.

Wrote the paper: JJB, AK, MAC.

All authors reviewed and approved the manuscript.

Funding:

This work was supported by grants from CIHR (PJT 399790), Human Frontier Science Program

Organization (CDA00009/2018 and RGY0072/2019), the SickKids Foundation and Canadian Institutes of

Health Research (CIHR) – Institute of Human Development, Child and Youth Health (NI19-1132R), and

Natural Sciences and Engineering Research Council of Canada (RGPIN-2017-06344) to MAC.

Competing Interests:

The authors have nothing to disclose.

Figure 1. Chemogenetic activation of the DP/DTT increases anxiety-like behavior in the open field and elevated plus maze tests.

A-B. Experimental timeline for the OFT. Mice previously injected with hM3D, hM4D or control mCherry constructs in the DP/DTT were injected with the DREADD agonist C21 one hour prior to being placed in an open field. The time spent in the center zone (red) and the outer zone (white) was measured. **C.** The time spent in the center zone of the open field was significantly reduced in hM3D mice compared to mCherry and hM4D mice (no difference between mCherry and hM4D, p=0.965). **D.** The amount of time spent in the outer zone of the open field arena was significantly greater in the hM3D mice compared to mCherry and hM4D mice (mCherry versus hM4D, p=0.938). **E.** The total distance travelled during the test did not differ between groups (one-way ANOVA, $F_{2,38}$=2.292, p=0.115). **F.** Representative track and heat maps. **G-H.** Experimental timeline for the EPM test. Mice previously injected with hM3D, hM4D or control mCherry constructs in the DP/DTT were injected with the DREADD agonist C21 one hour prior to being placed in an elevated plus maze. The time spent in the open arms vs closed arms was measured. **I.** The time spent in the open arms of the elevated plus maze was significantly reduced in hM3D mice compared to mCherry and hM4D mice (no difference between mCherry and hM4D mice, p=0.993). **J.** The number of open arm entries was significantly reduced in hM3D mice compared to hM4D mice (no difference between hM3D and mCherry, p=0.136; or hM4D and mCherry, p=0.430). **K.** The total distance travelled during the test did not differ between groups (one-way ANOVA, $F_{2,38}$=0.022, p=0.98). While we found that hM4D males travelled less than hM4D females (see results), separation of the data by sex did not reveal group differences (One-way ANOVA, females $F_{2,16}$= 0.3571, p= 0.7051; males $F_{2,19}$= 0.093, p= 0.9117). **L.** Representative track and heat maps. **M**. Schematic of IL viral targeting. Mice previously injected with hM3D or control mCherry constructs into the IL were injected with the DREADD agonist C21 one hour prior to undergoing the OFT (**N-O**) or EPM test (**P-R**). **N**. IL-hM3D mice spent the same

amount of time in the center zone of the OFT as IL-mCherry mice (t_{14}=0.2502, p=0.8061). **O**. OFT
distance travelled did not differ between groups (t_{14}=1.689, p=0.1134). **P**. IL-hM3D mice spent the same
amount of time in the open arms of the EPM as IL-mCherry mice (t_{14}=1.916, p=0.076). **Q**. IL-hM3D and
IL-mCherry mice displayed the same number of open arm entries in the EPM (t_{14}=0.8721, p=0.3979). **R**.
EPM distance travelled did not differ between groups (t_{14}=1.999, p=0.0654). *p<0.05, **p<0.01. Male
(square) and female (circle) individual datapoints are displayed for transparency (see text for details).

Figure 2. Chemogenetic activation of the DP/DTT increases immobility in the tail suspension test, but have no effect on forced swim behaviors.

A-B. TST experimental timeline. Mice previously injected with hM3D, hM4D or control mCherry constructs in the DP/DTT were injected with the DREADD agonist C21 one hour prior to undergoing the TST. **C.** Representative raster plots for each mouse indicating periods of immobility (black) and mobility (white) over the tail suspension test duration. **D.** The average percent of time spent immobile during the TST was significantly greater in hM3D mice compared to mCherry and hM4D mice (but not different between mCherry and hM4D, p=0.838). **E-F.** Effects of IL DREADD soma manipulation on the TST. Mice previously injected with hM3D or control mCherry constructs in the IL were injected with the DREADD agonist C21 one hour prior to undergoing the TST. **E.** Representative raster plots for each mouse indicating periods of immobility (black) and mobility (white) over the TST duration. **F.** IL-hM3D mice spent the same amount of time immobile in the TST as IL-mCherry mice (t_{14}=1.102, p=0.2892). **G-H.** FST experimental timeline. Mice previously injected with hM3D, hM4D or control mCherry constructs in the

DP/DTT were injected with the DREADD agonist C21 one hour prior to undergoing the forced swim test.

I. Representative raster plots for each mouse indicating periods of immobility (black) and mobility (white)

over the FST duration. **J.** There was no effect of treatment on percent immobility in the forced swim test

(one-way ANOVA, $F_{2,38}$=1.722, p=0.192). While we found that hM3D females had a lower percent time

spent immobile compared to males (see results), separation of the data by sex did not reveal group

differences (One-way ANOVA, females $F_{2,16}$= 2.816, p= 0.0896; males $F_{2,19}$=1.042, p= 0.3722). *p<0.05.

Male (square) and female (circle) individual datapoints are displayed for transparency (see text for

details).

Figure 3. Chemogenetic inhibition of the DP/DTT has modest effect on auditory fear acquisition, but activation of the DP/DTT facilitates within-session extinction, extinction retrieval and increases serum corticosterone levels.

A-E. DP/DTT manipulation during auditory fear acquisition. **A.** Experimental timeline. Mice previously injected with hM3D, hM4D or control mCherry constructs in the DP/DTT were injected with the DREADD agonist C21 one hour prior to undergoing auditory fear conditioning. During acquisition, mice were placed in a sound-attenuated fear conditioning chamber and received 6 tone-shock pairings (0.5mA). Twenty-four hours later, auditory fear memory was evaluated in a different context in the absence of shocks. **B.** During fear acquisition, DP/DTT manipulations did not affect average tone freezing (one-way ANOVA, $F_{2,38}=2.828$, $p=0.072$). **C.** hM4D mice displayed significantly more freezing than hM3D mice at

tone presentations 4 and 6 during acquisition. Two-way rmANOVA found no main effect of treatment
$F_{2,38}$=2.828, p=0.072, but a significant main effect of tone ($F_{5,190}$=100.6, p<0.001, and no treatment by
tone interaction: $F_{10,190}$=1.633, p=0.998). **D-E.** During fear retrieval, DP/DTT manipulations did not affect
freezing during average tone (**D**; one-way ANOVA, $F_{2,38}$ =0.507, p=0.607). Individual tone analysis (**E**)
showed no main effect of treatment (Two-way rmANOVA, $F_{2,38}$=0.507, p=0.607), but a significant main
effect of tone ($F_{5,190}$=4.00, p=0.002, no treatment by tone interaction: $F_{10,190}$=1.213, p=0.285), likely driven
by within-session extinction. ✱ indicates a significant difference between hM3D and hM4D groups
(p<0.05). **F-L.** DP/DTT manipulation during fear extinction. **F.** Experimental timeline. Mice previously
injected with hM3D, hM4D or control mCherry constructs in the DP/DTT (no prior behavior) underwent
fear acquisition (as previously; 6 tone-shock pairings). The following day, mice were injected with the
DREADD agonist C21 one hour prior to undergoing extinction training. During extinction training, mice
were placed in an alternate context (context B) and received 12 tone presentations. The following day,
fear memory was evaluated during extinction retrieval, in the absence of C21. During extinction retrieval,
mice were placed in context B and received 5 tone presentations. **G-H.** During auditory fear acquisition,
mice from all groups displayed similar average tone (**G**; one-way ANOVA $F_{2,31}$ = 0.5886, p=0.5612) but
decreased freezing in tone 6 in the hM3D group (**H**; Two-way rmANOVA significant effect of tone
$F_{5,155}$=74.69, p<0.0001 and interaction $F_{10,155}$=1.984, p=0.0385; but no main effect of treatment
$F_{2,31}$=0.8356, p=0.4431; tone 6 mCherry vs hM3D p=0.0023 and hM3D vs hM4D p=0.0109) freezing. **I-J.**
During extinction training, 1 hour after C21 treatment, hM3D mice showed decreased average tone
freezing (**I**; Two-way rmANOVA main effect of treatment and tone but no interaction $F_{22,341}$=1.072,
p=0.3758) and tone-by-tone (**J**) freezing compared to hM4D and mCherry control mice. **K-L.** During
extinction retrieval, hM3D mice froze less compared to hM4D and mCherry control mice in the average
tone (**K**) and tone-by-tone (**L**; Two-way rmANOVA significant main effect of treatment and tone but no
interaction $F_{8,124}$=0.6037, p=0.7733) analyses. Minimal sex differences were found in this dataset (see
Supplementary Materials). **M.** Experimental timeline for serum corticosterone assay. Mice previously
injected with hM3D, hM4D or control mCherry constructs in the DP/DTT were injected with the DREADD
agonist C21 and euthanized 90 minutes later. Blood was extracted at the time of perfusion for
corticosterone measurements. **N.** Serum corticosterone was significantly higher in hM3D mice compared
to mCherry mice (but no change between mCherry and hM4D, p=0.440, hM4D vs hM3D, p=0.103). ✱
indicates a significant group difference between mCherry and hM3D conditions; ✱ indicates a significant
group difference between hM3D and hM4D conditions; ✱ indicates a significant difference between
hM3D and all other groups. *p<0.05. Male (square) and female (circle) individual datapoints are
displayed for transparency.

Figure 4. Mapping of DP/DTT downstream projections and characterization of activity-tagged DP/DTT neurons following the OFT.

A-C. Representative projections of the DP/DTT in anterior, intermediate, and posterior sections. **A.** In anterior sections, we observed notable viral expression in the indusium griseum (IG), septum, bed nucleus of the stria terminalis (BNST), dorsal endopiriform nucleus (DEn), nucleus reuniens (RE) and

paraventricular thalamus (PVT). **B.** In intermediate sections, we observed continued viral expression in
the PVT and RE. Viral expression was also observed in the paraventricular nucleus (PVN) submedius
thalamic nucleus (Sub), dorsomedial hypothalamus (DMH), lateral hypothalamus (LH), and posterior
hypothalamus (PH). **C.** In the most posterior sections we examined, viral expression was detected in the
periaqueductal gray (PAG), retromammillary nuclei (RM), PVT, LH and PH. In posterior brainstem
sections, we also observed DP/DTT projections to the raphe nuclei (data not shown). **D-F**.
Characterization of neurochemical profile and downstream projections patterns from activity-tagged
DP/DTT neurons following the OFT. **D**. Experimental timeline. TRAP2 mice were infused with AAV
expressing a DIO-YFP construct in the DP/DTT. Seven days later, animals underwent the OFT and were
injected with 4HT tamoxifen immediately afterwards. Ten days later, animals were perfused and their
brains processed for immunohistochemistry against GFP, glutamate receptor 2/3 (GluR2/3) and
glutamate decarboxylase 67 (GAD67). **E**. Representative image of DP/DTT activity tagged neurons
(green). Scale bars:400μm (left), 100μm (right inset). **F**. Left: Representative image showing DP/DTT
OFT activity-tagged neurons (green) co-stained with GluR2/3 (magenta; arrows) or GAD67 (cyan; arrow
heads). Scale bar:50μm. Right: Distribution of DP/DTT OFT activity-tagged neurons indicates that these
cells predominantly co-label with GluR2/3 (67.9%), with 13.9% of neurons co-labeled with GAD67, 18.2%
negative for either marker, and a minority of cells (2.5%) co-labeled for both GluR2/3 and GAD67 (n=6).
483

Figure 5. Optogenetic activation of the DP/DTT-DMH pathway acutely suppresses freezing but has no effect on affective behavior.

A. Experimental timeline. Mice were injected with ChR2 or control GFP AAV constructs in the DP/DTT and implanted with an optic fiber in the DMH. Ten days later, animals underwent behaviour, starting with the OFT. Three days later, animals went through the EPM and, following another 3 days, one of two auditory fear conditioning protocols (see below). **B-D**. Behavior in the OFT. **B**. Light activation did not affect the time spent in the center of the open field between groups (Two-way rmANOVA significant main effect of stimulation $F_{1,18}$= 5.657, p=0.0287; but no main effect of treatment $F_{1, 18}$=0.1118, p=0.7420 or interaction $F_{1,18}$=0.02443, p=0.8775; no significant posthoc tests, p>0.17). **C**. Light activation did not affect the time spent in the outer zone of the open field between groups (Two-way rmANOVA significant main effect of stimulation $F_{1,18}$= 6.549, p=0.0197; but no main effect of treatment $F_{1, 18}$=0.11240, p=0.7288 or interaction $F_{1,18}$=0.1744, p=0.6811; no significant posthoc tests, p>0.097). **D**. The total distance travelled during the test was not affected by light activation (Two-way rmANOVA no main effect of treatment $F_{1, 18}$=2.724, p=0.1162; stimulation $F_{1,18}$= 3.054, p=0.0976; or interaction $F_{1,18}$=2.728, p=0.1160). **E-G**. Behavior in the EPM. Light activation did not affect the time spent in the open arms of

the EPM (**E**; Two-way rmANOVA no main effect of treatment $F_{1,14}$=0.9766, p=0.3398; stimulation $F_{1,14}$=
2.631, p=0.1271; or interaction $F_{1,14}$=0.0100, p=0.9217), the number of entries to the open arms of the
EPM (**F**; we note a non-significant trend toward a decrease in open arm entries in the EPM for the ChR2
animals: Two-way rmANOVA significant effect of stimulation $F_{1,14}$= 10.53, p=0.0059; but no main effect of
treatment $F_{1,14}$=3309, p=0.5743 or interaction $F_{1,14}$=0.1645, p=0.6912; Šídák's multiple comparisons test,
ChR2 p=0.057, GFP p=0.097) or the total distance travelled (**G**; Two-way rmANOVA no main effect of
treatment $F_{1,14}$=1.207, p=0.2904; stimulation $F_{1,14}$= 0.0676, p=0.7986; or interaction $F_{1,14}$=0.00511,
p=0.0940). **H-L**. Effect of DP/DTT-DMH activation on fear extinction and extinction retrieval. **H**.
Experimental timeline for the fear extinction cohort. Three days after the EPM, mice were trained in
auditory fear conditioning as described previously (6 tone-shock pairings). We note a significant effect of
tone on fear training in this cohort (data not shown; Two-way rmANOVA significant effect of tone $F_{3.537,}$
$_{35.37}$=50.62, $p<0.0001$; but no main effect of treatment $F_{1,10}$=1.792, p=0.2103 or interaction $F_{5,50}$=0.5963,
p=0.7029). The following day, mice underwent extinction training in context B (12 tones, as previously)
with light activation for the duration of each tone. On the next day, they underwent extinction retrieval in
context B (5 tones, same protocol as before) in the absence of light. **I**. During extinction training,
DP/DTT-DMH optogenetic activation reduced average freezing (**I**) and freezing across tones (**J**). **K-L**.
During extinction retrieval in the absence of light, average tone freezing (**K**; t_{10}=0.1392, p= 0.8921) and
freezing across tones (**L;** Two-way rmANOVA no significant main effect of treatment $F_{1,10}$=0.0193,
p=0.8921; tone $F_{2.707,27.07}$=2.575, $p=0.0798$ or interaction $F_{4,40}$=1.592, p=0.1953) were similar between
groups. **M-O**. Effect of DP/DTT-DMH activation on baseline fear after conditioning. **M**. Experimental
timeline for the fear baseline cohort. Three days after the EPM, a separate cohort of mice was trained in
auditory fear conditioning as previously (6 tone-shock pairings). The following day, mice were placed in
context B in the absence of tones, and after 2 minutes of baseline exploration the light was turned on for
2 minutes. **N**. DP/DTT-DMH optogenetic activation decreased freezing in the absence of tones. Two-way
rmANOVA significant main effect of stimulation and interaction, but not virus $F_{1,11}$=0.006, p=0.9396;
Šídák's multiple comparisons test, light OFF minus Light ON ChR2 p=0.0011, GFP p=0.7607. **O**.
DP/DTT-DMH optogenetic activation increased rearing instances. Two-way rmANOVA significant main
effect of stimulation and interaction, but not virus $F_{1,11}$=0.8904, p=0.3656; Šídák's multiple comparisons
test, light OFF minus Light ON ChR2 p=0.0008 GFP p=0.8609. *p<0.05. Male (square) and female
(circle) individual datapoints are displayed for transparency.

Supplementary Figure S1. Viral expression in the DP/DTT
A-B. Experimental timeline. C57BL/6J mice were injected with AAV5-hSyn-mCherry, AAV5-hSyn-hM3D-mCherry or AAV5-hSyn-hM4D-mCherry into the DP/DTT. Following 2 weeks, mice underwent a battery of behavioral tests comprised of the open field test (OFT), elevated plus maze (EPM), tail suspension test (TST), forced swim test (FST) and auditory fear conditioning (Fear) with a washout period of three days between each test. **C.** Representative images of an mCherry control virus-infused brain stained for mCherry (red) and the nuclear dye Hoechst (blue). **D-F.** Diagrams depicting overlaid viral spread for all mCherry (**D**), hM4D (**E**) and hM3D-injected (**F**) mice. Scale bar = 500μm.

Supplementary Figure S2. Additional behavioral metrics for DP/DTT soma manipulations on affective behavior.

A. Experimental timeline for the OFT. Mice previously injected with hM3D, hM4D or control mCherry constructs in the DP/DTT were injected with the DREADD agonist C21 one hour prior to being placed in an open field. The time spent in the center zone (red) and the outer zone (white) was measured. **B.** Time spent in the center of the open field arena was significantly reduced in hM3D mice compared to mCherry mice during minutes 2, 3, 4, and 5. The hM3D mice also spent less time in the center of the arena compared to hM4D mice during minute 2 (two way-rmANOVA, $F_{2,38}=7.795$, $p=0.002$, Tukey's post hoc tests: minute 2, hM3D mice versus mCherry mice, $p=0.047$ and hM3D versus hM4D mice $p=0.039$; minute 3, 4 and 5, hM3D versus mCherry, $p=0.03$, 0.031 and 0.04 respectively). ✱ indicates a significant difference between hM3D and mCherry groups; ✱ indicates a significant difference between hM3D and hM4D groups. **C**. Experimental timeline for the EPM test. Mice previously injected with hM3D, hM4D or control mCherry constructs in the DP/DTT were injected with the DREADD agonist C21 one hour prior to being placed in an elevated plus maze. **D**. hM3D mice spent significantly less time in the open arms of the elevated plus maze compared to mCherry mice (minute 1) (two-way repeated measures ANOVA (rmANOVA), main effect of treatment $F_{2,38}=5.867$, $p=0.006$ and minute $F_{4,152}=13.45$, $p<0.001$ but no interaction $F_{8,152}=0.711$, $p=0.681$; Tukey's post hoc tests: minute 1, hM3D versus mCherry, $p=0.008$; minute 3. **E**. The amount of time spent in the closed arms of the elevated plus maze was significantly greater in the hM3D mice compared to mCherry and hM4D mice (one-way ANOVA, $F_{2,38}=4.48$, $p<0.018$, Tukey's post hoc tests, hM3D versus mCherry, $p=0.052$, hM3D versus hM4D, $p=0.028$, no difference between the mCherry control and hM4D groups: $p=0.89$). **F**. The number of closed arm entries during the

test did not differ between groups ($F_{2,38}$=2.730, p=0.078). ✳ indicates a significant group difference
between mCherry and hM3D conditions. **G**. TST experimental timeline. Mice previously injected with
hM3D, hM4D or control mCherry constructs in the DP/DTT were injected with the DREADD agonist C21
one hour prior to undergoing the TST. **H**. Minute-by-minute analysis of immobility on the TST found that
hM3D mice spent significantly more time immobile than the hM4d mice during the third minute of the test.
A two-way repeated measures ANOVA found a significant main effect of treatment ($F_{2,38}$=5.077,
p=0.011), but no main effect of minute ($F_{3,114}$=1.553, p=0.205) or treatment by minute interaction
($F_{6,114}$=0.829, p=0.555). Tukey's post hoc test minute 3: hM3D mice versus hM4D mice, p=0.0275; ✳
indicates a significant difference between hM3D and hM4D groups. **I.** The longest continuous period of
immobility did not differ between groups (one-way ANOVA $F_{2, 38}$=1.874, p=0.167). **J.** The number of
mobility-immobility transitions (mobility bouts) did not differ between groups (one-way ANOVA
$F_{2,38}$=0.614, p=0.547). **K**. FST experimental timeline. Mice previously injected with hM3D, hM4D or
control mCherry constructs in the DP/DTT were injected with the DREADD agonist C21 one hour prior to
undergoing the FST. **L.** Minute-by-minute analysis of immobility on the FST. Percent immobility did not
differ between groups (Two-way rmANOVA, no main effect of treatment on percent immobility
$F_{2,38}$=1.723, p=0.192, no main effect of minute $F_{3,114}$=1.266, p=0.289, and no treatment by minute
interaction $F_{6,114}$=1.485, p=0.189). **M.** We noted a non-significant trend of treatment effect on the longest
bout of immobility (one-way ANOVA, $F_{2,38}$=3.22, p=0.051). **N.** The number of mobility-immobility
transitions (mobility bouts) did not differ between groups (one-way ANOVA, $F_{2,38}$=1.722, p=0.192).
*p<0.05. Male (square) and female (circle) individual datapoints are displayed for transparency (see text
for details).

585
Supplementary Figure S3. Confirmation of C21 effects on DP/DTT pyramidal neurons.
A-C. Validation of DP/DTT soma manipulations using cFos immunohistochemistry. **A.** Experimental
timeline for cFos analysis. Mice previously injected with hM3D, hM4D or control mCherry constructs in
the DP/DTT were injected with the DREADD agonist C21 and euthanized 90 minutes later. **B.**
Representative photomicrographs showing c-Fos (green) and mCherry (red) expression in the DP/DTT
following C21 treatment. Note the significant number of c-Fos$^+$ neurons in the hM3D mice. Hoechst
33342 (blue) was used as a nuclear marker. The lateral ventricle (dense blue stain on the right of each
photomicrograph) was used as a landmark for the DP/DTT. **C.** hM3D mice had significantly more c-Fos+
neurons in the DP/DTT than mCherry and hM4D mice (one-way ANOVA, $F_{2,12}$=143.9, p<0.001, Tukey's
post hoc tests, hM3D versus mCherry, p<0.001). Furthermore, mCherry mice also had significantly more
c-Fos+ neurons in the DP than hM4D mice (hM4D versus mCherry, p=0.030, hM3D versus hM4D,
p<0.001). Scale bar: 100µm. Male (square) and female (circle) individual datapoints are displayed for
transparency. **D-K.** Validation of DP/DTT soma manipulations using slice electrophysiology. **D.**
Experimental design. AAV constructs encoding hM3D-mCherry or hM4D-mCherry were injected into the
DP and slices were made at least 14 days later. **E.** Experimental timeline for electrophysiological
recordings. Slices were made and allowed to recover for 60 minutes prior to collecting baseline
recordings in mCherry$^+$ neurons. After a stable baseline recording of depolarization-induced firing, C21 (1
µM) was bath applied for at least 10 minutes and the last 5 minutes of the recording period were
compared to the baseline. **F.** Representative traces of action potentials recorded from an hM3D$^+$ neuron
at baseline and after application of C21. **G.** The number of action potentials evoked by somatic current
injection (100pA) in hM3D$^+$ transfected neurons was significantly increased following bath application of
C21 (paired t-test, t_{10}=4.604, p=0.001; n=11 cells, 4 mice). **H.** The resting membrane potential (RMP) of
hM3D$^+$ neurons became significantly more depolarized following bath application of C21 (paired t-test,
t_{10}=2.77, p=0.02). **I.** Representative traces of action potentials recorded from an hM4D$^+$ neuron at
baseline and after application of C21. **J.** The number of action potentials evoked by somatic current
injection (85pA) in hM4D$^+$ neurons was significantly decreased following bath application of C21 (paired
t-test, t_5=3.42, p=0.019; n=6 cells, 3 mice). **K.** The RMP of hM4D$^+$ neurons became significantly more
hyperpolarized following bath application of C21 (paired t-test, t_5=3.24, p=0.023). *p<0.05, ***p<0.001

615
Supplementary Figure S4. Validation of optogenetic manipulation of the DP/DTT-DMH pathway.
A. Experimental design. AAV constructs encoding GFP or ChR2 were injected into the DP/DTT and an
optical fiber was placed in the DMH. **B.** Representative image showing ChR2-EYFP expression in the DP
(left, scale bar=500µm) and the DP axonal projections to the DMH where the optical fiber was placed
(mid, scale bar=500µm). A higher-power inset of the DP/DTT-DMH projections and fiber location are
shown on the right (scale bar=200µm). **C.** Slices were prepared at least one week after behavior was
completed. **D-E**. Validation of DP/DTT optic activation in the DMH using slice electrophysiology. A light
pulse (5ms) triggers monosynaptic excitatory and inhibitory responses in patched DMH neurons of ChR2
mice (bottom), but not GFP mice (top). While Kataoka et al. demonstrated that their DP/DTT-DMH
effects were glutamatergic [68], future work is needed to determine the contribution of DP/DTT-triggered
excitation and inhibition in the DMH to our reported effects.

Supplemental Methods:

<u>Animals</u>

Adult male and female C57BL/6J and Fos$^{2A\text{-}iCreER}$ (TRAP2) mice (2-4 months old; Jackson laboratory)
were maintained on a 12 hour light-dark cycle (lights on at 7:00 A.M.), with experimental procedures
performed during the light phase. Mice were bred in-house and weaned on postnatal day 21 with same-
sex siblings (2-4 per cage). Food and water were available *ad libitum*. All experimental procedures were
approved by the Animal Care Committee at the University of Toronto.

<u>Stereotaxic surgery and viral injections</u>

Mice were injected with a cocktail of ketamine (100mg/kg) and xylazine (5mg/kg) to induce anesthesia.
Once mice were fully anesthetized, the head was shaved and swabbed with iodine followed by 70%
ethanol. Tear gel (Alcon) was applied to each eye to prevent dehydration. Mice were then secured in a
rodent stereotaxic apparatus (Stoelting) using ear bars. Body temperature was maintained at 37°C
throughout surgery with a far infrared homeothermic warming pad with a temperature feedback module
(RightTemp® Jr, Kent Scientific). A scalpel was used to make an incision down the midline of the scalp,
the connective tissue was excised, and the skull was cleaned with sterile phosphate buffered saline
(PBS, pH=7.4). An autoclaved cotton tip applicator was briefly submerged in 30% hydrogen peroxide and
then gently applied to the skull to identify bregma. A cordless micro drill (#58610, Stoelting) was used to
make craniotomies over the DP/DTT (+1.85 anterior-posterior, ±0.30 medial-lateral, -3.52mm dorso-
ventral, in reference to bregma) or IL (+1.9 anterior-posterior, ±0.26 medial-lateral, -2.9 dorso-ventral, in
reference to bregma). Virus was delivered with a 500nL Neuros Syringe (#65457-02, Hamilton Company)
attached to a probe holder on the stereotaxic apparatus (#751873, Harvard Apparatus). A total of 40-
100nL of virus was injected at a rate of 20nL per minute. The needle stayed in place for 5 minutes after
each injection to allow for diffusion of the virus before being slowly removed from the brain. We used the
following adeno-associated viruses (AAVs): AAV5-hSyn-mCherry (2.3 x 10^{13} vg/mL, Addgene #114472),
AAV5-hSyn-hM4D(Gi)-mCherry (8.6 x 10^{12} vg/mL, Addgene #50475), and AAV5-hSyn-hM3D(Gq)-
mCherry (1.9 x 10^{13} vg/mL, Addgene #50474), AAV1-CaMKIIa-hChR2(H134R)-EYFP (2.3 x 10^{13},
Addgene #26969), AAV1-CAG-GFP (1.2 x 10^{13} vg/mL, Addgene #37825) and AAV5-Ef1α-DIO-EYFP
(4.0 x 10^{12} vg/mL, UNC vector core). For optogenetic experiments, an optic fiber was implanted over the
DMH (-1.62 anterior-posterior, ±0.30 medial-lateral, -5.00 mm dorso-ventral, in reference to bregma).
Ketoprofen (1 mg/kg, s.c.) was used to reduce discomfort. Upon completing the viral injections, the skull
was cleaned with sterile PBS and the scalp was sutured with Vetbond tissue adhesive (3M). For
optogenetic experiments, the optic fiber was anchored by bone screws (HRS Scientific) and secured to
the skull using Metabond cement (Parkell, cat#S380). Warmed physiological saline (0.5 to 0.7mL, s.c.)

was injected at the end of each surgery to support hydration. Mice were placed in a clean cage located

on a heating pad for post-surgical recovery and returned to the colony room once fully ambulatory.

Ketoprofen (1 mg/kg, s.c.) was injected 24 and 48hr intervals after surgery to reduce post-surgical

discomfort.

<u>Tamoxifen injections</u>

4-hydroxytamoxifen (4-OHT; Cat# H4278, Sigma Aldrich) was mixed with ethanol (Cat# 6590-16, Fisher

Scientific) and Cremaphor (Cat# C5135, Sigma Aldrich) to make a 10mg/mL stock solution, which was

diluted 1:1 in 0.1M PBS and injected i.p. at a dose of 25mg/kg. TRAP2 mice were injected with AAV5-

Ef1α-DIO-EYFP in the DP/DTT and one week later underwent OFT. Immediately after the OFT, mice

were injected with 4-OHT and returned to the home cage. Animals were perfused 10 days later and their

brains processed for immunohistochemistry.

<u>Behavioral testing</u>

For experiments spanning Figures 1, 2 and 3A-E, following a 2-week recovery period after surgery,

cohorts of mice underwent the following behavioral tests interspersed by a 3-day washout period: 1)

open field test (OFT), 2) elevated plus maze (EPM), 3) tail suspension test (TST), 4) forced swim test

(FST) and 5) auditory fear conditioning. Behavioral task order was defined by degree of stress, starting

from the least to greater stress-inducing tasks to minimize carryover effects across tasks [134].

Correlation analyses between individual behavior in the OFT and EPM and subsequent freezing levels

during fear training or retrieval in hM3D and hM4D animals showed no significant differences (p>0.18,

data not shown), with the exception of a significant inverse correlation between the time spent in the

center zone of the open field and fear memory retrieval in the hM4D group (R^2=0.49, p=0.01). In these

experiments, the hM3D group showed a higher number of animals (7/15) with very weak spread to the

NAc compared to the hM4D group (2/11). Importantly, hM3D animals with or without NAc spread showed

no significant difference in behavior in the OFT, EPM, TST or auditory fear (p>0.33, data not shown),

suggesting this is not a major contributor to our effects. Five cohorts of mice were used in these

experiments, and we observed consistent effects of DP/DTT manipulation across all cohorts.

For experiments in Figure 3F-L, animals underwent fear conditioning following recovery from surgery

without prior behavior. Six cohorts of mice were used in these experiments, and we observed consistent

effects of DP/DTT manipulation across all cohorts. In the experiments featured in Figure 5, all cohorts

underwent the following behavioral tests interspersed by a 3-day washout period: 1) OFT, 2) EPM, 3)

auditory fear training, followed the next day by either (a) fear extinction training and extinction retrieval

(Figure 5H-L) or (b) baseline freezing testing (Figure 5M-O; see details below). Three cohorts of mice

were used in each of these experimental designs, and we observed consistent effects across all cohorts.

The DREADD agonist compound 21 (C21; 2mg/kg, 0.5mg/mL, i.p.) was injected 1hr before the start of

each behavioral test. Mice remained in their homecage until behavioral testing began. Detailed methods

for each behavioral test are described below. Upon completing behavioral testing, mice were returned to

their homecage and left undisturbed until the next test. All behavioral testing apparatuses were cleaned

with 70% ethanol followed by ddH_2O between subjects. Behavior was automatically scored using ANY-

maze software (Stoelting) (OFT, EPM), Depression Behavior Scorer (DBscorer) [135] (TST, FST) or

VideoFreeze [136,137] (Med Associates) (fear).

Open Field Test

Exploratory and anxiety-like behaviors were measured in the OFT as previously described [138–141]

with minor modifications. Mice were placed in the corner of a brightly illuminated open field arena (30cm

length x 30cm width x 30cm height) and an overhead monochrome video camera (DMK 22AUC03, The

Imaging Source) recorded behavior over the 8-minute test duration with ANY-maze software (version

7.1, Stoelting). The primary measures of interest included the time spent in the center zone of the maze

(10cm x 10cm), time spent in the outer zone, and the total distance travelled during the test. Track plots

and heat maps were generated using ANY-maze software. Optogenetic stimulation of the DP/DTT-DMH

pathway (473 nm, 20 Hz, 5ms pulses) occurred after 3 minutes of baseline activity with the light off, and

lasted for 3 minutes.

Elevated Plus Maze

Exploratory and anxiety-like behaviors were evaluated in the EPM as previously described [141,142]

with minor modifications. The EPM apparatus was comprised of two open and two closed arms of

identical dimensions (35cm length x 10cm width) raised 51.5cm above the floor. The closed arms had

30cm tall walls, whereas the open arms had a 0.1cm raised ledge to prevent mice from falling off the

apparatus during testing. At the start of each test, mice were placed in the middle of the platform (10cm

length x 10cm width) with their head facing towards one of the closed arms. A Logitech C920 1080P

webcam was placed directly above the EPM and recorded behavior over the 5-minute test duration via Logi Capture Software (v 2.06.12, Logitech). Video recordings were loaded in ANY-maze software and EPM behavior was analyzed offline. The primary measures of interest included the time mice spent in the open and closed arms of the EPM and the time spent in the middle platform. The software also tracked the number of entries into each zone and the total distance travelled during the test. Track plots and heatmaps were generated using ANY-maze software. Optogenetic stimulation of the DP/DTT-DMH pathway (473 nm, 20 Hz, 5ms pulses) occurred after 3 minutes of baseline activity with the light off, and lasted for 3 minutes.

Tail Suspension Test

The TST was used to measure passive coping behavior as previously described [143,144]. Mice were suspended approximately 45 cm above the floor by their tails for 6 minutes. Laboratory tape was placed <1cm from the tip of each tail and secured to a horizontal stainless-steel rod attached to a laboratory retort stand. Pairs of mice were positioned approximately 40cm apart from each other during testing. A Logitech C920 1080P webcam was placed horizontally in front of the mice and recorded behavior over the test duration. The final 4 minutes of the TST was analyzed using the open-source software DBscorer, which performs automated and unbiased behavioral scoring [135]. This software converts each video frame to a binary image and compares the percent area difference between each frame, which are averaged for every second of the test duration. When the percent area difference is below a user-defined threshold value, the animal is considered immobile. Pilot experiments found that a 0.6% area threshold value was consistent with manually scored videos. The data was automatically scored in 60s time bins and consists of the percent time spent immobile, the longest continuous bout of immobility during the test (in seconds), and the number of mobility-immobility transitions. The software also automatically generated raster plots of immobility over the test duration for each mouse.

Forced Swim Test

The FST was used to measure depression-like behavior [145–147]. Mice were individually placed in a 1L beaker (14.5cm tall) that was filled with approximately 750mL of warmed tap water (25°C). A Logitech C920 1080P webcam was placed horizontally in front of the beaker and recorded behavior for 6 minutes. The final 4 minutes of the FST was analyzed [148,149] using DBscorer with an area threshold value of 1.0% [135]. Videos were scored in 60s time bins and measured the percent of time spent immobile, the longest continuous bout of immobility (in seconds), and the number of mobility-immobility transitions. Raster plots were also generated to indicate periods of mobility and immobility throughout the 4-minute analysis period. Upon completing the FST, mice were dried off with a towel, placed in a fresh cage on a heating blanket for 30 minutes, and returned to the colony room. The water in each beaker was changed between subjects.

Auditory Fear Conditioning

Auditory fear conditioning was used to evaluate whether manipulations of the DP/DTT influenced cognition. Auditory fear training occurred in a stainless-steel fear conditioning apparatus (32cm wide, 25.5cm high, 25.5cm deep) that contained shock-grid floors (36 rods, 2mm diameter; Context A). The fear conditioning apparatus was located within a sound-attenuated chamber (63.5cm wide, 36.8cm, high, 74.9cm deep; catalog #NIR-022MD, Med Associates). On the training day, a two-minute acclimation period was provided to assess baseline freezing behavior. A total of 6 auditory tones (2 kHz, 80 dB, 25s) were presented with a 60s interval and co-terminated with a 2s foot shock (0.5mA), adapted from our previous studies [64,137]. Upon completing the training session, mice were placed in their home cage and returned to the colony room until the following day.

Fear memory was evaluated 24 hours after the training session. The fear conditioning apparatus was modified for memory retrieval by installing a curved white plastic insert along the walls and a white plastic insert over the shock-grid floors (Context B). Baseline freezing behavior was evaluated in the first two minutes of the test. A total of 6 auditory tones (same parameters as training) were presented every minute to evaluate fear memory. Mice were returned to their home cage upon completing the fear memory test.

For experiments manipulating DP/DTT activity during extinction, separate cohorts of animals underwent surgery and ten days later were trained in auditory fear training as described above (no prior behavioral testing). Upon completing the training session, mice were placed in their home cage and returned to the colony room. The following day, for extinction training, animals were placed in context B and after a two-minute acclimation period were exposed to 12 tones (2 kHz, 80 dB, 25s) with a 35s interval. Mice were returned to their home cage and the following day underwent extinction retrieval. At extinction retrieval, animals were placed in context B and, and after a two-minute acclimation period, were exposed to 5 tones (2 kHz, 80 dB, 25s) with a 35s interval. For all DP/DTT soma extinction experiments, animals were perfused for histological confirmation of targeting, and all animals displaying viral spread to IL were excluded from the dataset.

For the experiments examining the effect of optogenetic stimulation of the DP/DTT-DMH pathway during extinction training, animals previously trained in OFT and EPM underwent fear training as described previously. The next day, during extinction training (same protocol as above), light stimulation (473 nm, 20 Hz, 5ms pulses) occurred for the duration of each of the 12 tones. Mice were returned to their home cage and the following day underwent extinction retrieval (same parameters as above) in the absence of light stimulation.

For the experiment looking at the effect of DP/DTT-DMH stimulation on baseline freezing, a separate cohort of animals previously trained on OFT and EPM (but not fear) underwent fear conditioning as

described previously in the absence of light stimulation. The following day, animals were placed in
context B and, after 2 minutes with the light off, underwent light stimulation (473 nm, 20 Hz, 5ms pulses)
for 2 minutes in the absence of tones. Rearing was hand-scored by an experimenter blind to
experimental conditions, whereby a rearing instance was counted when a mouse elevated both front
paws off the floor (supported and unsupported rearing [123] were conflated in our analysis). In this
cohort, some animals showed spread into the LS, but an excluded animal in which only the LS was
targeted did not show the same behavioural profile as the DP/DTT-DMH targeted animals. Nevertheless,
future experiments must definitively exclude a contribution of the LS to these behavioral effects.

Freezing behavior was scored automatically with Med Associates VideoFreeze software as previously
described [136,137]. Due to interference of the patch cords with automated freezing counts, in all trials
with patch cords freezing was hand-scored and cross-checked by an experimenter blind to experimental
conditions.

Perfusions and sectioning

Mice were euthanized 90 minutes after the fear memory retrieval test. Mice were injected with avertin
(250mg/kg, i.p.) to induce deep anesthesia. Once unresponsive to tail and toe pinches, mice were
transcardially perfused with 0.1M phosphate-buffered saline (PBS), followed by cold 4%
paraformaldehyde fixative (PFA). The brains were extracted and stored in PFA overnight at 4°C. Brains
were sectioned at 50μm (1 in 6 series) in the coronal plane with a vibratome (model VT1000, Leica).
Sections were stored at -20°C in a cryoprotectant solution comprised of 60% glycerol and 0.01% sodium
azide in 0.1M PBS.

Serum corticosterone

A separate cohort of mice with no previous behavioral testing history (n=5 per group) was used to
determine the effect of DP/DTT manipulations on serum corticosterone. The DP/DTT was injected with
mCherry, hM4D, or hM3D viral constructs using identical parameters to those described earlier. After a
2-week recovery period, mice were weighed and acclimated to a procedure room for at least one hour.
Mice were injected with C21 (2mg/kg, 0.5mg/mL, i.p.) and immediately returned to their home cage. Mice
remained in their home cage for 90 minutes undisturbed and were then euthanized as described above.
Immediately prior to perfusion, blood was extracted from the heart (~0.3 to 0.5mL) with a 23-gauge
needle attached to a 1mL syringe. The blood was immediately transferred to a 1.5mL Eppendorf tube
and allowed to coagulate at room temperature for 30 minutes. Blood samples then underwent
centrifugation to isolate serum (10,000 rpm, 10 minutes). The isolated serum was extracted with a 1mL
pipette, transferred into a fresh microcentrifuge tube, and flash frozen with liquid nitrogen. Samples were
stored at -20°C until use. Serum corticosterone was evaluated with a colorimetric competitive enzyme
immunoassay kit (#ADI-900-097, Enzo Life Sciences) following the manufacturer's instructions. Briefly,

serum samples (1 in 40 dilution) were run in duplicate and read at 405nm with 570nm correction on a

Biotek Synergy Neo2 Multi-Mode Plate Reader (Agilent) and analyzed using Biotek Gen5 software

(version 3.08, Agilent).

<u>Immunofluorescence</u>

Immunofluorescence staining was performed on free-floating sections. Sections were incubated with any

of the following primary antibodies: polyclonal rabbit anti-mCherry (1:2000, catalog #ab167453, Abcam;

RRID: AB_2571870), polyclonal chicken anti-GFP (1:3000, catalog #13970, Abcam; RRID: AB_371416),

polyclonal rabbit anti-GluR2/3 (1:75, catalog # AB1506, Millipore; RRID: AB_90710), monoclonal mouse

anti-GAD67 (1:250, catalog # MAB5406, Millipore; RRID: AB_2278725) overnight at 4°C on a rotary

shaker under gentle agitation. On the following day, sections were washed in 0.1M PBS and then

incubated with the following secondary antibodies: goat anti-rabbit Alexa Fluor 568 (1:500, catalog

#A11011, Thermo Fisher Scientific; RRID: AB_143157), goat anti-mouse Alexa Fluor 633 (1:200, catalog

SAB4600138, Millipore), goat anti-chicken Alexa Fluor 488 (1:500, catalog # A11039, Thermo Fisher

Scientific; RRID: AB_2534096), for 2hrs at room temperature. Hoechst 33342 (1:1000 diluted in 0.1M

PBS, Thermo Fisher Scientific) was used as a counterstain. Sections were then rinsed in 0.1M, mounted

onto gelatin-coated slides, and allowed to air dry for 30 minutes. The mounted sections were then

coverslipped with Citifluor anti-fade mounting medium (catalog #17970, Electron Microscopy Sciences)

or permafluor antifade medium (catalog # TA030FM, Fisher Scientific) and stored in a microscope slide

storage book at 4°C until imaging.

To determine whether C21 modified DP/DTT activity across treatment groups, we processed a subset of

PFC sections for the immediately early gene c-Fos. We chose sections from mice used in the serum

corticosterone experiment because they did not undergo any behavioral testing which could influence

patterns of c-Fos expression. Immunofluorescence staining for c-Fos was performed as previously

described [136]. Sections were rinsed in 0.1M PBS and then incubated in 1% H_2O_2 in 0.1M PBS to

quench endogenous peroxidase activity. Sections were incubated overnight at 4°C with monoclonal

rabbit anti c-Fos (1:8000, catalog #226 008, Synaptic Systems; RRID: AB_2891278) and monoclonal rat

anti-mCherry (1:3000, catalog #M11217, Thermo Fisher Scientific; RRID: AB_2536611) diluted in 0.1M

Tris-buffered saline and 0.5% (w/v) Roche Blocking Reagent (catalog #11096176001, Sigma-Aldrich).

On the following day, sections were rinsed in 0.1M PBS and incubated in goat anti-rat Alexa Fluor 568

secondary antibody (1:500, catalog #A11077, Thermo Fisher Scientific; RRID: AB_2534121) for 1 hr.

Sections were then rinsed in 0.1M PBS and incubated in donkey anti-rabbit horseradish peroxidase-

conjugated secondary antibody (1:500, catalog #711-036-152, Jackson ImmunoResearch; RRID:

AB_2340590) for 1hr. Tyramide signal amplification was then performed with a fluorescein tyramide

conjugate (1:400) diluted in 0.1M borate buffer containing 0.01% H_2O_2. Hoechst 33342 was used as a

counterstain using the same parameters described above. Sections were then mounted onto gelatin-
coated slides, air dried, and coverslipped with Citifluor anti-fade mounting medium as described above.

<u>Image acquisition & quantification</u>

Images were acquired on a Nikon Eclipse Ni-U epifluorescence microscope connected to a computer
running NIS-Elements software (v 5.11.03, Nikon). Immunofluorescence signal was visualized with a
LED illumination system (X-Cite 120 LED Boost, Excelitas Technologies) and 10x Plan-Apochromat
objective. Images were captured with a Nikon DS-Qi2 digital camera. A total of 3 images per mouse were
used to evaluate viral expression in the DP/DTT for the mCherry, hM4D-mCherry, and hM3D-mCherry
cohorts. We evaluated viral expression via the mCherry tag in sections located approximately +1.9mm to
+1.7mm anterior-posterior (relative to bregma). Photomicrographs were superimposed on plates from the
Paxinos Mouse Brain atlas and the spread of viral expression was drawn using the polygonal tool in
Adobe Photoshop (version 24.1). Viral expression area for each mouse was represented using a semi-
transparent color setting, as this helped visualize the areas with the most concentrated viral expression
and viral spread across subjects.

For the DP/DTT TRAP neurochemical analysis, images were acquired using a Leica Stellaris 5 laser
scanning confocal microscope. Confocal 1um Z-stack images of the DP/DTT were obtained and
manually counted using LASX Office (version 1.4.5 27713) software (Leica).

Quantification of c-Fos expression in the DP/DTT was performed with ImageJ Software (version 1.53e)
using procedures described previously [136]. Briefly, a minimum of 3 PFC sections per subject were
used to count the number of c-Fos immunoreactive cells in the DP/DTT. The average number of cells per
section was calculated by summing the total number of c-Fos+ cells across both hemispheres and
dividing by the number of sections that were analyzed for each subject.

All figures were made in Adobe Photoshop. When brightness and/or contrast adjustments were made,
these changes were applied to all photomicrographs.

<u>Slice electrophysiology</u>

We used slice electrophysiology to confirm the effect of the DREADD agonist C21 on hM3D- and hM4D-
transfected neurons in the DP/DTT and the effect of optical stimulation on ChR2 and GFP transfected
DP/DTT axon terminals projecting to neurons in the DMH. Acute brain slices containing the mPFC were
prepared from adult male and female C57BL/6J mice at least 14 days after AAV injection. For
optogenetic experiments, electrophysiology recordings were conducted at least 1 week after the animals
underwent behavioral testing.

Mice were deeply anesthetized with isoflurane and decapitated. The brain was quickly removed and
placed in ice-cold sucrose based cutting solution containing the following (in mM): 180 sucrose, 2.5 KCl,

1.25 NaH_2PO_4, 25 $NaHCO_3$, 1 $CaCl_2$, 2 $MgCl_2$, 2 Na^+ pyruvate and 0.4 L-Ascorbic acid with 95% O_2 and

5% CO_2. The brain was cut in the coronal plane (300 μm thick) using a vibratome (VT1000S, Leica).

Brain slices containing the mPFC were allowed to recover for 30 min in a holding chamber containing a

recovery solution at 30 °C, made up of 50% sucrose-based cutting solution and 50% artificial

cerebrospinal fluid (ACSF). The ACSF was composed of (in mM): 120 NaCl, 2.5 KCl, 1.25 NaH_2PO_4, 25

$NaHCO_3$, 11 glucose, 2 $CaCl_2$, 1 $MgCl_2$, 2 Na^+ pyruvate and 0.4 L-Ascorbic acid with 95% O_2 and 5%

CO_2. Slices then underwent an additional 30 min recovery in regular ACSF at room temperature before

patch-clamp recordings. After recovery, individual slices were transferred to a submersion chamber,

perfused at 2 mL/min with ACSF at room temperature, and mounted on an upright microscope (Eclipse

FN1, Nikon) that was equipped with a water-immersion long-working distance objective (40x),

epifluorescence filters, and an infrared video camera [DAGE IR-2000].

For DREADD experiments, current-clamp recordings were obtained from mCherry+ pyramidal neurons in

layers 2/3 or 5 of the DP/DTT with borosilicate pipettes (3–5 MΩ) that were filled with intracellular solution

containing the following (in mM): 126 K D-Gluconate, 5 KCl, 10 HEPES, 4 MgATP, 0.3 NaGTP, 10 Na-

phosphocreatine. Baseline data was acquired by measuring resting membrane potential (RMP) and the

number of action potentials evoked by a long (1s) depolarizing current pulse. Somatic current pulses

were calibrated in the baseline period to trigger approximately 3-4 action potentials (hM3D-mCherry$^+$) or

7-8 action potentials (hM4D-mCherry$^+$ pyramidal neurons). After 5 minutes of stable neuronal discharge

induced by a somatic depolarization, C21 (1 μM, HelloBio) [150] was bath applied for at least 10 minutes,

with the last 5 min being compared to the baseline period.

For optogenetic experiments, voltage clamp recordings were obtained from DMH neurons with

borosilicate pipettes (3-5 MΩ) filled with intracellular solution containing the following (in mM): 130 Cs-

methanosulfonate, 10 HEPES, 0.5 EGTA, 8 NaCl, 4 Mg-ATP, 0.4 Na-GTP, 10 Na-phosphocreatine, 1 N-

ethyl lidocaine (QX-314). DP/DTT axon terminals were stimulated using a TTL-pulsed microscope

objective-coupled light-emitting diodes (460 nm, ~1.36 mW/mm2; Prizmatix). Evoked optical EPSCs and

IPSCs were recorded at -70mV and 0mV from both ChR2 and GFP control transfected animals,

respectively. All light-induced stimulation was conducted at 0.1 Hz to avoid inducing synaptic plasticity.

Electrophysiological data was acquired using a MultiClamp 700B amplifier (Molecular Devices). Signals

were low-pass filtered at 2kHz and digitized at 20 kHz using Digidata 1550A and pClamp10 software

(Molecular devices). Series resistance of the recorded neurons was regularly monitored and only cells

with stable series resistance (less than 20% change of the initial value) and stable holding current were

included in the experimental data.

<u>Statistical analysis</u>

All results are presented as the mean ± standard error of the mean (SEM). Statistical comparisons were

made in Prism 9.1 (GraphPad) with statistical significance ($p<0.05$) denoted on all graphs with an

asterisk. Non-statistically significant results are reported in the figure legends due to space limits. All data

were analyzed for sex differences, with non-statistically significant results reported below. To facilitate

interpretation of the data, and in the absence of major differences within and across tasks, we pooled

male and female mice, with individual data points presented as squares (males) and circles (females) for

transparency.

For within subject comparisons, a paired t-test was used. Comparisons of independent groups were

made using a one-way ANOVA, followed by Tukey's Post Hoc test when appropriate. Parametric data

with multiple comparisons were analyzed using a two-way repeated measures ANOVA, followed by

Tukey's or Šídák's Post Hoc test with corrections for multiple comparisons. Normality of parametric

datasets were confirmed by the D'Agostino and Pearson normality test (Prism 9.1).

<u>Sex differences</u>

Potential sex differences were examined for each dataset using two-way ANOVAs followed by Šídák's

multiple comparisons test, and reported in the main text when significant. To facilitate interpretation of

the data, and in the absence of major differences within and across tasks, we pooled male and female

mice for each dataset, with individual data points presented as squares (males) and circles (females).

For the OFT, we found no effect of sex on the time spent in the center zone (Two-way ANOVA,

$F_{1,35}=1.065$, $p=0.309$), outer zone ($F_{1,35}=0.553$, $p=0.462$), or total distance travelled ($F_{1,35}=2.351$,

$p=0.134$). For the EPM, we found no sex differences in the time spent in the open arms ($F_{1,35}=3.069$,

$p=0.009$), time spent in the closed arms ($F_{1,35}=1.928$, $p=0.174$), open arm entries ($F_{1,35}=0.386$, $p=0.538$),

or closed arm entries ($F_{1,35}=0.988$, $p=0.327$). We found no sex differences in the TST (Two-way

ANOVAs, time immobile $F_{1,35}=1.712$, $p=0.199$; longest immobility bout $F_{1,35}=0.061$, $p=0.807$; number of

mobility-immobility transitions $F_{1,35}=2.044$, $p=0.162$). In the FST, hM3D females spent significantly less

time immobile than hM3D males and hM4D females (Two-way ANOVA, $F_{1,35}=6.561$, $p=0.015$; Sidak's

multiple comparisons test, hM3D females versus hM3D males $p=0.0093$; hM3D females versus hM4D

females $p=0.0482$; mCherry male versus female, $p=0.264$; hM4D male versus female, $p=0.998$). We did

not observe any sex differences in the longest bout of immobility ($F_{1,35}=2.696$, $p=0.107$), or the number of

mobility-immobility transitions ($F_{1,35}=3.369$, $p=0.075$) in the FST. In auditory fear acquisition, while we

found a significant main effect of sex on average freezing to tone (Two-way ANOVA, $F_{1,35}=7.786$,

$p=0.009$). Post-hoc tests found that female hM4D mice froze significantly more than male hM3D mice

($p=0.0479$), but no other sex differences were observed (all p values >0.110). There were no sex

differences in average tone freezing (Sidak's multiple comparisons test, average tone freezing: mCherry

p=0.126, hM3D p=0.351, hM4D p=0.505). We also failed to see sex differences in baseline freezing

(two-way ANOVA, $F_{1,35}$=0.179, p=0.675). We found no sex differences in auditory fear memory retrieval

(Two-way ANOVA, average baseline freezing $F_{1,35}$=0.397, p=0.533, average tone freezing $F_{1,35}$=0.075,

p=0.786).

Overall, we did not see sex differences in baseline OFT or EPM behavior. While studies in rats show

consistent sex differences [151] in the EPM [151–157] and OFT [98,100,101,103,104 but see 105,106]

which are often replicated in mice [160–164], several mouse studies show reversals or no sex

differences [151,165–168], particularly in the OFT [151,160,168,169]. Although our lack of sex

differences in the OFT replicates similar findings in C57BL6 mice [160,168,169], we unexpectedly found

sex differences only in hM4D mice in the EPM. Similarly, our sex differences in the TST were only found

among DREADD injected animals. While we cannot explain the lack of sex differences in the matched

controls in these experiments, factors such as age [170,171] and estrous cycle [172,173] are known to

modulate sex differences in affective behavior, and could have contributed to our findings. Although we

have not seen major sex differences in the behaviors explored here, one study reported sex-specific

increased recruitment of the DP in female mice following auditory fear conditioning [174]. Due to the well

documented sex and gender differences in the prevalence of anxiety and posttraumatic stress (PTSD)

disorders [175–177], as well as in sympathetic and autonomic function [178–181], this raises the

intriguing possibility of sex differences in the long-term consequences of stress-induced activation of DP

pathways.

References:

1. Knight RT, Grabowecky MF, Scabini D. Role of human prefrontal cortex in attention control. Adv Neurol. 1995;66:21–26.

2. Rossi AF, Pessoa L, Desimone R, Ungerleider LG. The prefrontal cortex and the executive control of attention. Exp Brain Res. 2009;192:489–497.

3. Bechara A, Tranel D, Damasio H. Characterization of the decision-making deficit of patients with ventromedial prefrontal cortex lesions. Brain. 2000;123 (Pt 1:2189–2202.

4. Euston DR, Gruber AJ, McNaughton BL. The role of medial prefrontal cortex in memory and decision making. Neuron. 2012;76:1057–1070.

5. Bahmani Z, Clark K, Merrikhi Y, Mueller A, Pettine W, Isabel Vanegas M, et al. Prefrontal Contributions to Attention and Working Memory. Curr Top Behav Neurosci. 2019;41:129–153.

6. Vogel P, Hahn J, Duvarci S, Sigurdsson T. Prefrontal pyramidal neurons are critical for all phases of working memory. Cell Rep. 2022;39:110659.

7. Yizhar O, Fenno LE, Prigge M, Schneider F, Davidson TJ, Shea DJO, et al. Neocortical excitation / inhibition balance in information processing and social dysfunction. Nature. 2011;477:171–178.

8. Ko J. Neuroanatomical Substrates of Rodent Social Behavior: The Medial Prefrontal Cortex and Its Projection Patterns. Front Neural Circuits. 2017;11:41.

9. Yizhar O, Levy DR. The social dilemma: prefrontal control of mammalian sociability. Curr Opin Neurobiol. 2021;68:67–75.

10. Kuga N, Abe R, Takano K, Ikegaya Y, Sasaki T. Prefrontal-amygdalar oscillations related to social behavior in mice. Elife. 2022;11.

11. Etkin A, Egner T, Kalisch R. Emotional processing in anterior cingulate and medial prefrontal cortex. Trends Cogn Sci. 2011;15:85–93.

12. Price JL, Drevets WC. Neural circuits underlying the pathophysiology of mood disorders. Trends Cogn Sci. 2012;16:61–71.

13. Adhikari A, Topiwala M a, Gordon J a. Synchronized activity between the ventral hippocampus and the medial prefrontal cortex during anxiety. Neuron. 2010;65:257–269.

14. Adhikari A. Distributed circuits underlying anxiety. Front Behav Neurosci. 2014.

15. Likhtik E, Stujenske JM, Topiwala MA, Harris AZ, Gordon JA. Prefrontal entrainment of amygdala activity signals safety in learned fear and innate anxiety. Nat Neurosci. 2014;17:106–113.

16. Adhikari A, Topiwala MA, Gordon JA. Single units in the medial prefrontal cortex with anxiety-related firing patterns are preferentially influenced by ventral hippocampal activity. Neuron. 2011;71:898–910.

17. Yu H, Chen L, Lei H, Pi G, Xiong R, Jiang T, et al. Infralimbic medial prefrontal cortex signalling to calbindin 1 positive neurons in posterior basolateral amygdala suppresses anxiety- and depression-like behaviours. Nat Commun. 2022;13:1–14.

18. Sullivan RM, Gratton A. Behavioral effects of excitotoxic lesions of ventral medial prefrontal cortex in the rat are hemisphere-dependent. Brain Res. 2002;927:69–79.

19. Shah AA, Treit D. Excitotoxic lesions of the medial prefrontal cortex attenuate fear responses in the elevated-plus maze, social interaction and shock probe burying tests. Brain Res. 2003;969:183–194.

1017   20.   Lacroix L, Spinelli S, Heidbreder CA, Feldon J. Differential role of the medial and lateral prefrontal
1018   cortices in fear and anxiety. Behav Neurosci. 2000;114:1119–1130.

1019   21.   Suzuki S, Saitoh A, Ohashi M, Yamada M, Oka JI, Yamada M. The infralimbic and prelimbic
1020   medial prefrontal cortices have differential functions in the expression of anxiety-like behaviors in
1021   mice. Behav Brain Res. 2016;304:120–124.

1022   22.   Cao J, Liu X, Liu JX, Zhao S, Guo YX, Wang GY, et al. Inhibition of glutamatergic neurons in layer
1023   II/III of the medial prefrontal cortex alleviates paclitaxel-induced neuropathic pain and anxiety. Eur
1024   J Pharmacol. 2022;936.

1025   23.   Parfitt GM, Nguyen R, Bang JY, Aqrabawi AJ, Tran MM, Seo DK, et al. Bidirectional Control of
1026   Anxiety-Related Behaviors in Mice: Role of Inputs Arising from the Ventral Hippocampus to the
1027   Lateral Septum and Medial Prefrontal Cortex. Neuropsychopharmacology. 2017;42:1715–1728.

1028   24.   Jinks AL, McGregor IS. Modulation of anxiety-related behaviours following lesions of the prelimbic
1029   or infralimbic cortex in the rat. Brain Res. 1997;772:181–190.

1030   25.   Pati S, Sood A, Mukhopadhyay S, Vaidya VA. Acute pharmacogenetic activation of medial
1031   prefrontal cortex excitatory neurons regulates anxiety-like behaviour. J Biosci. 2018;43:85–95.

1032   26.   Adhikari A, Lerner TN, Finkelstein J, Pak S, Jennings JH, Davidson TJ, et al. Basomedial
1033   amygdala mediates top-down control of anxiety and fear. Nature. 2015;527:179–185.

1034   27.   Yin J Bin, Liang SH, Li F, Zhao WJ, Bai Y, Sun Y, et al. dmPFC-vlPAG projection neurons
1035   contribute to pain threshold maintenance and antianxiety behaviors. J Clin Invest. 2020;130:6555–
1036   6570.

1037   28.   Soumier A, Sibille E. Opposing effects of acute versus chronic blockade of frontal cortex
1038   somatostatin-positive inhibitory neurons on behavioral emotionality in mice.
1039   Neuropsychopharmacology. 2014;39:2252–2262.

1040   29.   Calhoon GG, Tye KM. Resolving the neural circuits of anxiety. Nat Neurosci. 2015;18:1394–1404.

1041   30.   Wang GQ, Cen C, Li C, Cao S, Wang N, Zhou Z, et al. Deactivation of excitatory neurons in the
1042   prelimbic cortex via Cdk5 promotes pain sensation and anxiety. Nat Commun. 2015;6:7660.

1043   31.   Sesack SR, Deutch a Y, Roth RH, Bunney BS. Topographical organization of the efferent
1044   projections of the medial prefrontal cortex in the rat: an anterograde tract-tracing study with
1045   Phaseolus vulgaris leucoagglutinin. J Comp Neurol. 1989;290:213–242.

1046   32.   Miller EK, Cohen JD. An Integrative Theory of Prefrontal Cortex Function. Annu Rev Neurosci.
1047   2001;24:167–202.

1048   33.   Gabbott PLA, Warner TA, Jays PRL, Salway P, Busby SJ. Prefrontal cortex in the rat: Projections
1049   to subcortical autonomic, motor, and limbic centers. J Comp Neurol. 2005;492:145–177.

1050   34.   Hoover WB, Vertes RP. Anatomical analysis of afferent projections to the medial prefrontal cortex
1051   in the rat. Brain Struct Funct. 2007;212:149–179.

1052   35.   Ährlund-Richter S, Xuan Y, van Lunteren JA, Kim H, Ortiz C, Pollak Dorocic I, et al. A whole-brain
1053   atlas of monosynaptic input targeting four different cell types in the medial prefrontal cortex of the
1054   mouse. Nat Neurosci. 2019;22:657–668.

1055   36.   Anastasiades PG, Carter AG. Circuit organization of the rodent medial prefrontal cortex. Trends
1056   Neurosci. 2021;44:550–563.

1057   37.   Riga D, Matos MR, Glas A, Smit AB, Spijker S, Van den Oever MC. Optogenetic dissection of
1058   medial prefrontal cortex circuitry. Front Syst Neurosci. 2014;8:230.

1059   38.   Bittar TP, Labonte B. Functional Contribution of the Medial Prefrontal Circuitry in Major Depressive

Disorder and Stress-Induced Depressive-Like Behaviors. Front Behav Neurosci. 2021;15:699592.

39. Carlén M. What constitutes the prefrontal cortex? Science (80-). 2017;358:478–482.

40. Le Merre P, Ahrlund-Richter S, Carlen M. The mouse prefrontal cortex: Unity in diversity. Neuron. 2021;109:1925–1944.

41. Xu P, Chen A, Li Y, Xing X, Lu H. Medial prefrontal cortex in neurological diseases. Physiol Genomics. 2019;51:432–442.

42. George O, Koob GF. Individual differences in prefrontal cortex function and the transition from drug use to drug dependence. Neurosci Biobehav Rev. 2010;35:232–247.

43. Riga D, Matos MR, Glas A, Smit AB, Spijker S, Van den Oever MC. Optogenetic dissection of medial prefrontal cortex circuitry. Front Syst Neurosci. 2014;8:230.

44. Chini M, Hanganu-Opatz IL. Prefrontal Cortex Development in Health and Disease: Lessons from Rodents and Humans. Trends Neurosci. 2021;44:227–240.

45. Xu P, Chen A, Li Y, Xing X, Lu H. Medial prefrontal cortex in neurological diseases. Physiol Genomics. 2019;51:432–442.

46. Bishop S, Duncan J, Brett M, Lawrence AD. Prefrontal cortical function and anxiety: controlling attention to threat-related stimuli. Nat Neurosci. 2004;7:184–188.

47. Myers-Schulz B, Koenigs M. Functional anatomy of ventromedial prefrontal cortex: implications for mood and anxiety disorders. Mol Psychiatry. 2012;17:132–141.

48. Park J, Moghaddam B. Impact of anxiety on prefrontal cortex encoding of cognitive flexibility. Neuroscience. 2017;345:193–202.

49. Andersen SL, Teicher MH. Stress, sensitive periods and maturational events in adolescent depression. TINS. 2008;31:183–191.

50. Chai XJ, Whitfield-Gabrieli S, Shinn AK, Gabrieli JDE, Nieto Castãón A, McCarthy JM, et al. Abnormal medial prefrontal cortex resting-state connectivity in bipolar disorder and schizophrenia. Neuropsychopharmacology. 2011;36:2009–2017.

51. Blumberg HP, Stem E, Martinez D, Ricketts S, de Asis J, White T, et al. Increased anterior cingulate and caudate activity in bipolar mania. Bipolar Disord Sci Ment Heal. 2019:177–184.

52. Kenwood MM, Kalin NH, Barbas H. The prefrontal cortex, pathological anxiety, and anxiety disorders. Neuropsychopharmacology. 2022;47:260–275.

53. Shah AA, Sjovold T, Treit D. Inactivation of the medial prefrontal cortex with the GABAA receptor agonist muscimol increases open-arm activity in the elevated plus-maze and attenuates shock-probe burying in rats. Brain Res. 2004;1028:112–115.

54. Bi LL, Wang J, Luo ZY, Chen SP, Geng F, Chen YH, et al. Enhanced excitability in the infralimbic cortex produces anxiety-like behaviors. Neuropharmacology. 2013;72:148–156.

55. Saitoh A, Ohashi M, Suzuki S, Tsukagoshi M, Sugiyama A, Yamada M, et al. Activation of the prelimbic medial prefrontal cortex induces anxiety-like behaviors via N-Methyl-D-aspartate receptor-mediated glutamatergic neurotransmission in mice. J Neurosci Res. 2014;92:1044–1053.

56. Ferenczi E a., Zalocusky K a., Liston C, Grosenick L, Warden MR, Amatya D, et al. Prefrontal cortical regulation of brainwide circuit dynamics and reward-related behavior. Science (80-). 2016;351:aac9698–aac9698.

57. Fuchikami M, Thomas A, Liu R, Wohleb ES, Land BB, DiLeone RJ, et al. Optogenetic stimulation of infralimbic PFC reproduces ketamine's rapid and sustained antidepressant actions. Proc Natl

1102   Acad Sci U S A. 2015;112:8106–8111.

58.   Hare BD, Shinohara R, Liu RJ, Pothula S, DiLeone RJ, Duman RS. Optogenetic stimulation of medial prefrontal cortex Drd1 neurons produces rapid and long-lasting antidepressant effects. Nat Commun. 2019;10:1–12.

59.   Slattery DA, Neumann ID, Cryan JF. Transient inactivation of the infralimbic cortex induces antidepressant-like effects in the rat. J Psychopharmacol. 2011;25:1295–1303.

60.   Chen YH, Wu JL, Hu NY, Zhuang JP, Li WP, Zhang SR, et al. Distinct projections from the infralimbic cortex exert opposing effects in modulating anxiety and fear. J Clin Invest. 2021;131.

61.   Vidal-Gonzalez I, Vidal-Gonzalez B, Rauch SL, Quirk GJ. Microstimulation reveals opposing influences of prelimbic and infralimbic cortex on the expression of conditioned fear. Learn Mem. 2006;13:728–733.

62.   Sierra-Mercado D, Padilla-Coreano N, Quirk GJ. Dissociable roles of prelimbic and infralimbic cortices, ventral hippocampus, and basolateral amygdala in the expression and extinction of conditioned fear. Neuropsychopharmacology. 2011;36:529–538.

63.   Arruda-Carvalho M, Clem RL. Prefrontal-amygdala fear networks come into focus. Front Syst Neurosci. 2015;9:145.

64.   Arruda-Carvalho M, Clem RL. Pathway-selective adjustment of prefrontal-amygdala transmission during fear encoding. J Neurosci. 2014;34:15601–15609.

65.   Quirk GJ, Russo GK, Barron JL, Lebron K. The role of ventromedial prefrontal cortex in the recovery of extinguished fear. J Neurosci. 2000;20:6225–6231.

66.   Sotres-Bayon F, Sierra-Mercado D, Pardilla-Delgado E, Quirk GJ. Gating of fear in prelimbic cortex by hippocampal and amygdala inputs. Neuron. 2012;76:804–812.

67.   Akhter F, Haque T, Sato F, Kato T, Ohara H, Fujio T, et al. Projections from the dorsal peduncular cortex to the trigeminal subnucleus caudalis (medullary dorsal horn) and other lower brainstem areas in rats. Neuroscience. 2014;266:23–37.

68.   Kataoka N, Shima Y, Nakajima K, Nakamura K. A central master driver of psychosocial stress responses in the rat. Science (80-). 2020;367:1105–1112.

69.   Nakamura K, Morrison SF. Central sympathetic network for thermoregulatory responses to psychological stress. Aut Neurosci. 2022;237:102918.

70.   Nakamura K, Nakamura Y, Kataoka N. A hypothalamomedullary network for physiological responses to environmental stresses. Nat Rev Neurosci. 2022;23:35–52.

71.   Diorio D, Viau V, Meaney MJ. The role of the medial prefrontal cortex (cingulate gyrus) in the regulation of hypothalamic-pituitary-adrenal responses to stress. J Neurosci. 1993;13:3839–3847.

72.   Radley JJ, Arias CM, Sawchenko PE. Regional differentiation of the medial prefrontal cortex in regulating adaptive responses to acute emotional stress. J Neurosci. 2006;26:12967–12976.

73.   Tavares RF, Correa FM. Role of the medial prefrontal cortex in cardiovascular responses to acute restraint in rats. Neuroscience. 2006;143:231–240.

74.   Radley JJ, Gosselink KL, Sawchenko PE. A discrete GABAergic relay mediates medial prefrontal cortical inhibition of the neuroendocrine stress response. J Neurosci. 2009;29:7330–7340.

75.   Olff M, Guzelcan Y, de Vries GJ, Assies J, Gersons BP. HPA- and HPT-axis alterations in chronic posttraumatic stress disorder. Psychoneuroendocrinology. 2006;31:1220–1230.

76.   Park J, Marvar PJ, Liao P, Kankam ML, Norrholm SD, Downey RM, et al. Baroreflex dysfunction

and augmented sympathetic nerve responses during mental stress in veterans with post-traumatic stress disorder. J Physiol. 2017;595:4893–4908.

77. Fonkoue IT, Marvar PJ, Norrholm S, Li Y, Kankam ML, Jones TN, et al. Symptom severity impacts sympathetic dysregulation and inflammation in post-traumatic stress disorder (PTSD). Brain Behav Immun. 2020;83:260–269.

78. Teed AR, Feinstein JS, Puhl M, Lapidus RC, Upshaw V, Kuplicki RT, et al. Association of Generalized Anxiety Disorder With Autonomic Hypersensitivity and Blunted Ventromedial Prefrontal Cortex Activity During Peripheral Adrenergic Stimulation: A Randomized Clinical Trial. JAMA Psychiatry. 2022;79:323–332.

79. Sullivan RM, Gratton A. Prefrontal cortical regulation of hypothalamic-pituitary-adrenal function in the rat and implications for psychopathology: side matters. Psychoneuroendocrinology. 2002;27:99–114.

80. Jaferi A, Bhatnagar S. Corticotropin-releasing hormone receptors in the medial prefrontal cortex regulate hypothalamic-pituitary-adrenal activity and anxiety-related behavior regardless of prior stress experience. Brain Res. 2007;1186:212–223.

81. Weinberg MS, Johnson DC, Bhatt AP, Spencer RL. Medial prefrontal cortex activity can disrupt the expression of stress response habituation. Neuroscience. 2010;168:744–756.

82. DeNardo LA, Liu CD, Allen WE, Adams EL, Friedmann D, Fu L, et al. Temporal evolution of cortical ensembles promoting remote memory retrieval. Nat Neurosci. 2019;22:460–469.

83. Allen WE, DeNardo LA, Chen MZ, Liu CD, Loh KM, Fenno LE, et al. Thirst-associated preoptic neurons encode an aversive motivational drive. Science (80-). 2017;357:1149–1155.

84. Hurley KM, Herbert H, Moga MM, Saper CB. Efferent projections of the infralimbic cortex of the rat. J Comp Neurol. 1991;308:249–276.

85. Fisk GD, Wyss JM. Descending projections of infralimbic cortex that mediate stimulation-evoked changes in arterial pressure. Brain Res. 2000;859:83–95.

86. Wang GQ, Cen C, Li C, Cao S, Wang N, Zhou Z, et al. Deactivation of excitatory neurons in the prelimbic cortex via Cdk5 promotes pain sensation and anxiety. Nat Commun. 2015;6.

87. Pizzagalli DA, Roberts AC. Prefrontal cortex and depression. Neuropsychopharmacology. 2022;47:225–246.

88. Challis C, Beck SG, Berton O. Optogenetic modulation of descending prefrontocortical inputs to the dorsal raphe bidirectionally bias socioaffective choices after social defeat. Front Behav Neurosci. 2014;8:1–14.

89. Miller OH, Bruns A, Ben Ammar I, Mueggler T, Hall BJ. Synaptic Regulation of a Thalamocortical Circuit Controls Depression-Related Behavior. Cell Rep. 2017;20:1867–1880.

90. Hamani C, MacHado DC, Hipólide DC, Dubiela FP, Suchecki D, MacEdo CE, et al. Deep brain stimulation reverses anhedonic-like behavior in a chronic model of depression: Role of serotonin and brain derived neurotrophic factor. Biol Psychiatry. 2012;71:30–35.

91. Kumar S, Black SJ, Hultman R, Szabo ST, DeMaio KD, Du J, et al. Cortical control of affective networks. J Neurosci. 2013;33:1116–1129.

92. Vialou V, Bagot RC, Cahill ME, Ferguson D, Robison AJ, Dietz DM, et al. Prefrontal Cortical Circuit for Depression- and Anxiety- Related Behaviors Mediated by Cholecystokinin : Role of □ FosB. J Neurosci. 2014;34:3878–3887.

93. Bagot RC, Parise EM, Peña CJ, Zhang H-X, Maze I, Chaudhury D, et al. Ventral hippocampal

afferents to the nucleus accumbens regulate susceptibility to depression. Nat Commun. 2015;6:7062.

94. Covington HE, Lobo MK, Maze I, Vialou V, Hyman JM, Zaman S, et al. Antidepressant effect of optogenetic stimulation of the medial prefrontal cortex. J Neurosci. 2010;30:16082–16090.

95. Perova Z, Delevich K, Li B. Depression of excitatory synapses onto parvalbumin interneurons in the medial prefrontal cortex in susceptibility to stress. J Neurosci. 2015;35:3201–3206.

96. Warden MR, Selimbeyoglu A, Mirzabekov JJ, Lo M, Thompson KR, Kim SY, et al. A prefrontal cortex-brainstem neuronal projection that controls response to behavioural challenge. Nature. 2012;492:428–432.

97. Ferguson BR, Gao WJ. Thalamic Control of Cognition and Social Behavior Via Regulation of Gamma-Aminobutyric Acidergic Signaling and Excitation/Inhibition Balance in the Medial Prefrontal Cortex. Biol Psychiatry. 2018;83:657–669.

98. Ouhaz Z, Ba-M'hamed S, Mitchell AS, Elidrissi A, Bennis M. Behavioral and cognitive changes after early postnatal lesions of the rat mediodorsal thalamus. Behav Brain Res. 2015;292:219–232.

99. Ouhaz Z, Ba-M'hamed S, Bennis M. Morphological, structural, and functional alterations of the prefrontal cortex and the basolateral amygdala after early lesion of the rat mediodorsal thalamus. Brain Struct Funct. 2017;222:2527–2545.

100. Beracochea D, Krazem A. Effects of mammillary body and mediodorsal thalamic lesions on elevated plus maze exploration. Neuroreport. 1991;2:793–796.

101. Li Y, Li S, Wei C, Wang H, Sui N, Kirouac GJ. Orexins in the paraventricular nucleus of the thalamus mediate anxiety-like responses in rats. Psychopharmacology (Berl). 2010;212:251–265.

102. Kirouac GJ. Placing the paraventricular nucleus of the thalamus within the brain circuits that control behavior. Neurosci Biobehav Rev. 2015;56:315–329.

103. Kooiker CL, Birnie MT, Baram TZ. The Paraventricular Thalamus: A Potential Sensor and Integrator of Emotionally Salient Early-Life Experiences. Front Behav Neurosci. 2021;15:1–9.

104. Barson JR, Leibowitz SF. GABA-induced inactivation of dorsal midline thalamic subregions has distinct effects on emotional behaviors. Neurosci Lett. 2015;609:92–96.

105. Zhao D, Wang D, Wang W, Dai J, Cui M, Wu M, et al. The altered sensitivity of acute stress induced anxiety-related behaviors by modulating insular cortex-paraventricular thalamus-bed nucleus of the stria terminalis neural circuit. Neurobiol Dis. 2022;174:105890.

106. Shin LM, Liberzon I. The neurocircuitry of fear, stress, and anxiety disorders. Neuropsychopharmacology. 2010;35:169–191.

107. Shi T, Feng S, Wei M, Zhou W. Role of the anterior agranular insular cortex in the modulation of fear and anxiety. Brain Res Bull. 2020;155:174–183.

108. Shi TY, Feng SF, Wei MX, Huang Y, Liu G, Wu HT, et al. Kainate receptor mediated presynaptic LTP in agranular insular cortex contributes to fear and anxiety in mice. Neuropharmacology. 2018;128:388–400.

109. Li H, Chen L, Li P, Wang X, Zhai H. Insular muscarinic signaling regulates anxiety-like behaviors in rats on the elevated plus-maze. Behav Brain Res. 2014;270:256–260.

110. Méndez-Ruette M, Linsambarth S, Moraga-Amaro R, Quintana-Donoso D, Méndez L, Tamburini G, et al. The role of the rodent insula in anxiety. Front Physiol. 2019;10:1–10.

111. Linley SB, Athanason AC, Rojas AKP, Vertes RP. Role of the reuniens and rhomboid thalamic

1230   nuclei in anxiety-like avoidance behavior in the rat. Hippocampus. 2021;31:756–769.

112.   Felix-Ortiz AC, Tye KM. Amygdala inputs to the ventral hippocampus bidirectionally modulate social behavior. J Neurosci. 2014;34:586–595.

113.   Felix-Ortiz AC, Beyeler A, Seo C, Leppla CA, Wildes CP, Tye KM. BLA to vHPC Inputs Modulate Anxiety-Related Behaviors. Neuron. 2013;79:658–664.

114.   Tye KM, Prakash R, Kim S-Y, Fenno LE, Grosenick L, Zarabi H, et al. Amygdala circuitry mediating reversible and bidirectional control of anxiety. Nature. 2011;471:358–362.

115.   Adhikari A, Lerner TN, Finkelstein J, Pak S, Jennings JH, Davidson TJ, et al. Basomedial amygdala mediates top-down control of anxiety and fear. Nature. 2015;527:179–185.

116.   Daviu N, Bruchas MR, Moghaddam B, Sandi C, Beyeler A. Neurobiological links between stress and anxiety. Neurobiol Stress. 2019;11:100191.

117.   Do-Monte FH, Manzano-Nieves G, Quiñones-Laracuente XK, Ramos-Medina L, Quirk GJ. Revisiting the Role of Infralimbic Cortex in Fear Extinction with Optogenetics. J Neurosci. 2015;35:3607–3615.

118.   Thompson BM, Baratta M V., Biedenkapp JC, Rudy JW, Watkins LR, Maier SF. Activation of the infralimbic cortex in a fear context enhances extinction learning. Learn Mem. 2010;17:591–599.

119.   Milad MR, Vidal-Gonzalez I, Quirk GJ. Electrical stimulation of medial prefrontal cortex reduces conditioned fear in a temporally specific manner. Behav Neurosci. 2004;118:389–394.

120.   Milad MR, Quirk GJ. Neurons in medial prefrontal cortex signal memory for fear extinction. 2002;420:713–717.

121.   Kim HS, Cho HY, Augustine GJ, Han JH. Selective Control of Fear Expression by Optogenetic Manipulation of Infralimbic Cortex after Extinction. Neuropsychopharmacology. 2016;41:1261–1273.

122.   Campos-Cardoso R, Desa Z, Fitzgerald B, Moore A, Duhon J, Landar V, et al. The mouse dorsal peduncular cortex encodes fear memory. Biorxiv. 2023;https://do.

123.   Sturman O, Germain PL, Bohacek J. Exploratory rearing: a context- and stress-sensitive behavior recorded in the open-field test. Stress. 2018;21:443–452.

124.   Borkar C, Stelly C, Fu X, Dorofeikova M, Le Q-S, Vutukuri R, et al. Top-down control of flight by a non-canonical cortico-amygdala pathway. BiorXiv. 2023;https://do.

125.   Figueiredo HF, Bruestle A, Bodie B, Dolgas CM, Herman JP. The medial prefrontal cortex differentially regulates stress-induced c-fos expression in the forebrain depending on type of stressor. Eur J Neurosci. 2003;18:2357–2364.

126.   Van Eden CG, Buijs RM. Functional neuroanatomy of the prefrontal cortex: Autonomic interactions. Prog Brain Res. 2000;126:49–62.

127.   Brake WG, Flores G, Francis D, Meaney MJ, Srivastava LK, Gratton A. Enhanced nucleus accumbens dopamine and plasma corticosterone stress responses in adult rats with neonatal excitotoxic lesions to the medial prefrontal cortex. Neuroscience. 2000;96:687–695.

128.   Crane JW, Ebner K, Day TA. Medial prefrontal cortex suppression of the hypothalamic-pituitary-adrenal axis response to a physical stressor, systemic delivery of interleukin-1β. Eur J Neurosci. 2003;17:1473–1481.

129.   Sullivan RM, Gratton A. Lateralized effects of medial prefrontal cortex lesions on neuroendocrine and autonomic stress responses in rats. J Neurosci. 1999;19:2834–2840.

130. Feldman S, Conforti N. Modifications of adrenocortical responses following frontal cortex simulation in rats with hypothalamic deafferentations and medial forebrain bundle lesions. Neuroscience. 1985;15:1045–1047.

131. Buijs RM, Van Eden CG. The integration of stress by the hypothalamus, amygdala and prefrontal cortex: Balance between the autonomic nervous system and the neuroendocrine system. Prog Brain Res. 2000;126:117–132.

132. Choi DC, Furay AR, Evanson NK, Ulrich-Lai YM, Nguyen MMN, Ostrander MM, et al. The role of the posterior medial bed nucleus of the stria terminalis in modulating hypothalamic-pituitary-adrenocortical axis responsiveness to acute and chronic stress. Psychoneuroendocrinology. 2008;33:659–669.

133. Spencer SJ, Buller KM, Day TA. Medial prefrontal cortex control of the paraventricular hypothalamic nucleus response to psychological stress: Possible role of the bed nucleus of the stria terminalis. J Comp Neurol. 2005;481:363–376.

134. Bertholomey ML, Nagarajan V, Smith DM, Torregrossa MM. Sex- and age-dependent effects of chronic corticosterone exposure on depressive-like, anxiety-like, and fear-related behavior: Role of amygdala glutamate receptors in the rat. Front Behav Neurosci. 2022;16:1–18.

135. Nandi A, Virmani G, Barve A, Marathe S. DBscorer: An open-source software for automated accurate analysis of rodent behavior in forced swim test and tail suspension test. ENeuro. 2021;8.

136. Botterill JJ, Khlaifia A, Walters BJ, Brimble MA, Scharfman HE, Arruda-Carvalho M. Off-Target Expression of Cre-Dependent Adeno- Associated Viruses in Wild-Type C57BL/6J Mice. ENeuro. 2021;8:1–16.

137. Samifanni R, Zhao M, Cruz-Sanchez A, Satheesh A, Mumtaz U, Arruda-Carvalho M. Developmental emergence of persistent memory for contextual and auditory fear in mice. Learn Mem. 2021;28:414–421.

138. Arruda-Carvalho M, Wu W, Cummings KA, Clem RL. Optogenetic examination of prefrontal-amygdala synaptic development. J Neurosci. 2017. 2017. https://doi.org/10.1523/JNEUROSCI.3097-16.2017.

139. Seibenhener ML, Wooten MC. Use of the open field maze to measure locomotor and anxiety-like behavior in mice. J Vis Exp. 2015:e52434.

140. Ueno H, Takahashi Y, Suemitsu S, Murakami S, Kitamura N, Wani K, et al. Effects of repetitive gentle handling of male C57BL/6NCrl mice on comparative behavioural test results. Sci Rep. 2020;10:3509.

141. Botterill JJ, Vinod KY, Gerencer KJ, Teixeira CM, LaFrancois JJ, Scharfman HE. Bidirectional regulation of cognitive and anxiety-like behaviors by dentate gyrus mossy cells in male and female mice. J Neurosci. 2021;41:2475–2495.

142. Komada M, Takao K, Miyakawa T. Elevated plus maze for mice. J Vis Exp. 2008. 2008. https://doi.org/10.3791/1088.

143. Steru L, Chermat R, Thierry B, Simon P. The tail suspension test: a new method for screening antidepressants in mice. Psychopharmacol. 1985;85:367–370.

144. Ueno H, Takahashi Y, Murakami S, Wani K, Matsumoto Y, Okamoto M, et al. Effect of simultaneous testing of two mice in the tail suspension test and forced swim test. Sci Rep. 2022;12:9224.

145. Arruda-Carvalho M, Sakaguchi M, Akers KG, Josselyn S a, Frankland PW. Posttraining ablation of adult-generated neurons degrades previously acquired memories. J Neurosci. 2011;31:15113–15127.

1317   146.   Porsolt RD, Bertin A, Jalfre M. Behavioral despair in mice: a primary screening test for
1318   antidepressants. Arch Int Pharmacodyn Ther. 1977;229:327–336.

1319   147.   Can A, Dao DT, Arad M, Terrillion CE, Piantadosi SC, Gould TD. The mouse forced swim test. J
1320   Vis Exp. 2012:e3638.

1321   148.   Kara NZ, Stukalin Y, Einat H. Revisiting the validity of the mouse forced swim test: Systematic
1322   review and meta-analysis of the effects of prototypic antidepressants. Neurosci Biobehav Rev.
1323   2018;84:1–11.

1324   149.   Can A, Dao DT, Arad M, Terrillion CE, Piantadosi SC, Gould TD. The mouse forced swim test. J
1325   Vis Exp. 2011:1–6.

1326   150.   Thompson KJ, Khajehali E, Bradley SJ, Navarrete JS, Huang XP, Slocum S, et al. DREADD
1327   Agonist 21 Is an Effective Agonist for Muscarinic-Based DREADDs in Vitro and in Vivo. ACS
1328   Pharmacol Transl Sci. 2018;1:61–72.

1329   151.   Archer J. Rodent Sex Differences in Emotional and Related Behavior I Ethology and
1330   Neurophysiology Group , School of Biological Sciences , University of Sussex ,. Neurophysiology.
1331   1975;479:451–479.

1332   152.   Johnston AL, File SE. Sex differences in animal tests of anxiety. Physiol Behav. 1991;49:245–250.

1333   153.   Knight P, Chellian R, Wilson R, Behnood-Rod A, Panunzio S, Bruijnzeel AW. Sex differences in
1334   the elevated plus-maze test and large open field test in adult Wistar rats. Pharmacol Biochem
1335   Behav. 2021;204:173168.

1336   154.   Börchers S, Krieger JP, Asker M, Maric I, Skibicka KP. Commonly-used rodent tests of anxiety-
1337   like behavior lack predictive validity for human sex differences. Psychoneuroendocrinology.
1338   2022;141.

1339   155.   Scholl JL, Afzal A, Fox LC, Watt MJ, Forster GL. Sex differences in anxiety-like behaviors in rats.
1340   Physiol Behav. 2019;211:112670.

1341   156.   Simpson J, Kelly JP. An investigation of whether there are sex differences in certain behavioural
1342   and neurochemical parameters in the rat. Behav Brain Res. 2012;229:289–300.

1343   157.   Ramos A, Kangerski AL, Basso PF, Da Silva Santos JE, Assreuy J, Vendruscolo LF, et al.
1344   Evaluation of Lewis and SHR rat strains as a genetic model for the study of anxiety and pain.
1345   Behav Brain Res. 2002;129:113–123.

1346   158.   Bishnoi IR, Ossenkopp KP, Kavaliers M. Sex and age differences in locomotor and anxiety-like
1347   behaviors in rats: From adolescence to adulthood. Dev Psychobiol. 2021;63:496–511.

1348   159.   Keeley RJ, Bye C, Trow J, McDonald RJ. Strain and sex differences in brain and behaviour of
1349   adult rats: Learning and memory, anxiety and volumetric estimates. Behav Brain Res.
1350   2015;288:118–131.

1351   160.   Baumgartner NE, Biraud MC, Lucas EK. Sex differences in socioemotional behavior and changes
1352   in ventral hippocampal transcription across aging in C57Bl/6J mice. Neurobiol Aging.
1353   2023;130:141–153.

1354   161.   Painsipp E, Wultsch T, Shahbazian A, Edelsbrunner M, Kreissl MC, Schirbel A, et al. Experimental
1355   gastritis in mice enhances anxiety in a gender-related manner. Neuroscience. 2007;150:522–536.

1356   162.   Tucker LB, Burke JF, Fu AH, McCabe JT. Neuropsychiatric Symptom Modeling in Male and
1357   Female C57BL/6J Mice after Experimental Traumatic Brain Injury. J Neurotrauma. 2017;34:890–
1358   905.

1359   163.   Hendershott TR, Cronin ME, Langella S, McGuinness PS, Basu AC. Effects of environmental

enrichment on anxiety-like behavior, sociability, sensory gating, and spatial learning in male and female C57BL/6J mice. Behav Brain Res. 2016;314:215–225.

164. Rodgers RJ, Cole JC. Influence of social isolation, gender, strain, and prior novelty on plus-maze behaviour in mice. Physiol Behav. 1993;54:729–736.

165. Tucker LB, McCabe JT. Behavior of male and female C57Bl/6J mice is more consistent with repeated trials in the elevated zero maze than in the elevated plus maze. Front Behav Neurosci. 2017;11:1–8.

166. Võikar V, Kõks S, Vasar E, Rauvala H. Strain and gender differences in the behavior of mouse lines commonly used in transgenic studies. Physiol Behav. 2001;72:271–281.

167. Kageyama R, Ohtsuka T, Shimojo H, Imayoshi I. Dynamic Notch signaling in neural progenitor cells and a revised view of lateral inhibition. Nat Neurosci. 2008;11:1247–1251.

168. An XL, Zou JX, Wu RY, Yang Y, Tai FD, Zeng SY, et al. Strain and sex differences in anxiety-like and social behaviors in C57BL/6J and BALB/cJ mice. Exp Anim. 2011;60:111–123.

169. Bolivar VJ, Caldarone BJ, Reilly AA, Flaherty L. Habituation of activity in an open field: A survey of inbred strains and F 1 hybrids. Behav Genet. 2000;30:285–293.

170. Imhof JT, Coelho ZMI, Schmitt ML, Morato GS, Carobrez AP. Influence of gender and age on performance of rats in the elevated plus maze apparatus. Behav Brain Res. 1993;56:177–180.

171. Estanislau C, Morato S. Behavior ontogeny in the elevated plus-maze: prenatal stress effects. Int J Dev Neurosci. 2006;24:255–262.

172. Walf AA, Koonce C, Manley K, Frye CA. Proestrous compared to diestrous wildtype , but not estrogen receptor beta knockout , mice have better performance in the spontaneous alternation and object recognition tasks and reduced anxiety-like behavior in the elevated plus and mirror maze. 2009;196:254–260.

173. Frye CA, Petralia SM, Rhodes ME. Estrous cycle and sex differences in performance on anxiety tasks coincide with increases in hippocampal progesterone and 3α,5α-THP. Pharmacol Biochem Behav. 2000. 2000. https://doi.org/10.1016/S0091-3057(00)00392-0.

174. du Plessis KC, Basu S, Rumbell TH, Lucas EK. Sex-Specific Neural Networks of Cued Threat Conditioning: A Pilot Study. Front Syst Neurosci. 2022;16:1–11.

175. McLean CP, Asnaani A, Litz BT, Hofmann SG. Gender differences in anxiety disorders: Prevalence, course of illness, comorbidity and burden of illness. J Psychiatr Res. 2011;45:1027–1035.

176. Kessler RC, Sonnega A, Nelson CB, Bromet E. Posttraumatic Stress Disorder in the National Comorbidity Survey. Arch Gen Psychiatry. 1995;52.

177. Bangasser DA, Cuarenta A. Sex differences in anxiety and depression: circuits and mechanisms. Nat Rev Neurosci. 2021;22:674–684.

178. Dart AM, Du XJ, Kingwell BA. Gender, sex hormones and autonomic nervous control of the cardiovascular system. Cardiovasc Res. 2002;53:678–687.

179. Hinojosa-Laborde C, Chapa I, Lange D, Haywood J. Gender differences in sympathetic nervous system regulation. Clin Exp Pharmacol Physiol. 1999;26:122–126.

180. Barsha G, Denton KM, Mirabito Colafella KM. Sex- and age-related differences in arterial pressure and albuminuria in mice. Biol Sex Differ. 2016;7:1–15.

181. Colafella KMM, Denton KM. Sex-specific differences in hypertension and associated cardiovascular disease. Nat Rev Nephrol. 2018;14:185–201.