1

3        The *Drosophila drop-dead* gene is required for eggshell integrity

5              Short title: *drop-dead* and the *Drosophila* eggshell

Tayler D. Sheahan[1,#a], Amanpreet Grewal[1], Laura E. Korthauer[1,#b], Edward M.

Blumenthal[1*]

[1] Department of Biological Sciences, Marquette University, Milwaukee, Wisconsin,

United States of America

[#a]Current Address: Pittsburgh Center for Pain Research and Department of

Neurobiology, University of Pittsburgh, Pittsburgh, Pennsylvania, United States of

America

[#b]Current Address: Department of Psychiatry and Human Behavior, Alpert Medical

School of Brown University, Providence, Rhode Island, United States of America

*Corresponding author

E-mail: edward.blumenthal@marquette.edu (EMB)

## Abstract

The eggshell of the fruit fly *Drosophila melanogaster* is a useful model for understanding the synthesis of a complex extracellular matrix. The eggshell is synthesized during mid-to-late oogenesis by the somatic follicle cells that surround the developing oocyte. We previously reported that female flies mutant for the gene *drop-dead* (*drd*) are sterile, but the underlying cause of the sterility remained unknown. In this study, we examined the role of *drd* in eggshell synthesis. We show that eggs laid by *drd* mutant females are fertilized but arrest early in embryogenesis, and that the innermost layer of the eggshell, the vitelline membrane, is abnormally permeable to dye in these eggs. In addition, the major vitelline membrane proteins fail to become crosslinked by nonreducible bonds, a process that normally occurs during egg activation following ovulation, as evidenced by their solubility and detection by Western blot in laid eggs. In contrast, the Cp36 protein, which is found in the outer chorion layers of the eggshell, becomes crosslinked normally. To link the *drd* expression pattern with these phenotypes, we show that *drd* is expressed in the ovarian follicle cells beginning in mid-oogenesis, and, importantly, that all *drd* mutant eggshell phenotypes could be recapitulated by selective knockdown of *drd* expression in the follicle cells. To determine whether *drd* expression was required for the crosslinking itself, we performed *in vitro* activation and crosslinking experiments. The vitelline membranes of control egg chambers could become crosslinked either by incubation in hyperosmotic medium, which activates the egg chambers, or by exogenous peroxidase and hydrogen peroxide. In contrast, neither treatment resulted in the crosslinking of the vitelline membrane in *drd* mutant egg chambers. These results indicate that *drd* expression in the follicle cells is necessary for vitelline

46    membrane proteins to serve as substrates for peroxidase-mediated cross-linking at

47    the end of oogenesis.

## Introduction

Animal epithelial cells produce an extracellular matrix (ECM) that must perform many roles, including as a structural support, barrier, and source of signaling molecules [1–3]. The eggshell of the insect *Drosophila melanogaster* is a model ECM consisting of five layers of protein, lipid, and carbohydrate [4]. Among its functions, the *Drosophila* eggshell serves as physical protection and a selective permeability barrier, provides patterning signals for the oocyte and developing embryo, and binds pheromones that prevent cannibalism by conspecific larvae [5–7]. Eggshell components are primarily synthesized by the follicle cells, a layer of somatic epithelial cells that surround the germline nurse cells and oocyte; together these three cell types make up the basic unit of oogenesis, the egg chamber.

The innermost layer of the eggshell is the proteinaceous vitelline membrane (VM). It is composed of six related structural proteins encoded by the genes *Vm26Aa*, *Vm26Ab*, *Vm26Ac*, *Vm32E*, *Vm34Ca*, and *Vml*, as well as other less abundant proteins [4,8–12]. While most VM components are produced by the follicle cells, at least three proteins, encoded by *fs(1)Nasrat* (*fs(1)N*), *fs(1)polehole* (*fs(1)ph*) and *closca* (*clos*), are secreted by the oocyte and become incorporated into the developing VM [13–15]. VM components are synthesized during mid-oogenesis (stages 8-11 of the 14 stages of oogenesis), followed by chorion components in stages 11-12 [4,16,17].

Following their synthesis and secretion, the proteins of the VM become cross-linked, forming a stable and insoluble matrix. The VM proteins are cross-linked to each other by disulfide bonds during the early stages of eggshell formation [11,18]. Immediately following ovulation and egg activation, VM proteins become cross-linked by non-reducible bonds, at least some of which are dityrosine bonds [19]. The non-

reducible cross-linking of the VM occurs in a matter of minutes as the egg moves

down the oviduct; soluble VM proteins are never detected in freshly laid eggs

[20,21]. While the formation of dityrosine bonds is typically catalyzed by a peroxidase

[22–25], the enzyme responsible for VM crosslinking has not been identified.

The structural integrity of the *Drosophila* VM can be disrupted by mutations in

several genes. Mutation of many of the genes encoding VM structural proteins

causes gross VM abnormalities and collapse of the eggs [26–28], as do mutations in

the cadherin *Cad99C* [29,30], which is localized to microvilli on the apical surface of

the follicle cells, and the eggshell components *yellow-g* and *yellow-g2* [8,31]. Other

mutations, in the genes encoding the minor eggshell components Nudel, Palisade

(Psd), Clos, Fs(1)ph, and Fs(1)N, result in a disruption in VM protein cross-linking

without altering overall VM integrity to the extent of causing eggs to collapse

[13,14,20,32,33], however all of these mutations result in female sterility.

In this paper, we studied the role of the *drop-dead* (*drd*) gene in oogenesis.

*drd* encodes a putative integral membrane protein of unknown function with

homology to prokaryotic acyltransferases [34]. Mutation of *drd* causes a wide range

of phenotypes, including female sterility, early adult death and neurodegeneration,

defective food movement through the gut, and absence of a peritrophic matrix from

the midgut [35–39]. The basis for female sterility has not previously been reported.

Here we demonstrate that *drd* expression in the follicle cells is required for non-

reducible cross-linking of the VM.

# Materials and methods

## *Drosophila* stocks and maintenance

All fly stocks were maintained on standard cornmeal-yeast-agar food
(http://flystocks.bio.indiana.edu/Fly_Work/media-recipes/molassesfood.htm) at 25°C
on a 12h:12h light-dark cycle. For RNAi experiments, a *UAS-Dcr-2* transgene was
included in the genetic background of the flies in order to boost RNAi efficiency; the
*drd$^{GD15915}$ UAS-Dcr-2* and *UAS-Dcr-2 drd$^{GD3367}$* lines were created previously by
recombination between VDRC stocks *w$^{1118}$;P{GD3367}v37404* (FBst0461992) and
*w$^{1118}$; P{GD15915}v51184* (FBst0469325) and Bloomington stock *w$^{1118}$; P{UAS-Dcr-
2.D}2* (FBst0024650, RRID:BDSC_24650) [40,41]. The *w*;*
*P{w$^{+mW.hs}$GAL4=GawB}CY2* stock (FBti0007266, referred to as *CY2-GAL4*) was
provided by Dr. Celeste Berg. Other stocks (*w$^{1118}$; P{UAS-GFP.nls}14*
(FBst0004775, RRID:BDSC_4775), *P{w$^{+mW.hs}$=GawB}T155* (FBst0005076,
RRID:BDSC_5076, referred to as *T155-GAL4*), *w* ovo$^{D1}$ v$^{24}$*
*P{w$^{+mW.hs}$=FRT(w$^{hs}$)}101/C(1)DX, y$^1$ f$^1$; P{ry$^{+t7.2}$=hsFLP}38* (FBst0001813,
RRID:BDSC_1813), and *y$^1$ w* v$^{24}$ P{w$^{+mW.hs}$=FRT(w$^{hs}$)}101* (FBst0001844,
RRID:BDSC_1844)), were obtained from the Bloomington *Drosophila* Stock Center.
Creation of the *drd-GAL4* driver transgene has been reported previously [36]. The
genes and alleles referenced in this work include *drd* (FBgn0260006), *drd$^{lwf}$*
(FBal0193421), *drd$^1$* (FBal0003113). Stocks were not outcrossed prior to this study.

## *drd$^1$* sequencing

Whole-fly RNA was prepared from Canton S and *drd$^1$* homozygous adults
using Trizol reagent (ThermoFisher Scientific, Waltham, MA). RNA was treated with

118  DNase (ThermoFisher Scientific, Waltham, MA) and cDNA was synthesized (qScript

119  cDNA supermix, Quantabio, Beverly, MA). Primers for amplification of the exon 8/9

120  junction were: CG5652 6a 5' GAT CGC TG GTG TTT GTT TT 3' and CG5652 6b

121  5' TTC GCT GGG GAT CAC TAA AC 3'.

## Egg-laying assay

Groups of 1-3 homozygous $drd^1/drd^1$ or $drd^{lwf}/drd^{lwf}$ females were mated with

Canton S males and placed on either regular food or food supplemented with yeast

paste. Flies were transferred to new vials daily until they died, and the number of

eggs laid was recorded. Because groups of three flies were assayed together in

early experiments, we analyzed the data twice—once assuming that all eggs were

laid by a single fly (model 1) and once assuming that egg-laying was distributed

evenly among all flies in a vial (model 2). The conclusions about the proportion of

flies that laid eggs were the same in both analyses.

## Analysis of embryogenesis

Females were mated with sibling males, and eggs were collected overnight

(16.5-18.5 hr) on apple juice agar plates supplemented with yeast paste [42]. Flies

were removed and the eggs were allowed to develop for an additional 2-6 hr. Eggs

were then covered in halocarbon 700 oil (Sigma-Aldrich, St. Louis, MO) and scored

for collapsed vs turgid. Turgid eggs were scored for fertilized vs unfertilized, and

fertilized eggs were scored for pregastrulation vs postgastrulation [42].

## Generation of *drd* germline clones

Germline mitotic clones mutant for *drd* were generated using the FLP/FRT-

dominant female sterile technique as described [43]. The *drd$^{lwf}$* allele was first

recombined onto the same chromosome as an FRT site by crossing *w drd$^{lwf}$* x *y w v*

*P{FRT}101*. Following the establishment of a stock carrying this recombinant

chromosome, *w drd$^{lwf}$ P{FRT}101/FM7a* females were crossed with *w ovo$^{D1}$ v*

*P{FRT}101; hs-FLP* males. The resulting *w drd$^{lwf}$ P{FRT}101/w ovo$^{D1}$ v P{FRT}101;*

*hs-FLP/+* embryos were heat shocked for 2 hr at 37°C to induce FLPase expression

and mitotic recombination, raised to adulthood, and crossed with Canton S males to

assay for fertility. Control embryos of the same genotype were not heat-shocked.

Male progeny of the germline clones were collected and their lifespan measured to

confirm the presence of the *drd$^{lwf}$* mutation.

## Visualization of *drd* expression pattern

155       *yw drd-GAL4/FM7i-GFP* females were crossed with *w; UAS-GFP.nls* males.

156   Female progeny were crossed with sibling males. Ovaries from females 3-6 days

157   post-eclosion were dissected in insect Ringers and separated into individual egg

158   chambers. Samples were imaged on a Nikon A1 Confocal Microscope (Nikon,

159   Tokyo, Japan) with NIS-Elements AR software (Nikon).

## Neutral red permeability assay

Eggs were collected on apple-juice agar plates for 2-19 hr, placed into a

stainless steel mesh basket, and rinsed with PBS. Eggs were dechorionated by

gently shaking in a 50% bleach solution for 3 min followed by rinsing with PBS;

exposure to bleach was only 1 min for *drd$^1$/drd$^1$* and *drd$^1$/FM7c*. The dechorionated

eggs were counted, stained with 5 mg/mL neutral red (VWR, Radnor, PA) in PBS for

10 min, rinsed with PBS, and scored as stained or unstained. No correlation was

observed between the duration of the egg collection and the staining results. See S1

Text for detailed protocol.

## Western blot analysis

Egg chambers were dissected in PBS or eggs were collected on apple-juice

agar plates, and samples were homogenized in 80µL of 20mM Tris-HCl (pH7.5),

0.15 M NaCl, 100 mM DTT. Samples were then heated at 100°C for 5 min,

centrifuged (14,000g, 1 min), and the resulting pellet discarded. One quarter volume

5x SDS-PAGE loading buffer was added to each sample, and they were again

heated for 5 min at 100°C and stored at -20°C until further use. Prior to

electrophoresis, samples were treated with 5% β-mercaptoethanol and heated for 3

min at 100°C. For any gel, the same amount of egg chamber equivalents of each

sample was loaded, typically ranging from 1-5 egg chamber equivalents. Following

separation via SDS-PAGE (12%, Mini PROTEAN 3 System, Bio-Rad, Hercules, CA),

proteins were transferred to PVDF membrane for 1hr using a Genie electroblot

chamber (Idea Scientific, Minneapolis, MN). Membranes were then washed for 10

min in PBS and blocked overnight in PBS/0.05% Tween-20 (PBS-T)/ 5% nonfat dry

milk at 4°C. After blocking, two 5 min washes in PBS-T were conducted prior to 1 hr

incubation in primary antibody diluted in PBS-T/1% BSA. Membranes were then

washed in PBS-T, once for 15 min, and four times for 5 min, followed by a 1 hr

incubation in secondary antibody (ECL HRP-linked donkey anti-rabbit IgG, 1:10,000,

Cytiva Life Sciences, Marlborough, MA). Again one 15 min and four 4 min washes in

PBS-T were conducted and antibody signals were detected via chemiluminescence

(ECL Prime Western Blotting System, Cytiva Life Sciences, Marlborough, MA).

Primary polyclonal rabbit antibodies were provided by Dr. Gail Waring and were

previously characterized antibodies against Cp36 and Vm26Ab [9].

## Immunostaining

*drd¹* heterozygous and homozygous females were collected on the day of

eclosion on placed on yeast paste with sibling males for two days. Ovaries were

immunostained as described [15], except that fixation was performed with 4%

paraformaldehyde in PBS rather than formaldehyde in PBS/Triton X-100. Anti-

Vm26Ab antibody was used at 1:5000, and the secondary antibody was Alexafluor

488 goat anti-rabbit IgG (1:400) (Invitrogen, Carlsbad, CA). Samples were imaged

on a Nikon A1 Confocal Microscope (Nikon, Tokyo, Japan) with NIS-Elements AR

software (Nikon). Images were analyzed in ImageJ v2.9.0, and the analyzer was

blind to genotype. VM staining intensity was determined in stage 9 and 10A egg

chambers by measuring mean pixel brightness in a 1mm x 5 mm rectangle of the

anterior lateral oocyte margin. Within each genotype, there was no significant

correlation between either VM width or staining intensity and developmental stage

(as measured by the oocyte length/width ratio), allowing us to pool data across

developmental stages for comparison between genotypes.

## *In vitro* egg activation

To stimulate egg production, *drd¹/FM7c* and *drd¹/drd¹* females were placed on

yeast paste and mated with sibling males 3-5 days before dissection. Egg activation

*in vitro* was performed using the method of Page and Orr-Weaver [44]. Stage 14 egg

chambers were dissected in isolation buffer (55 mM NaOAc, 40 mM KOAc, 110 mM

sucrose, 1.2 mM $MgCl_2$, 1 mM $CaCl_2$, 100 mM Hepes, pH 7.4 (NaOH)). Egg

chambers were then incubated for 10 min in hypo-osmotic activating buffer (3.3 mM

$NaH_2PO_4$, 16.6 mM $KH_2PO_4$, 10 mM NaCl, 50 mM KCl, 5% PEG 8000, 2 mM $CaCl_2$,

pH 6.4 (1:5 NaOH: KOH)) and then transferred into modified Zalokar's buffer for 30

min (9 mM $MgCl_2$, 10 mM $MgSO_4$, 2.9 mM $NaH_2PO_4$ 0.22 mM NaOAc, 5 mM

glucose, 27 mM glutamic acid, 33 mM glycine, 2 mM malic acid, 7 mM $CaCl_2$, pH 6.8

(1:1 NaOH: KOH)). To test for eggshell crosslinking, egg chambers were then

incubated in 50% bleach for 5 min and scored as intact, leaky, or completely

dissolved. In some experiments, hydrogen peroxide (0.006-0.06%) was included in

both the activating and Zalokar's buffer.

In a second series of experiments, the activating buffer incubation was

omitted. Egg chambers were dissected and manually dechorionated in isolation

buffer and then incubated for 30 min in Zalokar's buffer containing 4.5% hydrogen

peroxide and 1 mg/ml horseradish peroxidase (VWR, Radnor, PA) before bleaching

as above.

## 232  Data analysis and statistics

233  Data were graphed and analyzed using GraphPad Prism v9 for Windows (GraphPad

234  Software, San Diego, CA, www.graphpad.com). A p value of $<0.05$ was used as the

235  threshold for statistical significance.

# Results

## Identification of the *drd¹* mutation

The experiments in this study utilize flies carrying the two most severe alleles of *drd*: *drd^lwf* and *drd¹*. The mutation in the latter of these alleles not been molecularly characterized, although we previously reported that there were no alterations in the protein coding sequence [34]. Further sequencing of the final 5 introns and the ends of the first three large introns revealed six differences between *drd¹* and wild-type. One of these, a T to A transversion in the final intron, is predicted to create a strong ectopic splice acceptor site and result in the inclusion of an additional 10 nucleotides in the spliced transcript (S1 Fig) (www.fruitfly.org/seq_tools/splice.html) [45]. This aberrant splicing of the *drd* transcript in *drd¹* mutants was confirmed by RT-PCR. Virtual translation of the mutant transcript predicted that the final 76 amino acids of the Drd protein are replaced with a novel sequence of 45 amino acids in the *drd¹* mutant.

## Sterility of *drd* mutant females

We have previously reported that females homozygous for severe *drd* alleles are sterile and rarely lay eggs, and their ovaries contain very few vitellogenic egg chambers [34]. We hypothesized that these phenotypes could be an effect of the starvation observed in *drd* mutant flies, as opposed to a direct phenotype of *drd* mutation. Consistent with this hypothesis, feeding females a high protein diet (yeast paste) stimulated egg-laying in a subset of flies. Yeast paste increased the percentage of *drd^lwf* females, but not *drd¹* females, that laid eggs, and it increased by

more than 30-fold the median number of eggs laid during their lifetime by those *drd¹*

and *drd^{lwf}* homozygous females that laid eggs (S2 Fig).

Despite the improvement in egg-laying, *drd* homozygous females fed with

yeast paste remained sterile. Eggs were collected overnight from mated *drd¹*

heterozygotes and homozygotes and allowed to develop for an additional two hr

before examination for progression through embryogenesis. As shown in Table 1, a

significantly higher number of eggs laid by homozygotes were collapsed (p<0.0001,

Fisher's exact test). Of the turgid eggs, the large majority were fertilized, and the rate

of fertilization was not affected by the mother's genotype (p=0.30, Fisher's exact

test). However, virtually no fertilized eggs laid by homozygotes underwent

gastrulation, in contrast to embryos laid by heterozygotes (p<0.0001, Fisher's exact

test). Thus, the sterility of *drd* mutant females appears to result from early embryonic

arrest.

**Table 1. Sterility of *drd* mutant females**

| Maternal genotype (# of eggs) | *drd¹/FM7c* (164) | *drd¹/ drd¹* (293) |
|---|---|---|
| Collapsed (% of total) | 0.6% | 17.7% |
| Unfertilized (% of turgid eggs) | 7.5% | 11.2% |
| Pre-gastrulation (% of fertilized eggs) | 6.8% | 99.0% |
| Post-gastrulation (% of fertilized eggs) | 93.2% | 1.0% |

Table 1: Embryonic arrest in eggs laid by *drd¹* females. The second row indicates the

percent of all eggs that were collapsed or flaccid. The following row indicates the

percent of turgid eggs that were unfertilized (4 eggs from heterozygotes and 9 from

homozygotes could not be scored and were excluded). The final two rows indicate

the percent of fertilized eggs that were scored as pre-gastrulation and post-

gastrulation, respectively.

## Expression of *drd* in the egg chamber

To determine whether *drd* expression was required in the germline or soma

for female fertility, we created germline clones of *drd^lwf^*. Almost all of these females

(15/17) were fertile when crossed with wild-type males, while non-heat shocked

controls were all sterile (n=28), indicating that *drd* expression is not required in the

female germline for fertility. The male progeny of these clones were short-lived

(median lifespan of 4 days, n=55), as would be expected for males hemizygous for

*drd^lwf^*, confirming that the germline of these clones was mutant for *drd*.

Consistent with the germline clone analysis, driving a GFP reporter with a *drd-*

*GAL4* transgene resulted in labeling of all ovarian follicle cells (Fig 1). Visible

reporter expression appeared at stage 10B and persisted for the remainder of

oogenesis.

**Fig 1. Expression of *drd* in ovarian follicle cells.** The image shows a maximum

intensity projection of one stage 14 (top) and two stage 10B egg chambers in which

expression of a nuclear GFP reporter is driven by *drd-GAL4*. No staining was seen in

the nurse cell nuclei located in the anterior half of each stage 10B egg chamber (*).

No staining was seen in egg chambers from sibling control females lacking the *drd-*

*GAL4* driver (S3 Fig).

## Abnormal eggshell development in *drd* mutant females

Because a significant number of eggs laid by *drd* mutants were collapsed and

because *drd* expression was observed in the follicle cells and not the germline, we

next examined the integrity of the eggshell, a structure synthesized by the somatic

follicle cells. The integrity of the inner layer of the eggshell, the VM, was assayed by

staining dechorionated eggs with neutral red, a dye that is normally excluded by the

VM [20]. Virtually no eggs laid by *drd[1]* and *drd[lwf]* heterozygotes lysed during

dechorionation (1-3%) or were stained with neutral red (3-4%) (Fig 2). In contrast,

eggs laid by homozygotes of either *drd* allele showed a high susceptibility to lysis

during dechorionation (31-42%), and the large majority of surviving eggs were

permeable to neutral red (71-90%). A small fraction of eggs (4-11%) were not

dechorionated upon treatment with bleach and could not be assessed for dye

exclusion, but the abundance of such eggs was not affected by the genotype of the

mother (p=0.19, Chi-square test).

**Fig 2. Neutral red staining of eggs laid by *drd* homozygous and heterozygous**

**females.** (A) fraction of eggs that lysed during bleach treatment. (B) fraction of eggs

that were stained with neutral red after successful dechorionation. ****: significant

difference between eggs laid by heterozygotes vs homozygotes, p<0.0001, Fisher's

exact test. 135-249 eggs per genotype.

As a direct test of the incorporation of eggshell proteins into an insoluble

cross-linked matrix, we performed Western blots on lysates of staged egg chambers

and laid eggs. An antibody against the chorion protein Cp36 detected the expected

pattern of staining in samples from *drd[1]/FM7c* heterozygous females (Fig 3) [9].

Soluble Cp36 protein was not detected at stage 10 of oogenesis, which is before

chorion proteins such as Cp36 begin to be expressed, but Cp36 was detected in

stage 14 lysates, when the protein is present in the chorion but has not yet been fully

cross-linked. In lysates from eggs collected either 3 or 6 hr after deposition, Cp36

was not detectable, consistent with complete cross-linking of the protein into the

insoluble chorion. The same pattern of staining was observed in lysates from eggs

and egg chambers from *drd¹* homozygotes (Fig 3), indicating that crosslinking of the

chorion, or at least of the specific protein Cp36, is not affected by mutation of *drd*.

**Fig 3. Western blot of the chorion protein Cp36 in *drd* mutant and control egg**

**chambers and eggs.** Stage 10 and 14 egg chambers were dissected from *drd¹/drd¹*

mutant and *drd¹/FM7c* heterozygous females, and eggs laid by these females were

collected 0-3 and 0-6 hr after oviposition (lanes 3 and 7 and lanes 4 and 8). 4 eggs

or egg chambers per lane, 1:5000 primary antibody dilution.

In contrast to the chorion, crosslinking of VM proteins was clearly abnormal in

eggs laid by *drd* mutant homozygotes. Fig 4A shows the pattern of staining observed

in egg chambers and eggs from heterozygous controls, using an antibody raised

against the VM protein Vm26Ab [9]. This antibody has been reported to cross-react

with multiple VM proteins due to their high degree of sequence similarity [18,46], and

we typically observed multiple bands on Western blots. As expected, staining was

observed in lysates from stage 10 egg chambers, when the VM proteins are not yet

crosslinked, and from stage 14, when the VM proteins are crosslinked by disulfide

bonds but are soluble in the presence of reducing agents. In laid eggs from

heterozygotes, no soluble VM proteins were observed, consistent with the formation

of nonreducible dityrosine bonds among VM proteins during ovulation and egg

activation. In lysates of laid eggs from *drd¹* homozygotes, VM proteins were detected

on the Western blot, indicating that these proteins are not fully crosslinked in the

absence of *drd* expression. We observed solubility of VM proteins in laid eggs of *drd¹*

homozygotes in seven Western blots representing five biological replicates. The

decrease in signal seen in Fig 4A with increasing time after oviposition (compare

lanes 3 and 4) was not consistently observed.

**Fig 4. Western blots against vitelline membrane proteins in *drd* mutant and**

**control egg chambers and eggs.** (A) Western blot of samples treated with reducing

agents with an antibody raised against Vm26Ab. Samples include stage 10 and 14

egg chambers dissected from *drd¹/drd¹* mutant and *drd¹/FM7c* heterozygous

females, and eggs laid by these females were collected 0-3 and 0-6 hr after

oviposition. 4 eggs or egg chambers per lane, 1:10,000 primary antibody dilution. (B)

Western blot from egg chambers solubilized in the absence of reducing agent and

probed with an antibody against Vm26Ab. Samples include stage 10 and 14 egg

chambers dissected from *drd¹/drd¹* mutant and *drd¹/FM7c* heterozygous females. 2

egg chambers per lane, 1:10,000 primary antibody dilution.

To assay for the formation of disulfide bonds among VM proteins during

oogenesis, lysates were prepared from stage 10 and 14 egg chambers in the

absence of reducing agents. Mutation of *drd* did not alter the pattern of

immunostaining observed under these conditions (Fig 4B). Soluble protein was

detected at stage 10 but not at stage 14, indicating that VM proteins are cross-linked

by reducible bonds during oogenesis in both *drd* homozygotes and heterozygotes.

We performed immunostaining of stage 9 and 10A egg chambers against

Vm26Ab to determine whether the appearance of the developing VM is altered in *drd*

mutants (Fig 5A, B). We observed no significant effect of genotype on either the

staining intensity (p=0.14, Mann-Whitney test) or width (p=0.33, t-test) of the VM

(n=20-22 samples per genotype). There were also no obvious morphological

differences in the egg chambers between the two genotypes. However, *drd*[1]

homozygous egg chambers were slightly but significantly smaller than heterozygous

controls. We used the size ratio of anterior-posterior length/lateral width of each

oocyte as a measure of progression through oogenesis, as this ratio increases

during stages 9 and 10A (Fig 5C). Anterior-posterior length was significantly linearly

correlated with the size ratio for each genotype (p<0.0001, F test). The slope of the

relationship did not differ between the two genotypes (p=0.88) but the intercept did

differ (p=0.02), corresponding to a decrease in size of homozygous oocytes of

approximately 15%.

**Fig 5. Immunostaining of the VM**. Stage 10A egg chambers from a *drd*[1]

heterozygote (A) and homozygote (B) were stained with the anti-Vm26Ab antibody.

The rectangle indicates the region in which staining intensity and eggshell width

were measured. Oo: oocyte; nc: nurse cells; fc: follicle cells. (C) Plot of the size ratio

of anterior-posterior length/lateral width for each oocyte as a function of the anterior-

posterior length, showing the slight difference between *drd*[1] heterozygotes and

homozygotes.

# Follicle cell knockdown of *drd* recapitulates mutant

# phenotypes

Our finding that *drd* expression is required in the soma for female fertility,

coupled with the known role of the somatic follicle cells in the synthesis of the

eggshell, suggested that all of the *drd* mutant phenotypes related to fertility and

eggshell assembly could be associated with the expression of *drd* in the follicle cells.

To test this, we knocked down *drd* expression in the follicle cells, using two pan-

follicle cell GAL4 drivers, *CY2-GAL4* and *T155-GAL4*, and two inducible *drd* RNAi

transgenes, *drd^{GD3367}* and *drd^{GD15915}*, that we have previously shown to be effective

at knocking down *drd* expression [40]. Females heterozygous for a GAL4 driver and

an RNAi transgene (referred to as knockdown females) and sibling control females

that lacked the RNAi transgene were mated with sibling males, and their eggs were

scored for fertilization, embryonic development, eggshell integrity, and crosslinking.

As shown in Table 2, a greater percentage of eggs from knockdown females

were collapsed compared with sibling controls, and virtually all fertilized eggs from

knockdown females were arrested pre-gastrulation, as was seen in mutant females

(Fisher's exact test, $p < 0.0001$ for each genotype). The proportion of eggs laid by

knockdown females that was fertilized was not significantly different than that of

sibling controls for three of the four genotypes tested, with a small but significant

effect on fertilization for one combination of GAL4 driver and RNAi transgene.

Despite the near-universal arrest of eggs from knockdown females early in

oogenesis in this assay, we did occasionally observe larvae in vials of knockdown

females carrying *T155-GAL4* and either of the RNAi-transgenes, indicating that

these females were not fully sterile. Knockdown females carrying the *CY-GAL4*

driver, like *drd* mutant females, appeared to be completely sterile.

**Table 2. Sterility of *drd* knockdown females**

| Maternal genotype (# of eggs) | *w; UAS-Dcr-2 $drd^{GD3367}$/+; T155-GAL4/+* (188) | *w; $drd^{GD15915}$ UAS-Dcr-2/+; T155-GAL4/+* (155) | *w; UAS-Dcr-2 $drd^{GD3367}$/CY2-GAL4* (169) | *w; $drd^{GD15915}$ UAS-Dcr-2/CY2-GAL4* (206) |
|---|---|---|---|---|
| Collapsed (% of total) | 32.4%**** | 43.9%**** | 21.9%**** | 24.3%**** |
| Unfertilized (% of turgid eggs) | 6.1% | 10.0% | 7.8%* | 4.8% |
| Pre-gastrulation (% of fertilized eggs) | 100%**** | 98.6%**** | 100%**** | 96.4%**** |
| | *w; CyO/+; T155-GAL4/+* (153) | *w; CyO/+; T155-GAL4/+* (221) | *w; CY2-GAL4/CyO* (238) | *w; CY2-GAL4/CyO* (163) |
| Collapsed (% of total) | 0% | 0.5% | 0.4% | 0.6% |
| Unfertilized (% of turgid eggs) | 3.4% | 10.3% | 2.2% | 4.5% |
| Pre-gastrulation (% of fertilized eggs) | 0.7% | 1.0% | 1.3% | 2.7% |

Table 2: Embryonic arrest upon knockdown of *drd* expression in the follicle cells.

Knockdown females (top four rows) and sibling controls (bottom four rows) were

mated with sibling males, and eggs were collected and scored as in Table 1. The

number of uncollapsed but unscorable eggs that were excluded from further analysis

varied from 4-13 among each of the eight genotypes. ****: different from sibling

controls, $p < 0.0001$, Fisher's exact test, *: $p = 0.024$.

Tests of eggshell integrity by neutral red exclusion demonstrated that eggs

laid by follicle cell knockdown females showed a similar defect to those laid by *drd*

mutants. As shown in Fig 6, we observed a significant and consistent increase in the

proportion of eggs that were permeable to neutral red compared to those laid by

sibling controls (Fig 6C). The percent of eggs that lysed upon dechorionation was

also significantly different between knockdown and sibling control females for three

of the four genotypes (Fig 6A). Finally, and in contrast to the data from *drd* mutants,

we observed a small but significant increase in the fraction of eggs from knockdown

females that were successfully dechorionated relative to sibling controls in two of the

four genotypes (Fig 6B).

**Fig 6. Neutral red staining of eggs following *drd* knockdown.** Data from eggs

laid by *drd* knockdown (striped orange and green bars) and sibling control (solid

black bars) females. (A) fraction of eggs that lysed during bleach treatment. (B)

fraction of eggs that were successfully dechorionated. (C) fraction of eggs that were

stained with neutral red after dechorionation. Significant difference between eggs

laid by knockdown females and sibling controls by Fisher's exact test are indicated:

**** $p < 0.0001$, *** $p = 0.0006$, ** $p = 0.0016$, * $p = 0.015$. 195-225 eggs per genotype.

Crosslinking of VM proteins in laid eggs and stage 14 egg chambers from

knockdown and control females was assayed by Western blot with the antibody

against Vm26Ab as described above. A representative blot is shown in Fig 7.

Knockdown females carrying the *CY2-GAL4* driver showed a phenotype identical to

that of *drd* mutants: we consistently observed soluble VM proteins in both stage 14

egg chambers and laid eggs, while soluble VM proteins were only observed in the

stage 14 chambers of sibling controls (3 biological replicates with $drd^{GD3367}$ and 2

with $drd^{GD15915}$). In contrast, soluble VM proteins were detected in only some

samples of eggs laid by knockdown females carrying the *T155-GAL4* driver (2/4

biological replicates with $drd^{GD3367}$ and 1/3 with $drd^{GD15915}$).

**Fig 7. Western blots against vitelline membrane proteins in *drd* knockdown**

**egg chambers and eggs.** Blots were probed with an antibody raised against

Vm26Ab. Samples include stage 14 egg chambers dissected from, and eggs laid by,

females in which the inducible *drd* RNAi transgene, $drd^{GD3367}$, was driven by either

the *CY2-GAL4* or *T155-GAL4* driver, or sibling controls lacking the RNAi transgene.

Lanes are marked "st 14" for egg chambers and "e" for eggs. 2 eggs or egg

chambers per lane, 1:25,000 primary antibody dilution. Eggs were collected between

0-4.5 hr after oviposition.

# Cross-linking of VM in isolated egg chambers

To determine whether the *drd* cross-linking defect persists in isolated egg

chambers, we dissected stage 14 egg chambers and activated them *in vitro* by

exposure to hypo-osmotic medium as previously reported [44]. VM cross-linking was

assayed by then incubating the egg chambers in 50% bleach (Fig 8A). All egg

chambers dissected from $drd^1$/FM7c control females were successfully cross-linked

following hypo-osmotic treatment. In contrast, all egg chambers dissected from $drd^1$

homozygotes completely dissolved in bleach. Additional egg chambers were

dissected from homozygotes and treated with 0.006-0.06% hydrogen peroxide

during treatment with hypo-osmotic medium and for 30 min afterwards, and all these

egg chambers also dissolved completely in bleach. It is noteworthy that the *drd*

mutant phenotype is more severe in egg chambers activated *in vitro* than *in vivo*, as

most eggs laid by *drd* homozygous females remain intact upon treatment with bleach

(see Fig 2 above).

**Fig 8. Results of *in vitro* egg activation.** The figure indicates the number of egg

chambers dissected from each female genotype that remained intact, became leaky,

or dissolved in 50% bleach after treatment with hypo-osmotic medium (A) or

hydrogen peroxide and horseradish peroxidase (B).

We then attempted to directly cross-link the VM of stage 14 egg chambers by

treatment with hydrogen peroxide and peroxidase. Stage 14 egg chambers were

dissected, manually dechorionated, and treated with peroxide/peroxidase without

exposure to hypo-osmotic medium (Fig 8B). Egg chambers dissected from

*drd$^1$/FM7c* control females and treated with 1 mg/ml peroxidase and 4.5% hydrogen

peroxide all survived a subsequent challenge with bleach. This cross-linking was not

due to inadvertent activation of the egg chambers during dechorionation, as omitting

the hydrogen peroxide from the incubation resulted in all egg chambers dissolving in

bleach. Egg chambers from *drd$^1$* homozygotes treated with peroxide/peroxidase as

above all dissolved in bleach.

# Discussion

We have demonstrated that expression of *drd* in the ovarian follicle cells is

both necessary and sufficient for female fertility and the progression of embryonic

development beyond gastrulation. Germline clones in which *drd* expression is

restricted to somatic cells are fertile, while knockdown of *drd* expression specifically

in the follicle cells recapitulates the mutant phenotype. These results are consistent

with those of Kim et al, who reported expression of *drd* in the follicle cells [47].

Furthermore, we have shown that *drd* expression in the follicle cells is required for

proper development of the VM layer of the eggshell. In the absence of such

expression, many eggs collapse, and the remaining eggs have a fragile VM that fails

to act as a permeability barrier. Several major VM proteins—specifically those

recognized by the antibody used in these studies—remain soluble in the presence of

reducing agents, indicating that they have not been incorporated into the insoluble

network of cross-linked proteins seen in the wild-type VM. The variable solubility

phenotype observed with knockdown of *drd* with the *T155-GAL4* driver is consistent

with the fertility of some of these knockdown females and suggests that *T155-GAL4*

is not as effective as the *CY2-GAL4* driver in knocking down *drd* expression. The

difference in strength of these two driver lines has been reported previously [48].

The solubility of VM proteins in eggs laid by *drd* mutants indicates a defect in

the peroxidase-mediated cross-linking that normally occurs upon egg activation while

the egg is transiting down the oviduct. We have no evidence that egg activation itself

is defective, as the eggs were able to complete the early stages of embryogenesis.

Our data don't address the connection between defective VM cross-linking and

embryonic arrest, and it remains possible that embryonic arrest is an independent

*drd* phenotype. However, this same pairing of phenotypes has been reported with

another eggshell mutant. Females mutant for *psd*, which encodes a minor VM

protein, also lay eggs with a VM cross-linking defect and that arrest pre-gastrulation

and show a chromatin margination phenotype similar to that induced by anoxia [32].

Thus, it is possible that defective VM cross-linking is a direct cause of early

developmental arrest.

A published microarray study of ovarian gene expression has reported that

*drd* expression in the egg chamber begins at stage 8 of oogenesis, peaks at stages

10A and 10B, and then declines [16]. The timing of *drd* expression therefore parallels

the synthesis and secretion of VM proteins by the follicle cells. The pattern of

reporter expression shown in Fig 1 is consistent with the microarray data. We

observed GFP fluorescence starting at stage 10B and persisting through the rest of

oogenesis; one would expect both a delay between the onset of GFP expression and

significant accumulation in the follicle cell nuclei and persistence of the protein after

gene expression is downregulated. The Drd protein is unlikely to be a component of

the VM, as it is predicted to be an integral membrane protein and is reported to be

localized to an intracellular compartment [34,47]. Drd is also unlikely to be directly

involved in the cross-linking process, which occurs at the end of oogenesis when *drd*

expression is very low. Rather, our data suggest that the failure of VM proteins to

become cross-linked in *drd* mutants could be due to an absence of some

modification of the VM proteins in the follicle cells prior to secretion. The results of

our final experiment are consistent with this hypothesis. Incubation of stage 14 egg

chambers with peroxide and peroxidase resulted in cross-linking of the VM in egg

chambers from wild-type but not mutant females. Thus, VM proteins synthesized and

secreted from *drd* mutant follicle cells appear to be poor substrates for peroxidase-

mediated cross-linking for reasons still to be determined.

One interesting finding from our final experiment is that egg chambers from

*drd* mutant females dissolve immediately in bleach after activation *in vitro* with

hypoosmotic medium. In contrast, eggs laid by *drd* mutant females mainly survive

bleaching, even though their VMs are permeable to neutral red. The contrast

between these two results indicates that hypoosmotic treatment *in vitro* does not fully

recapitulate the activation process *in vivo* even though the two processes appear to

give identical results in wild-type flies [44]. The *drd* mutant female could prove to be

a useful system for identifying additional factors that influence eggshell maturation

during ovulation and oviposition.

In addition to characterizing the effect of *drd* mutations on oogenesis, we have

identified the molecular defect in the severe *drd$^1$* allele as a point mutation in the final

intron that disrupts the normal splicing of exons 8 and 9. The aberrant splicing

replaces the final 76 residues of the 827 amino acid Drd protein with a novel

sequence. Because *drd$^1$* is phenotypically identical to *drd$^{lwf}$*, a nonsense mutation

that eliminates all but the first 180 amino acids, our finding highlights the importance

of the C-terminal of Drd in protein function, stability, or localization. In contrast, the

*drd$^{W3}$* and *In(1)drd$^{x1}$* alleles, which eliminate the first exon and at least the first 125

amino acids, are phenotypically less severe [34].

The biochemical function of the Drd protein remains unknown. However, this

study highlights two themes that are emerging from our studies of this gene. First,

*drd* expression appears to be required in a number of different epithelial tissues,

including the ovarian follicle cells for oogenesis, the anterior midgut for digestive

function, and the respiratory tracheae for brain integrity. Second, *drd* expression is

required for the formation of extracellular barrier structures, as *drd* mutants show

defects both in the eggshell and in the peritrophic matrix of the midgut. Given the

amount of information known about eggshell formation, the ovary is an excellent

system for further studies of *drd* function.

# Acknowledgments

582    We thank Dr. Celeste Berg for the CY2-GAL4 stock, Drs. Gail Waring and Anita

583    Manogaran for antibodies and helpful discussions, and Anika Benske for technical

584    assistance. Stocks obtained from the Bloomington Drosophila Stock Center (NIH

585    P40OD018537) and the Vienna Drosophila Resource Center were used in this study.

# References Cited

1. Yue B. Biology of the extracellular matrix. J Glaucoma. 2014;23: S20–S23.
doi:10.1097/ijg.0000000000000108

2. Broadie K, Baumgartner S, Prokop A. Extracellular matrix and its receptors in
Drosophila neural development. Dev Neurobiol. 2011;71: 1102–1130.
doi:10.1002/dneu.20935

3. Walma DAC, Yamada KM. The extracellular matrix in development. Development.
2020;147: dev175596. doi:10.1242/dev.175596

4. Waring GL. Morphogenesis of the eggshells in Drosophila. Int Rev Cytol.
2000;198: 67–108. doi:10.1016/s0074-7696(00)98003-3

5. Merkle JA, Wittes J, Schüpbach T. Signaling between somatic follicle cells and the
germline patterns the egg and embryo of Drosophila. Curr Top Dev Biol. 2019;140:
55–86. doi:10.1016/bs.ctdb.2019.10.004

6. Narasimha S, Nagornov KO, Menin L, Mucciolo A, Rohwedder A, Humbel BM, et
al. Drosophila melanogaster cloak their eggs with pheromones, which prevents
cannibalism. PLoS Biol. 2019;17: e2006012. doi:10.1371/journal.pbio.2006012

7. Stein DS, Stevens LM. Maternal control of the Drosophila dorsal–ventral body
axis. Wiley Interdiscip Rev: Dev Biol. 2014;3: 301–330. doi:10.1002/wdev.138

8. Fakhouri M, Elalayli M, Sherling D, Hall JD, Miller E, Sun X, et al. Minor proteins
and enzymes of the Drosophila eggshell matrix. Dev Biol. 2006;293: 127–141.
doi:10.1016/j.ydbio.2006.01.028

9. Pascucci T, Perrino J, Mahowald AP, Waring GL. Eggshell assembly in
Drosophila: processing and localization of vitelline membrane and chorion proteins.
Dev Biol. 1996;177: 590–598. doi:10.1006/dbio.1996.0188

10. Fargnoli J, Waring GL. Identification of vitelline membrane proteins in Drosophila
melanogaster. Dev Biol. 1982;92: 306–314. doi:10.1016/0012-1606(82)90177-4

11. Andrenacci D, Cernilogar FM, Taddei C, Rotoli D, Cavaliere V, Graziani F, et al.
Specific domains drive VM32E protein distribution and integration in Drosophila
eggshell layers. J Cell Sci. 2001;114: 2819–2829. doi:10.1242/jcs.114.15.2819

12. Alatortsev VE. New genes for vitelline membrane proteins in Drosophila. Mol
Biol. 2006;40: 330–332. doi:10.1134/s002689330602021x

13. Jiménez G, González-Reyes A, Casanova J. Cell surface proteins Nasrat and
Polehole stabilize the Torso-like extracellular determinant in Drosophila oogenesis.
Gene Dev. 2002;16: 913–918. doi:10.1101/gad.223902

14. Ventura G, Furriols M, Martín N, Barbosa V, Casanova J. closca, a new gene
required for both Torso RTK activation and vitelline membrane integrity. Germline
proteins contribute to Drosophila eggshell composition. Dev Biol. 2010;344: 224–
232. doi:10.1016/j.ydbio.2010.05.002

15. Furriols M, Casanova J. Germline and somatic vitelline proteins colocalize in
aggregates in the follicular epithelium of Drosophila ovaries. Fly. 2014;8: 113–119.
doi:10.4161/fly.29133

16. Tootle TL, Williams D, Hubb A, Frederick R, Spradling A. Drosophila eggshell
production: identification of new genes and coordination by Pxt. PLoS One. 2011;6:
e19943. doi:10.1371/journal.pone.0019943

17. Cavaliere V, Bernardi F, Romani P, Duchi S, Gargiulo G. Building up the
Drosophila eggshell: First of all the eggshell genes must be transcribed. Dev Dyn.
2008;237: 2061–2072. doi:10.1002/dvdy.21625

18. Manogaran A, Waring GL. The N-terminal prodomain of sV23 is essential for the
assembly of a functional vitelline membrane network in Drosophila. Dev Biol.
2004;270: 261–71. doi:10.1016/j.ydbio.2004.02.009

19. Petri WH, Wyman AR, Kafatos FC. Specific protein synthesis in cellular
differentiation III. The eggshell proteins of Drosophila melanogaster and their
program of synthesis. Dev Biol. 1976;49: 185–199. doi:10.1016/0012-
1606(76)90266-9

20. LeMosy EK, Hashimoto C. The Nudel protease of Drosophila is required for
eggshell biogenesis in addition to embryonic patterning. Dev Biol. 2000;217: 352–
361. doi:10.1006/dbio.1999.9562

21. Heifetz Y, Yu J, Wolfner MF. Ovulation triggers activation of Drosophila oocytes.
Dev Biol. 2001;234: 416–424. doi:10.1006/dbio.2001.0246

22. Mindrinos MN, Petri WH, Galanopoulos VK, Lombard MF, Margaritis LH.
Crosslinking of the Drosophila chorion involves a peroxidase. Wilhelm Roux Arch
Dev Biol. 1980;189: 187–196. doi:10.1007/bf00868677

23. Heinecke JW. Tyrosyl radical production by myeloperoxidase: a phagocyte
pathway for lipid peroxidation and dityrosine cross-linking of proteins. Toxicology.
2002;177: 11–22. doi:10.1016/s0300-483x(02)00192-0

24. Hall HG. Hardening of the sea urchin fertilization envelope by peroxidase-
catalyzed phenolic coupling of tyrosines. Cell. 1978;15: 343–355. doi:10.1016/0092-
8674(78)90003-x

25. Han Q, Li G, Li J. Purification and characterization of chorion peroxidase from
Aedes aegypti eggs. Arch Biochem Biophys. 2000;378: 107–115.
doi:10.1006/abbi.2000.1821

26. Bauer BJ, Waring GL. 7C female sterile mutants fail to accumulate early eggshell
proteins necessary for later chorion morphogenesis in Drosophila. Dev Biol.
1987;121: 349–358. doi:10.1016/0012-1606(87)90171-0

27. Hawley RJ, Waring GL. Cloning and analysis of the dec-1 female-sterile locus, a
gene required for proper assembly of the Drosophila eggshell. Gene Dev. 1988;2:
341–349. doi:10.1101/gad.2.3.341

28. Savant SS, Waring GL. Molecular analysis and rescue of a vitelline membrane
mutant in Drosophila. Dev Biol. 1989;135: 43–52. doi:10.1016/0012-1606(89)90156-
5

29. D'Alterio C, Tran DDD, Yeung MWYA, Hwang MSH, Li MA, Arana CJ, et al.
Drosophila melanogaster Cad99C, the orthologue of human Usher cadherin
PCDH15, regulates the length of microvilli. J Cell Biol. 2005;171: 549–58.
doi:10.1083/jcb.200507072

30. Schlichting K, Wilsch-Bräuninger M, Demontis F, Dahmann C. Cadherin Cad99C
is required for normal microvilli morphology in Drosophila follicle cells. J Cell Sci.
2006;119: 1184–95. doi:10.1242/jcs.02831

31. Claycomb JM, Benasutti M, Bosco G, Fenger DD, Orr-Weaver TL. Gene
amplification as a developmental strategy: isolation of two developmental amplicons
in Drosophila. Dev Cell. 2004;6: 145–55.

32. Elalayli M, Hall JD, Fakhouri M, Neiswender H, Ellison TT, Han Z, et al. Palisade
is required in the Drosophila ovary for assembly and function of the protective
vitelline membrane. Dev Biol. 2008;319: 359–369. doi:10.1016/j.ydbio.2008.04.035

33. Cernilogar FM, Fabbri F, Andrenacci D, Taddei C, Gargiulo G. Drosophila
vitelline membrane cross-linking requires the fs(1)Nasrat, fs(1)polehole and chorion
genes activities. Dev Genes Evol. 2001;211: 573–580. doi:10.1007/s00427-001-
0192-1

34. Blumenthal EM. Cloning of the neurodegeneration gene drop-dead and
characterization of additional phenotypes of its mutation. Fly. 2008;2: 180–8.

35. Peller CR, Bacon EM, Bucheger JA, Blumenthal EM. Defective gut function in
drop-dead mutant Drosophila. J Insect Physiol. 2009;55: 834–9.
doi:10.1016/j.jinsphys.2009.05.011

36. Conway S, Sansone CL, Benske A, Kentala K, Billen J, Broeck JV, et al.
Pleiotropic and novel phenotypes in the Drosophila gut caused by mutation of drop-
dead. J Insect Physiol. 2018;105: 76–84. doi:10.1016/j.jinsphys.2018.01.007

37. Sansone CL, Blumenthal EM. Neurodegeneration in drop-dead mutant
Drosophila melanogaster is associated with the respiratory system but not with
hypoxia. PLoS One. 2013;8: e68032. doi:10.1371/journal.pone.0068032

38. Buchanan RL, Benzer S. Defective glia in the Drosophila brain degeneration
mutant drop-dead. Neuron. 1993;10: 839–850. doi:10.1016/0896-6273(93)90200-b

39. Benzer S. From the gene to behavior. JAMA. 1971;218: 1015–1022.
doi:10.1001/jama.1971.03190200047010

40. Sansone CL, Blumenthal EM. Developmental expression of drop-dead is
required for early adult survival and normal body mass in Drosophila melanogaster.
Insect Biochem Mol Biol. 2012;42: 690–8. doi:10.1016/j.ibmb.2012.06.002

41. Dietzl G, Chen D, Schnorrer F, Su K-C, Barinova Y, Fellner M, et al. A genome-
wide transgenic RNAi library for conditional gene inactivation in Drosophila. Nature.
2007;448: 151–156. doi:10.1038/nature05954

42. Wieschaus E, Nusslein-Volhard C. Looking at embryos. In: Roberts DB, editor.
Drosophila: a practical approach. Oxford: IRL Press; 1986. pp. 199–227.

43. Chou T, Perrimon N. The autosomal FLP-DFS technique for generating germline
mosaics in Drosophila melanogaster. Genetics. 1996;144: 1673–1679.
doi:10.1093/genetics/144.4.1673

44. Page AW, Orr-Weaver TL. Activation of the meiotic divisions in Drosophila
oocytes. Dev Biol. 1997;183: 195–207. doi:10.1006/dbio.1997.8506

45. Reese MG, Eeckman FH, Kulp D, Haussler D. Improved Splice Site Detection in
Genie. J Comput Biol. 1997;4: 311–323. doi:10.1089/cmb.1997.4.311

46. Wu T, Manogaran AL, Beauchamp JM, Waring GL. Drosophila vitelline
membrane assembly: A critical role for an evolutionarily conserved cysteine in the
"VM domain" of sV23. Dev Biol. 2010;347: 360–368.
doi:10.1016/j.ydbio.2010.08.037

47. Kim JY, Jang W, Lee HW, Park E, Kim C. Neurodegeneration of Drosophila
drop-dead mutants is associated with hypoxia in the brain. Genes Brain Behav.
2012;11: 177–184. doi:10.1111/j.1601-183x.2011.00743.x

48. Queenan AM, Ghabrial A, Schüpbach T. Ectopic activation of torpedo/Egfr, a
Drosophila receptor tyrosine kinase, dorsalizes both the eggshell and the embryo.
Development. 1997;124: 3871–3880. doi:10.1242/dev.124.19.3871

# Supporting information

**726**

**727** **S1 File. Protocol for neutral red permeability assay.**

**728**

**729** **S2 File. Raw numerical data.** The six tabs in the spreadsheet contain underlying

**730** data for figure S2, table 1, figure 2, figure 5, table 2, and figure 6, respectively.

**731**

**732** **S1 Fig. Mutation in the *drd¹* allele.** Upper sequence shows the wild-type sequence

**733** of the final five bases of exon 8 (bold), all of intron 8, and the first five bases of exon

**734** 9 (bold). The lower sequence shows the same region in *drd¹*, with the single base

**735** change (green, underlined) and the new start of exon 9 (bold).

**736**

**737** **S2 Fig. Egg-laying by *drd* mutant females.** (A) Fraction of females in each

**738** condition that laid any eggs during their lifetime. Because some non yeast-fed

**739** females were assayed in groups of 2-3, the data were analyzed with two different

**740** assumptions regarding the distribution of egg-laying (see methods). Brackets

**741** indicate the effect of yeast feeding (two-sided Fisher's exact test). (B) Number of

**742** eggs laid per female, omitting data from those females that laid no eggs. Brackets

**743** indicate the effect of yeast feeding (Kruskal-Wallis test with Dunn's post-hoc multiple

**744** comparisons test). n=21-67 females per condition.

**745**

**746** **S3 Fig. Control for *drd* expression in the egg chamber. (**A) Maximum intensity

**747** projection of a stage 14 egg chamber in which expression of a nuclear GFP reporter

**748** is driven by *drd-GAL4*. (B) Control stage 14 egg chamber with the nuclear GFP

**749** reporter but no *drd-GAL4* driver.

**750**

751    **S1_raw_images. Images of Western blots used in figures 3, 4, and 7.**

753 **Figure 1**

756    **Figure 2**

759 **Figure 3**

Cp36

**Figure 4**

**Figure 5**

**Figure 6**

**Figure 7**

**Figure 8**

**Figure S1**

`wild-type`
**TAATT**GTAGGTATTCGCCCAGGTCTAGCATTTATAGAGAATTGTATTTTTTGTTGTTTTTTATTTGCTCAACGCAG**CTCGA**
*drd[1]*
**TAATT**GTAGGTATTCGCCCAGGTCTAGCATTTATAGAGAATTGTATTTTTTGTTGTTTTTTATT<u>A</u>G**CTCAACGCAG**CTCGA**
782

786 **Figure S3**