Long-Term Oral Tamoxifen Administration Decreases Brain Derived

Neurotrophic Factor in the Hippocampus of Female Long-Evans Rats

Laura E. Been[1], Amanda R. Halliday[1], Sarah M. Blossom[1], Elena M. Bien[1], Anya G.

Bernhard[1], Grayson E. Roth[1], Karina I. Domenech Rosario[1], Karlie B. Pollock[1], Petra E.

Abramenko[1], Leily M. Behbehani[1], Gabriel J. Pascal[1], and Mary Ellen Kelly[1,2]

[1] Haverford College, Department of Psychology and Neuroscience Program, Haverford,

PA, USA

[2] University of Pennsylvania, Neuroscience Program, Philadelphia, PA, USA

Abstract

Tamoxifen is a selective estrogen receptor modulator (SERM) that is commonly used as

an adjuvant drug therapy for estrogen receptor-positive breast cancers. While this drug

is effective at reducing the rate of cancer recurrence, many patients report unwanted

cognitive and affective side effects such as brain fog, confusion, memory impairment,

anxiety, and depression. Despite this, the impacts of chronic tamoxifen exposure on the

brain are poorly understood, and rodent models of tamoxifen exposure do not replicate

the chronic oral administration seen in patients. We therefore used long-term *ad lib*

consumption of medicated food pellets in adult female rats to model chronic tamoxifen

exposure in a clinically-relevant way. Gonadally-intact adult female Long-Evans Hooded

rats consumed tamoxifen medicated food pellets for approximately 12 weeks while

control animals received standard chow. At the conclusion of the experiment, animals

were euthanized, and blood and brain samples were collected for analyses. Blood

tamoxifen levels were measured using a novel ultra-performance liquid

chromatography-tandem mass spectrometry assay, which found that this administration

paradigm produced serum levels of tamoxifen similar to those in human patients. In the

brain, brain derived neurotrophic factor (BDNF) was visualized in the hippocampus

using immunohistochemistry and quantified using background-subtracted optical

densitometry. Chronic oral tamoxifen treatment resulted in a decrease in BDNF

expression across several regions of the hippocampus. Together, these findings provide

a novel method of modeling and measuring chronic oral tamoxifen exposure, and

suggest a putative mechanism by which tamoxifen may cause cognitive and behavioral

changes reported by patients.

<u>Introduction</u>

Tamoxifen is a common adjuvant drug therapy used to decrease the recurrence

risk of estrogen receptor-positive breast cancers (Jebahi et al., 2021; Kadakia & Henry,

2015; Novick et al., 2020). The mechanism of action of tamoxifen depends on its

location in body tissue and the local estradiol environment (Novick et al., 2020).

Generally, tamoxifen acts as an antagonist in the presence of endogenous estradiol, but

is agonistic and protects against the effects of estrogen depletion in the absence of

endogenous estradiol (Azizi-Malekabadi et al., 2015; Novick et al., 2020). Thus, it has

been termed a selective estrogen receptor modulator (SERM). Patients typically take

tamoxifen orally for 5-10 years following or concurrent with primary cancer treatment

(Kadakia & Henry, 2015).

While tamoxifen is effective at decreasing the risk of cancer recurrence due to its

antagonistic effects in breast tissue (Kadakia & Henry, 2015), people undergoing

chronic tamoxifen treatment report unwanted psychological side effects such as anxiety,

depression, brain fog, confusion, and memory impairment (Jebahi et al., 2021; Lee et

al., 2016; Novick et al., 2020; Underwood et al., 2018). Long-term tamoxifen therapy

has been shown to decrease performance on visuospatial, visual, and verbal memory

tasks (Boele et al., 2015; Castellon et al., 2004; Lejbak et al., 2010; Underwood et al.,

2018), decision making tasks (X. Chen et al., 2014; Lejbak et al., 2010), and overall

executive function (Boele et al., 2015; Jenkins et al., 2004). However, it has been

difficult to separate the effects of tamoxifen from the effects of primary cancer

treatments, other adjuvant therapies such as aromatase inhibitors (Breckenridge et al.,

2012) and the psychosocial effects of being a cancer survivor (Ahles et al., 2012).

Like in people, tamoxifen treatment in rodents can lead to affective and cognitive

changes; however, the impact of tamoxifen on cognitive-behavioral outcomes varies

depending on the hormone status of the subjects (Novick et al., 2020). For example,

tamoxifen administration to gonad-intact female rodents has been shown to increase

anxiety- and depressive-like behaviors (Azizi-Malekabadi et al., 2015), decrease

memory consolidation and retrieval ability on step-down avoidance tasks (D. Chen et

al., 2002), and decrease novel object recognition ability (Valvassori et al., 2017). In

ovariectomized rodents, on the other hand, tamoxifen administration is protective

against anxiety- and depressive-like behaviors (Azizi-Malekabadi et al., 2015) and

improves working memory ability (Velázquez-Zamora et al., 2012). Because tamoxifen

is more commonly used in pre-menopausal patients (Kadakia & Henry, 2015), its effects

on gonadally-intact female rodents are of particular interest. Notably, previous rodent

studies have not modeled chronic oral tamoxifen use and instead have used a variety of

tamoxifen administration methods including long-term injections (Cechinel-Recco et al.,

2012), short-term or one-time injections (D. Chen et al., 2002; Kight & McCarthy, 2017;

Velázquez-Zamora et al., 2012), and short-term gavages (Valvassori et al., 2017).

The hippocampus is a strong putative candidate to mediate some of the cognitive

side effects of tamoxifen therapy. First, the hippocampus expresses □- and β-estrogen

receptors (ER□ and ERβ, respectively) across its various subregions (Frick et al., 2015;

Harte-Hargrove et al., 2013; Spencer et al., 2008; Woolley, 1998). Research in rodents

has demonstrated that estradiol fluctuations impact hippocampal neurophysiology (Liu

et al., 2008; Woolley, 1998), which in turn affects depressive- and anxiety-like behaviors

(Green & Galea, 2008; Walf & Frye, 2006) as well as performance on spatial memory

tasks (Frick et al., 2015; Spencer et al., 2008). Within hippocampal neurons, Brain-

Derived Neurotrophic Factor (BDNF) may play an important role in mediating

tamoxifen's effects on cognition and emotion. Estrogen receptor activation influences

BDNF transcription via estrogen response elements (EREs) that bind to the *BDNF* gene

(Harte-Hargrove et al., 2013; Scharfman & MacLusky, 2006; Sohrabji & Lewis, 2006;

Spencer et al., 2008). Indeed, fluctuating levels of estradiol during the estrous cycle

lead to fluctuating levels of BDNF expression in the hippocampus (Harte-Hargrove et

al., 2013; Luine & Frankfurt, 2013; Scharfman & MacLusky, 2014). Finally, estradiol has

been shown to mediate both synaptogenesis (Sato et al., 2007) and LTP (Spencer et

al., 2008) through its interactions with BDNF.

Taken together, this evidence suggests tamoxifen's impact on cognition and

emotion may be mediated by BDNF expression in the hippocampus. We therefore

hypothesized that long-term oral tamoxifen administration would decrease BDNF

expression in the hippocampus. To test this hypothesis, we established a model of long-

term oral tamoxifen administration in adult, intact female Long-Evans rats. Using a novel

ultra-performance liquid chromatography-tandem mass spectrometry (LC-MS/MS)

detection method, we found that this long-term oral administration produced serum

levels of tamoxifen comparable to those from Tamoxifen administration in people

(Farrar & Jacobs, 2022). Brain tissue analyses found that tamoxifen-treated animals

had significantly lower BDNF expression in the DG, medial CA3, and CA1. These

results provide an important first step in characterizing the effect of long-term oral

tamoxifen administration on the hippocampus, and may increase our understanding of

the cognitive and affective side effects associated with long-term tamoxifen use.

Methods

Subjects

For all experiments, adult female Long-Evans Hooded rats were purchased from

Charles River Laboratories (Wilmington, Massachusetts) at 3-4 months of age. Animals

were housed in pairs in standard cages (17.7 x 9.4 in x 8.26 in; Allentown LLC,

Allentown, NJ, USA) containing standard TekFresh cellulose low-dust rat bedding

(Teklad 7099; Inotiv, West Lafayette, IN, USA), and a Plexiglas tube (Bio-Serv,

Flemington, NJ, USA) and wooden block (Bio-Serv, Flemington, NJ, USA) for

enrichment. Subjects were kept on a 12-hour light-dark cycle with *ad libitum* food and

water access.

Tamoxifen Self-Administration

Animals were divided into two experimental groups: tamoxifen (n = 20) and

control (n = 16). Animals in the tamoxifen group consumed tamoxifen *ad libitum* for 10-

13 weeks via medicated food pellets (Envigo, Indianapolis, IN) that were custom

developed to simulate the standard dose of tamoxifen prescribed to premenopausal

women diagnosed with breast cancer (20 mg/kg daily) while control animals received an

ingredient-matched rodent feed (16% protein, 55% carb, 3.4% fat) that did not contain

Tamoxifen (Envigo, Farrar & Jacobs, 2022). The composition of the feed was identical

between the two groups except for the addition of tamoxifen, red food coloring, and

sucrose to enhance palatability in the medicated food pellets (**Table 1**). Importantly, the

base pellet (2016, Teklad Global 16% Protein Rodent diet) to which the tamoxifen and

associated ingredients were added, and that was fed ad-lib to control rats, did not

contain alfalfa or soybean meal, thus minimizing the occurrence of phytoestrogens in control rats during the time they were housed at Haverford College.

Body weight was monitored for all animals, regardless of experimental group, throughout the drug administration period. Rats that lost more than 15% of their starting body weight were presented with food pellets mashed in a flavor-enhanced, nutritionally-fortified gel (DietGel Recovery, Clear H20, Portland, ME, USA) in addition to *ad libitum* access to chow to further enhance palatability and encourage ingestion until weight stabilized. Starting at six weeks into the experiment, all rats in the tamoxifen group were presented with this mash daily to maintain adequate health, while also ensuring adequate consumption of Tamoxifen. As an additional check on animal health, starting at 6 weeks, body condition was also assessed daily with the Body Condition Scale (see **Supplementary Figure 1**). All rats in the Tamoxifen group were assigned scores of 3 (well-conditioned) or above. Final weight checks were taken prior to euthanization.

Blood Analysis

Following 10-13 weeks of access to medicated or control pellets, animals were euthanized via intracardial perfusion (see below). Blood samples were collected from the inferior vena cava immediately prior to perfusion and stored in heparin-coated vacutainer blood collection tubes (BD Medical, Franklin Lakes, NJ, USA) on ice until centrifugation. Samples were centrifuged at 2300 g at 4°C for 20 min and serum was stored at -80 until assay.

An ultra-performance liquid chromatography-tandem mass spectrometry assay

with reversed-phase chromatographic separation employing a Waters XBridge C18

(100 x 2.1 mm, 100 Å, 3.5 µm) and a runtime of 4.5 minutes was used to quantify

tamoxifen levels. Rat plasma samples (100 µL) were extracted with acetonitrile

containing tamoxifen-d5 (5 ng/mL) and 0.1% formic acid. Two µL of the extract were

injected onto the UPLC-MS/MS system for analysis.

Tamoxifen and tamoxifen-d5 were separated by ultra-performance liquid

chromatography (UPLC) and detected using a triple quadrupole mass spectrometer

(API4000). Tandem mass spectrometery (MS-MS) parameters were optimized in the

positive ionization mode and multiple reaction monitoring (MRM) transition of m/z 372.3

$\rightarrow$ 72.0 was used for tamoxifen analysis. Tamoxifen-d5 (m/z 377.3 $\rightarrow$ 72.0) was used as

an internal standard. A calibration curve for tamoxifen was prepared in human plasma

over the linear range of 0.1 $\rightarrow$ 250 ng/mL (r^2 > 0.99, Figure 2, Table 2). Calibration

standards and blanks in human plasma were extracted by protein precipitation with 400

µL of 5 ng/mL of tamoxifen-d5 in acetonitrile. Five µL of extract was injected for LC-

MS/MS analysis. Tamoxifen and tamoxifen-d5 were separated on an XBridge C18

column (100 x 2.1 mm, 100 Å, 3.5 µm) using 0.1 % formic acid in water and 0.1 %

formic acid in acetonitrile as aqueous and organic mobile phases with a flow rate of 0.5

mL/min. Analytes were separated using a gradient elution, with retention time of 2.68

min for tamoxifen and tamoxifen-d5. The LC-MS Tamoxifen assay used here was based

on assays initially developed and validated in human plasma (Bobin-Dubigeon et al.,

2019; Rama Raju et al., 2015). Therefore, as a confirmatory assay, human plasma and

rat plasma were compared using 100 ng/mL concentrations of tamoxifen and found to

be comparable (**Table 3**).

Tissue Histology and Immunohistochemistry

At the conclusion of the experiment, subjects were sacrificed by intracardial

perfusion. Briefly, subjects were given an overdose of sodium pentobarbital

(Beuthanasia-D, 22 mg/100 g body weight, Merck Animal Health, Madison, NJ, USA)

and then transcardially perfused with approximately 250 mL of 25mM phosphate

buffered saline (PBS) followed by approximately 250 mL of 4% paraformaldehyde in

PBS. Brains were immediately extracted and post-fixed in 4% paraformaldehyde

overnight and then cryoprotected for 48 h in 30% sucrose in PBS. Coronal sections (35

μm) were sectioned at -20°C on a cryostat (Leica Biosystems, Wetzlar, Germany),

collected in a 1:6 series, and stored in cryoprotectant until immunohistochemical

processing.

For immunohistochemistry, hippocampal sections were removed from cryoprotectant

and rinsed 5 x 5 minutes in 25 mM PBS. After PBS washes, sections were incubated in

hydrogen peroxide (1:100) for 15 minutes to reduce endogenous peroxidase activity.

Sections were then washed 5 x 5 minutes in 25 mM PBS and before incubation in a

rabbit monoclonal primary antibody against brain-derived neurotrophic factor (1:3000,

Cat # ANT-010, Alomone Labs, Jerusalem, Israel) in 0.4% Triton-X 100 in PBS for 24

hours at room temperature. This antibody has previously been validated for use in

rodent tissue using inducible BDNF knockouts (Sun et al., 2018) and shBDNF treated

rats (Taliaz et al., 2010). This antibody recognizes both mature BNDF and pro-BDNF

(Taliaz et al., 2010).

After incubation in the primary antibody, sections were rinsed in PBS (5 x 5 minutes)

and incubated for one hour in a biotinylated secondary antibody (1:600, Biotin-SP

AffiniPure Goat Anti-Rabbit IgG, Jackson Immunoresearch Labs; 111-065-003) in 0.4%

Triton-X 100 in PBS. Sections were then rinsed in PBS (5 x 5 minutes) and incubated in

avidin-biotin complex with horseradish peroxidase (per manufacturer's instructions,

Vectastain Elite ABC-HRP Kit, Vector Laboratories, Burlingame, CA, USA). After 4 x 5

minute washes in PBS, sections were washed 2 x 5 minutes in a sodium acetate buffer

and then incubated for 10 minutes in 3,3'-diaminobenzidine HCl (0.2 mg/mL) in a nickel

sodium acetate buffer (0.025 g Nickel II Sulfate/mL NaOAc) and hydrogen peroxide

(0.83 µL/mL) solution.

Stained tissue was then washed in sodium acetate buffer (2 x 5 min), washed in

PBS (3 x 5 min) mounted on glass slides (SuperFrost Plus Microscope Slides,

ThermoFisher Scientific, Waltham, MA, USA) and left to air-dry overnight. After drying,

slides were dehydrated with alcohols, cleared with xylenes and coverslipped with

Permount (ThermoFisher Scientific, Waltham, MA, USA).

BDNF Densitometry Analysis in Hippocampus

Images were acquired at 10X magnification using a Nikon E200 brightfield

microscope with a color camera and Spot Basic software (Diagnostic Instruments,

Sterling Heights, MI, USA). Five evenly-spaced template images (S1, S2, S3, S4, S5)

were selected from the rat brain atlas (Paxinos & Watson, 2013) to capture the rostral to

caudal extent of the hippocampus. The CA3 was split into medial and lateral regions for

imaging and analysis in order to capture it fully. Twenty images were collected for each

animal: five images per hippocampus region (DG, CA1, mCA3, lCA3).

Densitometry analyses were performed by experimenters who were blind to the

condition of the subject using ImageJ software (imagej.net/software/fiji; Schindelin et al.,

2012). Mean values for background optical density were collected from 36x36 pixel

regions that did not contain BDNF-ir cells using the "mean gray value" tool. Mean optical

density values for regions of interest (ROIs) were calculated after tracing the

hippocampus region of interest with the polygon tool (**Figure 3**) and analyzing with the

"mean gray value" tool (**Figure 4**). ROIs were checked for consistency across animals

by an experimenter blind to the condition of the subjects. Background optical density

was subtracted from ROI optical density using the following equation: $(OD) = \log (255 /$

$(OD_{ROI} - OD_{background}))$. Because background levels were variable across sections, likely

due to individual differences in perfusions, we background-subtracted OD values

between sections with high background levels versus low background levels and found

no significant difference between groups. This suggests that: 1) our method was

effective at subtracting background and 2) differences between treatment groups were

not due to differences in background.

Statistical Analysis

All statistics were run using Jamovi (www.jamovi.org) with alpha values of 0.05.

Assumptions of normality were tested using the Shapiro-Wilk normality test. Repeated

measures factorial ANOVAs were used to look for interactions and main effects of drug

condition (tamoxifen vs. control) and time (week) on body weight. Significant effects

were explored using Bonferroni's multiple comparisons tests and the Geisser and

Greenhouse method was used to correct violations of sphericity. Two-way ANOVAs were

used to look for interactions and main effects of drug condition (tamoxifen vs. control)

and section level (1-5) on BDNF-ir density within each of the four ROIs (DG, CA1,

vCA1, and CA3). Individual brain regions were analyzed separately. Significant effects

were explicated using Fisher's LSD post-hoc comparisons. Individual samples were

excluded from analysis if tissue damage precluded OD analysis, or if there was no

adequate match for the template atlas plate. The relationship between serum tamoxifen

levels and BDNF-ir density was examined using simple correlations.

7
8 **Results**

9
Body Weight

There was a significant interaction between time and drug condition on body

weight (F(11,99 = 10.65, p < 0.0001). Post-hoc tests revealed that starting at Week 3,

Tamoxifen-fed rats weighed significantly more than control rats for the duration of the

experiment (all p < 0.001, **Figure 1**).

Plasma Tamoxifen Levels

Tamoxifen was not found in any control
samples. In tamoxifen-treated animals,
plasma concentrations ranged from 10.7
to 48.3 ng/mL (M = 28.1, SD = 11.2,
Table 4). Representative
chromatograms can be found in Figure
2.

BDNF Immunoreactivity
9
For each brain region of interest, the density of BDNF-immunoreactive (ir)

neurons in the DG, mCA3, lCA3, and CA1 was analyzed and compared between

groups. In the DG, there was a significant main effect of drug condition on the density of

BDNF-ir cells (F(1, 146) = 9.454, p = 0.003). Post-hoc tests revealed that BDNF-ir

density was lower in tamoxifen-treated animals than control animals in Section 4 (p =

0.03, **Figure 4A**). In CA1, there was a main effect of drug condition on the density of

BDNF-ir cells (F(1, 146) = 8.886, p = 0.003). Post hoc tests revealed that BDNF-ir

density was lower in tamoxifen-treated animals than control animals in section 2 (p =

0.04) and section 4 (p = 0.02, **Figure 4B**). In mCA3, there was a significant main effect

of drug condition on the density of BDNF-ir cells (F, 1, 144) = 11.94, p < 0.001). Post-

hoc tests revealed that BDNF-ir was lower in tamoxifen-treated animals than control

animals in Section 3 (p < 0.01, **Figure 4C**). There was also a significant main effect of

section on the density of BDNF-ir cells in mCA3 (F(4, 144) = 3.302, p = 0.01). Post-hoc

tests revealed that Section 2 differed from Section 4 (p = 0.02) and Section 5 (p = 0.03),

and Section 3 differed from Section 4 (p < 0.01) and Section 5 (p < 0.01). Finally, in

lCA3, there was no effect of drug condition (F(1, 145) = 0.071, p = 0.79) or section

(F(1,145) = 0.87, p = 0.48) on the density of BDNF-ir cells (**Figure 4D**).

Animals' plasma tamoxifen concentration at the time of sacrifice did not correlate

significantly with BDNF-ir cells in the DG (r = -.4221, p = .2244, **Figure 5A**) or CA1 (r = -

.2053, p = .5693, **Figure 5B**). In both medial and lateral CA3 (**Figure 5C and 5D**), there

was a trend towards increased plasma tamoxifen concentration being associated with

decreased BDNF-ir cells, although these did not reach significance (mCA3 r = -.6112, p

= .0604; lCA3 r = -.6168, p = .0575).

Discussion

Here we demonstrate for the first time that long-term oral tamoxifen

administration decreases BDNF expression in the hippocampus of adult female Long-

Evans Hooded rats. Notably, we show that a novel administration method, long-term *ad*

lib exposure to medicated food pellets, produces plasma tamoxifen levels that are

similar to a human dose (20 mg/kg; Farrar & Jacobs, 2022). Further, we used a novel

LC-MS/MS assay to detect tamoxifen levels in rat plasma. Together, these results

provide a new way to model and measure the effects of long-term oral tamoxifen

administration on the brain, which may be particularly useful for investigating

mechanisms of cognitive and behavioral changes reported by patients taking

Tamoxifen.

Voluntary oral administration via medicated food pellets is a clinically relevant,

non-invasive, and effective method for modeling the effects of tamoxifen in rodents.

Tamoxifen can readily cross the blood-brain-barrier (Lien et al., 1991; Pareto et al.,

2004), and oral self-administration is an attractive alternative to systemic injections

(Cechinel-Recco et al., 2012; D. Chen et al., 2002; Kight & McCarthy, 2017; Velázquez-

Zamora et al., 2012) or oral gavage (Valvassori et al., 2017) in that it more closely

models the metabolism of tamoxifen use in patient populations and eliminates stress

associated with restraint, repeated injections, or gavage as potential confounding

variables. Nonetheless, there are some challenges with this method. First, there were

likely differences in palatability between tamoxifen and control chow. Tamoxifen-fed

animals initially lost weight and required additional intervention (food pellets mashed in

flavor-enhanced gel) to prevent further weight loss, whereas control animals gained

weight across the course of the experiment. A similar study in mice found that 15 days

of exposure to tamoxifen chow also led to weight loss, particularly in female subjects,

and animals did not consistently eat Tamoxifen chow for the first 5 days of exposure

(Smith et al., 2022). However, an anorexic effect of tamoxifen itself cannot be ruled out,

as tamoxifen injections induce anorexia in rodents (Lampert et al., 2013; Gray et al.,

1993; López et al., 2006; Wade & Heller, 1993; Walker et al., 2011, p. 2006). While oral

gavage would introduce stress and is therefore not recommended, future experiments

may wish to explore mild food deprivation, pair feeding tamoxifen-treated and control

animals, or giving a daily oral dose of tamoxifen to eliminate the effects of feeding

differences and/or body weight as potential confounds on dependent variables. There

was also variability in individual animals' plasma tamoxifen concentrations. These

differences likely resulted from differences in voluntary consumption of tamoxifen chow

across animals, both throughout the experiment and specifically on the day of

euthanasia. Blood draws were collected approximately 60 minutes after the last

tamoxifen pellets were presented, but the amount of medicated pellet consumed, or the

timing of consumption within the one-hour window, were not measured. Future studies

should collect data about volume and timing of food intake, particularly prior to blood

collection, as well as additional bioassays that correlate with tamoxifen consumption

such as uterine weight (Fong et al., 2007).

Our finding that long-term oral tamoxifen administration decreased BDNF

expression in the hippocampus may point towards a putative mechanism underlying

cognitive side effects that patients taking tamoxifen report. Given the known association

of estrogen receptor activation and BDNF transcription (Harte-Hargrove et al., 2013;

Scharfman & MacLusky, 2006; Sohrabji & Lewis, 2006), it is possible that tamoxifen's

action at estrogen receptors in the brain could disrupt BDNF transcription, altering

plasticity and neuronal stability in the hippocampus. These changes to hippocampal

function could in turn impair cognitive function and lead to clinical symptoms like brain

fog, memory deficits, and alterations in mood/anxiety. In support of this idea, Smith et

al. (2022) found that short term Tamoxifen administration impacts hippocampal

neurogenesis. Specifically Tamoxifen injection is associated with decreased progenitor

cell proliferation in the dentate gyrus, although Tamoxifen ingestion is associated with

increased neuronal differentiation in the dentate gyrus. Thus, our finding that long-term

oral tamoxifen administration decreased BDNF density in the current study may be due

to a decrease in cell proliferation, or conversely, decreases in BDNF following

Tamoxifen treatment could contribute to changes in neurogenesis.

Brain BDNF is difficult to visualize due to its relatively low expression in mature

neurons. In the current study, we used an immunohistochemistry approach with a

nickel-enhanced chromogen reaction to amplify visualization. This method produces robust BDNF staining, but can produce higher levels of background staining, necessitating a background-subtracted densitometry approach to quantification rather than individual cell counts. Further, BDNF staining patterns differ across published experiments, likely due to differences in antibodies. Here, we report BDNF expression in the granular cell layer is reduced following long-term oral Tamoxifen administration, but found little BDNF expression in the mossy fiber layer. This matches the expression pattern previously shown with this antibody (Taliaz et al., 2010). However, previous papers that used different antibodies report BDNF staining in the mossy fiber layer, but only sparse staining in the granule cell layer (Conner et al., 1997; Dieni et al., 2012). It is possible that different antibodies label distinct pools of BDNF based on differences in the targeted epitope. The method of visualization and antibody choice are therefore important considerations when visualizing and quantifying BDNF in the brain.

Future studies using this model of long-term oral tamoxifen administration should include behavioral assays of cognition, mood, and anxiety. Our finding that long-term tamoxifen administration decreased BDNF in the DG, CA1, and medial CA3 suggests that pattern separation, spatial memory, associative memory, and episodic memory processes could be particularly affected (Bartsch et al., 2011; Cherubini & Miles, 2015; Kesner, 2007; Langston et al., 2010). Disparate findings in CA3 subregions further point towards tamoxifen's effects on cognitive function. For example, there is some literature to suggest that the dorsal CA3 is responsible for cognition and information processing, while the ventral region is more closely associated with stress and affect (Fanselow & Dong, 2010). A full behavioral profile of animals during long-term oral tamoxifen

treatment, such as further assays of cognition (e.g., object-in-place, attentional set

shifting), anxiety-like behaviors (e.g., defensive burying, light-dark box) and learned

helplessness or anhedonia (e.g., forced swim, tail suspension, splash test, sucrose

preference) would be useful to characterize tamoxifen's effects on behavior in this

model, and may suggest additional brain regions for exploration. Further, while this

study focused on the hippocampus, it is likely that tamoxifen impacts other estrogen-

sensitive brain regions that are associated with cognition, anxiety, and/or mood

disorders, such as the nucleus accumbens and amygdala. Tamoxifen could disrupt

BDNF transcription in these regions by interacting with estrogen receptors via the same

putative mechanism that we have identified in the hippocampus. The relationship

between Tamoxifen, hippocampal BDNF levels, and body weight also requires further

investigation, as BDNF is known to regulate energy balance via action in other parts of

the brain (for review see Xu & Xie, 2016).

In addition to a novel administration paradigm, we also used a novel, sensitive,

and reproducible method for the analysis of tamoxifen that is suitable for rat plasma.

This will be useful to future researchers interested in the physiological effects of oral

tamoxifen administration. While not significant, we did observe a trend towards

correlation between plasma tamoxifen concentration and BDNF density in mCA3 and

lCA3, supporting tamoxifen's potential role in decreasing BDNF transcription in the

hippocampus. It is worth noting that tamoxifen is also commonly used in neuroscience

research as a tool to induce Cre mechanisms. While these research applications are

markedly different in that the doses are generally much higher and administration is

acute (often a single injected dose) (Donocoff et al., 2020; Heffner, 2011) our research

suggests that tamoxifen administration affects BDNF expression, and researchers using

tamoxifen in their work should consider its potential off-target effects on neurobiology.

Tamoxifen remains the gold-standard medication for preventing recurrence of

estrogen receptor-positive breast cancers. Its ongoing use as an adjuvant therapy will

likely continue despite common affective and cognitive side effects; tamoxifen

treatments will enable breast cancer survivors to live longer and healthier lives. Further

identifying and understanding the neural mechanisms underlying these side effects,

however, could lead to promising interventions that would mitigate unwanted

psychological changes without risking cancer recurrence. It is important that

researchers and clinicians continue to explore ways to improve psychiatric health in at-

risk and under-researched patient populations. Cancer survivors, women and other

gender minorities in particular, are particularly vulnerable to a range of affective,

anxiety, and cognitive disorders (Ahles et al., 2012; Di Giacomo et al., 2016; Fafouti et

al., 2010). It is imperative that we find ways to support all aspects of their health as they

navigate treatment.

Acknowledgements

This work was funded in part by Haverford College Department of Psychology. The

authors wish to thank Shanice Edwards, Christina Vedar, Abigail Cohen, Athena Zuppa,

and Ganesh Moorthy from the Children's Hospital of Philadelphia Research Institute

Bioanalytical Core for developing and conducting the mass spectrometry assays. Some

of the mass spectrometry data were previously presented as a poster at the American

Society for Mass Spectrometry Conference on Mass Spectrometry and Allied Topics in

Philadelphia, PA in November of 2021. The authors also wish to thank Dr. Louise

Charkoudian for her assistance interpreting the LC-MS data.

References

Ahles, T. A., Root, J. C., & Ryan, E. L. (2012). Cancer- and Cancer Treatment–
Associated Cognitive Change: An Update on the State of the Science. *Journal of Clinical Oncology*, *30*(30), 3675–3686. https://doi.org/10.1200/JCO.2012.43.0116

Azizi-Malekabadi, H., Pourganji, M., Zabihi, H., Saeedjalali, M., & Hosseini, M. (2015). Tamoxifen antagonizes the effects of ovarian hormones to induce anxiety and depression-like behavior in rats. *Arquivos de Neuro-Psiquiatria*, *73*, 132–139. https://doi.org/10.1590/0004-282X20140221

Bartsch, T., Döhring, J., Rohr, A., Jansen, O., & Deuschl, G. (2011). CA1 neurons in the human hippocampus are critical for autobiographical memory, mental time travel, and autonoetic consciousness. *Proceedings of the National Academy of Sciences*, *108*(42), 17562–17567. https://doi.org/10.1073/pnas.1110266108

Bobin-Dubigeon, C., Campone, M., Rossignol, E., Salaun, E., Amiand, M.-B., & Bard, J.-M. (2019). New UPLC–MS/MS assay for the determination of tamoxifen and its metabolites in human plasma, application to patients. *Future Science OA*, *5*(5), FSO374. https://doi.org/10.2144/fsoa-2018-0113

Boele, F. W., Schilder, C. M. T., de Roode, M.-L., Deijen, J. B., & Schagen, S. B. (2015). Cognitive functioning during long-term tamoxifen treatment in postmenopausal women with breast cancer. *Menopause*, *22*(1), 17–25. https://doi.org/10.1097/GME.0000000000000271

Breckenridge, L. M., Bruns, G. L., Todd, B. L., & Feuerstein, M. (2012). Cognitive limitations associated with tamoxifen and aromatase inhibitors in employed breast cancer survivors. *Psycho-Oncology*, *21*(1), 43–53.

https://doi.org/10.1002/pon.1860

Castellon, S. A., Ganz, P. A., Bower, J. E., Petersen, L., Abraham, L., & Greendale, G. A. (2004). Neurocognitive Performance in Breast Cancer Survivors Exposed to Adjuvant Chemotherapy and Tamoxifen. *Journal of Clinical and Experimental Neuropsychology*, *26*(7), 955–969. https://doi.org/10.1080/13803390490510905

Cechinel-Recco, K., Valvassori, S. S., Varela, R. B., Resende, W. R., Arent, C. O., Vitto, M. F., Luz, G., de Souza, C. T., & Quevedo, J. (2012). Lithium and tamoxifen modulate cellular plasticity cascades in animal model of mania. *Journal of Psychopharmacology*, *26*(12), 1594–1604. https://doi.org/10.1177/0269881112463124

Chen, D., Wu, C. F., Shi, B., & Xu, Y. M. (2002). Tamoxifen and toremifene cause impairment of learning and memory function in mice. *Pharmacology Biochemistry and Behavior*, *71*(1), 269–276. https://doi.org/10.1016/S0091-3057(01)00656-6

Chen, X., Li, J., Chen, J., Li, D., Ye, R., Zhang, J., Zhu, C., Tian, Y., & Wang, K. (2014). Decision-making impairments in breast cancer patients treated with tamoxifen. *Hormones and Behavior*, *66*(2), 449–456. https://doi.org/10.1016/j.yhbeh.2014.07.005

Cherubini, E., & Miles, R. (2015). The CA3 region of the hippocampus: How is it? what is it for? how does it do it? *Frontiers in Cellular Neuroscience*, *9*. https://www.frontiersin.org/articles/10.3389/fncel.2015.00019

Conner, J. M., Lauterborn, J. C., Yan, Q., Gall, C. M., & Varon, S. (1997). Distribution of Brain-Derived Neurotrophic Factor (BDNF) Protein and mRNA in the Normal Adult Rat CNS: Evidence for Anterograde Axonal Transport. *The Journal of*

Neuroscience, *17*(7), 2295–2313. https://doi.org/10.1523/JNEUROSCI.17-07-02295.1997

Dieni, S., Matsumoto, T., Dekkers, M., Rauskolb, S., Ionescu, M. S., Deogracias, R., Gundelfinger, E. D., Kojima, M., Nestel, S., Frotscher, M., & Barde, Y.-A. (2012). BDNF and its pro-peptide are stored in presynaptic dense core vesicles in brain neurons. *Journal of Cell Biology*, *196*(6), 775–788. https://doi.org/10.1083/jcb.201201038

Di Giacomo, D., Cannita, K., Ranieri, J., V, C., Passafiume, D., & Ficorella, C. (2016). Breast cancer and psychological resilience among young women. *Journal of Psychopatology*, *3*, 1–5.

Donocoff, R. S., Teteloshvili, N., Chung, H., Shoulson, R., & Creusot, R. J. (2020). Optimization of tamoxifen-induced Cre activity and its effect on immune cell populations. *Scientific Reports*, *10*(1), Article 1. https://doi.org/10.1038/s41598-020-72179-0

Fafouti, M., Paparrigopoulos, T., Zervas, Y., Rabavilas, A., Malamos, N., Liappas, I., & Tzavara, C. (2010). Depression, Anxiety and General Psychopathology in Breast Cancer Patients: A Cross-sectional Control Study. *In Vivo*, *24*(5), 803–810.

Fanselow, M. S., & Dong, H.-W. (2010). Are The Dorsal and Ventral Hippocampus functionally distinct structures? *Neuron*, *65*(1), 7. https://doi.org/10.1016/j.neuron.2009.11.031

Farrar, M. C., & Jacobs, T. F. (2022). Tamoxifen. In *StatPearls*. StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK532905/

Fong, C. J., Burgoon, L. D., Williams, K. J., Forgacs, A. L., & Zacharewski, T. R. (2007).

Comparative temporal and dose-dependent morphological and transcriptional

uterine effects elicited by tamoxifen and ethynylestradiol in immature,

ovariectomized mice. *BMC Genomics*, *8*(1), 151. https://doi.org/10.1186/1471-

2164-8-151

Frick, K. M., Kim, J., Tuscher, J. J., & Fortress, A. M. (2015). Sex steroid hormones

matter for learning and memory: Estrogenic regulation of hippocampal function in

male and female rodents. *Learning & Memory*, *22*(9), 472–493.

https://doi.org/10.1101/lm.037267.114

Gray, J. M., Schrock, S., & Bishop, M. (1993). Estrogens and antiestrogens: Actions

and interactions with fluphenazine on food intake and body weight in rats.

American Journal of Physiology-Regulatory, Integrative and Comparative

Physiology, *264*(6), R1214–R1218.

https://doi.org/10.1152/ajpregu.1993.264.6.R1214

Green, A. D., & Galea, L. A. M. (2008). Adult hippocampal cell proliferation is

suppressed with estrogen withdrawal after a hormone-simulated pregnancy.

Hormones and Behavior, *54*(1), 203–211.

https://doi.org/10.1016/j.yhbeh.2008.02.023

Harte-Hargrove, L. C., MacLusky, N. J., & Scharfman, H. E. (2013). Brain-derived

neurotrophic factor–estrogen interactions in the hippocampal mossy fiber

pathway: Implications for normal brain function and disease. *Neuroscience*, *239*,

46–66. https://doi.org/10.1016/j.neuroscience.2012.12.029

Heffner, C. (2011, July 25). *Intraperitoneal Injection of Tamoxifen for Inducible Cre-*

Driver Lines. The Jackson Laboratory. https://www.jax.org/research-and-faculty/resources/cre-repository/tamoxifen

Jebahi, F., Sharma, S., Bloss, J. E., & Wright, H. H. (2021). Effects of tamoxifen on cognition and language in women with breast cancer: A systematic search and a scoping review. *Psycho-Oncology*, *30*(8), 1262–1277. https://doi.org/10.1002/pon.5696

Jenkins, V., Shilling, V., Fallowfield, L., Howell, A., & Hutton, S. (2004). Does hormone therapy for the treatment of breast cancer have a detrimental effect on memory and cognition? A pilot study. *Psycho-Oncology*, *13*(1), 61–66. https://doi.org/10.1002/pon.709

Kadakia, K. C., & Henry, N. L. (2015). Adjuvant Endocrine Therapy in Premenopausal Women with Breast Cancer. *Clinical Advances in Hematology & Oncology : H&O*, *13*(10), 663–672.

Kesner, R. P. (2007). A behavioral analysis of dentate gyrus function. In H. E. Scharfman (Ed.), *Progress in Brain Research* (Vol. 163, pp. 567–576). Elsevier. https://doi.org/10.1016/S0079-6123(07)63030-1

Kight, K. E., & McCarthy, M. M. (2017). Sex differences and estrogen regulation of BDNF gene expression, but not propeptide content, in the developing hippocampus. *Journal of Neuroscience Research*, *95*(1–2), 345–354. https://doi.org/10.1002/jnr.23920

Lampert, C., Arcego, D. M., Laureano, D. P., Diehl, L. A., Da Costa Lima, I. F., Krolow, R., Pettenuzzo, L. F., Dalmaz, C., & Vendite, D. (2013). Effect of chronic administration of tamoxifen and/or estradiol on feeding behavior, palatable food

and metabolic parameters in ovariectomized rats. *Physiology & Behavior*, *119*, 17–24. https://doi.org/10.1016/j.physbeh.2013.05.026

Langston, R. F., Stevenson, C. H., Wilson, C. L., Saunders, I., & Wood, E. R. (2010). The role of hippocampal subregions in memory for stimulus associations. *Behavioural Brain Research*, *215*(2), 275–291. https://doi.org/10.1016/j.bbr.2010.07.006

Lee, P. E., Tierney, M. C., Wu, W., Pritchard, K. I., & Rochon, P. A. (2016). Endocrine treatment-associated cognitive impairment in breast cancer survivors: Evidence from published studies. *Breast Cancer Research and Treatment*, *158*(3), 407–420. https://doi.org/10.1007/s10549-016-3906-9

Lejbak, L., Vrbancic, M., & Crossley, M. (2010). Endocrine therapy is associated with low performance on some estrogen-sensitive cognitive tasks in postmenopausal women with breast cancer. *Journal of Clinical and Experimental Neuropsychology*, *32*(8), 836–846. https://doi.org/10.1080/13803391003596389

Lien, E. A., Solheim, E., & Ueland, P. M. (1991). Distribution of tamoxifen and its metabolites in rat and human tissues during steady-state treatment. *Cancer Research*, *51*(18), 4837–4844.

Liu, F., Day, M., Muñiz, L. C., Bitran, D., Arias, R., Revilla-Sanchez, R., Grauer, S., Zhang, G., Kelley, C., Pulito, V., Sung, A., Mervis, R. F., Navarra, R., Hirst, W. D., Reinhart, P. H., Marquis, K. L., Moss, S. J., Pangalos, M. N., & Brandon, N. J. (2008). Activation of estrogen receptor-β regulates hippocampal synaptic plasticity and improves memory. *Nature Neuroscience*, *11*(3), Article 3. https://doi.org/10.1038/nn2057

López, M., Lelliott, C. J., Tovar, S., Kimber, W., Gallego, R., Virtue, S., Blount, M., Vázquez, M. J., Finer, N., Powles, T. J., O'Rahilly, S., Saha, A. K., Diéguez, C., & Vidal-Puig, A. J. (2006). Tamoxifen-Induced Anorexia Is Associated With Fatty Acid Synthase Inhibition in the Ventromedial Nucleus of the Hypothalamus and Accumulation of Malonyl-CoA. *Diabetes*, *55*(5), 1327–1336. https://doi.org/10.2337/db05-1356

Luine, V., & Frankfurt, M. (2013). Interactions between estradiol, BDNF and dendritic spines in promoting memory. *Neuroscience*, *239*, 34–45. https://doi.org/10.1016/j.neuroscience.2012.10.019

Novick, A. M., Scott, A. T., Epperson, C. N., & Schneck, C. D. (2020). Neuropsychiatric Effects of Tamoxifen: Challenges and Opportunities. *Frontiers in Neuroendocrinology*, *59*, 100869. https://doi.org/10.1016/j.yfrne.2020.100869

Pareto, D., Alvarado, M., Hanrahan, S. M., & Biegon, A. (2004). In vivo occupancy of female rat brain estrogen receptors by 17beta-estradiol and tamoxifen. *NeuroImage*, *23*(3), 1161–1167. https://doi.org/10.1016/j.neuroimage.2004.07.036

Paxinos, G., & Watson, C. (2013). *The Rat Brain in Stereotaxic Coordinates* (7th ed.). Academic Press. https://www.elsevier.com/books/the-rat-brain-in-stereotaxic-coordinates/paxinos/978-0-12-391949-6

Rama Raju, K. S., Taneja, I., Singh, S. P., Tripathi, A., Mishra, D. P., Hussain, K. M., Gayen, J. R., Singh, S. K., & Wahajuddin, M. (2015). Simultaneous determination of centchroman and tamoxifen along with their metabolites in rat plasma using LC-MS/MS. *Bioanalysis*, *7*(8), 967–979. https://doi.org/10.4155/bio.14.253

Sato, K., Akaishi, T., Matsuki, N., Ohno, Y., & Nakazawa, K. (2007). β-Estradiol induces synaptogenesis in the hippocampus by enhancing brain-derived neurotrophic factor release from dentate gyrus granule cells. *Brain Research*, *1150*, 108–120. https://doi.org/10.1016/j.brainres.2007.02.093

Scharfman, H. E., & MacLusky, N. J. (2006). Estrogen and brain-derived neurotrophic factor (BDNF) in hippocampus: Complexity of steroid hormone-growth factor interactions in the adult CNS. *Frontiers in Neuroendocrinology*, *27*(4), 415–435. https://doi.org/10.1016/j.yfrne.2006.09.004

Scharfman, H. E., & MacLusky, N. J. (2014). Differential regulation of BDNF, synaptic plasticity and sprouting in the hippocampal mossy fiber pathway of male and female rats. *Neuropharmacology*, *76*, 696–708. https://doi.org/10.1016/j.neuropharm.2013.04.029

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D. J., Hartenstein, V., Eliceiri, K., Tomancak, P., & Cardona, A. (2012). Fiji: An open-source platform for biological-image analysis. *Nature Methods*, *9*(7), Article 7. https://doi.org/10.1038/nmeth.2019

Smith, B. M., Saulsbery, A. I., Sarchet, P., Devasthali, N., Einstein, D., & Kirby, E. D. (2022). Oral and Injected Tamoxifen Alter Adult Hippocampal Neurogenesis in Female and Male Mice. *ENeuro*, *9*(2), ENEURO.0422-21.2022. https://doi.org/10.1523/ENEURO.0422-21.2022

Sohrabji, F., & Lewis, D. K. (2006). Estrogen–BDNF interactions: Implications for neurodegenerative diseases. *Frontiers in Neuroendocrinology*, *27*(4), 404–414.

https://doi.org/10.1016/j.yfrne.2006.09.003

Spencer, J. L., Waters, E. M., Romeo, R. D., Wood, G. E., Milner, T. A., & McEwen, B. S.

(2008). Uncovering the Mechanisms of Estrogen Effects on Hippocampal Function.

Frontiers in Neuroendocrinology, *29*(2), 219–237.

https://doi.org/10.1016/j.yfrne.2007.08.006

Sun, C., Krimm, R., & Hill, D. L. (2018). Maintenance of Mouse Gustatory Terminal Field

Organization Is Dependent on BDNF at Adulthood. *The Journal of Neuroscience*,

38(31), 6873–6887. https://doi.org/10.1523/JNEUROSCI.0802-18.2018

Taliaz, D., Stall, N., Dar, D. E., & Zangen, A. (2010). Knockdown of brain-derived

neurotrophic factor in specific brain sites precipitates behaviors associated with

depression and reduces neurogenesis. *Molecular Psychiatry*, *15*(1), 80–92.

https://doi.org/10.1038/mp.2009.67

The jamovi project (2022). jamovi (Version 2.3) [Computer Software]. Retrieved from

https://www.jamovi.org

Underwood, E. A., Rochon, P. A., Moineddin, R., Lee, P. E., Wu, W., Pritchard, K. I., &

Tierney, M. C. (2018). Cognitive sequelae of endocrine therapy in women treated

for breast cancer: A meta-analysis. *Breast Cancer Research and Treatment*,

168(2), 299–310. https://doi.org/10.1007/s10549-017-4627-4

Valvassori, S. S., Borges, C. P., Varela, R. B., Bavaresco, D. V., Bianchini, G., Mariot,

E., Arent, C. O., Resende, W. R., Budni, J., & Quevedo, J. (2017). The different

effects of lithium and tamoxifen on memory formation and the levels of

neurotrophic factors in the brain of male and female rats. *Brain Research*

Bulletin, *134*, 228–235. https://doi.org/10.1016/j.brainresbull.2017.08.006

Velázquez-Zamora, D. A., Garcia-Segura, L. M., & González-Burgos, I. (2012). Effects

of selective estrogen receptor modulators on allocentric working memory

performance and on dendritic spines in medial prefrontal cortex pyramidal

neurons of ovariectomized rats. *Hormones and Behavior*, *61*(4), 512–517.

https://doi.org/10.1016/j.yhbeh.2012.01.010

Wade, G. N., & Heller, H. W. (1993). Tamoxifen mimics the effects of estradiol on food

intake, body weight, and body composition in rats. *American Journal of*

Physiology-Regulatory, Integrative and Comparative Physiology, *264*(6), R1219–

R1223. https://doi.org/10.1152/ajpregu.1993.264.6.R1219

Walf, A. A., & Frye, C. A. (2006). A Review and Update of Mechanisms of Estrogen in

the Hippocampus and Amygdala for Anxiety and Depression Behavior.

Neuropsychopharmacology, *31*(6), Article 6.

https://doi.org/10.1038/sj.npp.1301067

Walker, E. A., Foley, J. J., Clark-Vetri, R., & Raffa, R. B. (2011). Effects of repeated

administration of chemotherapeutic agents tamoxifen, methotrexate, and 5-

fluorouracil on the acquisition and retention of a learned response in mice.

Psychopharmacology, *217*(4), 539–548. https://doi.org/10.1007/s00213-011-

2310-8

Woolley, C. S. (1998). Estrogen-mediated structural and functional synaptic plasticity in

the female rat hippocampus. *Hormones and Behavior*, *34*(2), 140–148.

https://doi.org/10.1006/hbeh.1998.1466

Xu, B., & Xie, X. (2016). Neurotrophic factor control of satiety and body weight. *Nature*

Reviews Neuroscience, *17*(5), 282–292. https://doi.org/10.1038/nrn.2016.24

Figures and Tables

<mark>**Figure 1. Body Weight.** Body weight was tracked throughout the course of the</mark>

<mark>experiment. Starting at Week 6, Tamoxifen pellets were mashed in a flavor-enhanced</mark>

<mark>gel in addition to *ad libitum* presentation to prevent further weight loss. Repeated</mark>

<mark>measures factorial ANOVAs were used to look for interactions and main effects of drug</mark>

<mark>condition (TAM vs. CON) and time (Week) on body weight. There was a significant</mark>

<mark>interaction between drug condition and time on body weight. Post-hoc tests found that</mark>

<mark>starting at Week 3, TAM animals weighed significantly more than CON animals. Data</mark>

<mark>presented as mean ± standard error, ***significant interaction, $p < 0.001$; # significant</mark>

<mark>post hoc test, $p < 0.001$.</mark>

2 **Figure 2. Mass Spectrometry Analysis of Tamoxifen.** Tamoxifen and tamoxifen-d5

3 were separated by ultra-performance liquid chromatography (UPLC) and detected using

4 a triple quadrupole mass spectrometer. Tandem mass spectrometry (MS-MS)

5 parameters were optimized in the positive ionization mode and the multiple reaction

monitoring (MRM) transition of m/z 372.3 72.0 was used for tamoxifen analysis.

Tamoxifen-d5 (m/z 377.3 72.0) was used as an internal standard. A) A calibration curve

for tamoxifen was prepared in human plasma and was linear over the range of 0.1 –

250 ng/mL. Analytes were separated using a gradient elution, with retention time of 2.68

min for tamoxifen and tamoxifen-d5. B) Representative chromatogram from Tamoxifen-

treated animal of Tamoxifen (left) and Tamoxifen-d5 (right). C) Representative

chromatogram from control-treated rat (standard chow) of Tamoxifen (left) and

Tamoxifen-d5 (right) in plasma.

<mark>Figure 3. BDNF Expression in Hippocampal Subregions.</mark> Representative

photomicrographs showing immunohistochemical localization of BDNF in the dentate

gyrus (DG), CA1, <mark>medial CA3 (mCA3) and lateral CA3 (lCA3)</mark>. For each region of

interest, five sections (S1-S5) capturing the rostral to caudal extent of the hippocampus

were analyzed. Regions of interest are identified on the corresponding template atlas

plate and traced in cyan on the photomicrograph.

Figure 4. Quantification of BDNF Optical Density in Hippocampal Subregions. For each brain region of interest, immunohistochemical localization of BDNF was quantified using background-subtracted optical density in five sections capturing the rostral to caudal extent of the hippocampus. Significant interactions and/or main effects of drug condition (Tamoxifen, TAM vs. Control, CON) and section number (1-5) on BDNF optical density were detected using two-way ANOVAs. A) In DG, there was a significant main effect of drug condition on BDNF-ir density. Post-hoc tests revealed that BDNF-ir density was lower in TAM animals than CON animals in Section 4. B) In CA1, there was a main effect of drug condition on the density of BDNF-ir cells. Post hoc tests revealed that BDNF-ir density was lower in TAM animals than CON animals in section 2. C) In

mCA3, there was also a significant main effect of drug condition on the density of

BDNF-ir cells. Post-hoc tests revealed that BDNF-ir was lower in TAM animals than

CON animals in Section 3. D) Finally, in lCA3, there were no interactions or main effects

of drug condition or section level on BDNF immunoreactivity. Data are presented as

mean ± standard error, asterisks indicate significant omnibus tests, * p < 0.05; ** p <

0.01; *** p < 0.001. Pound signs indicate significant post-hoc differences between drug

conditions, # p < 0.05; ## p < 0.01; ### p < 0.001

Figure 5. Plasma Tamoxifen and Brain BDNF Correlations. The relationship

between plasma tamoxifen levels and BDNF expression in hippocampal subregions

was investigated using simple correlations. In all four brain regions of interest, plasma

tamoxifen was negatively correlated with brain BDNF levels. Although none of these

correlations reached significance, trends towards significance were detected in C)

mCA3 and D) ICA3. Pearson's correlation coefficient (r) and p values for each brain

area presented.

1   <mark>**Table 1**. **Formula Composition of Medicated and Non-Medicated Pellets.** *for</mark>

2   detailed formulation of 2016 Teklad Global 16% Protein Rodent Diet, see Inotiv.com.

	Tamoxifen Diet (g/Kg)	Control Diet (g/Kg)
2016, Teklad Global 16% Protein Rodent Diet*	949.75	1000
Sucrose	49.96	0.00
Tamoxifen USP	.04	0.00
Red Food Color	0.25	0.00

2 **Table 2. Summary of Tamoxifen calibration curve in plasma.** A calibration curve for

3 tamoxifen was prepared in human plasma and was linear over the range of 0.1 – 250

4 ng/mL with coefficient of regression, $r^2 > 0.99$. Data presented as Mean, Standard

5 Deviation (SD), Accuracy (%), and Coefficient of Variation (CV %).

Calibration Curve (n=6)	Tamoxifen Concentration (ng/mL)									
	0.1	0.5	1	2.5	5.0	10	25	50	100	250
1	0.108	0.538	1.00	2.61	5.11	10.5	29.4	46.4	108	270
2	0.117	0.533	1.11	2.61	4.84	10.3	27.4	48.4	107	253
3	0.104	0.484	0.95	2.69	4.79	10.2	29.2	54.0	109	247
4	0.090	0.479	0.88	2.08	4.25	9.55	25.2	47.9	95.6	266
5	0.091	0.566	0.922	2.49	5.06	10.5	27.9	52.9	92.1	234
6	0.092	0.467	0.841	2.18	4.25	9.38	24.8	46.5	97.2	230
Mean	0.100	0.511	0.949	2.44	4.72	10.1	27.3	49.4	101	250
SD	0.011	0.040	0.096	0.253	0.382	0.487	1.95	3.29	7.35	16.3
Accuracy (%)	100	102	94.9	97.7	94.3	101	109	98.7	101	100
CV (%)	11.1	7.79	10.1	10.4	8.09	4.84	7.15	6.66	7.25	6.52

6
7

Table 3. Human plasma and rat plasma comparison. The Tamoxifen assay was

initially developed in human plasma. Human plasma and rat plasma were compared

using 100 ng/mL concentrations of tamoxifen. Results were comparable between

human and rat plasma. Concentration (ng/mL), Data presented as Concentration

(ng/mL), Mean, Standard Deviation (SD), Coefficient of Variation (% CV), and Accuracy

(%).

	Concentration (ng/mL)	Mean ± SD	% CV	% Accuracy
Human Plasma	100	101 ± 7.35	7.25	101
Rat Plasma	100	97.62 ± 2.01	2.06	97.6

1   **Table 4. Plasma Tamoxifen Concentration.** Tamoxifen levels (ng/mL) in TAM-treated

2   rats (n = 10).

Subject	Measured Tamoxifen Concentration (ng/mL)
TAM 1	48.3
TAM 2	33.8
TAM 3	26.5
TAM 4	35.0
TAM 5	22.0
TAM 6	34.2
TAM 9	16.3
TAM 10	10.7
TAM 11	35.0
TAM 12	19.3

1   **Supplementary Figure 1. Body Condition Scoring.** In addition to weight, Body

2   Condition (BC) Scoring was used as a measure of animal health. Adapted from:

3   Hickman D, Swan M. 2010. Use of a Body Condition Score Technique to Assess Health

4   Status in a Rat Model of Polycystic Kidney Disease.

BC 1

Rat is emaciated

- Segmentation of vertebral column prominent if not visible
- Little or no flesh cover over dorsal pelvis, pins prominent if not visible
- Segmentation of caudal vertebrae prominent

BC 2

Rat is under-conditioned

- Segmentation of vertebral column prominent
- Thin flesh cover over dorsal pelvis, little subcutaneous fat, pins easily palpable
- Thin flesh cover over caudal vertebrae, segmentation palpable with slight pressure

BC 3

Rat is well-conditioned

- Segmentation of vertebral column easily palpable
- Moderate subcutaneous fat store over pelvis, pins easily palpable with slight pressure
- Moderate fat store around tail base, caudal vertebrae may be palpable but not segmented

BC 4

Rat is over-conditioned

☐ Segmentation of vertebral column palpable with slight pressure

☐ Thick subcutaneous fat store over dorsal pelvis, pins of pelvis palpable with firm pressure

☐ Thick fat store over tail base, caudal vertebrae not palpable

BC 5

Rat is obese

☐ Segmentation of vertebral column palpable with firm pressure, may be a continuous column

☐ Thick subcutaneous fat store over dorsal pelvis, pins of pelvis not palpable with firm pressure

☐ Thick fat store over tail base, caudal vertebrae not palpable